insects

Special Issue Reprint

RNAi in Insect Physiology

Edited by
Jisheng Liu and Luc Swevers

mdpi.com/journal/insects

MDPI

RNAi in Insect Physiology

RNAi in Insect Physiology

Guest Editors

Jisheng Liu
Luc Swevers

MDPI

Basel • Beijing • Wuhan • Barcelona • Belgrade • Novi Sad • Cluj • Manchester

Guest Editors

Jisheng Liu
School of Life Sciences
Guangzhou University
Guangzhou
China

Luc Swevers
Institute of Biosciences
& Applications
National Centre for Scientific
Research "Demokritos"
Athens
Greece

Editorial Office
MDPI AG
Grosspeteranlage 5
4052 Basel, Switzerland

This is a reprint of the Special Issue, published open access by the journal *Insects* (ISSN 2075-4450), freely accessible at: https://www.mdpi.com/journal/insects/special_issues/H0T9U70D3V.

For citation purposes, cite each article independently as indicated on the article page online and as indicated below:

Lastname, A.A.; Lastname, B.B. Article Title. *Journal Name* **Year**, *Volume Number*, Page Range.

ISBN 978-3-7258-7464-4 (Hbk)
ISBN 978-3-7258-7465-1 (PDF)
https://doi.org/10.3390/books978-3-7258-7465-1

© 2026 by the authors. Articles in this reprint are Open Access and distributed under the Creative Commons Attribution (CC BY) license. The reprint as a whole is distributed by MDPI under the terms and conditions of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) license (https://creativecommons.org/licenses/by-nc-nd/4.0/).

Contents

About the Editors . . . . . **vii**

Jisheng Liu and Luc Swevers
RNAi in Insect Physiology: Unlocking Mechanisms and Pioneering Sustainable Pest Control
Reprinted from: *Insects* **2026**, *17*, 333, https://doi.org/10.3390/insects17030333 . . . . . **1**

Chaoshan Liang, Jiahui Song, Kangkang Xu, Fanghao Wan, Guy Smagghe and Wenjia Yang
Functional Characterization of Akt, a Serine/Threonine Kinase, in *Tuta absoluta*: A Key Regulator of Molting, Hormones, and Reproduction with Potential for RNAi Pest Control
Reprinted from: *Insects* **2026**, *17*, 183, https://doi.org/10.3390/insects17020183 . . . . . **6**

Kangkang Xu, Yue Wang, Shuyan Yan, Fanghao Wan, Guy Smagghe and Wenjia Yang
Identification and Functional Characterization of Chitinase Genes During Larva–Pupa–Adult Transitions in *Tuta absoluta*
Reprinted from: *Insects* **2026**, *17*, 114, https://doi.org/10.3390/insects17010114 . . . . . **23**

Man-Hong Cheng, Kai-Yun Fu, Wei Zhou, Ji-Feng Shi and Wen-Chao Guo
Silencing *Miniature* Gene Disrupts Elytral and Hindwing Structures in *Leptinotarsa decemlineata*
Reprinted from: *Insects* **2025**, *16*, 700, https://doi.org/10.3390/insects16070700 . . . . . **39**

Xiaodong Deng, Changhao He, Chunmei Xue, Dianlong Xu, Juncai Li and Xiaowen Fei
Silencing the *cyp314a1* and *cyp315a1* Genes in the *Aedes albopictus* 20E Synthetic Pathway for Mosquito Control and Assessing Algal Blooms Induced by Recombinant RNAi *Microalgae*
Reprinted from: *Insects* **2025**, *16*, 1033, https://doi.org/10.3390/insects16101033 . . . . . **53**

Xinyi Liu, Guoqiang Zhao, Hui Liu, Yuxuan Mao, Meng Xu, Jing Wu, et al.
Elimination of Ultraviolet Light-Mediated Attraction Behavior in *Culex* Mosquitoes via dsRNA-Mediated Knockdown of Opsins
Reprinted from: *Insects* **2025**, *16*, 997, https://doi.org/10.3390/insects16100997 . . . . . **84**

Yiyun Wei, Xinyao Gu, Fengling Si, Xiaojie Chen, Liang Qiao, Hongxing Yan and Bin Chen
Transcriptome and Functional Analyses Revealed the Carboxylesterase Genes Involved in Pyrethroid Resistance in *Anopheles sinensis* (Diptera: Culicidae)
Reprinted from: *Insects* **2025**, *16*, 938, https://doi.org/10.3390/insects16090938 . . . . . **99**

Xue Han, Yulong Jia, Changchun Dai, Xiaoyun Wang, Jian Liu and Zhenqi Tian
Expression of Heat Shock Protein 90 Genes Induced by High Temperature Mediated Sensitivity of *Aphis glycines* Matsumura (Hemiptera: Aphididae) to Insecticides
Reprinted from: *Insects* **2025**, *16*, 772, https://doi.org/10.3390/insects16080772 . . . . . **116**

Kai Li, Tongtong Chen, Yuliang Li, Kai Sun, Kun Pang, Xiaoping Yu and Peiying Hao
Risk Assessment of RNAi-Based Potential Pesticide ds*NlAtg3* and Its Homologues for *Nilaparvata lugens* and Non-Target Organisms
Reprinted from: *Insects* **2025**, *16*, 225, https://doi.org/10.3390/insects16020225 . . . . . **128**

Qitong Huang, Qian Ma, Xiaobin Liu, Keyan Zhu-Salzman and Weining Cheng
Characterization and Functional Analysis of Small Heat Shock Protein Genes (*Hsp22.2* and *Hsp26.7*) in *Sitodiplosis mosellana* Diapause
Reprinted from: *Insects* **2025**, *16*, 649, https://doi.org/10.3390/insects16070649 . . . . . **142**

Bin Tang, Yuxin Ge, Yongkang Liu, Liwen Guan, Ye Han, Yang Zhu, et al.
Effects of Trehalase on the Gene Expression of the Reproductive Regulation Pathway Network and Triglyceride Metabolism in *Nilaparvata lugens* (Stål)
Reprinted from: *Insects* **2025**, *16*, 725, https://doi.org/10.3390/insects16070725 . . . . . **158**

Jisheng Liu, Weijian Chen, Minchun Lai, Jiahua Chen and Luc Swevers
Activation of *BmToll9-1* in Silkworm (*Bombyx mori*) Larval Midgut by *Escherichia coli* and Regulation of Growth
Reprinted from: *Insects* **2025**, *16*, 621, https://doi.org/10.3390/insects16060621 . . . . . **172**

Zelong Yang, Yan Lu, Zhuo Jiang, Xilin Jiao, Han Lin, Wanning Jiang, et al.
Development of a Stage- and Species-Specific RNAi System for Molecular Insights in *Trichogramma* Wasps
Reprinted from: *Insects* **2025**, *16*, 673, https://doi.org/10.3390/insects16070673 . . . . . **184**

Lei Wang, Yujie Lu and Zongpei Zhao
Odorant Binding Proteins in *Tribolium castaneum*: Functional Diversity and Emerging Applications
Reprinted from: *Insects* **2025**, *16*, 1250, https://doi.org/10.3390/insects16121250 . . . . . **200**

Hemant Kumar, Nikita Gal'chinsky, Verma Sweta, Nikita Negi, Roman Filatov, Anamika Chandel, et al.
Perspectives of RNAi, CUADb and CRISPR/Cas as Innovative Antisense Technologies for Insect Pest Control: From Discovery to Practice
Reprinted from: *Insects* **2025**, *16*, 746, https://doi.org/10.3390/insects16070746 . . . . . **218**

About the Editors

Jisheng Liu

Jisheng Liu is a Professor at the School of Life Sciences at Guangzhou University. He is a high-level talent in Guangzhou and a Pearl River Rising Star in Science and Technology. He completed his Ph.D. studies in Applied Biological Sciences at Ghent University. His research interests include innate immunity and RNA interference (RNAi) in model insects, such as the silkworm *Bombyx mori*, the red flour beetle *Tribolium casteneum*, and other insect species.

Luc Swevers

Luc Swevers is Research Director at the Institute of Biosciences & Applications, National Centre for Scientific Research "Demokritos" in Athens, Greece. He completed his Ph.D. studies in Zoology at K.U. Leuven, Belgium. His research interests include RNA interference (RNAi) and antiviral defence mechanisms in the silkworm, *Bombyx mori*.

insects

MDPI

Editorial

RNAi in Insect Physiology: Unlocking Mechanisms and Pioneering Sustainable Pest Control

Jisheng Liu [1,*] and Luc Swevers [2]

[1] School of Life Sciences, Guangzhou University, Guangzhou 510006, China
[2] Institute of Biosciences and Applications, National Centre for Scientific Research Demokritos, 15341 Athens, Greece
* Correspondence: jisheng.liu@gzhu.edu.cn

1. Unveiling the RNAi Landscape: From Core Mechanisms to Applied Frontiers

The advent of RNA interference (RNAi) in 1998, marked by the discovery of gene silencing triggered by double-stranded RNA (dsRNA) in nematodes, opened a transformative chapter in molecular biology [1]. Today, RNAi has matured into an indispensable tool for elucidating gene function, while its potential as a targeted, eco-friendly pest control strategy is reshaping entomological research and agricultural practices [2]. This Special Issue of Insects, dedicated to "RNAi in Insect Physiology", stands as a testament to the dynamic progress of RNAi research, bridging fundamental insights into insect physiology with groundbreaking applications in pest management.

At its core, RNAi encompasses a sophisticated network of gene silencing pathways, driven by small RNAs, including small interfering RNAs (siRNAs), microRNAs (miRNAs), and PIWI-interacting RNAs (piRNAs). These pathways collectively fine-tune gene expression in eukaryotic cells, offering precise molecular control over biological processes [3]. What makes RNAi particularly compelling in entomology is its unique mechanism of sequence-specific gene silencing, a feature that not only revolutionizes gene functional studies but also addresses the urgent need for sustainable pest control as traditional chemical pesticides face growing challenges of resistance and environmental harm [4].

This Special Issue was conceived to capture the full spectrum of RNAi research in insects with a dual focus: advancing our understanding of gene function in insect physiology and accelerating the translation of RNAi into practical pest control solutions. The 14 contributions featured here collectively paint a vivid picture of RNAi's power, unlocking the molecular underpinnings of insect development, reproduction, and stress adaptation, while simultaneously demonstrating its viability as a next-generation pest management tool.

2. Highlights of the Special Issue: Bridging Mechanism and Impact

This Special Issue brings together 12 cutting-edge research articles and 2 insightful review articles, collectively showcasing the multifaceted power of RNAi in unraveling insect physiology and driving innovative pest control strategies.

On the front of gene function and pest control target discovery, key studies make significant strides. Research on the tomato leafminer *Tuta absoluta* identifies Akt, a serine/threonine kinase, as a central regulator of molting, hormones, and reproduction, highlighting its potential as an RNAi pest control target (Contribution 1). Complementing this, another work on the same pest characterizes 11 chitinase genes critical for larva–pupa–adult transitions, establishing them as promising targets for RNAi-based insecticidal

development (Contribution 2). For the Colorado potato beetle *Leptinotarsa decemlineata*, silencing the *Miniature* gene effectively disrupts elytral and hindwing structures, offering a novel RNAi target to curb its dispersal and resistance (Contribution 3).

The issue also advances RNAi application in vector and pest management. For *Aedes albopictus*, a key Dengue fever vector, silencing genes in the 20E synthetic pathway demonstrate effective mosquito control, while assessing the environmental impacts of recombinant RNAi microalgae delivery (Contribution 4). In *Culex* mosquitoes, dsRNA-mediated knockdown of opsins eliminates UV-light-mediated attraction, enhancing trap efficacy (Contribution 5). Additionally, a study on *Anopheles sinensis*, a major malaria vector, reveals the involvement of carboxylesterase genes in pyrethroid resistance through transcriptome and functional analyses, providing a molecular basis for combating insecticide resistance in disease vectors (Contribution 6). Research on the soybean aphid *Aphis glycines* links high-temperature-induced heat shock protein 90 expression to insecticide sensitivity (Contribution 7), and studies on the brown planthopper *Nilaparvata lugens* conduct risk assessments for RNAi-based pesticides, ensuring safety for both target pests and non-target organisms (Contribution 8).

Beyond control, the issue deepens understanding of insect physiological adaptation and molecular mechanisms. Work on the wheat pest *Sitodiplosis mosellana* (Diptera: Cecidomyiidae) characterizes small heat shock protein genes involved in diapause, shedding light on stress adaptation (Contribution 9). For the brown planthopper, the study clarifies how trehalase regulates reproductive pathways and triglyceride metabolism, explaining its high reproductive capacity (Contribution 10). In the silkworm *Bombyx mori*, research reveals how *Escherichia coli* activates *BmToll9-1* to regulate growth, advancing insights into insect immune regulation (Contribution 11), a study that has completed the systematic investigation of the silkworm *BmToll9s* gene [5–8]. Moreover, a breakthrough study in *Trichogramma* wasps reports a stage- and species-specific RNAi system, overcoming technical barriers for molecular studies in these minute parasitoids (Contribution 12).

The two review articles provide critical context and forward-looking perspectives. One focuses on odorant binding proteins in *Tribolium castaneum*, exploring their functional diversity beyond olfaction and emerging applications in pest control (Contribution 13). The other offers a comprehensive overview of RNAi, contact unmodified antisense DNA biotechnology (CUADb), and clustered regularly interspaced short palindromic repeats (CRISPR) and CRISPR-associated proteins (CRISPR/Cas) as innovative silencing technologies, outlining their synergies and potential to transform pest management from discovery to practice (Contribution 14).

Together, these contributions bridge fundamental RNAi mechanisms with tangible impact, advancing both basic entomological research and practical pest control solutions.

3. Looking Ahead: The Promise of RNAi in Entomology

As we reflect on the contributions in this Special Issue, it is clear that RNAi is not just a research tool; it is a catalyst for innovation in entomology [9]. The studies presented here have expanded our understanding of insect physiology, identified novel targets for pest control, and developed new methodologies to harness RNAi's potential. However, the journey is far from complete.

Looking ahead, several critical directions demand further exploration. First, improving the efficiency and specificity of RNAi delivery remains a key challenge, particularly for insects with diverse physiological barriers [2]. Advances in nanotechnology, microalgae-based delivery, and other innovative platforms will be crucial to overcoming these hurdles [10,11]. Second, understanding the off-target effects and environmental impacts of RNAi-based pest control strategies is essential to ensure their safe and sustain-

https://doi.org/10.3390/insects17030333

able deployment [12]. Third, integrating RNAi with other emerging technologies, such as CRISPR/Cas, synthetic biology, and ecological modeling, will unlock new synergies, enabling more comprehensive and resilient pest management solutions [13]. Fourth, dsRNA, as a pathogen-associated molecular pattern molecule, has the potential to interact with other immune defense mechanisms, which may complicate its use in pest control [14].

Moreover, as climate change and global trade continue to alter pest distributions and dynamics, the need for adaptive, environmentally friendly pest control strategies has never been greater. RNAi, with its specificity, scalability, and potential for integration into integrated pest management systems, is uniquely positioned to meet this challenge, embodying the core vision of One Health by safeguarding the health of agricultural ecosystems, thereby bridging crop protection with broader environmental and public health priorities [15].

4. Closing Remarks

This Special Issue on "RNAi in Insect Physiology" captures a pivotal moment in the evolution of RNAi research, a moment where fundamental insights converge with practical applications to address some of the most pressing challenges in entomology. The work presented here not only advances our knowledge of insect molecular biology but also paves the way for a new era of sustainable pest management, one that prioritizes ecological balance and human well-being.

We hope that this collection will inspire further research, foster collaboration across disciplines, and accelerate the translation of RNAi from the laboratory to the field. As the field of entomology continues to evolve, RNAi will undoubtedly remain at the forefront of innovation, driving discoveries that protect crops, safeguard public health, and preserve ecosystems.

We thank all our readers for engaging with this Special Issue and invite the scientific community to continue pushing the boundaries of RNAi research. Together, we can harness the power of this remarkable technology to build a more sustainable future for agriculture and the environment.

Funding: This research received no external funding.

Data Availability Statement: No datasets were generated or analyzed.

Acknowledgments: We sincerely thank all of the authors for their valuable contributions, the reviewers for their constructive evaluations, and the editorial staff of Insects for their support throughout the preparation of this Special Issue.

Conflicts of Interest: The authors declare no conflicts of interest.

List of Contributions:

1. Liang, C.; Song, J.; Xu, K.; Wan, F.; Smagghe, G.; Yang, W. Functional Characterization of Akt, a Serine/Threonine Kinase, in *Tuta absoluta*: A Key Regulator of Molting, Hormones, and Reproduction with Potential for RNAi Pest Control. *Insects* **2026**, *17*, 183. https://doi.org/10.3390/insects17020183.
2. Xu, K.; Wang, Y.; Yan, S.; Wan, F.; Smagghe, G.; Yang, W. Identification and Functional Characterization of Chitinase Genes During Larva-Pupa-Adult Transitions in *Tuta absoluta*. *Insects* **2026**, *17*, 114. https://doi.org/10.3390/insects17010114.
3. Cheng, M.H.; Fu, K.Y.; Zhou, W.; Shi, J.F.; Guo, W.C. Silencing *Miniature* Gene Disrupts Elytral and Hindwing Structures in *Leptinotarsa decemlineata*. *Insects* **2025**, *16*, 700. https://doi.org/10.3390/insects16070700.
4. Deng, X.; He, C.; Xue, C.; Xu, D.; Li, J.; Fei, X. Silencing the *cyp314a1* and *cyp315a1* Genes in the *Aedes albopictus* 20E Synthetic Pathway for Mosquito Control and Assessing Algal Blooms Induced by Recombinant RNAi *Microalgae*. *Insects* **2025**, *16*, 1033. https://doi.org/10.3390/insects16101033.

5. Liu, X.; Zhao, G.; Liu, H.; Mao, Y.; Xu, M.; Wu, J.; Li, L.; Zhai, Z.; Wu, P. Elimination of Ultraviolet Light-Mediated Attraction Behavior in *Culex* Mosquitoes via dsRNA-Mediated Knockdown of Opsins. *Insects* **2025**, *16*, 997. https://doi.org/10.3390/insects16100997.
6. Wei, Y.; Gu, X.; Si, F.; Chen, X.; Qiao, L.; Yan, H.; Chen, B. Transcriptome and Functional Analyses Revealed the Carboxylesterase Genes Involved in Pyrethroid Resistance in *Anopheles sinensis* (Diptera: Culicidae). *Insects* **2025**, *16*, 938. https://doi.org/10.3390/insects16090938.
7. Han, X.; Jia, Y.; Dai, C.; Wang, X.; Liu, J.; Tian, Z. Expression of Heat Shock Protein 90 Genes Induced by High Temperature Mediated Sensitivity of *Aphis glycines* Matsumura (Hemiptera: Aphididae) to Insecticides. *Insects* **2025**, *16*, 772. https://doi.org/10.3390/insects16080772.
8. Li, K.; Chen, T.; Li, Y.; Sun, K.; Pang, K.; Yu, X.; Hao, P. Risk Assessment of RNAi-Based Potential Pesticide ds*NlAtg3* and Its Homologues for *Nilaparvata lugens* and Non-Target Organisms. *Insects* **2025**, *16*, 225. https://doi.org/10.3390/insects16020225.
9. Huang, Q.; Ma, Q.; Liu, X.; Zhu-Salzman, K.; Cheng, W. Characterization and Functional Analysis of Small Heat Shock Protein Genes (*Hsp22.2* and *Hsp26.7*) in *Sitodiplosis mosellana* Diapause. *Insects* **2025**, *16*, 649. https://doi.org/10.3390/insects16070649.
10. Tang, B.; Ge, Y.; Liu, Y.; Guan, L.; Han, Y.; Zhu, Y.; Hu, G.; Wu, Y. Effects of Trehalase on the Gene Expression of the Reproductive Regulation Pathway Network and Triglyceride Metabolism in *Nilaparvata lugens* (Stål). *Insects* **2025**, *16*, 725. https://doi.org/10.3390/insects16070725.
11. Liu, J.; Chen, W.; Lai, M.; Chen, J.; Swevers, L. Activation of *BmToll9-1* in Silkworm (*Bombyx mori*) Larval Midgut by *Escherichia coli* and Regulation of Growth. *Insects* **2025**, *16*, 621. https://doi.org/10.3390/insects16060621.
12. Yang, Z.; Lu, Y.; Jiang, Z.; Jiao, X.; Lin, H.; Jiang, W.; Du, W.; Zhang, X.; Peng, Z.; Zhang, J.; et al. Development of a Stage- and Species-Specific RNAi System for Molecular Insights in *Trichogramma* Wasps. *Insects* **2025**, *16*, 673. https://doi.org/10.3390/insects16070673.
13. Wang, L.; Lu, Y.; Zhao, Z. Odorant Binding Proteins in *Tribolium castaneum*: Functional Diversity and Emerging Applications. *Insects* **2025**, *16*, 1250. https://doi.org/10.3390/insects16121250.
14. Kumar, H.; Gal'chinsky, N.; Sweta, V.; Negi, N.; Filatov, R.; Chandel, A.; Ali, J.; Oberemok, V.; Laikova, K. Perspectives of RNAi, CUADb and CRISPR/Cas as Innovative Antisense Technologies for Insect Pest Control: From Discovery to Practice. *Insects* **2025**, *16*, 746. https://doi.org/10.3390/insects16070746.

References

1. Fire, A.; Xu, S.; Montgomery, M.K.; Kostas, S.A.; Driver, S.E.; Mello, C.C. Potent and specific genetic interference by double-stranded RNA in Caenorhabditis elegans. *Nature* **1998**, *391*, 806–811. [CrossRef] [PubMed]
2. Liu, J.; He, Q.; Lin, X.; Smagghe, G. Recent progress in nanoparticle-mediated RNA interference in insects: Unveiling new frontiers in pest control. *J. Insect Physiol.* **2025**, *167*, 104884. [CrossRef] [PubMed]
3. Siomi, H.; Siomi, M.C. On the road to reading the RNA-interference code. *Nature* **2009**, *457*, 396–404. [CrossRef] [PubMed]
4. Li, Z.; Liu, Y.; Liang, Y.; Pan, T.; Liu, J. RNA interference-based pesticides: Mechanism, application, and commercialization in sustainable pest management. *Pestic. Biochem. Physiol.* **2026**, *219*, 107034. [CrossRef] [PubMed]
5. Liu, J.; Yang, W.; Liao, W.; Huang, Y.; Chen, W.; Bu, X.; Huang, S.; Jiang, W.; Swevers, L. Immunological function of *Bombyx Toll9-2* in the silkworm (*Bombyx mori*) larval midgut: Activation by *Escherichia coli*/lipopolysaccharide and regulation of growth. *Arch. Insect Biochem. Physiol.* **2024**, *116*, e22130. [CrossRef] [PubMed]
6. Liu, J.; Chen, W.; Situ, J.; Li, J.; Chen, J.; Lai, M.; Huang, F.; Li, B. BmToll9-1 is a positive regulator of the immune response in the silkworm *Bombyx mori*. *Insects* **2024**, *15*, 643. [CrossRef] [PubMed]
7. Liu, J.; Chen, W.; Chen, S.; Li, S.; Swevers, L. Similarly to *BmToll9-1*, *BmToll9-2* Is a Positive Regulator of the Humoral Immune Response in the Silkworm, *Bombyx mori*. *Insects* **2024**, *15*, 1005. [CrossRef] [PubMed]
8. Feng, Z.; Li, Z.; He, Q.; Deng, B.; Ma, H.; Chen, W.; Li, X.; Zhang, J.; Swevers, L.; Liu, J. Transcriptional activation of *BmToll9-2* to exogenous dsRNA in the larvae of *Bombyx mori*. *J. Insect Physiol.* **2025**, *165*, 104860. [CrossRef] [PubMed]
9. Zhu, K.Y.; Palli, S.R. Mechanisms, applications, and challenges of insect RNA interference. *Annu. Rev. Entomol.* **2020**, *65*, 293–311. [CrossRef] [PubMed]
10. Liu, J.; Yang, Q.; Yang, Y.; Lin, X. Larval RNA Interference in Silkworm Bombyx mori through Chitosan/dsRNA Nanoparticle Delivery. *J. Vis. Exp.* **2024**, e67360. [CrossRef] [PubMed]
11. Liu, J.; Yang, Y.; Yang, Q.; Lin, X.; Liu, Y.; Li, Z.; Swevers, L. Successful oral RNA interference efficiency in the silkworm *Bombyx mori* through nanoparticle-shielded dsRNA delivery. *J. Insect Physiol.* **2025**, *161*, 104749. [CrossRef]

12. Christiaens, O.; Sweet, J.; Dzhambazova, T.; Urru, I.; Smagghe, G.; Kostov, K.; Arpaia, S. Implementation of RNAi-based arthropod pest control: Environmental risks, potential for resistance and regulatory considerations. *J. Pest Sci.* **2022**, *95*, 1–15. [CrossRef]
13. Faber, N.R.; Ashok, K.; Venkatesan, T.; Wertheim, B.; Bulgarella, M. Leveraging advances in RNAi and CRISPR for improved biological pest control. *Curr. Opin. Insect Sci.* **2026**, *73*, 101453. [CrossRef] [PubMed]
14. Feng, M.; Liu, J.; Wang, L.; Swevers, L. DsRNA as pathogen-associated molecular pattern in innate immunity and multiple functions of the RNAi machinery complicate the use of RNAi in pest control. *Front. Insect Sci.* **2025**, *5*, 1749008. [CrossRef]
15. Liu, J.; Smagghe, G. RNAi-Based Pesticides: Genetic Innovations for Sustainable Crop Protection and One Health. *Adv. Genet.* **2026**, *7*, e00063. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

https://doi.org/10.3390/insects17030333

insects

MDPI

Article

Functional Characterization of Akt, a Serine/Threonine Kinase, in *Tuta absoluta*: A Key Regulator of Molting, Hormones, and Reproduction with Potential for RNAi Pest Control

Chaoshan Liang [1], Jiahui Song [1], Kangkang Xu [1,*], Fanghao Wan [1,2], Guy Smagghe [3,4] and Wenjia Yang [1,*]

[1] Guizhou Key Laboratory of Agricultural Biosecurity, Key Laboratory of Surveillance and Management of Invasive Alien Species in Guizhou Province, College of Biological and Environmental Engineering, Guiyang University, Guiyang 550005, China; liangcs11@163.com (C.L.); songjiahui199701@126.com (J.S.); wanfanghao@caas.cn (F.W.)
[2] Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China
[3] Institute of Entomology, Guizhou University, Guiyang 550025, China; guysma9@gmail.com
[4] Department of Biology, Vrije Universiteit Brussel (VUB), 1050 Brussels, Belgium
* Correspondence: kkxu1988@163.com (K.X.); yangwenjia10@126.com (W.Y.)

Simple Summary

The tomato leafminer is a globally destructive pest that inflicts substantial damage on tomato crops, causing significant economic losses. A comprehensive understanding of the biological mechanisms regulating its growth and reproduction is crucial for the development of sustainable pest management strategies. This study focused on a key gene, *Akt*, which acts as a central switch in the insect's growth and reproductive systems. We identified and characterized the *Akt* gene (*TaAkt*) and then reduced its activity using RNA interference (RNAi). Silencing of *TaAkt* resulted in severe developmental abnormalities: numerous pupae failed to metamorphose into adults, exhibiting weakened body structure, and experienced disruptions in hormone levels that control molting and growth. In female insects, *TaAkt* knockdown led to underdeveloped ovaries, fewer eggs, and lower hatching success. These findings underscore the role of *TaAkt* in the regulation of growth and reproduction in the tomato leafminer. Targeting this single gene presents a promising approach for the development of innovative, environmentally sustainable pest control strategies that reduce both the survival and reproductive ability of these insects without relying on chemical pesticides.

Abstract

Insect insulin signaling plays a central role in regulating development, metamorphosis, and reproduction, yet its mechanistic functions in the tomato leafminer, *Tuta absoluta*, a globally significant pest, remain poorly understood. This study aimed to elucidate the role of the serine/threonine kinase Akt (*TaAkt*) in coordinating metamorphosis and female reproductive processes. The *TaAkt* gene was cloned and characterized, and its spatiotemporal expression was analyzed across various developmental stages and tissues. RNA interference (RNAi) was employed to knock down *TaAkt* in late pupae and newly emerged females, followed by assessment of pupal-adult eclosion, chitin metabolism, 20-hydroxyecdysone (20E) titer, ovarian development, juvenile hormone (JH) levels, vitellogenin synthesis, and fecundity. Knockdown of *TaAkt* significantly reduced 20E titers and downregulated the expression of ecdysone biosynthesis and signaling genes, leading to pupal mortality, defective molting, and reduced chitin content. In adult females, *TaAkt* silencing impaired ovarian growth, decreased JH levels, suppressed vitellogenin production, and reduced egg number and hatching rates. These findings demonstrate that *TaAkt* exerts pleiotropic control over both

metamorphic and reproductive processes in *T. absoluta*. The study identifies *TaAkt* as a promising molecular target for RNAi-based pest management strategies, offering a potential approach to simultaneously suppress survival and reproductive capacity in this economically important pest.

Keywords: tomato leafminer; *Tuta absoluta*; *Akt* gene; insulin signaling pathway; RNA interference; molting; reproduction; juvenile hormone; sustainable pest management

1. Introduction

The South American tomato leafminer, *T. absoluta* (Lepidoptera: Gelechiidae), is one of the most destructive invasive insect pests threatening global tomato production. Native to Peru, *T. absoluta* has rapidly expanded its geographical range over the past two decades and is now established in more than 100 countries across South America, Europe, Africa, and Asia [1–3]. The extraordinary invasive success of *T. absoluta* is attributed to a combination of biological traits, including high fecundity, short generation time, strong environmental adaptability, and a cryptic larval feeding habit within plant tissues, which collectively reduce exposure to natural enemies and control measures [4]. As a consequence, management of this pest remains exceptionally challenging.

The current control strategies rely predominantly on chemical insecticides. However, intensive and prolonged insecticide use has led to the rapid evolution of resistance in *T. absoluta* populations to multiple classes of compounds, including organophosphates, pyrethroids, diamides, and spinosyns [5]. This escalating resistance crisis, together with increasing concerns regarding environmental contamination and non-target effects, underscores the urgent need for alternative, mechanism-based control strategies. From a physiological perspective, a promising avenue lies in identifying and disrupting key endocrine and metabolic pathways that regulate insect growth, development, and reproduction. Elucidating such mechanisms in *T. absoluta* is therefore not only of fundamental interest in insect physiology but also critical for the development of innovative and sustainable pest control approaches.

Nutrient availability and metabolic state are primarily sensed through the insulin/insulin-like growth factor signaling (IIS) pathway, an evolutionarily conserved regulatory system that links environmental conditions to developmental decisions [6]. In insects, IIS is activated by insulin-like peptides (ILPs) that bind to the insulin receptor (InR), triggering a phosphorylation cascade involving phosphoinositide 3-kinase (PI3K) and downstream kinases. This pathway regulates diverse physiological processes, including carbohydrate and lipid metabolism, protein synthesis, cell growth, body size determination, lifespan, and reproductive output [7–10]. A central component of this cascade is Protein Kinase B (Akt), a serine/threonine kinase that serves as a key molecular hub translating insulin signals into coordinated physiological responses.

Akt exerts its regulatory role by phosphorylating multiple downstream targets, including Forkhead box-containing protein (FoxO), phosphoinositide-dependent kinase (PDK), and Target of Rapamycin (TOR) complex, thereby promoting anabolic metabolism and inhibiting catabolic and stress-response pathways under nutrient-rich conditions [11]. Accumulating evidence across insect taxa demonstrates that Akt is indispensable for normal growth, development, and reproduction. In the silkworm *Bombyx mori*, *BmAkt* regulates ecdysteroid biosynthesis and modulates molting processes, while also enhancing detoxification capacity in response to imidacloprid exposure by inducing downstream detoxification enzyme genes [12,13]. In *Drosophila melanogaster*, genetic manipulation of *Akt* profoundly

https://doi.org/10.3390/insects17020183

affects organismal size and metabolism: RNA interference (RNAi)-aided knockdown leads to reduced body weight, cell number, and tissue size, whereas *Akt* overexpression causes excessive cellular proliferation [14,15]. Similarly, in the red flour beetle *Tribolium castaneum*, silencing *TcAkt* results in severe reproductive defects, including reduced fecundity and egg hatchability [16]. In disease-vector mosquitoes such as *Anopheles stephensi* and *Aedes aegypti*, fat body–specific overexpression of *Akt* not only extends lifespan but also alters immune physiology, leading to reduced *Plasmodium falciparum* infection intensity [17]. Together, these studies highlight Akt as a multifunctional regulator integrating nutrition, endocrine signaling, metabolism, immunity, and reproduction.

Against this background, a critical unresolved biological question is how insulin signaling, via Akt, integrates nutritional cues with endocrine regulation to coordinate molting and female reproduction during key developmental transitions in *T. absoluta*. We hypothesize that Akt functions as a central physiological regulator linking IIS with JH and ecdysteroid signaling, vitellogenesis, and cuticle metabolism during the pupal-adult transition. Disruption of *Akt* signaling is therefore expected to impair hormonal balance, structural remodeling, and reproductive capacity. To address this hypothesis, the present study focuses on the functional characterization of the *T. absoluta Akt* gene (*TaAkt*). Specifically, *TaAkt* was cloned, sequenced, and analyzed, and its spatial and temporal expression patterns were determined. RNAi was employed to suppress *TaAkt* expression in female pupae, and the resulting effects on adult emergence, ovarian development, and fertility were systematically evaluated. Furthermore, we examined how *TaAkt* knockdown influences JH and 20E synthesis and signal transduction, vitellogenin expression, and chitin metabolism. By integrating molecular, endocrine, and developmental analyses, this study provides new insights into the physiological role of *TaAkt* in regulating molting and female reproduction in *T. absoluta*. These findings not only advance our understanding of insect growth and reproductive physiology but also identify Akt as a potential target for RNAi-based control strategies against this globally important pest.

2. Materials and Methods

2.1. Experimental Insects

The experimental insects were collected from an organic agricultural base in Kunming City, Yunnan Province, China. The population was maintained for multiple generations in an experimental sunlight room under controlled conditions: temperature of 26 ± 1 °C, relative humidity of 60 ± 5%, and a photoperiod of 16 h light followed by 8 h darkness, as reported before [18]. Larvae of *T. absoluta* were reared on fresh tomato seedlings, which were replaced regularly to ensure adequate nutrition. Adult moths were provided with 10% (*w*/*v*) honey solution as a food source. Insect development and survival were monitored routinely throughout the rearing period.

2.2. Total RNA Extraction and cDNA Synthesis

Samples from different developmental stages, including larvae, female pupae, and female adults, as well as various tissues, were collected for total RNA extraction to support gene cloning and quantitative real-time PCR (qPCR) analyses. All samples were immediately frozen in liquid nitrogen and subsequently ground to a fine powder using a high-speed, low-temperature tissue grinder. Total RNA was extracted following the manufacturer's instructions of the TransZol Up Kit (TransGen Biotech, Beijing, China). RNA integrity was assessed by 1.0% agarose gel electrophoresis, and RNA concentration and purity were determined using a NanoDrop 2000 spectrophotometer (NanoDrop Technologies, Wilmington, MA, USA). High-quality RNA samples were used for subsequent cDNA synthesis and downstream analyses.

https://doi.org/10.3390/insects17020183

2.3. Cloning and Sequence Analysis of TaAkt

The *Akt* gene sequences from *B. mori* (GenBank accession number: XM_038017116.2) and *Helicoverpa armigera* (GenBank accession number: JN399217.1) were retrieved from the National Center for Biotechnology Information (NCBI) database (accessed on 18 December 2025). The putative *T. absoluta Akt* gene (*TaAkt*) was identified by comparative analysis against the *T. absoluta* transcriptome database (Accession number: SRR13065833) using TBtools version 2.156. Gene-specific primers were designed using Primer Premier 5.0 software (Table S1), and *TaAkt* was amplified by polymerase chain reaction (PCR).

PCR reactions were performed in a 50 μL volume containing 25.0 μL of 2× EasyTaq® PCR SuperMix (+dye) (TransGen Biotech, Beijing, China), 21.0 μL of ddH_2O, 2.0 μL of cDNA template, and 1.0 μL each of forward and reverse primers (10 μM). The PCR amplification conditions were as follows: initial denaturation at 94 °C for 3 min; 35 cycles of denaturation at 95 °C for 30 s, annealing at 55 °C for 30 s, and extension at 72 °C for 2 min; and a final extension at 72 °C for 10 min. PCR products were verified by agarose gel electrophoresis, ligated into pGEM-T Easy vector (Promega, Madison, WI, USA), and transformed into *Escherichia coli* Trans5α competent cells (TransGen Biotech, Beijing, China). Positive monoclonal colonies were selected and cultured, and the resulting plasmids were extracted and sent to Guangzhou Ruibo Xingke Biotechnology (Guangzhou, China) for sequencing. Purified plasmids were subsequently used for double-stranded RNA (dsRNA) synthesis.

Sequence homology analysis and conserved domain identification of TaAkt protein were performed using Basic Local Alignment Search Tool (BLAST) (https://blast.ncbi.nlm.nih.gov/Blast.cgi, accessed on 18 December 2025). The physical and chemical properties of the deduced TaAkt protein were predicted using the Compute pI/MW tool (https://web.expasy.org/compute_pi/, accessed on 18 December 2025). Transmembrane domains were analyzed with TMHMM2.0 (https://services.healthtech.dtu.dk/service.php?TMHMM-2.0, accessed on 18 December 2025), and signal peptide prediction was conducted using SignalP 4.1 (https://services.healthtech.dtu.dk/service.php?SignalP-4.1, accessed on 18 December 2025). Phylogenetic analysis was performed using the neighbor-joining method in MEGA version 7.0, with 1000 bootstrap replicates to assess branch reliability [19].

2.4. Developmental and Tissue-Specific Expression Analysis of TaAkt

Fourteen developmental samples were collected to analyze the temporal expression pattern of *TaAkt*, covering the period from the first instar larva to the adult stage. These included first- to fourth-instar larvae (1L–4L), female pupae from day 1 to day 7 (P1–P7), and 1- to 3-day-old adults (A1–A3). Each developmental group consisted of 30 individuals, with three independent biological replicates. For spatial expression analysis, seven tissues, namely head, midgut, fat body, ovary, abdomen, wings, and epidermis, were dissected from adult females after anesthesia with an appropriate amount of CO_2. Fifty individuals were dissected per biological replicate to ensure sufficient tissue quantity, and three biological replicates were performed. All samples were immediately frozen in liquid nitrogen and stored at −80 °C until RNA extraction.

Relative expression levels of *TaAkt* were quantified by qPCR using gene-specific primers (Table S1). *T. absoluta* elongation factor 1-alpha (*TaEF1α*; GenBank accession number: MZ054826) was used as the internal reference gene [20]. The qPCR reactions were performed in a total volume of 20 μL, containing 1 μL of each of forward and reverse primers (10 μM), 10 μL of TransStart® Green qPCR SuperMix (TransGen Biotech, Beijing, China), 1 μL of cDNA template, and 7 μL of ddH_2O. The amplification program consisted of an initial denaturation at 95 °C for 3 min, followed by 41 cycles of 95 °C for 5 s and 55.9 °C for 30 s. A melting curve analysis was conducted from 60 to 95 °C to confirm amplification

https://doi.org/10.3390/insects17020183

specificity. Each reaction was performed with three technical replicates. Relative gene expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method [21].

2.5. RNAi Analysis

Specific dsRNA primers targeting *TaAkt* were meticulously designed using the dsRNAEngineering platform (https://dsrna-engineer.cn/, accessed on 12 June 2025) to minimize off-target effects [22] (Table S1). dsRNA targeting *TaAkt* (ds*TaAkt*) and green fluorescent protein (ds*GFP*, used as a control) were synthesized using the TranscriptAid T7 High Yield Transcription Kit (Thermo Fisher Scientific, Wilmington, DE, USA). The synthesized dsRNA was purified and quantified, then diluted to a final concentration of 2000 ng/μL. For RNAi experiments, 400 ng of dsRNA was injected into each insect using a Nanoliter 2010 microinjector (World Precision Instruments, Sarasota, FL, USA). Injections were performed into the dorsal vessel at the second abdominal segment of 2-day-old female pupae. Following injection, pupae were left undisturbed for 1 h to allow complete absorption of dsRNA, then transferred to rearing containers containing sterile soil. The soil surface was moistened with sterile water, and insects were maintained in an artificial climate chamber under standard rearing conditions. Individual development and survival were monitored daily. Each treatment group consisted of 50 individuals, with three independent biological replicates. To evaluate RNAi efficiency, 30 individuals were collected at 12, 24, 48, and 72 h post-injection for qPCR analysis. Mortality was recorded for up to 7 days after injection in the remaining individuals. Representative lethal phenotypes were documented using a Keyence VHX-6000 digital stereomicroscope (Keyence, Osaka, Japan).

2.6. Effects of TaAkt Knockdown on Pupal-Adult Molting and Expression of Molting-Related Genes

To investigate the role of *TaAkt* in pupal-adult molting, changes in 20E titer and chitin metabolism were analyzed following RNAi treatment. At 48 h after injection of ds*TaAkt* or ds*GFP*, samples (0.05 g per replicate) were collected, with three biological replicates per treatment. Samples were homogenized in 1× phosphate-buffered saline (PBS; pH 7.2–7.4) at a tissue mass-to-buffer volume ratio of 1:10 (*w*/*v*). The homogenates were centrifuged at 10,000× *g* for 20 min at 4 °C, and the supernatants were collected for subsequent analyses. The 20E titer was determined using an Insect Enzyme-linked Immunosorbent Assay (ELISA) Kit (Mlbio, Shanghai, China) following the manufacturer's instructions. Absorbance was measured at 450 nm using a microplate reader after equilibration of reagents to room temperature. Chitin content was assessed using a Chitinase Activity Detection Kit (Solarbio, Beijing, China) according to the manufacturer's protocol. Absorbance was measured at 540 nm using a SpectraMax L microplate reader (Molecular Devices, Sunnyvale, CA, USA). Chitinase activity was calculated based on the standard curve generated for each assay.

To further elucidate the molecular effects of *TaAkt* knockdown on molting, total RNA extracted at 48 h post-injection was used as a template for qPCR analysis. The expression levels of genes involved in ecdysteroid synthesis (*TaCYP306a1*, *TaCYP307a1*, *TaCYP314a1*, and *TaCYP315a1*), ecdysone signal transduction (*TaEcR*, *TaUSP*, *TaE75*, *TaHR3*, *TaHR38*, and *TaFTZ-F1*), and chitin metabolism (*TaCDA1*, *TaCHT5*, *TaUAP*, and *TaCHS*) were quantified to assess the regulatory role of *TaAkt* in pupal-adult transition.

2.7. Effects of TaAkt Knockdown on Ovarian Development and Fertility of T. absoluta

To assess the effects of *TaAkt* knockdown on ovarian development, ovaries were dissected from 2-day-old female adults under a stereomicroscope in 1× phosphate-buffered saline (PBS; Solarbio, Beijing, China). Ovarian morphology was photographed using a Keyence VHX-6000 digital stereomicroscope (Keyence, Osaka, Japan). The length of ovarian tubules and developing oocytes was measured to evaluate ovarian development.

https://doi.org/10.3390/insects17020183

To determine the effects of *TaAkt* silencing on female fertility, ds*GFP*- and ds*TaAkt*-injected pupae were monitored until adult emergence. Female adults within 6 h of eclosion on day 1 were individually transferred to separate rearing containers supplied with fresh tomato leaves. For the mating assays, three males that emerged on the same day were introduced into each container. Mating behavior and oviposition were observed daily, and tomato leaves were replaced every 24 h until female death. The oviposition period (number of days of egg laying per female) was recorded. Eggs deposited on removed leaves were counted daily. The leaves containing eggs were transferred to a rearing chamber and maintained under standard conditions to determine egg hatchability. Total egg number and hatching rate were calculated for each female. Each treatment consisted of 30 mating pairs, with three independent biological replicates.

2.8. Effects of TaAkt Knockdown on JH Synthesis and Vitellogenin Content

To quantify changes in JH titers following *TaAkt* silencing, female pupae from the control (ds*GFP*) and treatment (ds*TaAkt*) groups were collected 48 h post-injection. Each sample weighed 0.05 g, with three biological replicates per group. Samples were rapidly frozen in liquid nitrogen, homogenized in 500 µL of 1× PBS, and centrifuged at 3000× *g* for 20 min at 4 °C. The supernatants were transferred to clean 1.5 mL microcentrifuge tubes for analysis. JH titers were determined using an Insect JH ELISA Kit (Mlbio, Shanghai, China) according to the manufacturer's instructions. Standard, blank, and sample wells were prepared in 96-well plates, and absorbance was measured at 450 nm using a SpectraMax L microplate reader (Molecular Devices, Sunnyvale, CA, USA). The standard curve regression equation was generated using a standard sample provided in the kit, and JH titers were calculated accordingly.

To examine the effect of *TaAkt* knockdown on vitellogenin content, female adults at 1 day post-eclosion were collected from both ds*GFP* and ds*TaAkt* groups. Samples (0.05 g per replicate) were analyzed using an Insect Vitellogenin (VTG) ELISA Kit (Mlbio, Shanghai, China) following the manufacturer's protocol, with three biological replicates. In parallel, insects were collected 48 h after dsRNA injection for gene expression analysis. The relative transcript levels of vitellogenesis-related genes (*TaVg* and *TaVgR*) and JH synthesis and signaling pathway genes (*TaKr-h1*, *TaMet*, and *TaJHAMT*) were quantified by qPCR. Each treatment included 20 individuals per biological replicate, with three biological replicates.

2.9. Statistical Analysis

All data are presented as mean ± standard error (SE) and were analyzed using SPSS 20.0 software (IBM, Chicago, IL, USA). A one-way analysis of variance (ANOVA) was applied to assess differences in temporal and spatial expression patterns, followed by appropriate post hoc tests when necessary. Comparisons between two groups were performed using Student's *t*-test. Survival data were analyzed via the Kaplan–Meier method, and survival curves were generated using GraphPad Prism 8.0.1 (GraphPad Software, La Jolla, CA, USA). Statistical significance was defined at $p < 0.05$, with high significance at $p < 0.01$, and extreme significance at $p < 0.001$.

3. Results

3.1. Sequence and Phylogenetic Analysis of TaAkt

The open reading frame of the *TaAkt* gene (GenBank accession number: PP920559) comprises 1539 bp, which encodes a protein of 512 amino acid residues, with a predicted molecular mass of 57.17 kDa and a theoretical isoelectric point of 6.27. Bioinformatic analysis revealed the presence of two potential N-glycosylation sites, located at amino acid positions 97–99 (NFS) and 319–321 (NIT), as well as 50 predicted phosphorylation

https://doi.org/10.3390/insects17020183

sites distributed throughout the protein sequence. Domain prediction analysis indicated that TaAkt possesses the characteristic structural features of the protein kinase B (Akt) family. These include an N-terminal pleckstrin homology (PH) domain spanning amino acids 17–120, which is essential for membrane localization and phosphoinositide binding, followed by a conserved serine/threonine protein kinase catalytic domain (S_TKC; amino acids 171–428) and a serine/threonine-type protein kinase extension domain (S_TK_X; amino acids 428–492) (Figure 1A). Phylogenetic analysis based on full-length Akt amino acid sequences from multiple insect species demonstrated that Akt proteins cluster into a single well-supported clade, indicating a high degree of evolutionary conservation across insects. Within this clade, TaAkt showed the closest phylogenetic relationship to Akt from *Bicyclus anynana*, another lepidopteran species (Figure 1B).




Figure 1. Domain architecture and phylogenetic analysis of *TaAkt* from *T. absoluta*. (**A**) Schematic representation of the conserved domains of the TaAkt protein, including the pleckstrin homology (PH) domain, the serine/threonine protein kinase catalytic domain (S_TKC), and the serine/threonine-type protein kinase extension domain (S_TK_X). (**B**) Phylogenetic tree of insect Akt proteins is constructed using the neighbor-joining method with 1000 bootstrap replicates. TaAkt is highlighted with a red line, and GenBank accession numbers for each species are listed in the tree.

3.2. Spatiotemporal Expression Analysis of TaAkt

The *TaAkt* gene was expressed at all examined developmental stages of *T. absoluta*, although transcript abundance varied significantly across development. Overall, *TaAkt* expression increased prior to molting and declined following each molting event. After adult emergence, *TaAkt* transcript levels progressively decreased with increasing adult age. Notably, *TaAkt* expression was highest in fourth-instar larvae, first-day female pupae, and first-day adults, with expression levels approximately 19-, 11-, and 8.4-fold higher,

https://doi.org/10.3390/insects17020183

respectively, than those observed in 3-day-old adults. In contrast, relatively low expression levels were detected in 6-day-old female pupae, 2-day-old adults, and 3-day-old adults. These differences in temporal expression were statistically significant (one-way ANOVA: $F_{(13,28)} = 50.48$, $p < 0.001$; Figure 2A).




Figure 2. Spatiotemporal expression patterns of *TaAkt* in *T. absoluta*. (**A**) Relative expression levels of *TaAkt* at different developmental stages. 1L-4L, first- to fourth-instar larvae; P1–P7, female pupae from day 1 to day 7; A1–A3, 1- to 3-day-old adults. (**B**) Relative expression levels of *TaAkt* in different tissues. HD, head; MG, midgut; OV, ovary; FB, fat body; AN, abdomen; WG, wing; EP, epidermis. Different uppercase letters above bars indicate significant differences among samples ($p < 0.05$, one-way ANOVA).

Spatial expression analysis revealed that *TaAkt* was ubiquitously expressed in all examined tissues but exhibited pronounced tissue-specific differences. The highest expression levels were observed in the wings and head, where *TaAkt* transcript abundance was approximately 63- and 28-fold higher, respectively, than that in the epidermis. The epidermis showed the lowest expression level among all tissues analyzed. Tissue-specific expression differences were highly significant (one-way ANOVA: $F_{(6,14)} = 194.65$, $p < 0.001$; Figure 2B).

3.3. TaAkt Knockdown on Pupal-Adult Transition in T. absoluta

Compared with the ds*GFP* control, *TaAkt* transcript levels decreased by 86%, 81%, 93%, and 88% at 12, 24, 48, and 72 h post-injection, respectively, confirming efficient gene silencing (12 h: $t = 9.51$, $p < 0.001$; 24 h: $t = 8.49$, $p < 0.001$; 48 h: $t = 57.12$, $p < 0.001$; 72 h: $t = 10.87$, $p < 0.001$; Figure 3A). Following *TaAkt* knockdown, pupal survival and successful adult emergence were markedly impaired. Within 7 days after injection, 98% of pupae in the ds*GFP* control group successfully completed the pupal-adult molt and emerged normally. In contrast, only 58% of pupae injected with ds*TaAkt* were able to molt and emerge successfully, indicating a significant reduction in survival during the pupal-adult transition (Figure 3B). Two distinct lethal phenotypes were observed among ds*TaAkt*-treated pupae (Figure 3C). Approximately 24% of pupae died during the pupal stage and exhibited severe cuticular shrinkage (P1 phenotype). An additional 18% of pupae initiated but failed to complete the pupal-adult molt and subsequently died (P2 phenotype). These results demonstrate that *TaAkt* is essential for successful pupal development and adult emergence in *T. absoluta*.

https://doi.org/10.3390/insects17020183




Figure 3. Effects of *TaAkt* knockdown on pupal development and survival of *T. absoluta*. (**A**) Relative expression levels of *TaAkt* at 12, 24, 48, and 72 h after injection with ds*TaAkt* or ds*GFP*. (**B**) Effects of *TaAkt* knockdown on survival rate during the pupal-adult transition. (**C**) Representative lethal phenotypes observed in female pupae following *TaAkt* silencing. Significant differences between ds*TaAkt* and ds*GFP* groups were determined using Student's *t*-test (** $p < 0.01$, *** $p < 0.001$). P1, pupal death characterized by cuticular shrinkage and darkening; P2, failure to complete the pupal-adult molt, resulting in death.

3.4. Effects of TaAkt Knockdown on 20E and Chitin Metabolism

At 48 h post-dsRNA injection, the 20E titer in ds*TaAkt*-treated insects was 15.83 ng/mL, representing a significant 58% reduction compared with 35.08 ng/mL in the ds*GFP* control group ($t = 15.82$, $p < 0.001$; Figure 4A). Consistent with these findings, the transcript levels of 20E synthesis genes, including *TaCYP306a1* ($t = 13.80$, $p = 0.005$), *TaCYP307a1* ($t = 49.42$, $p < 0.005$), *TaCYP314a1* ($t = 72.25$, $p < 0.005$), and *TaCYP315a1* ($t = 43.44$, $p < 0.005$), were significantly downregulated. Similarly, expression levels of 20E signaling pathway genes (*TaEcR*, *TaUSP*, *TaE75*, *TaHR3*, *TaHR38*, and *TaFTZ-F1*) were markedly decreased following *TaAkt* silencing (all $p < 0.005$; Figure 4B).

Knockdown of *TaAkt* also affected chitin metabolism. Chitin content in ds*TaAkt*-treated insects was 11.03 µmol/mL, representing a 37% reduction compared with 17.56 µmol/mL in the control group ($t = 3.26$, $p = 0.031$; Figure 4C). Analysis of chitin metabolism-related genes revealed significant downregulation of *TaCDA1* (−44%; $t = 5.26$, $p = 0.006$), *TaCHT5* (−82%; $t = 12.98$, $p < 0.001$), and *TaCHS* (−64%; $t = 20.96$, $p < 0.001$), whereas *TaUAP* expression showed a slight increase (Figure 4D).

https://doi.org/10.3390/insects17020183




Figure 4. Effects of *TaAkt* knockdown on 20E and chitin metabolism in *T. absoluta*. (**A**) 20E titer following ds*TaAkt* treatment. (**B**) Relative expression of 20E synthesis and signaling pathway genes at 48 h after *TaAkt* knockdown. (**C**) Chitin content after 48 h of *TaAkt* silencing. (**D**) Relative expression of chitin metabolism-related genes following 48 h of *TaAkt* knockdown. Significant differences between ds*TaAkt* and ds*GFP* control groups were determined using Student's *t*-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

3.5. Effects of TaAkt Silencing on Ovarian Development and Fertility

Ovaries from ds*GFP*-injected females developed normally, with well-formed ovarian tubes containing numerous mature eggs rich in vitellogenin, indicative of healthy reproductive status. In contrast, ovaries from ds*TaAkt*-injected females exhibited abnormal development: ovarian tubes were reduced in size, and most oocytes remained small and white, suggesting impaired vitellogenesis (Figure 5A). The average ovarian tube length in the ds*TaAkt* group was 2550 μm, representing a 48% reduction compared with 4930 μm in the control group. Similarly, egg length was significantly reduced to 271 μm after *TaAkt* knockdown, a 43% decrease relative to 477 μm in the ds*GFP* group (ovarian tube: $t = 12.55$, $p < 0.001$; egg length: $t = 11.72$, $p < 0.001$; Figure 5B).

Reproductive performance was also impaired. The average oviposition period of ds*TaAkt*-treated females was 6.70 days, significantly shorter than the 10.45 days observed in controls ($t = 14.59$, $p < 0.001$; Figure 5C). Moreover, ds*TaAkt*-injected females laid fewer eggs, with an average of 66.4 eggs per female, compared with 157.3 eggs per female in the ds*GFP* group, representing a 58% reduction. Egg hatching rate was similarly decreased, from 86% in controls to 46% in the ds*TaAkt* group, a 46% reduction (number of eggs: $t = 31.33$, $p < 0.001$; hatching rate: $t = 29.82$, $p < 0.001$; Figure 5D).

https://doi.org/10.3390/insects17020183




Figure 5. Effects of *TaAkt* knockdown on ovarian development and fertility in *T. absoluta*. (**A**) Representative ovarian phenotypes of 2-day-old female adults following ds*GFP* or ds*TaAkt* injection. (**B**) Quantitative measurements of ovarian tube length and egg length after *TaAkt* knockdown. (**C**) Average oviposition period of female adults following ds*TaAkt* treatment. (**D**) Effects of *TaAkt* silencing on fecundity (eggs laid per female) and egg hatching rate. Significant differences between ds*TaAkt* and ds*GFP* groups were determined using Student's *t*-test (* $p < 0.05$, ** $p < 0.01$).

3.6. Effects of TaAkt Silencing on JH Synthesis, JH Signaling, and Vitellogenin Content

Forty-eight hours after dsRNA injection, the JH titer in ds*TaAkt*-treated female pupae was 40.50 pg/mL, representing a 25% decrease compared with 53.85 pg/mL in the ds*GFP* control group ($t = 2.92$, $p = 0.043$; Figure 6A). Silencing of *TaAkt* significantly suppressed the expression of JH pathway genes. Transcript levels of *TaKr-h1*, *TaMet*, and *TaJHAMT* were reduced by 83%, 99%, and 72%, respectively, compared with controls (*TaKr-h1*: $t = 94.43$, $p < 0.001$; *TaMet*: $t = 33.25$, $p < 0.001$; *TaJHAMT*: $t = 33.61$, $p < 0.001$; Figure 6B).

Consistent with these effects on JH signaling, vitellogenin content in 1-day-old adult females was significantly decreased to 2.59 μg/mL, a 28% reduction relative to 3.60 μg/mL in the control group ($t = 4.15$, $p = 0.014$; Figure 6C). Furthermore, the relative transcript levels of *TaVg* and its receptor *TaVgR* were significantly downregulated by 27% and 34%, respectively, 48 h post-ds*TaAkt* injection (*TaVg*: $t = 7.46$, $p = 0.002$; *TaVgR*: $t = 9.98$, $p < 0.001$; Figure 6D).

https://doi.org/10.3390/insects17020183




Figure 6. Effects of *TaAkt* knockdown on JH signaling and vitellogenin (VTG) in *T. absoluta*. (**A**) JH titer 48 h after ds*TaAkt* injection. (**B**) Relative expression levels of JH signaling pathway genes (*TaKr-h1*, *TaMet*, and *TaJHAMT*) following *TaAkt* knockdown. (**C**) Vitellogenin content in 1-day-old adult females after ds*TaAkt* treatment. (**D**) Relative transcript levels of *TaVg* and *TaVgR* in females 48 h post-ds*TaAkt* injection. Significant differences between ds*TaAkt* and ds*GFP* control groups were determined using Student's *t*-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

4. Discussion

Insect insulin signaling and its central kinase, Akt, have been extensively studied in both model and non-model species [15,17]. Such investigations not only advance our understanding of fundamental developmental and metabolic processes in insects but also provide convenient experimental systems for exploring insulin-related mechanisms relevant to human physiology. *Akt* genes have been identified across a wide range of insect taxa, including Diptera [23–25], Orthoptera [26,27], Lepidoptera [13,28,29], Hemiptera [30–32], Coleoptera [33–35], Blattodea [36], and Hymenoptera [37,38]. In the present study, we cloned the ORF of *TaAkt* from the tomato leafminer, *T. absoluta*. The gene encodes a 512-amino-acid protein with high homology to Akt orthologs in other insects. *TaAkt* contains three conserved domains characteristic of the PKB/Akt family: the Pleckstrin homology (PH), Ser/Thr protein kinase catalytic (S_TKc), and Ser/Thr-type kinase (S_TK_X) domains, as well as a conserved hydrophobic motif (F-X-X-F-[S/T]-Y) spanning residues 485–490, identical to that reported in *H. armigera*, *Lasioderma serricorne*, and *B. mori* [13,29,33]. These conserved features are critical for Akt phosphorylation and interaction with upstream kinases such as PDK [11]. The high conservation across insect taxa underscores the functional

https://doi.org/10.3390/insects17020183

importance of Akt in key physiological processes and suggests that insights gained from *T. absoluta* may be extrapolated to other pest species.

JH and 20E are pivotal regulators of insect development and reproduction, coordinating processes such as molting, pupation, eclosion, and oviposition through temporally controlled titer fluctuations [39,40]. The insulin signaling pathway is positioned upstream of these hormonal cascades, integrating nutritional and developmental cues to modulate hormone synthesis and downstream signaling [41]. Akt, as a key transducer in this pathway, communicates insulin signals to diverse downstream targets, affecting gene expression and cellular metabolism [13]. Disruption of Akt function has been reported to impair both metamorphosis and reproductive output across multiple insect species [30,33,42].

In the present study, RNAi-mediated silencing of *TaAkt* during late pupal development partially blocked adult eclosion, a process primarily regulated by ecdysone. Knockdown of *TaAkt* resulted in a significant decrease in 20E titers and downregulation of ecdysone biosynthesis genes (*TaCYP306a1*, *TaCYP307a1*, *TaCYP314a1*, and *TaCYP315a1*), as well as downstream signaling components (*TaECR*, *TaUSP*, *TaE75*, *TaHR3*, *TaHR38*, and *TaFTZ-F1*). These findings are consistent with observations in *H. armigera*, where silencing of *HaPDK* reduced 20E titers and disrupted midgut remodeling [43], and in *B. mori*, where enhanced Akt phosphorylation stimulates ecdysteroid secretion, while PI3K inhibition reduces it [41]. Phenotypically, *TaAkt* knockdown induced pupal melanization, high mortality, and failure to complete the pupal-adult molt, similar to defects observed in *Hyphantria cunea*, *D. melanogaster*, and *Maruca vitrata* [44–46]. Silencing of *TaAkt* also affected chitin content and the expression of chitin metabolism-related genes (*TaCDA1*, *TaCHT5*, *TaUAP*, and *TaCHS*), which are essential for cuticle remodeling during eclosion [18]. These results collectively indicate that *TaAkt* orchestrates pupal-adult metamorphosis through coordinated regulation of ecdysone synthesis, signaling, and chitin metabolism.

Beyond development, insulin signaling plays a central role in insect reproductive physiology. In female *T. absoluta*, silencing *TaAkt* impeded ovarian growth, reducing the length of ovarioles and oocytes, decreasing egg production and hatching rates, and significantly lowering vitellogenin content and transcription of *Vg* and *VgR* genes. These results suggest that *TaAkt* regulates reproduction via both vitellogenin synthesis in the fat body and its uptake in the ovary. Similar outcomes have been reported in *Cyrtorhinus lividipennis*, *M. vitrata*, and *C. lividipennis* [30,47,48], whereas increased Akt activity in the fat body of *A. stephensi* enhances *Vg* expression and female fertility [23]. Moreover, knockdown of *TaAkt* decreased the expression of JH synthesis (*TaJHAMT*) and signaling genes (*TaMet* and *TaKr-h1*), corroborating previous findings in *L. serricorne*, *Liposcelis entomophila*, and *Frankliniella occidentalis*, where the depletion of *Akt* or *PDK* impaired JH production and disrupted reproductive processes [33,42,49]. Likewise, knockdown of other insulin pathway components, such as *ILP2*, *InR*, *TOR*, and *PDK*, reduces JH synthesis across various insect taxa [50–53]. Collectively, these findings reinforce the centrality of Akt in integrating nutritional signals, JH synthesis, and vitellogenin-mediated reproduction.

A key innovation of this study lies in elucidating the pleiotropic function of *TaAkt*, simultaneously coordinating molting and reproduction. By regulating 20E production, chitin metabolism, JH signaling, and vitellogenin synthesis, *TaAkt* disruption produces combined developmental and reproductive phenotypes, leading to high mortality and dramatically reduced fecundity. This dual role mirrors observations in *C. lividipennis*, *Colaphellus bowringi*, and *M. vitrata* [30,48,54], highlighting a conserved physiological strategy across insect species. The pleiotropic nature of Akt, embedded within the IIS pathway, allows it to phosphorylate multiple downstream targets involved in nutrient sensing, protein synthesis, and hormonal regulation [55]. Such multi-target potential enhances the attractiveness of Akt as an RNAi-based pest control target, where a single intervention

https://doi.org/10.3390/insects17020183

simultaneously impairs survival and reproductive success, potentially reducing pest populations more efficiently than strategies targeting single-function genes [56,57]. Furthermore, the requirement for compensatory mutations to restore fitness across multiple physiological axes increases the evolutionary cost of resistance, potentially slowing its development [58].

From an applied perspective, our study provides critical conceptual and practical insights for sustainable agriculture. *T. absoluta* is a globally invasive pest causing extensive damage to tomato crops. Targeting *TaAkt* via RNAi or other molecular interventions could enable selective suppression of both survival and reproduction, offering an innovative, environmentally friendly alternative to chemical pesticides. The ability to disrupt hormone signaling and vitellogenesis simultaneously could reduce crop damage while minimizing off-target effects on beneficial insects. These findings thus bridge fundamental insect physiology with applied pest management. Nevertheless, several gaps and limitations remain. First, the molecular mechanisms linking Akt to hormone biosynthesis and chitin metabolism require deeper mechanistic dissection, including identification of intermediate kinases, transcription factors, and tissue-specific regulatory networks. Second, the ecological efficacy of *TaAkt* RNAi under field conditions, including delivery strategies, stability, and potential off-target effects, has not yet been evaluated. Third, while our study focused on females, potential sex-specific differences in Akt function remain unexplored. Finally, interactions between insulin signaling and other pathways, such as TOR, FoxO, and ecdysone-independent growth regulators, merit further investigation to fully understand systemic regulatory networks. Consequently, we propose that future research directions should include: (1) elucidation of Akt-mediated transcriptional networks across various tissues; (2) development and optimization of RNAi delivery systems suitable for greenhouse and field applications; (3) assessment of the selectivity and non-target safety of Akt-based interventions; and (4) evaluation of potential synergistic effects with other hormonal or metabolic pathways for integrated pest management strategies. Addressing these areas will enhance both fundamental insect endocrinology and applied pest control technologies.

In summary, this study demonstrates that *TaAkt* serves as a pivotal regulator of both metamorphosis and reproduction in *T. absoluta*, integrating hormonal, metabolic, and developmental pathways. The dual impact of *TaAkt* RNAi on survival and reproductive fitness highlights its potential as a high-value molecular target for sustainable and selective pest management strategies. By linking mechanistic understanding to practical applications, these findings contribute to the advancement of innovative RNAi-based biopesticides and provide a foundation for translational research in agricultural pest control.

5. Conclusions

This study successfully cloned and characterized the ORF of the serine/threonine kinase *TaAkt* from *T. absoluta*. Functional analyses reveal that *TaAkt* regulates female emergence, ovarian development, and reproductive capacity by modulating 20E synthesis, chitin metabolism, JH production, and vitellogenin synthesis. RNAi-mediated knockdown of *TaAkt* disrupts pupal-adult metamorphosis, reduces JH titers, impairs vitellogenin accumulation, and leads to decreased fecundity. These findings provide novel mechanistic insights into insulin signaling in insect physiology and identify *Akt* as a promising target for RNAi-based pest management in sustainable agriculture. Future studies should focus on optimizing delivery strategies, assessing field efficacy, and evaluating ecological safety to translate these molecular insights into practical pest control solutions.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/insects17020183/s1: Table S1: Primers used in this study.

https://doi.org/10.3390/insects17020183

Author Contributions: Conceptualization, K.X. and W.Y.; methodology, C.L. and J.S.; validation, C.L.; investigation, C.L. and J.S.; data curation, C.L. and K.X.; writing—original draft preparation, K.X.; writing—review and editing, G.S. and W.Y.; supervision, F.W.; funding acquisition, W.Y. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Natural Science Foundation of Guizhou Province (QKHJC-ZK-2022-003), the Program of High-level Innovation Talents in Guizhou Province (GCC-2023-008), and the Program for Natural Science Research in Guizhou Education Department (QJJ-2023-024).

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding authors.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Biondi, A.; Guedes, R.N.C.; Wan, F.H.; Desneux, N. Ecology, worldwide spread, and management of the invasive south American tomato pinworm, *Tuta absoluta*: Past, present, and future. *Annu. Rev. Entomol.* **2018**, *63*, 239–258. [CrossRef] [PubMed]
2. Desneux, N.; Wajnberg, E.; Wyckhuys, K.A.G.; Burgio, G.; Arpaia, S.; Narváez-Vasquez, C.A. Biological invasion of European tomato crops by *Tuta absoluta*: Ecology, geographic expansion and prospects for biological control. *J. Pest Sci.* **2010**, *83*, 197–215. [CrossRef]
3. Júnior, P.A.S.; Kumar, L.; Silva, R.S.D.; Picanco, M.C. Global geographic distribution of *Tuta absoluta* as affected by climate change. *J. Pest Sci.* **2019**, *92*, 1373–1385.
4. Colmenarez, Y.C.; Vasquez, C.; Bueno, A.D.B.; Cantor, F.; Hidalgo, E.; Corniani, N. Sustainable nanagement of the invasive *Tuta absoluta* (Lepidoptera: Gelechiidae): An overview of case studies from Latin American countries participating in plantwise. *J. Integr. Pest Manag.* **2022**, *13*, pmac012. [CrossRef]
5. Phan, N.T.; Biddinger, D.J.; Rajotte, E.G.; Smagghe, G.; Reddy, G.V.P.; Ren, Z.X.; Joshi, N.K. Pesticide use in integrated pest and pollinator management framework to protect pollinator health. *Pest Manag. Sci.* **2025**, *81*, 1691–1696. [CrossRef]
6. Nässel, D.R.; Broeck, J.V. Insulin/IGF signaling in *Drosophila* and other insects: Factors that regulate production, release and post-release action of the insulin-like peptides. *Cell Mol. Life Sci.* **2016**, *73*, 271–290. [CrossRef]
7. Van de Velde, S.; Badisco, L.; Claeys, I.; Verleyen, P.; Chen, X.; Vanden Bosch, L.; Vanden Broeck, J.; Smagghe, G. Insulin-like peptides in *Spodoptera littoralis* (Lepidoptera): Detection, localization and identification. *Gen. Comp. Endocrinol.* **2007**, *153*, 72–79. [CrossRef]
8. Arik, A.J.; Hun, L.V.; Quicke, K.; Piatt, M.; Ziegler, R.; Scaraffia, P.Y.; Badgandi, H.; Riehle, M.A. Increased Akt signaling in the mosquito fat body increases adult survivorship. *FASEB J.* **2015**, *29*, 1404–1413. [CrossRef]
9. Lin, X.Y.; Yu, N.; Smagghe, G. Insulin receptor regulates food intake through sulfakinin signaling in the red flour beetle, *Tribolium castaneum*. *Peptides* **2016**, *80*, 89–95. [CrossRef]
10. Lin, X.Y.; Smagghe, G. Roles of the insulin signaling pathway in insect development and organ growth. *Peptides* **2019**, *122*, 169923. [CrossRef] [PubMed]
11. Badisco, L.; Wielendaele, P.V.; Broeck, J.V. Eat to reproduce: A key role for the insulin signaling pathway in adult insects. *Front. Physiol.* **2013**, *4*, 202. [CrossRef] [PubMed]
12. Gu, S.H.; Yeh, W.L.; Young, S.C.; Lin, P.L.; Li, S. TOR signaling is involved in PTTH-stimulated ecdysteroidogenesis by prothoracic glands in the silkworm, *Bombyx mori*. *Insect Biochem. Mol. Biol.* **2012**, *42*, 296–303. [CrossRef]
13. Mao, T.T.; Li, F.C.; Fang, Y.L.; Wang, H.; Chen, J.; Li, M.X.; Lu, Z.T.; Qu, J.W.; Li, J.X.; Hu, J.H.; et al. Effects of chlorantraniliprole exposure on detoxification enzyme activities and detoxification-related gene expression in the fat body of the silkworm, *Bombyx mori*. *Ecotoxicol. Environ. Saf.* **2019**, *176*, 58–63. [CrossRef]
14. Weinkove, D.; Neufeld, T.P.; Twardzik, T.; Waterfield, M.D.; Leevers, S.J. Regulation of imaginal disc cell size, cell number and organ size by *Drosophila* class I(A) phosphoinositide 3-kinase and its adaptor. *Curr. Biol.* **1999**, *9*, 1019–1029. [CrossRef]
15. Zhu, W.X.; Duan, Y.W.; Chen, J.Q.; Merzendorfer, H.; Zou, X.; Yang, Q. SERCA interacts with chitin synthase and participates in cuticular chitin biogenesis in *Drosophila*. *Insect Biochem. Mol. Biol.* **2022**, *145*, 103783. [CrossRef] [PubMed]
16. Parthasarathy, R.; Palli, S.R. Molecular analysis of nutritional and hormonal regulation of female reproduction in the red flour beetle, *Tribolium castaneum*. *Insect Biochem. Mol. Biol.* **2011**, *41*, 294–305. [CrossRef]
17. Hun, L.V.; Cheung, K.W.; Brooks, E.; Zudekoff, R.; Luckhart, S.; Riehle, M.A. Increased insulin signaling in the *Anopheles stephensi* fat body regulates metabolism and enhances the host response to both bacterial challenge and *Plasmodium falciparum* infection. *Insect Biochem. Mol. Biol.* **2021**, *139*, 103669. [CrossRef]
18. Zhou, Y.F.; Zhang, Y.; Xu, K.K.; Liu, R.Y.; Liu, W.B.; Ma, H.; Yang, W.J. *Chitin deacetylase 1* gene as an optimal RNAi-based target for controlling the tomato leaf miner *Tuta absoluta*. *Insects* **2024**, *15*, 838. [CrossRef]

https://doi.org/10.3390/insects17020183

19. Tamura, K.; Stecher, G.; Peterson, D.S.; Filipski, A.; Kumar, S. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* **2013**, *30*, 2725–2729. [CrossRef]
20. Yan, X.; Zhang, Y.B.; Xu, K.K.; Wang, Y.W.; Yang, W.J. Selection and validation of reference genes for gene expression analysis in *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae). *Insects* **2021**, *12*, 589. [CrossRef] [PubMed]
21. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **2001**, *25*, 402–408. [CrossRef] [PubMed]
22. Chen, Y.; Shi, Y.F.; Wang, Z.G.; An, X.; Wei, S.Y.; Andronis, C.; Vontas, J.; Wang, J.J.; Niu, J.Z. dsRNAEngineer: A web-based tool of comprehensive dsRNA design for pest control. *Trends Biotechnol.* **2025**, *43*, 969–983. [CrossRef] [PubMed]
23. Hun, L.V.; Luckhart, S.; Riehle, M.A. Increased Akt signaling in the fat body of *Anopheles stephensi* extends lifespan and increases lifetime fecundity through modulation of insulin-like peptides. *J. Insect Physiol.* **2019**, *118*, 103932. [CrossRef]
24. Du, X.; Wang, Y.; Wang, J.; Liu, X.; Chen, J.; Kang, J.; Yang, X.; Wang, H. d-Chiro-Inositol extends the lifespan of male *Drosophila melanogaster* better than d-Pinitol through insulin signaling and autophagy pathways. *Exp. Gerontol.* **2022**, *165*, 111856. [CrossRef]
25. Char, A.B.; Trammell, C.E.; Fawcett, S.; Chauhan, M.; Debebe, Y.; Céspedes, N.; Paslay, R.A.; Ahlers, L.R.H.; Patel, D.; Luckhart, S.; et al. Sustained antiviral insulin signaling during West Nile virus infection results in viral mutations. *Front. Cell Infect. Microbiol.* **2024**, *14*, 1492403. [CrossRef]
26. Li, S.; Wang, J.; Cui, D.N.; Hao, K.; Chen, J.; Tu, X.B.; Zhang, Z.H. Antagonism between PTP1B and PTK Mediates Adults' Insulin-Like Signaling Regulation of Egg Diapause in the Migratory Locust. *Insects* **2021**, *12*, 253. [CrossRef] [PubMed]
27. Chang, B.H.; Cui, B.Y.; Ullah, H.; Li, S.; Hao, K.; Tu, X.B.; Wang, G.J.; Nong, X.Q.; McNeill, M.R.; Huang, X.B.; et al. Role of PTP/PTK trans activated insulin-like signalling pathway in regulation of grasshopper (*Oedaleus asiaticus*) development. *Environ. Sci. Pollut. Res. Int.* **2019**, *26*, 8312–8324. [CrossRef]
28. Sun, R.R.; Jiang, L.W.; Chen, W.L.; Xu, Y.H.; Yi, X.; Zhong, G.H. Azadirachtin exposure inhibit ovary development of *Spodoptera litura* (Lepidoptera: Noctuidae) by altering lipids metabolism event and inhibiting insulin signaling pathways. *Ecotoxicol. Environ. Saf.* **2023**, *262*, 115151. [CrossRef] [PubMed]
29. Li, H.Y.; Wang, T.; Yang, Y.P.; Geng, S.L.; Xu, W.H. TGF-β signaling regulates p-Akt levels via PP2A during diapause entry in the cotton bollworm, *Helicoverpa armigera*. *Insect Biochem. Mol. Biol.* **2017**, *87*, 165–173. [CrossRef]
30. Hu, K.; Jin, R.; Liu, J.Q.; Zhu, J.; Dai, W.; Wang, Y.; Li, Y.; Liu, F. Functional characterization of the InR/PI3K/AKT signaling pathway in female reproduction of the predatory bug *Cyrtorhinus lividipennis* (Hemiptera: Miridae). *J. Econ. Entomol.* **2024**, *117*, 1655–1664. [CrossRef]
31. Yu, J.T.; Zhao, W.; Chen, X.F.; Lu, H.; Xiao, Y.; Li, Q.; Luo, L.; Kang, L.; Cui, F. A plant virus manipulates the long-winged morph of insect vectors. *Proc. Natl. Acad. Sci. USA* **2024**, *121*, e2315341121. [CrossRef]
32. Sun, C.X.; Li, Z.X. Biosynthesis of aphid alarm pheromone is modulated in response to starvation stress under regulation by the insulin, glycolysis and isoprenoid pathways. *J. Insect Physiol.* **2021**, *128*, 104174. [CrossRef] [PubMed]
33. Xu, K.K.; Yan, Y.; Yan, S.Y.; Xia, P.L.; Yang, W.J.; Li, C.; Yang, H. Disruption of the Serine/Threonine Kinase Akt Gene Affects Ovarian Development and Fecundity in the Cigarette Beetle, *Lasioderma serricorne*. *Front. Physiol.* **2021**, *12*, 765819. [CrossRef] [PubMed]
34. Zhou, M.; Pan, B.Y.; Guan, L.W.; Wang, Y.Y.; Xu, K.K.; Wang, S.G.; Tang, B.; Li, C. Comparative transcriptomic and metabolomics analysis of modified atmosphere responses in *Tribolium castaneum* (Coleoptera: Tenebrionidae). *J. Insect Sci.* **2024**, *24*, 8. [CrossRef]
35. Han, J.S.; Choi, S.Y.; Choi, R.Y.; Park, K.W.; Kang, K.Y.; Lee, M.K. Anti-muscle atrophy effect of fermented *Tenebrio molitor* larvae extract by modulating the PI3K-Akt-mTOR/FoxO3α pathway in mice treated with dexamethasone. *Biomed. Pharmacother.* **2024**, *178*, 117266. [CrossRef] [PubMed]
36. Luo, Y.; Hou, W.X.; Zhao, S.S.; Cheng, Y.L.; Zhang, K.J.; Lin, L.G.; Li, S.; Pei, X.J. An ovarian insulin-like peptide specifically regulates energy allocation and oocyte development in nutrition-restricted cockroaches. *BMC Biol.* **2025**, *23*, 226. [CrossRef]
37. Ament, S.A.; Corona, M.; Pollock, H.S.; Robinson, G.E. Insulin signaling is involved in the regulation of worker division of labor in honey bee colonies. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 4226–4231. [CrossRef]
38. Yan, H.; Opachaloemphan, C.; Carmona-Aldana, F.; Mancini, G.; Mlejnek, J.; Descostes, N.; Sieriebriennikov, B.; Leibholz, A.; Zhou, X.F.; Ding, L.; et al. Insulin signaling in the long-lived reproductive caste of ants. *Science* **2022**, *377*, 1092–1099. [CrossRef]
39. Dubrovsky, E.B. Hormonal cross talk in insect development. *Trends Endocrinol. Metab.* **2005**, *16*, 6–11. [CrossRef]
40. Wu, Z.X.; Yang, L.B.; He, Q.J.; Zhou, S.T. Regulatory Mechanisms of Vitellogenesis in Insects. *Front. Cell Dev. Biol.* **2021**, *8*, 593613. [CrossRef]
41. Gu, S.H.; Young, S.C.; Lin, J.L.; Lin, P.L. Involvement of PI3K/Akt signaling in PTTH-stimulated ecdysteroidogenesis by prothoracic glands of the silkworm, *Bombyx mori*. *Insect Biochem. Mol. Biol.* **2011**, *41*, 197–202. [CrossRef]
42. Lu, T.; Lu, Y.J.; Wang, L.; Liu, Z.P.; Miao, S.Y.; Tai, Y.J.; Yang, B.B. The serine/threonine kinase Akt gene affects fecundity by reducing Juvenile hormone synthesis in *Liposcelis entomophila* (Enderlein). *Pestic. Biochem. Physiol.* **2023**, *196*, 105583. [CrossRef]
43. Pan, J.; Di, Y.Q.; Li, Y.B.; Chen, C.H.; Wang, J.X.; Zhao, X.F. Insulin and 20-hydroxyecdysone oppose each other in the regulation of *phosphoinositide-dependent kinase-1* expression during insect pupation. *J. Biol. Chem.* **2018**, *293*, 18613–18623. [CrossRef] [PubMed]

https://doi.org/10.3390/insects17020183

44. Yan, L.Q.; Du, H.; Li, Y.; Li, X.; Sun, L.L.; Cao, C.W. Identification and characterization of key genes in insulin signaling pathway as molecular targets for controlling the fall webworm, *Hyphantria cunea*. *Pest Manag. Sci.* **2023**, *79*, 899–908. [CrossRef]
45. Shakhmantsir, I.; Massad, N.L.; Kennell, J.A. Regulation of cuticle pigmentation in drosophila by the nutrient sensing insulin and TOR signaling pathways. *Dev. Dyn.* **2014**, *243*, 393–401. [CrossRef] [PubMed]
46. Al-Baki, A.; Jung Kyo, J.; Kim, Y. Alteration of insulin signaling to control insect pest by using transformed bacteria expressing dsRNA. *Pest Manag. Sci.* **2020**, *76*, 1020–1030. [CrossRef]
47. Han, B.F.; Zhang, T.T.; Feng, Y.J.; Liu, X.P.; Zhang, L.S.; Chen, H.Y.; Zeng, F.R.; Wang, M.Q.; Liu, C.X.; Li, Y.Y.; et al. Two insulin receptors coordinate oogenesis and oviposition via two pathways in the green lacewing, *Chrysopa pallens*. *J. Insect Physiol.* **2020**, *123*, 104049. [CrossRef] [PubMed]
48. Al-Baki, M.A.; Lee, D.W.; Jung, J.K.; Kim, Y. Insulin signaling mediates previtellogenic development and enhances juvenile hormone-mediated vitellogenesis in a lepidopteran insect, *Maruca vitrata*. *BMC Dev. Biol.* **2019**, *19*, 14. [CrossRef]
49. Qiu, X.Y.; Huang, W.Q.; Yue, W.B.; Li, D.Y.; Zhi, J.R. Response of the serine/threonine kinase AKT and phosphoinositide-dependent kinase PDK in *Frankliniella occidentalis* (Thysanoptera: Thripidae) to three kinds of foods and their regulation of reproductive function. *Insect Mol. Biol.* **2024**, *33*, 372–386. [CrossRef]
50. Fu, K.Y.; Zhu, T.T.; Guo, W.C.; Ahmat, T.; Li, G.Q. Knockdown of a putative insulin-like peptide gene *LdILP2* in *Leptinotarsa decemlineata* by RNA interference impairs pupation and adult emergence. *Gene* **2016**, *581*, 170–177. [CrossRef]
51. Pan, X.; Pei, Y.; Zhang, C.; Huang, Y.; Chen, L.; Wei, L.; Li, C.; Dong, X.; Chen, X. Effect of insulin receptor on juvenile hormone signal and fecundity in *Spodoptera litura* (F.). *Insects* **2022**, *13*, 701. [CrossRef] [PubMed]
52. P'erez-Hedo, M.; Rivera-Perez, C.; Noriega, F.G. The insulin/TOR signal transduction pathway is involved in the nutritional regulation of juvenile hormone synthesis in *Aedes aegypti*. *Insect Biochem. Mol. Biol.* **2013**, *43*, 495–500. [CrossRef] [PubMed]
53. Yan, S.Y.; Song, J.H.; Yang, W.J.; Liu, W.X.; Zhang, G.F.; Wan, F.H.; Desneux, N.; Zhang, Y.B. Dual function of the *Tuta absoluta* 3-phosphoinositide-dependent protein kinase-1 in pupa ecdysis and adult reproduction. *J. Pest Sci.* **2025**, *98*, 1063–1073. [CrossRef]
54. Zhang, Y.K.; Zhang, H.X.; An, H.M.; Wang, K.; Zhu, F.; Liu, W.; Wang, X.P. Key roles of insulin receptor InR1 in initiating reproductive diapause in males of the cabbage beetle *Colaphellus bowringi* (Coleoptera: Chrysomelidae). *Pest Manag. Sci.* **2024**, *80*, 3852–3860. [CrossRef]
55. Cai, M.J.; Zhao, W.L.; Jing, Y.P.; Song, Q.; Zhang, X.Q.; Wang, J.X.; Zhao, X.F. 20-hydroxyecdysone activates Forkhead box O to promote proteolysis during *Helicoverpa armigera* molting. *Development* **2016**, *143*, 1005–1015. [CrossRef]
56. Xu, Q.Q.; Shang, F.; Feng, S.Y.; Xie, Q.P.; Zhang, W.; Wang, Z.G.; Wang, J.J. Design the fusion double-strand RNAs to control two global sap-sucking pests. *Pestic. Biochem. Physiol.* **2024**, *205*, 106114. [CrossRef]
57. Smagghe, G.; Palli, S.R.; Swevers, L. *RNAi in Agriculture: Basic Science to Applications-from Bioinformatics and Laboratory Assays over Regulatory Issues to Field Uses*; Springer: Dordrecht, The Netherlands, 2025; Volume 715.
58. Zhang, H.; Li, H.C.; Miao, X.X. Feasibility, limitation and possible solutions of RNAi-based technology for insect pest control. *Insect Sci.* **2013**, *20*, 15–30. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

https://doi.org/10.3390/insects17020183

insects

MDPI

Article

Identification and Functional Characterization of Chitinase Genes During Larva–Pupa–Adult Transitions in *Tuta absoluta*

Kangkang Xu [1,†], Yue Wang [1,†], Shuyan Yan [1,2], Fanghao Wan [2], Guy Smagghe [3,4] and Wenjia Yang [1,*]

[1] Guizhou Key Laboratory of Agricultural Biosecurity, Key Laboratory of Surveillance and Management of Invasive Alien Species in Guizhou Province, College of Biological and Environmental Engineering, Guiyang University, Guiyang 550005, China; kkxu1988@163.com (K.X.); yanshuyan27@126.com (S.Y.)
[2] Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China; wanfanghao@caas.cn
[3] Institute of Entomology, Guizhou University, Guiyang 550025, China; guysma9@gmail.com
[4] Department of Biology, Vrije Universiteit Brussel (VUB), 1050 Brussels, Belgium
* Correspondence: yangwenjia10@126.com
† These authors contributed equally to this work.

Simple Summary

The South American tomato leaf miner, *Tuta absoluta*, is a highly destructive invasive pest of solanaceous crops worldwide. Intensive reliance on chemical insecticides has led to widespread resistance and heightened environmental concerns, underscoring the need for species-specific and physiologically informed alternatives. Chitinases, which mediate chitin turnover during molting and metamorphosis, represent promising molecular targets for RNA interference (RNAi)-based control strategies. Therefore, this study aims to systematically characterize the chitinase family genes in *T. absoluta* and investigate their roles in larva–pupa–adult transitions through molecular characterization, spatiotemporal expression, and RNAi analyses. These findings reveal that four key chitinases (*TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF*) are indispensable for the molting physiology and metamorphic progression of *T. absoluta*, thereby providing potential molecular targets for leaf miner pest management.

Abstract

Chitinases are crucial in the molting process of insects and represent potential targets for the development of RNA interference (RNAi)-based insecticidal strategies. In this study, we identified and characterized 11 chitinase genes (*TaChts*) in *T. absoluta*, each harboring at least one GH18 catalytic domain. Phylogenetic analysis placed these proteins into 11 established groups (I-X and h), revealing conserved lineage-specific patterns. Spatiotemporal expression profiling showed that most *TaChts* peak during key developmental transitions. *TaCht1*, *TaCht2*, *TaCht3*, *TaCht5*, *TaCht6*, *TaCht7*, *TaCht10*, *TaCht-h*, and *TaIDGF* were predominantly expressed in the integument, whereas *TaCht8* and *TaCht11* showed gut-enriched expression, indicating functional specialization. Stage-tailored RNAi assays demonstrated that silencing *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF* caused pronounced mortality in both larvae and pupae. In larvae, delivery of dsRNA using carbon quantum dots (CQDs) disrupted molting: knockdown of *TaCht7*, *TaCht10*, and *TaIDGF* prevented shedding of the old cuticle, whereas *TaCht5* silencing produced severe cuticular shrinkage and blackening. Hematoxylin and eosin staining revealed detachment of epidermal cells from the cuticle, providing cytological evidence of disrupted cuticle–epidermis remodeling. In pupae, microinjection of dsRNA induced cuticular collapse, melanization, and eclosion failure due to persistent adhesion of the pupal case, accompanied by marked reductions in chitinase activity. Together, our findings highlight the critical roles of four key chitinases in endocuticular turnover and

metamorphic progression in *T. absoluta*, offering mechanistic insight into chitin-mediated developmental processes. The identified genes represent strong candidates for RNAi-based, species-specific pest management strategies against this globally significant insect.

Keywords: *Tuta absoluta*; chitinase; molting and metamorphosis; RNA interference; carbon quantum dots; cuticle remodeling; developmental physiology; pest management strategies

1. Introduction

The South American tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae), is an invasive pest originating from Peru that has rapidly become one of the most destructive species affecting solanaceous crops worldwide [1–3]. Its larvae mine leaves, stems, and fruits across all plant growth stages, generating serpentine tunnels that directly compromise photosynthesis and yield while predisposing damaged tissues to secondary infection by pathogens [4]. Presently, management relies heavily on chemical insecticides; however, chronic and indiscriminate use has produced widespread resistance, ecological contamination, and pest resurgence [5,6]. These constraints highlight the urgent need for physiologically informed, species-specific, and environmentally compatible alternatives to conventional chemical control.

RNA interference (RNAi) has emerged as a promising molecular tool for pest management due to its capacity to induce post-transcriptional gene silencing through exogenous double-stranded RNA (dsRNA), thereby disrupting essential physiological processes and causing aberrant development or mortality [7–10]. For insect physiologists, RNAi also provides an unparalleled experimental approach for resolving gene function and dissecting the molecular underpinnings of development, endocrinology, cuticle physiology, and metamorphic transitions. RNAi exhibits high specificity and efficiency in its mechanism of action; however, the effective delivery of dsRNA to target sites remains a significant challenge in practical applications. To address this issue, the utilization of nanomaterials as delivery carriers presents a promising strategy to overcome critical obstacles, such as degradation, cellular penetration, and targeted transport of dsRNA. The range of nanomaterials is diverse, including chitosan, liposomes, inorganic materials, synthetic polymers, and peptides, each with unique properties that enhance RNAi efficacy [11]. Among these, carbon nanoparticles have emerged as a promising carrier for dsRNA delivery, offering advantages such as low toxicity, high biocompatibility, and intrinsic fluorescence, which facilitates precise tracking of delivery [12]. Carbon quantum dots (CQDs) have been extensively used to form nanoparticles with dsRNA via electrostatic interactions, thereby gaining prominence in insect RNAi research. CQDs, along with other nanoparticle types, have proven effective in delivering dsRNA for the management of plant pathogens and pests [13]. Due to their robust endosomal escape capabilities, CQD nanoparticles have been identified as the most efficient carriers for inducing systemic RNAi through oral administration in *Chilo suppressalis* [14]. Furthermore, carbon dots derived from waste candle soot have been employed to deliver dsRNA targeting storage proteins and vitellogenin, resulting in a significant disruption of growth and reproduction in *Spodoptera litura* [15]. Collectively, these studies underscore the distinct advantages of CQDs in terms of biocompatibility, traceability, and delivery efficiency, highlighting their potential for future agricultural applications.

Chitin biology is central to these developmental processes. As a major structural polysaccharide of the insect exoskeleton and midgut peritrophic matrix, chitin is essential for maintaining cuticle integrity, preventing desiccation and pathogen invasion, and

https://doi.org/10.3390/insects17010114

shaping the gut environment for digestion and nutrient absorption [16]. Because insects do not store chitin, its cyclic degradation during molting and morphogenesis must be precisely coordinated through the regulated expression of chitinases (Chts). These enzymes hydrolyze β-1,4-glycosidic linkages in chitin and are indispensable for apolysis, ecdysis, pupation, and adult emergence [17–19]. The evolutionary conservation of chitin metabolic pathways, coupled with their absence in plants and vertebrates, makes chitinase genes not only tractable for molecular physiological studies but also highly selective targets for RNAi-based intervention [20]. Physiological investigations in multiple insects have revealed the essential and stage-specific roles of different Cht family members. In *Myzus persicae*, knockdown of *MpCht3*, *MpCht5*, *MpCht7*, *MpCht10*, and *MpCht11-2* disrupts nymphal molting and causes high mortality [21]. In *Musca domestica*, silencing *MdCht9* significantly reduces chitin content, induces over 50% lethality, and produces wing deformities, underscoring the role of chitin remodeling throughout metamorphosis [22]. In *Sogatella furcifera*, RNAi against several *Cht* genes decreases female fecundity [23]. Comparable effects, including molting defects, growth retardation, and death, are reported in *Leptinotarsa decemlineata* [24], *Phenacoccus solenopsis* [25], and *Bactrocera dorsalis* [26]. Beyond RNAi, chitinase genes have also been exploited to enhance viral biocontrol agents; incorporation of a *Spodoptera exigua Cht* gene into *Hyphantria cunea* nucleopolyhedrovirus markedly increases pathogenicity toward larval hosts [27]. Together, these findings demonstrate the centrality of chitinases in developmental physiology and highlight their potential in novel pest-management strategies.

Despite extensive work in other lepidopterans, the chitinase gene family of *T. absoluta*, including gene repertoire, expression dynamics, and functional roles during molting and metamorphosis, remains unexplored. Such knowledge is crucial not only for designing RNAi-based control tools but also for understanding the physiological mechanisms that underpin the species' rapid development, high survivorship, and ecological adaptability. Therefore, this study provides the first comprehensive identification and bioinformatic characterization of the chitinase gene family in *T. absoluta*. We examine temporal and tissue-specific expression patterns across key developmental transitions, with a focus on physiological processes related to cuticle turnover, metamorphic remodeling, and digestive system restructuring. To functionally dissect these genes, we employ stage-appropriate RNAi delivery strategies: (1) microinjection into pupae, which is an immobile, rigid stage well suited for precise dsRNA administration, to evaluate effects on metamorphic progression, pupal development, and adult eclosion; and (2) a carbon quantum dot (CQD)-mediated soaking method for feeding larvae, enabling efficient dsRNA uptake to assess consequences for larval survival, cuticle integrity, and molting success. The latter also offers a potential platform for oral RNAi delivery in future applied contexts. We believe that through these integrated molecular and physiological approaches, we elucidate the indispensable roles of multiple chitinase genes in *T. absoluta* development and metamorphosis. Our findings not only advance the basic physiological understanding of chitin metabolism in a globally important invasive species but also provide a mechanistic foundation and technical pathway for designing stage-targeted RNAi strategies. Such strategies may contribute to more sustainable, biologically grounded management options for *T. absoluta*, aligning with current demands for precision pest control that minimizes environmental impact.

2. Materials and Methods

2.1. Insect Rearing

A laboratory colony of *T. absoluta* was initially established from individuals collected in 2023 from Kunming City, Yunnan Province, China. The colony was continu-

https://doi.org/10.3390/insects17010114

ously maintained on tomato plants (*Solanum lycopersicum* cv. Provence) under controlled environmental conditions of 26 ± 1 °C, 60 ± 5% relative humidity, and a photoperiod of 16 h light followed by 8 h darkness, as reported before [28]. These conditions ensured stable development and synchronized life stages for downstream physiological and molecular experiments.

2.2. RNA Isolation and cDNA Synthesis

Total RNA was extracted from all developmental stages and tissues using the TaKaRa MiniBEST Universal RNA Extraction Kit (TaKaRa, Dalian, China), following the manufacturer's protocol. RNA concentration and purity were determined using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA), and RNA integrity was evaluated via electrophoresis on 1% agarose gels. First-strand cDNA synthesis was performed using 1 μg of total RNA and the One-Step gDNA Removal and cDNA Synthesis SuperMix (TransGen Biotech, Beijing, China). The resulting cDNA samples were stored at −80 °C for subsequent use.

2.3. Identification and Phylogenetic Analysis

To identify *chitinase* (*Cht*) family genes in *T. absoluta*, raw RNA-sequencing reads were initially obtained from the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) database (Accession number: SRR13065833). A de novo transcriptome assembly was conducted using the SOAPdenovo-Trans version 1.04 [29]. Subsequently, a local BLAST search (https://blast.ncbi.nlm.nih.gov/Blast.cgi, accessed on 5 November 2025) was performed against all assembled unigenes utilizing TBtools version 2.156. Reference chitinase sequences from other lepidopteran species, including *Bombyx mori* and *Plutella xylostella*, were retrieved from the NCBI database and used as query sequences. This search resulted in the identification of eleven putative *Cht* family genes in *T. absoluta* (Table S1). Candidate open reading frames (ORFs) were PCR-amplified using gene-specific primers (Table S2), cloned into the pTOPO-TA/Blunt vector (Aidlab Biotech, Beijing, China), and transformed into *Escherichia coli DH5α* competent cells (Yeasen Biotech, Shanghai, China). Positive clones were sequenced by CoWin Biotech (Taizhou, China). The ORF sequences were verified with ORF finder (http://www.ncbi.nlm.nih.gov/gorf/gorf.html, accessed on 5 December 2025). Domain structures were predicted by using SMART (https://smart.embl-heidelberg.de/, accessed on 5 December 2025) and the NCBI Conserved Domain Search (https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi, accessed on 5 December 2025).

To examine evolutionary relationships among chitinases, amino acid sequences from Lepidoptera, Coleoptera, Diptera, and Hemiptera were collected from GenBank (Table S3). Multiple protein sequence alignments were generated using ClustalW in MEGA 7, after which a phylogenetic tree was constructed using the neighbor-joining method with 1000 bootstrap replicates to evaluate node confidence [30]. This comparative analysis enabled classification of *T. absoluta Cht* genes into established subfamilies and provided insight into their evolutionary divergence.

2.4. Developmental and Tissue-Specific Expression Analysis

Expression profiling was conducted across major developmental stages, including first- through fourth-instar larvae, prepupae, pupae (1–7 days old), and adults (1–5 days old). Six larval tissues, including head, integument, fat body, Malpighian tubules, foregut, and midgut, were dissected from fourth-instar larvae under a stereo microscope (Olympus, Tokyo, Japan). Total RNA extraction and cDNA synthesis followed the procedures outlined in Section 2.2. Quantitative real-time PCR (qPCR) was performed on a CFX-96 real-time detection system (Bio-Rad, Hercules, CA, USA). Each reaction (20 μL) contained 10 μL of

https://doi.org/10.3390/insects17010114

TransStart® Top Green qPCR SuperMix (TransGen Biotech, Beijing, China), 1 μL of cDNA template, 1 μL of each of forward and reverse primers (Table S2), and 7 μL of nuclease-free water. Thermal cycling conditions included an initial denaturation at 95 °C for 30 s, followed by 41 cycles of 95 °C for 5 s and 55.9 °C for 30 s. Melting curve analysis (60–95 °C) confirmed primer specificity. Relative expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method [31], with *elongation factor 1-alpha* (*TaEF1α*; GenBank accession number: MZ054826) serving as the internal reference [32].

2.5. Functional Analysis of Chitinase Genes via RNAi

To investigate the functional roles of chitinase family members during *T. absoluta* development, dsRNA targeting specific *TaCht* genes was synthesized using the TranscriptAid T7 High Yield Transcription Kit (Thermo Fisher Scientific, Wilmington, DE, USA). The ds*GFP* served as the negative control. For pupal-stage RNAi assays, 2-day-old pupae were immobilized and microinjected with 500 ng of dsRNA into the dorsal vessel of the second dorsal segment using a Nanoliter 2010 microinjector (World Precision Instruments, Sarasota, FL, USA). Each treatment contained 30 pupae, with four biological replicates. Injected pupae were transferred to sterile soil moistened with sterile water and maintained under standard conditions until adult emergence or mortality assessment.

RNAi at the larval stage employed a CQD-mediated soaking method to enhance dsRNA uptake, and the CQDs were prepared as previously described [33]. Uniformly developed third-instar larvae were starved for 4 h before treatment. Larvae were then immersed in a dsRNA-CQD mixture prepared at a dsRNA: CQD ratio of 2:1, yielding a final dsRNA concentration of 1000 ng/μL in the soaking solution. After 10 min of incubation, the mixture was removed with a micropipette, and larvae were transferred onto fresh tomato leaves for rearing under the same conditions described in Section 2.1. To assess knockdown efficiency, 20 larvae were randomly selected at 72 h post-treatment for qPCR analysis. The remaining larvae were monitored every 24 h for 8 days to document survival rates and morphological abnormalities. Morphological observations were conducted using a Keyence digital microscope (Keyence, Osaka, Japan), allowing detailed assessment of cuticle defects, molting failure, and developmental arrest.

2.6. Chitinase Activity Assay

To evaluate how *TaCht* silencing affects endogenous chitinase activity in *T. absoluta* pupae, a quantitative enzymatic assay was conducted using the Chitinase Assay Kit (Solarbio, Beijing, China). Pupal samples were collected 24 h after dsRNA injection. Approximately 0.1 g of pupal tissue was homogenized in 1 mL of cold extraction buffer, followed by centrifugation at 10,000× *g* for 20 min at 4 °C. The supernatant was collected for chitinase activity measurement. A standard curve was prepared using serial dilutions of *N*-acetyl-*D*-glucosamine. Chitinase activity was calculated based on the concentration of *N*-acetylglucosamine produced in the reaction and expressed as units per gram (U/g) of tissue. One unit (U) of enzyme activity was defined as the amount of enzyme capable of generating 1 μg of *N*-acetylglucosamine per hour at 37 °C. The calculation used was: Chitinase activity (U/g) = 2.5 × C/W, where C is the *N*-acetylglucosamine concentration (μg/mL) determined from the standard curve and W is the tissue mass (g). Each treatment included three biological replicates, each with three technical repeats.

2.7. Frozen Sectioning and H&E Staining

To investigate structural alterations in the larval cuticle following chitinase gene knockdown, frozen sectioning followed by hematoxylin-eosin (H&E) staining was performed. Larvae were collected 24 h after dsRNA treatment and fixed overnight in 4% paraformaldehyde at room temperature. After dehydration, samples were embedded in

https://doi.org/10.3390/insects17010114

frozen embedding medium and placed in a cryostat microtome for 30 min before sectioning. Tissue sections (10 μm) were mounted onto adhesive slides. Sections were stained with hematoxylin for 60 s, rinsed, and differentiated in 0.1% hydrochloric acid-ethanol until nuclei appeared purple. Bluing was performed in PBS/warm water/1% ammonia for 50 s, followed by eosin staining for 60 s. The slides were dehydrated in 95% ethanol twice for 5 min each, cleared in xylene twice for 5 min each, and examined using a LSM 900 confocal laser-scanning microscope (Zeiss, Oberkochen, Germany).

2.8. Statistical Analysis

Survival curves were analyzed using the Kaplan–Meier method, and differences between ds*TaCht*-treated and control groups were tested using the Log-rank test. For pairwise comparisons between RNAi-treated and control samples, Student's *t*-test was used (* $p < 0.05$; ** $p < 0.01$). All statistical analyses were performed in SPSS v26.0, and figures were generated using GraphPad Prism v8.3.

3. Results

3.1. Molecular Cloning and Sequence Analysis of Chitinase Genes

From the *T. absoluta* transcriptome database, 11 full-length cDNA sequences belonging to the chitinase gene family were successfully cloned and identified: *TaCht1*, *TaCht2*, *TaCht3*, *TaCht5*, *TaCht6*, *TaCht7*, *TaCht8*, *TaCht10*, *TaCht11*, *TaCht-h*, and *TaIDGF*. The ORFs of these genes ranged from 1182 to 8667 bp. Domain prediction analyses revealed that all TaCht proteins contain conserved glycosyl hydrolase family 18 (GH18) catalytic domains (Figure 1A). Several chitinases contained multiple GH18 copies, with TaCht7 harboring two and TaCht10 possessing five such domains, whereas the remaining proteins each contained one. Signal peptides were predicted at the N-termini of all chitinases except TaCht6 and TaCht11, suggesting that most TaChts may be secreted enzymes. Chitin-binding domains (CBDs) were present in TaCht3, TaCht5, TaCht7, TaCht8, and TaCht10, with TaCht3 containing two CBDs and TaCht10 containing seven, indicating potentially strong associations with chitinous substrates. Phylogenetic analysis clustered the 11 TaChts into distinct, well-supported groups corresponding to classical insect chitinase subfamilies (Figure 1B). TaCht5 belonged to Group I, TaCht10 to Group II, TaCht7 to Group III, TaCht8 to Group IV, TaIDGF to Group V, TaCht6 to Group VI, TaCht2 to Group VII, TaCht11 to Group VIII, TaCht1 to Group IX, and TaCht3 to Group X. TaCht-h, a lineage-specific chitinase found only in Lepidoptera, was placed within the lepidopteran-specific h-group.

3.2. Developmental and Tissue-Specific Expression Patterns

Expression profiling across developmental stages showed that *TaCht1*, *TaCht2*, *TaCht5*, *TaCht8*, *TaCht10*, *TaCht11*, *TaCht-h*, and *TaIDGF* were highly expressed during the adult stage (Figure 2A). Among these, *TaCht1*, *TaCht2*, *TaCht5*, and *TaCht-h* exhibited peak expression on the first day post-eclosion, while *TaCht8* and *TaCht11* reached maximal expression on day three. *TaCht3*, *TaCht6*, and *TaCht7* displayed predominantly pupal-biased expression, with *TaCht3* showing a gradual increase from days 4 to 7, and *TaCht6* and *TaCht7* peaking on days 5 and 6, respectively.

https://doi.org/10.3390/insects17010114




Figure 1. Domain and phylogenetic analysis of chitinase proteins in *T. absoluta*. (**A**) Domain architectures of eleven chitinases in *T. absoluta*. Domain structures were predicted using SMART and the NCBI Conserved Domain Search based on deduced amino acid sequences. Signal peptides are shown in blue, catalytic domains in purple, and chitin-binding domains in green. (**B**) Phylogenetic analysis and group classification of chitinases (Chts) from *Anopheles gambiae* (Ag), *Bombyx mori* (Bm), *Culex quinquefasciatus* (Cq), *Danaus plexippus* (Dp), *Drosophila melanogaster* (Dm), *Nilaparvata lugens* (Nl), *Ostrinia furnacalis* (Of), *Phthorimaea operculella* (Po), *Plutella xylostella* (Px), *Spodoptera litura* (Sl), *Tribolium castaneum* (Tc), and *Tuta absoluta* (Ta). *T. absoluta* chitinases are indicated by red dots. Phylogenetic tree was constructed by MEGA 7 using the neighbor-joining method. Bootstrap analyses with 1000 replications were conducted, and only node support values of >70% are shown. The GenBank accession numbers for these chitinases are listed in Table S3.

Tissue-specific profiles indicated that most genes, including *TaCht1*, *TaCht2*, *TaCht3*, *TaCht5*, *TaCht6*, *TaCht7*, *TaCht10*, *TaCht-h*, and *TaIDGF*, were preferentially expressed in the integument, supporting their involvement in cuticle turnover during molting (Figure 2B). In contrast, *TaCht8* and *TaCht11* showed highest expression in the gut, suggesting functional roles in peritrophic matrix remodeling or digestive processes. *TaCht-h* and *TaIDGF* also exhibited elevated expression in the Malpighian tubules, while *TaCht3* displayed notably high expression in the head, potentially indicating additional roles in mouthpart remodeling or neural-associated structures.

https://doi.org/10.3390/insects17010114




Figure 2. Spatiotemporal expression profiles of chitinase genes in *T. absoluta*. (**A**) Temporal expression patterns of *TaCht* genes across developmental stages. Whole bodies from seventeen stages were used for qPCR analysis. L1–L4: first- to fourth-instar larvae; PP: prepupae; P1–P7: pupae 1 to 7 days old; AD1–AD5: adults 1 to 5 days old. (**B**) Tissue expression profiles of *TaChts* in fourth-instar larvae. cDNA was prepared from the head (HD), integument (IN), fat body (FB), Malpighian tubules (MT), foregut (FG), and midgut (MG). Table S4 presents the expression values of each gene across various developmental stages and tissues.

3.3. Silencing Efficiency and Mortality Induced by RNAi of Chitinase Genes

RNAi-mediated knockdown of individual *TaCht* genes was performed to investigate their essential biological roles. During the larval stage, third-instar larvae were exposed to CQD-assisted dsRNA soaking (Figure 3A). The qPCR analyses at 72 h post-treatment verified that transcript levels of all 11 *TaCht* genes were significantly reduced (Figure 3B). Survival analysis revealed that silencing *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF* resulted in substantial larval mortality, namely 67%, 60%, 73%, and 67%, respectively, while knockdown of the remaining seven chitinase genes did not significantly affect larval survival (Figure 3D).

https://doi.org/10.3390/insects17010114




Figure 3. Effect of *TaCht* gene knockdown on survival in *T. absoluta*. (**A**) Schematic of the RNAi strategy: third-instar larvae ingest leaves pre-soaked with a nanomaterial/dsRNA complex for *TaCht* silencing, whereas 2-day-old pupae receive direct microinjection of dsRNA. (**B**) Relative expression of *TaChts* in third-instar larvae 3 days after ds*TaChts* or ds*GFP* soaking. (**C**) Relative expression of *TaChts* in 2-day-old pupae 3 days after ds*TaChts* or ds*GFP* microinjection. (**D**) Effect of *TaCht* silencing on 8-day survival of third-instar larvae after dsRNA soaking. (**E**) Effect of *TaCht* silencing on 8-day survival of 2-day-old pupae after dsRNA microinjection. Significance was determined using Student's *t*-test (** $p < 0.01$, * $p < 0.05$, ns: no significance). Mean survival time was calculated using the Kaplan–Meier method.

In the pupal stage, microinjection of dsRNA into 2-day-old pupae successfully induced strong transcript suppression of all target genes at 72 h (Figure 3C). Consistent with larval results, silencing of *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF* significantly increased pupal mortality, whereas silencing the other *TaCht* genes produced no significant effects

https://doi.org/10.3390/insects17010114

(Figure 3E). These findings demonstrate that these four genes play indispensable roles during both larval and pupal development, while other chitinases may have redundant or stage-limited functions.

3.4. Effects of Silencing TaCht5, TaCht7, TaCht10, and TaIDGF on Larval–Pupal Ecdysis

To clarify how these four genes contribute to larval–pupal transition, third-instar larvae were treated with dsRNA-CQD complexes. In the ds*GFP* control group, larvae successfully completed larval–pupal ecdysis (Figure 4A). In contrast, ds*TaCht5*-treated larvae displayed severe shrinkage, cuticular melanization, and ultimately death without successful molting. Silencing *TaCht7*, *TaCht10*, or *TaIDGF* produced similar phenotypes, including pronotal cracking, failed separation of the abdominal integument, and incomplete apolysis that prevented pupation (Figure 4A).




Figure 4. Effects of silencing *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF* on larval–pupal development in *T. absoluta*. (**A**) Defective larval phenotypes following knockdown of *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF*. (**B**) Hematoxylin and eosin staining of abdominal cuticle sections after injection with dsRNA targeting *TaCht5*, *TaCht7*, *TaCht10*, *TaIDGF* or ds*GFP*. ec: epidermal cells; lc: larval cuticle; pc: pupal cuticle.

The observed mortality rates, namely 67% for ds*TaCht5*, 60% for ds*TaCht7*, 73% for ds*TaCht10*, and 67% for ds*TaIDGF*, corroborated the phenotypic severity of molting defects. To further examine cuticular changes, H&E staining revealed that control larvae exhibited clear separation between the new and old cuticle layers prior to ecdysis. In contrast, dsRNA-treated larvae showed detachment of epidermal cells from the cuticle, irregular chitin organization, and failed cuticle splitting (Figure 4B), explaining the lethal molting failure induced by gene silencing.

https://doi.org/10.3390/insects17010114

3.5. Effects of Silencing TaCht5, TaCht7, TaCht10, and TaIDGF on Pupal–Adult Transition

To evaluate the functions of these genes in pupal metamorphosis, dsRNAs were microinjected into 2-day-old pupae and developmental outcomes were observed. After ds*TaCht5* injection, 67% of pupae underwent cuticular melanization and died before eclosion, and an additional 13% initiated eclosion but could not shed the pupal case, resulting in death. Silencing *TaCht7* caused 59% melanized, shrunken pupae and 23% adults that reached eclosion but failed to detach from their pupal cases. In ds*TaCht10*-treated pupae, 55% exhibited shrinkage and melanization, and 15% died during incomplete eclosion. Silencing *TaIDGF* caused 61% early pupal mortality due to melanization and 25% mortality during incomplete adult emergence. The total mortality rates were 81% (*TaCht5*), 84% (*TaCht7*), 76% (*TaCht10*), and 84% (*TaIDGF*), while all ds*GFP*-injected pupae successfully emerged as normal adults (Figure 5A). Chitinase activity assays showed that 72 h after dsRNA injection, enzymatic activity decreased by 82% in ds*TaCht5*-, 80% in ds*TaCht7*-, 85% in ds*TaCht10*-, and 80% in ds*TaIDGF*-treated pupae compared to controls (Figure 5B).




Figure 5. Effects of silencing *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF* on pupal–adult development in *T. absoluta*. (**A**) Defective pupal phenotypes following knockdown of *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF*. (**B**) Changes in chitinase activity after injection of dsRNA targeting *TaCht5*, *TaCht7*, *TaCht10*, *TaIDGF* or ds*GFP*. Significance was determined using Student's *t*-test (** $p < 0.01$).

https://doi.org/10.3390/insects17010114

4. Discussion

Chitinases are essential enzymes involved in chitin degradation, playing indispensable roles in insect growth, molting, and metamorphosis. Consequently, the identification and characterization of insect chitinases are crucial for uncovering developmental mechanisms while simultaneously guiding the design of innovative, physiology-based pest control tools. In this study, eleven chitinase genes were identified from the transcriptome database of *T. absoluta*, representing a moderate gene family size relative to other insects. Comparable chitinase gene numbers have been reported in *Acyrthosiphon pisum* and *B. mori*, both with nine genes [34,35], *Spodoptera frugiperda* with eleven [36], and *P. xylostella* with thirteen [37]. In contrast, species such as *L. decemlineata*, *S. furcifera*, and *Aedes albopictus* contain 12, 19, and 20 chitinase genes, respectively [24,38,39]. These interspecific differences highlight lineage-specific expansions potentially associated with distinct ecological niches, cuticle chemistries, or metamorphic strategies.

Phylogenetic analysis confirmed that the identified *T. absoluta* chitinases follow the conventional classification for Lepidoptera [37], placing all TaChts into groups I-X and h. The conservation of GH18 catalytic domains across all identified genes, accompanied by diverse domain architectures [40], suggests functional partitioning of chitinolytic processes across tissues and developmental stages. This structural diversification aligns with the increasing appreciation within insect physiology that chitinase paralogs evolve distinct biochemical properties to meet the variable mechanical and physiological demands of molting, gut homeostasis, immune defense, and cuticle remodeling.

The expression patterns of chitinase genes across developmental stages and tissues in insects typically reveal strong specialization. Previous investigations, for example, in *S. frugiperda* and *Diaphorina citri*, have demonstrated integument-specific expression of several *Cht* paralogs (*SfCht2*, *SfCht3*, *SfCht10*), midgut-enriched expression of *Cht8* subfamily members, and stage-specific peaks aligned with larval–larval or larval–pupal molts [36,41]. The *T. absoluta* expression profiles observed here reinforce these patterns, offering new insights into cuticle physiology within this highly invasive pest. Elevated expression of *TaCht1*, *TaCht2*, *TaCht5*, *TaCht7*, *TaCht10*, *TaCht-h*, and *TaIDGF* during molting, coupled with their enrichment in the integument, strongly suggests involvement in cuticular turnover. Conversely, gut-specific expression of *TaCht8* and *TaCht11* indicates potential roles in peritrophic matrix remodeling or digestive physiology, opening avenues for further functional dissection of midgut-specific chitin synthases and hydrolases.

A major innovation of this study lies in the development and application of stage-tailored RNAi delivery systems, addressing a long-standing limitation in insect functional genomics. Physiological barriers such as gut nucleases, cuticle hardening, and feeding cessation have historically constrained the efficacy of RNAi in holometabolous insects [7]. By deploying nanocarrier CQDs during the larval stage, which is when feeding and gut permeability favor dsRNA uptake, and microinjection during the nonfeeding pupal stage, which is when direct hemocoelic access is required, this work demonstrates a robust methodology capable of overcoming these barriers [8]. The dual approach not only enhances gene-silencing efficiency but also provides a blueprint for integrating nanoparticle-assisted delivery with conventional techniques across metamorphosis, marking a significant achievement in insect physiological experimentation. Notably, CQDs can be synthesized from cost-effective carbon precursors and readily conjugated with dsRNA, suggesting the technical feasibility of large-scale production [42,43]. Within this context, CQD-dsRNA complexes hold potential for adaptation to foliar applications, such as direct spraying onto plant surfaces, thereby exposing feeding larvae to dsRNA in a manner similar to other nano-enabled RNAi strategies documented for crop protection [44]. However, these applications are currently speculative and necessitate comprehensive validation.

https://doi.org/10.3390/insects17010114

The functional assays focusing on the larval–pupal and pupal–adult transitions revealed striking phenotypic effects associated with four genes: *TaCht5*, *TaCht10*, *TaCht7*, and *TaIDGF*. The absence of phenotypes following the silencing of *TaCht1*, *TaCht2*, *TaCht3*, *TaCht6*, *TaCht8*, and *TaCht11* suggests functional redundancy or roles limited to physiological processes not critical for molting survival. These findings parallel studies in *S. furcifera* where several *Cht* paralogs showed no lethal phenotypes upon RNAi despite detectable expression [38], emphasizing the need for broader integrative analyses of gene networks, compensatory metabolic pathways, and hormonal regulation. In contrast, the lethal molting defects observed after silencing *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF* underscore the essential functions of these chitinases in cuticular degradation. Massive evidence across insect orders supports the fundamental involvement of *Cht5*, *Cht7*, and *Cht10* families in ecdysis-driven remodeling [45]. Orthologous phenotypes include failed nymph–adult transition in *S. furcifera* [38], abnormal pupal eclosion in *P. xylostella* [37], wing-curling defects in *C. suppressalis* [46], compromised cuticle integrity and flight morphology in *D. melanogaster* [47], and larval mortality in *Agrotis ipsilon* due to suppressed chitin degradation [48]. The parallels in *T. absoluta*, which included melanization, integument shrinkage, arrested ecdysis, and fatal entrapment within exuviae, confirm that chitinase-mediated cuticle turnover is physiologically conserved but anatomically tuned to species-specific cuticle architectures. Microscopic H&E analysis provided direct evidence linking gene silencing to cellular disruptions, revealing failures in epidermal-cuticular detachment, a key prerequisite for successful apolysis. Coupled with significant reductions in enzymatic activity following dsRNA treatment, these findings create a mechanistic bridge connecting gene expression, enzyme function, and whole-organism outcomes. Such multi-level validation represents a major strength of this study, emphasizing integrative approaches bridging molecular, cellular, and organismal physiology.

Next to the fundamental insights, we believe that the applied implications of these findings are substantial. Chitinases central to metamorphosis, particularly *TaCht5*, *TaCht7*, *TaCht10*, and *TaIDGF*, represent high-value RNAi targets for next-generation biopesticides. The demonstrated efficiency of CQD-assisted RNAi uptake in larvae offers a feasible route for field-adapted formulations, and the species specificity of chitinase sequences minimizes risks to beneficial arthropods. Moreover, because chitin metabolism is highly conserved yet absent in plants and vertebrates, targeting these genes avoids hazards associated with off-target effects typical of chemical insecticides. However, several limitations and knowledge gaps remain. First, while this study documents lethal phenotypes, it does not quantify potential compensatory upregulation of other chitinases or related hydrolases following RNAi, a process increasingly recognized in RNAi-mediated gene networks. Second, upstream endocrine regulation, particularly 20E and juvenile hormone signaling, was not explored, limiting interpretation of how *TaCht* expression integrates into larger metamorphic circuits. Third, the efficiency of CQD-dsRNA complexes under natural environmental conditions (UV degradation, microbial activity, plant leaf surface chemistry) remains untested. Finally, although microinjection offers strong mechanistic insight, it is impractical for field application, necessitating the development of environmentally stable larval-delivery formulations. Based on this, we suggest that future directions should include: (1) integrative hormonal and transcriptomic profiling during silencing to map chitinase regulation within the broader endocrine landscape; (2) CRISPR-based knockout studies to validate essential functions without RNAi variability; (3) structural enzymology to identify catalytic residues and potential inhibitors; (4) evaluation of nanoparticle-based oral RNAi in greenhouse and field conditions; and (5) exploration of synergistic pest control strategies, combining RNAi with fungal entomopathogens, botanical compounds, or NPV-based approaches [22]. Collectively, our study provides new mechanistic insights into the

https://doi.org/10.3390/insects17010114

physiology of insect molting and metamorphosis, advances methodological innovation for functional genomics in holometabolous insects, and establishes chitinase pathways as promising and practical targets for next-generation pest management strategies.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/insects17010114/s1, Table S1: Putative nucleotide sequences of eleven chitinase family genes in *Tuta absoluta*. Table S2: Primers used in this study. Table S3: Scientific names of chitinase sequences from different insect species used in constructing the phylogenetic tree. Table S4: Developmental and tissue-specific expression profiles of eleven *TaCht* genes in *Tuta absoluta*.

Author Contributions: Conceptualization, K.X.; methodology, Y.W.; investigation, K.X. and Y.W.; data curation, K.X., S.Y. and Y.W.; writing—original draft preparation, K.X. and S.Y.; writing—review and editing, G.S. and W.Y.; supervision, F.W.; funding acquisition, W.Y. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Natural Science Foundation of Guizhou Province (QKHJC-ZK-2022-003), the Program of Excellent Innovation Talents in Guizhou Province (GCC-2023-071), and the Program for Natural Science Research in Guizhou Education Department (QJJ-2023-024).

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Materials. Further inquiries can be directed to the corresponding authors.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Desneux, N.; Luna, M.G.; Guillemaud, T.; Urbaneja, A. The invasive South American tomato pinworm, *Tuta absoluta*, continues to spread in Afro-Eurasia and beyond: The new threat to tomato world production. *J. Pest Sci.* **2011**, *84*, 403–408. [CrossRef]
2. Biondi, A.; Guedes, R.N.C.; Wan, F.H.; Desneux, N. Ecology, worldwide spread, and management of the invasive south American tomato pinworm, *Tuta absoluta*: Past, present, and future. *Annu. Rev. Entomol.* **2018**, *63*, 239–258. [CrossRef] [PubMed]
3. Zhang, G.F.; Ma, D.Y.; Wang, Y.S.; Gao, Y.H.; Liu, W.X.; Zhang, R.; Fu, W.J.; Xian, X.Q.; Wang, J.; Kuang, M.; et al. First report of the South American tomato leafminer, *Tuta absoluta* (Meyrick), in China. *J. Integr. Agric.* **2020**, *19*, 1912–1917. [CrossRef]
4. Desneux, N.; Han, P.; Mansour, R.; Arno, J.; Brevault, T.; Campos, M.R.; Chailleux, A.; Guedes, R.N.C.; Karimi, J.; Konan, K.A.J.; et al. Integrated pest management of *Tuta absoluta*: Practical implementations across different world regions. *J. Pest Sci.* **2022**, *95*, 17–39. [CrossRef]
5. Guedes, R.N.C.; Campos, M.R.; Haddi, K.; Bielza, P.; Siqueira, H.A.A.; Tsagkarakou, A.; Vontas, J.; Nauen, R. Insecticide resistance in the tomato pinworm *Tuta absoluta*: Patterns, spread, mechanisms, management and outlook. *J. Pest Sci.* **2019**, *92*, 1329–1342. [CrossRef]
6. Wang, M.H.; Ismoilov, K.; Liu, W.X.; Bai, M.; Bai, X.S.; Chen, B.; Chen, H.L.; Chen, H.S.; Dong, Y.C.; Fang, K.; et al. *Tuta absoluta* management in China: Progress and prospects. *Entomol. Gen.* **2024**, *44*, 269–278. [CrossRef]
7. Adeyinka, O.; Riaz, S.; Toufiq, N.; Yousaf, I.; Bhatti, M.; Batcho, A.; Olajide, A.; Nasir, I.; Tabassum, B. Advances in exogenous RNA delivery techniques for RNAi-mediated pest control. *Mol. Biol. Rep.* **2020**, *47*, 6309–6319. [CrossRef]
8. Yan, S.; Ren, B.Y.; Shen, J. Nanoparticle-mediated double-stranded RNA delivery system: A promising approach for sustainable pest management. *Insect Sci.* **2021**, *28*, 21–34. [CrossRef]
9. Oberemok, V.V.; Laikova, K.; Andreeva, O.A.; Gal'chinsky, N.V. Oligonucleotide insecticides and RNA-based insecticides: 16 years of experience in contact using of the next generation pest control agents. *J. Plant Dis. Protect.* **2024**, *131*, 1837–1852. [CrossRef]
10. Smagghe, G.; Palli, S.R.; Swevers, L. *RNAi in Agriculture: Basic Science to Applications-From Bioinformatics and Laboratory Assays over Regulatory Issues to Field Uses*; Springer: Dordrecht, The Netherlands, 2025; 715p, ISBN 978-3-031-81548-5.
11. Liu, J.S.; He, Q.Y.; Lin, X.F.; Smagghe, G. Recent progress in nanoparticle-mediated RNA interference in insects: Unveiling new frontiers in pest control. *J. Insect Physiol.* **2025**, *167*, 104884. [CrossRef]
12. Arjunan, N.; Thiruvengadam, V.; Sushil, S.N. Nanoparticle-mediated dsRNA delivery for precision insect pest control: A comprehensive review. *Mol. Biol. Rep.* **2024**, *51*, 355. [CrossRef]
13. Xing, Y.; Jiang, H.; Cai, L. Engineered nanotransporters for efficient RNAi delivery in plant protection applications. *J. Integr. Plant Biol.* **2025**, *67*, 1223–1245. [CrossRef]
14. Wang, K.X.; Peng, Y.C.; Chen, J.S.; Peng, Y.; Wang, X.S.; Shen, Z.H.; Han, Z.J. Comparison of efficacy of RNAi mediated by various nanoparticles in the rice striped stem borer (*Chilo suppressalis*). *Pestic. Biochem. Physiol.* **2020**, *165*, 104467. [CrossRef]

https://doi.org/10.3390/insects17010114

15. Pandey, R.; Chauhan, R.; Saurabh, S.; Shukla, A.K.; Jamal, F.; Singh, S.P.; Singh, P.K.; Mishra, M. The impact of carbon NPs on the accumulation of storage proteins and the generation advancement of the polyphagous insect pest tobacco cutworm *Spodoptera litura* (Fabricius). *Environ. Sci. Nano* **2024**, *11*, 2428–2446. [CrossRef]
16. Zhu, K.Y.; Merzendorfer, H.; Zhang, W.; Zhang, J.; Muthukrishnan, S. Biosynthesis, turnover, and functions of chitin in insects. *Annu. Rev. Entomol.* **2016**, *61*, 177–196. [CrossRef]
17. Liu, X.J.; Cooper, A.M.W.; Zhang, J.Z.; Zhu, K.Y. Biosynthesis, modifications and degradation of chitin in the formation and turnover of peritrophic matrix in insects. *J. Insect Physiol.* **2019**, *114*, 109–115. [CrossRef]
18. Merzendorfer, H. Insect chitin synthases: A review. *J. Comp. Physiol. B.* **2006**, *176*, 1–15. [CrossRef] [PubMed]
19. Tharanathan, R.N.; Kittur, F.S. Chitin-the undisputed biomolecule of great potential. *Crit. Rev. Food Sci. Nutr.* **2003**, *43*, 61–87. [CrossRef]
20. Kramer, K.J.; Muthukrishnan, S. Insect chitinases: Molecular biology and potential use as biopesticides. *Insect Biochem. Mol. Biol.* **1997**, *27*, 887–900. [CrossRef] [PubMed]
21. Zhang, C.; He, L.; Ding, B.; Yang, H. Identification and functional characterization of the chitinase and chitinase-like gene family in *Myzus persicae* (Sulzer) during molting. *Pest Manag. Sci.* **2025**, *81*, 327–339. [CrossRef] [PubMed]
22. Zhao, W.J.; Li, Y.; Jiao, Z.L.; Su, P.P.; Yang, L.B.; Sun, C.Q.; Xiu, J.F.; Shang, X.L.; Guo, G. Function analysis and characterisation of a novel chitinase, *MdCht9*, in *Musca domestica*. *Insect Mol. Biol.* **2024**, *33*, 157–172. [CrossRef]
23. Zeng, Q.H.; Gong, M.; Yang, H.; Chen, N.N.; Lei, Q.; Jin, D.C. Effect of four chitinase genes on the female fecundity in *Sogatella furcifera* (Horváth). *Pest Manag. Sci.* **2024**, *80*, 1912–1923. [CrossRef] [PubMed]
24. Chen, Y.; Tang, H.; Zhou, W.; Li, C.; Chen, Y.N.; Zhang, Q.; Fu, K.Y.; Guo, W.; Shi, J.F. Identification of chitinase genes and roles in the larval-pupal transition of *Leptinotarsa decemlineata*. *Pest Manag. Sci.* **2024**, *80*, 282–295. [CrossRef]
25. Omar, M.A.A.; Ao, Y.; Li, M.; He, K.; Xu, L.; Tong, H.; Jiang, M.; Li, F. The functional difference of eight chitinase genes between male and female of the cotton mealybug, *Phenacoccus solenopsis*. *Insect Mol. Biol.* **2019**, *28*, 550–567. [CrossRef]
26. Liu, S.H.; Li, H.F.; Yang, Y.; Yang, R.L.; Yang, W.J.; Jiang, H.B.; Dou, W.; Smagghe, G.; Wang, J.J. Genome-wide identification of chitinase and chitin deacetylase gene families in the oriental fruit fly, *Bactrocera dorsalis* (Hendel). *Comp. Biochem. Physiol. D Genom. Proteom.* **2018**, *27*, 13–22. [CrossRef]
27. Shinoda, T.; Kobayashi, J.; Matsui, M.; Chinzei, Y. Cloning and functional expression of a chitinase cDNA from the common cutworm, *Spodoptera litura*, using a recombinant baculovirus lacking the virus-encoded chitinase gene. *Insect Biochem. Mol. Biol.* **2001**, *31*, 521–532. [CrossRef]
28. Zhou, Y.F.; Zhang, Y.; Xu, K.K.; Liu, R.Y.; Liu, W.B.; Ma, H.; Yang, W.J. *Chitin deacetylase 1* gene as an optimal RNAi-based target for controlling the tomato leaf miner *Tuta absoluta*. *Insects* **2024**, *15*, 838. [CrossRef] [PubMed]
29. Xie, Y.L.; Wu, G.X.; Tang, J.B.; Luo, R.B.; Patterson, J.; Liu, S.L.; Huang, W.H.; He, G.Z.; Gu, S.C.; Li, S.K.; et al. SOAPdenovo-Trans: De novo transcriptome assembly with short RNA-Seq reads. *Bioinformatics* **2014**, *30*, 1660–1666. [CrossRef] [PubMed]
30. Kumar, S.; Stecher, G.; Tamura, K. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* **2016**, *33*, 1870–1874. [CrossRef]
31. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **2001**, *25*, 402–408. [CrossRef]
32. Yan, X.; Zhang, Y.B.; Xu, K.K.; Wang, Y.W.; Yang, W.J. Selection and validation of reference genes for gene expression analysis in *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae). *Insects* **2021**, *12*, 589. [CrossRef]
33. Yang, N.; Tang, J.X.; Lai, J.; Chen, L.H.; Xu, Y.Q.; Bian, Q.; Zhang, Z.; Zhang, H.H.; Qian, K. Growth-promoting mechanism and biological activity of weed-derived carbon dots on Chinese cabbage and peas. *Ind. Crop Prod.* **2025**, *228*, 120836. [CrossRef]
34. Nakabachi, A.; Shigenobu, S.; Miyagishima, S. Chitinase-like proteins encoded in the genome of the pea aphid, *Acyrthosiphon pisum*. *Insect Mol. Biol.* **2010**, *19*, 175–185. [CrossRef]
35. Pan, Y.; Lü, P.; Wang, Y.; Yin, L.J.; Ma, H.X.; Ma, G.H.; Chen, K.P.; He, Y.Q. In silico identification of novel chitinase-like proteins in the silkworm, *Bombyx mori*, genome. *J. Insect Sci.* **2012**, *12*, 150. [CrossRef] [PubMed]
36. Zhang, Y.; Chen, R.; Lv, X.Y.; Zeng, L.N.; Gao, Z.X.; Wang, X.Y.; Liu, H.M. Knockdown of chitinase genes enhances larvae susceptibility to *Beauveria bassiana* in *Spodoptera frugiperda*. *Pestic. Biochem. Physiol.* **2026**, *216*, 106796. [CrossRef]
37. Zhu, B.; Shan, J.Q.; Li, R.; Liang, P.; Gao, X.W. Identification and RNAi-based function analysis of chitinase family genes in diamondback moth, *Plutella xylostella*. *Pest Manag. Sci.* **2019**, *75*, 1951–1961. [CrossRef] [PubMed]
38. Yang, X.B.; Zhou, C.; Long, G.Y.; Yang, H.; Chen, C.; Jin, D.C. Characterization and functional analysis of chitinase family genes involved in nymph–adult transition of *Sogatella furcifera*. *Insect Sci.* **2021**, *28*, 901–916. [CrossRef]
39. An, S.; Liu, W.J.; Fu, J.W.; Zhang, Z.; Zhang, R.L. Molecular identification of the chitinase genes in *Aedes albopictus* and essential roles of *AaCht10* in pupal-adult transition. *Parasites Vectors* **2023**, *16*, 120. [CrossRef] [PubMed]
40. Zhu, Q.; Arakane, Y.; Banerjee, D.; Beeman, R.W.; Kramer, K.J.; Muthukrishnan, S. Domain organization and phylogenetic analysis of the chitinase-like family of proteins in three species of insects. *Insect Biochem. Mol. Biol.* **2008**, *38*, 452–466. [CrossRef]

https://doi.org/10.3390/insects17010114

41. Wu, Z.Z.; Zhang, W.Y.; Lin, Y.Z.; Li, D.Q.; Shu, B.S.; Lin, J.T. Genome-wide identification, characterization and functional analysis of the chitianse and chitinase-like gene family in *Diaphorina citri*. *Pest Manag. Sci.* **2022**, *78*, 1740–1748. [CrossRef] [PubMed]
42. Kang, C.; Huang, Y.; Yang, H.; Yan, X.F.; Chen, Z.P. A review of carbon dots produced from biomass wastes. *Nanomaterials* **2020**, *10*, 2316. [CrossRef] [PubMed]
43. Das, S.; Debnath, N.; Cui, Y.J.; Unrine, J.; Palli, S.R. Chitosan, carbon quantum dot, and silica nanoparticle mediated dsRNA delivery for gene silencing in *Aedes aegypti*: A comparative analysis. *ACS Appl. Mater. Interfaces* **2015**, *7*, 19530–19535. [CrossRef] [PubMed]
44. Chen, C.; Imran, M.; Feng, X.Y.; Shen, X.H.; Sun, Z.K. Spray-induced gene silencing for crop protection: Recent advances and emerging trends. *Front. Plant Sci.* **2025**, *16*, 1527944. [CrossRef] [PubMed]
45. Lee, Y.; Muthukrishnan, S.; Kramer, K.J.; Sakamoto, T.; Tabunoki, H.; Arakane, Y.; Noh, M.Y. Functional importance of groups I and II chitinases in cuticle chitin turnover during molting in a wood-boring beetle, *Monochamus alternatus*. *Pestic. Biochem. Physiol.* **2023**, *194*, 105496. [CrossRef]
46. Su, C.; Tu, G.; Huang, S.; Yang, Q.; Shahzad, M.F.; Li, F. Genome-wide analysis of chitinase genes and their varied functions in larval moult, pupation and eclosion in the rice striped stem borer, *Chilo suppressalis*. *Insect Mol. Biol.* **2016**, *25*, 401–412. [CrossRef]
47. Dong, W.; Gao, Y.H.; Zhang, X.B.; Moussian, B.; Zhang, J.Z. Chitinase 10 controls chitin amounts and organization in the wing cuticle of *Drosophila*. *Insect Sci.* **2020**, *27*, 1198–1207. [CrossRef]
48. Li, D.Q.; Zhang, J.Q.; Yang, Y.; Liu, J.; Lu, J.J.; Ren, M.F.; Abbas, M.; Zhu, K.Y.; Zhang, J.Z. Identification and RNAi-based functional analysis of chitinase family genes in *Agrotis ipsilon*. *Pest Manag. Sci.* **2022**, *78*, 4278–4287. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

https://doi.org/10.3390/insects17010114

insects

MDPI

Article

Silencing *Miniature* Gene Disrupts Elytral and Hindwing Structures in *Leptinotarsa decemlineata*

Man-Hong Cheng [1,†], Kai-Yun Fu [1,†], Wei Zhou [2], Ji-Feng Shi [2,*] and Wen-Chao Guo [1,*]

[1] Institute of Plant Protection Xinjiang Academy of Agricultural Sciences/Key Laboratory of Integrated Pest Management on Crops in Northwestern Oasis, Ministry of Agriculture/Xinjiang Key Laboratory of Agricultural Biosafety, Urumqi 830091, China; chengmh0325@163.com (M.-H.C.); fukaiyun000@foxmail.com (K.-Y.F.)

[2] College of Sericulture, Textile, and Biomass Sciences, Southwest University, Chongqing 400715, China; zorroice@yeah.net

* Correspondence: shijifeng@swu.edu.cn (J.-F.S.); gwc1966@163.com (W.-C.G.)

† These authors contributed equally to this work.

Simple Summary: The Colorado potato beetle (*Leptinotarsa decemlineata*) is a destructive pest that causes significant damage to potato crops. The beetle's wings are crucial for its flight, and survival, and protection against environmental stressors, such as pesticides, which also make controlling this pest difficult. The *Miniature* (*Mi*) gene plays an essential role in the development of the wings and their protective outer covering (cuticle) in *Drosophila melanogaster*. In this study, we investigated the function of the *LdMi* gene in *L. decemlineata* using RNA interference to suppress its expression. Our results demonstrate that knockdown of the *LdMi* gene caused severe deformities in the elytra and hindwings of the beetle, with significant changes in the chemical composition of the wings, weakening their structural integrity. This study highlights the critical role of the *LdMi* gene in wing development and provides valuable insights into advancing pest control strategies.

Abstract

The Colorado potato beetle (*Leptinotarsa decemlineata*, CPB) is a major pest in potato crops, notorious for its rapid dispersal and insecticide resistance, which are enabled by its robust elytra and flight-capable hindwings. The *Miniature* (*Mi*) gene, encoding a protein with a zona pellucida (ZP) domain, is involved in wing development and cuticle integrity, yet its functional role in beetles remains underexplored. In this study, we cloned and characterized the *LdMi* gene in the CPB and investigated its function using RNA interference (RNAi), morphological analyses, and spectroscopy. *LdMi* encodes a 146.35 kDa transmembrane protein with a conserved ZP domain, clusters with coleopteran homologs, and exhibits relative conservation across insect species. Expression profiling showed high *LdMi* transcript levels in the hindwings, the elytra, and the pupal stages. RNAi knockdown in fourth-instar larvae resulted in severe eclosion defects, including malformed wings and reduced adult weight. Scanning electron microscopy (SEM) revealed disrupted elytral patterns and deformed hindwing veins in knockdown individuals. Spectroscopic analyses using Fourier-transform infrared (FTIR) and Raman spectroscopy indicated a reduction in protein–chitin crosslinking and diminished hydrogen bonding, suggesting compromised cuticular integrity. These results highlight the essential role of *LdMi* in cuticle formation and the surface morphology of the elytra and hindwings, offering new insights into ZP domain proteins in insects.

Keywords: miniature; elytra; hindwing veins; *Leptinotarsa decemlineata*

1. Introduction

The Colorado potato beetle (*Leptinotarsa decemlineata*, CPB) is a notorious defoliator of potato crops and poses a serious threat to global potato production. Originally native to North America, the CPB has expanded its range across Europe, Asia, and Africa, causing significant crop losses [1–4]. In China, the beetle was first recorded in Xinjiang Province in the early 1990s and has since spread into Inner Mongolia, Heilongjiang, and Jilin, endangering additional potato-growing regions [5,6]. The beetle's hindwings enable its strong flight capability, facilitating migration at speeds of 40–50 km per year, while the rigid elytra protect against environmental stress and act as a barrier to insecticides, reducing the effectiveness of chemical control methods [7–11]. This combination of flight mobility and elytral resistance presents considerable challenges for effective pest management.

The zona pellucida (ZP) domain is a conserved protein motif essential for insect development, particularly in remodeling the apical compartment, maintaining cuticle integrity, and facilitating wing morphogenesis [12,13]. ZP domain proteins are widely expressed in the epithelial tissues during development, significantly contributing to structural stability [12–15]. For example, in *Drosophila melanogaster*, ZP proteins like Dusky-like (Dyl) are essential for forming sensory bristles and wing hairs [16]. Additionally, the ZP proteins Papillote and Piopio (Pio) are involved in cell adhesion to the apical extracellular matrix and the organization of microtubules [17]. Research on other insects, such as *Henosepilachna vigintioctopunctata* and *Bombyx mori*, further emphasizes the diverse roles of ZP proteins in maintaining cuticle integrity and wing architecture [18,19]. In *Tribolium castaneum*, the ZP protein Dusky (Dy) functions upstream of Forked (Fork) in wing morphogenesis [20,21], while Dyl operates downstream of Blimp-1 through Shavenbaby (Svb) to regulate larval epidermal pigmentation and metamorphosis [22]. Moreover, knockdown of *TcSvb* significantly reduces the expression of related ZP domain proteins, such as Singed (Sg) [22].

The *Miniature* (*Mi*) gene, a member of the ZP domain protein family, is closely associated with cuticle strength and membrane resilience, both essential for insects' flight performance and survival [12]. In *D. melanogaster*, the *DmMi* gene is critical to the differentiation and maturation of the wings' epithelial cells. Studies have shown that *DmMi* and its homolog *Dusky* encode transmembrane proteins that localize to the apical membranes of wing epithelial cells, playing a significant role in cuticle formation and maintaining cellular integrity [23]. Furthermore, *DmMi* is associated with the neurohormone bursicon and forms a stable complex that facilitates wing cell maturation. Its expression is correlated with wing expansion and apoptosis in epithelial cells, with delayed wing expansion observed in *Drosophila* mutants lacking *Miniature* [24]. *DmMi* is also critical for post-eclosion wing maturation, influencing both apoptosis and the epithelial-to-mesenchymal transition (EMT) during wing development. The absence of *Miniature* compromises wing integrity and disrupts hormone diffusion within the extracellular matrix, leading to defects in wing morphology and function [25]. Despite these findings, the role of Mi in other insect species has yet to be reported.

In this study, we cloned and characterized the *LdMi* gene in *L. decemlineata* and assessed its expression across various developmental stages and tissues using quantitative real-time PCR (qRT-PCR). To investigate the function of *LdMi*, we employed RNA interference (RNAi), which revealed that the knockdown of *LdMi* significantly disrupted the morphology and structure of both the elytra and the hindwings. Additionally, we utilized scanning electron microscopy (SEM) and FTIR/Raman spectroscopy to evaluate the structural and compositional changes following the knockdown. Our findings provide

valuable insights into the role of the *Mi* gene in beetle wing development and deepen our understanding of the functional diversity of ZP domain proteins across insect lineages.

2. Materials and Methods

2.1. Insect Collection and Rearing

L. decemlineata adults were collected from potato fields in Urumqi, Xinjiang (43.82° N, 87.61° E), during spring emergence and reared under the standard conditions [26]. The beetles were maintained at 28 ± 1 °C and 50–60% humidity under a 14:10 light–dark cycle, with fresh potato leaves provided as food. Eggs hatched within seven days, and the larvae passed through four instars: the first three lasted 2 days each, and the fourth lasted 3.5 days. The fourth-instar larvae then burrowed into the soil to pupate and emerged as adults.

2.2. Cloning of the LdMi Gene

The *LdMi* gene sequence was obtained from *L. decemlineata* genome (https://www.hgsc.bcm.edu/arthropods/colorado-potato-beetle-genome-project, accessed on 1 January 2024, PRJNA854273) and transcriptome databases (PRJNA464380). Total RNA was extracted using TRIzol reagent (Invitrogen, New York, NY, USA) according to the manufacturer's instructions. RNA quantity and quality were assessed using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, New York, NY, USA), with purity determined by the OD260/280 and OD260/230 ratios. Gene verification was conducted via PCR using the primers listed in Table S1, followed by 5′- and 3′-RACE to obtain the full-length *LdMi* sequence. The RACE reactions were performed using the SMARTer RACE cDNA amplification kit (Takara Bio, Dalian, China). The full *LdMi* sequence was submitted to GenBank (Accession No. PQ374228).

The open reading frames (ORFs) and corresponding amino acid sequences were identified, with their theoretical isoelectric points, and the molecular weights were calculated using the ExPASy online tool (https://web.expasy.org/peptide_mass/, accessed on 1 January 2025). The exon–intron organization was predicted by aligning the ORFs with genomic sequences and visualized using Adobe Illustrator CS6. The domain analysis was performed using the SMART tool (http://smart.embl-heidelberg.de/ (accessed on 1 January 2025)). A phylogenetic tree was constructed using MEGA6 with neighbor-joining and 1000 bootstrap replicates. The sequence accession numbers for the ZP domain proteins from various insects are listed in Table S2.

2.3. Quantitative Real-Time PCR (qRT-PCR)

qRT-PCR was used to quantify the *LdMi* gene expression and evaluate the RNAi knockdown efficiency in *L. decemlineata*. Gene-specific primers (Table S1) were designed using Beacon Designer 7 (Premier Biosoft International, Palo Alto, CA, USA). cDNA was synthesized from various developmental stages and adult tissues to determine spatiotemporal expression profiles. For the analysis of the RNAi knockdown efficiency, the larvae were fed potato leaves treated with ds*LdMi* or ds*GFP* suspensions for three days, and samples were collected on day 3 to assess the knockdown levels compared to those in the controls.

Total RNA was extracted from the collected samples and reverse-transcribed into cDNA. qRT-PCR was conducted under the previously established conditions [26], using SYBR Green fluorescence for quantification. The gene expression levels were normalized to the geometric mean of the reference genes (RP18, RP4, ARF1, and ARF4), with the data analyzed via the $2^{-\Delta\Delta CT}$ method. Each experiment was performed in triplicate.

2.4. dsRNA Preparation

Synthesis of the dsRNA targeting the *LdMi* transcripts was conducted utilizing the T7 dual promoter system, as per the established protocols [27]. Two non-overlapping regions of the *LdMi* gene were selected to ensure RNAi efficiency. Production occurred in the RNase III-deficient *E. coli strain HT115*, with induction using 0.1 mM IPTG. A dsRNA construct targeting enhanced green fluorescent protein (ds*GFP*) served as a negative control. Following induction, bacterial cultures were processed through centrifugation at 5000× *g* for 10 min, and the resulting pellets were resuspended in 0.05 M phosphate-buffered saline (PBS, pH = 7.4) to their original volume, with the final concentration of dsRNA at approximately 0.5 μg/mL for the bioassays.

2.5. dsRNA Delivery and RNAi Bioassays

The introduction of dsRNA into the *L. decemlineata* larvae followed the previously described methods [28]. Potato leaves were immersed in a dsRNA-containing bacterial suspension for five seconds, air-dried for two hours, and then used for larval feeding. The control groups were fed PBS- and ds*GFP*-treated leaves. Fourth-instar larvae, starved for four hours post-molting, were placed into feeding dishes. with ten larvae per dish and twelve replicates per treatment.

On day three, knockdown efficiency was evaluated through RT-qPCR in three replicates. The remaining larvae were fed untreated leaves to analyze the adult phenotypes. Once their prepupal behavior had been observed, the larvae were transferred into soil-filled cups for pupation. Seven days after pupation, the emerged adults were collected, and any eclosion defects, along with abnormalities in the elytra and the hindwings, were recorded and compared to the control groups.

2.6. Morphological Analyses of the Wings

Following *LdMi* knockdown, eclosion abnormalities, including deformities in the elytra and hindwings, were observed. No significant differences were found between the sexes, so females were selected for further analysis. Wing morphology was assessed by measuring the elytral and hindwing area, elytral areal density (defined as the elytron weight per area), and elytral thickness, following the established methods [28]. The elytra and hindwings from 4-day-old adults were imaged using a Leica DVM6 microscope (Wetzlar, Germany), and the area analysis was conducted using ImageJ software (Fiji version, Win64). The mid-elytral thickness was measured using a Mitutoyo micrometer, achieving a precision of ±0.001 mm. Although each elytron and hindwing was assessed independently, the statistical comparison revealed negligible differences between the left and right elytra, permitting pooling of the datasets for subsequent analyses.

2.7. SEM Analysis of the Elytra and Hindwings

The surface features of the dorsal and ventral sides of the elytra, the dorsal surface of the hindwings, and the hindwing veins in *L. decemlineata* following *LdMi* knockdown were examined using a Phenom scanning electron microscope (PHENOM SCIENTIFIC, Eindhoven, The Netherlands). To improve the conductivity, samples were coated with gold before imaging under high-vacuum conditions. Various regions of each elytron and hindwing were analyzed in detail to document the full range of structural abnormalities and fragility caused by *LdMi* suppression.

2.8. Spectroscopy Analysis

The chemical composition of both the elytra and the hindwings from female beetles with *LdMi* knockdown was analyzed using Fourier-transform infrared (FTIR) and confocal Raman spectrometry on micronized powder, obtained four days post-eclosion. Before spectral acquisition, lyophilized samples were ground into a fine powder and compressed into potassium bromide (KBr) pellets. The FTIR spectra were recorded using a Jasco 460Plus spectrometer (Jasco, Tokyo, Japan), operating at a resolution of 4 cm^{-1} over the range of 4000 to 400 cm^{-1}, with 100 accumulations for each measurement. A Raman analysis was conducted using a WiTec Alpha 300R microscope (WiTec, Ulm, Germany), covering the spectral acquisition from 3500 to 100 cm^{-1} at a resolution of less than 0.5 cm^{-1}. This system allowed for non-destructive biochemical profiling using 785 nm laser excitation, performing 100 scans of 500 ms each. All of the spectral data were processed for baseline correction, smoothing, and intensity normalization. Chitin derived from shrimp served as the reference standard for peak identification.

2.9. The Data Analysis

The data were analyzed using GraphPad Prism 10 software and presented as the mean ± SE. A one-way ANOVA was employed to assess the statistical significance across groups, followed by Tukey's post hoc test for multiple comparisons. p-values were classified as $p < 0.05$, $p < 0.01$, and $p < 0.001$. Since no significant differences were detected between the two dsRNAs targeting different regions of the *LdMi* gene, the data were combined for the subsequent analysis.

3. Results

3.1. Characterization of the LdMi Gene in L. decemlineata

The *LdMi* gene fragments were identified through a comprehensive bioinformatics search across multiple transcriptome databases [29] and genomes [6,30] of *L. decemlineata*, utilizing the amino acid sequences of the *T. castaneum* and *D. melanogaster* Mi proteins as queries. Following molecular cloning and RACE techniques, we successfully obtained a full-length cDNA of 2195 nucleotides. The ORF comprises 1782 nucleotides, which putatively encodes a protein of 146.35 kDa and 593 amino acid residues.

Alignment of the full-length cDNA with the genomic DNA of *L. decemlineata* revealed the structure of the *LdMi* gene, consisting of seven exons (Figure 1A). Subsequent sequence alignment among the Mi proteins from *T. castaneum*, *B. mori*, *Apis mellifera*, *D. melanogaster*, and Anopheles gambiae indicated that Mi encodes a transmembrane protein characterized by a ZP domain (Figure 1B).

Phylogenetic analysis of the Mi proteins and other functionally characterized ZP-domain-containing proteins from various insect species, including Dy, Dyl, Pio, Sg, Fork, Svb, and Blimp-1, revealed distinct clustering patterns. The *Ld*Mi protein grouped closely with the other insect Mi proteins into a well-supported clade, suggesting a shared evolutionary origin and a conserved functional role across these species. Notably, *Ld*Mi clustered tightly with the *Tc*Mi protein from *T. castaneum* within the Coleoptera group. Additionally, the Mi proteins were more closely related to their paralogues, Dy and Dyl, than to the other ZP-domain-containing proteins. This implies that *Ld*Mi may share similar physiological functions with Dy and Dyl while potentially taking on new roles (Figure 1C).




Figure 1. Gene structure, protein alignment, and phylogenetic analysis of the *Miniature* (Mi) protein. (**A**) A schematic of the exon–intron structure of the *LdMi* gene, where exons are shown as green-filled boxes and introns are indicated by lines. (**B**) Multiple sequence alignment of the Mi proteins highlights the ZP domain and the transmembrane region, delineated by black boxes and green lines, respectively. The shading intensity, from light to dark, reflects increasing sequence similarity. Alignment gaps were introduced to optimize the comparison. (**C**) A phylogenetic tree showing the evolutionary relationships of the Mi proteins and additional functionally characterized ZP domain proteins in insects, constructed using the neighbor-joining method with 1000 bootstrap replications. The tree includes eight groups: *Miniature* (Mi), Dusky (Dy), Dusky-like (Dyl), Piopio (Pio), Singed (Sg), Forked (Fork), Shavenbaby (Svb), and Blimp-1. The *L. decemlineata* miniature protein (*Ld*Mi) is indicated by a green dot.

3.2. The Expression of LdMi Across Developmental Stages and Tissues

Before proceeding with the functional analysis of *LdMi*, we evaluated its mRNA levels across various developmental stages and tissues using quantitative real-time PCR (qRT-PCR) (Figure 2). During *L. decemlineata*'s developmental stages, *LdMi* was expressed in all of the samples analyzed, exhibiting a consistent expression pattern. Notably, the expression levels were lowest immediately after molting (day 0) and increased significantly during the mid-development stages of each instar (day 1) for the first to third instars and at the late stage of the fourth instar (day 3.5), with the first instar showing the highest expression on day 1 (Figure 2A). High expression levels were also observed during the pupal stage and at the egg stage. Tissue-specific analyses in adults revealed that the *LdMi* expression was highest in the hindwings, followed by the elytra and the epidermis (Figure 2B).




Figure 2. The transcription patterns of *LdMi* in *L. decemlineata*. (**A**) The temporal expression of *LdMi* was analyzed using cDNA templates prepared from pooled samples across various developmental stages, including mixed eggs (1–3 days old), first- to fourth-instar larvae (collected daily), wandering larvae, prepupae, pupae, and newly emerged adult males and females (day 1). The green circular dots indicate biological triplicates. (**B**) The tissue-specific expression was measured by quantifying the relative transcript levels in the epidermis, gut, fat body, hemocytes, elytra, and hindwings of day-1 adults, as well as the testes and ovaries of day-4 adults. Three independent pools of 20–30 individuals were analyzed for each sample, and qRT-PCR was employed, with technical triplicates. The $2^{-\Delta\Delta Ct}$ method was used to calculate the relative expression, normalizing against eggs (temporal analysis) or hemocytes (tissue analysis). The data represent the mean ± standard error (SE) of three biological replicates, with differing letters indicating significant differences ($p < 0.05$).

3.3. The Effects of LdMi Knockdown on Adult Eclosion and Wing Morphology in L. decemlineata

To investigate the function of *LdMi*, dsRNA specific to the *LdMi* gene was synthesized using *Escherichia coli HT115* and administered to newly molted fourth-instar larvae through the ingestion of treated potato leaves. The larvae were fed dsRNA-soaked leaves for three days, followed by fresh potato leaves until eclosion. ds*GFP* was used as the control. The qRT-PCR analysis indicated a significant reduction in *LdMi* mRNA levels following three days of continuous dsRNA intake (Figure 3A). The larvae treated with *LdMi* dsRNA showed a notable decrease in larval weight by day 4 (Figure S1). While normal pupation was observed (Figure S2), both male and female adults that emerged exhibited significantly lower weights compared to those in the ds*GFP*-treated group (Figures 3C and S3). Furthermore, the eclosion rates declined significantly, primarily due to deformities in both the elytra and the hindwings (Figure 3B,D). Morphometric measurements indicated significant reductions in elytral area, areal density, and thickness (Figures 3D–F and S4), as well as a marked decrease in hindwing area (Figure 3G).




Figure 3. The impact of RNAi-mediated knockdown of *LdMi* on adult development and wing morphology in *L. decemlineata*. Fourth-instar larvae were fed potato leaves treated with ds*GFP* or ds*LdMi* for three days, followed by feeding with fresh leaves until pupation. (**A**) The transcript levels of *LdMi* were assessed on day 3 post-feeding, with green hollow circles representing biological replicates. Although *LdMi* knockdown did not affect pupation, it significantly increased the proportion of adults with deformities (**B**). Letters a, b, and c indicate the results from the one-way ANOVA, showing significant differences between the proportions of normally emerged and deformed females under the ds*LdMi* and ds*GFP* treatments. Female adults exhibited a reduced body weight on day 1 post-emergence (**C**). Knockdown individuals displayed morphological abnormalities, including irregularly shaped elytra and underdeveloped hindwings (**D**). The measurements of elytral area (**E**), areal density (**F**), and hindwing area (**G**) were significantly reduced. Asterisks denote statistically significant differences (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

3.4. Ultrastructural and Compositional Changes in the Elytra and Hindwings Following LdMi Knockdown

The SEM analysis revealed clear morphological differences between the control (ds*GFP*) and *LdMi*-knockdown beetles (Figure 4). In the ds*GFP* group, the dorsal surface of the elytra exhibited smooth, well-defined polygonal patterns (Figure 4(a1,a2)), while the ventral surface contained uniformly arranged microstructures (Figure 4(a3,a4)). In contrast, *LdMi* knockdown resulted in irregular dorsal surfaces with collapsed regions

(Figure 4(b1,b2)) and poorly organized ventral structures (Figure 4(b3,b4)). In the hindwings, ds*GFP*-treated individuals had smooth, raised veins and evenly distributed microtrichia across the membrane (Figure 4(c1–c3)). However, the *LdMi*-knockdown beetles exhibited flattened veins and sparse or missing microtrichia (Figure 4(d1–d3)), suggesting impaired wing development.




Figure 4. Ultrastructural defects in the elytra and hindwings of *L. decemlineata* following *LdMi* knockdown. In the ds*GFP*-treated beetles, the elytra displayed a regular hexagonal pattern on the dorsal surface (**a1**,**a2**) and a smooth, uniform ventral surface (**a3**,**a4**), with magnified views provided in panels (**a2**,**a4**). Hindwings exhibited a scale-like pattern with raised veins (**c1**–**c3**). In contrast, the ds*LdMi*-treated beetles showed a disrupted elytral structure, with a rough dorsal surface (**b1**,**b2**) and an irregular ventral surface (**b3**,**b4**). Knockdown individuals exhibited disorganized veins, flattened profiles, and blunted spiny structures in their hindwings (**d1**–**d3**). The boundaries of the hindwing veins are marked by yellow dashed lines.

To investigate the molecular and nanostructural changes associated with these morphological abnormalities further, FTIR and Raman spectroscopy analyses were performed on the elytra and hindwings of the *LdMi*-knockdown beetles (Figure 5). These spectra were compared to those for commercial chitin samples and the ds*GFP* control beetles to assess the impact of *LdMi* knockdown on the cuticular composition and structural integrity.

The FTIR absorption profiles of the ds*GFP* elytra and hindwings showed characteristic peaks for chitin (1550–1560 cm^{-1} for amide II, N-H bending and C-N stretching; 1650–1660 cm^{-1} for amide I, C=O stretching), proteins (notable peaks at 1550–1560 cm^{-1} and 1307 cm^{-1} for amide III), and lipids (2800–3000 cm^{-1}) (Figure 5A) [31–33]. In contrast, the *LdMi*-knockdown samples displayed significant reductions in these peaks, especially in the amide II and amide III bands, as well as a marked decline in the O-H stretching peak at 3268 cm^{-1}. These alterations indicate a disruption in hydrogen bonding and other structural interactions crucial for maintaining the mechanical integrity of the exoskeleton. Additionally, significant reductions across the fingerprint region (550–1300 cm^{-1}) highlighted the degradation of chitin, proteins, and other structural components (Figure 5A) [34].

Raman spectroscopy further confirmed these findings. The control samples exhibited characteristic peaks at 1158 cm^{-1} and 1515 cm^{-1}, corresponding to saccharide ring and amide bond vibrations, which are associated with the basic chitin and protein structures [33,35,36]. In contrast, the *LdMi*-knockdown samples showed intensified peaks at these positions (Figure 5B). This suggests abnormal overpacking and crystallization defects within the molecular structure, indicating significant changes in the organizational integrity of the cuticular components.




Figure 5. FTIR (**A**) and Raman (**B**) spectra analysis of compositional changes in *LdMi*-suppressed elytra and hindwings. (**A**) The gray dashed lines in the FTIR spectra highlight peaks for chitin and protein at 1650–1660 cm^{-1} (amide I), 1550–1560 cm^{-1} (amide II), and 1307 cm^{-1} (amide III). The black line indicates the reference standard for peak identification of chitin. Blue dotted lines indicate hydrogen bonding peaks at 3448 cm^{-1} (N-H stretching) and 3268 cm^{-1} (O-H stretching), linked to mechanical integrity. (**B**) In the Raman spectra, gray dashed lines mark peaks at 1158 cm^{-1} (saccharide rings) and 1515 cm^{-1} (amide bonds), reflecting chitin and protein structures.

Together, these results highlight the crucial role of *LdMi* in maintaining the structural and molecular integrity of the elytra and the hindwings. *LdMi* knockdown leads to significant changes in both the ultrastructure and composition of these structures, impairing their mechanical properties and development.

4. Discussion

In this study, we cloned and characterized the *LdMi* gene, a member of the zona pellucida (ZP) domain protein family, in *L. decemlineata* [23–25]. RNAi-mediated silencing of *LdMi* resulted in wing abnormalities in adults, underscoring its critical role in wing morphogenesis. Below, we discuss the functional significance of *LdMi* in the context of wing development and cuticle formation.

4.1. Mi Is Conserved and Crucial for Wing Development

Multiple lines of evidence support the role of *LdMi* as a key regulator of wing development in *L. decemlineata*. The phylogenetic analysis revealed that *Ld*Mi belongs to a distinct clade of insect ZP domain proteins, closely related to Dy and Dyl (Figure 1C), which have previously been implicated in cuticle formation and epithelial morphogenesis [18,21,23]. This evolutionary relationship suggests conserved functional roles within this protein subfamily.

Tissue-specific expression profiling revealed that *LdMi* is predominantly expressed in the elytra and hindwing epidermis (Figure 2B), consistent with the findings in *D. melanogaster*, where *DmMi* orthologs are active in the developing pupal wings [23].

Moreover, RNAi-mediated knockdown of *LdMi* resulted in significant wing deformities, including reductions in their size and structural density (Figure 3). These findings are consistent with other studies in *Drosophila* species where Mi proteins regulated wing size and cellular structure [23,25]. For instance, mutations in *Mi* genes in *D. melanogaster* and *Drosophila virilis* lead to a 1.5-fold reduction in wing surface area [37,38]. In *DmMi* loss-of-function mutants, wing size is significantly reduced and the cuticle appears darker compared to these properties in the wild type [24]. Additionally, wing expansion and epithelial apoptosis are delayed, while the overexpression of Mi accelerates wing development [24], reinforcing the requirement for conserved Mi proteins for wing tissue organization.

Interestingly, despite its early-stage expression, *LdMi* knockdown did not visibly impact embryonic or larval epidermal development, suggesting that *LdMi* functions in a stage-specific manner, primarily during cuticle maturation and wing expansion. These observations are consistent with the findings in *Drosophila* and suggest that Mi may have additional roles, yet to be fully understood, during different developmental stages.

4.2. Mi Regulates Wing Surface Architecture and Cuticle Composition

In addition to gross morphological changes, silencing *LdMi* also disrupted the microstructure of the wing cuticle. SEM revealed collapsed and disorganized surface regions in both the elytra and the hindwings of the *LdMi*-knockdown beetles, including flattened veins and a marked reduction in microtrichia (Figure 4). These disruptions indicate a compromised apical extracellular matrix (ECM), which likely affects the integrity of the cuticulin envelope and its association with the cytoskeleton.

Similar morphological defects in wing structure and vein formation have been observed in *Drosophila* miniature (Mi) mutants. These mutants exhibit a reduced epidermal cell size, resulting in smaller wings, and impaired epidermal flattening during pupal development, preventing proper wing expansion [37,38]. This phenotype points to disrupted ECM organization, particularly within the cuticulin envelope, which is essential for wing shaping and mechanical support [23,39]. Disruption of Mi protein function compromises the coordination between the apical membrane, the cytoskeleton, and the ECM, resulting in defective cuticle formation and impaired wing expansion, as observed in both beetles and *Drosophila* [23,24,40].

Biochemical analyses using FTIR and Raman spectroscopy corroborated these structural findings further (Figure 5). Reductions in the amide I, II, and III peaks and attenuated signals for N–H and O–H stretching suggest a loss of protein–chitin crosslinking and hydrogen bonding—both critical for maintaining cuticle rigidity. Decreased intensity in the fingerprint region also indicates impaired chitin fiber organization, which may compromise the mechanical resilience of the wing cuticle.

These findings are consistent with previous reports on other insect models. For example, *Dy* and *Dyl*, which are structurally related to *LdMi*, have been shown to regulate chitin deposition and cuticular structure in *D. melanogaster* [16,23], *Harmonia vigintioctop-*

unctata [18], and *Tribolium castaneum* [20,41]. Knockdown of these genes leads to similar epidermal and cuticular defects, supporting the broader role of ZP domain proteins in cuticle biogenesis and epidermal differentiation.

5. Conclusions

This study demonstrate that *Ld*Mi is a crucial regulator of wing development and cuticular integrity in *L. decemlineata*. RNA-interference-mediated silencing of *LdMi* resulted in severe morphological deformities in both the elytra and the hindwings, leading to abnormal wing structures, a reduced adult body weight, and impaired eclosion. Spectroscopic analyses showed that these morphological defects were linked to compromised protein–chitin crosslinking and weakened hydrogen bonding, highlighting the gene's role in maintaining both biochemical stability and mechanical strength.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/insects16070700/s1. Figure S1. Effect of *LdMi* gene knockdown on larval weight. Figure S2. Effect of *LdMi* knockdown on pupation rate. Figure S3. Effect of *LdMi* knockdown on male body weight. Figure S4. Effect of *LdMi* knockdown on elytral thickness. Table S1. Primers used in RT-PCR, RACE, ORF verification, dsRNA synthesis, and qRT-PCR. Table S2. GenBank accession numbers of representative ZP domain proteins.

Author Contributions: The experiments were designed by M.-H.C., K.-Y.F., J.-F.S. and W.-C.G.; the RNAi and TEM analyses was implemented by K.-Y.F. and J.-F.S.; software was used by M.-H.C. and J.-F.S.; and the figures were designed and created by M.-H.C., W.Z. and J.-F.S. The statistical analyses were implemented by M.-H.C.; resources were provided by K.-Y.F. and W.-C.G.; writing of the manuscript in the form of original draft preparation was carried out by M.-H.C. and J.-F.S.; writing in the form of review and editing was carried out by M.-H.C., K.-Y.F. and J.-F.S.; project administration was carried out by K.-Y.F., W.-C.G. and J.-F.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the Major Science and Technology Projects in Xinjiang (No. 2023A02006), the Project of Fund for Stable Support to Agricultural Sci-Tech Renovation (No. xjnkywdzc-2023004), and the Natural Science Foundation Project of Chongqing (No. CSTB2022NSCQ-MSX0663).

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Materials. Further inquiries can be directed to the corresponding authors.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Lomakin, J.; Huber, P.A.; Eichler, C.; Arakane, Y.; Kramer, K.J.; Beeman, R.W.; Kanost, M.R.; Gehrke, S.H. Mechanical properties of the beetle elytron, a biological composite material. *Biomacromolecules* **2011**, *12*, 321–335. [CrossRef]
2. Orlowski, G.; Mroz, L.; Kadej, M.; Smolis, A.; Tarnawski, D.; Karg, J.; Campanaro, A.; Bardiani, M.; Harvey, D.J.; Mendez, M.; et al. Breaking down insect stoichiometry into chitin-based and internal elemental traits: Patterns and correlates of continent-wide intraspecific variation in the largest European saproxylic beetle. *Environ. Pollut.* **2020**, *262*, 114064. [CrossRef] [PubMed]
3. Kadoic Balasko, M.; Mikac, K.M.; Bazok, R.; Lemic, D. Modern Techniques in Colorado Potato Beetle (*Leptinotarsa decemlineata* Say) Control and Resistance Management: History Review and Future Perspectives. *Insects* **2020**, *11*, 581. [CrossRef] [PubMed]
4. Alyokhin, A.; Baker, M.; Mota-Sanchez, D.; Dively, G.; Grafius, E. Colorado Potato Beetle Resistance to Insecticides. *Am. J. Potato Res.* **2008**, *85*, 395–413. [CrossRef]
5. Li, C.; Liu, H.; Huang, F.; Cheng, D.F.; Wang, J.J.; Zhang, Y.H.; Sun, J.R.; Guo, W.C. Effect of temperature on the occurrence and distribution of colorado potato beetle (Coleoptera: Chrysomelidae) in China. *Environ. Entomol.* **2014**, *43*, 511–519. [CrossRef] [PubMed]
6. Yan, J.; Zhang, C.; Zhang, M.; Zhou, H.; Zuo, Z.; Ding, X.; Zhang, R.; Li, F.; Gao, Y. Chromosome-level genome assembly of the *Colorado potato beetle*, *Leptinotarsa decemlineata*. *Sci. Data* **2023**, *10*, 36. [CrossRef]

7. Alyokhin, A.; Mota-Sanchez, D.; Baker, M.; Snyder, W.E.; Menasha, S.; Whalon, M.; Dively, G.; Moarsi, W.F. The Red Queen in a potato field: Integrated pest management versus chemical dependency in *Colorado potato beetle* control. *Pest Manag. Sci.* **2015**, *71*, 343–356. [CrossRef]
8. Timani, K.; Bastarache, P.; Morin, P.J. Leveraging RNA Interference to Impact Insecticide Resistance in the Colorado Potato Beetle, *Leptinotarsa decemlineata*. *Insects* **2023**, *14*, 418. [CrossRef]
9. Chen, Y.H.; Cohen, Z.P.; Bueno, E.M.; Christensen, B.M.; Schoville, S.D. Rapid evolution of insecticide resistance in the *Colorado potato beetle*, *Leptinotarsa decemlineata*. *Curr. Opin. Insect Sci.* **2023**, *55*, 101000. [CrossRef]
10. Scott, I.M.; Tolman, J.H.; MacArthur, D.C. Insecticide resistance and cross-resistance development in *Colorado potato beetle Leptinotarsa decemlineata* Say (Coleoptera: Chrysomelidae) populations in Canada 2008–2011. *Pest Manag. Sci.* **2015**, *71*, 712–721. [CrossRef]
11. Cingel, A.; Savic, J.; Lazarevic, J.; Cosic, T.; Raspor, M.; Smigocki, A.; Ninkovic, S. Extraordinary Adaptive Plasticity of Colorado Potato Beetle: "Ten-Striped Spearman" in the Era of Biotechnological Warfare. *Int. J. Mol. Sci.* **2016**, *17*, 1538. [CrossRef] [PubMed]
12. Litscher, E.S.; Wassarman, P.M. Zona Pellucida Proteins, Fibrils, and Matrix. *Annu. Rev. Biochem.* **2020**, *89*, 695–715. [CrossRef]
13. Fernandes, I.; Chanut-Delalande, H.; Ferrer, P.; Latapie, Y.; Waltzer, L.; Affolter, M.; Payre, F.; Plaza, S. Zona pellucida domain proteins remodel the apical compartment for localized cell shape changes. *Dev. Cell* **2010**, *18*, 64–76. [CrossRef]
14. Jovine, L.; Qi, H.; Williams, Z.; Litscher, E.S.; Wassarman, P.M. A duplicated motif controls assembly of zona pellucida domain proteins. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 5922–5927. [CrossRef]
15. Jazwinska, A.; Affolter, M. A family of genes encoding zona pellucida (ZP) domain proteins is expressed in various epithelial tissues during Drosophila embryogenesis. *Gene Expr. Patterns GEP* **2004**, *4*, 413–421. [CrossRef] [PubMed]
16. Adler, P.N.; Sobala, L.F.; Thom, D.; Nagaraj, R. dusky-like is required to maintain the integrity and planar cell polarity of hairs during the development of the Drosophila wing. *Dev. Biol.* **2013**, *379*, 76–91. [CrossRef]
17. Bokel, C.; Prokop, A.; Brown, N.H. Papillote and Piopio: Drosophila ZP-domain proteins required for cell adhesion to the apical extracellular matrix and microtubule organization. *J. Cell Sci.* **2005**, *118*, 633–642. [CrossRef] [PubMed]
18. Zhang, Y.; Tan, Q.; Lin, M.; Shen, C.; Jin, L.; Li, G. Dusky-like Is Critical for Morphogenesis of the Cellular Protuberances and Formation of the Cuticle in *Henosepilachna vigintioctopunctata*. *Biology* **2023**, *12*, 866. [CrossRef]
19. Zou, Y.L.; Ding, X.; Zhang, L.; Xu, L.F.; Liang, S.B.; Hu, H.; Dai, F.Y.; Tong, X.L. Bmmp influences wing morphology by regulating anterior-posterior and proximal-distal axes development. *Insect Sci.* **2022**, *29*, 1569–1582. [CrossRef]
20. Li, C.; Li, B.; Ma, S.; Lu, P.; Chen, K. Dusky works upstream of Four-jointed and Forked in wing morphogenesis in *Tribolium castaneum*. *Insect Mol. Biol.* **2017**, *26*, 677–686. [CrossRef]
21. Li, C.; Yang, L.; Wang, Y.; Du, H.; Zhang, J.; Lu, Y.; Li, B.; Chen, K. Functional analysis of zona pellucida domain protein Dusky in *Tribolium castaneum*. *Insect Sci.* **2022**, *29*, 388–398. [CrossRef] [PubMed]
22. Li, C.; Yun, X.; Li, B. Dusky-like is required for epidermal pigmentation and metamorphosis in *Tribolium castaneum*. *Sci. Rep.* **2016**, *6*, 20102. [CrossRef]
23. Roch, F.; Alonso, C.R.; Akam, M. Drosophila miniature and dusky encode ZP proteins required for cytoskeletal reorganisation during wing morphogenesis. *J. Cell Sci.* **2003**, *116*, 1199–1207. [CrossRef] [PubMed]
24. Bilousov, O.O.; Katanaev, V.L.; Demydov, S.V.; Kozeretska, I.A. The downregulation of the *Miniature* gene does not replicate *Miniature* loss-of-function phenotypes in *Drosophila melanogaster* wing to the full extent. *Cytol. Genet.* **2013**, *47*, 124–127. [CrossRef]
25. Bilousov, O.O.; Kozeretska, I.A.; Katanaev, V.L. Role of the gene Miniature in Drosophila wing maturation. *Genesis* **2012**, *50*, 525–533. [CrossRef] [PubMed]
26. Chen, Y.; Tang, H.; Zhou, W.; Li, C.; Chen, Y.N.; Zhang, Q.; Fu, K.Y.; Guo, W.C.; Shi, J.F. Identification of chitinase genes and roles in the larval-pupal transition of *Leptinotarsa decemlineata*. *Pest Manag. Sci.* **2024**, *80*, 282–295. [CrossRef]
27. Shi, J.F.; Mu, L.L.; Chen, X.; Guo, W.C.; Li, G.Q. RNA interference of chitin synthase genes inhibits chitin biosynthesis and affects larval performance in *Leptinotarsa decemlineata* (Say). *Int. J. Biol. Sci.* **2016**, *12*, 1319–1331. [CrossRef]
28. Shi, J.F.; Cheng, M.H.; Zhou, W.; Zeng, M.Z.; Chen, Y.; Yang, J.X.; Wu, H.; Ye, Q.H.; Tang, H.; Zhang, Q.; et al. Crucial roles of specialized chitinases in elytral and hindwing cuticles construction in *Leptinotarsa decemlineata*. *Pest Manag. Sci.* **2024**, *80*, 4437–4449. [CrossRef]
29. Kumar, A.; Congiu, L.; Lindstrom, L.; Piiroinen, S.; Vidotto, M.; Grapputo, A. Sequencing, *De Novo* assembly and annotation of the Colorado Potato Beetle, *Leptinotarsa decemlineata*, Transcriptome. *PLoS ONE* **2014**, *9*, e86012. [CrossRef]
30. Schoville, S.D.; Chen, Y.H.; Andersson, M.N.; Benoit, J.B.; Bhandari, A.; Bowsher, J.H.; Brevik, K.; Cappelle, K.; Chen, M.M.; Childers, A.K.; et al. A model species for agricultural pest genomics: The genome of the *Colorado potato beetle*, *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). *Sci. Rep.* **2018**, *8*, 1931. [CrossRef]
31. Triunfo, M.; Tafi, E.; Guarnieri, A.; Salvia, R.; Scieuzo, C.; Hahn, T.; Zibek, S.; Gagliardini, A.; Panariello, L.; Coltelli, M.B.; et al. Characterization of chitin and chitosan derived from *Hermetia illucens*, a further step in a circular economy process. *Sci. Rep.* **2022**, *12*, 6613. [CrossRef] [PubMed]

32. Kaya, M.; Baran, T.; Erdogan, S.; Mentes, A.; Ozusaglam, M.A.; Cakmak, Y.S. Physicochemical comparison of chitin and chitosan obtained from larvae and adult *Colorado potato beetle* (*Leptinotarsa decemlineata*). *Mater. Sci. Eng. C* **2014**, *45*, 72–81. [CrossRef]
33. Jantzen da Silva Lucas, A.; Quadro Oreste, E.; Leao Gouveia Costa, H.; Martin Lopez, H.; Dias Medeiros Saad, C.; Prentice, C. Extraction, physicochemical characterization, and morphological properties of chitin and chitosan from cuticles of edible insects. *Food Chem.* **2021**, *343*, 128550. [CrossRef]
34. Vargas, W.E.; Avendano, E.; Hernandez-Jimenez, M.; Azofeifa, D.E.; Libby, E.; Solis, A.; Barboza-Aguilar, C. Photonic Crystal Characterization of the Cuticles of *Chrysina chrysargyrea* and *Chrysina optima* Jewel Scarab Beetles. *Biomimetics* **2018**, *3*, 30. [CrossRef] [PubMed]
35. Dimzon, I.K.; Knepper, T.P. Degree of deacetylation of chitosan by infrared spectroscopy and partial least squares. *Int. J. Biol. Macromol.* **2015**, *72*, 939–945. [CrossRef]
36. Kumirska, J.; Czerwicka, M.; Kaczynski, Z.; Bychowska, A.; Brzozowski, K.; Thoming, J.; Stepnowski, P. Application of spectroscopic methods for structural analysis of chitin and chitosan. *Mar. Drugs* **2010**, *8*, 1567–1636. [CrossRef]
37. Dobzhansky, T. The influence of the quantity and quality of chromosomal material on the size of the cells in drosophila melanogaster. *W. Roux' Archiv f. Entwicklungsmechanik* **1929**, *115*, 363–379. [CrossRef]
38. Kozeretska, I.A.; Gubenko, I.S.; Gorb, S.N. New unusual miniature-like wing mutation in *Drosophila virilis*. *J. Morphol.* **2004**, *261*, 270–275. [CrossRef] [PubMed]
39. Newby, L.M.; White, L.; DiBartolomeis, S.M.; Walker, B.J.; Dowse, H.B.; Ringo, J.M.; Khuda, N.; Jackson, F.R. Mutational analysis of the Drosophila miniature-dusky (m-dy) locus: Effects on cell size and circadian rhythms. *Genetics* **1991**, *128*, 571–582. [CrossRef]
40. Ren, N.; Zhu, C.; Lee, H.; Adler, P.N. Gene expression during Drosophila wing morphogenesis and differentiation. *Genetics* **2005**, *171*, 625–638. [CrossRef]
41. Bilousov, O.O.; Katanaev, V.L.; Kozeretska, I.A. Miniature as a hypothetical regulatory protein of the Bursicon/Rickets signaling cascade in the wing epithelia of *Drosophila melanogaster*. *Biopolym. Cell* **2012**, *28*, 288–291. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Silencing the *cyp314a1* and *cyp315a1* Genes in the *Aedes albopictus* 20E Synthetic Pathway for Mosquito Control and Assessing Algal Blooms Induced by Recombinant RNAi *Microalgae*

Xiaodong Deng [1,2], Changhao He [1], Chunmei Xue [3], Dianlong Xu [2,4,5], Juncai Li [1] and Xiaowen Fei [1,*]

[1] Department of Biochemistry and Molecular Biology & Key Laboratory of Tropical Translational Medicine of Ministry of Education, College of Basic Medical Sciences, Hainan Medical University, Haikou 571199, China
[2] Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Science & Key Laboratory of Biology and Genetic Resources of Tropical Crops of Hainan Province, Hainan Institute for Tropical Agricultural Resources, Haikou 571101, China
[3] College of Life Sciences and Agriculture, Jiamusi University, Jiamusi 154007, China
[4] Hainan Provincial Key Laboratory for Functional Components Research and Utilization of Marine Bio-Resources, Haikou 571101, China
[5] Zhanjiang Experimental Station, Chinese Academy of Tropical Agricultural Science, Zhanjiang 524013, China
* Correspondence: xiaowen.fei@muhn.edu.cn

Simple Summary: This study explores a novel approach to controlling *Aedes albopictus*, a key vector for dengue fever, by using RNA interference (RNAi) mediated by genetically modified microalgae. We constructed double-stranded RNA (dsRNA) expression vectors that target the *cyp314a1* and *cyp315a1* genes of *Ae. albopictus*, and we transformed two species of algae, *Chlamydomonas reinhardtii* and *Chlorella vulgaris*, to express these dsRNAs. The recombinant RNAi algal strains were then fed to *Ae. albopictus* larvae, resulting in high mortality rates, disrupted pupation and reduced emergence. The algae triggered an increase in reactive oxygen species (ROS) and enzyme activity in the larvae, indicating redox stress. In simulated field experiments, RNAi recombinant *Chlorella* significantly reduced mosquito populations, and improved water quality by reducing nitrogen, phosphorus and others. Furthermore, with the release of RNAi recombinant *Chlorella* into the test water, the biotic community restructuring dominated by resource competition caused by algal bloom, as well as the proliferation of anaerobic bacteria and the decline of aerobic bacteria triggered by anaerobic conditions, are the main trends in the changes in the test water. This study is an important addition to the use of RNAi recombinant microalgae as a biocide.

Abstract

As one of the key vectors for the transmission of Dengue fever, *Aedes albopictus* is highly ecologically adaptable. The development of environmentally compatible biological defence and control technologies has therefore become an urgent need for vector biological control worldwide. This study constructed and used double-stranded RNA (dsRNA) expression vectors targeting the *cyp314a1* and *cyp315a1* genes of *Ae. albopictus* to transform *Chlamydomonas reinhardtii* and *Chlorella vulgaris*, achieving RNA interference (RNAi)-mediated gene silencing. The efficacy of the RNAi recombinant algal strain biocide against *Ae. albopictus* was evaluated by administering it to *Ae. albopictus* larvae. The results showed that the oral administration of the *cyp314a1* and *cyp315a1* RNAi recombinant *C. reinhardtii*/*C. vulgaris* strains was lethal to *Ae. albopictus* larvae and severely affected their pupation and emergence. The recombinant algal strains triggered a burst of ROS (Reactive Oxygen Species) in the mosquitoes' bodies, resulting in significant increases in the activities of the superoxide dismutase (SOD), peroxiredoxin (POD) and catalase (CAT), as well as significant

upregulation of the mRNA levels of the CME pathway genes in larvae. In the simulated field experiment, the number of *Ae. albopictus* was reduced from 1000 to 0 in 16 weeks by the RNAi recombinant *Chlorella*, which effectively controlled the population of mosquitoes. Meanwhile, the levels of nitrogen (N), phosphorus (P), nitrate, nitrite, ammonia and COD (Chemical Oxygen Demand) in the test water decreased significantly. High-throughput sequencing analyses of 18S rDNA and 16S rDNA showed that, with the release of RNAi recombinant *Chlorella* into the test water, the biotic community restructuring dominated by resource competition caused by algal bloom, as well as the proliferation of anaerobic bacteria and the decline of aerobic bacteria triggered by anaerobic conditions, are the main trends in the changes in the test water. This study is an important addition to the use of RNAi recombinant microalgae as a biocide.

Keywords: CYP314A1 and CYP315A1; *Chlamydomonas reininatus*; *Chlorella*; RNAi; *Aedes albopictus*; 18S rDNA and 16S rDNA high-throughput sequencing

1. Introduction

Dengue fever is an acute arboviral disease primarily transmitted through *Aedes* mosquito bites [1]. According to World Health Organization (WHO) surveillance data (2023 update), reveal a dramatic tenfold increase in reported cases, from 500,000 in 2000 to 5.2 million in 2023. This increase has been accompanied by over 5000 dengue-related fatalities, representing the highest mortality figures documented since systematic monitoring began [2–4].

The primary vectors of Dengue fever are *Aedes* mosquitoes, notably *Ae. aegypti* and *Ae. albopictus*. These species alsotransmit Zika virus disease, Yellow fever, Rift Valley fever, and Chikungunya fever [5–7]. *Ae. albopictus*, in particular, demonstrates high ecological plasticity and has become the main dengue vector in China. It is distributed across a wide range of climates, from temperate regions (up to 45° north latitude) to tropical ecosystems.

Using biotechnological interventions to suppress vector populations represents an inevitable paradigm shift in the control of mosquito-borne diseases. Current advancements focus on two main strategies: Wolbachia-based population replacement and transgenic mosquito approaches that induce male sterility through genetic modification [8–12]. However, scalability remains challenging due to the substantial costs associated with the mass rearing of mosquitoes and the sex sorting protocols required for field implementation [13,14]. Microalgae-mediated biocontrol is a promising alternative that leverages natural trophic relationships in mosquito larval habitats. Microalgae such as *Chlorella* spp., *Scenedesmus* spp., *Spirulina* spp. and *Chlamydomonas* spp. are essential dietary components for developing *Aedes* larvae, and can be cultivated on a large scale at low cost. These microalgae offer distinct operational advantages. Compared to Wolbachia-infected or transgenic male-sterile *Ae. aegypti* strains, algal-based interventions are more economically feasible and easier to control environmentally. Strategic deployment in contained aquatic ecosystems enables rapid algal dominance, thus gradually reducing the number of Aedes mosquitoes in the area. This novel approach to vector suppression is particularly promising in interrupting the transmission networks of arboviruses such as dengue, Zika and yellow fever.

We have used microalgae expressing RNA interference fragments of the Aedes *HR3* (*Hormone receptor 3*), *3HKT* (*3-Hydroxykynurenine transaminase*) and *CHSA* (*Chitin synthase A*) genes to effectively control *Aedes* populations [15–17]. However, it has been shown that prolonged selection pressure for silencing against specific gene targets leads to tolerance (drug resistance) in mosquitoes to the targeted RNAi-interfering algal strains [18–21]. Cytochrome P450 (CYP450) is an ancient and widely functional superfamily of monooxygenases [22]. The genes encoding these enzymes are among the largest gene superfamilies in plants, animals and fungi and are essential for development, growth, and reproduction [23,24]. CYP450 are present in almost all organisms, including some viruses, with a few exceptions, such as *Escherichia coli*. Insects possess only four CYP450 clades, which is significantly fewer than the ten found in humans [25,26]. The functions of CYP450 in insects are diverse and include metabolic detoxification, growth and development regulation, and environment adaptation. In metabolic detoxification, insects adapt to the chemical defences of host plants by catalysing the hydroxylation or oxidation of toxic plant xenobiotics to produce less toxic or non-toxic compounds via CYP450 [27–35]. CYP450 can also degrade insecticides [36,37]. For instance, RNAi silencing of the MusiDN2722 CYP450 gene in *Megalurothrips usitatus* significantly increased susceptibility to acetamiprid, confirming the direct role of CYP450 in insecticide detoxification [38]. Additionally, insect CYP450s participate in the biosynthesis of ecdysone to regulate larval moulting, metamorphosis, and embryonic development [39–42]. These CYP450s include neverland (CYP307A1), spookier (CYP307A2), phantom (CYP306A1), disembodied (CYP302A1), shadow (CYP315A1), and shade (CYP314A1). CYP315A1 and CYP314A1 jointly catalyse the final conversion of cholesterol into the physiologically active 20-hydroxyecdysterone (20E) [43,44]. Given the crucial role of ecdysone in inset development and metamorphosis, this study selected the key enzymes CYP314A1 and CYP315A1 in the 20E anabolic pathway of *Aedes* mosquitoes as RNAi targets. The dsRNAs of these enzymes were then recombined into the genomes of *Chlorella vulgaris* and *Chlamydomonas reinhardtii*, and the microalgae were fed to *Aedes* mosquito larvae in order to inhibit the growth of *Aedes* mosquito populations.

This study aims to utilise low-cost microalgae by releasing recombinant strains into waters adjacent to mosquito breeding sites to control local mosquito populations, while concurrently assessing the environmental impact of algal blooms induced by these engineered microalgae. However, due to governmental regulations on genetically modified organisms (GMOs), the direct release of recombinant microalgae into natural water bodies (e.g., ponds and lakes) was prohibited. Consequently, we established a contained experimental area that simulated wild-type aquatic conditions in order to replicate microalgal growth and reproduction dynamics, while also monitoring of mosquito population numbers. From September 2022 to June 2023, we systematically monitored phytoplankton, zooplankton, and prokaryotic communities in the simulated aquatic environment using water quality assays and 18S and 16S rDNA high-throughput sequencing. The population densities and structural profiles of aquatic organisms were analysed under *Chlorella* bloom conditions. These findings provide foundational data for the future practical application of recombinant microalgae in the control of mosquito populations and the prevention of mosquito-borne diseases such as dengue fever.

2. Methods

2.1. Bioinformatic Analysis

The CYP314A1 and CYP315A1 proteins in *Ae. albopictus* and other organisms were identified using the *Ae. albopictus* CYP314A1 and CYP315A1 protein sequence as seed sequences in BLASTP (Version 2.13.0, Bethesda, MA, USA) searches of the Vectorbase (https://vectorbase.org/vectorbase/app, accessed on 7 May 2024) and NCBI (National

Center for Biotechnology Information) (https://www.ncbi.nlm.nih.gov/, accessed on 7 May 2024) databases. An online InterProScan analysis (https://www.ebi.ac.uk/interpro/search/sequence/, accessed on 16 April 2025) was carried out to identify the domains of the CYP314A1 and CYP315A1 proteins. A phylogenetic tree was then constructed using MEGA11 software (Version 11, Temple University, Philadelphia, PA, USA) based on the alignment of the CYP314A1 and CYP315A1 orthologue in *Ae. albopictus* and other organisms. The tree was generated using the neighbour-joining (NJ) method with Poisson correction and pairwise deletion parameters. Based on sequence information obtained from online databases, a schematic representation of CYP314A1 and CYP315A1 homologous protein structures was created. To identify CYP450 homologous proteins, conserved domain sequences of *Ae. albopictus* CYP450, including WxxxR (Helix-C), GxE/DTT/S (Helix-I), ExxR (Helix-K), PxxFxPE/DRF (PERF), and PFxxGxRxCxG/A (heme-binding domain), were queried in the PFAM (http://pfam.xfam.org/, accessed on 16 April 2025), NCBI and VactorBase (https://vectorbase.org/vectorbase/app, accessed on 15 April 2025) databases After filtering out redundant, 100% identical transcripts from the protein-coding sequence, those encoding at least 300 amino acids were retained. All CYP450 amino acid sequences were collected and annotated using InterPro, and multiple sequences were aligned using ClustalW (Version 11, Dublin, Ireland) and MEGA11. The JTT model was selected for the above processed sequences using maximum likelihood tree. After constructing the tree, iTOL was used to classify it by clan.

2.2. Mosquito Maintenance

Mosquitoes rearing was performed as previously reported [15–17]. Local wild type *Ae. albopictus* were captured in Haikou, China, and propagated for five generations in the laboratory, which were kept in an insectary at the Chinese Academy of Tropical Agricultural Science. The mosquitoes were kept at a temperature of 25 °C and a relative humidity of 80%, with a 12 h cycle of darkness and light. The mosquito cages were also equipped with a sponge saturated with sugar water to provide nourishment. Chicken blood was used to feed the female mosquitoes. Following oviposition on moist filter paper, the eggs were harvested and kept in a dry state. The eggs were then transferred to incubation water containing rat food to facilitate hatching.

2.3. Microalgae Strains and Plasmids

The *Chlorella vulgaris* HOC5 strain was isolated from local waters in Hainan, China, and cultured in TAP medium. *Chlamydomonas reinhardtii* CC425 was purchased from the Freshwater Algae Culture Collection at the Institute of Hydrobiology (FACHB), Chinese Academy of Sciences. Liquid cultures were maintained at 25 °C in a shaker at 180 rpm under continuous light (150 μmol/(m^2·s)) [16]. Plasmid pMD18-T was constructed locally by our research group and stored in our laboratory. The pMaa7 IR/XIR RNAi expression vector was purchased from the *Chlamydomonas* centre at Duke University in Durham, NC, USA.

2.4. Construction of RNAi Vectors

Total RNA was extracted from *Ae. albopictus* larvae using TRIzol reagent (Takara, Shiga, Japan) according to the manufacturer's protocol [15]. Complementary DNA (cDNA) was then synthesised from the total RNA using oligo-dT primers (Shenggong, Shanghai, China) and served as the template for PCR amplification. The *cyp314a1* (AALFPA_063667) and *cyp315a1* (AALFPA_079547) DNA fragment were PCR-amplified using the primers listed in Supplementary Table S1, respectively. These primers target the corresponding *cyp314a1* and *cyp315a1* CDS regions from 390 to 805 and 5 to 324, respectively. The amplified fragments were then inserted into the pMD18-T plasmid backbone (pMD-cyp314a1/cyp315a1).

These fragments were then digested with HindIII/BamHI and XbaI/SalI, before being inserted into the corresponding cloning sites of the pT282 plasmid, which contained the inverted repeat sequence of the respective genes. The RNAi (pMaa7 IR/XIR) and intermediate (pT282-cyp314a1/cyp315a1) vectors were both digested using EcoRI and then ligated to produce the RNAi recombinant vectors, pMaa7 IR/cyp314a1IR and pMaa7 IR/cyp315a1IR [45].

2.5. Microalgae Transfection

The *Chlamydomonas/Chlorella* was grown, collected by centrifugation, and resuspended in electroporation buffer. For transfection, 2 μg of either pMaa7 IR/cyp314a1IR or pMaa7 IR/cyp315a1IR was mixed with the 100 μL recipient *Chlamydomonas/Chlorella* cells (2×10^3 cells μL^{-1}) and the mixture was then incubated on ice for 15 min. The mixture was then transferred into an electroporation cuvette and subjected to an electric pulse in an electrometer (BTX, ECM 630, Shelton, WA, USA) with a 1600 V/cm voltage and a pulse time of 1 ms. The suspension was then transferred to TAP medium containing 60 mmol/L sorbitol and allowed to recover overnight. Finally, the cells were harvested by centrifugation at 10,000× *g* for 3 min and plated on TAP agar plates containing 10 μg/mL paromomycin. The plates were incubated at 23 °C under continuous illumination of 100 μmol m^{-2} s^{-1}. Algal colonies appeared after 5–10 days [46,47].

2.6. Mosquito Feeding Experiments

After the RNAi vector, pMaa7IR/cyp314a1IR and pMaa7IR/cyp315a1IR had been transferred into *Chlamydomonas/Chlorella*. The transgenic microalgae were characterised using PCR to determine if the *cyp314a1* or *cyp315a1* inverted repeat expression cassette had been integrated into the *Chlamydomonas/Chlorella* genome. The positive clones were used to feed the mosquitoes in the laboratory. As previously described, ten and 300 mosquito feeding experiments were performed in the insectary [15,16]. In the 10-mosquito feeding experiment, the mosquitoes were divided into experimental and control groups. Each group contained 10 L1 larvae in 5 mL water supplemented with 75 mg fresh microalgae. The larvae in the experimental groups were fed the *cyp314a1* and *cyp315a1* RNAi transgenic *Chlamydomonas/Chlorella* strains, respectively. In contrast, the larvae in the control groups were fed *C. reinhardtii* CC425 or *C. vulgaris* HOC5, water, feed, and the empty plasmid pMaa7IR/XIR transgenic strain. The experiments were performed in triplicate, recording the mortality, pupation, and adult emergence rates.

In the 300-mosquito feeding experiment, 300 L1 larvae were placed into 50 mL of water supplemented with 750 mg of fresh microalgae. The larvae feeding the recombinant *Chlamydomonas/Chlorella* were set as the test treatments, whereas the larvae fed with *C. reinhardtii* CC425 or *C. vulgaris* HOC5, water, and feed were used as controls. The triplicate experiments included records of mortality, pupation, and adult emergence rates.

To measure body size, five third-instar mosquito larvae were randomly selected from each treatment and control group. Larva was placed a stereomicroscope (Keyence VHX-1000 Super Depth 3D Stereomicroscope, Osaka, Japan), and body length and width were measured using the built-in scale. Body width was determined at the widest segment, and body length was measured from the anterior margin of the head to the tip of the abdomen. The values represented are the mean measurements from the five larvae.

2.7. Measurement of Larval Superoxide Dismutase (SOD), Peoxidase (POD) and Catalase (CAT) Activities

The larvae were fed either *C. reinhardtii* CC425 or *C. vulgaris* HOC5 (used as controls). Some larvae were also fed recombinant *Chlamydomonas* CC425-CYP314A1 and CC425-CYP315A1, while others were fed recombinant *Chlorella* HOC5-CYP314A1 and HOC5-CYP315A1. Ten to fifteen L4 larvae were sampled from each group, mixed with 5 mmol/L phosphate-buffer solution (pH 7.8, Shenggong, Shanghai, China), ground and homogenised on ice, then centrifuged at 10,000× *g* for 20 min at 4 °C. The soluble protein content was determined using the G-250 method (Bradford Protein Concentration Assay Kit, Beyotime Biotechnology, Shanghai, China). SOD activity was measured using an SOD activity assay kit (Beyotime Biotechnology, Shanghai, China), POD activity was measured using a peroxidase (POD) activity assay kit (Beijing Solarbio Technology Co., Ltd., Beijing, China), and CAT activity using a catalase assay kit (Beyotime Biotechnology, Shanghai, China) [48–51].

2.8. The 20-Hydroxyecdysone(20E) Compensation Test

A total of six groups were tested. In each group, 30 *Ae. albopictus* larvae were fed a 15 mg/mL algal solution. 300 μL of 20E (1 mg/mL) was added to the relevant groups, and the mosquitoes were fed for 15 days [52,53]. Mosquitoes fed *C. reinhardtii* CC425 and *C. reinhardtii* CC425+20E were used as control groups. The treatment groups comprised mosquitoes fed *cyp314a1* and *cyp315a1* RNAi transgenic *Chlamydomonas* lines and mosquitoes fed *cyp314a1* and *cyp315a1* RNAi transgenic *Chlamydomonas* lines with the addition of 20E. The mortality rate, pupation rate and eclosion rate were recorded for each group. Total RNA was extracted from L4 larvae, and the expression levels of the relevant target genes (*cyp314a1* and *cyp315a1*) were detected in the larvae by RT-qPCR. The experiment was repeated three times.

2.9. RT-qPCR

For the reverse transcription quantitative polymerase chain reaction (PCR), 20–30 L3 larvae were collected and pooled from each treatment. Total RNA was isolated from the larvae using TRIzol reagent (Takara, Shiga, Japan), following the manufacturer's instructions [54]. Single-stranded cDNA was synthesised from the total RNA using oligo-dT primers (Shenggong, Shanghai, China). Real-time PCR was performed using a BioRad iCycler iQ Real-Time PCR Detection System (Bio-Rad, Hercules, CA, USA) and SYBR Green as a fluorescent dye. Primers targeting the *Aedes* RPS17 gene were used as internal controls [55]. The primers used to amplify the *cyp314a1* and *cyp315a1* target genes, as well as the primers used to amplify the genes involved in the clathrin-mediated endocytosis (CME) pathway, are listed in the Supplementary Table S1. The amplification rate (Ct) of each transcript was calculated using the PCR baseline subtraction method and iCycler software (version 2.3, Bio-Rad, Hercules, CA, USA) (constant fluorescence level). The cycle threshold (Ct) was determined in triplicate and the relative fold differences were calculated using the relative quantification method ($2^{-\Delta\Delta CT}$) [56]. Gene expression was determined relative to the endogenous control.

2.10. Simulated Field Trials

Simulated field trials were performed as described in the previous study [16]. An unused factory building on the outskirts of Haikou City was chosen for the experiment. The mosquitoes were reared in polyester mesh cages with a mesh size of 0.4 mm and a capacity of 1.5 m^3 [57]. Four 1000 L buckets for microalgae production were placed in the cages, which were fitted with LED lights. An aeration pump was used to ensure constant water circulation. The buckets were then filled with 10 L of cultivated microalgae. From 10

L to 200 L, sufficient light and aeration were maintained for the *Chlorella* culture. Jinniu Ling reservoir water was then added to bring the volume to 800 L, and the incubation process continued until the concentration of *Chlorella* cells reached 2×10^6–2×10^7 cells/mL. Around 1000 L1 larvae were placed in each bucket. A 10% sucrose solution was prepared for the male adults, while the egg-laying females were given chicken blood. The number of adult Aedes mosquitoes was counted weekly, in a similar manner to how the abundance of *Chlorella* in the water was determined. Blood and sugar water were discontinued one day prior to counting the mosquitoes to temporarily reduce their physical strength. Mosquito counting was conducted between 3:00 and 6:00 a.m. A dim LED flashlight was used to illuminate the cages during the procedure [16].

Water quality parameters were analysed in accordance with the Environmental Quality Standards for Surface Water (GB 3838-2002) [58]. Concentrations of ammonium nitrogen (NH_3-N), nitrate (NO_3^-), nitrite (NO_2^-) and chemical oxygen demand (COD) were quantified using a Hach DR3900 spectrophotometer (Hach Company, Loveland, CO, USA). Nitrogen (N) levels were determined through alkaline persulfate digestion followed by UV spectrophotometric detection, while phosphorus (P) concentrations were measured using the ammonium molybdate spectrophotometric method [59,60].

2.11. Sample Preparation and DNA Extraction of the Test Water

An 18S and 16S rDNA high-throughput sequencing analysis was performed on the water from simulated-field trials to detect the effect of recombinant RNAi *Chlorella* on eukaryotic and prokaryotic organisms in the test water. After feeding *Ae. albopictus* for 30 days, a 1 L water sample was collected from the bucket containing the mosquitoes and filtered through a 0.22 μm polycarbonate membrane (Millipore, Burlington, MA, USA) under a 30 kPa vacuum to collect the plankton. The membranes were stored at −80 °C until analysis. Genomic DNA was extracted using a modified Winnepenninckx et al. method [61].

2.12. Amplification and Sequencing of the Hypervariable Region

The V4 hypervariable region of the 18S rDNA and the V3-V4 region of the 16S rDNA were selected as targets for high-throughput sequencing analysis. The target regions were amplified using the primers listed in Supplementary Table S1 [62,63]. For the 18S rDNA sequencing, sequence libraries were constructed using the NEBNext Ultra DNA Library Prep Kit for Illumina (San Diego, CA, USA) in accordance with the manufacturer's protocol. The libraries were then sequenced on the Illumina HiSeq2500 platform. For 16S rDNA sequencing, the purified PCR products were sequenced on an Illumina MiSeq (Model MiSeq, Illumina, San Diego, CA, USA) platform. Data optimisation was conducted using the Chen and Jiang approach [64]. Low-quality sequences (i.e., those less than 200 base pairs in length) and sequencing primer adapters located at the 3′ ends of the primers were eliminated. After identifying chimeric sequences, high-quality sequences exhibiting 97% sequence identity were grouped into OTUs. The taxonomic categorisation of the OTUs was determined using USEARCH (version 5.2.236, Robert C. Edgar, San Francisco, CA, USA). Alpha diversity analysis was facilitated by calculating indices such as Shannon, Observed Species and Chao1 using QIIME software (version 1.9.1, University of Colorado, Boulder, CO, USA). Differences in community structure between samples or groups were analysed using principal coordinate analysis (PCoA) plots generated with R software (version 2.15.3, R Foundation for Statistical Computing, Vienna, Austria) [65,66].

2.13. Statistical Analyses

Data analysis was performed using the Statistical Package for the Social Sciences (SPSS, version 26.0, IBM, Chicago, IL, USA) and results are presented as the mean ± standard deviation (S.D.). Duncan's multiple range test was performed to examine significant differences between means. Values with a p-value of less than 0.05 were considered statistically significant. Asterisks indicate statistical significance in all cases: * $p < 0.05$; ** $p < 0.01$. Error bars show the standard deviation [67]. The 95% confidence interval (CI) was calculated using IBM SPSS Statistics 26 (version 26.0, IBM, Chicago, IL, USA). For PERMANOVA F-test, differences in microbial community structure among treatments were assessed using a permutational multivariate analysis of variance based on Bray–Curtis dissimilarity matrices. Grouping assignment (treatments) was used as the explanatory factor. The F-statistic was derived by comparing between-group variance to within-group variance, and significance was determined through 999 permutations of sample labels [67].

3. Results

3.1. CYP314A1 and CYP315A1 Proteins

A BlastP search was performed using *Ae. albopictus* CYP314A1 and CYP315A1 as source sequences to retrieve a large number of amino acid sequences. Several representative CYP314A1 and CYP315A1 orthologues from different species were selected for further analysis (Supplementary Tables S2 and S3), including characterisation of the structural domains of the nucleic acids and proteins (Supplementary Figures S1 and S2) and phylogenetic analyses. Analysis of the CYP314A1/CYP315A1 protein sequences using the online tool Interproscan (https://www.ebi.ac.uk/interpro/search/sequence/, accessed on 16 April 2025) revealed that all p450 domains were present. Results from protein sequence alignment showed that all p450 domains of CYP314A1 and CYP315A1 contained the following functional regions: WxxxR (Helix-C), GxE/DTT/S (Helix-I), ExxR (Helix-K), PxxFxPE/DRF (PERF), and PFxxGxRxCxG/A (heme-binding domain) (Supplementary Figures S3 and S4). Analysis of the CYP450 gene family in *Ae. albopictus* revealed that the species contains 81 genes distributed among clan 2 (nine genes), clan 3 (52 genes), and clan 4 (27 genes), as well as the MITO family (seven genes). The genes involved in ecdysone synthesis belonged to clan 2 (e.g., *cyp306a1*, *cyp307a1*, and *cyp18a1*) and the MITO family (e.g., *cyp302a1*, *cyp315a1*, and *cyp314a1*) (Supplementary Figure S5 and Table S4).

3.2. Cyp314a1 and cyp315a1 RNAi Recombinant Chlamydomonas/Chlorella Are Lethal to Ae. albopictus

The target genes selected for RNAi were *cyp314a1* and *cyp315a1* (AALFPA_063667 and AALFPA_079547), which are key enzymes in the *Ae. albopictus* 20E synthesis pathway. The *cyp314a1* and *cyp315a1* RNAi interference fragments were amplified using *Ae. albopictus* cDNA as a template. The amplified fragments were approximately 416 bp and 319 bp long, respectively. These fragments were then cloned forward and reverse into the pMaa7IR/XIR vector to create the recombinant RNAi vectors pMaa7IR/cyp314a1IR and pMaa7IR/cyp315a1IR. These RNAi vectors were subsequently transformed into *C. reinhardtii* CC425 and *C. vulgaris* HOC5. Several positive recombinant algal strains were identified through the implementation of PCR, and these were then employed in subsequent experiments (Supplementary Figure S6).

In the laboratory *Ae. albopictus* larvae feeding assay, larvae fed with the RNAi recombinant *C. reinhardtii* CYP314A(1-3) and CYP315A(1-3) began dying on the second day, with mortality rates of 93.3–100% (95% CI: 88.22–100%) and 96.7–100% (95% CI: 94.11–100%) on the 10th day, respectively. Conversely, no mortality was observed in the larvae fed *C. reinhardtii* CC425 and feed; only 16.7% and 30% of larvae died when fed water and the empty plasmid Maa7IR/XIR transgenic line, respectively. These results suggest that the oral administration of *cyp314a1* and *cyp315a1* RNAi recombinant *C. reinhardtii* is lethal to *Ae. albopictus* larvae (Figure 1).




Figure 1. Mortality of *Ae. albopictus* fed with *cyp314a1* and *cyp315a1* RNAi recombinant *Chlamydomonas*. (**A**) Mortality of *Ae. albopictus* fed with *cyp314a1* RNAi recombinant *Chlamydomonas*; (**B**) Mortality of *Ae. albopictus* fed with *cyp315a1* RNAi recombinant *Chlamydomonas*; (**C**) Mortality of *Ae. albopictus* fed with *cyp314a1* RNAi recombinant *Chlamydomonas* on days 3, 7 and 10 after feeding; (**D**) Mortality of *Ae. albopictus* fed with *cyp315a1* RNAi recombinant *Chlamydomonas* on days 3, 7 and 10 after feeding.Water: larvae fed with water; Feed: larvae fed with feed; CC425: larvae fed with *C. reinhardtii CC425*; Maa7: larvae fed with empty plasmid pMaa7IR/XIR transgenic *C. reinhardtii CC425*; CYP314A-1 to CYP314A-3: larvae fed with pMaa7IR/cyp314a1IR transgenic *C. reinhardtii CC425* lines. CYP315A-1 to CYP315A-3: larvae fed with pMaa7 IR/cyp315a1IR transgenic *C. reinhardtii CC425* lines. Each treated or control group contained ten *Aedes* larvae, and the experiments were performed in triplicate for 10 days. Data are expressed as the mean ± SD (n = 3), and significant differences ($p < 0.05$, Duncan's multiple range test) are indicated by different letters.

3.3. Effects of RNAi Recombinant Chlorella on Larval Development, and Redox Stress and 20E Compensation Assay

Approximately 300 L1 *Ae. albopictus* larvae in each treatment group were subjected to a 25-day feeding experiment. The larvae fed with the recombinant *C. reinhardtii* CC425-CYP314A1 and CC425-CYP315A1 began to die on the second day and 93.67% (95% CI: 91.19–96.15%) and 91.56% (95% CI: 88.89–94.22%) died within 25 days, respectively. In the control groups, 5.11% (95% CI: 2.72–7.51%), 0.22% (95% CI: 0–0.69%), 0.67% (95% CI: 0.67–0.67%) and 13.78% (95% CI: 11.67–15.85%) of the larvae died when fed water, feed, *C. reinhardtii* CC425 and transgenic *C. reinhardtii* pMaa7IR/XIR, respectively (Figure 2A). Similarly, the larvae fed recombinant *Chlorella* HOC5-CYP314A1 and HOC5-CYP315A1

began dying on the second day, with 90.44% (95% CI: 79.8–100%) and 83.11% (95% CI: 73.09–93.13%) dying within 25 days, respectively. By contrast, in the control groups, only 5.11% (95% CI: 2.72–7.51%), 0.22% (95% CI: 0–0.69%), 0.78% (95% CI: 0–1.74%) and 15.67% (95% CI: 12.35–18.89%) of larvae died when fed water, feed, *Chlorella* HOC5 and pMaa7IR/XIR transgenic *Chlorella*, respectively (Figure 2B).




Figure 2. Mortality, pupation, and eclosion rates of *Ae. albopictus* fed with *cyp314a1* and *cyp315a1* RNAi recombinant *Chlamydomonas* (**A**,**B**) and *Chlorella* (**C**,**D**). Water: larvae fed with water; Feed: larvae fed with feed; CC425: larvae fed with *C.reinhardtii CC425*; HOC5: larvae fed with wild *C. vulgaris* HOC5; CC425-Maa7 and HOC5-Maa7: larvae fed with an empty Maa7IR/XIR plasmid in transgenic *Chlorella or Chlamydomonas*; CC425-CYP314A1 and CC425-CYP315A1: larvae fed with *cyp314a1* and *cyp315a1* RNAi recombinant *Chlamydomonas* and *Chlorella* lines. HOC5-CYP314A1 and HOC5-CYP315A1: larvae fed with *cyp314a1* and *cyp315a1* RNAi recombinant *Chlorella* lines. Each treated or control group contained 300 *Aedes* larvae, and the experiments were performed in triplicate over 25 days. Data are expressed as the mean ± SD (n = 3), and significant differences ($p < 0.05$, Duncan's multiple range test) are indicated by different letters.

Larvae fed the control diet began pupating on day 4, and all had completed pupation by day 12. Those fed *C. reinhardtii* CC425 began pupating on day 5. By day 11, 99.33% (95% CI: 99.33–99.33%) of the larvae had pupated. Larvae fed pMaa7IR/XIR transgenic *C. reinhardtii* CC425 began pupating on day 7, with 86.22% (95% CI: 84.13–88.31%) having pupated by day 16. Larvae fed on recombinant *Chlamydomonas* CC425-CYP314A1 and CC425-CYP315A1 began to pupate on day 6. By day 14, 6.33% (95% CI: 3.85–8.81%) and 8.44% (95% CI: 5.78–11.11%) of the larvae had pupated, respectively (Figure 2A). Similarly, the larvae fed the recombinant *Chlorella* HOC5-CYP314A1 and HOC5-CYP315A1 began to pupate on day 7. By day 25, 8.22% (95% CI: 0.19–16.62%) and 15.44% (95% CI: 8.54–22.34%) of the larvae had pupated, respectively. In the control groups, 0%, 99.78% (95% CI: 99.31–100%), 99.22% (95% CI: 98.26–100%) and 84.00% (95% CI: 80.21–87.8%) of the larvae pupated when fed water, feed, *Chlorella* HOC5 and pMaa7IR/XIR transgenic *Chlorella*, respectively (Figure 2B).

The emergence rates of adult mosquitoes showed that those fed feed began to emerge on day 5, with 98.11% of pupae emerging within 25 days. Those fed *C. reinhardtii* CC425 began to emerge as adults on day 6, with 97.78% (95% CI: 97.31–98.25%) of pupae ultimately emerging. However, Aedes pupae emerged as adults at lower rates of 2.89% (95% CI: 0–6.34%) and 5.11% (95% CI: 0.12–10.1%) when fed recombinant *Chlamydomonas* CC425-CYP314A1 and CC425-CYP315A1 strains, respectively (Figure 2A). Similarly, only 2.89% (95% CI: 1.94–3.84%) and 7.11% (95% CI: 4.21–10.01%) of *Aedes* pupae emerged as adults when fed the recombinant *Chlorella* HOC5-CYP314A1 and HOC5-CYP315A1 strains, respectively. In the control groups, adult mosquito emergence was observed to be 0%, 98.11% (95% CI: 96.38–99.84%), 96.89% (95% CI: 95.16–98.62%) and 82.56% (95% CI: 80.16–84.95%) when the larvae were fed water, feed, *Chlorella* HOC5 or pMaa7IR/XIR transgenic *Chlorella*, respectively (Figure 2B).

The *Ae. albopictus* fed with feed had the longest L3 larval body length (5.56 mm/5.89 mm), followed by mosquitoes fed with *C. reinhardtii* CC425 or *C. vulgaris* HOC5, with body lengths of 4.63 mm and 4.35 mm, respectively. The body length of mosquitoes fed with water was the shortest (1.84 mm), whereas the body length of the mosquitoes fed with recombinant lines was significantly lower ($p < 0.05$) than that of the control group (fed with feed, *C. reinhardtii* CC425 or *C. vulgaris* HOC5) (Figure 3A–D). In terms of body width, the *Ae. albopictus* fed with feed had the widest L3 larval body width (0.80 mm), followed by mosquitoes fed with *C. reinhardtii* CC425 or *C. vulgaris* HOC5, with body widths of 0.59 mm and 0.57 mm, respectively. The body width of mosquitoes fed with water was the narrowest (0.21 mm), whereas the body width of other mosquitoes fed with recombinant strains was significantly lower ($p < 0.05$) than that of the control group (fed with feed, *C. reinhardtii* CC425 or *C. vulgaris* HOC5) (Figure 3A–D). A notable observation was that *Ae. albopictus* fed recombinant *Chlorella* exhibited a reduction in body length and width when compared with *Ae. albopictus* fed recombinant *C. reinhardtii*. (Figure 3A–D).

Activity assays of superoxide dismutase (SOD), peroxiredoxin (POD) and catalase (CAT) in *Ae. albopictus* larvae revealed significantly higher enzyme activity in larvae fed *cyp314a1* and *cyp315a1* RNAi recombinant *Chlamydomonas* or *Chlorella* than in the control group (fed *C. reinhardtii* CC425, or *C. vulgaris*) (Figure 4).

Results from the 20E compensation assay revealed that *Ae. albopictus* fed on recombinant *Chlamydomonas* supplemented with 20E exhibited a substantial reduction in larval mortality, as well as a notable increase in pupation and emergence rates (Figure 5A). Expression analysis of *cyp314a1* and *cyp315a1* revealed that mRNA levels of both target genes were significantly reduced ($p < 0.05$) in *Ae. albopictus* fed either *cyp314a1* or *cyp315a1* RNAi recombinant *Chlamydomonas*, regardless of whether 20E was supplemented (Figure 5B).




Figure 3. Length and width of larvae. The length and width of L3 larvae from each treatment were measured. (**A**,**C**), larvae fed with recombinant *Chlamydomonas* strains; (**B**,**D**), larvae fed with recombinant *Chlorella* strains. (**C**,**D**), 1, larvae fed water; 2, larvae fed feed; 3, larvae fed *C. reinhardtii* CC425 (**C**) or *C. vulgaris* HOC5 (**D**); 4, larvae fed with empty plasmid Maa7IR/XIR transgenic *Chlamydomonas* (**C**) or *Chlorella* (**D**) strain; 5 and 6, larvae fed with recombinant *Chlamydomonas* (**C**) or *Chlorella* (**D**) strain. Water: water was fed to the larvae; Feed: larvae fed feed; CC425: larvae fed *C. reinhardtii* CC425; HOC5: larvae fed wild *Chlorella vulgaris* HOC5; CC425-Maa7 and HOC5-Maa7: larvae fed with empty plasmid Maa7IR/XIR transgenic *Chlorella* or *Chlamydomonas* strains; CC425-CYP314A1 and CC425-CYP315A1: larvae fed with recombinant *Chlamydomonas* strains. HOC5-CYP314A1 and HOC5-CYP315A1: larvae fed with recombinant *Chlorella* strains. Data are expressed as the mean ± SD (n = 3), and significant differences ($p < 0.05$, Duncan's multiple range test) are indicated by different letters.




Figure 4. Analysis of CAT, POD and SOD enzyme activities in *Aedes* larvae after feeding on recombinant *Chlamydomonas* (**A**) and *Chlorella* (**B**) strains. CC425: larvae fed with *C. reinhardtii* CC425; HOC5: larvae fed with *C. vulgaris* HOC5; CC425-CYP314A1 and CC425-CYP315A1: larvae fed with recombinant *Chlamydomonas* strains; HOC5-CYP314A1 and HOC5-CYP315A1: larvae fed with recombinant *Chlorella* strains. Data are expressed as the mean ± SD (n = 3), and significant differences ($p < 0.05$, Duncan's multiple range test) are indicated by different letters.




Figure 5. 20E compensation test. (**A**) Mortality, pupation, and eclosion rates of *Ae. albopictus* fed with *cyp314a1* and *cyp315a1* RNAi recombinant *Chlamydomonas* with or without 20E; (**B**) The relative *cyp314a1* and *cyp315a1* mRNA levels in L3 larvae fed with recombinant *Chlamydomonas* with or without 20E. CC425: larvae fed *C. reinhardtii* CC425; CC425+20E: larvae fed with C. *reinhardtii* CC425 and 20E; CC425-CYP314A1 and CC425-CYP315A1: larvae fed with recombinant *Chlamydomonas* strains. CC425-CYP314A1+20E and CC425-CYP315A1+20E: larvae fed with the recombinant *Chlamydomonas* strains and 20E. Data are expressed as the mean ±SD (n = 3), and significant differences ($p < 0.05$, Duncan's multiple range test) are indicated by different letters.

3.4. The Expression of the RNAi Target Genes and the Genes in the Clathrin-Mediated Endocytosis (CME) Pathway

Real-time PCR was used to examine *cyp314a1* and *cyp315a1* gene expression in *Ae. albopictus* larvae fed with recombinant *Chlamydomonas* and *Chlorella* strains. Larvae fed with *C. reinhardtii* CC425 or *C. vulgaris* HOC5 were used as controls. The expression levels of the *cyp314a1* and *cyp315a1* in larvae fed the recombinant *Chlamydomonas* or *Chlorella* strains were significantly lower ($p < 0.05$) than in the control larvae (Figure 6A,B). The

most substantial decline in gene expression was observed in larvae fed the *cyp314a1* RNAi recombinant *Chlamydomonas* or *Chlorella* strains (82.5% and 80.6% decreases, respectively). These observations collectively indicate that the *cyp314a1* and *cyp315a1* genes in *Ae. albopictus* were effectively silenced by the recombinant *Chlamydomonas* or *Chlorella* strains (Figure 6A,B).




Figure 6. The mRNA levels of genes in *Ae. albopictus* L3 larvae fed with recombinant *Chlamydomonas*/*Chlorella* dsRNA/shRNA. The relative *cyp314a1* and *cyp315a1* mRNA levels in L3 larvae fed with recombinant *Chlamydomonas* (**A**) or *Chlorella* (**B**). The relative *flot-1*, *flot2*, *Rab7*, *iqf*, *Chc*, *iqfR*, *AP50*, *Arf72*, *vhaSFD*, *Dynamin* and *vha16* mRNA levels in *Ae. albopictus* L3 larvae fed with *cyp314a1* dsRNA/shRNA recombinant *Chlorella* (**C**). The relative *flot-1*, *flot2*, *Rab7*, *iqf*, *Chc*, *iqfR*, *AP50*, *Arf72*, *vhaSFD*, *Dynamin* and *vha16* mRNA levels in *Ae. albopictus* L3 larvae fed with *cyp315a1* dsRNA/shRNA recombinant *Chlorella* (**D**). CC425: larvae fed with *C.reinhardtii* CC425; HOC5: larvae fed with wild *C. vulgaris* HOC5; CC425-CYP314A1 and CC425-CYP315A1: larvae fed with recombinant *Chlamydomonas* strains; HOC5-CYP314A1 and HOC5-CYP315A1: larvae fed with recombinant *Chlorella* strains. HOC5-CYP314A1(dsRNA/shRNA) and HOC5-CYP315A1(dsRNA/shRNA): larvae fed with recombinant *Chlorella* containing dsRNA/shRNA. Expression of CME-related genes: Clathrin adapter protein 50 (*AP50*), Clathrin heavy chain (*Chc*), Liquid facets (*lqf*), Liquid facets related protein (*lqfR*), *Dynamin*, Vacuolar H + ATPase 16 kDa (*vha16*), V-type proton ATPase subunit H (*VhaSFD*), Ras-like GTPase (*Rab7*), ADP-ribosylation factor 72 (*Arf72*), Flotillin-1 (*Flot-1*) and Flotillin-2 (*Flot-2*). Results are expressed as the mean ± SE of three independent replicates. For (**A**–**D**), significant differences ($p < 0.05$, Duncan's multiple range test) are shown by different letters.

Furthermore, an evaluation was conducted of the levels of several gene transcripts involved in the clathrin-mediated endocytosis (CME) pathway (Supplementary Table S5). These include coat assembly proteins (clathrin adapter AP50 and clathrin heavy chain [Chc]), vesicle formation and budding proteins (liquid facets [lqf], liquid facets-related [lqfR], and dynamin), early endosome proteins (Vha16 and VhaSFD), endosome maturation protein Rab7 and ADP-ribosylation factor-like 1 (Arf72A). Two genes of the flotillin-mediated endocytosis pathway are also included *flot-1* and *flot-2*. The results demonstrated that the mRNA levels of *flot-2*, *Rab7*, *Chc*, *Dynamin* and *vha16* were significantly increased in larvae fed with *cyp314a1* shRNA or recombinant *Chlorella* dsRNA (Figure 6C). Furthermore, the mRNA levels of *flot2*, *Rab7*, *Chc*, *lqfR*, *Dynamin* and *vha16* were also found to be significantly increased in larvae fed with *cyp315a1* shRNA or recombinant *Chlorella* dsRNA (Figure 6D). These results suggested that knockdown of *cyp314a1* or *cyp315a1* influences the expression of CME pathway-related genes more than feeding with ds/shRNA-free *Chlorella* does.

3.5. Simulated-Field Trials

Large-scale simulated field tests were conducted over a period of 16 weeks. The results demonstrated that the *Ae. albopictus* population fed on Jinniu Ling reservoir water (JNL) increased from 1000 to a maximum of 5063 individuals over 10 weeks. After this, the population slowly decreased to around 4497 individuals after 16 weeks. *Ae. albopictus* raised in Jinniu Ling reservoir water supplemented with wild-type *C. vulgaris* HOC5 (JNL-HOC5) increased to a maximum of 6895 individuals after 10 weeks and then decreased slowly to around 6574 individuals after 16 weeks. In the treatment groups where the water was supplemented with recombinant *Chlorella* HOC5-CYP314A1 or HOC5-CYP315A1, the quantity of *Ae. albopictus* increased in the first three weeks and then decreased from 1100 (JNL-CYP314A1) or 1090 (JNL-CYP315A1) to zero after 13–15 weeks (Figure 7A–C). These results demonstrate the ability of recombinant *Chlorella* HOC5-CYP314A1 and HOC5-CYP315A1 to suppress the *Ae. albopictus* population under simulated field conditions.

To understand the proliferation of RNAi recombinant *Chlorella* in natural water bodies and its effect on water nutrients, a series of water quality tests and *Chlorella* density measurements were conducted on each group of water samples. Following a rapid logarithmic growth phase (days 5–7), the recombinant algal strains entered a stabilisation period (days 8–11), after which they entered a period of reduced growth (days 12–90). The removal rates of nitrogen (N), phosphorus (P), nitrate, nitrite, ammonia and chemical oxygen consumption (COD) from the water were found to be positively correlated with the density of *Chlorella* (Figure 7D). On day 30, the removal rates of N, P, nitrate, nitrite, ammonia, and COD by *C. vulgaris* HOC5, RNAi recombinant *Chlorella* HOC5-CYP314A1 and HOC5-CYP315A1 were 86.96–90.51%, 85.39–91.38%, 91.97–93.03%, 80.88–90.54%, 86.79–89.25% and 92.28–93.32%, respectively (Figure 7E). These values corresponded to water samples containing nitrogen, phosphorus, nitrate, nitrite, ammonia, and COD at concentrations of 2.96–3.95 mg/L, 0.07–0.13 mg/L, 2.14–2.44 mg/L, 0.007–0.013 mg/L, 0.03–0.04 mg/L and 31.08–35.56 mg/L, respectively (Figure 7F). By contrast, the removal rates of N, P, nitrate, nitrite, ammonia, and COD in the control Jinniu Ling Reservoir water samples were only 3.9%, 8.89%, 0.04%, 2.41%, 11.66% and 2.73%, respectively. These levels corresponded to concentrations of 37.71 mg/L, 0.81 mg/L, 32.96 mg/L, 0.08 mg/L, 0.31 mg/L and 496.65 mg/L, respectively. By adding RNAi recombinant *Chlorella*, the levels of N, P, nitrate, nitrite, ammonia and COD in the severely eutrophic Jinniu Ling Reservoir water were reduced to Class V levels ($N \leq 2.0$, $P \leq 0.4$, nitrate ≤ 10, nitrite ≤ 1, ammonia ≤ 2.0, and COD ≤ 40), in accordance with China's environmental quality standard for surface water (GB 3838-2002) (Figure 7E,F) [58].




Figure 7. Simulated-field trials in the suburban of Haikou. Test water was taken from Jinniu Ling reservoir (**A**). 200 L of *C. vulgaris* HOC5 were mixed with 600 L of the Jinniu Ling reservoir water in the barrel. Approximately 1000 L1 larvae were placed in each cage, and the number of adult *Ae. albopictus* was counted once a week (**B**). The population and survival rates of *Ae. albopictus* in the JNL

(control), JNL-HOC5 (control), JNL-CYP314A1 and JNL-CYP315A1 treatment groups are shown in (**C**). Relative abundance of *Chlorella* in the JNL-HOC5 (control), JNL-CYP314A1 and JNL-CYP315A1 treatment groups (**D**). The removal rates of nitrogen, phosphorus, nitrate, nitrite, ammonia, and COD by RNAi recombinant *Chlorella* (**E**). Nitrogen, phosphorus, nitrate, nitrite, ammonia, and COD content in the JNL-HOC5 (control), JNL-CYP314A1 and JNL-CYP315A1 treatment groups (**F**). JNL: The mosquito lived in water from the Jinniu Ling reservoir alone. JNL-HOC5: The mosquito lived in water from the Jinniu Ling reservoir that was supplemented with wild *Chlorella* HOC5. JNL-CYP314A1 and JNL-CYP315A1: the mosquito lived in water from the Jinniu Ling reservoir that was supplemented with recombinant *Chlorella* HOC5-CYP314A1 and HOC5-CYP315A1.

3.6. Plankton Abundance and Diversity in Simulated Field Test Waters

To evaluate the impact of RNAi recombinant *Chlorella* on aquatic biological communities, high-throughput analyses of 18S rDNA and 16S rDNA was performed on water samples containing the recombinant *Chlorella* strains HOC5-314A1 and HOC5-315A1. For the 18S rDNA sequencing, 988,356 qualified tags were obtained from the samples after several quality controls steps. After removing unclassified and unique tags from the dataset, the total number of high-quality tags was 818,788. These sequences were then clustered into OTUs with 97% concordance and a total of 1009 OTUs were obtained. Similarly, 1,125,735 qualified tags were obtained from the 16S rDNA sequencing after several quality controls. The total number of high-quality tags was 761,131 and 1213 OTUs were obtained.

To assess and compare planktonic biodiversity across samples, we computed richness and diversity indices, including Chao1, observed species and Shannon indices, for phytoplankton, zooplankton, and prokaryote communities. These indices were based on OTUs clustered at 97% sequence similarity, followed by α-diversity analysis (Figure 8). Compared with the control group (JNL group), the phytoplankton Chao1 index and observed species index in the JNL-CYP314A1 treatment group (supplemented with RNAi recombinant *Chlorella* strain HOC5-CYP314A) showed no significant differences. The zooplankton Chao1 index and observed species index exhibited an upward trend but did not reach statistical significance. In contrast, the zooplankton Chao1 index and observed species index in the JNL-CYP315A1 treatment group (supplemented with recombinant strain HOC5-CYP315A1) decreased significantly ($p < 0.05$) compared with the JNL group (Figure 8B). With respect to prokaryotic communities, the JNL-CYP315A1 group displayed a significant increase ($p < 0.05$) in the observed species index relative to the JNL control group (Figure 8C).

Shannon index analysis revealed that, compared with the JNL group, the phytoplankton community diversity in the JNL-CYP314A1 treatment group decreased significantly ($p < 0.05$), whereas zooplankton diversity increased markedly (Figure 8A,B). In addition, both phytoplankton and prokaryotic diversity in the JNL-CYP315A1 group declined significantly ($p < 0.05$) compared with the JNL group (Figure 8A,C). These findings indicate that the supplementation of recombinant *Chlorella* altered the patterns of aquatic biodiversity in the Jinniu Ling Reservoir.

The NMDS and PCA analyses of the eukaryotic communities revealed that JNL (pink points) was clearly separated from the other groups, indicating that the baseline water community structure differed substantially from that of the treatments with added *Chlorella*. JNL-HOC5 (green points) clustered closely with JNL-CYP314A1 (purple points) and JNL-CYP315A1 (blue points), suggesting that the community structure became more similar after the addition of *Chlorella*. However, differences still remained between the RNAi-transgenic *Chlorella* treatments (JNL-CYP314A1, JNL-CYP315A1) and the wild-type HOC5. The PCA results showed that the first principal component (Axis1) explained 75.19% of the variance, indicating that the major community differences were determined by the addition of *Chlorella* and its genotype (Figure 9A,B).




Figure 8. Box plots showing the differences between groups in the Chao1 index, the observed species index and the Shannon index for phytoplankton (**A**), zooplankton (**B**) and prokaryotes (**C**). JNL: water samples from Jinniu Ling Reservoir; JNL-HOC5: water samples from Jinniu Ling Reservoir with wild *Chlorella* HOC5 added; JNL-CYP314A1: water samples from Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP314A1 added; JNL-CYP315A1: water samples from Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP315A1 added. Data are expressed as the mean ± SD (n = 3), and significant differences ($p < 0.05$, Duncan's multiple range test) are indicated by different letters.

The NMDS and PCA analyses of the prokaryotic communities demonstrated that JNL was distinctly separated from the other groups, indicating that the addition of *Chlorella* had a pronounced effect on the prokaryotic community. JNL-HOC5 overlapped to some extent with JNL-CYP314A1 and JNL-CYP315A1, but the RNAi-treated groups (JNL-CYP314A1, JNL-CYP315A1) displayed a more dispersed distribution, suggesting that the effects of recombinant *Chlorella* on prokaryotic communities were more complex and variable. The PCA results showed that the Axis1 explained 36.3% of the variance and the Axis2 explained 28.15%, indicating that multiple factors contributed to the differences in the prokaryotic community rather than a single dominant factor (Figure 9C,D).




Figure 9. NMDS2 and PCA analyses of eukaryote (**A**,**B**) and prokaryote (**C**,**D**) JNL: water samples from Jinniu Ling Reservoir; JNL-HOC5: water samples from Jinniu Ling Reservoir with wild *Chlorella* HOC5 added; JNL-CYP314A1: water samples from Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP314A1 added; JNL-CYP315A1: water samples from Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP315A1 added.

3.7. Changes in Plankton in Simulated Field Test Waters

A total of ten groups of eukaryotic microalgae were identified at the phylum level in the Jinniu Ling Reservoir (JNL, control) sample, including *Chlorophyta*, *Chrysophyta*, *Cryptophyta*, *Bacillariophyta*, *Pyrrophyta*, *Streptophyta*, *Ochrophyta*, *Phragmoplastophyta*, *Glaucophyta* and *Haptophyta* (Supplementary Table S6). Of these, 366 OTUs within 83 genera were identified in JNL. *Chlorophyta* exhibited the highest number of OTUs. A total of 120 OTUs were identified within 28 genera of *Chlorophyta*. The second largest class was *Chrysophyta*, comprising 72 OTUs within 15 genera. *Cryptophyta*, *Bacillariophyta* and *Pyrrophyta* also exhibited a relatively high number of OTUs. By contrast, *Phragmoplastophyta*, *Glaucophyta* and *Haptophyta* exhibited lower numbers of OTUs (Supplementary Table S6). In the Jinniu Ling Reservoir water plus *Chlorella* HOC5 group (JNL-HOC5), 356 OTUs belonging to 86 genera were observed. *Chlorophyta* exhibited the highest number of OTUs, with 126 across 27 genera. As shown in Supplementary Table S6, *Chrysophyta* was the second largest phylum, comprising 63 OTUs across 17 genera. In the JNL-CYP314A1 treatment

group, 383 OTUs belongs to 92 genera were observed. This was a greater number than in the JNL or JNL-HOC5 control groups. Of these, the number of *Cryptophyta*, *Bacillariophyta* and *Streptophyta* species increased. However, the JNL-CYP315A1 treatment group exhibited a mere 334 OTUs across 80 genera, falling short of the OTUs and genera observed in the control groups (JNL, or JNL-HOC5). Regarding species abundance, *Chlorophyta*, *Bacillariophyta*, *Pyrrophyta*, *Chrysophyta* and *Cryptophyta* were found to be more prevalent in the treatment groups (JNL-CYP314A1 and JNL-CYP315A1) and in the control group (JNL, JNL-HOC5). Compared with the control group, the relative abundance of *Bacillariophyta*, *Pyrrophyta* and *Chrysophyta* decreased in the treatment group, while the relative abundance of *Chlorophyta* increased (Figure 10A).

Analysis at the Phytoplankton genus level showed that adding recombinant *Chlorella* increased its abundance in the JNL-CYP314A1 and JNL-CYP315A1 treatment groups to 75.80% and 82.41%, respectively, compared to the control JNL group at 12.25%. The abundance of most phytoplankton was decreased in the JNL-CYP314A1 and JNL-CYP315A1 treatment groups, except for *Chlorella*, *Cryptomonas* and *Teleaulax*, whose abundance increased (Figures 10B and 11A). For example, the abundance of *Aulacoseira*, *Cyclotella*, *Fragilaria*, *Skeletonema*, *Thalassiosira*, *Discostella* and *Nitzschia* in the *Bacillariophyta* decreased. The abundance of *Mychonastes*, *Tetraedron*, *Coelastrella*, *Monoraphidium*, *Desmodesmus*, *Scenedesmus* and *Golenkinia* in the *Chlorophyta* also decreased. *Rhodomonas* in the *Cryptophyta*, *Ceratium* in the *Dinophyta*, and *Dinobryon*, *Ochromonas* and *Chromulina* in the *Chrysophyta*, also decreased (Figures 10B and 11A).

Analysis of zooplankton at the phylum level revealed that *Ciliophora*, *Cercozoa*, *Arthropoda* and *Rotifera* occupied the top positions in terms of species number and abundance (Supplementary Table S7 and Figure 10C). Specifically, *Ciliophora* exhibited abundances of 87.40%, 88.01%, 76.35% and 92.03% in JNL, JNL-HOC5, JNL-CYP314A1 and JNL-CYP315A1 water, respectively. Analysis of the normalised zooplankton genus-level data showed that the genera *Tintinnidium*, *Entodinium* and *Dileptus* occupied the top three positions in terms of species abundance. *Tintinnidium* was the most prevalent species in JNL, JNL-HOC5, JNL-CYP314A1 and JNL-CYP315A1 water, with abundances of 61.56%, 45.87%, 38.28% and 49.99%, respectively (Figure 10C).

Analysis of differences in species abundance revealed higher abundances of *Cilicophora* (*Histriculus*, *Cyclidium*, *Polyplastron*, *Histiobalantium* and *Vorticella*) and *Protozoa* (*Heteromita*, *Reclinomonas* and *Bodomorpha*) in treatment groups JNL-CYP314A1 and JNL-CYP315A1 than in the control group, JNL. In contrast, the *Cilicophora* (*Pseudouroleptus*, *Trochilia*, *Bryometopus*, *Cryptocaryon*, *Halteria*, *Didinium*, *Dileptus*, *Paracercomonas* and *Epispathidium*), and rumen-specialised groups of ruminants (*Entodinium*, *Ophryoscolex* and *Dasytricha*), as well as other protists such as *Colpodella*, *Cytolphosis*, *Arcuospathidium* and *Tintinnidium* (which consume diatoms and dinoflagellates), were reduced in abundance (Figures 10C,D and 11B).




Figure 10. Relative abundance of phytoplankton (**A**,**B**), zooplankton (**C**,**D**) and prokaryotes (**E**,**F**) at the phylum (**A**,**C**,**E**) and genus (**B**,**D**,**F**) levels in simulated field test waters. JNL: water samples from the Jinniu Ling Reservoir; JNL-HOC5: water samples from the Jinniu Ling Reservoir with wild *Chlorella* HOC5 added; JNL-CYP314A1: water samples from the Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP314A1 added; JNL-CYP315A1: water samples from the Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP315A1 added. *Chlorella* increased its abundance in the JNL-CYP314A1 and JNL-CYP315A1 treatment groups to 75.80% and 82.41%, respectively, compared to the control JNL group at 12.25% (**B**). *Ciliophora* exhibited abundances of 87.40%, 88.01%, 76.35% and 92.03% in JNL, JNL-HOC5, JNL-CYP314A1 and JNL-CYP315A1 water, respectively (**C**). *Tintinnidium* was the most prevalent species in JNL, JNL-HOC5, JNL-CYP314A1 and JNL-CYP315A1 water, with abundances of 61.56%, 45.87%, 38.28% and 49.99%, respectively (**D**). *Cyanobium_PCC-6307* was the most prevalent species in the JNL, JNL-HOC5, JNL-CYP314A1 and JNL-CYP315A1 water samples, with concentrations of 25.57%, 10.68%, 8.65% and 8.19%, respectively (**F**).




Figure 11. Heat map analysis of phytoplankton at genus level (**A**), zooplankton at genus level (**B**), prokaryotes at phylum level (**C**), and prokaryotic at genus level (**D**) in simulated field test water. JNL: water samples from the Jinniu Ling Reservoir; JNL-HOC5: water samples from the Jinniu Ling Reservoir with wild *Chlorella* HOC5 added; JNL-CYP314A1: water samples from the Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP314A1 added; JNL-CYP315A1: water samples from the Jinniu Ling Reservoir with RNAi recombinant *Chlorella* HOC5-CYP315A1 added.

3.8. Changes in Prokaryotes in Simulated Field Test Waters

Analysis of the water samples using 16S rDNA high-throughput sequencing revealed the presence of high concentrations of *Cyanobacteria*, *Proteobacteria*, *Firmicutes*, *Bacteroidota*, *Actinobacteriota* and *Actinobacteria* in the control (JNL and JNL-HOC5) and treatment (JNL-CYP314A1 and JNL-CYP315A1) groups (Figure 10E). *Cyanobacteria* were the most prevalent of these, with proportions of 47.57%, 39.96%, 37.78% and 30.10% in JNL, JNL-HOC5, JNL-CYP314A1 and JNL-CYP315A1, respectively. At the genus level, the top three positions in terms of abundance were occupied by *Cyanobium_PCC-6307*, *Paenibacillus*, and *CL500-29_marine_group*. As shown in Figure 10F, *Cyanobium_PCC-6307* was the most prevalent species in the JNL, JNL-HOC5, JNL-CYP314A1 and JNL-CYP315A1 water samples, with concentrations of 25.57%, 10.68%, 8.65% and 8.19%, respectively.

Cyanobium_PCC-6307 belongs to the family *Cyanobiaceae*, a group of photosynthetic, autotrophic prokaryotes that are widely distributed in freshwater, marine and wetland environments. It has a thylakoid structure typical of cyanobacteria and fixes carbon through photosynthesis, while also being involved in nitrogen metabolism. It is one of the dominant bacterial genera in natural or artificial waters and is an indicator species of eutrophication [68–71]. The abundance of *Cyanobium_PCC-6307* in the JNL-CYP314A1 and JNL-CYP315A1 waters decreased as the N and P contents decreased. *Paenibacillus* is a genus of Gram-positive bacteria widely distributed in the natural environment. It belongs to the phylum *Firmicutes*, the class Bacilli, and the family *Paenibacillaceae*. The genus currently comprises over 150 species, around 20 of which have the ability to fix nitrogen, secrete organic acids to dissolve insoluble phosphorus in water, degrade organic pollutants and reduce the eutrophication of water [72–75]. The results showed that the *Paenibacillus* abundance in treated JNL-CYP314A1 and JNL-CYP315A1 water increased significantly (19.92% and 21.54%, respectively) compared to the control (JNL, 6.79%) (Figure 10F). The *CL500-29_marine_group* belongs to the phylum *Actinobacteriota*, which contains the dominant bacteria in freshwater and offshore environments. The *CL500-29_marine_group* is an important phylum of *Actinobacteria* with significant ecological functions. It can efficiently utilise a variety of carbon sources, such as amino acids, carboxylic acids and lipids. Its abundance is positively correlated with lake carbon concentration and nitrogen and phosphorus removal efficiency, suggesting that it is closely related to water purification capacity [76–80]. The sequencing results obtained in this study align with the following observations: the introduction of recombinant *Chlorella* into the treatment group let to lower concentrations of carbon, nitrogen and phosphorus in the water, as well as a decrease in the abundance of the CL500-29_marine_group.

Heat map-based analysis of differences in the abundance of prokaryotic species at the phylum level after normalisation revealed that the abundances of *Gemmatimonadota*, *Acidobacteriota*, *Elusimicrobiota*, *Armatimonadota*, *Crenarchaeota*, *Entotheonellaeota*, *Kryptonia*, *Chloroflexi*, *Methylomirabilota*, *Latescibacteria*, *Nitrospirota*, *Myxococcota* and *Latescibacterota* were higher in the treatment groups (JNL-CYP314A1 and JNL-CYP315A1) than in the control group (JNL). Conversely, the abundances of *Cyanobacteria*, *Actinobacteria*, *Halobacterota*, *Kapabacteria*, *Fusobacteriota*, *Gemmatimonadetes*, *Actinobacteriota*, *Bdellovibrionota*, *Fibrobacterota*, *Planctomycetota*, *Verrucomicrobiota*, and *Deinococcota* were markedly lower in the treatment groups than in the control group (Figures 10E and 11C). At the genus level, the analysis revealed that the abundances of *Vogesella*, *Limnohabitans*, *Pseudarcicella*, *Aquabacterium*, *Rheinheimera*, *Escherichia-Shigella*, *Providencia*, *Gaiella*, *Haliangium*, *Pseudomonas*, *Candidatus*, *Methylopumilus*, and *Paenibacillus* were higher in the treatment groups (JNL-CYP314A1 and JNL-CYP315A1) than in the control group (JNL). By contrast, the abundance of the following bacteria decreased in the treatment groups compared to the control group: *Mycobacterium*, *Flavobacterium*, *Anaeromyxobacter*, *Sphaerotilus*,

Fluviicola, *Cetobacterium*, *Cyanobium_PCC-6307*, *Polynucleobacter*, *Pseudohongiella*, *Sediminibacterium*, *Acidibacter*, *Dinghuibacter*, *Roseomonas*, *Rubrivivax*. CL500-29_marine_group, *Limnobacter* and *Chthoniobacter* were lower in the treatment groups than in the control group (Figures 10F and 11D).

4. Discussion

Ae. albopictus is recognised as a vector of significant public health vector, primarily transmitting viruses such as the Dengue, Yellow fever, Chikungunya and Zika viruses. These diseases cause substantial human mortality each year [81]. In the absence of effective therapies or vaccines, control the vectors remains the most effective strategy for preventing the spread of these pathogens. Extensive research has demonstrated that RNAi-based biopesticides represent a new generation of pesticides and are a powerful tool for controlling insect populations. Compared to other strategies, RNAi biopesticides are more environmentally friendly. However, they also face challenges related to target-site resistance, as evidenced by the emergence of resistance issues following the widespread adoption of transgenic insect-resistant cotton [82–85]. While the high efficiency and specificity of RNAi biopesticides have garnered significant attention, the environmental impacts of their large-scale deployment and application have not yet been comprehensively evaluated [86,87].

In this study, the *cyp314a1* and *cyp315a1* genes of *Ae. albopictus* were selected as RNAi targets. CYP450 is an ancient and multifunctional superfamily of monooxygenases [22]. Insects possess four CYP450 clans with roles in metabolic detoxification, growth regulation and development [25,26]. CYP450 enzymes are involved in ecdysone biosynthesis. Specifically, shadow (*sad*, CYP315A1) and shade (*shd*, CYP314A1) catalyse the final steps of cholesterol conversion into the biologically active 20-hydroxyecdysone (20E) [88–90]. As they are critical to ecdysone synthesis, the *cyp314a1* and *cyp315a1* genes are predominantly found in arthropods, suggesting they have minimal impact on humans, higher animals and plants. Our analysis of *Ae. albopictus* CYP450 genes revealed that the species harbours 81 such genes, distributed across Clan 2 (9 genes), Clan 3 (52 genes), Clan 4 (27 genes) and MITO (7 genes). Genes involved in ecdysone synthesis belong to Clan 2 (e.g., *cyp306a1*, *cyp307a1* and *cyp18a1*) and the MITO family (e.g., *cyp302a1*, *cyp314a1* and *cyp315a1*) (Supplementary Figure S5).

Observations demonstrated that *Ae. albopictus* larvae fed with *cyp314a1* or *cyp315a1* recombinant RNAi *Chlamydomonas* exhibited mortality rates of 93.3–100% or 96.7–100%, respectively, within a 10-day period (Figure 1). In a feeding trial involving 300 mosquitoes, larvae fed *cyp314a1* or *cyp315a1* recombinant RNAi *Chlamydomonas* showed mortality rates of 93.67% and 91.56% over 25 days, respectively. Those fed *cyp314a1* or *cyp315a1* recombinant RNAi *Chlorella* exhibited mortality rates of 90.44% and 83.11%, respectively (Figure 2). These results confirm that the oral administration of *cyp314a1* or *cyp315a1* RNAi *Chlamydomonas/Chlorella* has a lethal effect on *Ae. albopictus* larvae. Pupation rates were 6.33% and 8.44% for larvae fed with *cyp314a1* or *cyp315a1* RNAi recombinant *Chlamydomonas*, respectively, and 8.22% and 15.44% for larvae fed with *cyp314a1* or *cyp315a1* RNAi recombinant *Chlorella*, respectively (Figure 2). With regard to adult emergence, the emergence rates of larvae fed with *cyp314a1* or *cyp315a1* RNAi recombinant *Chlamydomonas* were 2.89% and 5.11%, respectively, while the emergence rates of those fed with *cyp314a1* or *cyp315a1* RNAi recombinant *Chlorella* were 2.89% and 7.11%, respectively (Figure 2). These findings suggest that recombinant algal strains expressing *Cyp314a1* or *Cyp315a1* RNAi significantly disrupt the pupation and adult emergence processes of *Ae. albopictus*.

Using *Ae. albopictus* larvae as a model, we observed that feeding them *cyp314a1* or *cyp315a1* RNAi recombinant algal strains significantly increased the enzymatic activities

of SOD, POD and CAT compared to control groups (fed *C. reinhardtii* CC425 or *C. vulgaris* HOC5), indicating an enhanced redox stress (Figure 4). The 20E rescue assay revealed that supplementing *cyp314a1* or *cyp315a1* RNAi recombinant *Chlamydomonas* with exogenous 20E substantially reduced larval mortality and restored pupation and adult emergence rates (Figure 5A). 20E regulates processes such as moulting, metamorphosis, and tissue apoptosis, and is the key hormone required for maintaining the transition from larva to adult. Deficiency or reduction of 20E leads to developmental arrest and moulting failure in larvae. Redox stress induces cellular damage, mitochondrial disfunction, and energy metabolism disorders. The combined effects of these two factors ultimately result in larval death.

Studies have demonstrated that the dsRNA and shRNA of *cyp314a1* or *cyp315a1* likely penetrate the interior of mosquitoes via the clathrin-mediated endocytosis (CME) pathway in intestinal epithelial cells. In contrast, invertebrates such as nematodes use the SID pathway to transport dsRNA into and between cells [91]. However, insects primarily absorb exogenous dsRNA through CME [92,93]. Following the knockout of the *cyp314a1* or *cyp315a1* gene, the mRNA levels of certain key CME pathway genes increased (see Figure 6C,D), corroborating the findings of Abbasi et al. (2020) regarding *Ae. aegypti* [94].

Subsequent simulated-field trials revealed that RNAi-recombinant *Chlorella* reduced the *Ae. albopictus* population from 1000 to zero within 16 weeks (see Figure 7A–C). It also reduced the levels of nitrogen, phosphorus, ammonia, nitrates, nitrites and COD in the experimental water (Jinniu Ling Reservoir), upgrading it to China's Class V surface water standard (GB 3838-2002) (see Figure 7E,F) [58].

To understand the effect of RNAi recombinant *Chlorella* on the test water's biological populations, 18S and 16S rDNA high-throughput sequencing analyses were performed. The results showed that the rapid reproduction of the recombinant *Chlorella* caused a bloom that consumed nutrients such as C, N and P at a rapid rate. Competition for nutrient salts directly inhibits the growth of other algae, such as diatoms (e.g., *Aulacoseira*, *Cyclotella* and *Thalassiosira)* and green algae (e.g., *Scenedesmus* and *Coelastrella*) [95–97]. Competition for light resources results in decreased light transmittance (increased turbidity), leading to the decline of photosynthesis-dependent non-planktonic algae (e.g., the attached diatom *Fragilaria* and the green algae *Desmodesmus*, *Tetraedron* and *Monoraphidium*) due to insufficient light [68]. However, motile species such as *Cryptomonas* and *Teleaulax* can acquire light energy through vertical migration, giving them a competitive advantage (see Figures 10A,B and 11A) [98].

During the algal bloom triggered by recombinant RNAi in *Chlorella*, the water body's dissolved oxygen (DO) levels exhibited significant diurnal fluctuations: photosynthesis resulted in DO supersaturation during the day, whereas respiration caused hypoxic stress at night. These extreme redox oscillations periodically stressed aerobic zooplankton (*Didinium* and *Halteria*) physiologically, driving the dynamic remodelling of the microbial community structure. The drastic decrease in DO at night significantly inhibited the proliferation of the aerobic groups *Cyanobium PCC-6307*, *Limnobacter*, *Chthoniobacter* and *Actinobacteria*. Meanwhile, the abundance of facultative and specialised anaerobic microorganisms such as *Methylomirabilota* and *Nitrospirota* increased. The biomass of the strict anaerobe *Chloroflexi* and *Elusimicrobiota* increased significantly compared to the control due to the enhanced anaerobic microenvironment in the bottom sediment (Figure 11C) [99–101]. DOM and POM, produced by the decomposition of *Chlorella* residues during the decline of the algal bloom, provided preferential carbon sources for heterotrophic flora, including *Pseudarcicella*, *Aquabacterium*, *Vogesella*, *Rheinheimera*, *Pseudomonas*, *Paenibacillus* and *Haliangium*, as well as the *Myxococcota*. The release of nitrogen and phosphorus derived from the algae, activates the metabolic activities of the nitrifying bacterium *Crenarchaeota*, the denitrifying bacterium

Nitrospirota and the sulfur-oxidising bacterium *Gemmatimonadota*. Conversely, *Cyanobacteria*, *Planctomycetes* and *Halobacteria* decreased in abundance due to inferior competition for carbon and nitrogen (Figures 10E,F and 11C,D) [102–106].

Analysis based on alpha diversity data showed that adding the wild type of *C. vulgaris* HOC5 slightly altered the biotope but did not significantly reduce diversity (similar Chao1 and Shannon indices) or impact community stability overall. However, the alpha diversity (e.g., observed species and Shannon index) of the JNL-CYP315A treatment group differed significantly from that of the control JNL group. This indicates that the RNAi recombinant algal strain significantly altered the structure of the biological community, resulting in fewer zooplankton species and decreased diversity (see Figure 8). The significant difference in observed species and Shannon index between the JNL-CYP314A1 and JNL-CYP315A1 treatment groups suggests that the difference in biological impact on the surrounding environment may be due to variations between the CYP314A1 and CYP315A1 RNAi recombinant algal strains. Analysis of the dsRNA target fragments revealed multiple differential fragments between the two strains. BLAST comparison of fragments belonging to the CYP315A1 dsRNA target region in the NCBI database showed that multiple fragments were homologous ($\geq$80%) with some of the less abundant zooplankton species. This may explain the reduction in zooplankton species in the JNL-CYP315A treatment group (Figure 8B and Supplementary Table S8).

In conclusion, changes in environmental water bodies are mainly due to the release of RNAi recombinant *Chlorella* and community reconfiguration due to resource competition (e.g., nitrogen–phosphorus competition between *Chlorella* and diatoms), which is induced by algal blooms. Other factors include anaerobic bacterial proliferation and aerobic bacterial decline caused by anaerobic conditions.

While this study has made some progress, the impact of exogenous DNA fragments (particularly antibiotic resistance genes) carried by RNAi recombinant *Chlorella* on environmental organisms (including those in the direct and indirect food chains) requires further evaluation. Lake algal blooms generally go through a cycle of 'recovery-emergence-decay-dormancy' [107]. During the bloom stage, the productivity of phytoplankton increases significantly [108], whereas during the decay and decomposition stage, the apoptosis of algal cells releases large quantities of sugars, proteins, and lignin into the water. These substances then become an important source of DOM in the water [109]. However, this study only examined the biotope composition and structure during the RNAi recombinant *Chlorella* bloom, and did not consider what happened after the bloom ended. Future studies will explore this to clarify the environmental impact of RNAi recombinant *Chlorella* algal blooms. Current evidence suggests that the environmental impact of such blooms is limited and similar to that of green algal blooms in urban eutrophic lakes. As the bloom enters the decay and dormancy stage, the biological populations in the water are restructured as nutrients are depleted, which increases species diversity and restores the population balance.

5. Conclusions

In this study, we selected the *cyp314a1* and *cyp315a1*, two genes involved in the ecdysone biosynthesis pathway of *Ae. albopictus*, as targets for RNA interference (RNAi). The recombinant *Chlamydomonas*/*Chlorella* mosquito biocide obtained through genetic transformation exhibited high lethality towards *Ae. albopictus* larvae. Simulated field trials of the recombinant *Chlorella* biocide effectively controlled the *Ae. albopictus* population. The recombinant *Chlorella* induced a bloom, altering the population structure of the water and inhibiting the growth of most phytoplankton through trophic competition. The anaerobic environment created by the bloom subsequently triggered the proliferation of anaerobic

bacteria and the decline of aerobic bacteria. This study is an important addition to research on the use of recombinant microalgae as mosquito biocides.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/insects16101033/s1. Table S1: Primers used in this study; Table S2: Origin, taxonomy and GenBank accession number of the CYP314A1 orthologues used in the present study; Table S3: Origin, taxonomy and GenBank accession number of the CYP315A1 orthologues used in this study; Table S4: *Ae. albopictus* CYP450 information; Table S5: *Ae. albopictus* endocytic genes and related sequences; Table S6: The number of genera and OTUs in different groups of phytoplankton at the phylum level; Table S7: The number of genera and OTUs in different groups of zooplankton at the phylum level; Table S8: Comparison of sequence homology between *Ae. albopictus cyp315a1* dsRNA region and zooplankton target genes; Figure S1: Phylogenetic relationship of CYP314A1 and analysis of their conserved domains; Figure S2: Phylogenetic relationship of CYP315A1 and analysis of their conserved domains; Figure S3: The alignment of the amino acid sequences of CYP314A1 in *Ae. albopictus* and its orthologues in the *Arthropoda*; Figure S4: The alignment of the amino acid sequences of CYP315A1 in *Ae. albopictus* and its orthologues in the *Arthropoda*; Figure S5: Phylogeny of *Ae. albopictus* cytochrome P450; Figure S6: PCR electrophoresis results of recombinant plasmid pMaa7IR/cyp314a1IR(A) and pMaa7IR/cyp315a1IR (B) transgenic algal lines; Figure S7: Sequence of *cyp314a1* and *cyp315a1* shRNA and predicted shRNA secondary structure.

Author Contributions: X.D. and X.F. conceived and designed the study. C.H. and X.D. performed the experiments. X.D., C.X. and J.L. analysed data. D.X., X.D. and X.F. prepared all the figures. X.F. wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by the Key Research and Development Projects of Hainan Province (ZDYF2022SHFZ314), the National Natural Science Foundation of China (82260669), Hainan Medical University Academic Enhancement Support Program Project (XSTS2025151), Hainan Medical University Science and Innovation Program (KC200574), and Financial Fund of the Ministry of Agriculture and Rural Affairs, P. R. China (NFZX2024).

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information Files.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Waggoner, J.J.; Gresh, L.; Vargas, M.J. Viremia and Clinical Presentation in Nicaraguan Patients Infected with Zika Virus, Chikungunya Virus, and Dengue Virus. *Clin. Infect. Dis.* **2016**, *63*, 1584–1590. [CrossRef]
2. Bhatt, S. The global distribution and burden of dengue. *Nature* **2013**, *496*, 504–507. [CrossRef]
3. WHO. *Ten Threats to Global Health in 2019*; WHO: Geneva, Switzerland, 2019.
4. Xiang, B.; Gao, P.; Kang, Y.F. Importation of Zika virus in China: A significant risk in southern China. *J. Infect.* **2017**, *74*, 328–330. [CrossRef] [PubMed]
5. Chen, J.; Lu, H.Z. Yellow fever in China is still an imported disease. *Biosci. Trends* **2016**, *10*, 158–162. [CrossRef] [PubMed]
6. Liu, W.; Sun, F.J.; Tong, Y.G. Rift Valley fever virus imported into China from Angola. *Lancet Infect. Dis.* **2016**, *16*, 1226. [CrossRef] [PubMed]
7. Wu, D.; Wu, J.; Zhang, Q.L. Chikungunya outbreak in Guangdong province, China, 2010. *Emerg. Infect. Dis.* **2012**, *18*, 493–495. [CrossRef]
8. Hemingway, J. Averting a malaria disaster: Will insecticide resistance derail malaria control? *Lancet* **2016**, *387*, 1785–1788. [CrossRef]
9. Van, H. Global trends in the production and use of DDT for control of malaria and other vector-borne diseases. *Malar. J.* **2012**, *11*, 401.
10. Bian, G.; Zhou, G.; Lu, P.; Xi, Z. Replacing a native Wolbachia with a novel strain results in an increase in endosymbiont load and resistance to dengue virus in a mosquito vector. *PLoS Negl. Trop. Dis.* **2013**, *7*, e2250. [CrossRef]
11. Xue, L.; Fang, X.; Hyman, J.M. Comparing the effectiveness of different strains of Wolbachia for controlling chikungunya, dengue fever, and zika. *PLoS Negl. Trop. Dis.* **2018**, *12*, e0006666. [CrossRef]

12. Flores, H.A.; Neill, S.L. Controlling vector-borne diseases by releasing modified mosquitoes. *Nat. Rev. Microbiol.* **2018**, *16*, 508–518. [CrossRef] [PubMed]
13. Bargielowski, I.; Alphey, L.; Koella, J.C. Cost of mating and insemination capacity of a genetically modified mosquito *Aedes aegypti* OX513A compared to its wild type counterpart. *PLoS ONE* **2011**, *6*, e26086. [CrossRef] [PubMed]
14. Curtis, Z.; Matzen, K.; Neira, O.M. Assessment of the Impact of Potential Tetracycline Exposure on the Phenotype of *Aedes aegypti* OX513A: Implications for Field Use. *PLoS Negl. Trop. Dis.* **2015**, *9*, e0003999. [CrossRef] [PubMed]
15. Fei, X.; Zhang, Y.; Ding, L.; Xiao, S.; Xie, X.; Li, Y.; Deng, X. Development of an RNAi-based microalgal larvicide for the control of *Aedes aegypti*. *Parasit. Vectors* **2021**, *14*, 387. [CrossRef]
16. Fei, X.; Huang, X.; Li, Z.; Li, X.; He, C.; Xiao, S.; Li, Y.; Zhang, X.; Deng, X. Effect of marker-free transgenic Chlamydomonas on the control of Aedes mosquito population and on plankton. *Parasit. Vectors* **2023**, *16*, 18. [CrossRef]
17. Fei, X.; Xiao, S.; Huang, X.; Li, Z.; Li, X.; He, C.; Li, Y.; Zhang, X.; Deng, X. Control of Aedes mosquito populations using recombinant microalgae expressing short hairpin RNAs and their effect on plankton. *PLoS Negl. Trop. Dis.* **2023**, *17*, e0011109. [CrossRef]
18. Li, X.; Zhang, H. Endocytic pathway mediates refractoriness of insect *Bactrocera dorsalis* to RNA interference. *Sci. Rep.* **2015**, *5*, 8700. [CrossRef]
19. Xu, L.; Xu, S.; Sun, L. Synergistic action of the gut microbiota in environmental RNA interference in a leaf beetle. *Microbiome* **2021**, *9*, 98. [CrossRef]
20. Zhang, J.; Khan, S.A.; Hasse, C. Full crop protection from an insect pest by expression of long double-stranded RNAs in plastids. *Science* **2015**, *347*, 991–994. [CrossRef]
21. Xu, W.; Zhang, M.; Li, Y. Complete protection from *Henosepilachna vigintioctopunctata* by expressing long double-stranded RNAs in potato plastid. *J. Integr. Plant Biol.* **2023**, *65*, 1003–1011. [CrossRef]
22. Nelson, D.R.; Goldstone, J.V.; Stegeman, J.J. The cytochrome P450 genesis locus: The origin and evolution of animal cytochrome P450s. *Philos. Trans. R. Soc. B Biol. Sci.* **2013**, *368*, 20120474. [CrossRef]
23. Ventura, T.; Bose, U.; Fitzgibbon, Q.P.; Smith, G.G.; Shaw, P.N.; Cummins, S.F.; Elizur, A. *CYP*450s analysis across spiny lobster metamorphosis identifies a long sought missing link in crustacean development. *J. Steroid Biochem. Mol. Biol.* **2017**, *171*, 262–269. [CrossRef]
24. Dermauw, W.; Van, L.T.; Feyereisen, R. Diversity and evolution of the P450 family in arthropods. *Insect Biochem. Mol. Biol.* **2020**, *127*, 103490. [CrossRef] [PubMed]
25. Feyereisen, R. Evolution of insect P450. *Biochem. Soc. Trans.* **2006**, *34*, 1252–1255. [CrossRef]
26. Feyereisen, R. Arthropod CYPomes illustrate the tempo and mode in P450 evolution. *Biochim. Biophys. Acta* **2011**, *1814*, 19–28. [CrossRef]
27. Zhang, H.; Gao, H.; Lin, X.; Yang, B.; Wang, J.; Yuan, X.; Zhang, Z.; He, T.; Liu, Z. Akt-FoxO signaling drives co-adaptation to insecticide and host plant stresses in an herbivorous insect. *J. Adv. Res.* **2025**, *75*, 53–64. [CrossRef]
28. Mao, Y.; Cai, W.; Wang, J.; Hong, G.; Tao, X.; Wang, L. Silencing a cotton bollworm P450 monooxygenase gene by plant-mediated RNAi impairs larval tolerance of gossypol. *Nat. Biotechnol.* **2007**, *25*, 1307–1313. [CrossRef] [PubMed]
29. Ma, K.; Li, F.; Tang, Q.; Liang, P.; Liu, Y.; Zhang, B. CYP4CJ1-mediated gossypol and tannic acid tolerance in *Aphis gossypii* Glover. *Chemosphere* **2019**, *219*, 961–970. [CrossRef] [PubMed]
30. Bass, C.; Hebsgaard, M.; Hughes, J. Genomic resources for the brown planthopper, *Nilaparvata lugens*: Transcriptome pyrosequencing and microarray design. *Insect Sci.* **2011**, *19*, 1–12. [CrossRef]
31. Bass, C.; Carvalho, R.; Oliphant, L.; Puinean, A.; Field, L.; Nauen, R. Overexpression of a cytochrome P450 monooxygenase, CYP6ER1, is associated with resistance to imidacloprid in the brown planthopper, *Nilaparvata lugens*. *Insect Mol. Biol.* **2011**, *20*, 763–773. [CrossRef]
32. Mao, K.; Zhang, X.; Ali, E.; Liao, X.; Jin, R.; Ren, Z. Characterization of nitenpyram resistance in *Nilaparvata lugens* (Stal). *Pestic. Biochem. Physiol.* **2019**, *157*, 26–32. [CrossRef]
33. Matsuda, K.; Ihara, M.; Sattelle, D. Neonicotinoid insecticides: Molecular targets, resistance, and toxicity. *Annu. Rev. Pharmacol. Toxicol.* **2020**, *60*, 241–255. [CrossRef] [PubMed]
34. Sun, X.; Gong, Y.; Ali, S.; Hou, M. Mechanisms of resistance to thiamethoxam and dinotefuran compared to imidacloprid in the brown planthopper: Roles of cytochrome P450 monooxygenase and a P450 gene CYP6ER1. *Pestic. Biochem. Physiol.* **2018**, *150*, 17–26. [CrossRef] [PubMed]
35. Zhang, X.; Liao, X.; Mao, P.; Yang, D.; Li, E.; Alia, E. The role of detoxifying enzymes in field-evolved resistance to nitenpyram in the brown planthopper *Nilaparvata lugens* in China. *Crop Prot.* **2017**, *94*, 106–114. [CrossRef]
36. Tan, Q.M.; Chen, W.W.; Li, H.H.; Liao, S.C.; Yi, G.Q.; Mei, Y.; Luo, J.; Tan, H.H.; Li, X.S. Adipokinetic hormone signaling regulates cytochrome P450-mediated chlorantraniliprole sensitivity in *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Pest. Manag. Sci.* **2022**, *78*, 2618–2628. [CrossRef]

37. Ullah, F.; Gul, H.; Tariq, K.; Desneux, N.; Gao, X.; Song, D. Functional analysis of cytochrome P450 genes linked with acetamiprid resistance in melon aphid, *Aphis gossypii*. *Pestic. Biochem. Physiol.* **2020**, *170*, 104687. [CrossRef]
38. Chen, W.; Li, Z.; Zhou, C.; Ali, A.; Ali, S.; Wu, J. RNA interference in cytochrome P450 monooxygenase (CYP) gene results in reduced insecticide resistance in *Megalurothrips usitatus* Bagnall. *Front. Physiol.* **2023**, *14*, 1130389. [CrossRef]
39. Wang, X.P.; Ye, M.Z.; Tu, W.T.; Zhao, X.F. 20-hydroxyecdysone via upregulating 4EBP expression and inhibiting its phosphorylation represses cell proliferation to permit insect larval molting. *Insect Sci.* **2025**, *in press*. [CrossRef]
40. Yuan, D.; Zhou, S.; Liu, S.; Li, K.; Zhao, H.; Long, S.; Liu, H.; Xie, Y.; Su, Y.; Yu, F.; et al. The AMPK-PP2A axis in insect fat body is activated by 20-hydroxyecdysone to antagonize insulin/IGF signaling and restrict growth rate. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 9292–9301. [CrossRef]
41. Guo, S.; Tian, Z.; Wu, Q.W.; King-Jones, K.; Liu, W.; Zhu, F.; Wang, X.P. Steroid hormone ecdysone deficiency stimulates preparation for photoperiodic reproductive diapause. *PLoS Genet.* **2021**, *17*, e1009352. [CrossRef]
42. Nian, X.; Wang, B.; Holford, P.; Beattie, G.; Tan, S.; Yuan, W.; Cen, Y.; He, Y.; Zhang, S. Neuropeptide Ecdysis-Triggering Hormone and Its Receptor Mediate the Fecundity Improvement of '*Candidatus* Liberibacter Asiaticus'-Infected *Diaphorina citri* Females and CLas Proliferation. *Adv. Sci.* **2025**, *12*, e2412384. [CrossRef]
43. Moulos, P.; Alexandratos, A.; Nellas, I.; Dedos, S.G. Refining a steroidogenic model: An analysis of RNA-seq datasets from insect prothoracic glands. *BMC Genom.* **2018**, *19*, 537. [CrossRef]
44. Liu, S.; He, C.; Liang, J.; Su, Q.; Hua, D.; Wang, S.; Wu, Q.; Xie, W.; Zhang, Y. Molecular characterization and functional analysis of the Halloween genes and CYP18A1 in *Bemisia tabaci* MED. *Pestic. Biochem. Physiol.* **2020**, *167*, 104602. [CrossRef]
45. Rohr, J.; Sarkar, N.; Balenger, S.; Jeong, B.R.; Cerutti, H. Tandem inverted repeat system for selection of effective transgenic RNAi strains in Chlamydomonas. *Plant J.* **2004**, *40*, 611–621. [CrossRef]
46. Wang, Y.L.; Xu, Z.D.; Jiang, S.J.; Huang, K.Y. Screening of *Chlorella* cell wall-deficient mutants and establishment of transformation system. *Acta Hydrobiol. Sin.* **2016**, *40*, 370–377.
47. Liu, J.; He, X.C.; Chen, X.; Ma, B.; Wei, H.T.; Yang, M.F. An efficient method for electrotransformation of Chlamydomonas reinhardtii. *J. Beijing Univ. Agric.* **2019**, *34*, 9–13.
48. Bradford, M.M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **1976**, *72*, 248–254. [CrossRef] [PubMed]
49. Ukeda, H. Novel spectrophotometric assay for superoxide dismutase (SOD) activity using water-soluble tetrazolium salts. *Biosci. Biotechnol. Biochem.* **2002**, *66*, 941–943.
50. Chance, B.; Maehly, A.C. Assay of catalases and peroxidases. *Methods Enzymol.* **1955**, *2*, 764–775.
51. Beers, R.F.; Sizer, I.W. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J. Biol. Chem.* **1952**, *195*, 133–140. [CrossRef]
52. Guo, S. 20-Hydroxyecdysone regulates larval-pupal transition via insulin-like signaling in the mosquito *Aedes aegypti*. *Insect Biochem. Mol. Biol.* **2020**, *127*, 103477.
53. Zhang, Q. Ecdysone signaling mediates the trade-off between immunity and reproduction in *Aedes albopictus*. *PLoS Pathog.* **2021**, *17*, e1009632.
54. Chomczynski, P.; Sacchi, N. The single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction: Twenty-something years on. *Nat. Protoc.* **2006**, *1*, 581–585. [CrossRef]
55. Dzaki, N. Evaluation of reference genes at different developmental stages for quantitative real-time PCR in *Aedes aegypti*. *Sci. Rep.* **2017**, *7*, 43618. [CrossRef] [PubMed]
56. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative PCR and the $2-\Delta\Delta CT$ method. *Methods* **2001**, *25*, 402–408. [CrossRef]
57. Mysore, K.; Li, P.; Wang, C.W.; Hapairai, L.K.; Scheel, N.D.; Realey, J.S.; Sun, L. Characterization of a yeast interfering RNA larvicide with a target site conserved in the synaptotagmin gene of multiple disease vector mosquitoes. *PLoS Negl. Trop. Dis.* **2019**, *13*, e0007422. [CrossRef]
58. *GB 3838-2002*; Environmental Quality Standards for Surface Water. Ministry of Ecology and Environment of China; Environmental Science Press: Beijing, China, 2002.
59. American Public Health Association (APHA). *Standard Methods for the Examination of Water and Wastewater*, 23rd ed.; APHA: Washington, DC, USA, 2017.
60. Liu, Y. Comparative analysis of water quality testing methods between GB and ISO standards. *Environ. Monit. China* **2015**, *31*, 45–52.
61. Winnepenninckx, B.; Backeljau, T.; Wachter, R. Extraction of high molecular weight DNA from mollusks. *Trends Genet.* **1993**, *9*, 407. [PubMed]
62. Cheung, M.K.; Au, C.H.; Chu, K.H.; Kwan, H.S.; Wong, C.K. Composition and genetic diversity of picoeukaryotes in subtropical coastal waters as revealed by 454 pyrosequencing. *ISME J.* **2010**, *4*, 1053–1059. [CrossRef]

63. Zimmermann, J.; Jahn, R.; Gemeinholzer, B. Barcoding diatoms: Evaluation of the V4 subregion on the 18S rRNA gene, including new primers and protocols. *Org. Divers. Evol.* **2011**, *11*, 173–192. [CrossRef]
64. Chen, H.; Jiang, W. Application of high-throughput sequencing in understanding human oral microbiome related with health and disease. *Front. Microbiol.* **2014**, *5*, 508. [CrossRef]
65. Yuan, X.L.; Cao, M.; Liu, X.M.; Du, Y.M.; Shen, G.M.; Zhang, Z.F.; Li, J.H.; Zhang, P. Composition and genetic diversity of the *Nicotiana tabacum* microbiome in different topographic areas and growth periods. *Int. J. Mol. Sci.* **2018**, *19*, 3421. [CrossRef]
66. Lozupone, C.; Hamady, M.; Knight, R. Unifrac-an online tool for comparing microbial community diversity in a phylogenetic context. *BMC Bioinform.* **2006**, *7*, 371. [CrossRef]
67. Altman, D.G.; Bland, J.M. Standard deviations and standard errors. *Br. Med. J.* **2005**, *331*, 903. [CrossRef]
68. Komárek, J.; Kaštovský, J.; Jezberová, J. Taxonomic classification of cyanoprokaryotes (cyanobacterial genera) using a polyphasic approach. *Preslia* **2014**, *86*, 295–335.
69. Callieri, C. Picocyanobacteria in freshwater ecosystems: Biogeographic patterns and community assembly mechanisms. *J. Limnol.* **2012**, *71*, 1–16.
70. Paerl, H.W.; Otten, T.G. Harmful cyanobacterial blooms: Causes, consequences, and controls. *Microb. Ecol.* **2013**, *65*, 995–1010. [CrossRef]
71. Wu, Y. Dynamics of cyanobacterial communities and their environmental drivers in constructed wetlands. *Environ. Sci. Technol.* **2019**, *53*, 8854–8862.
72. Ash, C. Molecular identification of rRNA group 3 bacilli (Ash, Farrow, Wallbanks and Collins) using a PCR probe test. Proposal for the creation of a new genus Paenibacillus. *Antonie Van Leeuwenhoek* **1993**, *64*, 253–260. [CrossRef]
73. Grady, E.N. Current knowledge and perspectives of Paenibacillus: A review. *Microb. Cell Factories* **2016**, *15*, 203. [CrossRef]
74. Huang, X. Biodegradation of phenanthrene by Paenibacillus sp. PRNK-6: Pathway characterization and biosurfactant production. *Chemosphere* **2019**, *231*, 529–538.
75. Zhao, Y. Paenibacillus strains as bioagents for eutrophication control: Nutrient removal and microbial community modulation. *Environ. Res.* **2021**, *200*, 111751.
76. Newton, R.J. A guide to the natural history of freshwater lake bacteria. *Microbiol. Mol. Biol. Rev.* **2011**, *75*, 14–49. [CrossRef]
77. Salcher, M.M. In situ substrate preferences of abundant bacterioplankton populations in a prealpine freshwater lake. *ISME J.* **2011**, *5*, 896–907. [CrossRef] [PubMed]
78. Henson, M.W. Expanding the diversity of bacterioplankton isolates and modeling seasonal niche preferences of freshwater Actinobacteria. *mSystems* **2018**, *3*, e00227-18.
79. Ghai, R. Metagenomics uncovers a new group of low GC and ultra-small marine Actinobacteria. *Sci. Rep.* **2014**, *4*, 2471. [CrossRef]
80. Liao, J. Ecological drivers of the microbial communities in interconnected freshwater and marine ecosystems. *Microbiome* **2021**, *9*, 217.
81. Paupy, C. *Aedes albopictus*, an arbovirus vector: From the darkness to the light. *Microbes Infect.* **2009**, *11*, 1177–1185. [CrossRef]
82. Zhu, K.Y.; Palli, S.R. Mechanisms, applications, and challenges of insect RNA interference. *Annu. Rev. Entomol.* **2020**, *65*, 293–311. [CrossRef]
83. Tabashnik, B.E. Insect resistance to Bt crops: Lessons from the first billion acres. *Nat. Biotechnol.* **2013**, *31*, 510–521. [CrossRef] [PubMed]
84. Marcombe, S. Insecticide resistance in *Aedes albopictus*: Current status and mechanisms. *PLoS Neglected Trop. Dis.* **2019**, *13*, e0007615.
85. Bachman, P.M. Ecological risk assessment for RNA interference-based plant-protective products: An overview of challenges and proposed science-based considerations. *Regul. Toxicol. Pharmacol.* **2016**, *81*, 108–120.
86. Joga, M.R. RNA interference-based insecticidal crops: Potential effects on non-target species. *Bioscience* **2016**, *66*, 657–665.
87. Benelli, G.; Beier, J.C. Current vector control challenges in the fight against mosquito-borne diseases. *Acta Trop.* **2017**, *174*, 91–96. [CrossRef]
88. Gilbert, L.I. The regulation of ecdysteroid synthesis in insects: A multi-gene family approach. *Insect Biochem. Mol. Biol.* **2004**, *34*, 1245–1252.
89. Rewitz, K.F. The insect neuropeptide PTTH activates receptor tyrosine kinase Torso to initiate metamorphosis. *Science* **2007**, *316*, 894–897. [CrossRef]
90. Yoshiyama, T. Neverland is a conserved cholesterol-binding protein required for ecdysone synthesis in insects. *Dev. Biol.* **2006**, *298*, 555–570.
91. Feinberg, E.H. Transport of dsRNA into cells by the transmembrane protein SID-1. *Science* **2003**, *301*, 1545–1547. [CrossRef]
92. Xiao, D. Clathrin-dependent endocytosis plays a predominant role in cellular uptake of double-stranded RNA in the red flour beetle. *Insect Biochem. Molec.* **2015**, *60*, 68–77. [CrossRef]
93. Cappelle, K. The involvement of clathrin-mediated endocytosis and two Sid-1-like transmembrane proteins in double-stranded RNA uptake in the Colorado potato beetle midgut. *Insect Mol. Biol.* **2016**, *25*, 315–323. [CrossRef]

94. Abbasi, R. A novel paperclip double-stranded RNA structure demonstrates clathrin-independent uptake in the mosquito *Aedes aegypti*. *Insect Biochem. Mol. Biol.* **2020**, *127*, 103492. [CrossRef]
95. Smith, V.H.; Crews, T. Ecological impacts of genetically modified microalgae: A review. *Appl. Phycol.* **2014**, *26*, 1–14.
96. Litchman, E.; Klausmeier, C.A. Trait-based community ecology of phytoplankton. *Annu. Rev. Ecol. Evol. Syst.* **2008**, *39*, 615–639. [CrossRef]
97. Grover, J.P. Phytoplankton competition and coexistence: Intrinsic ecosystem dynamics and human impacts. *Hydrobiologia* **2012**, *698*, 5–25.
98. Jones, R.I.; Gowen, R.J. Vertical distribution and temporal variation of Cryptophyceae in a stratified lake. *J. Phycol.* **1990**, *26*, 472–478.
99. Buchan, A. Master recyclers: Features and functions of bacteria associated with phytoplankton blooms. *Nat. Rev. Microbiol.* **2014**, *12*, 686–698. [CrossRef]
100. Ettwig, K.F. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* **2010**, *464*, 543–548. [CrossRef]
101. Tang, X. Metagenomic insights into the microbial community and nutrient cycling in the subsurface layer of a stratified eutrophic reservoir. *Microb. Ecol.* **2018**, *76*, 125–138.101.
102. Kraft, B. Oxygen and nitrogen production by an ammonia-oxidizing archaeon. *Science* **2022**, *375*, 97–100. [CrossRef] [PubMed]
103. Liu, J. Gemmatimonadota as a key player in organic sulfur cycling in freshwater sediments. *ISME Commun.* **2022**, *2*, 1–11.
104. Shi, L. Phytoplankton-derived organic matter shapes microbial community structure and function in eutrophic lakes. *Environ. Microbiol.* **2020**, *22*, 5163–5178.
105. Peacock, J.P. Resource partitioning drives microbial niche differentiation in a eutrophic lake. *mSystems* **2021**, *6*, e01222-20.
106. Mu, D.S. Genomic insights into the ecological role of Myxococcota in degrading algal-derived polysaccharides. *Front. Microbiol.* **2020**, *11*, 591738.
107. Zhu, J.; Zou, N.; Zhong, H. Spatiotemporal characteristics and its environmental application of dissolved organic matter (DOM) in sediments from Chaohu Lake, a eutrophic lake. *Acta Sci. Circumstantiae* **2020**, *40*, 2528–2538.
108. Knoll, L.B.; Vanni, M.J.; Renwick, W.H. Temperate reservoirs are large carbon sinks and small CO_2 sources: Results from high-resolution carbon budget. *Glob. Biogeochem. Cycles* **2013**, *27*, 52–64. [CrossRef]
109. Villacorte, L.O.; Ekowati, Y.; Neu, T.R. Characterisation of algal organic matter produced by bloom-forming marine and freshwater algae. *Water Res.* **2015**, *73*, 216–230. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Elimination of Ultraviolet Light-Mediated Attraction Behavior in *Culex* Mosquitoes via dsRNA-Mediated Knockdown of Opsins

Xinyi Liu †, Guoqiang Zhao †, Hui Liu, Yuxuan Mao, Meng Xu, Jing Wu, Lijiao Li, Zongzhao Zhai * and Pa Wu *

Hunan Provincial Key Laboratory of Animal Intestinal Function and Regulation, College of Life Sciences, Hunan Normal University, Changsha 410081, China; xinyiliu599@sina.com (X.L.); zhaogq@hunnu.edu.cn (G.Z.); 0807lh@sina.com (H.L.); myx2025@sina.com (Y.M.); xumengaa@sina.com (M.X.); jingwu202508@163.com (J.W.); lilijiao202508@163.com (L.L.)

* Correspondence: zongzhao.zhai@foxmail.com (Z.Z.); wupa@hunnu.edu.cn (P.W.)

† These authors contributed equally to this work.

Simple Summary: Mosquitoes use ultraviolet light as a navigation cue, but the detailed behavioral characteristics and molecular mechanisms underlying this response remain poorly understood. In this study, we examined how mosquitoes respond to ultraviolet light under controlled laboratory conditions using three distinct experimental assays. We found that mosquitoes were strongly attracted to low-intensity ultraviolet light, but showed no preference between high-intensity ultraviolet light and darkness. Using video tracking and automated computational analysis of their movement, we further demonstrated that mosquitoes were more active under ultraviolet light and preferred areas illuminated with low-intensity ultraviolet light over dark regions. Through intrathoracic injection of dsRNA, we identified three genes essential for mosquito attraction to ultraviolet light. These results reveal, for the first time, the genetic mechanisms governing mosquito attraction to ultraviolet light. Our findings not only provide deeper insight into mosquito light-guided behavior but may also improve the design of light-based control strategies, opening new avenues for reducing mosquito populations and curbing disease transmission.

Abstract

It is widely recognized that mosquitoes are attracted to ultraviolet (UV) light traps in field and semi-field trials. However, the specific characteristics of mosquito behavioral responses to UV light remain poorly defined. Moreover, the molecular mechanisms underlying their phototactic behavior remain unexplored. Here, we characterized mosquito photobehavior under UV light in a laboratory setting using three experimental apparatuses. Our findings indicate that mosquitoes exhibit strong attraction to low-intensity UV light, yet show no preference between high-intensity UV light and darkness. Video recordings and automatic analyses of photobehavior under low-intensity UV light revealed that mosquitoes preferred the window illuminated by UV light over an unilluminated window and were more active when exposed to UV light. Through RNA interference (RNAi)-mediated knockdown of opsins highly expressed in the adult stage of *Culex quinquefasciatus*, we identified *CqOpsin3*, *CqOpsin5*, and *CqOpsin6* as crucial mediators of UV phototaxis. This study provides methods for characterizing mosquito photobehavior under UV light in the laboratory, and represents the first mechanistic investigation into UV light-mediated attractive behavior in mosquitoes.

Keywords: *Aedes*; *Culex quinquefasciatus*; mosquito; opsin; ultraviolet light

1. Introduction

Mosquitoes pose major public health problems worldwide by transmitting a variety of arboviruses, including dengue virus, Zika virus, yellow fever virus, and Japanese encephalitis virus [1,2]. Light traps, commonly used in field surveillance to monitor mosquito populations, are also frequently employed in households for mosquito control and to reduce arbovirus transmission [3,4]. Despite the widespread use of visible light traps to capture mosquitoes in the field, Liu, et al. [5] revealed that both diurnal and nocturnal mosquitoes are photonegative to visible light, a finding supported by other studies that recommend its removal from light traps [6,7]. Ultraviolet (UV) light, which lies beyond the human visual spectrum (also known as black light), can be perceived by many insects including mosquitoes [8]. Although UV light traps are widely used in field collections, mosquito photobehavior under UV light remains poorly understood.

In *Drosophila*, the behavioral responses to UV and visible light, as well as their interaction, have been extensively studied under controlled conditions [9,10]. In contrast, studies on mosquito responses to light have primarily relied on field-based methods, evaluating trapping efficiency by comparing the catch rates of light traps randomly placed in the field or semi-field conditions [3,11–13]. Recent studies have characterized mosquito behavioral responses to the visible spectrum under laboratory conditions. A wind-tunnel assay has been used to investigate the visual responses of *Aedes* mosquitoes to black objects in combination with other sensory cues including odor [14]. Our previous study further characterized the behavioral responses of *Aedes* and *Culex* mosquitoes to white light using an experimental apparatus modified from *Drosophila* photobehavior assay [5]. Despite these advances in studying visible light, mosquito behavioral responses to UV light under controlled laboratory conditions remain relatively poorly understood. Using a custom-designed arena, one study compared the circadian regulation of UV light-evoked attraction and avoidance behaviors in *Aedes* and *Anopheles* mosquitoes, but did not include *Culex* species [15]. The immediate behavioral responses of mosquitoes to UV light, as well as the specific sensory receptors mediating these responses, remain to be elucidated.

Bioinformatic analyses of assembled mosquito genomes, including those of *Aedes aegypti* (*Ae. aegypti*), *Anopheles gambiae*, and *Culex quinquefasciatus* (*Cx. quinquefasciatus*), have predicted a range of opsin genes and their potential sensitive wavelengths [16,17]. In *Cx. quinquefasciatus,* thirteen opsin genes have been identified by bioinformatic analysis, including eight predicted to be long-wavelength-sensitive, one UV-sensitive, and one short-wavelength-sensitive [16]. Molecular evolutionary studies further revealed that seven long-wavelength opsins in *Cx. quinquefasciatus* are homologous to the *Drosophila* Rh6 gene, likely resulting from a series of duplications and diversification [17]. However, among the mosquitoes in the Culicidae family analyzed, only one opsin in each mosquito species is homologous to the UV-sensitive Rh3 and Rh4 genes in *Drosophila* [17]. In addition to bioinformatic analyses, studies have systematically mapped the expression patterns of opsin genes in *Ae. aegypti* [18–21]. For instance, *Opsin1* is broadly expressed in R1–6 and R8 photoreceptors, while *Opsin2* and *Opsin8* are expressed in a non-overlapping pattern in R7 photoreceptor cells [20,21]. Additionally, *Opsin9* is co-expressed with either *Opsin2* or *Opsin8* in all R7 cells, as well as a subset of R8 cells [18]. Beyond expression profiling, functional studies using gene perturbation approaches have begun to dissect the roles of opsins in the behavioral response to visible light in *Aedes* mosquitoes [5,14]. Despite these advances, significant knowledge gaps remain. The expression patterns and molecular functions of opsin genes in *Culex* mosquitoes remain unexplored. Moreover, the molecular mechanisms underlying UV light perception remain uncharacterized across all mosquito species.

In this study, we characterized the photobehavior of *Cx. quinquefasciatus* under a gradient of UV light intensities, and identified *CqOpsin3*, *CqOpsin5*, and *CqOpsin6* as crucial factors in controlling its phototactic behavior toward UV light.

2. Materials and Methods

2.1. Mosquitoes

Mosquito colonies of *Aedes albopictus* (*Ae. albopictus*) (eggs purchased from Guangzhou Wolbaki Biotechnology Co., Ltd., Guangzhou, China), *Cx. quinquefasciatus* (eggs kindly provided by Y. Huang, Hunan Provincial Center for Disease Control and Prevention), and the Liverpool strain of *Ae. aegypti* (LVP, obtained from Prof. G.H. Wang, Institute of Zoology, Chinese Academy of Sciences, Beijing, China) were maintained under standard insectary conditions. Larvae were maintained in dechlorinated tap water and fed daily with a sterile liver powder suspension (60 g/L; CM0077, Oxoid, Tokyo, Japan). Pupae were collected using transfer pipettes and placed in bowls inside mesh cages for adult emergence. Adults were housed in 40 cm × 40 cm × 40 cm mesh cages with unlimited access to water and sugar sources (raisins). All developmental stages were maintained in incubators (HWS-1000, Ningbojiangnan, Ningbo, China) under standard insectary conditions (27 °C, 75% relative humidity, 12 h:12 h light/dark cycle). All behavioral assays were conducted with 4–10 days post-emergence adult female mosquitoes, as females are the host-seeking sex responsible for blood feeding and virus transmission, making them the biologically relevant target for investigating UV light attraction.

Based on the established ecological and morphological traits of *Cx. quinquefasciatus*, we conducted field collections targeting typical larval habitats of this species. Specifically, we surveyed sewage ditches, contaminated puddles, septic tanks, fertilizing buckets, and other organically polluted aquatic environments located in and around campus areas, as well as in nearby villages and farmland. Larvae and pupae were collected from these sites using a sieve, transferred to plastic containers, and transported to the laboratory. After emergence, adults were identified as *Cx. quinquefasciatus* based on key morphological characteristics and kept in mesh cages with unlimited access to water and sugar (raisins). All field-collected mosquitoes were maintained in a room exposed to natural sunlight through a window. The ambient temperature and humidity in the room were similar to outdoor conditions.

2.2. Mosquito Photobehavior

2.2.1. Y-Maze Photobehavior Assay

A Y-maze assay, which we previously designed to study mosquito behavioral responses to visible light [5], was adopted to test mosquito photobehavior under UV light. A computer-controlled Tunable Light Source (CME-TLSX300F, Microenery, Beijing, China) provided UV illumination to one of the two long arms of the Y-maze (Figure 1A) with light intensity measured by a UV Radiation Meter (UV340B, Sanpometer, Shenzhen, China). The total number of female mosquitoes tested in each group is presented in Table S1. For each biological replicate, we performed two technical replicates with the light source alternated between the two long arms to control for any arm-specific bias. Mosquitoes that remained in the release arm without making a choice were excluded from analysis. The preference index (PI) was calculated as follows: PI = (Number of mosquitoes in the UV illuminated arm − Number of mosquitoes in darkness)/(Number of mosquitoes in the UV illuminated arm + Number of mosquitoes in darkness). PI > 0 indicates that more mosquitoes preferred the UV illuminated area, while the opposite is true for PI < 0.




Figure 1. Phototaxis of female mosquitoes under UV light with a gradient of intensity. (**A**) Schematic diagram of the Y-maze assay. The arrowhead indicates the mosquito release point. (**B**–**E**) Preference for UV-lit versus shaded area. Behavioral response of laboratory-reared *Cx. quinquefasciatus* (**B**), laboratory-reared *Ae. albopictus* (**C**), laboratory-reared *Ae. aegypti* (**D**) and field-collected *Cx. quinquefasciatus* (**E**) to a gradient of UV light intensities. (**B**–**E**) Values represent mean ± SEM of ≥3 biological replicates (two technical replicates each). The total number of female mosquitoes tested in each group is presented in Table S1. Statistical significance of phototaxis relative to chance was computed using a one-sample *t*-test or Wilcoxon signed-rank test. Significance: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, ns: not significant.

2.2.2. Tube Photobehavior Assay

This assay was modified from a previously established tube assay [5] designed to investigate mosquito photobehavior under visible light, using the same experimental apparatus. While overhead white LED light provided illumination through the glass tube walls in the previous assay [5], UV light was directionally introduced from one open end of the glass tube without passing through the glass wall in this assay (Figure 2A). Prior to the experiment, mosquitoes were released into the glass tube and allowed to acclimate for 10 min (min), during which they distributed evenly within the tube. After the acclimation period, mosquitoes were exposed to UV light and allowed 5 min to make their choice. Light intensity was measured with a UV Radiation Meter (UV340B, Sanpometer, Shenzhen,

China). PI = (N_near − N_far)/(N_near + N_far), where N_near represents the number of mosquitoes in the half of the tube closest to the UV light source, and N_far represents the number of mosquitoes in the half of the tube farthest from the UV light source. PI > 0 indicates that more mosquitoes preferred the UV illuminated area, while the inverse is true for PI < 0.




Figure 2. Phototaxis of female mosquitoes under UV light of different wavelengths. (**A**) Schematic diagram of the tube assay. (**B**,**C**) Assessment of mosquito behavioral response to UV light using a tube assay. Mosquito photobehavior under 345 nm (**B**) and 395 nm (**C**) UV light at an intensity of 50 μW/cm^2. (**B**,**C**) Values represent mean ± SEM of ≥3 biological replicates (two technical replicates each). The total number of female mosquitoes tested in each group is presented in Table S1. Statistical significance of phototaxis relative to chance was computed using a one-sample *t*-test or Wilcoxon signed-rank test. Significance: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, ns: not significant.

2.2.3. Automatic Analysis of Mosquito Photobehavior

This assay was designed to allow automated object-tracking analysis. The test cage was composed of acrylic panels measuring 30 cm × 10 cm × 15 cm (length × width × height) and featured two meshed windows (9 cm × 9 cm) on one panel (Figure 3A). UV light at 345 nm, emitted from a computer-controlled Tunable Light Source (CME-TLSX300F, Microenery, Beijing, China) provided illumination for one of the two meshed windows. To control for positional bias, the meshed window illuminated by UV light was alternated after each trial. The white opaque test cage contained one transparent acrylic panel on the top to allow infrared light of 850 nm (SMD2835-300, Ledlightsworld, Shenzhen, China) to provide illumination for video recording in darkness. A camera (a2A1920-160 μm BAS, Basler, Ahrensburg, Germany) equipped with an infrared filter (central wavelength of 850 nm, BP850-35.5, Basler, Germany) was positioned opposite the meshed window. Recordings were made at 1920 × 800 resolution and 25 fps with manual focus and exposure settings using Pylon Viewer 8.0.0 software. Female mosquitoes were acclimated in the test cage for at least 24 h before testing with continuous access to water and sugar (raisins). For the photobehavior assay, each trial involved the release of eighty untreated mosquitoes. Mosquitoes were exposed to UV light of the indicated intensity and wavelength from a computer-controlled Tunable Light Source (CME-TLSX300F, Microenery, Beijing, China) at one of the two meshed windows for 2 min. Following a 30 min rest period in darkness, a technical replication was performed with the UV light illumination switched to the other window. To ensure consistency and minimize circadian variability, all behavioral assays were conducted between Zeitgeber Time (ZT) 6 and ZT12 for *Cx. quinquefasciatus*, and between ZT1 and ZT10 for *Ae. albopictus* and *Ae. aegypti*.




Figure 3. Video recording and automatic analysis of mosquito photobehavior under UV light. (**A**) Schematic diagram of video recording setup. Mosquitoes were exposed to 345 nm UV light from one of the two meshed windows, and their behavior was video recorded for 2 min. (**B**) Representative video frame taken from an experiment in which females were exposed to 345 nm UV light at 50 $\mu W/cm^2$. (**C**–**F**) Photo preference between darkness and 345 nm UV light at 50 $\mu W/cm^2$. Shown are the preference index (**C**), the total number of mosquito tracks (**D**), the cumulative distance moved by mosquitoes (**E**) and the average dwell time per mosquito track (**F**) on UV-lit versus unlit mesh window. (**C**–**F**) Values represent mean ± SEM of ≥3 biological replicates (two technical replicates each). The total number of female mosquitoes tested in each group is presented in Table S1. (**C**) Data were analyzed using one sample *t*-test. (**D**–**F**) Group comparisons were performed using unpaired *t*-test. Significance: * $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$, ns: not significant.

We used an automated object-tracking and preference index calculation script based on MATLAB (R2022b, MathWorks, Natick, MA, USA), modified from Chandel, et al. [22], to analyze mosquitoes in recorded video. The workflow of the object-tracking script includes the following steps: (1) reading the recorded video and manually selecting two regions of interest (corresponding to the two meshed windows); (2) randomly extracting 100 frames from the video to calculate the modal pixel value, which is used to generate a background model; (3) identifying mosquitoes using the background model and a white-and-black model. The background model identifies moving objects by analyzing the absolute differences in pixel values between each frame and the background. The white-and-black model detects stationary mosquitoes which appear as black blobs against the white background; (4) assigning IDs to each tracked mosquito for motion analysis; (5) calculating the PI and other behavioral metrics. The overall PI for a given time period was calculated as the average of the PI values from each frame. For each frame, PI was defined as: PI = (Number

of mosquitoes on UV-illuminated window − Number of mosquitoes on the window in darkness)/(Number of mosquitoes on UV-illuminated window + Number of mosquitoes on the window in darkness). PI > 0 indicates that more mosquitoes preferred the UV illuminated area, while the inverse is true for PI < 0.

Track number was calculated as the sum of IDs in the respective window within the 2 min recorded time. The cumulative distance mosquitoes moved on a given mesh window was calculated as the sum of the distances traveled by each track on that window. The distance of each track was calculated as the sum of the Euclidean distances between two adjacent coordinates of the track. The average dwell time per track was calculated by dividing the total time spent by all mosquitoes in that window by the number of mosquito tracks. The total time spent by each mosquito in a respective window was calculated by dividing the number of video frames the mosquito spent in that window by the video frame rate.

2.3. RNA Extraction and Quantification of Gene Expression

The RNA expression profile of *Cx. quinquefasciatus* and *Ae. aegypti* were analyzed at selected developmental stages, including fourth-instar larvae, pupae and adults aged 5 to 7 days post-emergence. Tissues from these stages were homogenized for RNA extraction. Female adult mosquitoes on the third day post double-strand RNA (dsRNA) injection were also homogenized for RNA extraction. RNA extraction and complementary DNA (cDNA) synthesis were performed using RNAiso Plus (9190, Takara, Beijing, China) and the PrimeScript RT reagent kit (RR037A, Takara, Beijing, China), respectively. To analyze opsin gene expression in *Ae. albopictus* and *Cx. quinquefasciatus*, RNA samples for quantitative real-time polymerase chain reaction (RT-qPCR) were collected within the same circadian time window as that used for behavioral assays.

Gene expression was analyzed by RT-qPCR with the ChamQ SYBR Color qPCR Master Mix (Q421, Vazyme, Nanjing, China) on a Quantagene q225 thermal cycler system (q225, Kubo Technology, Beijing, China). The primers for RT-qPCR are listed in Supplementary Table S2. Gene expression levels were normalized to the reference genes Rpl8 (*LOC6031076*) in *Cx. quinquefasciatus* and actin (*AAEL011197*) in *Ae. aegypti*.

2.4. Gene Silencing in Mosquitoes

Gene silencing was performed using synthetized and purified dsRNA prepared with the MEGAscript T7 transcription kit (AM1334, Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. Female adult mosquitoes, aged 0–5 days post-eclosion and not blood-fed, were anesthetized at 4 °C and placed on a CO_2 gas plate. Using the Glass Micromotor Syringe Pump (R-480, RWD, Shenzhen, China), 300 nL of solution containing 1 μg of dsRNA was injected into the thorax of each mosquito. Behavioral assays were conducted 3 days post-injection, and silencing efficiency was validated by RT-qPCR. Mosquitoes injected with double-strand green fluorescent protein (dsGFP) were used as the control group. The primers used in this assay are listed in Supplementary Table S3.

2.5. Statistical Analysis

Statistical analyses and graphing were performed using GraphPad Prism 8 (Prism, La Jolla, CA, USA). A schematic summary of assay type, mosquito species, light wavelength, light intensity, biological replicates, total number of mosquitoes tested, and test duration is provided in Supplementary Table S1. All datasets were assessed for normality. To determine whether PI significantly differed from chance, we applied a one-sample *t*-test (for normal distribution data) or a Wilcoxon signed-rank test (for non-normal distributions). Comparisons between two groups were conducted using an unpaired *t*-test (normal distribution) or a Mann–Whitney test (non-normal distribution). Multiple group comparisons

were performed using one-way ANOVA with Tukey's post hoc test (normal distribution) or a Kruskal–Wallis test with Dunn's post hoc test (non-normal distribution). The p value was used to evaluate the null hypothesis that any observed effect was due to chance or that there was no difference between the means of the two groups. Statistical significance was set at the level of specifically * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, ns: not significant. Details of the test methods are summarized in Tables S4 and S5. In figures, datasets labeled with different lowercase letters indicate statistically significant differences between groups.

2.6. Ethical Note

The study has been approved by the Hunan Normal University Ethics Committee (Protocol 188/2022).

3. Results

3.1. Mosquitoes Preferred Low Intensity UV over Darkness

Previous studies have assessed mosquito responses to UV light using light traps in field or semi-field trials by evaluating attraction efficiency based on the number of mosquitoes caught per trap [3,11–13]. However, to investigate the mechanisms underlying mosquito phototaxis in response to UV light, a controlled laboratory assay needs to be established. We employed a Y-maze assay (Figure 1A), adapted from a well-established *Drosophila* light/dark assay used to study photobehavior [23], to characterize mosquito photobehavior under UV light. The preference index was calculated following an approach similar to that used in *Drosophila* phototaxis studies [9]: PI = $(N_{UV} - N_D)/(N_{UV} + N_D)$, where N_{UV} and N_D represent the number of mosquitoes on the UV-illuminated side and the dark side, respectively. We conducted a Y-maze assay in which mosquitoes could choose between darkness and a gradient of UV light intensities. Laboratory-reared *Cx. quinquefasciatus* and *Ae. albopictus* consistently preferred UV light at intensities ranging from 50 $\mu W/cm^2$ to 1000 $\mu W/cm^2$ over darkness, but did not show a significant preference between darkness and UV light at 1500 $\mu W/cm^2$ (Figure 1B,C). The *Ae. aegypti* Liverpool strain (LVP) was more sensitive to UV light, preferring intensities between 5 $\mu W/cm^2$ and 50 $\mu W/cm^2$ over darkness, but exhibited no significant preference between darkness and UV light at 150 $\mu W/cm^2$, 500 $\mu W/cm^2$, and 1000 $\mu W/cm^2$ (Figure 1D). While UV light is widely used for mosquito trapping in the field, we investigated whether field-collected mosquitoes displayed similar photobehavior. We collected larvae and pupae of *Cx. quinquefasciatus* from the field and tested the emerged adults using the Y-maze assay. Consistently, field-collected *Cx. quinquefasciatus* preferred low-intensity UV light below 1000 $\mu W/cm^2$, but no significant phototaxis was observed at 1500 $\mu W/cm^2$ (Figure 1E). These results indicate that mosquito phototaxis is affected by illumination intensity. Accordingly, subsequent experiments with *Cx. quinquefasciatus* and *Ae. albopictus* were conducted using UV light intensities below 1000 $\mu W/cm^2$, while *Ae. aegypti* was tested at 50 $\mu W/cm^2$.

To further validate the observation that mosquitoes exhibit positive phototaxis toward low-intensity UV light, we conducted a tube assay using a 60 cm glass tube illuminated by UV light at one end (Figure 2A). UV light at 50 $\mu W/cm^2$ attracted *Cx. quinquefasciatus* and *Ae. albopictus* to the near-UV end of the tube but did not affect the distribution of *Ae. aegypti* (Figure 2B). Subsequently, we tested UV light at 150 $\mu W/cm^2$ measured at the near-UV end of the tube. Consistently, *Cx. quinquefasciatus* and *Ae. albopictus* that were initially randomly distributed in the tube moved toward the UV light within 5 min after light exposure (Figure S1A). While the abovementioned experiments evaluated mosquito phototaxis with 345 nm UV light, we further investigated mosquito photobehavior under 395 nm UV light. When tested with a Y-maze assay, all three mosquito species preferred

the UV light-illuminated environment at 395 nm over an unilluminated one (Figure S1B). In the tube assay, 395 nm UV light of 50 $\mu W/cm^2$ attracted *Cx. quinquefasciatus* and *Ae. albopictus* to the near-UV end of the tube, whereas it failed to modulate the distribution of *Ae. aegypti* (Figure 2C). Collectively, these results demonstrated that *Cx. quinquefasciatus* and *Ae. albopictus* were significantly attracted to low-intensity UV light across different spectra and experimental setups. By contrast, the behavioral response of *Ae. aegypti* to UV light was context-dependent.

3.2. UV Light Affects Mosquito Preference and Activity

To further illustrate mosquito photobehavior under UV light, we developed an additional behavioral assay to enable video recording and automated analysis of mosquito behavior. Custom-designed opaque acrylic cages with two mesh windows on one panel (Figure 3A) were used for subsequent study in a dark room. Eighty acclimatized female mosquitoes, housed in the assay cage for over 24 h, were exposed to UV light (50 $\mu W/cm^2$) emanating from one of the two windows for 2 min, while their behavior was recorded (Figure 3B and Supplementary Video S1). We used a custom video-tracking program modified from a previous study [22] to automatically analyze mosquito behavior. Compared with traditional manual measurement at set intervals, the automated program enabled continuous recording and analysis of mosquito activity throughout the 2 min experimental window (Supplementary Video S2). We calculated the average PI for the 2 min video by summing the PI of each frame and dividing by the total number of frames, with PI of each frame calculated as $PI = (N_{UV} - N_D)/ (N_{UV} + N_D)$, where N_{UV} and N_D represent the number of mosquitoes on the UV-illuminated side and the dark side, respectively. Consistent with the Y-maze assay, the results showed that mosquitoes significantly preferred the UV-illuminated area over the unilluminated area (Figure 3C). To further characterize mosquito photobehavior under UV light, we calculated three additional metrics: the number of mosquito tracks at each window, the cumulative distance mosquitoes moved on the respective mesh window across all tracks, and the average dwell time per mosquito track on the respective mesh window. The number of mosquito tracks as well as the total distance of tracks in the UV-illuminated window was significantly larger compared to those in the unilluminated window (Figure 3D,E). By contrast, laboratory-reared *Ae. albopictus* and *Cx. quinquefasciatus* mosquitoes spent significantly longer time per track in darkness than at the UV-illuminated window (Figure 3F). These results suggest that mosquitoes were more active under UV illumination, while mosquitoes in darkness were predominantly stationary. UV light of 150 $\mu W/cm^2$ drew similar conclusions (Figure S2). Collectively, our results show that mosquitoes preferred UV light and exhibited enhanced activity.

3.3. CqOpsin3, CqOpsin5, and CqOpsin6 Mediate Phototactic Behavior to UV Light in Cx. quinquefasciatus

Opsins are a class of transmembrane proteins that initiate photon-induced signaling pathways in both invertebrates and vertebrates [16]. Our previous study identified *opsin1* as a critical regulator of photonegative behavior toward visible light [5]. Here, we investigated which opsin regulates the positive phototaxis of *Aedes* mosquitoes toward UV light. RT-qPCR analysis identified four opsin genes highly expressed in the adult stage of *Ae. aegypti* (Figure 4A). We knocked down these four opsins at the transcript level through intrathoracic injection of dsRNA, resulting in a reduction in their relative abundance of over 74% (Figure S3). However, silencing these genes had no significant effect on the photopositive behavior of *Ae. aegypti* toward low-intensity UV light (Figure 4B).




Figure 4. Photobehavior of opsin-silenced *Ae. aegypti*. (**A**) Expression of *Ae. aegypti* opsin genes in fourth-instar larvae (4th), pupae, and adults, normalized to *Ae. aegypti* Actin (*AAEL011197*). (**B**) The photobehavior of opsin-silenced *Ae. aegypti* assessed using a Y-maze assay under 345 nm UV light at an intensity of 10 $\mu W/cm^2$. Values represent mean ± SEM of ≥3 biological replicates (two technical replicates each). The total number of female mosquitoes tested in each group is presented in Table S1. Statistical significance of phototaxis relative to chance was computed using a one-sample *t*-test or Wilcoxon signed-rank test. Multiple group comparisons were analyzed using the non-parametric Kruskal–Wallis test, followed by Dunn's test. Significance: ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Different lowercase letters indicate statistically significant differences between groups.

We next investigated which opsins regulate the positive phototaxis of *Culex* mosquitoes toward UV light. Bioinformatic analysis predicted thirteen opsin genes in *Cx. quinquefasciatus* [16]. To elucidate the mechanisms underlying photopositive responses to low-intensity UV light, we first profiled the expression patterns of these thirteen opsin genes in *Cx. quinquefasciatus* using RT-qPCR (Figure 5A). The results revealed that *CqOpsin2*, *CqOpsin3*, *CqOpsin5*, *CqOpsin6* and *CqOpsin11* were highly expressed in adults (Figure 5A). To assess their functional roles in low-intensity UV light responses, we conducted RNAi-based screening by intrathoracic injection of double-stranded RNA targeting each of the five highly expressed opsins. This approach efficiently knocked down all five opsins at the transcript level, reducing their relative abundance by more than 59% (Figure S4). Knockdown of *CqOpsin5* and *CqOpsin6* completely abolished the photopositive behavior of *Cx. quinquefasciatus* toward UV light at 500 $\mu W/cm^2$, whereas silencing *GFP* (negative control), *CqOpsin2* and *CqOpsin11* had no discernable effect (Figure 5B). Additionally, silencing *CqOpsin3* significantly reduced photopositive behavior under 500 $\mu W/cm^2$ UV light (Figure 5B). Furthermore, knockdown of *CqOpsin3*, *CqOpsin5* and *CqOpsin6* completely abolished the photopositive behavior under 50 $\mu W/cm^2$ UV light (Figure 5C). To validate the roles of these opsins in UV light-mediated attraction in field-derived *Cx. quinquefasciatus*, we silenced the five adult-expressed opsins in adults that had emerged from field-collected larvae and pupae. Consistently, silencing *CqOpsin3*, *CqOpsin5* and *CqOpsin6* significantly abolished the photopositive behavior toward UV light in field-collected mosquitoes (Figure 5D). Moreover, we further confirmed these findings using a tube assay (Figure 5E). Collectively, these results demonstrate that *CqOpsin3*, *CqOpsin5* and *CqOpsin6* play vital roles in UV light-mediated attraction behavior in *Cx. quinquefasciatus*.




Figure 5. Photobehavior of opsin-silenced *Cx. quinquefasciatus*. (**A**) Expression of *Cx. quinquefasciatus* opsin genes in fourth-instar larvae (4th), pupae, and adults, normalized to *Cx. quinquefasciatus* Rpl8 (*LOC6031076*). (**B**–**D**) Photobehavior of opsin-silenced *Cx. quinquefasciatus* assessed using the Y-maze assay. Effects of opsin knockdown on photobehavior of laboratory-reared *Cx. quinquefasciatus* under 345 nm UV light of 500 μW/cm^2 (**B**) and 50 μW/cm^2 (**D**). Effects of opsin gene knockdown on the photobehavior of field-collected *Cx. quinquefasciatus* under 345 nm UV light of 500 μW/cm^2. (**E**) Photobehavior of opsin-silenced *Cx. quinquefasciatus* evaluated using a tube assay under 345 nm UV light at an intensity of 500 μW/cm^2. (**B**–**E**) Values represent mean ± SEM of ≥3 biological replicates (two technical replicates each). The total number of female mosquitoes tested in each group is presented in Table S1. Statistical significance of phototaxis relative to chance was computed using a one-sample *t*-test or Wilcoxon signed-rank test. Different lowercase letters indicate statistically significant differences between groups. (**B**–**E**) Multiple group comparisons were analyzed using a one-way ANOVA, followed by Tukey's multiple comparisons test. (**C**) Multiple group comparisons were analyzed using the non-parametric Kruskal–Wallis test, followed by Dunn's test. Significance: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, ns: not significant.

4. Discussion

It is widely acknowledged that mosquitoes are attracted to UV light; however, the molecular mechanisms underlying this phenomenon remain unexplored. To address this gap, we characterized mosquito photobehavior under UV light using a Y-maze assay and further analyzed their behavior in a test cage with an automatic tracking program. Through dsRNA-mediated knockdown, we identified *CqOpsin3*, *CqOpsin5* and *CqOpsin6* as essential regulators of UV light-induced phototaxis.

Our results showed that UV light at low-intensities is an attractive cue for mosquitoes, whereas exposure to high-intensities of UV light (above 1500 μW/cm^2) failed to elicit attrac-

tion. A comparable phenomenon has been reported in *Drosophila* [24]. It is well-established that adult flies are attracted to UV light within minutes of its onset [25]. However, a recent study indicated that UV light intensity influences the decision between positive phototaxis and avoidance behavior [24]. This avoidance response is thought to reflect an evolutionary adaptation to avoid the detrimental effects of short-wavelength light [24,26].

Another notable finding of our study is that *Ae. aegypti* exhibited heightened sensitivity to UV light, being attracted only at intensities below 50 $\mu W/cm^2$, unlike other tested mosquito species. Although *Ae. aegypti* and *Ae. albopictus* are closely related evolutionarily, with their corresponding opsins sharing high sequence similarity [5], their phototactic responses to UV light between 150 $\mu W/cm^2$ and 1500 $\mu W/cm^2$ diverged significantly. Whether this differential arises from differential opsin expression, variation in neural processing, or contributions from non-visual sensory pathways remains to be elucidated. Indeed, our study is limited in that we did not assess the photobehavior of field-collected *Ae. aegypti,* and all experiments were conducted under controlled laboratory conditions. Further validation of this species-specific behavioral response in the field could provide important theoretical insights for optimizing UV light traps to more precisely target particular mosquito populations.

Phylogenetic analyses predicted *AaegOpsin8* to be UV-sensitive [16]; however, individual knockdown of this gene, as well as each of the other highly expressed adult-stage opsins, did not affect UV attraction in *Ae. aegypti* in our assays. One possible explanation is functional redundancy, whereby another yet unidentified UV-sensitive opsin may compensate for the loss of *AaegOpsin8*. To test this hypothesis, generating double or triple opsin knockout mutants using genetic engineering approaches would be a useful direction for future research.

Bioinformatic analysis predicted only one UV-sensitive opsin in *Cx. quinquefasciatus* [16], whereas our study identified three opsins that are essential for photopositive behavior toward UV light. Although *Opsin5* and *Opsin6* were predicted to be long-wavelength-sensitive opsins [16], our functional data demonstrate their requirement for UV phototaxis; nevertheless, their exact mechanism of action remains to be elucidated. Notably, the *Drosophila* opsin Rh1 exhibits broad-spectrum sensitivity to blue and green light, alongside a secondary peak in the UV range resulting from a bound carotenoid-derived pigment [27,28]. Therefore, characterizing the spectral properties of *Culex* opsins using techniques such as microspectrophotometry (MSP) and electroretinography (ERG) with ectopic expression of each opsin in mosquito photoreceptor cells could provide valuable information on mosquito photobehavior under UV light in addition to our behavioral tests.

Besides opsins, cryptochromes (CRY), which are key regulators of circadian rhythm in response to blue or UV-A light, have also been shown to modulate UV light-evoked avoidance/attraction behavior in *Drosophila* [26,29]. In fruit flies, attraction under low-intensity UV light is primarily mediated by opsins in photoreceptor cells, whereas avoidance under high-intensity UV light is regulated through CRY and Rh7-dependent pathways [24]. Whether similar mechanisms involving cryptochromes and other non-opsin phototransduction molecules influence photobehavior in mosquitoes remains an open question that warrants further investigation.

In conclusion, the behavioral response of mosquitoes to UV light follows a dosage-dependent pattern, highlighting the need to optimize UV intensity for field collections. Our study represents the first mechanistic research into UV light-mediated attractive behavior, revealing the vital role of *CqOpsin3*, *CqOpsin5*, and *CqOpsin6* in UV light phototaxis of *Cx. quinquefasciatus*. These findings not only advance our understanding of mosquito photosensory biology but also provide a functional foundation for developing more efficient and species-specific light-based control strategies.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/insects16100997/s1, Figure S1: The photobehavior of female mosquitoes under different wavelengths of UV light; Figure S2: Video recordings and automated analysis were conducted to assess mosquito photobehavior under 345 nm UV light at an intensity of 150 $\mu W/cm^2$; Figure S3: RNAi-mediated knockdown efficiency of *Ae. aegypti* opsin genes; Figure S4: RNAi-mediated knockdown efficiency of *Cx. quinquefasciatus* opsin genes; Table S1: Summary of experimental conditions for all behavioral assays; Table S2: qPCR primers of opsin genes used in expression profile; Table S3: Primers for dsRNA synthesis and detection; Table S4: Details of the test methods used for each sample analysis; Table S5: The results of multiple comparisons test for testing among different groups; Video S1: Representative recording of mosquito photobehavior; Video S2: Automated analysis of mosquito activity.

Author Contributions: Conceptualization, P.W.; methodology, P.W.; software, G.Z.; validation, X.L., G.Z. and H.L.; formal analysis, X.L. and G.Z.; investigation, X.L., G.Z., H.L., Y.M., M.X., J.W. and L.L.; resources, Z.Z. and P.W.; data curation, X.L. and G.Z.; writing—original draft preparation, P.W.; writing—review and editing, X.L., G.Z. and P.W.; visualization, X.L.; supervision, Z.Z. and P.W.; project administration, P.W.; funding acquisition, P.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Science Fund for Distinguished Young Scholars of Hunan Province (2022JJ10039) and the Science and Technology Program of Hunan Province (2019RS1036 and 2021RC3095).

Data Availability Statement: The data presented in this study are openly available in the Zenodo data repository at 10.5281/zenodo.14942954, with the URL: https://zenodo.org/records/14942954?preview=1&token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6IjY3YzgwZThjLWIxYTAtNDg2MC04MTNlLTJkMTJkMDAwMGMxNyIsImRhdGEiOnt9LCJyYW5kb20iOiIyMTAxOTJhOTdmOGZhM2JlNDc0Y2U3OTRkZDY4ZTE3ZCJ9.SauOXEheZWkIThKnsAwjMLR9Ro_InnekWZg-omxso8zNI0dfqORLcmSH2mJc3OhIViB0T393oOtqc6KFMMNUxA, accessed on 19 September 2025, and reference number [30]. The code for automated analysis is available in the GitHub (https://github.com/zgq2020/Liu-et-al.-2025, version 1.0, accessed on 1 August 2025).

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

Ae. aegypti	*Aedes aegypti*
Ae. albopictus	*Aedes albopictus*
Cx. quinquefasciatus	*Culex quinquefasciatus*
dsGFP	Double-strand green fluorescent protein
dsRNA	Double-strand RNA
min	Minute
PI	Preference index
RNAi	RNA interference
RT-qPCR	Quantitative real-time polymerase chain reaction
UV	Ultraviolet
ZT	Zeitgeber time

References

1. Zhu, Y.; Zhang, C.; Zhang, L.; Yang, Y.; Yu, X.; Wang, J.; Liu, Q.; Wang, P.; Cheng, G. A human-blood-derived microRNA facilitates flavivirus infection in fed mosquitoes. *Cell Rep.* **2021**, *37*, 110091. [CrossRef] [PubMed]
2. Wu, P.; Yu, X.; Wang, P.; Cheng, G. Arbovirus lifecycle in mosquito: Acquisition, propagation and transmission. *Expert Rev. Mol. Med.* **2019**, *21*, e1. [CrossRef]

3. Lhaosudto, S.; Ngoen-Klan, R.; Meunworn, V.; Kongmee, M.; Hii, J.; Chareonviriyaphap, T. Comparison of different spectral ranges of UV-LED lighting for outdoor mosquito trapping in forested area in Thailand. *J. Med. Entomol.* **2024**, *61*, 1510–1518. [CrossRef]
4. Lee, H.I.; Seo, B.Y.; Shin, E.H.; Burkett, D.A.; Lee, J.K.; Shin, Y.H. Efficiency evaluation of Nozawa-style black light trap for control of anopheline mosquitoes. *Korean J. Parasitol.* **2009**, *47*, 159–165. [CrossRef]
5. Liu, X.; Yang, S.; Yao, Y.; Wu, S.; Wu, P.; Zhai, Z. Opsin1 regulates light-evoked avoidance behavior in Aedes albopictus. *BMC Biol.* **2022**, *20*, 110. [CrossRef]
6. Harker, A.; Fairbanks, K.; Dewsnup, M.A.; Faraji, A.; Bibbs, C.S. Blinded by the light: Does heat or light enhance wild mosquito (Diptera: Culicidae) attraction to CO_2-baited traps in the Great Salt Lake area? *J. Med. Entomol.* **2024**, *61*, 802–807. [CrossRef] [PubMed]
7. Boze, B.G.V.; Renfro, K.; Markowski, D.; Lozano-Fuentes, S. Effect of Incandescent Light on Collection of West Nile Virus Vectors Using CDC Miniature Light Traps in Northern Colorado. *J. Insect Sci.* **2021**, *21*, 10. [CrossRef]
8. Shimoda, M.; Honda, K.-I. Insect reactions to light and its applications to pest management. *Appl. Entomol. Zool.* **2013**, *48*, 413–421. [CrossRef]
9. Yamaguchi, S.; Desplan, C.; Heisenberg, M. Contribution of photoreceptor subtypes to spectral wavelength preference in Drosophila. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 5634–5639. [CrossRef]
10. Gao, S.; Takemura, S.Y.; Ting, C.Y.; Huang, S.; Lu, Z.; Luan, H.; Rister, J.; Thum, A.S.; Yang, M.; Hong, S.T.; et al. The neural substrate of spectral preference in Drosophila. *Neuron* **2008**, *60*, 328–342. [CrossRef]
11. Pimnon, S.; Ngoen-Klan, R.; Sumarnrote, A.; Chareonviriyaphap, T. UV Light-Emitting-Diode Traps for Collecting Nocturnal Biting Mosquitoes in Urban Bangkok. *Insects* **2022**, *13*, 526. [CrossRef]
12. Mwanga, E.P.; Ngowo, H.S.; Mapua, S.A.; Mmbando, A.S.; Kaindoa, E.W.; Kifungo, K.; Okumu, F.O. Evaluation of an ultraviolet LED trap for catching Anopheles and Culex mosquitoes in south-eastern Tanzania. *Parasites Vectors* **2019**, *12*, 418. [CrossRef]
13. Kim, D.Y.; Leepasert, T.; Bangs, M.J.; Chareonviriyaphap, T. Semi-field evaluation of novel chemical lures for Aedes aegypti, Culex quinquefasciatus, and Anopheles minimus (Diptera: Culicidae) in Thailand. *Parasites Vectors* **2021**, *14*, 606. [CrossRef] [PubMed]
14. Zhan, Y.; Alonso San Alberto, D.; Rusch, C.; Riffell, J.A.; Montell, C. Elimination of vision-guided target attraction in Aedes aegypti using CRISPR. *Curr. Biol.* **2021**, *31*, 4180–4187. [CrossRef] [PubMed]
15. Baik, L.S.; Nave, C.; Au, D.D.; Guda, T.; Chevez, J.A.; Ray, A.; Holmes, T.C. Circadian Regulation of Light-Evoked Attraction and Avoidance Behaviors in Daytime- versus Nighttime-Biting Mosquitoes. *Curr. Biol.* **2020**, *30*, 3252–3259.e3253. [CrossRef] [PubMed]
16. Giraldo-Calderón, G.I.; Zanis, M.J.; Hill, C.A. Retention of duplicated long-wavelength opsins in mosquito lineages by positive selection and differential expression. *BMC Evol. Biol.* **2017**, *17*, 84. [CrossRef]
17. Feuda, R.; Goulty, M.; Zadra, N.; Gasparetti, T.; Rosato, E.; Pisani, D.; Rizzoli, A.; Segata, N.; Ometto, L.; Stabelli, O.R. Phylogenomics of Opsin Genes in Diptera Reveals Lineage-Specific Events and Contrasting Evolutionary Dynamics in Anopheles and Drosophila. *Genome Biol. Evol.* **2021**, *13*, evab170. [CrossRef] [PubMed]
18. Hu, X.; Whaley, M.A.; Stein, M.M.; Mitchell, B.E.; O'Tousa, J.E. Coexpression of spectrally distinct rhodopsins in Aedes aegypti R7 photoreceptors. *PLoS ONE* **2011**, *6*, e23121. [CrossRef]
19. Rocha, M.; Kimler, K.J.; Leming, M.T.; Hu, X.; Whaley, M.A.; O'Tousa, J.E. Expression and light-triggered movement of rhodopsins in the larval visual system of mosquitoes. *J. Exp. Biol.* **2015**, *218*, 1386–1392. [CrossRef]
20. Hu, X.; Leming, M.T.; Metoxen, A.J.; Whaley, M.A.; O'Tousa, J.E. Light-mediated control of rhodopsin movement in mosquito photoreceptors. *J. Neurosci.* **2012**, *32*, 13661–13667. [CrossRef]
21. Hu, X.; England, J.H.; Lani, A.C.; Tung, J.J.; Ward, N.J.; Adams, S.M.; Barber, K.A.; Whaley, M.A.; O'Tousa, J.E. Patterned rhodopsin expression in R7 photoreceptors of mosquito retina: Implications for species-specific behavior. *J. Comp. Neurol.* **2009**, *516*, 334–342. [CrossRef]
22. Chandel, A.; DeBeaubien, N.A.; Ganguly, A.; Meyerhof, G.T.; Krumholz, A.A.; Liu, J.; Salgado, V.L.; Montell, C. Thermal infrared directs host-seeking behaviour in Aedes aegypti mosquitoes. *Nature* **2024**, *633*, 615–623. [CrossRef]
23. Choe, K.M.; Clandinin, T.R. Thinking about visual behavior; learning about photoreceptor function. *Curr. Top. Dev. Biol.* **2005**, *69*, 187–213. [CrossRef]
24. Baik, L.S.; Recinos, Y.; Chevez, J.A.; Au, D.D.; Holmes, T.C. Multiple Phototransduction Inputs Integrate to Mediate UV Light-evoked Avoidance/Attraction Behavior in Drosophila. *J. Biol. Rhythms* **2019**, *34*, 391–400. [CrossRef]
25. Zhu, E.Y.; Guntur, A.R.; He, R.; Stern, U.; Yang, C.H. Egg-laying demand induces aversion of UV light in Drosophila females. *Curr. Biol.* **2014**, *24*, 2797–2804. [CrossRef]
26. Baik, L.S.; Recinos, Y.; Chevez, J.A.; Holmes, T.C. Circadian modulation of light-evoked avoidance/attraction behavior in Drosophila. *PLoS ONE* **2018**, *13*, e0201927. [CrossRef]

27. Stark, W.S.; Stavenga, D.G.; Kruizinga, B. Fly photoreceptor fluorescence is related to UV sensitivity. *Nature* **1979**, *280*, 581–583. [CrossRef]
28. Sharkey, C.R.; Blanco, J.; Leibowitz, M.M.; Pinto-Benito, D.; Wardill, T.J. The spectral sensitivity of Drosophila photoreceptors. *Sci. Rep.* **2020**, *10*, 18242. [CrossRef] [PubMed]
29. Baik, L.S.; Fogle, K.J.; Roberts, L.; Galschiodt, A.M.; Chevez, J.A.; Recinos, Y.; Nguy, V.; Holmes, T.C. CRYPTOCHROME mediates behavioral executive choice in response to UV light. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 776–781. [CrossRef] [PubMed]
30. Liu, X.; Zhao, G.; Liu, H.; Mao, Y.; Xu, M.; Wu, J.; Li, L.; Zhai, Z.; Wu, P. Elimination of ultraviolet light-mediated attraction behavior in Culex mosquitoes via dsRNA-mediated knockdown of Opsins. *Zenodo* **2025**. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Transcriptome and Functional Analyses Revealed the Carboxylesterase Genes Involved in Pyrethroid Resistance in *Anopheles sinensis* (Diptera: Culicidae)

Yiyun Wei †, Xinyao Gu †, Fengling Si, Xiaojie Chen, Liang Qiao, Hongxing Yan and Bin Chen *

Chongqing Key Laboratory of Vector Control and Utilization, Institute of Entomology and Molecular Biology, College of Life Sciences, Chongqing Normal University, Chongqing 401331, China; weiyiyun1988@163.com (Y.W.); gxy576692235@163.com (X.G.); sifengling1217@163.com (F.S.); chenxiaojie092355@163.com (X.C.); qiaoliangswu@163.com (L.Q.); 18716278039@163.com (H.Y.)

* Correspondence: bin.chen@cqnu.edu.cn

† These authors contributed equally to this work.

Simple Summary: Pyrethroid resistance in *Anopheles sinensis*, a major malaria vector in Asia, severely compromises insecticide-based vector control. This study identified carboxylesterases as important factors in the mechanism of resistance to deltamethrin. Transcriptomic comparison of deltamethrin-resistant and susceptible strains revealed four upregulated carboxylesterase (CCE) genes. Based on prior evidence, we evaluated eight candidate CCEs, confirming significant upregulation of five genes (*AsAe9*, *AsAe10*, *AsAce1*, *AsAce2*, and *AsBe4*) in resistant strains via quantitative real-time PCR (qRT-PCR). The gene silencing mediated by RNA interference (RNAi) reduced the knockdown time and increased the mortality rates of resistant mosquitoes when exposed to deltamethrin. This study revealed carboxylesterase genes involved in pyrethroid resistance in *An. sinensis*, providing critical insights into resistance mechanisms and management strategies in mosquitoes.

Abstract

Carboxylesterases (CCEs) have been demonstrated to be involved in pyrethroid resistance in insect species. This study aims to investigate CCE-mediated resistance mechanisms in *Anopheles sinensis*, a major malaria vector. Through comparative transcriptomics of a deltamethrin-resistant strain (CQ-LR) versus susceptible strain (WX-LS) of *An. sinensis*, we identified differentially expressed CCE genes across five developmental stages, five tissues, and three time points post-blood-meal. Four candidate genes (*AsAe9*, *AsAe10*, *AsAce2*, *AsUn5*) showed significantly upregulated expression. Subsequent qRT-PCR validation across four field-derived resistant strains (WX-LR, AH-LR, YH-LR, CQ-LR) and the susceptible strain confirmed significant upregulation of *AsAe9*, *AsAe10*, *AsAce1*, *AsAce2*, and *AsBe4* in more than two resistant populations. RNAi-based functional validation showed that silencing *AsAe10* or *AsBe4* in the WX-LR strain significantly decreased knockdown time and raised 24 h mortality upon diagnostic deltamethrin exposure, with *AsAe10* silencing having the strongest effect. This study identifies CCE genes involved in deltamethrin resistance in *An. sinensis*, providing valuable insights into the resistance mechanisms of pyrethroid and a theoretical basis for mosquito resistance management.

Keywords: *Anopheles sinensis*; pyrethroid resistance; carboxylesterases; RNAi; functional analyses

1. Introduction

Anopheles sinensis, a primary vector for malaria and filariasis, exhibits extensive distribution across China and other East Asian nations [1–3]. The vivax malaria outbreak demonstrates strong epidemiological association with elevated vectorial capacity in *An. sinensis* for *Plasmodium vivax* [4]. Current strategies for controlling *An. sinensis* predominantly rely on pyrethroid-based interventions—insecticide-treated nets and indoor residual spraying—alongside supplemental organophosphate, organochlorine, and carbamate applications [5]. Alarmingly, the development of mosquito resistance to insecticides, particularly to pyrethroids, has compromised disease control efficacy [6–9]. Field-derived *An. sinensis* populations from Hainan, China, have developed resistance to DDT, cypermethrin, and malathion [10].

Target-site insensitivity and enhanced metabolic detoxification represent two principal mechanisms underlying insecticide resistance in mosquitoes [3,11]. Carboxylesterases, cytochrome P450s, and glutathione S-transferases in mosquitoes are typical enzymes correlated with pyrethroid resistance [11–15]. In recent years, our research has focused on *An. sinensis* to elucidate the mechanisms of insecticide resistance. Multiple gene families in mosquitoes have been confirmed to be involved in insecticide resistance mechanisms [16–21], including cytochromes P450, glutathione S-transferases, ATP-binding cassette transporters, and carboxylesterases. Carboxylesterases, members of the α/β-hydrolase fold superfamily [22], mediate specific hydrolysis of carboxyl ester bonds in organophosphates and pyrethroids [23–25]. For example, carboxylesterase Ha006a facilitates metabolic detoxification of pyrethroids (fenvalerate, λ-cyhalothrin, and deltamethrin) and organophosphates (paraoxon ethyl, profenofos, and chlorpyrifos) in *Helicoverpa armigera* [26]. Similarly, carboxylesterase RpCarE can hydrolyze isoprocarb and cyhalothrin in *Rhopalosiphum padi* [27]. Notably, Boest1 contributes to malathion and bifenthrin detoxification in *Bradysia odoriphaga* [28].

Gene mutation and overexpression represent two primary mechanisms for carboxylesterase-mediated pyrethroid resistance [29]. Specific mutations significantly enhance hydrolysis efficiency: *CarE001C* (H423I/R322L) in *H. armigera* elevates fenvalerate hydrolysis, while *E3* (W251L/F309L) in *Lucilia cuprina* increases permethrin degradation [30,31]. In pyrethroid-resistant *Musca domestica*, *MdαE7* mutations are accompanied by elevated expression and gene amplification, enhancing pyrethroid hydrolysis [32,33]. Overexpression of carboxylesterase genes further contributes to pyrethroid resistance. Upregulated expression of esterase genes in fenvalerate-resistant *H. armigera* midgut/fat body correlates with enhanced activity [34]. Constitutive/inductive *PxαE14* overexpression facilitates multi-pyrethroid detoxification in *Plutella xylostella* [35]. Studies have revealed significantly elevated carboxylesterase expression in pyrethroid-resistant mosquito populations, exemplified by upregulated *CPIJ018231*, *CPIJ018232*, and *CPIJ018233* in permethrin-resistant *Culex quinquefasciatus* [36], overexpressed *CCEae3a* in deltamethrin-resistant *Aedes aegypti* [37], and increased *COEAE1F* expression in deltamethrin-resistant *Anopheles gambiae* [38]. Moreover, increased esterase activity was observed in deltamethrin-resistant *Cx. pipiens pallens* and *Aedes aegypti* strains [39,40]. Correlative analysis by Chang et al. identified carboxylesterases as contributors to deltamethrin/permethrin resistance in *An. sinensis* [3], while Wu et al. found upregulated carboxylesterase genes in deltamethrin-resistant strains [16]. Collectively, these findings establish a correlation between pyrethroid resistance and carboxylesterase overexpression in mosquitoes, suggesting potential involvement of *An. sinensis* carboxylesterases in pyrethroid resistance. However, identification and functional analyses of specific pyrethroid resistance-associated carboxylesterase genes in *An. sinensis* remain limited.

In this study, comparative transcriptomics of susceptible and resistant *An. sinensis* strains were utilized to identify CCE genes exhibiting upregulated expression. Based on these findings and prior evidence of CCE-mediated resistance mechanisms, we identified

candidate CCE genes of *An. sinensis* associated with deltamethrin resistance. Functionally, qRT-PCR was performed to confirm five upregulated genes across different resistant strains, and their causal roles in resistance were further elucidated through RNA interference (RNAi) and bioassays. These results enhance our mechanistic understanding of CCE-facilitated pyrethroid resistance and provide a theoretical basis for optimizing mosquito control strategies.

2. Materials and Methods

2.1. Mosquito Strains

The susceptible strain WX-LS, originally sourced from Wuxi, Jiangsu Province, China, has been maintained at the Institute of Entomology and Molecular Biology, Chongqing Normal University, China, under insecticide-free conditions. The resistant strain WX-LR was derived from WX-LS through deltamethrin selection. The pyrethroid-resistant strains CQ-LR, AH-LR, and YN-LR were established as near-isogenic lines via backcrossing for over ten generations between laboratory-susceptible strains and resistant populations collected from Chongqing, Anhui, and Yunnan, China, respectively. All five *An. sinensis* strains were reared separately under controlled conditions of 27 ± 1 °C and 70% ± 10% relative humidity. The susceptibility of adult female mosquitoes was evaluated three days post-emergence using the WHO standard bioassay protocol with filter papers impregnated with a diagnostic concentration of 0.05% deltamethrin (CAS: 52918-63-5, Sigma-Aldrich, St. Louis, MO, USA) [5]. The LC_{50} of deltamethrin for the laboratory-susceptible strain was 0.0067 mg/L. After a one-hour exposure to 0.05% deltamethrin-treated filter papers, the laboratory-resistant strains exhibited mortality rates of 14.29% for WX-LR, 13.33% for YN-LR, 17.14% for CQ-LR, and 11.43% for AH-LR following a 24 h recovery period.

2.2. Identification of Pyrethroid Resistance-Associated CCEs Using RNA-Seq

Transcriptomic analysis was performed on *An. sinensis*, encompassing both susceptible and resistant strains. Specimens were procured from diverse developmental phases, including fourth-instar larvae, female and male pupae, and adult females and males. Additionally, tissues from adult mosquitoes, namely the antennae, cuticle, fat body, Malpighian tubules, and midgut, were also collected. For blood-fed females, samples were gathered at 1, 3, and 12 h post-feeding. Three biological replicates were used to ensure reliability. The sequencing was conducted by Novogene Co., Ltd. (Beijing, China) using the Illumina HiSeq™ 2000 platform (Illumina, San Diego, CA, USA). Subsequently, following previously described methods [41,42], the data were analyzed by the Institute of Entomology and Molecular Biology, Chongqing Normal University. CCE genes associated with pyrethroid resistance in *An. sinensis* were identified based on the transcriptional profiles of distinct developmental stages, tissues, and blood-fed females from both deltamethrin-resistant (CQ-LR) and susceptible (WX-LS) strains. The sequencing reads were aligned to the *An. sinensis* genome using TopHat [43]. The expression levels of the CCE genes were quantified in FPKM (fragments per kilobase of transcript per million mapped reads) by employing Cufflinks [44]. Additionally, Cufflinks was utilized to assess the differential expression of transcripts between the resistant and susceptible strains. A gene was deemed significantly differentially expressed if it satisfied the following criteria: $|\log_2$ (fold change in FPKM)$| \geq 1$ and p-value ≤ 0.05. These thresholds enabled the identification of potential CCE genes implicated in pyrethroid resistance. The differential expression patterns of the CCE genes were visualized using GraphPad Prism 10.0 software (GraphPad Software Inc., San Diego, CA, USA).

2.3. qRT-PCR Verification of Pyrethroid Resistance-Associated CCE Genes

Previous research has demonstrated that five genes (*CarE001G*, *Mdα E7*, *Pxαe28*, and *Ldace1*) in *H. armigera*, *M. domestica*, *P. xylostella*, and *Leptinotarsa decemlineata* are involved in insecticide metabolism or are resistance-conferring [45–48]. Their corresponding homologs in *An. sinensis* have been identified as *AsBe4*, *AsAe4*, *AsUn4*, and *AsAce1*. These genes, along with the genes identified through RNA-seq analysis in this study, were selected as candidate target genes for validation via qRT-PCR. Total RNA was extracted using the Trizol Reagent Kit (Thermo Fisher Scientific Inc., Waltham, MA, USA) in accordance with the manufacturer's instructions. The cDNA template was synthesized from the total RNA using the PrimeScript™ RT Reagent Kit with gDNA Eraser (TaKaRa, Biotech Co., Ltd., Dalian, China). The qRT-PCR assays were conducted using an iTaq™ SYBR® Green Supermix (Bio-Rad, Hercules, CA, USA). Each treatment was replicated three times biologically. The *ribosomal protein L49* (*RPL49*) of *An. sinensis* was employed as the reference gene. The primers for qRT-PCR were designed using Primer Premier 5.0 (Supplementary Files S1). The relative expression levels were determined using the $2^{-\Delta\Delta Ct}$ method [49].

2.4. RNAI-Based Validation of Pyrethroid Resistance-Related CCE Genes

To elucidate the involvement of specific CCE genes in the pyrethroid resistance of *An. sinensis*, RNA interference (RNAi) was employed in conjunction with deltamethrin bioassays on female mosquitoes from the WX-LR strain. Primers for dsRNA synthesis were designed based on the cDNA sequences of the five CCE genes and *enhanced green fluorescent protein* (EGFP) using Primer Premier 5.0 (Supplementary Files S1). The dsRNA fragments were synthesized in vitro using the T7 RiboMAX Express RNAi System (Promega (Beijing) Biotech Co., Ltd., Beijing, China), with *EGFP* dsRNA serving as a negative control. The dsRNA concentrations were measured using Nano-Drop™ 1000 (Thermo Fisher Scientific Inc., Waltham, MA, USA). An 800 ng amount of the dsRNA fragments was injected into the hemocoel at the abdomen region between the second and third segment of the pupa at 10 h post-pupation of *An. sinensis*. The efficiency of gene silencing by dsRNA was evaluated using qRT-PCR in female adult mosquitoes three hours after emergence. The bioassay procedure adhered to the WHO standard insecticide susceptibility testing protocol [5]. Vigorous female adults were transferred to holding tubes for 1 h pre-exposure. Moribund/dead mosquitoes were discarded. Paper impregnated with 0.05% deltamethrin was secured to the inner surface of the exposure tubes. Following retraction of the sliding mesh partition, mosquitoes were gently blown into exposure tubes. Mosquitoes were exposed for 1 h with knockdown counts recorded at 10 min intervals. Survivors were transferred to holding tubes with 10% sugar water for 24 h recovery prior to mortality assessment. A mosquito was classified as dead or knocked down if it was immobile or unable to stand or take off. The knockdown curves of each treatment were fitted and the half-knockdown time (KT_{50}) was calculated using GraphPad Prism 10.0 software.

2.5. Statistical Analysis

In this study, the disparities in gene expression levels and biological assay data across diverse treatments were analyzed. One-way analysis of variance (ANOVA) was initially conducted to assess the overall differences in expression levels among the distinct populations or developmental stages of *An. sinensis*, and Tukey's multiple comparison tests were employed to discern the pairwise differences. The difference in knockdown rates between the control and treatment groups during the 1 h exposure to deltamethrin was analyzed using the Log-rank (Mantel–Cox) test. These analyses were performed using GraphPad Prism version 10.0 software.

3. Results and Discussion

3.1. Identification of Carboxylesterase Genes Associated with Pyrethroid Resistance in An. sinensis

3.1.1. Identification of CCEs via Comparative Transcriptomics

The overexpression of esterase genes in mosquitoes facilitates their insecticide resistance [12,36,37,50,51]. In this study, RNA-seq was employed to profile differential expression of *CCE* genes in *An. sinensis* across five developmental stages, five tissue types, and three post-blood meal time points in paired deltamethrin-resistant (CQ-LR) and susceptible (WX-LS) strains. Developmental stage-specific profiling revealed that *AsAe10* and *AsBe3* exhibited significantly upregulated expression in four stages, excluding the fourth instar of larva (Figure 1a). The majority of overexpressed genes belonged to the *Ae* or *Be* classes, including *AsAe9*. In contrast, the measured *CCE* genes in the fourth larva exhibited significant downregulation in expression levels (Figure 1a). Tissue-specific profiling identified significantly upregulated *CCE* genes in *An. sinensis*: five in both the antennae and cuticle, four in the Malpighian tubules and midgut, and three in the fat body (Figure 1b). *AsAe10* exhibited significant overexpression across all five examined tissues, whereas *AsBe2* showed significant upregulation in three tissues (Figure 1b). Widespread upregulation was also observed for *Ae*-class genes, with *AsAe9* showing specific overexpression in antennae. The newly identified *AsAce2* exhibited upregulated expression in the fat body (Figure 1b). Specific temporal patterns of *CCE* gene upregulation emerged post-blood meal: four genes at 1 h, twelve genes at 3 h, and three genes at 12 h (Figure 1c). *Asle1* and *AsAe10* exhibited significant upregulation across two distinct temporal intervals. *AsAce2* and *AsUn5* had significant overexpression at 3 h after blood meal (Figure 1c). As shown in Figure 1, *AsAe9* and *AsAe10* exhibited significant upregulation across the three experiments, and *AsAce2* was overexpressed in both tissue-specific and post-blood meal assays. Notably, the homolog of *AsAe10* in *H. armigera*, *CarE001H*, has been functionally validated as a key gene involved in pyrethroid detoxification [52]. Similarly, the ortholog of *AsAce2* in *P. xylostella* (*Pxαe18*) plays an analogous role in detoxification processes [46]. Therefore, we preliminarily identify *AsAe9, AsAe10*, *AsAce2*, and *AsUn5* as genes associated with deltamethrin resistance in *An. sinensis*.

Developmental stage- and tissue-specific upregulation of carboxylesterase genes is involved in resistance mechanisms in insects. For instance, in a pyrethroid-resistant strain of *Cx. quinquefasciatus*, two esterase genes exhibit upregulated expression levels in both larval and adult stages relative to the susceptible strain [15]. Conversely, another study revealed that resistant larvae—but not adults—of the same species demonstrate significant overexpression of three distinct carboxylesterase genes [36]. This study proposes that constitutive upregulation of *CCE* genes in pupal and adult stages of the resistant *An. sinensis* strain reflects stage-specific selection pressure. Given that insecticide treatments target advanced developmental stages, these alterations in expression profiles could be associated with adaptive genetic responses to insecticide exposure, thereby driving resistance evolution. The midgut and fat body are typically organs for tissue-specific carboxylesterase expression, which is critical for metabolic detoxification [53–58]. In *P. xylostella*, the fipronil resistance-associated carboxylesterase genes *Pxae22* and *Pxae31* exhibit upregulated expression in fourth-instar larvae and the midgut [56]. Wu et al. documented 1.9-fold elevated esterase activity in a fenvalerate-resistant *H. armigera* strain (1690-fold resistance), concomitant with 2-to-90-fold upregulated expression of four esterase genes in the midgut and fat body tissues relative to a susceptible strain [34]. Our earlier investigation revealed spatiotemporal overexpression patterns of pyrethroid resistance-associated *P450* genes in *An. sinensis*: elevated in resistant strains specifically within pupae/adults, the adult midgut/cuticle, and at 1- and 3 h post-blood meal [57]. Malpighian tubule-specific gene expression in mosquitoes facilitates dual roles in blood meal digestion and xenobiotic

detoxification [58]. Reports indicate that *Anopheles* with pyrethroid resistance mediated by *kdr* mutations or *P450* overexpression exhibit altered blood-feeding capabilities [59,60]. Conversely, following a single blood meal, pyrethroid-resistant *An. funestus* exhibits significantly enhanced resistance to permethrin [61]. Researchers consider that insecticide detoxification mechanisms involved in resistance may be activated by a prior blood meal, thereby enhancing the resistance phenotype before insecticide exposure. Moreover, the blood meal sustains DDT and permethrin resistance during mosquito development, likely due to blood meal-induced elevation in glutathione S-transferase activity [62]. We hypothesize that, compared to the susceptible strain, significant blood-induced upregulation of carboxylesterase expression in the resistant mosquito strain may enhance blood meal metabolism. This could boost their fitness and stabilize insecticide resistance inheritance. Consequently, we measured carboxylesterase expression post-blood feeding in both resistant and susceptible mosquitoes in this study. Our integrated transcriptomic approach, validated in *An. gambiae*, *An. sinensis*, and *M. domestica* [38,63,64], consistently identifies carboxylesterase genes as important actors in pyrethroid resistance. Notably, transcriptomic analysis of the deltamethrin-resistant *An. sinensis* strain established by Zhou et al. did not detect overexpressed *CCE* genes [64]. This discrepancy with our findings is likely attributable to the strain's geographic origin (Hubei, China) and differential resistance gene dominance.

3.1.2. Validation of Differential Expression for CCEs via qRT-PCR

Many studies on mosquito resistance mechanisms validate the necessity for qRT-PCR confirmation of expression differences in candidate genes between resistant and susceptible strains [15,16,57]. qRT-PCR has been previously employed to identify pyrethroid resistance-associated *P450* and *GST* genes in *An. sinensis* [19,57]. The differential expression of eight genes—comprising four previously reported genes (*AsAe4*, *AsAce1*, *AsBe4*, *AsUn4*) and four genes newly identified in this study (*AsAe9*, *AsAe10*, *AsAce2*, *AsUn5*)—was quantified via qRT-PCR in deltamethrin-resistant and susceptible *An. sinensis* strains.

While *AsAe4* showed no significant upregulation in four *An. sinensis*-resistant strains compared with the susceptible WX-LS strain (Figure 2), its homolog *MdαE7* in resistant *M. domestica* is involved in permethrin detoxification [47,65]. Additionally, the *P. xylostella* homolog *PxEst-6* exhibits competence in binding and metabolizing bifenthrin, cyfluthrin, cypermethrin, and λ-cyhalothrin [66]. This functional difference likely reflects species-specific variations in pyrethroid detoxification metabolism mediated by the target gene. Significant upregulation of *AsAe9*, *AsAce1*, and *AsBe4* was detected in two *An. sinensis* resistant strains (Figure 2). *Ace1* shows upregulated expression in a pyrethroid-treated *L. decemlineata* strain, which may be implicated in insecticide resistance mechanisms [45]. A homolog of *AsBe4* (carboxylesterase *CarE001G* in *H. armigera*) exhibits metabolic activity toward β-cypermethrin, λ-cyhalothrin, and fenvalerate in vitro [48]. *AsBe4* (corresponding to *Bemisia tabaci* carboxylesterase *BTbe5*) demonstrates upregulation following imidacloprid exposure, and RNAi-mediated gene silence functionally validates its role in the insecticidal mechanism [67]. Elevated relative expression of *AsAe10* was observed across all four resistant strains, especially in the WX-LR strain (Figure 2), consistent with RNA-seq data from this study. The homolog of *AsAe10* in *H. armigera* (*CarE001H*) has been functionally validated as a key metabolic detoxification gene involved in pyrethroid catabolism [52]. *CCEae3A* (homolog of *Ae10*) exhibits significant upregulation in the *Ae. aegypti* temephos-resistant strain, and its overexpression in transgenic *An. gambiae* confers resistance to α-cypermethrin [51]. *AsAce2* exhibited significantly upregulated expression in four *An. sinensis* resistant strains (Figure 2). Its homolog *Pxae18* is overexpressed in chlorpyrifos-resistant *P. xylostella*, with RNAi confirming resistance involvement [46].

AsUn4 and AsUn5 exhibited significant upregulation in the AH-LR and WX-LR strains, respectively (Figure 2). The homolog of *AsUn4* in *P. xylostella* (*PxαE28*) exhibits significant upregulation in chlorpyrifos-resistant strains, and RNAi-mediated gene silencing restores insecticide susceptibility, confirming its functional role in resistance [46]. Consequently, based on comparative transcriptomics and qRT-PCR validation, *AsAe9*, *AsAe10*, *AsAce1*, *AsAce2*, and *AsBe4* were selected as deltamethrin resistance-associated carboxylesterase genes for subsequent functional analysis.




Figure 1. RNA sequencing analyses of differentially expressed *CCE* genes in deltamethrin-resistant *An. sinensis* strain (CQ-LR) versus susceptible strain (WX-LS). (**a**) Developmental stage-specific expression; (**b**) tissue-specific expression; (**c**) temporal dynamics of gene expression post-blood meal. Vertical dotted lines mark the ± 1 of $\log_2$ (fold changes in FPKM value) values on the *x*-axis, and horizontal dotted lines denote the *p*-value = 0.05 of $-\log_{10}$ (*p*-value) on the *y*-axis. Criteria for significantly upregulated genes: $|\log_2 (\text{fold change in FPKM})| \geq 1$ and *p*-value ≤ 0.05.




Figure 2. Validation of expression levels for eight selected carboxylesterase genes in deltamethrin-resistant (WX-LR, AH-LR, YN-LR, CQ-LR) and susceptible (WX-LS) *An. sinensis* strains. (**a**) *AsAe4*; (**b**) *AsAe9*; (**c**) *AsAe10*; (**d**) *AsAce1*; (**e**) *AsAce2*; (**f**) *AsBe4*; (**g**) *AsUn4*; (**h**) *AsUn5*. The genes' relative expression levels are presented as the mean ± SD (standard deviation) from three biological replicates with three technical replicates each. Columns marked with the different lowercase letters indicate significantly differential expression across the strains ($p < 0.05$).

3.2. Function Analysis of Carboxylesterase Genes in Pyrethroid Resistance of An. sinensis

3.2.1. Determination of Time Points for CCE Silencing

Functional validation was performed through RNAi silencing to confirm the involvement of *AsAe9*, *AsAe10*, *AsAce1*, *AsAce2*, and *AsBe4* in pyrethroid resistance in *An. sinensis*. To determine the optimal developmental stage for dsRNA injection, the specific expression profiles of these five genes in pupae and adults in both pyrethroid-resistant (WX-LR) and susceptible (WX-LS) strains were acquired using qRT-PCR (Figure 3). All five genes exhibited expression through the pupal and adult stages, but had various expression levels across the different periods. In the WX-LS strain, *AsAe9*, *AsAce1*, and *AsAce2* exhibited maximal relative expression levels at P_{last} (terminal pupal stage), whereas *AsAe10* and *AsBe4* demonstrated divergent peak timing (Figure 3). In the WX-LR strain, three target genes (*AsAe10*, *AsAce2*, *AsBe4*) achieved peak expression at A_{72} (adults 72 h post-emergence). Compared to the susceptible strain, all five genes had significant upregulation in at least one adult stage of WX-LR ($p < 0.05$), with three genes concurrently overexpressed during pupal development (Figure 3). The five genes in both strains exhibited the lowest expression levels at the P_{10} stage (10 h post-pupation), with no significantly differential expression between strains ($p > 0.05$, Figure 3). This expression convergence established P_{10} as the optimal developmental stage for dsRNA microinjection in subsequent RNAi-mediated functional validation.

3.2.2. Functional Analysis of CCE Genes in Pyrethroid Resistance

RNAi experiments targeting the genes *AsAe9*, *AsAe10*, *AsAce1*, *AsAce2*, and *AsBe4* in the deltamethrin-resistant strain WX-LR revealed that the relative expression levels of the first four genes were significantly reduced ($p < 0.01$; Figure 4A–D), with silencing efficiencies of 79.8%, 51.8%, 55.4%, and 66.3%, respectively. In contrast, silencing of *AsBe4* did not result in a significant reduction in expression, showing a 30.4% decrease (Figure 4E). In the subsequent bioassay, silencing of *AsAe10*, *AsAce1*, *AsAce2*, and *AsBe4* resulted in significantly shortened knockdown times ($p < 0.05$; Figure 4b–e) in resistant *An. sinensis* following 1 h of exposure to 0.05% deltamethrin. The most significant reduction in knockdown time occurred in the *AsAe10*-silenced group, followed by *AsBe4* (Figure 4b–e). Integrated analysis of the knockdown data revealed that silencing each of the five genes resulted in a reduction in KT_{50} in resistant mosquitoes. Compared to the control groups, gene-silenced *An. sinensis* exhibited reductions in KT_{50} values: *AsAe9* and *AsAe10* decreased from a common control value of 55.12 min (95% fiducial limit [FL]: 53.73–56.66) to 51.09 min (95% FL: 50.36–51.82) and 43.83 min (95% FL: 40.91–46.74), while *AsAce1*, *AsAce2*, and *AsBe4*—sharing a common control value of 57.43 min (95% FL: 56.77–58.15)—declined to 49.09 min (95% FL: 46.27–52.08), 50.08 min (95% FL: 46.16–54.26), and 48.02 min (95% FL: 43.63–52.23), respectively. The non-overlapping 95% fiducial limits between the silenced groups and the control groups indicate a statistically significant reduction in knockdown time. The most significant decrease in KT_{50} was observed in the *AsAe10*-silenced group, which is consistent with the findings described above. RNAi-mediated silencing of five carboxylesterase genes in the pyrethroid-resistant strain enhanced deltamethrin susceptibility, resulting in increased mortality rates of 40.3% (*AsAe9*), 25.3% (*AsAe10*), 9.5% (*AsAce1*), 24.9% (*AsAce2*), and 14.2% (*AsBe4*) after 1 h exposure to deltamethrin followed by 24 h recovery (Figure 4I–V). Silencing of *AsAe9*, *AsAe10*, and *AsBe4* significantly compromised the resistance phenotype, with *AsAe10* exhibiting the most pronounced effect ($p < 0.05$, Figure 4I–V). In conclusion, among the five genes investigated, although silencing of several genes led to reduced knockdown time and increased mortality in the resistant strain after deltamethrin exposure, only silencing of *AsAe10* and *AsBe4* resulted in statistically significant changes in both endpoints. Notably, *AsAe10* silencing produced the most substantial phenotypic alteration. Furthermore, the expression of *AsAe10* was

significantly upregulated in the resistant strain, suggesting that it may be strongly associated with pyrethroid resistance in *An. sinensis*. Its overexpression could potentially enhance the metabolic detoxification of deltamethrin in mosquitoes. Interestingly, *AsBe4* silencing also significantly compromised resistance despite its relatively lower silencing efficiency, suggesting its potentially critical role in the resistance phenotype.




Figure 3. Developmental stage-specific expression profiling of five carboxylesterase genes in deltamethrin-resistant and susceptible *An. sinensis* strains. (**a**): *AsAe9*; (**b**) *AsAe10*; (**c**) *AsBe4*; (**d**) *AsAce1*; (**e**) *AsAce2*. The relative expression levels of genes were normalized to 0 h post-pupation of WX-LS, and are shown as the mean ± SD from three biological replicates in qRT-PCR analysis. Columns marked with homogeneous uppercase (for WX-LS) or lowercase (for WX-LR) letters indicate no significant difference in expression across developmental stages ($p > 0.05$). Asterisks indicate significant inter-strain differential expression at matched timepoints (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; ****: $p < 0.0001$). Pupal phases: P_0 (0 h post-pupation), P_{10} (10 h post-pupation), P_{20} (20 h post-pupation), P_{30} (30 h post-pupation), P_{last} (terminal pupal stage). Adult stages: A_0 (0 h post-emergence, newly emerged adults), A_3 (3 h post-emergence), A_6 (6 h post-emergence), A_9 (9 h post-emergence), A_{24} (24 h post-emergence), A_{72} (72 h post-emergence).




Figure 4. Functional validation of the carboxylesterase genes *AsAe9* (**A**,**a**,**I**), *AsAe10* (**B**,**b**,**II**), *AsAce1* (**C**,**c**,**III**), *AsAce2* (**D**,**d**,**IV**), and *AsBe4* (**E**,**e**,**V**) of *An. sinensis* in its deltamethrin resistance. (**A**–**E**) The expression levels of *AsAe9*, *AsAe10*, *AsAce1*, *AsAce2*, and *AsBe4*post-dsRNA injection (qRT-PCR; mean ± SD). (**a**–**e**) Time–knockdown profiles of *An. sinensis* during 0.05% deltamethrin exposure. (**I**–**V**) Mortality of *An. sinensis* after 24 h recovery following 0.05% deltamethrin exposure (mean ± SD). The groups injected with dsRNA of enhanced green fluorescent protein (*dsEGFP*) were negative controls. The increases in knockdown number were recorded at 10 min intervals for 1 h. The asterisks indicate significant differences between groups (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; ****: $p < 0.0001$).

Insect esterases can hydrolyze pyrethroids, thereby contributing to detoxification. Enhanced resistance phenotypes correlate with gene mutations that elevate both the expression and hydrolytic activity of these enzymes [68,69]. In *An. gambiae*, transcriptional upregulation of carboxylesterase genes significantly correlates with deltamethrin resistance [38]. In the deltamethrin-resistant *Cx. pipiens pallens*, carboxylesterase activity is associated with resistance, with resistance declining concomitantly upon relaxation of deltamethrin selection pressure [39]. In *Ae. aegypti*, resistance to permethrin and deltamethrin has been linked to elevated carboxylesterase activity [40,70], with *CCEae3a* overexpression correlating with deltamethrin resistance [37]. Germline transformation of *Aedes aegypti* with the *An. gambiae CCEae3a* gene—overexpressed in the donor *Ae. Aegypti* strain—elevates enzyme activity and significantly increases α-cypermethrin resistance [51]. In the permethrin-resistant strain of *Cx. quinquefasciatus*, *α-esterase* and *esterase B1* exhibit significant upregulation, respectively, compared to the susceptible stain [15]. Concurrently, three carboxylesterase genes show significant overexpression in the resistant strain, collectively indicating enhanced metabolic detoxification capacity [36]. Carboxylesterase upregulation and enhanced activity represent an important pyrethroid resistance mechanism in mosquitoes. This study confirms that the significantly upregulated carboxylesterase genes in resistant *An. sinensis* are involved in deltamethrin resistance. Similar roles of carboxylesterases in organophosphorus insecticide resistance have been demonstrated. In *Ae. aegypti* and *Aedes albopictus*, temephos resistance is involved in increased carboxylesterase activity, with resistant strains exhibiting significant overexpression of *CCEae6a*, and concomitant gene amplification [71,72].

RNAi-mediated silencing provides a robust approach for functionally validating the contribution of upregulated carboxylesterase genes to insecticide resistance, a methodology extensively applied in resistance mechanism research [73]. RNAi-based functional validation has elucidated the roles of cytochrome P450 in *An. gambiae*, the GST in *Ae. aegypti*, and G-protein-coupled receptor (GPCR) genes in *Cx. quinquefasciatus* permethrin resistance [74–76]. In *P. xylostella*, the carboxylesterase genes *PxαE8*, *PxαE14*, and *PxEst-6* exhibit tissue-specific expression in the midgut or Malpighian tubules, and their association with pyrethroid resistance has been validated through RNAi-mediated gene silencing [35,66,77]. Pyrethroid-induced carboxylesterase upregulation (*EoCarE592* in *Ectropis obliqua*; four genes in *Grapholita molesta*) enhances insect susceptibility upon RNAi-mediated gene silencing [78,79]. Additionally, researchers also have considered carboxylesterase-mediated sequestration of organophosphates as an important mechanism for mosquito resistance [80], and this sequestration is closely associated with its overexpression [81]. Currently, novel RNAi-based insecticides have been commercialized for agricultural pest control [82,83]. Multiple studies have confirmed the feasibility of applying RNAi technology for mosquito management [84], with ongoing research focused on identifying mosquito-specific target genes and optimizing dsRNA delivery methods [85,86]. Prospectively, leveraging the insecticide resistance-associated genes identified in *An. sinensis* in this study could enable the development of RNAi biopesticides. These agents would reduce carboxylesterase-mediated metabolism of pyrethroids in mosquitoes, thereby delaying resistance evolution. Furthermore, synergistic application with conventional insecticides could extend their effective lifespan.

This study advances our understanding of pyrethroid resistance mechanisms in *An. sinensis*, and provides a theoretical foundation for optimizing mosquito control strategies. The carboxylesterase genes identified in this research offer molecular tools for resistance management. For *AsAe10*, future studies will bridge mechanism and application. Bidirectional in vivo validation via transgenic *An. sinensis* (overexpression/knockout) will delineate carboxylesterase-mediated resistance pathways, while integrated in vitro model-

ing (molecular docking/eukaryotic expression) will analyze metabolic dynamics. It is also a significant strategy to develop synergistic *dsRNA*–insecticide combinations for delaying vector resistance.

4. Conclusions

This study integrated transcriptomic screening and functional validation to identify carboxylesterase genes involved in deltamethrin resistance in *Anopheles sinensis*. Based on comparative transcriptomic analyses between resistant and susceptible strains, *AsAe9*, *AsAe10*, *AsAce2*, and *AsUn5* were identified as significantly upregulated carboxylesterase genes exhibiting differential spatiotemporal expression patterns and post-blood meal induction profiles. qRT-PCR confirmed significant overexpression of *AsAe10* and *AsAce2* in all four resistant strains, with *AsAe9*, *AsAce1*, and *AsBe4* elevated in two strains. RNA interference targeting the five carboxylesterase genes revealed that silencing of *AsAe10* and *AsBe4* significantly shortened the knockdown time and increased 24 h mortality in the resistant strain upon exposure to 0.05% deltamethrin, with *AsAe10* exhibiting a more pronounced effect. This study identifies these genes as contributors to pyrethroid resistance in *An. sinensis*, provides further insight into the resistance mechanisms, and offers a theoretical basis for refining resistance management strategies against mosquitoes.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/insects16090938/s1. Supplementary File S1: Primer sequences and coding sequences of genes.

Author Contributions: Conceptualization, B.C., Y.W., F.S. and L.Q.; methodology, B.C., Y.W., F.S. and L.Q.; software, Y.W., X.G. and H.Y.; investigation, X.G. and X.C.; data curation, X.G. and X.C.; writing—original draft preparation, Y.W.; writing—review and editing, B.C. and Y.W.; visualization, Y.W., X.G. and H.Y.; supervision, B.C.; project administration, B.C. and Y.W.; funding acquisition, B.C. and Y.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Natural Science Foundation of China (grant number 31672363, 31872262), the National Key Program of Science and Technology Foundation Work of China (grant number 2015FY210300), and the Science and Technology Research Program of Chongqing Municipal Education Commission (grant number KJQN202200516).

Data Availability Statement: The data presented in the study are available in "FigShare" at https://www.doi.org/10.6084/m9.figshare.29975380.

Acknowledgments: The authors would like to express their gratitude to Junfu Chen from Chongqing Normal University for help in the visualization of this study.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Rueda, L.M.; Zhao, T.; MA, Y.J.; Gao, Q.; Ding, Z.G.; Khuntirat, B.; Sattabongkot, J.; Wilkerson, R.C. Updated distribution records of the *Anopheles (Anopheles) hyrcanus* species-group (Diptera: Culicidae) in China. *Zootaxa* **2007**, *1407*, 43–55. [CrossRef]
2. Sinka, M.E.; Bangs, M.J.; Manguin, S.; Rubio-Palis, Y.; Chareonviriyaphap, T.; Coetzee, M.; Mbogo, C.M.; Hemingway, J.; Patil, A.P.; Temperley, W.H.; et al. A global map of dominant malaria vectors. *Parasites Vectors* **2012**, *5*, 69. [CrossRef]
3. Chang, X.; Zhong, D.; Fang, Q.; Hartsel, J.; Zhou, G.; Shi, L.; Fang, F.; Zhu, C.; Yan, G. Multiple resistances and complex mechanisms of *Anopheles sinensis* mosquito: A major obstacle to mosquito-borne diseases control and elimination in China. *PLoS Negl. Trop. Dis.* **2014**, *8*, e2889. [CrossRef]
4. Pan, J.; Zhou, S.; Zheng, X.; Huang, F.; Wang, D.; Shen, Y.; Su, Y.; Zhou, G.; Liu, F.; Jiang, J. Vector capacity of *Anopheles sinensis* in malaria outbreak areas of central China. *Parasites Vectors* **2012**, *5*, 136. [CrossRef]
5. World Health Organization. *Test Procedures for Insecticide Resistance Monitoring in Malaria Vector Mosquitoes*, 2nd ed.; World Health Organization: Geneva, Switzerland, 2016; pp. 6–35. Available online: https://iris.who.int/handle/10665/250677 (accessed on 30 June 2018).

6. Hemingway, J.; Field, L.; Vontas, J. An overview of insecticide resistance. *Science* **2002**, *298*, 96–97. [CrossRef]
7. Zaim, M.; Guillet, P. Alternative insecticides: An urgent need. *Trends Parasitol.* **2002**, *18*, 161–163. [CrossRef] [PubMed]
8. Butler, D. Mosquitoes score in chemical war. *Nature* **2011**, *475*, 19. [CrossRef]
9. Kelvin, A.A. Outbreak of Chikungunya in the Republic of Congo and the global picture. *J. Infect. Dev. Ctries* **2011**, *5*, 441–444. [CrossRef]
10. Sun, D.; Wang, G.; Zeng, L.; Li, S.; He, C.; Hu, X.; Wang, S. Extensive resistance of *Anopheles sinensis* to insecticides in malaria-endemic areas of Hainan Province, China. *Am. J. Trop. Med. Hyg.* **2017**, *97*, 295–298. [CrossRef]
11. Liu, N. Insecticide resistance in mosquitoes: Impact, mechanisms, and research directions. *Annu. Rev. Entomol.* **2015**, *60*, 537–559. [CrossRef] [PubMed]
12. Raymond, M.; Chevillon, C.; Guillemaud, T.; Lenormand, T.; Pasteur, N. An overview of the evolution of overproduced esterases in the mosquito Culex pipiens. *Philos. Trans. R. Soc. B Biol. Sci.* **1998**, *353*, 1707–1711. [CrossRef]
13. Feyereisen, R. Insect cytochrome P450. In *Comprehensive Molecular Insect Science*; Gilbert, L., Iatrou, K., Gill, S.S., Eds.; Elsevier B.V.: Amsterdam, The Netherlands, 2005; Volume 4, pp. 1–77.
14. Marcombe, S.; Mathieu, R.B.; Pocquet, N.; Riaz, M.-A.; Poupardin, R.; Sélior, S.; Darriet, F.; Reynaud, S.; Yébakima, A.; Corbel, V.; et al. Insecticide resistance in the dengue vector *Aedes aegypti* from Martinique: Distribution, mechanisms and relations with environmental factors. *PLoS ONE* **2012**, *7*, e30989. [CrossRef] [PubMed]
15. Ramkumar, G.; Muthusamy, R.; Narayanan, M.; Shivakumar, M.S.; Kweka, E.J. Overexpression of cytochrome P450 and esterase genes involved in permethrin resistance in larvae and adults of *Culex quinquefasciatus*. *Parasitol. Res.* **2023**, *122*, 3205–3212. [CrossRef]
16. Wu, X.; Xu, B.; Si, F.; Li, J.; Yan, Z.; Yan, Z.; He, X.; Chen, B. Identification of carboxylesterase genes associated with pyrethroid resistance in the malaria vector *Anopheles sinensis* (Diptera: Culicidae). *Pest Manag. Sci.* **2018**, *74*, 159–169. [CrossRef]
17. Yan, Z.; He, Z.; Yan, Z.; Si, F.; Zhou, Y.; Chen, B. Genome-wide and expression-profiling analyses suggest the main cytochrome P450 genes related to pyrethroid resistance in the malaria vector, *Anopheles sinensis* (Diptera Culicidae). *Pest. Manag. Sci.* **2018**, *74*, 1810–1820. [CrossRef]
18. He, Q.; Yan, Z.; Si, F.; Zhou, Y.; Fu, W.; Chen, B. ATP-binding cassette (ABC) transporter genes involved in pyrethroid resistance in the malaria vector *Anopheles sinensis*: Genome-wide identification, characteristics, phylogenetics, and expression profile. *Int. J. Mol. Sci.* **2019**, *20*, 1049. [CrossRef]
19. Tao, F.; Si, F.; Hong, R.; He, X.; Li, X.; Qiao, L.; He, Z.; Yan, Z.; He, S.; Chen, B. Glutathione S-transferase (GST) genes and their function associated with pyrethroid resistance in the malaria vector *Anopheles sinensis*. *Pest Manag. Sci.* **2022**, *78*, 4127–4139. [CrossRef]
20. Si, F.; Qiao, L.; He, Q.; Zhou, Y.; Yan, Z.; Chen, B. HSP superfamily of genes in the malaria vector *Anopheles sinensis*: Diversity, phylogenetics and association with pyrethroid resistance. *Malar. J.* **2019**, *18*, 132. [CrossRef]
21. Liu, B.; Qiao, L.; He, Q.; Zhou, Y.; Ren, S.; Chen, B. Genome-wide identification, characterization and evolution of cuticular protein genes in the malaria vector *Anopheles sinensis* (Diptera: Culicidae). *Insect. Sci.* **2018**, *25*, 739–750. [CrossRef] [PubMed]
22. Cygler, M.; Schrag, J.D.; Sussman, J.L.; Harel, M.; Silman, I.; Gentry, M.K.; Doctor, B.P. Relationship between sequence conservation and three-dimensional structure in a large family of esterases, lipases, and related proteins. *Protein Sci.* **1993**, *2*, 366–382. [CrossRef] [PubMed]
23. Huang, J.; Qiao, C. Mechanism and application of insect detoxification enzymes in bioremediation of pesticide contamination. *J. Agro-Environ. Sci.* **2002**, *21*, 285–287. Available online: https://www.aes.org.cn/nyhjkxxb/ch/reader/view_abstract.aspx?file_no=8574&flag=1 (accessed on 30 June 2002).
24. Oakeshott, J.G.; Devonshire, A.L.; Claudianos, C.; Sutherland, T.D.; Horne, I.; Campbell, P.M.; Ollis, D.L.; Russell, R.J. Comparing the organophosphorus and carbamate insecticide resistance mutations in cholin-and carboxyl-esterases. *Chem. Biol. Interact.* **2005**, *157*, 269–275. [CrossRef]
25. Zhan, H.; Huang, Y.; Lin, Z.; Bhatt, P.; Chen, S. New insights into the microbial degradation and catalytic mechanism of synthetic pyrethroids. *Environ. Res.* **2020**, *182*, 109138. [CrossRef] [PubMed]
26. Kaur, H.; Rode, S.; Lonare, S.; Demiwal, P.; Narasimhappa, P.; Arun, E.; Kumar, R.; Das, J.; Ramamurthy, P.C.; Sircar, D.; et al. Heterologous expression, biochemical characterization and prospects for insecticide biosensing potential of carboxylesterase Ha006a from *Helicoverpa armigera*. *Pestic. Biochem. Physiol.* **2024**, *200*, 105844. [CrossRef] [PubMed]
27. Wang, K.; Huang, Y.; Li, X.; Chen, M. Functional analysis of a carboxylesterase gene associated with isoprocarb and cyhalothrin resistance in *Rhopalosiphum padi* (L.). *Front. Physiol.* **2018**, *9*, 992. [CrossRef] [PubMed]
28. Ding, Q.; Xu, X.; Sang, Z.; Wang, R.; Ullah, F.; Gao, X.; Song, D. Characterization of the insecticide detoxification carboxylesterase *Boest1* from *Bradysia odoriphaga* Yang et Zhang (Diptera: Sciaridae). *Pest Manag. Sci.* **2022**, *78*, 591–602. [CrossRef]
29. Zhang, Y.; Ma, X.; Han, Y.; Wang, L.; Liu, Z.; Guo, H.; Fang, J. Transcript-level analysis of detoxification gene mutation-mediated chlorpyrifos resistance in *Laodelphax striatellus* (Hemiptera: Delphacidae). *J. Econ. Entomol.* **2019**, *112*, 1285–1291. [CrossRef]

30. Xu, J.; Chang, Y.; Lu, M.; Tie, Y.; Dong, Y.; Chen, G.; Ma, Z.; Liu, X.; Li, Y. Two single mutations in carboxylesterase 001C improve fenvalerate hydrolase activity in *Helicoverpa armigera*. *Pestic. Biochem. Physiol.* **2021**, *179*, 104969. [CrossRef]
31. Heidari, R.; Devonshire, A.L.; Campbell, B.E.; Dorrian, S.J.; Oakeshott, J.G.; Russell, R.J. Hydrolysis of pyrethroids by carboxylesterases from *Lucilia cuprina* and *Drosophila melanogaster* with active sites modified by in vitro mutagenesis. *Insect Biochem. Mol. Biol.* **2005**, *35*, 597–609. [CrossRef] [PubMed]
32. Feng, X. Functional Characterization of Carboxylesterases in Pyrethroid Resistant House Fly, Musca Domestica. PhD Thesis, Auburn University, Auburn, AL, USA, 2018. Available online: https://www.proquest.com/openview/0d65cf2c4f2c2640ed6c3a2986ef362a/1?cbl=18750&diss=y&pq-origsite=gscholar (accessed on 4 August 2018).
33. Zhang, L.; Shi, J.; Shi, X.; Liang, P.; Gao, J.; Gao, X. Quantitative and qualitative changes of the carboxylesterase associated with beta-cypermethrin resistance in the housefly, *Musca domestica* (Diptera: Muscidae). *Comp. Biochem. Physiol. B.* **2010**, *156*, 6–11. [CrossRef]
34. Wu, S.; Yang, Y.; Yuan, G.; Campbell, P.M.; Teese, M.G.; Russell, R.J.; Oakeshott, J.G.; Wu, Y. Overexpressed esterases in a fenvalerate resistant strain of the cotton bollworm. *Helicoverpa armigera. Insect Biochem. Mol. Biol.* **2011**, *41*, 14–21. [CrossRef] [PubMed]
35. Li, R.; Zhu, B.; Hu, X.; Shi, X.; Qi, L.; Liang, P.; Gao, X. Overexpression of *PxαE14* contributing to detoxification of multiple insecticides in *Plutella xylostella* (L.). *J. Agric. Food Chem.* **2022**, *70*, 5794–5804. [CrossRef]
36. Gong, Y.; Li, M.; Li, T.; Liu, N. Molecular and functional characterization of three novel carboxylesterases in the detoxification of permethrin in the mosquito, *Culex quinquefasciatus*. *Insect Sci.* **2022**, *29*, 199–214. [CrossRef]
37. Goindin, D.; Delannay, C.; Gelasse, A.; Ramdini, C.; Gaude, T.; Faucon, F.; David, J.-P.; Gustave, J.; Vega-Rua, A.; Fouque, F. Levels of insecticide resistance to deltamethrin, malathion, and temephos, and associated mechanisms in *Aedes aegypti* mosquitoes from the Guadeloupe and Saint Martin islands (French West Indies). *Infect. Dis. Poverty* **2017**, *6*, 38. [CrossRef] [PubMed]
38. Oumbouke, W.A.; Pignatelli, P.; Barreaux, A.M.G.; Tia, I.Z.; Koffi, A.A.; Alou, L.P.A.; Sternberg, E.D.; Thomas, M.B.; Weetman, D.; 'Guessan, R.N. Fine scale spatial investigation of multiple insecticide resistance and underlying target-site and metabolic mechanisms in *Anopheles gambiae* in central Côte d'Ivoire. *Sci. Rep.* **2020**, *10*, 15066. [CrossRef] [PubMed]
39. Shi, L.; Hu, H.; Ma, K.; Zhou, D.; Yu, J.; Zhong, D.; Fang, F.; Chang, X.; Hu, S.; Zou, F.; et al. Development of resistance to pyrethroid in *Culex pipiens pallens* population under different insecticide selection pressures. *PLoS Negl. Trop. Dis.* **2015**, *9*, e0003928. [CrossRef]
40. Aponte, H.A.; Penilla, R.; Dzul-Manzanilla, F.; Che-Mendoza, A.; López, A.D.; Solis, F.; Manrique-Saide, P.; Ranson, H.; Lenhart, A.; McCall, P.J.; et al. The pyrethroid resistance status and mechanisms in *Aedes aegypti* from the Guerrero state, Mexico. *Pestic. Biochem. Physiol.* **2013**, *107*, 226–234. [CrossRef]
41. Chen, B.; Zhang, Y.; He, Z.; Li, W.; Si, F.; Tang, Y.; He, Q.; Qiao, L.; Yan, Z.; Fu, W.; et al. De novo transcriptome sequencing and sequence analysis of the malaria vector *Anopheles sinensis* (Diptera: Culicidae). *Paras. Vector.* **2014**, *7*, 314. [CrossRef]
42. Zhang, Y.; Lan, Y.; Chen, B. ASDB: A comprehensive omics database for *Anopheles sinensis*. *Genomics* **2021**, *113*, 976–982. [CrossRef]
43. Trapnell, C.; Pachter, L.; Salzberg, S.L. TopHat: Discovering splice junctions with RNA-Seq. *Bioinformatics* **2009**, *25*, 1105–1111. [CrossRef]
44. Trapnell, C.; Williams, B.A.; Pertea, G.; Mortazavi, A.; Kwan, G.; van Baren, M.J.; Salzberg, S.L.; World, B.J.; Pachter, L.P. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **2010**, *28*, 511–515. [CrossRef]
45. Lü, F.G.; Fu, K.Y.; Li, Q.; Guo, W.C.; Ahmat, T.; Li, G.Q. Identification of carboxylesterase genes and their expression profiles in the Colorado potato beetle *Leptinotarsa decemlineata* treated with fipronil and cyhalothrin. *Pestic. Biochem. Physiol.* **2015**, *122*, 86–95. [CrossRef]
46. Xie, M.; Ren, N.; You, Y.; Chen, W.; Song, Q.; You, M. Molecular characterization of two α-esterase genes involving chlorpyrifos detoxification in the diamondback moth, *Plutella xylostella*. *Pest Manag. Sci.* **2017**, *73*, 1204–1212. [CrossRef]
47. Feng, X.; Liu, N. Functional characterization of carboxylesterases in insecticide resistant house flies, *Musca domestica*. *J. Vis. Exp.* **2018**, *138*, 58106. [CrossRef]
48. Bai, L.; Zhao, C.; Xu, J.; Feng, C.; Li, Y.; Dong, Y.; Ma, Z. Identification and biochemical characterization of carboxylesterase 001G associated with insecticide detoxification in *Helicoverpa armigera*. *Pestic. Biochem. Physiol.* **2019**, *157*, 69–79. [CrossRef]
49. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **2001**, *25*, 402–408. [CrossRef]
50. Paton, M.G.; Karunaratne, S.H.P.; Giakoumaki, E.; Roberts, N.; Hemingway, J. Quantitative analysis of gene amplification in insecticide-resistant *Culex mosquitoes*. *Biochem. J.* **2000**, *346*, 17–24. [CrossRef] [PubMed]
51. Poulton, B.C.; Fraser, C.; Amalia, A.; Sattelle, D.B.; Lycett, G.J. *Aedes aegypti* CCEae3A carboxylase expression confers carbamate, organophosphate and limited pyrethroid resistance in a model transgenic mosquito. *PLoS Negl. Trop. Dis.* **2024**, *18*, e0011595. [CrossRef] [PubMed]

52. Li, Y.; Bai, L.; Zhao, C.; Xu, J.; Sun, Z.; Dong, Y.; Li, D.; Liu, X.; Ma, Z. Functional characterization of two carboxylesterase genes involved in pyrethroid detoxification in *Helicoverpa armigera*. *J. Agric. Food Chem.* **2020**, *68*, 3390–3402. [CrossRef]
53. Li, H.; Buczkowski, G.; Mittapalli, O.; Xie, J.; Wu, J.; Westerman, R.; Schemerhorn, B.J.; Murdock, L.L.; Pittendrigh, B.R. Transcriptomic profiles of *Drosophila melanogaster* third instar larval midgut and responses to oxidative stress. *Insect Mol. Biol.* **2008**, *17*, 325–339. [CrossRef] [PubMed]
54. Arrese, E.; Soulages, J. Insect fat body: Energy, metabolism, and regulation. *Annu. Rev. Entomol.* **2010**, *55*, 207–225. [CrossRef]
55. Mao, K.; Ren, Z.; Li, W.; Cai, T.; Qin, X.; Wan, H.; Jun, B.; He, S.; Li, J. Carboxylesterase genes in nitenpyram-resistant brown planthoppers, *Nilaparvata lugens*. *Insect Sci.* **2021**, *28*, 1049–1060. [CrossRef]
56. Ren, N.; Xie, M.; You, Y.; Li, J.; Chen, W.; Cheng, X.; You, M. Fipronil-resistance mediated by carboxylesterases in the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae). *Acta Entomol. Sin.* **2015**, *58*, 288–296. [CrossRef]
57. Guo, Y.; Si, F.; Han, B.; Qiao, L.; Chen, B. Identification and functional validation of P450 genes associated with pyrethroid resistance in the malaria vector *Anopheles sinensis* (Diptera: Culicidae). *Acta Trop.* **2024**, *260*, 107413. [CrossRef]
58. Piermarini, P.M.; Esquivel, C.J.; Denton, J.S. Malpighian tubules as novel targets for mosquito control. *Int. J. Environ. Res. Public Health* **2017**, *14*, 111. [CrossRef] [PubMed]
59. Diop, M.M.; Chandre, F.; Rossignol, M.; Porciani, A.; Chateau, M.; Moiroux, N.; Pennetier, C. Sub-lethal insecticide exposure affects host biting efficiency of Kdr-resistant *Anopheles gambiae*. *Peer Commun. J.* **2021**, *1*, E28. [CrossRef]
60. Nouage, L.; Elanga-Ndille, E.; Binyang, A.; Tchouakui, M.; Atsatse, T.; Ndo, C.; Kekeunou, S.; Wondji, C.S. Influence of GST- and P450-based metabolic resistance to pyrethroids on blood feeding in the major African malaria vector *Anopheles funestus*. *PLoS ONE* **2020**, *15*, e0230984. [CrossRef]
61. Spillings, B.L.; Coetzee, M.; Koekemoer, L.L.; Brooke, B.D. The effect of a single blood meal on the phenotypic expression of insecticide resistance in the major malaria vector *Anopheles funestus*. *Malar. J.* **2008**, *7*, 226. [CrossRef] [PubMed]
62. Oliver, S.V.; Brooke, B.D. The effect of multiple blood-feeding on the longevity and insecticide resistant phenotype in the major malaria vector *Anopheles arabiensis* (Diptera: Culicidae). *Parasites Vectors* **2014**, *7*, 390. [CrossRef] [PubMed]
63. Li, M.; William, R.R.; Zhang, L.; Scott, J.G.; Gao, X.; Kristensen, M.; Liu, N. A whole transcriptomal linkage analysis of gene co-regulation in insecticide resistant house flies, *Musca domestica*. *BMC Genom.* **2013**, *14*, 803. [CrossRef]
64. Zhou, D.; Liu, X.; Sun, Y.; Ma, L.; Shen, B.; Zhu, C. Genomic analysis of detoxification supergene families in the mosquito *Anopheles sinensis*. *PLoS ONE* **2015**, *10*, e0143387. [CrossRef]
65. Feng, X.; Liu, N. Functional analyses of house fly carboxylesterases involved in insecticide resistance. *Front. Physiol.* **2020**, *11*, 595009. [CrossRef]
66. Li, Y.; Sun, H.; Tian, Z.; Li, Y.; Ye, X.; Li, R.; Li, X.; Zheng, S.; Liu, J.; Zhang, Y. Identification of key residues of carboxylesterase PxEst-6 involved in pyrethroid metabolism in *Plutella xylostella* (L.). *J. Hazard. Mater.* **2021**, *407*, 124612. [CrossRef]
67. Xia, J.; Xu, H.; Yang, Z.; Pan, H.; Yang, X.; Guo, Z.; Yang, F.; Guo, L.; Sun, X.; Wang, S.; et al. Genome-wide analysis of carboxylesterases (COEs) in the whitefly, *Bemisia tabaci* (Gennadius). *Int. J. Mol. Sci.* **2019**, *20*, 4973. [CrossRef]
68. Devonshire, A.L.; Heidari, R.; Huang, H.Z.; Hammock, B.D.; Russell, R.J.; Oakeshot, J.G. Hydrolysis of individual isomers of fluorogenic pyrethroid analogs by mutant carboxylesterases from *Lucilia cuprina*. *Insect Biochem. Molec.* **2007**, *37*, 891–902. [CrossRef]
69. Li, Y.; Farnsworth, C.A.; Coppin, C.W.; Teese, M.G.; Liu, J.W.; Scott, C.; Zhang, X.; Russell, R.J.; Oakeshott, J.G. Organophosphate and pyrethroid hydrolase activities of mutant esterases from the cotton bollworm *Helicoverpa armigera*. *PLoS ONE* **2013**, *8*, e77685. [CrossRef]
70. Marcombe, S.; Fustec, B.; Cattel, J.; Chonephetsarath, S.; Thammavong, P.; Phommavanh, N.; David, J.P.; Corbel, V.; Sutherland, I.W.; Hertz, J.C.; et al. Distribution of insecticide resistance and mechanisms involved in the arbovirus vector *Aedes aegypti* in Laos and implication for vector control. *PLoS Negl. Trop. Dis.* **2019**, *13*, e0007852. [CrossRef]
71. Poupardin, R.; Srisukontarat, W.; Yunta, C.; Ranson, H. Identification of carboxylesterase genes implicated in temephos resistance in the dengue vector *Aedes aegypti*. *PLoS Negl. Trop. Dis.* **2014**, *8*, e2743. [CrossRef] [PubMed]
72. Grigoraki, L.; Lagnel, J.; Kioulos, I.; Kampouraki, A.; Morou, E.; Labbe, P.; Weill, M.; Vontas, J. Transcriptome profiling and genetic study reveal amplified carboxylesterase genes implicated in temephos resistance, in the Asian tiger mosquito *Aedes albopictus*. *PLoS Negl. Trop. Dis.* **2015**, *9*, e0003771. [CrossRef] [PubMed]
73. Scott, J.G.; Michel, K.; Bartholomay, L.C.; Siegfried, B.D.; Hunter, W.B.; Smagghe, G.; Zhu, K.Y.; Douglas, A.E. Towards the elements of successful insect RNAi. *J. Insect Physiol.* **2013**, *59*, 1212–1221. [CrossRef] [PubMed]
74. Lycett, G.J.; McLaughlin, L.A.; Ranson, H.; Hemingway, J.; Kafatos, F.C.; Loukeris, T.G.; Paine, M.J.I. *Anopheles gambiae* P450 reductase is highly expressed in oenocytes and in vivo knockdown increases permethrin susceptibility. *Insect Mol. Biol.* **2006**, *15*, 321–327. [CrossRef]
75. Lumjuan, N.; Rajatileka, S.; Changsom, D.; Wicheer, J.; Leelapat, P.; Prapanthadara, L.A.; Somboon, P.; Lycett, G.; Ranson, H. The role of the *Aedes aegypti* Epsilon glutathione transferases in conferring resistance to DDT and pyrethroid insecticides. *Insect Biochem. Mol. Biol.* **2011**, *41*, 203–209. [CrossRef]

76. Li, T.; Liu, L.; Zhang, L.; Liu, N. Role of G-protein-coupled receptor-related genes in insecticide resistance of the mosquito, *Culex quinquefasciatus*. *Sci. Rep.* **2014**, *4*, 6474. [CrossRef]
77. Li, R.; Sun, X.; Liang, P.; Gao, X. Characterization of carboxylesterase *Px*α*E8* and its role in multi-insecticide resistance in *Plutella xylostella* (L.). *J. Integr. Agric.* **2022**, *21*, 1713–1721. [CrossRef]
78. Peng, C.; Yin, H.; Liu, Y.; Mao, X.F.; Liu, Z.Y. RNAi mediated gene silencing of detoxification related genes in the *Ectropis oblique*. *Genes* **2022**, *13*, 1141. [CrossRef]
79. Li, J.; Jia, Y.; Zhang, D.; Li, Z.; Zhang, S.; Liu, X. Molecular identification of carboxylesterase genes and their potential roles in the insecticides susceptibility of *Grapholita molesta*. *Insect Mol. Biol.* **2023**, *32*, 305–315. [CrossRef]
80. Vaughan, A.M.; Hawkes, N.J.; Hemingway, J. Co-amplification explains linkage disequilibrium of two mosquito esterase genes in insecticide resistant *Culex quinquefasciatus*. *Biochem. J.* **1997**, *325*, 359–365. [CrossRef] [PubMed]
81. Wheelock, C.E.; Shan, G.; Ottea, J. Overview of carboxylesterases and their role in the metabolism of insecticides. *J. Pest Sci.* **2005**, *30*, 75–83. [CrossRef]
82. Yan, J.; Nauen, R.; Reitz, S.; Alyokhin, A.; Zhang, J.; Mota-Sanchez, D.; Kim, Y.; Palli, S.R.; Rondon, S.I.; Nault, B.A.; et al. The new kid on the block in insect pest management: Sprayable RNAi goes commercial. *Sci. China Life Sci.* **2024**, *67*, 1766–1768. [CrossRef]
83. Narva, K.; Toprak, U.; Alyokhin, A.; Groves, R.; Jurat-Fuentes, J.L.; Moar, W.; Nauen, R.; Whipple, S.; Head, G. Insecticide resistance management scenarios differ for RNA-based sprays and traits. *Insect Mol. Biol.* **2025**, *34*, 518–526. [CrossRef]
84. Airs, P.M.; Bartholomay, L.C. RNA Interference for Mosquito and Mosquito-Borne Disease Control. *Insects* **2017**, *8*, 4. [CrossRef] [PubMed]
85. Wiltshire, R.M.; Duman-Scheel, M. Advances in oral RNAi for disease vector mosquito research and control. *Curr. Opin. Insect Sci.* **2020**, *40*, 18–23. [CrossRef] [PubMed]
86. Girard, M.; Berthaud, V.; Martin, E.; Vallon, L.; Rebollo, R.; Vallier, A.; Vigneron, A.; Hay, A.; Moro, C.V.; Minard, G. Evaluation of non-invasive dsRNA delivery methods for the development of RNA interference in the Asian tiger mosquito *Aedes albopictus*. *J. Pest Sci.* **2025**, *98*, 581–596. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Expression of Heat Shock Protein 90 Genes Induced by High Temperature Mediated Sensitivity of *Aphis glycines* Matsumura (Hemiptera: Aphididae) to Insecticides

Xue Han, Yulong Jia, Changchun Dai, Xiaoyun Wang, Jian Liu and Zhenqi Tian *

Key Laboratory of Crop Pests in Northern Cold Regions of Heilongjiang Province, College of Plant Protection, Northeast Agricultural University, No. 600 Changjiang Road, Xiangfang District, Harbin 150030, China; hanxue3980@126.com (X.H.); jiayulong0706@163.com (Y.J.); changchundai@163.com (C.D.); wxy3236@126.com (X.W.); jliu@neau.edu.cn (J.L.)
* Correspondence: tianzq@neau.edu.cn

Simple Summary: Heat shock protein (HSP) genes are known to be activated in response to abiotic stresses. The *Hsp90* genes in *Aphis glycines* Matsumura (Hemiptera: Aphididae) (*AgHsp75*, *AgHsp83*, and *AgGrp94*) are significantly upregulated under both high-temperature conditions and insecticide exposure. The mortality of *A. glycines* was significantly increased after silencing *AgHsp90s*, indicating their critical role in stress tolerance. These findings suggest that targeting HSPs pathways may provide novel strategies for sustainable aphid management in the context of climate warming.

Abstract

Soybean aphid, *Aphis glycines* Matsumura (Hemiptera: Aphididae), is a major pest of soybean fields. While high-temperature stress induced by global warming can initially suppress aphid populations, these pests may eventually adapt, leading to more severe infestations and crop damage. Heat shock proteins (HSPs), which are upregulated in response to heat stress to protect aphid development, also confer tolerance to other abiotic stressors, including insecticides. To investigate the role of HSPs in insecticide resistance in *A. glycines*, we analyzed the expression profiles of three *AgHsp90* genes (*AgHsp75*, *AgHsp83*, and *AgGrp94*) following exposure to high temperatures and insecticides. Functional validation was performed using RNA interference (RNAi) to silence *AgHsp90* genes. Our results demonstrated that *AgHsp90* genes were significantly upregulated under both heat and insecticide stress conditions. Furthermore, after feeding on dsRNA of *AgHsp90* genes, mortality rates of *A. glycines* significantly increased when exposed to imidacloprid and lambda-cyhalothrin. This study provides evidence that *AgHsp90* genes play a crucial role in mediating thermal tolerance and insecticide resistance in *A. glycines*.

Keywords: *Aphis glycines*; heat shock protein; thermal stress; insecticide resistance

1. Introduction

With global climate change, the frequency of extreme weather events is increasing, and their duration is becoming prolonged [1]. Recent observations indicate a warming trend in the local climate of Harbin, northeastern China, with diurnal high temperatures frequently exceeding 35 °C for approximately seven consecutive days during summer and autumn in Heilongjiang Province [2]. This phenomenon significantly impacts poikilothermic animals, particularly insects [3]. High temperature exhibits negative effects on native

insect herbivores, including reduced survival rates, compromised reproductive capacity, and diminished food intake, ultimately impairing growth and development [4,5]. Extreme high temperatures (EHTs) can induce protein denaturation and heat damage, manifesting as altered phospholipid membrane fluidity and disrupted cellular homeostasis [6]. Furthermore, EHTs promote the accumulation of reactive oxygen species (ROS) in insects, leading to oxidative stress (OS) and resulting in damage to DNA, lipids, or proteins [7].

Insects have evolved complex heat tolerance mechanisms to cope with thermal stresses, including behavioral, physiological, biochemical, and molecular adaptations [8,9]. Among these, heat shock proteins (HSPs) serve as essential molecular chaperones that participate in diverse biological processes. They facilitate the refolding of misfolded proteins, thereby maintaining cellular proteostasis [10]. HSPs have been implicated as critical determinants of both insect thermotolerance and insecticide resistance, orchestrating stress response pathways [11,12]. Based on molecular weight and sequence homology, HSPs are classified into six superfamilies: small HSPs (sHSPs), HSP40, HSP60, HSP70, HSP90, and HSP100 [13,14]. While HSP70 has been extensively studied in insect thermoregulation, the description of HSP90 in thermal adaptation remains comparatively underexplored [15]. In this study, we focused on three members of the *AgHsp90s*: *AgHsp75*, *AgHsp83*, and *AgGrp94*. *AgHsp83* represents the canonical cytosolic *Hsp90* homolog. Although designated *AgHsp75* (75 kDa), it shares high homology in core domains (notably the conserved ATPase domain) with *AgHsp83* and contains signature *Hsp90* motifs, confirming its classification within the *Hsp90* family despite nominal molecular weight variations across species. *AgGrp94* (Glucose-Regulated Protein 94) denotes the endoplasmic reticulum (ER)-resident *Hsp90* isoform (also termed *Hsp90b1*), primarily involved in folding, assembly, and quality control of ER luminal proteins, and its expression is often induced by ER stress (e.g., unfolded protein response) [16]. Therefore, verifying the functions of these three genes can provide a basis for in-depth research on aphid *Hsp90s*.

The soybean aphid, *Aphis glycines*, is a major pest of cultivated soybean. It causes leaf shrinkage, plant dwarfing, poor root development, reduced pod number, and decreased hundred-grain weight by feeding on the phloem sap of soybean plants. Severe infestations can lead to plant wilting and mortality [17,18]. *Aphis glycines* can also transmit plant viruses, such as soybean mosaic virus [19], alfalfa mosaic virus [20], and potato virus Y [21]. Additionally, aphids excrete honeydew on plant leaves, leading to sooty mold, which obstructs leaf photosynthesis [18].

The escalating frequency and dosage of chemical insecticide applications have led to the development of robust insecticide resistance in insect populations. A growing body of evidence indicates that high-temperature stress significantly enhances insecticide resistance in insects [22–24]. Notably, recent studies suggest that HSP70 in aphids may serve as critical mediators linking thermal stress responses to insecticide resistance mechanisms [12,25,26]. For instance, the high-temperature induced upregulation of *Hsp70* genes in *Aphis gossypii* and *A. glycines* has been correlated with increased resistance to multiple insecticides, including acetamiprid, sulfoxaflor, and imidacloprid. Specifically, in *A. glycines*, imidacloprid treatment significantly elevated the relative expression of *Hsp75* (a member of *Hsp90s*), suggesting its potential involvement in insecticide resistance [27]. Nevertheless, the role of *Hsp90* genes in regulating insecticide sensitivity of *A. glycines* remains poorly characterized.

In this study, based on the results of transcriptome sequencing, three *Hsp90* genes were selected. We examined the expression patterns of *Hsp90* genes to elucidate their role in mediating the relationship between thermal stress and insecticide resistance in *A. glycines* using reverse transcription quantitative PCR (RT-qPCR). Furthermore, the role of *Hsp90* genes (*AgHsp75*, *AgHsp83*, and *AgGrp94*) in sensitivity of *A. glycines* to imidacloprid and lambda-cyhalothrin was identified using RNA interference (RNAi) technology. Our

findings provide novel insights into the physiological adaptation mechanisms of *A. glycines* under thermal stress and insecticide exposure and establish a theoretical framework for developing molecular biotechnology-based control strategies targeting this pest under elevated temperature conditions.

2. Materials and Methods

2.1. Aphid and Host Source

Aphis glycines were collected from a soybean field at Northeast Agricultural University (NEAU), Harbin, Heilongjiang Province, China (126.72° E, 45.74° N) in 2021. The soybean cultivar Heinong 51 (Fangyuan Agricultural Co., Ltd., Wuchang, Heilongjiang Province, China) was used for aphid rearing. A single apterous adult aphid was transferred to a soybean plant at the V2 growth stage. The aphids and plants were maintained in a nylon mesh insect-rearing cage (40 × 40 × 40 cm) at 25 ± 1 °C, 70 ± 5% RH, and a photoperiod of 16: 8 (L: D) h in an RTOP-D Intelligent Climate Chamber (Zhejiang TOP Cloud-agri Technology Co., Ltd., Hangzhou, China).

2.2. Total RNA Extraction, cDNA Synthesis, and Gene Cloning

The total RNA of aphids was extracted from aphids using Trizol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. RNA quality and concentration were assessed using a NanoDrop spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). First-strand cDNA was synthesized using 1 μg total RNA and the Fast First-Strand cDNA Synthesis Mix for RT kit (Goonie Biotech Co., Ltd., Guangzhou, China). The primers were designed based on the predicted ORFs using primer premier 5.0 (Premier Biosoft International, Palo Alto, CA, USA; Table S1). The amplification products were purified from 1% agarose gels using Thermo Scientific GeneJET Kit (Thermo Fisher Scientific, Waltham, MA, USA). The purified fragments were cloned into the pEASY-T3 Simple vector (TransGen Biotech, Beijing, China) and transformed into chemically competent Trans1-T1 Escherichia coli cells (TransGen Biotech, Beijing, China). Positive clones were selected using blue-white screening and subsequently sequenced by Sangon Biotech Co., Ltd. (Shanghai, China). Plasmids containing correctly sequenced inserts were used as templates for dsRNA synthesis.

2.3. Real-Time Quantitative PCR (RT-qPCR) Analysis

The quantitative real-time (qRT-) PCR was performed using the Hieff qPCR SYBR Green Master Mix kit (Yeasen Biotechnology Co., Ltd., Shanghai, China) on a Bio-Rad CFX Maestro system (BioRad Co., Ltd., Hercules, CA, USA). The thermal cycling conditions consisted of an initial denaturation at 95 °C for 5 min, followed by 40 cycles of 95 °C for 10 s, 60 °C for 30 s, and 72 °C for 20 s, with a final extension at 72 °C for 10 min. Melting curve analysis was performed within the 60–95 °C temperature range to confirm the consistency and specificity of each reaction product. The qRT-PCR primers of *AgHsp75*, *AgHsp83*, and *AgGrp94* were designed using Beacon Designer 7 and are listed in Table S1. The *EF1α*, previously validated as stably expressed in *A. glycines*, was used as the reference gene [28]. Relative gene expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method [29,30]. Each experiment included three biological replicates with three technical replicates per sample.

2.4. Gene Expression Responds to Thermal Treatments

In this study, the newly born nymphs were initially reared at 25 °C. The 1st, 2nd, 3rd, and 4th instar nymphs, along with 3-day-old adults, were transferred to 29 °C and 33 °C for 24 h thermal treatments, respectively. Each developmental stage was triplicated at each temperature treatment. Following thermal treatment, ten aphids from each treatment

group were collected into 1.5 mL centrifuge tubes, immediately frozen in liquid nitrogen, and subsequently stored at −80 °C until further analysis.

2.5. Gene Expression Responds to Insecticide Treatment

In this study, imidacloprid (active ingredient 98% *w*/*w*, Binnong Technology Co., Ltd., Binzhou, Shandong Province, China) and lambda-cyhalothrin (active ingredient 98% *w*/*w*, Binnong Technology Co., Ltd., Binzhou, Shandong Province, China) were dissolved in dimethylformamide (DMF) to prepare 100 mg/mL stock solutions. Then the stock solutions were diluted to 5 concentration gradients using 0.1% Triton X-100. A 0.1% Triton X-100 solution and ddH_2O were used as the controls. Each concentration was tested with three biological replicates. For the bioassay, soybean leaves were dipped in each solution and air dried. Twenty-one-day-old apterous adult *A. glycines* were transferred onto the treated leaves and maintained at 25 ± 1 °C. Aphid mortality was recorded after 24 h. The median lethal concentration (LC_{50}) for imidacloprid and lambda-cyhalothrin was calculated from mortality data using SPSS 27 (IBM Corp., Armonk, NY, USA, 2020).

One-day-old apterous adults were exposed to LC_{30} of imidacloprid for 24 h using the leaf-dip method. Treated aphids were then collected and immediately stored at −80 °C for subsequent RT-qPCR analysis of *ApHsp90* gene expression.

2.6. RNA Interference

The specific fragments of *AgHsp75*, *AgHsp83*, and *AgGrp94* were amplified from cDNA templates and subsequently used for dsRNA synthesis. The gene-specific primers containing T7 RNA polymerase promoter sequences are listed in Table S1. The dsRNA was synthesized using the MEGAscrip T7 Kit (Invitrogen, Carlsbad, CA, USA) following the manufacturer's protocol. Green fluorescent protein (GFP) dsRNA was synthesized as a negative control.

The RNAi was carried out according to the artificial diet rearing method as described by Wang et al. [31] with several modifications. The dsRNA (150 ng/μL) was mixed with 0.5 M sucrose solution to prepare the RNAi diet. Each experimental group consisted of 30 one-day-old apterous adult aphids, which were transferred to 4.5 cm diameter plastic tubes (2 open ends). One tube end was sealed with two layers of parafilm, while the other end was covered with gauze for ventilation. A 100 μL aliquot of the RNAi diet was applied to the center of the parafilm layer. A parallel control group was established using an artificial diet containing ds*GFP*. After 48 h of feeding, aphids were collected for RT-qPCR analysis to evaluate knockdown efficiency.

2.7. Bioassays

To explore the potential function of *AgHsp75*, *AgHsp83*, and *AgGrp94* in insecticide sensitivity, three-day-old adults were selected for experiments. Because these three *AgHsp90s* are expressed differently in response to high temperatures at different developmental stages, but they are all significantly highly expressed at three-day-old adults. The *A. glycines* that had been fed on dsRNA were exposed to LC_{50} concentrations of imidacloprid and lambda-cyhalothrin using the leaf-dip method, as mentioned above. Mortality was assessed after 24 h of exposure. Each bioassay was independently repeated three times (biological replicates), with 20 individuals tested per replication.

2.8. Data Analysis

The Shapiro–Wilk test was employed to test the normality of the data. In the absence of normality, the data were log10(x + 1) transformed. The relative expression levels of *AgHsp90s* at different developmental stages of *A. glycines* were analyzed using one-way ANOVA followed by Tukey's Honestly Significant Difference (HSD) test for post hoc multi-

ple comparisons. Student's *t*-test was used to analyze the differences between two samples, including RNA interference efficiency and the expression level of *AgHsp90s* after exposure to insecticides. The mortality rate after RNA interference was analyzed using the Mann–Whitney U test. The probit regression was used to simulate and fit the virulence regression equation. All statistical analyses were performed using SPSS 27 (IBM Corporation, Armonk, NY, USA).

3. Results

3.1. AgHsp90 Genes Profiling at Different Developmental Stages

The relative expression levels of *AgHsp75*, *AgHsp83*, and *AgGrp94* varied significantly across different developmental stages. Both *AgHsp75* ($F = 33.31$; df = 4; $p < 0.05$; Figure 1a) and *AgHsp83* ($F = 8.24$; df = 4; $p < 0.05$; Figure 1b) exhibited the highest during the 2nd and 4th instar nymphal stages, respectively, showing significant upregulation compared to the 1st instar nymphs. Notably, *AgGrp94* demonstrated its highest expression level specifically in the 2nd instar nymphs ($F = 96.39$; df = 4; $p < 0.05$; Figure 1c).




Figure 1. Developmental stage-specific expression patterns of three *AgHsp90* genes ((**a**) *AgHsp75*, (**b**) *AgHsp83*, and (**c**) *AgGrp94*) in *Aphis glycines*. Expression levels in instar nymphs (N1) were normalized to 1-fold. Data represent means ± SE (n = 3 biological replicates). Different lowercase letters indicate statistically significant differences among developmental stages (one-way ANOVA with Tukey's HSD test, $p < 0.05$).

3.2. AgHsp90 Genes Profiling in Response to Different Thermal Stresses

The relative expression levels of *AgHsp75*, *AgHsp83*, and *AgGrp94* were significantly upregulated following heat stress ($p < 0.05$; Figure 2). Heat stress induced a generally dose-dependent upregulation pattern, with expression levels increasing proportionally to temperature elevation during each developmental stage. Notably, the most pronounced induction of *Hsp90* gene expression occurred in the 1st instar nymphs and adult aphids. Specifically, the expression of *AgHsp75* at 33 °C exhibited a remarkable upregulation, reaching 53.15-fold and 24.29-fold relative to the control group ($p < 0.05$; Figure 2a).




Figure 2. Temperature-dependent expression of *AgHsp90* genes in *Aphis glycines* after 24 h exposure to different high temperatures. (**a**) *AgHsp75*, (**b**) *AgHsp83*, and (**c**) *AgGrp94*. The 25 °C group served as the control. Different lowercase letters indicate significant differences among multiple samples (one-way ANOVA with Tukey's HSD test, $p < 0.05$).

3.3. Expression Levels of AgHsp90 Genes Following Insecticide Exposure

The LC_{30} values for imidacloprid and lambda-cyhalothrin are presented in Table 1. Exposure to LC_{30} concentrations of imidacloprid and lambda-cyhalothrin significantly

upregulated *ApHsp90* gene expression ($p < 0.05$; Figure 3). Specifically, following 24 h of imidacloprid exposure, the expression levels of *AgHsp75*, *AgHsp83*, and *AgGrp94* increased by 7.87-fold, 3.52-fold, and 3.65-fold, respectively. Similarly, lambda-cyhalothrin exposure induced significant upregulation of *AgHsp90s*, with mRNA levels increasing to 1.41-fold, 1.33-fold, and 3.04-fold for *AgHsp75*, *AgHsp83*, and *AgGrp94*, respectively, compared to the control group.

Table 1. Resistance of *Aphis glycines* to different insecticides.

Insecticides	Slope (±SE)	LC_{30} (95% CL)(mg/L)	LC_{50} (95% CL)(mg/L)	χ^2	df	*p*
Imidacloprid	1.59 ± 0.33	2.30 (0.90–4.02)	4.91 (2.60–8.54)	2.93	5	0.72
Lambda-cyhalothrin	1.51 ± 0.42	0.79 (0.11–1.62)	1.75 (0.55–3.18)	0.45	5	0.99




Figure 3. Insecticide-induced expression of *AgHsp90* genes in adult *Aphis glycines* after 24 h exposure. (**a**) Imidacloprid, (**b**) Lambda-cyhalothrin. Untreated insects served as the control. The asterisk above bar indicates significant differences (Student's *t*-test, * $p < 0.05$, *** $p < 0.001$).

3.4. Effect of RNAi on AgHsp90 Gene Expression and Bioassays

In order to evaluate the interference efficiency, the relative expression levels of *AgHsp90* genes were examined at 24 h, 48 h, and 72 h post-feeding. The highest interference efficiency was observed at 48 h post-feeding, followed by a significant decline at 72 h ($p < 0.05$; Figure 4).




Figure 4. Expression of *AgHsp90* genes in adult *Aphis glycines* following dsRNA feeding. (**a**) *AgHsp75*, (**b**) *AgHsp83*, and (**c**) *AgGrp94*. The asterisk above bar indicates significant differences (Student's *t*-test, * $p < 0.05$, *** $p < 0.001$).

Based on the above results, we assessed insecticide sensitivity at the peak silencing time point (48 h). Mortality rates of *A. glycines* exposed to imidacloprid and lambda-cyhalothrin for 24 h after *AgHsp90s* silencing were significantly increased compared to control groups ($p < 0.05$; Figure 5). Following feeding ds*AgHsp75*, ds*AgHsp83*, and ds*AgGrp94*,

the mortality of *A. glycines* treated with imidacloprid increased by 37%, 22%, and 32%, respectively (Figure 5a). Similarly, mortality of *A. glycines* exposed to lambda-cyhalothrin increased by 22%, 18%, and 18%, respectively (Figure 5b).




Figure 5. Mortality rates of *Aphis glycines* after 24 h exposure to insecticides following RNAi knockdown. (**a**) Imidacloprid, (**b**) Lambda-cyhalothrin. The asterisk above bar indicates significant differences (Mann–Whitney U test, * $p < 0.05$).

4. Discussion

Global climate change is characterized by a significant increase in both the frequency and intensity of extreme high-temperature (EHT) events, which affect insects across multiple biological levels. EHTs can cause direct thermal damage while simultaneously inducing a cascade of molecular, biochemical, and physiological responses. HSPs represent a primary cellular defense mechanism, functioning to synthesize and accumulate specific chaperone molecules that prevent protein denaturation and cellular dysfunction under EHT conditions [32]. Previous studies have demonstrated that the transcriptional profiles of insect *Hsp* genes are developmentally regulated. Consistent with these findings, our results revealed distinct developmental expression patterns for the three *Hsp90* genes in *A. glycines* (Figure 1). Specifically, *AgHsp75* and *AgHsp83* exhibited peak expression levels during the N2 and N4 nymphal stages (Figure 1a,b), while *AgGrp94* showed highest expression at the N2 stage (Figure 1c). The expression patterns of *Hsp70* genes displayed similar developmental trends to those observed for *Hsp90* genes. In *A. gossypii*, expression level of *Hsp70* gene was significantly higher in the 2nd-instar nymphs compared to other developmental stages [11]. In contrast, *Hsp70* expression peaked in adult *Myzus persicae* [33]. In *Rhopalosiphum padi*, all four analyzed *Hsp* genes exhibited distinct developmental expression profiles [34]. Furthermore, *Hsp70* demonstrated stage-specific expression variation in *A. gossypii*, with highest levels in 4th-instar nymphs relative to 1st- and 2nd-instar nymphs and adults [35].

The expression of *Hsp* genes exhibits stage-specific responses to high-temperature stress. When *A. glycines* were exposed to 29 °C and 33 °C, all three *Hsp90* genes (*AgHsp75*, *AgHsp83*, and *AgGrp94*) demonstrated significant upregulation (Figure 2). Especially, the temperature-induced expression of *AgHsp90* genes was most pronounced in 1st-instar nymphs and adult aphids. This differential expression pattern may reflect distinct physiological requirements at different developmental stages. For instance, the 1st-instar nymphs are more thermally vulnerable; they may require the activation of higher levels of heat shock proteins to protect body tissues from heat damage [12,36]. The elevated expression in adults may be crucial for maintaining reproductive function under thermal stress [37,38]. These findings suggest that *Hsp90* genes play stage-specific adaptive roles in thermotolerance, which may contribute to surviving and reproducing across different developmental stages under temperature stress.

Given the diverse biological functions of HSPs, they serve as critical molecular chaperones that mediate cellular responses to various exogenous stresses, including high temperatures, insecticides, and other abiotic factors. For instance, the differential expression patterns of *AcHsp83a* and *AcHsp83b* in *Arma chinensis* suggest that these genes are pivotal for growth, development, and stress tolerance, particularly under extreme temperatures and UV-B radiation [39]. Consistent with previous findings, our study demonstrates that *Hsp75* expression in *A. glycines* is inducible by both thermal stress and insecticide exposure [27], further highlighting the multifaceted role of HSPs in stress adaptation. Due to the expression level of *Hsp90s* being different during different stages, there may be variations in insecticide resistance. Our results reveal that *Hsp90* genes are significantly upregulated in three-day-old adults *A. glycines* upon exposure to insecticides, underscoring their potential involvement in stress response pathways. Similarly, there are also many reports on its role in insecticide resistance. This observation aligns with extensive evidence documenting the role of HSP90 family members in insecticide resistance. For instance, *Sogatella furcifera* exhibits upregulation of seven *Hsp* genes under imidacloprid stress [40]. While similar responses have been reported in *Leptinotarsa decemlineata* [41], *Cydia pomonella* [42], and *Liposcelis bostrychophila* [43]. Notably, in *Anopheles arabiensis*, heat shock not only exacerbates existing pyrethroid resistance but also induces de novo resistance phenotypes [44]. Functionally, our study provides direct evidence that knockdown of *AgHsp90* genes significantly increases *A. glycines* susceptibility to insecticides, thereby confirming their contributory role in insecticide resistance. This finding is corroborated by similar observations in *Nilaparvata lugens*, where *Hsp70* knockdown significantly enhances insecticide sensitivity [26]. Collectively, these results emphasize that diverse HSP family members play indispensable roles in insect adaptation to both abiotic stresses and xenobiotic challenges, particularly in the context of insecticide resistance mechanisms.

Previous studies have suggested that organisms can exhibit cross-protection against multiple stressors, wherein exposure to one stressor induces resistance to another. For instance, *Bemisia tabaci* exposed to thiamethoxam exhibited reduced mortality and an increased lethal mean time (LT_{50}) [45]. Similarly, sublethal doses of neonicotinoid insecticides enhanced the thermal tolerance and survival rates of *Apis mellifera* [46]. In this study, we found that high temperatures upregulated *Hsp90* gene expression, which modulates the sensitivity of *A. glycines* to insecticides. These findings suggest that *Hsp90* genes play a regulatory role in cross-protection mechanisms. This phenomenon may extend to other insect species, as evidenced by studies in *Tetranychus cinnabarinus*, where *TcHsp90* was implicated in cross-protection against thermal stress and emamectin benzoate exposure [47]. Additionally, *Apolygus lucorum* exhibited *Hsp90*-mediated responses to combined temperature and insecticide stresses [48]. Collectively, these studies highlight the critical role of *Hsp90* genes in conferring cross-resistance between high temperatures and insecticides in insects.

The emergence of cross-resistance significantly complicates pest management strategies. In summer conditions characterized by high temperature and humidity, field populations of *A. gossypii* predominantly exhibit the summer morphotype, which is characterized by smaller body size. Research has demonstrated that the summer morphotype displays enhanced resistance to insecticides [49]. This resistance is further exacerbated by the extensive application of chemical insecticides, rendering the *A. gossypii* control increasingly challenging [50]. Liu et al. [12] proposed that the upregulated expression of *Hsp70* genes in *A. gossypii* plays a critical role in this phenomenon. *Hsp70* genes mainly resist environmental stress by recognizing and transiently binding exposed hydrophobic residues [51]. In contrast, *Hsp90* likely operates further upstream within the stress response network. Rather than directly interacting with insecticides, *Hsp90* functions as a critical regulator of

signal transduction cascades and a stabilizer of client proteins essential for survival under stress [52]. This functional distinction suggests potential synergies, where *Hsp90*-mediated stabilization enables *Hsp70* function. Notably, *A. glycines* exhibits parallel physiological responses to high temperatures, including the emergence of smaller summer morphotypes [53]. Based on these observations, we hypothesize that *Hsp90* genes, in addition to *Hsp70* genes, are also involved in regulating cross-resistance between high temperatures and insecticides in *A. glycines*. Given these findings, the control of *A. glycines* is expected to become progressively more difficult under ongoing climate warming. Consequently, elucidating whether *AgHsp90* genes serve as key regulators in the development of cross-resistance has important implications for the development of effective pest management strategies.

The interaction between insecticide toxicity and elevated temperature exhibits dynamic and complex responses. In the short term, high temperature often shows an antagonistic effect: on one hand, warming accelerates insecticide degradation (such as the environmental half-life of avermectin is significantly shortened) [54]; on the other hand, insects enhance their cellular protection capabilities through stress pathways, such as the induction of HSPs, thereby improving their tolerance to insecticides. Wang et al. [55] demonstrated that the LC_{50} of avermectin at 35 °C was significantly higher than at 25 °C in *Liriomyza trifolii*, supporting this antagonistic mechanism. However, under prolonged exposure, a synergistic effect gradually prevails. The combined stress of sub-lethal concentrations of insecticides and high temperature induces cross-adaptation, characterized by the coordinated upregulation of detoxification-related genes (P450, GST) and heat stress-related genes (*Hsp70*, *Hsp90*) [56]. Notably, avermectin-resistant strains have evolved constitutive heat resistance, exhibiting enhanced thermotolerance and an elevated critical maximum temperature for HSPs activation, along with increased baseline expression levels of heat shock proteins [57]. In our study, both high temperature and insecticide exposure induced the upregulation of *AgHsp90s*, suggesting that *A. glycines* may activate a shared stress-response pathway under these conditions. This finding implies the existence of an interactive resistance mechanism between high temperature and insecticides in *A. glycines*.

5. Conclusions

In conclusion, our results confirm that the three *Hsp90* genes in *A. glycines* not only play a role in the insect's high-temperature response but also play an important role in its sensitivity to insecticides. Our discovery provides a perspective on the defense mechanisms of aphids against high temperatures and other environmental pressures. These data also indicate that *AgHsp90* genes may be a potential target for pest management based on RNAi.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/insects16080772/s1. Table S1: The primers used for amplification, qRT-PCR, and dsRNA synthesis. Table S2: Developmental stage-specific expression patterns of three *AgHsp90* genes. Table S3: Temperature-dependent expression of *AgHsp90* genes in *Aphis glycines* after 24 h exposure to different high temperatures. Table S4: Insecticide-induced expression of *AgHsp90* genes in adult *Aphis glycines* after 24 h exposure. Table S5: Expression of *AgHsp90* genes in adult *Aphis glycines* following dsRNA feeding. Figure S1: The melt curve of *AgHsp90s* products. *AgHsp75* (a), *AgHsp83* (b), *AgGrp94* (c).

Author Contributions: Conceptualization, Z.T., C.D., X.W. and J.L.; methodology, Z.T.; investigation, Z.T., C.D., X.W., J.L., X.H. and Y.J.; software, X.H. and Y.J.; resources, Z.T. and X.W.; data curation, Z.T., X.H. and Y.J.; writing—original draft, Z.T., X.H., C.D. and J.L.; writing—review and editing, Z.T., X.H. and Y.J.; formal analysis, X.H. and Y.J.; visualization, X.H.; supervision, Z.T., X.W.; project administration, Z.T.; funding acquisition, Z.T. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the National Natural Science Foundation of China (32202301) and the "Young Talents" Project of Northeast Agricultural University (2023-KYYWF-1166).

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding author.

Acknowledgments: The authors would like to acknowledge the Key Laboratory of Crop Pests in Northern Cold Regions of Heilongjiang Province, College of Plant Protection, Northeast Agricultural University, for all the support given to this research.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. IPCC. *Global Warming of 1.5 °C*; Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P.R., Pirani, A., Eds.; Cambridge University Press: Cambridge, UK, 2018; 616p. [CrossRef]
2. Heilongjiang Meteorological Bureau. *Ambient Temperature Data of Harbin (2010–2020)*; Heilongjiang Meteorological Bureau: Harbin, China, 2020.
3. Gao, G.; Feng, L.; Perkins, L.E.; Sharma, S.; Lu, Z. Effect of the frequency and magnitude of extreme temperature on the life history traits of the large cotton aphid, *Acyrthosiphon gossypii* (Hemiptera: Aphididae): Implications for their population dynamics under global warming. *Entomol. Gen.* **2018**, *37*, 103–113. [CrossRef]
4. Abdel-Hady, A.A.A.; Ramadan, M.M.; Lü, J.; Hashem, A.S. High-temperature shock consequences on the red flour beetle (*Tribolium castaneum*) and the rice weevil (*Sitophilus oryzae*). *J. Therm. Biol.* **2021**, *100*, 103062. [CrossRef] [PubMed]
5. Liu, J.; Wang, C.; Desneux, N.; Lu, Y. Impact of temperature on survival rate, fecundity, and feeding behavior of two aphids, *Aphis gossypii* and *Acyrthosiphon gossypii*, when reared on cotton. *Insects* **2021**, *12*, 565. [CrossRef] [PubMed]
6. Bowler, K. Heat death in poikilotherms: Is there a common cause? *J. Therm. Biol.* **2018**, *76*, 77–79. [CrossRef] [PubMed]
7. Dampc, J.; Kula-Maximenko, M.; Molon, M.; Durak, R. Enzymatic defense response of apple aphid *Aphis pomi* to increased temperature. *Insects* **2020**, *11*, 436. [CrossRef] [PubMed]
8. Durak, R.; Dampc, J.; Kula-Maximenko, M.; Mołoń, M.; Durak, T. Changes in antioxidative, oxidoreductive and detoxification enzymes during development of aphids and temperature increase. *Antioxidants* **2021**, *10*, 1181. [CrossRef] [PubMed]
9. Saeidi, F.; Mikani, A.; Moharramipour, S. Thermal tolerance variations and physiological adjustments in a winter active and a summer active aphid species. *J. Therm. Biol.* **2021**, *98*, 102950. [CrossRef] [PubMed]
10. Shan, Q.; Ma, F.; Wei, J.; Li, H.; Ma, H.; Sun, P. Physiological functions of heat shock proteins. *Curr. Protein Pept. Sci.* **2020**, *21*, 751–760. [CrossRef] [PubMed]
11. Liang, P.; Guo, M.; Wang, D.; Li, T.; Li, R.; Li, D.; Cheng, S.; Zhen, C.; Zhang, L. Molecular and functional characterization of heat-shock protein 70 in *Aphis gossypii* under thermal and xenobiotic stresses. *Pestic. Biochem. Physiol.* **2024**, *199*, 105774. [CrossRef] [PubMed]
12. Liu, J.P.; Liu, P.; Wang, W.; Liang, G.M.; Lu, Y.H. Characterizing three heat shock protein 70 genes of *Aphis gossypii* and their expression in response to temperature and insecticide stress. *J. Agric. Food Chem.* **2025**, *73*, 2842–2852. [CrossRef] [PubMed]
13. Kim, K.; Kim, R.; Kim, S.H. Crystal structure of small heat-shock protein. *Nature* **1998**, *394*, 595–599. [CrossRef] [PubMed]
14. Feder, M.E.; Hofmann, G.E. Heat-shock proteins, molecular chaperones, and the stress response: Evolutionary and ecological physiology. *Annu. Rev. Physiol.* **1999**, *61*, 243–282. [CrossRef] [PubMed]
15. González-Tokman, D.; Córdoba-Aguilar, A.; Dáttilo, W.; Lira-Noriega, A.; Sánchez-Guillén, R.A.; Villalobos, F. Insect responses to heat: Physiological mechanisms, evolution and ecological implications in a warming world. *Biol. Rev.* **2020**, *95*, 802–821. [CrossRef] [PubMed]
16. Kampinga, H.H.; Hageman, J.; Vos, M.J.; Kubota, H.; Tanguay, R.M.; Bruford, E.A.; Cheetham, M.E.; Chen, B.; Hightower, L.E. Guidelines for the nomenclature of the human heat shock proteins. *Cell Stress Chaperones* **2009**, *14*, 105–111. [CrossRef] [PubMed]
17. Ragsdale, D.W.; Landis, D.A.; Brodeur, J.; Heimpel, G.E.; Desneux, N. Ecology and management of the soybean aphid in North America. *Annu. Rev. Entomol.* **2011**, *56*, 375–399. [CrossRef] [PubMed]
18. Wu, Z.; Schenk-Hamlin, D.; Zhan, W.; Ragsdale, D.; Heimpel, G. The soybean aphid in China: A historical review. *Ann. Entomol. Soc. Am.* **2004**, *97*, 209–218. [CrossRef]
19. Wang, R.-Y.; Ghabrial, S.A. Effect of aphid behavior on efficiency of transmission of soybean mosaic virus by the soybean-colonizing aphid, *Aphis glycines*. *Plant Dis.* **2002**, *86*, 1260–1264. [CrossRef] [PubMed]
20. Hill, J.H.; Alleman, R.; Hogg, D.B.; Grau, C.R. First report of transmission of soybean mosaic virus and alfalfa mosaic virus by *Aphis glycines* in the new world. *Plant Dis.* **2001**, *85*, 561. [CrossRef] [PubMed]
21. Davis, J.A.; Radcliffe, E.B.; Ragsdale, D.W. Soybean aphid, *Aphis glycines* Matsumura, a new vector of potato virus Y in potato. *Am. J. Potato Res.* **2005**, *82*, 197–201. [CrossRef]

22. Chen, B.; Feder, M.E.; Kang, L. Evolution of heat-shock protein expression underlying adaptive responses to environmental stress. *Mol. Ecol.* **2018**, *27*, 3040–3054. [CrossRef] [PubMed]
23. Dong, B.; Liu, X.-Y.; Li, B.; Li, M.-Y.; Li, S.-G.; Liu, S. A heat shock protein protects against oxidative stress induced by lambda-cyhalothrin in the green peach aphid *Myzus persicae*. *Pestic. Biochem. Physiol.* **2022**, *181*, 104995. [CrossRef] [PubMed]
24. Zhou, M.; Liu, Y.; Wang, Y.; Chang, Y.; Wu, Q.; Gong, W.; Du, Y. Effect of high temperature on abamectin and thiamethoxam tolerance in *Bemisia tabaci* MEAM1 (Hemiptera: Aleyrodidae). *Insects* **2024**, *15*, 399. [CrossRef] [PubMed]
25. Han, L.L.; Zhu, M.H.; Dong, T.Y.; Zhao, K.J.; Qu, Z.C.; Yang, L.; Han, X.X. Effects of heat shock and imidacloprid on the expressions of *Hsp70* and *Hsc70* mRNA in the *Aphis glycines* (Hemiptera: Aphididae). *Acta Entomol. Sin.* **2014**, *57*, 387–394.
26. Lu, K.; Chen, X.; Liu, W.; Zhang, Z.; Wang, Y.; You, K.; Li, Y.; Zhang, R.; Zhou, Q. Characterization of heat shock protein 70 transcript from *Nilaparvata lugens* (Stål): Its response to temperature and insecticide stresses. *Pestic. Biochem. Physiol.* **2017**, *142*, 102–110. [CrossRef] [PubMed]
27. Wang, X.-Y.; Li, S.-Y.; Yao, L.; Liu, Y.; Chen, Y.-R.; Yang, H.-J.; Fan, D. Stress expression of *AgHsp75* gene in soybean aphid (*Aphis glycines*) under imidacloprid and temperature stress. *Chin. J. Biol. Control* **2020**, *36*, 913–919. [CrossRef]
28. Wang, Z.; Zhang, H.; Zhang, Z.; Zhao, J.; Ma, F.; Zheng, M.; Yang, M.; Sang, X.; Ma, K.; Li, L. Selection of reference genes for normalization of qRT–PCR analysis in the soybean aphid *Aphis glycines* Matsumura (Hemiptera: Aphididae). *J. Econ. Entomo.* **2022**, *115*, 2083–2091. [CrossRef] [PubMed]
29. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. *Methods* **2001**, *25*, 402–408. [CrossRef] [PubMed]
30. Pfaffl, M.W. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **2001**, *29*, e45. [CrossRef] [PubMed]
31. Wang, L.; Zhu, J.; Cui, L.; Wang, Q.; Huang, W.; Ji, X.; Yang, Q.; Rui, C. Overexpression of ATP-binding cassette transporters associated with sulfoxaflor resistance in *Aphis gossypii* Glover. *Pest Manag. Sci.* **2021**, *77*, 4064–4072. [CrossRef] [PubMed]
32. Ma, C.-S.; Ma, G.; Pincebourde, S. Survive a warming climate: Insect responses to extreme high temperatures. *Annu. Rev. Entomol.* **2021**, *66*, 163–184. [CrossRef] [PubMed]
33. Chen, N.; Tan, J.-Y.; Wang, Y.; Qi, M.-H.; Peng, J.-N.; Chen, D.-X.; Liu, S.; Li, M.-Y. A heat shock protein 70 protects the green peach aphid (*Myzus persicae*) against high-temperature stress. *J. Asia-Pac. Entomol.* **2022**, *25*, 101992. [CrossRef]
34. Li, Y.; Zhao, Q.; Duan, X.; Song, C.; Chen, M. Transcription of four *Rhopalosiphum padi* (L.) heat shock protein genes and their responses to heat stress and insecticide exposure. *Comp. Biochem. Physiol. A* **2017**, *205*, 48–57. [CrossRef] [PubMed]
35. Cha, S.; Shim, J.-K.; Lee, K.-Y. Identification of glucose-regulated protein 78 (*grp78*) cDNA from *Aphis gossypii* and its expression during development, heat shock and nutritional ingestion. *Physiol. Entomol.* **2016**, *41*, 193–201. [CrossRef]
36. Li, H.; Du, Y. Molecular cloning and characterization of an *Hsp90/70* organizing protein gene from *Frankliniella occidentalis* (Insecta: Thysanoptera, Thripidae). *Gene* **2013**, *520*, 148–155. [CrossRef] [PubMed]
37. Li, H.; Tan, Y.; Zhao, X.; Chen, J.; Li, S.; Ye, J.; Hao, D. Accumulation of heat shock protein 20s in the ovary and testis of *Monochamus alternatus* protects reproduction against high temperatures. *Entomol. Gen.* **2023**, *43*, 1021–1030. [CrossRef]
38. Shu, Y.; Du, Y.; Wang, J. Molecular characterization and expression patterns of *Spodoptera litura* heat shock protein 70/90, and their response to zinc stress. *Comp. Biochem. Physiol. A* **2010**, *158*, 102–110. [CrossRef] [PubMed]
39. Meng, J.-Y.; Jin, X.; He, L.-C.; Zhang, X.-X.; Yang, C.-L.; Zhang, C.-Y. Cloning, Expression profiles of *Arma chinensis* heat shock protein genes *AcHsp83a* and *AcHsp83b*, and their response to high and low temperature and UV-B stress. *Acta Entomol. Sin.* **2023**, *66*, 1425–1434. [CrossRef]
40. Zhou, C.; Yang, H.; Wang, Z.; Long, G.-Y.; Jin, D.-C. Comparative transcriptome analysis of *Sogatella furcifera* (Horváth) exposed to different insecticides. *Sci. Rep.* **2018**, *8*, 8773. [CrossRef] [PubMed]
41. Dumas, P.; Morin, M.D.; Boquel, S.; Moffat, C.E.; Morin, P.J. Expression status of heat shock proteins in response to cold, heat, or insecticide exposure in the Colorado potato beetle *Leptinotarsa decemlineata*. *Cell Stress Chaperones* **2019**, *24*, 539–547. [CrossRef] [PubMed]
42. Yang, X.-Q.; Zhang, Y.-L.; Wang, X.-Q.; Dong, H.; Gao, P.; Jia, L.-Y. Characterization of multiple heat-shock protein transcripts from *Cydia pomonella*: Their response to extreme temperature and insecticide exposure. *J. Agric. Food Chem.* **2016**, *64*, 4288–4298. [CrossRef] [PubMed]
43. Feng, H.-Y.; Chen, Z.-D.; Jiang, S.-D.; Miao, Z.-Q.; Wang, J.-J.; Wei, D.-D. Characterization and expression of heat shock protein 70s in *Liposcelis bostrychophila*: Insights into their roles in insecticidal stress response. *J. Stored Prod. Res.* **2024**, *106*, 102289. [CrossRef]
44. Oliver, S.V.; Brooke, B.D. The effect of elevated temperatures on the life history and insecticide resistance phenotype of the major malaria vector *Anopheles arabiensis* (Diptera: Culicidae). *Malar. J.* **2017**, *16*, 73. [CrossRef] [PubMed]
45. Su, Q.; Li, S.; Shi, C.; Zhang, J.; Zhang, G.; Jin, Z.; Li, C.; Wang, W.; Zhang, Y. Implication of heat-shock protein 70 and UDP-glucuronosyltransferase in thiamethoxam-induced whitefly *Bemisia tabaci* thermotolerance. *J. Pest Sci.* **2018**, *91*, 469–478. [CrossRef]

46. Gonzalez, V.H.; Hranitz, J.M.; McGonigle, M.B.; Manweiler, R.E.; Smith, D.R.; Barthell, J.F. Acute exposure to sublethal doses of neonicotinoid insecticides increases heat tolerance in honey bees. *PLoS ONE* **2022**, *17*, e0240950. [CrossRef] [PubMed]
47. Feng, H.; Wang, L.; Liu, Y.; He, L.; Li, M.; Lu, W.; Xue, C. Molecular characterization and expression of a heat shock protein gene (*Hsp90*) from the carmine spider mite, *Tetranychus cinnabarinus* (Boisduval). *J. Insect Sci.* **2010**, *10*, 112. [CrossRef] [PubMed]
48. Sun, Y.; Sheng, Y.; Bai, L.; Zhang, Y.; Xiao, Y.; Xiao, L.; Tan, Y.; Shen, Y. Characterizing heat shock protein 90 gene of *Apolygus lucorum* (Meyer-Dür) and its expression in response to different temperature and insecticide stresses. *Cell Stress Chaperones* **2014**, *19*, 725–739. [CrossRef] [PubMed]
49. Wu, K.M.; Liu, Q.X. Study on Insecticide-Induced Resurgence of Cotton Aphid. *Acta Ecol. Sin.* **1992**, *4*, 341–347.
50. Koo, H.-N.; An, J.-J.; Park, S.-E.; Kim, J.-I.; Kim, G.-H. Regional susceptibilities to 12 insecticides of melon and cotton aphid, *Aphis gossypii* (Hemiptera: Aphididae) and a point mutation associated with imidacloprid resistance. *Crop Prot.* **2014**, *55*, 91–97. [CrossRef]
51. Silva, A.X.; Bacigalupe, L.D.; Luna-Rudloff, M.; Figueroa, C.C. Insecticide Resistance Mechanisms in the Green Peach Aphid *Myzus persicae* (Hemiptera: Aphididae) II: Costs and Benefits. *PLoS ONE* **2012**, *7*, 36810. [CrossRef] [PubMed]
52. Sangster, T.A.; Lindquist, S.; Queitsch, C. Under cover: Causes, effects and implications of Hsp90-mediated genetic capacitance. *Bioessays* **2004**, *26*, 348–362. [CrossRef] [PubMed]
53. Xu, G.Q.; Chen, Y.; Wang, X.Y.; Liu, P.B.; Xu, L.; Zhao, T.H. Adaptation of the dwarf forms of soybean aphid, *Aphis glycines* to environment and impact on soybean yield. *Chin. J. Appl. Entomol.* **2011**, *48*, 1638–1645.
54. De Beeck, L.O.; Verheyen, J.; Olsen, K.; Stoks, R. Negative effects of pesticides under global warming can be counteracted by a higher degradation rate and thermal adaptation. *J. Appl. Ecol.* **2017**, *54*, 1847–1855. [CrossRef]
55. Wang, Y.C.; Chang, Y.W.; Du, Y.Z. Temperature affects the tolerance of *Liriomyza trifolii* to insecticide abamectin. *Ecotox. Environ. Safe* **2021**, *218*, 112307. [CrossRef] [PubMed]
56. Wang, Y.C.; Chang, Y.W.; Du, Y.Z. Transcriptome analysis reveals gene expression differences in Liriomyza trifolii exposed to combined heat and abamectin exposure. *PeerJ* **2021**, *9*, e12064. [CrossRef] [PubMed]
57. Wang, Y.C.; Chang, Y.W.; Gong, W.R.; Hu, J.; Du, Y.Z. The development of abamectin resistance in *Liriomyza trifolii* and its contribution to thermotolerance. *Pest Manag. Sci.* **2024**, *80*, 2053–2060. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Risk Assessment of RNAi-Based Potential Pesticide ds*NlAtg3* and Its Homologues for *Nilaparvata lugens* and Non-Target Organisms

Kai Li †, Tongtong Chen †, Yuliang Li, Kai Sun, Kun Pang, Xiaoping Yu * and Peiying Hao *

Key Laboratory of Microbiological Metrology, Measurement & Bio-Product Quality Security, State Administration for Market Regulation, China Jiliang University, Hangzhou 310018, China; 17816121215@163.com (K.L.); nadinechentt@163.com (T.C.); 18853663892@163.com (Y.L.); sunkai@cjlu.edu.cn (K.S.); pangk@cjlu.edu.cn (K.P.)

* Correspondence: yuxiaoping19630306@163.com (X.Y.); haopeiy@163.com (P.H.)

† These authors contributed equally to this work.

Simple Summary: The brown planthopper (*Nilaparvata lugens*) is a serious pest insect of rice. RNA interference technology for controlling *N. lugens* has good prospects based on its advantages, such as high species specificity, low potential for environmental pollution, and obvious effects. However, it is necessary to select suitable target genes and design species-specific dsRNA fragments for RNAi so that it can achieve effective pest control and avoid the risk to non-target organisms. In this study, we evaluated three different RNA interfering fragments targeting an autophagy-related gene *Atg3* (ds*NlAtg3*-474×1, ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10) of *N. lugens* and also assessed their effects on non-target species. The results showed that all three ds*NlAtg3* fragments were effective against *N. lugens*. In addition, the effects of ds*NlAtg3*-47×10 (specifically designed against *N. lugens*) on another four organisms, including two natural enemies, *Dolomedes sulfureus* and *Tytthus chinensis*, were tested. The results showed that the fragment ds*NlAtg3*-47×10 had no significant effect on the survival or development of non-target organisms. Therefore, the fragment ds*NlAtg3*-47×10 has good potential to be developed as insecticide for controlling *N. lugens*.

Abstract: The brown planthopper (*Nilaparvata lugens*) is an insect pest of rice, which mainly feeds on the phloem sap of the leaf sheath. RNA interference (RNAi) has application prospects in pest control, but it is necessary to select target genes and design suitable dsRNA fragments for RNAi so that it can achieve effective pest control and avoid risks to non-target organisms. NlAtg3 is a key protein in the autophagy pathway of *N. lugens*. Three kinds of dsRNA fragments of the *NlAtg3* gene (ds*NlAtg3*-474×1, ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10) were designed to compare the RNAi efficiency and specificity against the target insect *N. lugens* and non-target organisms through microinjection. The results showed that the fragment ds*NlAtg3*-474×1 showed strong inhibitory effects on the survival of *N. lugens*, which resulted in the survival rate decreasing to zero on the fifth day, while the survival rate of a closely related species, *Sogatella furcifera*, dropped to 2.22%. In contrast, ds*NlAtg3*-47×10 specifically designed against *N. lugens* only showed slight or no inhibitory effects on *S. furcifera* and other non-target organisms such as *Drosophila melanogaster*, but still showed good lethal effects against *N. lugens*, with the survival rate dropping to 18.89% on the ninth day. In addition, after being fed *N. lugens* injected with ds*NlAtg3*-47×10 fragments, the survival rate of the natural enemies *Dolomedes sulfureus* and *Tytthus chinensis* did not show significant change, compared with those treated with the ds*GFP* control. Our results suggest that the *NlAtg3* gene can serve as a potential target for controlling *N. lugens*. Moreover, by designing suitable RNAi fragments, it is possible to avoid harm to non-target organisms while effectively inhibiting the target insect *N. lugens*.

Keywords: RNA interference; *Nilaparvata lugens*; non-target organism; natural enemy; biosafety

1. Introduction

The brown planthopper *Nilaparvata lugens* is an insect pest of rice [1]. *N. lugens* damages rice by sucking the phloem sap of the leaf sheath, laying eggs in rice tissues, and transferring a variety of rice viruses [2,3]. However, with the misuse of insecticides, *N. lugens* has developed resistance to chemical insecticides [4,5]. Breeding new insect-resistant rice varieties is an environmentally friendly method, but *N. lugens* feeding on an insect-resistant rice variety for several generations can generally adapt to this variety. RNA interference (RNAi) can inhibit mRNA expression levels of the target gene in organisms, including in some species of insect. Therefore, clarifying gene function and precisely manipulating specific gene expression in insects [6,7] has opened a new pathway for effective control of insect pests.

RNAi-based insect-resistant genetically engineered (IRGE) crops have shown good application prospects in pest control. In some corn varieties, RNAi via the dsRNA of *DvvSnf7*, *Cry3Bb1*, and *Cry34Ab1*/*Cry35Ab1* genes caused significant death of western corn rootworms [8,9]. Currently, transgenic rice expressing dsRNA has made significant breakthroughs in the management of *N. lugens*. Rice expressing dsRNA of the *Nlsid-1* and *Nlaub* genes of *N. lugens* has been demonstrated to inhibit the expression of target genes in the midgut [10]. It has been reported that rice plants with overexpression of *osa-miR162a* had enhanced resistance against *N. lugens* and reduced the reproductive capacity of adults [11]. However, some rice-mediated RNAi had little effect on the survival rate of brown planthopper. It has been reported that the expression of *NlGST1-1* and *GST* activity in *N. lugens* nymphs was significantly inhibited after the insects fed on transgenic rice expressing *NlGST1-1* dsRNA, but the lethal phenotype of *N. lugens* nymphs after feeding was hardly observed [12]. It generally takes a long time to acquire insect-resistant genetically engineered rice, so before constructing IRGE rice, it is necessary to screen suitable target genes for RNAi first through convenient methods like microinjection of dsRNA or feeding-based RNAi methods.

In addition to the selection of the target gene, the effect of RNAi is also affected by several other factors, including the length, the sequence composition, and the concentration of dsRNA fragments [13,14]. After entering the cells of an insect, dsRNA fragments are first cleaved into siRNA by Dicer-2 enzyme or Dicer-2 nuclease homologues. Fragments of siRNA are usually 19–24 nt, and their bases at the 3′ end protruding from the sticky end and the phosphate group at the 5′ end play key roles in their functions. However, when Dicer-2 enzymes in insects cleave exogenous dsRNA, short fragments of varying length or sequence are produced due to the irregular cleaving position, thus affecting RNAi efficiency [15,16]. Moreover, the dsRNA length for inducing effective RNAi varies among different insects [17]. In *Drosophila melanogaster* S2-cells, dsRNAs ranging from 21 to 592 bp in length all resulted in effective silencing of the target gene, irrespective of the dsRNA length, after being forcibly introduced into cells by transfection. In contrast, uptake of dsRNAs added to the medium was clearly length-dependent [18].

For insects, autophagy is essential for growth and development. Some studies have shown that the loss of *Atg9* leads to a shortened lifespan, motor deficits, and increased susceptibility to stress in *Drosophila*. *Atg9* loss also resulted in aberrant adult midgut morphology with dramatically enlarged enterocytes [19]. Abdul indicated that LAMP2A can potentiate autophagic flux in the *Drosophila* brain, leading to enhanced stress resistance

and neuroprotection [20]. *ATG3* is one of the key genes in the autophagy pathway, and encodes an E2 ubiquitin-like binding protein in the ATG2 system, promoting the extension of phagophore [21]. Our previous study showed that RNAi through microinjection of *dsNlAtg3* into the fifth-instar nymphs of *N. lugens* resulted in a perfect lethal effect, indicating its possible role in pest control [22]. However, the minimum effective length of a dsRNA fragment specific to *N. lugens*, and its effect on non-target organisms including natural enemies, are currently unavailable. Therefore, it is necessary to screen suitable dsRNA fragments of *ATG3* specific to *N. lugens* and explore their effects on non-target organisms for the construction of a rice-mediated RNAi system for controlling *N. lugens* in the future.

2. Materials and Methods

2.1. Insects

The experimental insects of *N. lugens* and *Sogatella furcifera* have been reared on TN1 rice over ten years. The *Dolomedes sulfureus* (Araneae: Pisauridae) was collected from the rice paddy of China Jiliang University, Hangzhou, China. The *Tytthus chinensis* (Heteroptera: Miridae) was collected from a rice paddy in Ningbo, China. The *Drosophila melanogaster* was purchased from the Qidong Fungene Biotechnology company (Qidong, JiangSu, China) and fed with a medium consisting of agar, yeast and cornmeal. The insects and rice plants were maintained at a temperature of 26 ± 2 °C, 75% ± 5% RH, and a photoperiod of 16L:8D.

2.2. Double-Stranded RNA Synthesis and Microinjection

In order to compare the RNAi efficiency and specification of non-concatemerized long dsRNA (NCL-dsRNA) with concatemerized dsRNAs (C-dsRNAs), three kinds of dsDNA templates (dsDNA-474×1, dsDNA-138×3 and dsDNA-47×10) were prepared accordingly for transcripting NCL-dsRNA (ds*NlAtg3*-474×1) and C-dsRNAs (ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10) against the *NlAtg3* gene (GenBank acc. no.: MF040142.1) from *N. lugens* (Figure 1). Total RNA was extracted and used as the template to synthesize cDNA as previously described [22]. PCR was used to synthesize dsDNA-474×1 using the primers in Supplementary Table S1. dsDNA-138×3 and dsDNA-47×10 (with a *Bam*H I site and T7 promotor at both ends) for synthesizing dsRNA of ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10 were prepared by Personalbio (Shanghai, China) using a chemical synthesis method. For dsDNA-138×3, the repeat unit was a 138 bp sequence, and the whole dsDNA template contained 3 repeats of the 138 bp unit (Figure 1B). Similarly, the whole dsDNA-47×10 template contained 10 repeats of the 47 bp unit. The pMD18-T vector was used to clone the dsDNA products, which were then separately introduced into *E. coli* JM109 and sent to the Ykang company (Hangzhou, China)for sequencing. Following the manufacturer's instructions, ds*NlAtg3*-474×1, ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10 were then synthesized using the MEGAscript® T7 High Yield Transcription Kit (Figure 1A, Thermo Fisher Scientific, Waltham, MA, USA). *GFP* (GenBank acc. no.: MF169984) was used to synthesize ds*GFP* (1155 bp-1811 bp) as the control. The sequence information of the *Atg3* gene, primers, *Bam*H I site and T7 promotor are listed in Supplementary Table S1.

Microinjection needles were processed by a PC-10 microelectrode tractor (NARISHIGE, Japan, Tokyo). The 5th-instar nymphs of *N. lugens* were injected with 50 nL of dsRNA (250 ng/μL) per insect, using a microinjector (EPPENDORF, Hamburg, Germany). The synthesized dsRNA was injected into the thorax at the site between the middle and hind legs of *N. lugens*. The injected insects were cultured on TN1 rice plants [23].




Figure 1. Schematic diagram of the location and repetition of different dsRNA fragments against *NlAtg3*. (**A**) Schematic diagram of the 474 bp sequence location selected for synthesizing ds*NlAtg3*-474×1, with the T7 promoter sequence and protective bases added. (**B**) Schematic diagram of the 138 bp sequence location selected for synthesizing ds*NlAtg3*-138×3. The cDNA template used for synthesizing the ds*NlAtg3*-138×3 fragment contains 3 repeats of a 138 bp sequence, T7 promoter and protective bases. (**C**) Schematic diagram of the 47 bp sequence location selected for synthesizing ds*NlAtg3*-47×10. The cDNA template used for synthesizing the ds*NlAtg3*-47×10 fragment contains 10 repeats of a 47 bp sequence, T7 promoter and protective bases.

2.3. Effect of RNA Interference on the Survival of N. lugens

To evaluate the effect of different dsRNA fragments on the survival of *N. lugens*, ds*NlAtg3*-474×1, ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10 were injected into the 5th-instar nymphs, and ds*GFP* was used as a control. Each treatment contained 3 replicates, and each replicate pooled 30 nymphs of *N. lugens*. The survival rate and other physiological indicators of *N. lugens* were observed and counted daily.

2.4. Gene Expression Analysis Using RT-qPCR

The RT-qPCR technique was performed to analyze the *NlAtg3* expression subsequent to the injection of different dsRNAs. For this purpose, primers (Supplementary Table S1) specific to synthesizing ds*NlAtg3*-474×1-qf and ds*NlAtg3*-474×1-qr were designed with the Primer Premier 5.0 software. The RT-qPCR reaction systems (20 μ) contained 10 μL of

TB green RemixExTaqII, forward and reverse primers (10 μM) (0.4 μL), cDNA as a template (2.0 μL), ROX reference dye (0.4 μL), and water (6.8 μL). The PCR reactions were performed using the Step One Plus real-time PCR system (Bio-Rad, Hercules, CA, USA). The reference gene was 18s rRNA (GenBank accession number: JN662398.1) of *N. lugens* (*Nl18S*). Each sample contains 3 technical replicates. The $2^{-\Delta\Delta Ct}$ method was used to detect the relative expression levels of genes [24].

2.5. Effect of RNAi on the Survival of Non-Target Organisms

The 5th-instar insects of *S. furcifera* were separately injected with ds*NlAtg3*-474×1, ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10 between 1 and 12 h after molting, and ds*GFP* was introduced by microinjection as a control. Through the thorax, at the site between the middle and hind legs, 250 ng of dsRNA per insect was injected, and the insects were transferred to feed on TN1 rice plants [25]. Each treatment was set up with three duplicates, each consisting of 30 *N. lugens* insects. The survival rates of each group were tracked daily. As a non-target insect, *D. melanogaster* larvae of the third instar were injected with 250 ng of ds*NlAtg3*-47×10 (injected into the larvae's abdomen as described above) [26]. Prior to injection, *D. melanogaster* was frozen on ice for 5–10 min. The injected larvae were transferred to the Drosophila medium, and their mortality, growth and development were observed and recorded.

2.6. Effect of RNAi on Natural Enemies Feeding dsRNA-Treated N. lugens

Three *N. lugens* insects, injected with ds*NlAtg3*-47×10, were first released in a tube, and 24 h later, one *D. sulfureus* was released into the tube to feed these injected *N. lugens*. During the following days, three additional injected *N. lugens* were added to each tube daily. In total, 10 tubes were prepared for ds*NlAtg3*-47×10, and 10 tubes were prepared for ds*GFP* as a control. The mortality of the *D. sulfureus* was observed and recorded every day. *T. chinensis* was treated as described above for *D. sulfureus*.

2.7. Statistical Analysis

For gene transcript levels, the data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test. The comparison of the survival curve was conducted by the Log-rank test. All statistical analyses were conducted on SPSS 22.0 software (http://www.spss.com, accessed on 6 September 2024).

3. Results

3.1. Spatiotemporal Expression Patterns of NlATG3 Gene in N. lugens

To verify the expression of the *NlATG3* gene in *N. lugens*, its spatial and temporal expression was determined by RT-qPCR. The results showed that *NlATG3* expression was relatively stable in nymphs, but was lower in female adults and significantly higher in male adults ($p < 0.01$) (Figure 2A). The expression level of *NlATG3* was the highest in the testis (2.02), followed by the cuticle (1.92) and the fat body (1.62), while it was the lowest in the ovary (0.40). It was medium in the head (set to 1), thorax (0.77), and gut (0.80) (Figure 2B). Therefore, the *NlATG3* gene was expressed in different tissues or parts at all developmental ages.




Figure 2. Spatiotemporal expression patterns of the *N. lugens NlATG3* gene. (**A**) Relative expression levels of the *NlATG3* gene at different developmental stages.1st: 1st-instar nymphs; 2nd: 2nd-instar nymphs; 3rd: 3rd-instar nymphs; 4th: 4th-instar nymphs; 5th: 5th-instar nymphs; Female: female adult; Male: male adult. (**B**) Relative expression of *NlATG3* gene in different tissue sites. The data in the figure represent the mean ± SEM of three biological replicates, using one-way ANOVA and multiple comparison test by Tukey's method, and different lowercase letters over the bars in the figure represent significant differences.

3.2. Sequence Analysis of dsNlAtg3 Fragments with Different Lengths

The alignment results showed that ds*NlAtg3*-474×1 of *N. lugens* had 84.07% similarity with that of *S. furcifera*, 51.43% with *D. melanogaster*, and 35.17% with *T. chinensis*. For ds*NlAtg3*-474×1 of *N. lugens*, the longest consecutive identical bases were 27, 16, and 13 in *S. furcifera*, *D. melanogaster*, and *T. chinensis*, respectively (Figure 3A). The ds*NlAtg3*-138×3 of *N. lugens* had 86.52% similarity with that *S. furcifera*, 55.78% with *D. melanogaster*, and 65.28% with *T. chinensis*, respectively. For ds*NlAtg3*-138×3 of *N. lugens*, the longest consecutive identical bases were 25, 8, and 13 in *S. furcifera*, *D. melanogaster*, and *T. chinensis*, respectively (Figure 3B). ds*NlAtg3*-47×10 of *N. lugens* had 78% similarity with that of *S. furcifera*, 32% with *D. melanogaster*, and 40% with *T. chinensis*, respectively. For ds*NlAtg3*-47×10 of *N. lugens*, the longest consecutive identical bases were 12, 5, and 3 in *S. furcifera*, *D. melanogaster*, and *T. chinensis*, respectively (Figure 3C). The sequence of *Atg3* in *D. sulfureus* is not yet available, and is not included in the alignment. However, low conservation is expected between *D. sulfureus* and *N. lugens*, taking *Atg3*-47×10 into consideration. As a spider, *D. sulfureus* has a relatively distant genetic relationship with the insect *N. lugens*, compared with the other insects *S. furcifera*, *D. melanogaster*, and *T. chinensis*. By reducing the longest consecutive identical sequence in dsRNA sequence, it is possible to avoid harm to non-target organisms.




Figure 3. Sequence alignment of different *NlAtg3* fragments with different species. (**A**) ds*NlAtg3*-474×1. (**B**) ds*NlAtg3*-138×3. (**C**) ds*NlAtg3*-47×10. The black color represents that all four sequences contain the same base. The number of identical bases in the four segments gradually decreases, and the color gradually becomes lighter. The *represent every ten-base spacer.

3.3. RNAi Efficiency of Different dsRNA Fragments on the Expression of NlAtg3 in N. lugens

The effect of RNAi on the expression of the target gene *NlAtg3* was analyzed by injecting fifth-instar *N. lugens* nymphs with 250 ng of dsRNA. RT-qPCR revealed that the mRNA expression of *NlAtg3* in all treated groups was significantly suppressed compared with the ds*GFP* control. On day 1, treatment with ds*NlAtg3*-474×1, ds*NlAtg3*-138×3, and ds*NlAtg3*-47×10 resulted in large decreases in the mRNA levels of 78.5%, 88.2%, and 93.7%, respectively (Figure 4). It also showed that dsRNA containing more concatemerized repeats had a better inhibitory effect at this stage, that is, ds*NlAtg3*-47×10 > ds*NlAtg3*-138×3 > ds*NlAtg3*-474×1. From day 2 on, the mRNA expression of the three treated groups did not show significant differences between different treatments, but dsRNA containing more ds*NlAtg3* repeats still showed a better inhibitory effect than that containing fewer

repeats on day 2. On day 4 post injection, the expression of the target gene in all treated groups remained at very low levels; treatment with ds*NlAtg3*-474×1, ds*NlAtg3*-138×3, and ds*NlAtg3*-47×10 decreased the expression by 95.33%, 97.09%, and 94.84%, respectively. These results showed that the expression of the target *NlAtg3* gene in *N. lugens* was successfully knocked down through RNAi.




Figure 4. Expression levels of *NlAtg3* mRNA in the 5th-instar nymphs of *N. lugens* after injection of different kinds of interference fragments. ds*GFP* was used as the control group, and Nl*18s* was used as the internal reference gene for gene expression normalization. n = 5 insects. Bars are the mean ± SEM from three independent experiments. The data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test. Different lowercase letters on the bars indicate significant differences ($p < 0.05$).

3.4. Effect of Different dsRNA Fragments on the Survival of N. lugens

The fifth-instar *N. lugens* larvae were separately injected with ds*NlAtg3*-474×1, ds*NlAtg3*-138×3, or ds*NlAtg3*-47×10, at 250 ng per insect. In the ds*NlAtg3*-474×1 group, nymph survival decreased significantly and the survival rate decreased to 13.91% on day 7, while the survival rate of the ds*GFP* control group still remained at a high level of 93.33% on day 4 (Figure 5A). After injection with ds*NlAtg3*-138×3, the survival rate decreased from day 1, and decreased to 10.51% on day 7 (Figure 5B). A strong effect was also achieved in nymphs injected with ds*NlAtg3*-47×10, with the survival rate reaching 18.89% on day 9. In contrast, the survival rate of the ds*GFP* control group decreased relatively slowly, to 70% on day 9 post injection (Figure 5C). This indicates that the length of the dsRNA fragments was related to the effect of RNAi on the survival of *N. lugens*.




Figure 5. Effect of injecting different lengths of dsRNA fragments on the survival rate of *N. lugens*. (**A**–**C**) Effect of injecting ds*NlAtg3*-474×1, ds*NlAtg3*-138×3 and ds*NlAtg3*-47×10 on the survival rate of *N. lugens*. The comparison of the survival curve was conducted by the Log-rank test. Error bars indicate the standard error (SE) from three independent experiments, with n = 30 insects per treatment. Histogram bars annotated with asterisks are significantly different (***, $p < 0.001$). Significant differences between means (multiple *t*-tests) are depicted.

3.5. Effects of Different dsNlAtg3 Fragments on the Closely Related Species S. furcifera

S. furcifera and *N. lugens* belong to the same family (Delphacidae). By injecting 250 ng of ds*NlAtg3*-474×1, ds*NlAtg3*-138×3, and ds*NlAtg3*-47×10 into each fifth-instar nymph of *S. furcifera*, the impact of different dsRNA fragments on the survival of *S. furcifera* was investigated. The results showed that these three kinds of dsRNA fragments had different effects on the survival rate of *S. furcifera*. After injecting ds*NlAtg3*-474×1, the survival rate declined rapidly to 2.22% on the fifth day post injection. Meanwhile, the survival rate of the ds*GFP* group remained at a high level of 79.26% (Figure 6A). The treatment of ds*NlAtg3*-138×3 achieved similar effects to that of ds*NlAtg3*-474×1 (Figure 6B). Nonetheless, no significant difference in the survival rate was observed between the ds*NlAtg3*-47×10 group (68.89%) and the ds*GFP* control group (78.89%) on day 7 (Figure 6C). This indicates that by designing suitable RNA interference fragments, it is possible to avoid harm to closely related species while effectively inhibiting the target pest *N. lugens*.




Figure 6. Effects of injecting ds*NlAtg3* fragments of different lengths on the survival rate of *S. furcifera*. (**A**) Effect of injecting ds*NlAtg3*-474×1 on the survival rate of white-back planthopperds. (**B**) Effect of injecting ds*NlAtg3*-137×3 on the survival rate of *S. furcifera*. (**C**) Effect of injecting ds*NlAtg3*-47×10 on the survival rate of *S. furcifera*. Injection of ds*GFP* is used for controls. The comparison of the survival curve was conducted by the Log-rank test. Error bars indicate the standard error (SE) from three independent experiments, with n = 30 per treatment. Histogram bars annotated with asterisks are significantly different (***, $p < 0.001$); no asterisks indicate no significant differences among the comparisons.

3.6. Effects of Short Concatemerized dsNlAtg3-47×10 on Non-Target Organism D. melanogaster

D. melanogaster belongs to another genus of insects, *Drosophila*, and has a relatively distant evolutionary relationship with *N. lugens*, compared with *S. furcifera*. After injection of ds*NlAtg3*-47×10, there was no substantial reduction in the survival rate of *Drosophila* (83.33%) compared to the dsGFP control group (86.11%) on the 11th day post injection (Figure 7A). On day 4 after ds*NlAtg3*-47×10 injections, the pupation rate of *Drosophila* reached 86.67%, and it was not significantly different from that of the control group injected with ds*GFP* (Figure 7B). On day 11 after injection of ds*NlAtg3*-47×10, the pupation rate of *D. melanogaster* reached 83.33%, while the pupation rate of its control group ds*GFP* was 86.11%, with no significant difference observed between the two groups in terms of survival rate (Figure 7C). These results suggest that ds*NlAtg3*-47×10 targeted against *N. lugens* had no significant effect on the growth and development of the non-target organism *D. melanogaster*.




Figure 7. Effects of ds*NlAtg3*-47×10 injection on fly survival and growth and development. (**A**) Survival rate. (**B**) Pupation rate. (**C**) Eclosion rate. Injection of dsGFP used as controls. The comparison of the survival curve was conducted by the Log-rank test. Error bars indicate the standard error (SE) from three independent experiments, with n = 20 per treatment. No asterisks indicate no significant differences among the comparisons.

3.7. Effects of dsNlAtg3-47×10 on Natural Enemies of N. lugens

The fifth-instar *N. lugens* nymphs were injected with 250 ng of ds*NlAtg3*-47×10, and 24 h later were fed to its natural enemies *D. sulfureus* and *T. chinensis*. After one week of feeding, it was found that no *D. sulfureus* had died in the ds*NlAtg3*-47×10 group (Figure 8A). The survival rate of the *T. chinensis* in the ds*NlAtg3*-47×10 group was only slightly decreased, with no significant difference found between the two groups (Figure 8B). This suggests that ds*NlAtg3*-47×10 has no obvious effect on the survival of its natural enemies *D. sulfureus* and *T. chinensis* through food chain transmission. These results suggest that ds*NlAtg3*-47×10 targeting against *N. lugens* is safe for the natural enemies *D. sulfureus* and *T. chinensis*.




Figure 8. Effects of ds*NlAtg3*-47×10 injection on *D. sulfureus* and *T. chinensis* survival. Injection of ds*GFP* is used for controls. (**A**) Effect of ds*NlAtg3*-47×10 injection on *D. sulfureus* survival. (**B**) Effect of ds*NlAtg3*-47×10 injection on *T. chinensis* survival. The comparison of the survival curve was conducted by the Log-rank test. No asterisks indicate no significant differences among the comparisons.

4. Discussion

In this study, an autophagy-related gene, *NlAtg3*, was used to synthesize dsRNA fragments with different lengths and sequence compositions. Among the three dsRNA fragments (ds*NlAtg3*-474×1, ds*NlAtg3*-138×3, ds*NlAtg3*-47×10), the ds*NlAtg3*-47×10 fragment was proved to be specific and efficient against *N. lugens*, but was safe for non-target organisms, including two natural enemies (*D. sulfureus* and *T. chinensis*) of *N. lugens*. Therefore, this study establishes a foundation for the subsequent construction and application of rice-mediated RNA interference systems for *N. lugens*.

Previous RNAi experiments on potato beetles showed that the interference efficiency of 200–700 bp dsRNA was higher than that of 60 bp dsRNA in a feeding experiment, but

there was no significant difference between dsRNA fragments with length varying from 200 bp to 700 bp [27]. Our work also supports that long dsRNA fragments (ds*NlAtg3*-474×1 and ds*NlAtg3*-138×3) covering a larger area of the target sequence achieved higher RNAi efficiency than that of the ds*NlAtg3*-47×10 fragment, which only covers a 47 bp area of the target sequence. In addition, transfection of dsRNA ranging from 21 to 592 bp in length resulted in effective target gene silencing in *Drosophila melanogaster* S2-cells, irrespective of the dsRNA length. In contrast, uptake of dsRNAs added to the medium was clearly length-dependent. Importantly, a diverse pool of short siRNAs (enzymatically generated 21-mer siRNAs) failed to enter S2 cells, and did not result in any significant silencing [18]. Currently, the number of 21-mer matches is considered a critical factor in the function of dsRNA transfected in cells, especially in its effects on non-target organisms [28,29]. In this study, the specific dsRNA fragment ds*NlAtg3*-47×10 showed a perfect interference effect on the target pest *N. lugens*, but did not cause significant lethality, neither to the closely related species *S. furcifera*, nor to the natural enemies *D. sulfureus* and *T. chinensis*. For ds*NlAtg3*-47×10 of *N. lugens*, the longest consecutive identical bases in *S. furcifera*, *D. melanogaster*, and *T. chinensis* were 12, 5, and 3, respectively, far fewer than 21 bp (Figure 3C). Therefore, by reducing the longest consecutive identical sequence in the dsRNA sequence, it is possible to avoid harming non-target organisms. Similarly, when conducting a non-target insect risk assessment on the *N. lugens* natural enemy species *C. lividipennis*, it was revealed that three dsRNA segments (ds*NlNa*, ds*NlAup5*, and ds*NlvATP-A*) specifically against *NlNa*, *NlAup5*, and *NlvATP-A* exhibited significant lethality towards *N. lugens*. However, when fed at high concentrations (10×), the survival rate, the spawning rate and adult survival time of non-target *C. lividipennis* did not show a significant impact [30]. It should be noted that very short dsRNA fragments may not be a good approach to reducing the non-target effect. They may bring other issues, for example, lost function for different populations with SNPs, or binding to unspecified genes in the target species and non-target species.

The evaluation criteria for non-target organism effects are important in studies on RNAi risk assessment. The selection of target genes, the dsRNA design, and the evaluation and investigation of the effects on non-target organisms can partly reduce the risk posed by RNAi-based biopesticides [31]. Tan et al. showed that no adverse effects on the growth and development were observed in either *Apis mellifera* L. larvae or adults at long and high *DvSnf7* dsRNA exposure levels [32]. Castellanos et al. demonstrated that feeding the parasitoid wasp *Telenomus*. *podisi* with wasp-specific dsRNA targeting the *vATPase A* and *actin-2* genes led to high mortality, demonstrating that dietary RNAi is functional in *T*. *podisi*. When feeding *T*. *podisi* with its host *Euschistus heros*-specific dsRNA targeting the same genes, no lethal or sublethal effects were observed [33]. Therefore, by reasonably designing and synthesizing suitable dsRNA, it is possible to acquire specific siRNA fragments and effectively improve the specificity and reduce the risk [33]. In future work, we will add some further comparisons with studies on other similar species, and above all, further investigate the effects on more non-target organisms.

Currently, transgenic plants expressing dsRNA have made significant breakthroughs in the management of insect pests. Three genes (*NlHT1*, *Nlcar* and *Nltry*) highly expressed in the midgut of *N*. *lugens* were used to develop dsRNA constructs for transforming rice, and some of the transcribed dsRNA was processed to siRNAs in the transgenic lines. When *N*. *lugens* fed on rice plants expressing dsRNA, the mRNA levels of the targeted genes in the midgut were reduced [10]. It has also been reported that rice plants with *osa-miR162a* overexpression had enhanced resistance to *N. lugens* and reduced the reproductive capacity of adults [11]. Some candidate corn varieties expressing the dsRNA of *DvvSnf7*, *Cry3Bb1* and *Cry34Ab1*/*Cry35Ab1* genes have been shown to cause significant mortality of western corn rootworm via the siRNA signaling pathway [8,9]. During a risk assessment of

siRNA-mediated IRGE rice, it was found that the IRGE rice targeting the *dib* 3′ untranslated sequence had significant resistance to its target pest *C. Suppressalis*, but was sensitive to non-target organisms, and the impact on *A. mellifera* could be ignored [34]. When Baum et al. fed western corn rootworm (WCR) with transgenic corn plants which express WCR dsRNAs, there was a significant reduction in WCR feeding damage [35]. Therefore, RNAi-based, genetically engineered insect-resistant rice carrying species-specific dsRNA like ds*NlAtg3*-47×10, is expected to have good application potential.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/insects16020225/s1.

Author Contributions: Conceptualization, P.H. and X.Y.; methodology, K.L. and T.C.; software, K.L. and T.C.; validation, K.L. and T.C.; investigation, K.L., T.C., Y.L., K.S. and K.P.; resources, P.H. and X.Y.; data curation, K.L. and T.C.; preparation of original draft, K.L., T.C. and P.H.; writing—review and editing, P.H., K.L. and X.Y.; project administration, K.L., T.C. and P.H.; funding acquisition, P.H. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by the Major Research plan of the Natural Science Foundation of Zhejiang Province, China (grant numbers LD25C140001), the NSFC (grant numbers U21A20223, 32472649), and the Zhejiang Province Leading Earth Goose Program (grant number 2022C02047).

Data Availability Statement: The raw data supporting the conclusions of this article will be made available by the authors on request.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Foissac, X.; Thi Loc, N.; Christou, P.; Gatehouse, A.M.R.; Gatehouse, J.A. Resistance to Green Leafhopper *(Nephotettix virescens*) and Brown Planthopper (*Nilaparvata lugens*) in Transgenic Rice Expressing Snowdrop Lectin (*Galanthus nivalis* Agglutinin; GNA). *J. Insect Physiol.* **2000**, *46*, 573–583. [CrossRef] [PubMed]
2. Cheng, J.A. Rice Planthopper Problems and Relevant Causes in China. In *Planthoppers: New Threats to the Sustainability of Intensive Rice Production Systems in Asia*; International Rice Research Institute: Manila, Philippines, 2009; Volume 157, p. 178.
3. Rivera, C.T.; Ou, S.H.; Iida, T.T. Grassy Stunt Disease of Rice and Its Transmission by the Planthopper *Nilaparvata lugens* Stal. *Plant Dis. Report.* **1966**, *50*, 453–456.
4. Fujii, T.; Sanada-Morimura, S.; Oe, T.; Ide, M.; Van Thanh, D.; Van Chien, H.; Van Tuong, P.; Loc, P.M.; Cuong, L.Q.; Liu, Z.; et al. Long-term Field Insecticide Susceptibility Data and Laboratory Experiments Reveal Evidence for Cross Resistance to Other Neonicotinoids in the Imidacloprid-resistant Brown Planthopper *Nilaparvata lugens*. *Pest. Manag. Sci.* **2020**, *76*, 480–486. [CrossRef]
5. Gorman, K.; Liu, Z.; Denholm, I.; Brüggen, K.-U.; Nauen, R. Neonicotinoid Resistance in Rice Brown Planthopper, *Nilaparvata lugens*. *Pest. Manag. Sci. Former. Pestic. Sci.* **2008**, *64*, 1122–1125. [CrossRef]
6. Fire, A.; Xu, S.; Montgomery, M.K.; Kostas, S.A.; Driver, S.E.; Mello, C.C. Potent and Specific Genetic Interference by Double-Stranded RNA in *Caenorhabditis elegans*. *Nature* **1998**, *391*, 806–811. [CrossRef]
7. Eaton, B.A.; Fetter, R.D.; Davis, G.W. Dynactin Is Necessary for Synapse Stabilization. *Neuron* **2002**, *34*, 729–741. [CrossRef]
8. Johnson, K.D.; Campbell, L.A.; Lepping, M.D.; Rule, D.M. Field Trial Performance of Herculex XTRA (Cry34Ab1/Cry35Ab1) and SmartStax (Cry34Ab1/Cry35Ab1 + Cry3Bb1) Hybrids and Soil Insecticides Against Western and Northern Corn Rootworms (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* **2017**, *110*, 1062–1069. [CrossRef]
9. Bachman, P.M.; Huizinga, K.M.; Jensen, P.D.; Mueller, G.; Tan, J.; Uffman, J.P.; Levine, S.L. Ecological risk assessment for DvSnf7 RNA: A plant-incorporated protectant with targeted activity against western corn rootworm. Regulatory toxicology and pharmacology. *Regul. Toxicol. Pharmacol.* **2016**, *81*, 77–88. [CrossRef]
10. Zha, W.; Peng, X.; Chen, R.; Du, B.; Zhu, L.; He, G. Knockdown of Midgut Genes by dsRNA-Transgenic Plant-Mediated RNA Interference in the Hemipteran Insect *Nilaparvata lugens*. *PLoS ONE* **2011**, *6*, e20504. [CrossRef]
11. Shen, W.; Cao, S.; Liu, J.; Zhang, W.; Chen, J.; Li, J.-F. Overexpression of an Osa-miR162a Derivative in Rice Confers Cross-Kingdom RNA Interference-Mediated Brown Planthopper Resistance without Perturbing Host Development. *Int. J. Mol. Sci.* **2021**, *22*, 12652. [CrossRef] [PubMed]

12. Yang, J.; Sun, X.-Q.; Zhu-Salzman, K.; Qin, Q.-M.; Feng, H.-Q.; Kong, X.-D.; Zhou, X.-G.; Cai, Q.-N. Host-Induced Gene Silencing of Brown Planthopper Glutathione S-Transferase Gene Enhances Rice Resistance to Sap-Sucking Insect Pests. *J. Pest. Sci.* **2021**, *94*, 769–781. [CrossRef]
13. Zhang, H.; Li, H.; Miao, X. Feasibility, Limitation and Possible Solutions of RNAi-based Technology for Insect Pest Control. *Insect Sci.* **2013**, *20*, 15–30. [CrossRef] [PubMed]
14. Katoch, R.; Sethi, A.; Thakur, N.; Murdock, L.L. RNAi for Insect Control: Current Perspective and Future Challenges. *Appl. Biochem. Biotech.* **2013**, *171*, 847–873. [CrossRef] [PubMed]
15. Ding, S.-W. RNA-Based Antiviral Immunity. *Nat. Rev. Immunol.* **2010**, *10*, 632–644. [CrossRef]
16. Nandety, R.S.; Kuo, Y.-W.; Nouri, S.; Falk, B.W. Emerging Strategies for RNA Interference (RNAi) Applications in Insects. *Bioengineered* **2015**, *6*, 8–19. [CrossRef]
17. Terenius, O.; Papanicolaou, A.; Garbutt, J.S.; Eleftherianos, I.; Huvenne, H.; Kanginakudru, S.; Albrechtsen, M.; An, C.; Aymeric, J.-L.; Barthel, A.; et al. RNA Interference in Lepidoptera: An Overview of Successful and Unsuccessful Studies and Implications for Experimental Design. *J. Insect Physiol.* **2011**, *57*, 231–245. [CrossRef]
18. Saleh, M.C.; Van Rij, R.P.; Hekele, A.; Gillis, A.; Foley, E.; O'Farrell, P.H.; Andino, R. The Endocytic Pathway Mediates Cell Entry of dsRNA to Induce RNAi Silencing. *Nat. Cell Biol.* **2006**, *8*, 793–802. [CrossRef]
19. Yi, S.; Wang, L.; Ho, M.S.; Zhang, S. The autophagy protein Atg9 functions in glia and contributes to parkinsonian symptoms in a Drosophila model of Parkinson's disease. *Neural Regen. Res.* **2024**, *19*, 1150–1155. [CrossRef]
20. Issa, A.R.; Sun, J.; Petitgas, C.; Mesquita, A.; Dulac, A.; Robin, M.; Mollereau, B.; Jenny, A.; Chérif-Zahar, B.; Birman, S. The lysosomal membrane protein LAMP2A promotes autophagic flux and prevents SNCA-induced Parkinson disease-like symptoms in the *Drosophila* brain. *Autophagy* **2021**, *14*, 1898–1910. [CrossRef]
21. Fang, D.; Xie, H.; Hu, T.; Shan, H.; Li, M. Binding Features and Functions of ATG3. *Front. Cell Dev. Biol.* **2021**, *9*, 685625. [CrossRef]
22. Ye, C.; Feng, Y.; Yu, F.; Jiao, Q.; Wu, J.; Ye, Z.; Zhang, P.; Sun, C.; Pang, K.; Hao, P.; et al. RNAi-mediated Silencing of the Autophagy-related GeneNIATG3 Inhibits Survival and Fecundity of the Brown Planthopper, *Nilaparvata lugens*. *Pest. Manag. Sci.* **2021**, *77*, 4658–4668. [CrossRef] [PubMed]
23. Liu, S.; Ding, Z.; Zhang, C.; Yang, B.; Liu, Z. Gene Knockdown by Intro-Thoracic Injection of Double-Stranded RNA in the Brown Planthopper, *Nilaparvata lugens*. *Insect Biochem. Mol. Biol.* **2010**, *40*, 666–671. [CrossRef] [PubMed]
24. Hu, K.; Tian, P.; Yang, L.; Qiu, L.; He, H.; Ding, W.; Li, Z.; Li, Y. Knockdown of *Methoprene-Tolerant* Arrests Ovarian Development in the *Sogatella furcifera* (Hemiptera: Delphacidae). *J. Insect Sci.* **2019**, *19*, 5. [CrossRef]
25. Wang, Z.; Yang, H.; Zhou, C.; Yang, W.; Jin, D.; Long, G. Molecular Cloning, Expression, and Functional Analysis of the Chitin Synthase 1 Gene and Its Two Alternative Splicing Variants in the White-Backed Planthopper, *Sogatella furcifera* (Hemiptera: Delphacidae). *Sci. Rep.* **2019**, *9*, 1087. [CrossRef]
26. Powell, M.; Pyati, P.; Cao, M.; Bell, H.; Gatehouse, J.A.; Fitches, E. Insecticidal Effects of dsRNA Targeting the Diap1 Gene in Dipteran Pests. *Sci. Rep.* **2017**, *7*, 15147. [CrossRef]
27. He, W. The Effects of dsRNA Molecule on the RNAi Efficiency Against *Leptinotarsa decemlineata* and Its Application in Plastid-Mediated RNAi for Pest Control. Doctoral Dissertation, Hubei University, Wuhan, China, 2022. [CrossRef]
28. Haller, S.; Widmer, F.; Siegfried, B.D.; Zhuo, X.; Romeis, J. Responses of two ladybird beetle species (Coleoptera: Coccinellidae) to dietary RNAi. *Pest. Manag. Sci.* **2019**, *75*, 2652–2662. [CrossRef]
29. Poreddy, S.; Li, J.; Baldwin, I.T. Plant-mediated RNAi silences midgut-expressed genes in congeneric lepidopteran insects in nature. *BMC Plant Biol.* **2017**, *17*, 199. [CrossRef]
30. Dang, C.; Zhang, Y.; Sun, C.; Li, R.; Wang, F.; Fang, Q.; Yao, H.; Stanley, D.; Ye, G. dsRNAs Targeted to the Brown Planthopper *Nilaparvata lugens*: Assessing Risk to a Non-Target, Beneficial Predator, *Cyrtorhinus lividipennis*. *J. Agric. Food Chem.* **2022**, *70*, 373–380. [CrossRef]
31. Luo, X.; Nanda, S.; Zhang, Y.; Zhou, X.; Yang, C.; Pan, H. Risk assessment of RNAi-based biopesticides. *New Crops* **2024**, *1*, 2949–9526. [CrossRef]
32. Tan, J.; Levine, S.L.; Bachman, P.M.; Jensen, P.D.; Mueller, G.M.; Uffman, J.P.; Meng, C.; Song, Z.; Richards, K.B.; Beevers, M.H. No impact of DvSnf7 RNA on honey bee (*Apis mellifera* L.) adults and larvae in dietary feeding tests. *Environ. Toxicol. Chem.* **2016**, *35*, 287–294. [CrossRef]
33. Castellanos, N.L.; Smagghe, G.; Taning, C.N.T.; Oliveira, E.E.; Christiaens, O. Risk assessment of RNAi-based pesticidesto non-target organisms: Evaluating the effects of sequence similarity in the parasitoid wasp *Telenomus podisi*. *Sci. Total Environ.* **2022**, *832*, 154746. [CrossRef]

34. Chen, J.; Wang, H.; Yang, X.; Chen, G.; Du, L.; Chen, H.; Li, Y.; Peng, Y.; Han, L. Consumption of miRNA-Mediated Insect-Resistant Transgenic Rice Pollen Does Not Harm *Apis mellifera* Adults. *J. Agric. Food Chem.* **2021**, *69*, 4234–4242. [CrossRef] [PubMed]
35. Baum, J.A.; Bogaert, T.; Clinton, W.; Heck, G.R.; Feldmann, P.; Ilagan, O.; Johnson, S.; Plaetinck, G.; Munyikwa, T.; Pleau, M.; et al. Control of Coleopteran Insect Pests through RNA Interference. *Nat. Biotechnol.* **2007**, *25*, 1322–1326. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Characterization and Functional Analysis of Small Heat Shock Protein Genes (*Hsp22.2* and *Hsp26.7*) in *Sitodiplosis mosellana* Diapause

Qitong Huang [1,2], Qian Ma [2], Xiaobin Liu [1], Keyan Zhu-Salzman [3,*] and Weining Cheng [2,*]

[1] Shandong Institute of Sericulture, Shandong Academy of Agricultural Sciences, Yantai 265503, China; 13760637249hqt@nwafu.edu.cn (Q.H.); bin0331@126.com (X.L.)
[2] Key Laboratory of Plant Protection Resources and Pest Management of Ministry of Education, College of Plant Protection, Northwest A&F University, Yangling 712100, China; maqian981017@163.com
[3] Department of Entomology, Texas A&M University, College Station, TX 77843, USA
* Correspondence: ksalzman@tamu.edu (K.Z.-S.); chengwn@nwsuaf.edu.cn (W.C.)

Simple Summary: Small heat shock proteins (sHsps) are crucial players not only in heat/cold adaptation in insects, but also in insect diapause. This study aimed to characterize two sHsp genes (*SmHsp22.2* and *SmHsp26.7*) from *S. mosellana*, a serious wheat pest that undergoes long obligatory larval diapause to survive temperature extremes during summer and winter, and determine their expression in association with diapause and thermal stress, as well as their roles in cold stress. The results showed that *SmHsp22.2* and *SmHsp26.7* were significantly upregulated during diapause including short-term heat/cold stress. RNAi-mediated knockdown of both genes significantly increased the mortality of *S. mosellana* larvae under cold stress. These results suggest that *SmHsp22.2* and *SmHsp26.7* form an integral part of adaptations to diapause in *S. mosellana*, and that their increased expression is essential for protecting *S. mosellana* from the adverse effects of low temperature.

Abstract: Small heat shock proteins (sHsps) play crucial roles in organismal adaptation to stress tolerance. *Sitodiplosis mosellana*, a devastating insect wheat pest, undergoes long obligatory larval diapause to survive temperature extremes during summer and winter. To elucidate the function of sHsps in this process, two sHsp-encoding genes (*SmHsp22.2* and *SmHsp26.7*) were characterized from *S. mosellana*, and their responsiveness to diapause and thermal stress, as well as their roles in cold stress, was analyzed. Both *SmHsp22.2* and *SmHsp26.7* possessed the canonical α-crystallin domain and lacked introns. Quantitative PCR indicated significant upregulation of *SmHsp22.2* and *SmHsp26.7* during diapause, especially in summer and winter. Notably, *SmHsp22.2* exhibited higher expression in summer relative to winter, whereas *SmHsp26.7* showed the opposite profile. Moreover, short-term heat shock ($\geq$35 °C) in over-summering larvae or cold shock (≤ -10 °C) in over-wintering larvae was found to trigger transcriptional upregulation of both genes, while prolonged temperature extremes (i.e., 45–50 °C or −15 °C) did not elicit a comparable response. RNA interference-mediated knockdown of both genes significantly increased the mortality of *S. mosellana* larvae under cold stress. These findings indicate the importance of both *SmHsps* in diapause and environmental adaptation in *S. mosellana*.

Keywords: small heat shock protein; diapause; *Sitodiplosis mosellana*; temperature stress; cold adaptation

1. Introduction

Diapause is a developmental tactic that ensures the survival of insects in extreme environments, synchronizing their life cycles with times of the year conducive to growth and reproduction [1,2]. Diapause can occur at various life stages according to the insect species [3–6], and involves significant physiological and metabolic shifts, including reductions in metabolic rates, augmentation of energy reserves, and enhanced resistance to stress [7,8]. Given its critical importance in insect growth and reproduction, unraveling the molecular underpinnings of diapause is paramount for formulating effective agricultural pest-control strategies [9].

Heat shock proteins (Hsps) are upregulated in response to various biotic and abiotic stresses, including pathogen invasion, heat, and cold [10–13]. Hsps function as molecular chaperones, and promote cellular protein homeostasis by binding and refolding stress-compromised proteins [14–16]. They are classified into various families according to evolutionary relationships and size, including the Hsp100, Hsp90, Hsp70, Hsp60, Hsp40, and small Hsp (sHsp) families [17–19].

sHsps are a large and ancient family of ATP-independent molecular chaperones with monomeric molecular weights between 12 and 43 kDa. Despite variations in their sequences and oligomer sizes, all sHsps contain a highly conserved core sequence of approximately 80–100 amino acids, the α-crystallin domain, which is bordered by non-conserved N- and C-terminal extensions [20,21]. The α-crystallin domain comprises a β-sandwich consisting of 7 to 8 antiparallel β-strands, and is responsible for the chaperone activity of sHsps [22]. Beyond their roles as chaperones, sHsps engage in a diverse array of biological processes, such as maintenance of the cytoskeleton [23], DNA repair [24], cellular apoptosis and autophagy [25], and membrane stability [26].

Recent studies have demonstrated the roles of sHsps in the regulation of insect diapause; however, their expression patterns differ markedly according to the insect species and sHsp family [27,28]. For example, in the blowfly *Lucilia sericata*, *Hsp23* transcript levels are not altered by diapause [29], in contrast to changes in *Hsp23* expression seen during pupal diapause in the flesh fly *Sarcophaga crassipalpis* [30] and larval diapause of the blowfly *Calliphora vicina* [31]. Furthermore, elevated expression of *sHsps* during diapause has been reported in a variety of insect species, including *Calanus finmarchicus Hsp22* [32], *Leguminivora glycinivorella Hsp19.8* and *18.9* [33], *Pieris melete Hsp19.5* and *20.0* [27], and *Ostrinia nubilalis Hsp20.1* [34]. In contrast, downregulation of *Sesamia nonagrioides Hsp20.8* and *Trogoderma granarium Hsp20.5* and *21.2* was observed during diapause [28,35]. These variations in expression suggest differences in sHsp function among insect species during diapause.

The orange wheat blossom midge *Sitodiplosis mosellana* (Géhin) (Diptera: Cecidomyiidae), a notorious pest of wheat across the northern hemisphere, causes substantial reductions in yield and economic damage during peak infestation years [36–38]. As a univoltine species, the mature larvae in northern China undergo obligatory diapause within cocoons in the soil during the 10 months from June to early April of the following year. The initiation of post-diapause development is triggered by the onset of increased ambient temperatures in the spring [39]. This prolonged diapause not only enables the survival of the insect during hot summer and cold winter temperatures but also ensures synchronization of its development with wheat growth cycles. While we have previously elucidated the expression patterns of *Hsp17.4* and *Hsp20.3* in *S. mosellana* diapause [40], the roles of other sHsps in regulating this process are still unclear.

In this study, we cloned and characterized two novel sHsp-encoding genes from the pre-diapause larvae of *S. mosellana*, and analyzed their expression profiles under conditions of diapause and thermal stress. In addition, the effects of *SmHsp22.2*/*Hsp26.7*

knockdown on cold tolerance in *S. mosellana* were also investigated. The findings offer valuable molecular insights into stress tolerance associated with diapause in *S. mosellana*.

2. Materials and Methods

2.1. Insect Collection

The *S. mosellana* larvae utilized in the study were collected from natural settings, as previously described [41]. Specifically, pre-diapause larvae were obtained by dissection of *S. mosellana*-infested wheat spikes collected from wheat fields on 25 May 2022. The wheat ears containing mature third-instar larvae were bulk-collected and transferred to a field insectary in Yangling, Shaanxi, China (34°16′ N, 108°4′ E), where they were placed on damp soil to promote entry into diapause. Successful diapause entry is characterized by the presence of a larval cocoon. Prior research has shown that larvae in cocoons collected from December onwards emerge successfully as adults upon exposure to temperatures of 25 °C, indicating the termination of diapause and transition into post-diapause quiescence by December [42]. Larvae at various stages, including both larvae in diapause and those in post-diapause quiescence (i.e., cocooned larvae), as well as post-diapause developmental stages (i.e., larvae emerging from cocoons), were collected in succession by monthly sieving of the soil in the insectary between late June 2022 to early April 2023. The collected larvae were immediately frozen in liquid nitrogen and subsequently stored at −80 °C until analysis.

2.2. RNA Extraction, cDNA Synthesis, and gDNA Isolation

Total RNA was extracted from the entire bodies of a cohort of 20 pre-diapause *S. mosellana* larvae using TRIzol reagent (TaKaRa, Dalian, China). RNA integrity was assessed with 1% agarose gel electrophoresis, with quantification using a spectrophotometer. One microgram of total RNA was then used as a template for reverse transcription into cDNA using a PrimeScript^{TM} RT Reagent Kit with gDNA Eraser (TaKaRa, Dalian, China), as directed. For the extraction of genomic DNA, the Biospin Insect Genomic DNA Extraction Kit was utilized following the manufacturer's instructions (Bioer Technology Co., Ltd., Hangzhou, China).

2.3. Cloning of the Opening Reading Frames of SmHsp22.7 and SmHsp26.7

Based on the de novo transcriptome data of *S. mosellana* larvae obtained previously, gene-specific primers targeting *Hsp22.2* and *Hsp26.7* (Table 1) were designed to amplify the open reading frames (ORFs). The RT-PCR amplification protocol consisted of a pre-denaturation step (95 °C, 3 min), followed by 32 cycles of denaturation (95 °C, 30 s), annealing (56 °C, 30 s), and extension (72 °C, 45 s), concluding with a final extension at 72 °C for 10 min. The resultant PCR products were fractionated on 1% agarose gels, and bands of the anticipated sizes were excised and purified using a gel extraction kit (Tiangen, Beijing, China). The purified DNA fragments were thereafter integrated into a pEASYR-Blunt Zero cloning vector and transferred into Trans1-T1 competent cells.

Table 1. Primer sequences used in this study.

Primer Name	Sequence (5′ to 3′)	Purpose
Hsp22.2 sense	CTAAAGTGAAGTAGAAAAAATGG	ORF and gDNA cloning
Hsp22.2 antisense	GCATCACATCTTTTACATTCC	
Hsp26.7 sense	ATGAAGTATTTCTCCGTTTTGG	
Hsp26.7 antisense	TTAGGCCTTTAGTTTTTCATCC	

Table 1. *Cont.*

Primer Name	Sequence (5′ to 3′)	Purpose
dsHsp22.2 sense	taatacgactcactatagggTGTTTCACGACACTTCAGCC	dsRNA synthesis
dsHsp22.2 antisense	taatacgactcactatagggGGCTCCAGTTTGTTGGATGT	
dsHsp26.7 sense	taatacgactcactatagggAGATTGTTGGCTCACTCGCT	
dsHsp26.7 antisense	taatacgactcactatagggCTTGGCGTTCACCACAATCG	
dsGFP sense	taatacgactcactatagggTGACCACCCTGACCTAC	
dsGFP antisense	taatacgactcactatagggTTGATGCCGTTCTTCTGC	
Hsp22.2 sense	ATTGCCATCGTTGTTCTG	qPCR
Hsp22.2 antisense	TCCATCTTCGGGTGTGCT	
Hsp26.7 sense	CGATTGTGGTGAACGCCAAG	
Hsp26.7 antisense	CAATTTGGCGCACGTTGGAT	
GAPDH sense	CCATCAAAGCAAGCAAGA	
GAPDH antisense	CAGCACGGAGCACAAGAC	

2.4. Bioinformatics

The functional domains in the sequences of *S. mosellana Hsp22.2* and *Hsp26.7* were predicted using the Conserved Domain Database (https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi, accessed on 15 October 2023), and the molecular weights and isoelectric points of the proteins were determined using the Compute pI/Mw tool (https://web.expasy.org/compute_pi/, accessed on 15 October 2023). Multiple sequence alignment was performed with DNAMAN software package (Version 8.0; Lynnon Bio-Soft, QC, Canada). Phylogenetic relationships among SmHsp22.2, SmHsp26.7, and sHsps from other insect taxa were assessed by the construction of a neighbor-joining phylogenetic tree using MEGA (version X) (https://www.megasoftware.net/, accessed on 15 May 2023). Secondary and tertiary protein structures were predicted using ESPrigt (http://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi, accessed on 15 October 2023) and the SWISS-MODEL web server (http://swissmodel.expasy.org/, accessed on 15 October 2023), respectively. The representation and visualization of amino acid chains in the protein tertiary structures, specifically the α-helices and β-strands, were performed using PyMOL version 2.5.2 (Schrödinger, LLC, New York, NY, USA).

2.5. Heat/Cold Shock Treatments

S. mosellana typically over-summers and over-winters as cocooned larvae at depths of 3 to 10 cm beneath the soil surface [43]. The climate of Yangling district in Shaanxi Province, China (34°16′ N, 108°4′ E) is characterized by temperatures below 30 °C in summer and above 0 °C in winter. However, the soil surface temperatures can reach or exceed 45 °C in summer and may fall to −15 °C in winter, which is above the average super-cooling point (−23.6 °C) for *S. mosellana* cocoons [44]. To clarify the potential roles of *SmHsp26.7* and *SmHsp22.2* under these circumstances, cocoons were subjected to heat treatment (35-50 °C) in August and cold treatment (−15-0 °C) in December as outlined below.

Groups of 20 cocoons newly collected at the end of August or December were placed individually in 1.5 mL polystyrene tubes. For heat shock treatment, the tubes with the August-collected cocoons were immersed in water baths set at 35, 40, 45, or 50 °C for 1 h, or in a 35 °C water bath for 15-90 min. Likewise, tubes containing the December-collected cocoons were exposed to incubator temperatures of 0, −5, −10, or −15 °C for 1 h or −10 °C for 15-90 min. Untreated larvae represented the control group. After treatment, the samples were immediately frozen in liquid nitrogen, followed by storage at −80 °C until further analysis. All treatments were conducted in triplicate with 20 individuals per replicate.

2.6. Quantitative Real-Time PCR (qRT-PCR) Analysis

qRT-PCR analyses were conducted to assess the responsiveness of the two *S. mosellana sHsp* genes to diapause and thermal stress. qRT-PCR parameters were as follows: *Hsp22.2* (sense primer Tm = 55 °C, antisense primer Tm = 58 °C; amplicon size= 163 bp); *Hsp26.7* (sense primer Tm = 60 °C, antisense primer Tm= 60 °C; amplicon size= 195 bp); *GAPDH* (sense primer Tm = 55 °C, antisense primer Tm = 58 °C; amplicon = 86 bp). Total RNA from various samples, including larvae at distinct diapause stages and diapause larvae treated with heat or cold stress, was isolated and reverse-transcribed into cDNA as described above. The qPCR reaction volume (20 μL) contained 10 μL of 2 × SuperReal PreMix Plus (TIANGEN, Beijing, China), 1.0 μL of synthesized cDNA, 0.8 μL of each of the sense and antisense gene-specific primers (10 μM) (Table 1), and 8.2 μL of nuclease-free water. Amplification was performed on an iQ5 real-time PCR system (Bio-Rad, Hercules, CA, USA) with steps comprising pre-denaturation (95 °C, 30 s), amplification (95 °C, 5 s; 60 °C, 32 s, 35 cycles), and melting curve analysis (95 °C, 15 s; 60 °C, 60 s; 40 °C, 30 s). Each reaction was conducted in triplicate with three replicates each. RNase-free water was employed as a negative control in place of the cDNA template to ensure the accuracy and reliability of the results. The relative expression levels of *S.mosellana Hsp22.2* and *Hsp26.7* were standardized against those of the reference gene, *GAPDH* (GenBank accession number: KR733066), expressed constitutively throughout diapause, and the relative mRNA levels were quantified using the $2^{-\Delta\Delta CT}$ method.

2.7. dsRNA Synthesis and RNA Interference

Double-stranded RNA (dsRNA)-specific primers with T7 promoter sequences for *Hsp26.7* and *Hsp22.2* (Table 1) were designed using SnapDragon dsRNA design software (https://www.flyrnai.org/snapdragon, accessed on 15 May 2023), and the corresponding gene fragments were obtained through RCR amplification as stated above. After recovery and purification of the PCR products, dsRNA synthesis was performed using the T7 High Yield Transcription Kit (Vazyme Biotech Co., Ltd., Nanjing, China), as directed, followed by further purification. A solution was prepared with 10 μL dsRNA product, 80 μL nuclease-free water, and 10 μL 3 M sodium acetate (pH 5.2). This was followed by the addition of 100 μL of phenol-chloroform, which was mixed gently and allowed to stand for 3 min before centrifugation at 12,000 rpm for 5 min. The supernatant was removed and an equal volume of isopropanol was added and incubated at −20 °C for 2 h. The material was centrifuged at 12,000 rpm for 15 min at 4 °C, and the supernatant was discarded. Next, 1 mL 75% ethanol was added, followed by centrifugation at 12,000 rpm for 15 min at 4 °C. After removal of the supernatant, the pellet was in nuclease-free water and allowed to air-dry. The synthesis and purification steps of double-strand green fluorescent protein (dsGFP) as the negative control were similar to those described above.

After quantification of the dsRNA concentration using a spectrophotometer Nanodrop2000c (Thermo Fisher Scientific, West Palm Beach, FL, USA), high-concentration dsRNAs were diluted in nuclease-free water to 10 μg/μL. Cocooned larvae that had been collected in January (post-diapause quiescent larvae) were selected for RNAi, as larvae express high levels of *SmHsp22.2* and *SmHsp26.7* at this stage (see Section 3). Based on preliminary laboratory combined analysis of post-RNAi gene-silencing efficiency and survival rates, the optimal RNAi parameters were determined to be 30 nL of 10 μg/μL double-stranded RNA (dsRNA). Thus, in this study, each cocooned larva was injected with 300 ng (30 nL) of dsRNA at the junction between the 5th and 6th abdominal segments using a Nanoject II Auto-Nanoliter Injector (Drummond Scientific Company, Broomall, PA, USA), with corresponding amounts of dsGFP and DEPC water serving as the control groups. Following injection, the larvae were placed in Petri dishes containing moistened filter paper

and incubated for 12, 24, or 48 h in an incubator at 24 ± 1 °C with 70% ± 5% relative humidity (RH) and a 16/8 h light/dark cycle. Subsequently, 20 treated individuals were selected to form one group, resulting in a total of 3 groups, to determine the gene-silencing efficiency via qPCR.

2.8. Cold Tolerance of Larvae After RNAi

Twenty-four hours post-RNA interference, the larvae were exposed to a temperature of −10 °C for 2 h in a low-temperature thermostat bath, as both genes exhibit peak expression levels at this temperature (see Results). The larvae were then returned to the incubator (24 ± 1 °C, 70 ± 5% relative humidity, 16:8 light:dark), and mortality was calculated every two days over a six-day period by assessing the status of the larvae. Larvae are classified as alive only if they simultaneously meet two criteria: (1) the ability to crawl after breaking the cocoon, and (2) clear responsiveness to gentle brush stimulation when motility appears ambiguous. Each group contained 40 individuals, and three separate replicates were conducted for each treatment.

2.9. Data Analysis

Data for different diapause stages, heat- or cold-shock treatments, and RNA interference were analyzed using one-way analysis of variance (ANOVA). Tukey's multiple range test was applied for pairwise comparisons to assess the significance of differences between treatments with $p < 0.05$ considered statistically significant. All statistical analyses were performed using SPSS version 20.0, and the results are presented as means ± standard error (SE). For mortality scoring, all data were analyzed by Kaplan–Meier survival analysis, and differences between survival curves were assessed with the log-rank test, considering $p < 0.05$ as statistically significant. Statistical computations were performed using GraphPad Prism 9.5.0 (GraphPad Software, San Diego, CA, USA).

3. Results

3.1. Characterization of SmHsp22.2 and SmHsp26.7 cDNAs

The whole open reading frames (ORFs) of the two sHsp genes, namely *SmHsp22.2* (accession No. PQ660685) and *SmHsp26.7* (accession No. PQ660686), were obtained from transcriptome data of *S. mosellana* larvae and confirmed using RT-PCR. The ORFs of *SmHsp22.2* and *SmHsp26.7* contained 582 and 711 nucleotide pairs, respectively, corresponding to 193 and 236 amino acid residues (Figure S1). Their calculated molecular weights were 22.2 kDa and 26.7 kDa, with isoelectric points (pI) of 6.98 and 8.90, respectively. A comparison of the cDNA and gDNA sequences indicated that both genes lacked introns.

Analysis of domains identified the presence of the canonical α-crystallin domain in both SmHsps (a.a. 75-155 in SmHsp22.2 and a.a. 121-199 in SmHsp26.7). In addition, a characteristic motif I/VXI/V, situated near the C-terminus, along with an arginine residue necessary for the structural stability and chaperone function of sHsps [27,45] (Figure 1) was also conserved.

BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi, accessed on 15 May 2023) analysis indicated that SmHsp22.2 showed the highest sequence identity of 53% with Hsp23 and Hsp18.4 from *Bactrocera dorsalis*, as well as 40-48% identity with *Chironomus riparius* Hsp27, and Hsp23 proteins from *Delia antiqua*, *Polypedilum vanderplanki*, and *Diamesa zernyi*. SmHsp26.7 displayed 57% identity with *B. dorsalis* Hsp23, and 43-55% identity with *B. dorsalis* Hsp18.4, *Chironomus riparius* Hsp27, and Hsp23 proteins from *D. antiqua*, *P. vanderplanki*, and *D. zernyi*. The two SmHsps were 50% identical (Figure 1). The phylogenetic analysis demonstrated the clustering of sHsps from the same species, with the two SmHsps

exhibiting their closest evolutionary ties to their counterparts in the Nematocera suborder of the Diptera (*C. riparius*), aligning with the canonical taxonomic relationships (Figure 2).




Figure 1. Multiple sequence alignment of SmHsp22.2 and SmHsp26.7 (red triangles) as well as sHsps from other insects. Identical and similar amino acids are distinguished by black and grey shading, respectively. The α-crystallin domain is denoted by a single underline, and the V/P/I motif is indicated by a double underline. Conserved arginine is denoted with a downward-pointing triangle. Insect species and GenBank accession numbers of sHsps shown are as follows: *Sitodiplosis mosellana* (SmHsp22.2, XMH29534; SmHsp26.7, XMH29535); *Delia antiqua* (DaHsp23, ADX36150.1); *Bactrocera dorsalis* (BdHsp18.4, ARQ14797.1; BdHsp23, XP_011198115.2); *Diamesa zernyi* (DzHsp23, UJQ69871.1); *Polypedilum vanderplanki* (PvHsp23, ADM13385.1); *Chironomus riparius* (CrHsp27, AGJ98435.1).




Figure 2. Phylogenetic neighbor-joining tree of SmHsp22.2 and SmHsp26.7 (red triangles) as well as sHsps from other insects.

The secondary structure predictions identified six β-strands and two α-helices in the α-crystallin domains of both SmHsp22.2 and SmHsp26.7 (Figure 1). The tertiary structures of the proteins were modeled using the zebrafish (*Danio rerio*) homolog (PDB ID: 3n3e.1A) and the human (*Homo sapiens*) homolog (PDB ID: 2Wj7.1.A) as templates, due to their high sequence identity with SmHsp22.2 (45.7%) and SmHsp26.7 (56.4%), respectively. Both SmHsps contained a conserved α-crystallin domain, while SmHsp22.2 exists as a dimer and SmHsp26.7 is found as a monomer. Each monomer contained six β-strands, collectively forming a compact β-sandwich composed of two antiparallel β-sheets (Figure 3).




Figure 3. Tertiary structure prediction of SmHsp22.2 (**A**) and SmHsp26.7 (**B**). Predicted amino acid chains, specifically the α-helices and β-strands, are illustrated in a rainbow spectrum using PyMOL.

3.2. Expression of SmHsp22.2 and Hsp26.7 During Diapause

To explore the relationship between *SmHsp26.7/Hsp22.2* expression and diapause, we analyzed the relative transcript levels of both genes in 3rd instar larvae of *S. mosellana* across four distinct physiological stages associated with diapause, namely, pre-diapause, diapause, post-diapause quiescence, and post-diapause development (representing larvae collected during May, June-November, December-February, and March to early April of the following year, respectively) (Figure 4). Expression levels of *SmHsp22.2* increased significantly after the initiation of diapause (June), persisting at high levels throughout diapause and the early-to-mid phase of post-diapause quiescence, and undergoing a sharp decline during the late post-diapause quiescence phase, ultimately reaching their lowest levels at the post-diapause developmental phase (Figure 4A).




Figure 4. Relative transcription levels of *SmHsp22.2* (**A**) and *SmHsp26.7* (**B**) in *Sitodiplosis mosellana* larvae in pre-diapause, diapause, post-diapause quiescence, and post-diapause development (representing larvae collected during May, June-November, December-February, and March to early April of the following year, respectively). The transcript abundance of each tested stage was normalized to that of the pre-diapausing larvae (arbitrarily set as 1). The results are denoted as means ± SE. Bars labeled with different letters denote significant differences (Tukey's multiple range test, $p < 0.05$).

Likewise, *SmHsp26.7* expression was low during the pre-diapause and post-diapause developmental stages, while peaking during the early-to-mid post-diapause quiescence stage (December and January), the coldest months of the year. In contrast to *SmHsp22.2*, *SmHsp26.7* expression increased markedly from July to August, surpassing the levels observed in other diapause stages (Figure 4B).

3.3. Expression of SmHsp22.2 and Hsp26.7 in Response to Heat Shock During Diapause

The expression profiles of two *SmHsps* were markedly similar in over-summering diapause larvae after exposure to different high-temperature treatments. Compared with the control group, the expression levels of both *SmHsp22.2* and *SmHsp26.7* increased significantly in the temperature range of 35 to 40 °C, peaking at 35 °C, with increases of approximately 2.87-fold and 3.26-fold, respectively. However, no significant increase was apparent at 45–50 °C in both *S. mosellana Hsp22.2/26.7* genes (Figure 5A,B).




Figure 5. Relative transcription levels of *SmHsp22.2* and *SmHsp26.7* in response to different high temperatures (35–50 °C) for 1 h (**A**,**B**) or 35 °C for different durations (0–120 min) (**C**,**D**) in over-summering diapausing larvae of *Sitodiplosis mosellana*. The transcript abundance of each treatment was normalized to that of untreated control (CK, 0 min) (arbitrarily set as 1). The results are denoted as means ± SE. Bars labeled with different letters denote significant differences (Tukey's multiple range test, $p < 0.05$).

The duration of treatment also influenced the mRNA expression levels of SmHsp22.2 and SmHsp26.7. At 35 °C, the expression of both SmHsp22.2 and SmHsp26.7 rose significantly at 15 min, culminating in peaks at 30 min and 60 min, respectively (Figure 5C,D).

3.4. Expression of SmHsp22.2 and Hsp26.7 in Response to Cold Shock During Diapause

The expression of *SmHsp22.2* and *SmHsp26.7* in over-wintering diapause larvae was also influenced by exposure to cold temperatures. At −5 °C, the expression of both *SmHsp22.2* and *SmHsp26.7* was significantly upregulated, peaking at −10 °C with approximate increases of 2.62-fold and 7.71-fold, respectively. Notably, the expression of *SmHsp22.2* was observed to decrease markedly, while that of *SmHsp26.7* showed no discernible change at 0 °C when compared to the control group (Figure 6A,B).




Figure 6. Relative transcription levels of *SmHsp22.2* and *SmHsp26.7* in response to different low temperatures (0–−15 °C) for 1 h (**A**,**B**) or −10 °C for different durations (0–120 min) (**C**,**D**) in overwintering diapausing larvae of *Sitodiplosis mosellana*. The transcript abundance of each treatment was normalized to that of untreated control (CK, 0 min) (arbitrarily set as 1). The results are denoted as means ± SE. Bars labeled with different letters denote significant differences (Tukey's multiple range test, $p < 0.05$).

At −10 °C, *SmHsp22.2* expression levels rose progressively with time and were significantly upregulated at 30 min, reaching their maximum at 60 min. The expression of *SmHsp26.7*, however, showed a maximum response between 30 and 60 min of cold treatment, with expression reaching its highest point at 30 min (Figure 6C,D).

3.5. Effects of SmHsp22.2 and SmHsp26.7 Knockdown on Cold Tolerance in S. mosellana

RNAi-mediated knockdown of *SmHsp22.2*/*SmHsp26.7* led to a marked decrease in transcription levels of both genes at three time points (Figure 7). After the treatment with *dsHsp22.2*, the expression of *SmHsp22.2* dropped by 45, 52, and 57%, at 12, 24, and 48 h, respectively, compared to the DEPC-water control, as well as 47, 51, and 56% at 12, 24, and 48 h, relative to the dsGFP control (Figure 7A). Similarly, *SmHsp26.7* expression was reduced by 49, 61, and 70% compared with the DEPC-water control, and 51, 62, and 68% relative to the dsGFP control, at the same respective time points (Figure 7B). Therefore, we conclude that the knockdown of both genes was successful.

To further investigate the roles of *SmHsp22.2* and *SmHsp26.7* in *S. mosellana* cold tolerance, larvae in which the genes had been silenced 24 h previously were placed in a low-temperature thermostat bath at −10 °C for 2 h, followed by mortality assessments every two days over a six-day period. The results showed that the survival rate of ds*Hsp22.2* or ds*Hsp26.7*-injected larvae was significantly lower than that of DEPC-water or ds*GFP* groups in 2, 4, and 6 days ($p < 0.0001$) (Figure 8), indicating the involvement of both genes in the cold resistance of *S. mosellana*.




Figure 7. Relative expression levels of *SmHsp22.2* (**A**) and *SmHsp26.7* (**B**) at different time points after RNAi. The transcript abundance of each time point was normalized to that of the DEPC water (arbitrarily set as 1). The results are denoted as means ± SE. Bars labeled with different letters denote significant differences (Tukey's multiple range test, $p < 0.05$).




Figure 8. Survival curves of *Sitodiplosis mosellana* larvae injected with dsRNA. The distinct survival curves correspond to various experimental groups. DEPC-treated water and ds*GFP* functioned as the blank control and negative control, respectively. Kaplan–Meier estimators were utilized for survival curve modeling, with pairwise comparisons performed using Log-rank tests. ****, $p < 0.0001$; ns, $p > 0.05$. Each group—DEPC-water, ds*GFP*, ds*Hsp22.2*, and ds*Hsp26.7*—comprised 120 specimens, with cumulative mortality rates of 23% (28/120), 25% (30/120), 65% (78/120), and 72% (86/120) over six days, respectively.

4. Discussion

In addition to the two documented sHsp proteins, Hsp17.4 and Hsp20.3, in *S. mosellana* [40], this study identified two novel sHsp members from this species, namely, *SmHsp22.2* and *SmHsp26.7*. Comparative analysis of the genomic and cDNA sequences of

SmHsp22.2 and *SmHsp26.7* revealed the absence of introns in both genes. Typically, sHsps have been broadly categorized into two types, namely, orthologous types with introns and species-specific types without introns [46]. Despite the prevalence of the species-specific sHsps in insects, intron-containing genes have been observed in several species, including *Tribolium castaneum* [47], *Choristoneura fumiferana* [48], and *Mythimna separata* [49]. The evolutionary and functional diversification of sHsps in insects may thus have resulted in the emergence of genes that are either devoid of introns or possess shorter introns, thereby enhancing their capacity to respond to unfavorable environmental conditions by facilitating rapid expression [50,51].

In common with known insect sHsp proteins, SmHsp22.2 and SmHsp26.7 were found to contain all the canonical signature motifs, specifically, an α-crystalline domain comprised of several β-strands that is responsible for the molecular chaperone functionality, and the "I/VXI/V" motif that is indispensable for sustaining oligomer stability and structural assembly [23,52]. In contrast to the conserved α-crystallin domain located near the C-terminus, the N-terminal regions of sHsps exhibit substantial diversity, contributing to their ability to interact with a variety of protein partners [20]. The phylogenetic analysis demonstrated marked clustering of sHsps within the same species, which is consistent with previous findings on fungi, insects, plants, and vertebrates [53], and implies that sHsps have likely evolved through gene duplications occurring after species divergence.

Given the distinctive responsiveness and functionality of sHsp members in reacting to stress during insect diapause [3,27,28], we examined the expression patterns of two novel sHsp-encoding genes in a natural population of *S. mosellana*, rather than the commonly utilized laboratory-reared populations employed in most molecular studies on diapause. Analysis of the field-collected *S. mosellana* showed significant upregulation of both *SmHsp22.2* and *SmHsp26.7* during diapause, particularly in summer and winter (Figure 4), suggesting their potential roles in protection and cold tolerance in diapause larvae. Similar results have also been reported in other insect species, where *Hsp19.5* and *Hsp20.0* from *P. melete*, as well as *Hsp23* from *Delia antiqua*, exhibited increased expression in over-wintering and over-summering diapause pupae [27,54]. These results suggest that modulations in sHsp expression form an integral part of diapause in insects.

In natural settings, insects frequently encounter complex and severe environmental conditions, and an important adaptation strategy in insects under these conditions is modulation of the expression levels of sHsps, which exhibit varied transcriptional responses when subjected to temperature fluctuations [14,55]. The present results indicated that two *S. mosellana sHsps* were significantly upregulated in both heat-stressed over-summering larvae and cold-stressed over-wintering larvae, although this induction was transient (Figures 5 and 6). It is plausible that *SmHsps* offer temporary protection for *S. mosellana* during diapause when subjected to extreme temperatures at the soil surface due to agricultural practices such as tillage [56,57]. Notably, the significant downregulation of *Hsp22.2* and *Hsp26.7* expression beyond a threshold temperature (Figures 5 and 6) suggests that thermal stress causes cellular damage exceeding the protective capacity of sHsps [58], impairing their synthesis and underscoring the limited ability of insects to mitigate cellular injury through sHsp upregulation.

Overall, the transcriptional upregulation of sHsp genes in insects is closely associated with cold tolerance [14,59]. For instance, in over-wintering diapause pupae of *Sarcophaga crassipalpis*, *Hsp23* expression was found to be significantly elevated [60], and its knockdown via RNAi markedly influenced pupal survival at low temperatures [30]. Similarly, suppression of *Hsp19.0* expression in larval *Chilo suppressalis* was observed to markedly reduce their viability when exposed to subzero temperatures of −11 °C [61]. Furthermore, downregulation of *Hsp14.9*, *Hsp19.9*, *Hsp20.3*, and *Hsp24.0* in *Haemaphysalis longicornis* led

to a significant decrease in their survival rates under low-temperature stress at −14 °C [13]. In this study, a temperature of −10 °C was selected to analyze the functions of *S. mosellana Hsp22.2* and *Hsp26.7*, as the peak expression levels of both genes occur at this temperature (Figure 6). After injection of *dsHsp22.2/26.7*, the larvae exhibited significantly lower survival rates compared to the DEPC-water and dsGFP control groups (Figure 8), suggesting that both sHsp genes are essential for protecting *S. mosellana* from the adverse effects of low temperature. These findings highlight potential molecular targets for RNA interference (RNAi)-based pest management strategies. Nonetheless, practical application necessitates further refinement of delivery vectors and comprehensive ecological risk assessments prior to field deployment. Thus, there is still a long way to go before *Sitodiplosis mosellana* can be effectively controlled.

5. Conclusions

The diapause-induced upregulation of *SmHsp22.2* and *SmHsp26.7*, their induction upon sublethal heat (35–40 °C) and moderate cold (−10 °C), and gene-specific RNAi-induced cold sensitivity are strong results that enhance knowledge about sHsp function in insect stress physiology. These findings have potential practical relevance to predicting the dynamics of pest populations under climate variability and indicate that *SmHsps* could be targeted with RNAi-based biocontrol in wheat systems; however, field validation is necessary.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/insects16070649/s1, Figure S1. Nucleic acid and deduced amino acid sequences of *SmHsp22.2* and *SmHsp26.7* in *Sitodiplosis mosellana*. Initiation codons (ATG) and termination codons (TAA) were indicated by boxes. The α-crystallin domain was highlighted by shading. I/VXI/V motif was indicated with a single underline.

Author Contributions: Conceptualization, W.C. and K.Z.-S.; methodology, Q.H.; investigation, Q.H.; formal analysis, Q.H., Q.M., X.L., and W.C.; writing—original draft preparation, Q.H., and W.C.; writing—review and editing, W.C. and K.Z.-S. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Natural Science Foundation of Shandong Province (ZR2023QC051), Shandong Academy of Agricultural Sciences Agricultural Science and Technology Innovation Engineering (CXGC2024F19, CXGC2025F18), National Natural Science Foundation of China (31371933), Open Project of Shaanxi Laboratory for Agriculture in Arid Areas (2024ZY-JCYJ-02-39).

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Materials. Further inquiries can be directed to the corresponding author.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Zonato, V.; Collins, L.; Pegoraro, M.; Tauber, E.; Kyriacou, C.P. Is diapause an ancient adaptation in *Drosophila*? *J. Insect Physiol.* **2017**, *98*, 267–274. [CrossRef]
2. Schebeck, M.; Lehmann, P.; Laparie, M.; Bentz, B.J.; Ragland, G.J.; Battisti, A.; Hahn, D.A. Seasonality of forest insects: Why diapause matters. *Trends Ecol. Evol.* **2024**, *39*, 757–770. [CrossRef] [PubMed]
3. Yang, M.Y.; Wang, Z.C.; Wang, R.; Zhang, X.M.; Li, M.Y.; Xin, J.J.; Qin, Y.S.; Zhang, C.; Meng, F.L. Transcriptomic and proteomic analyses of the mechanisms of overwintering diapause in soybean pod borer (*Leguminivora glycinivorella*). *Pest Manag. Sci.* **2020**, *76*, 4248–4257. [CrossRef]
4. Guo, S.; Tian, Z.; Wu, Q.W.; King-Jones, K.; Liu, W.; Zhu, F.; Wang, X.P. Steroid hormone ecdysone deficiency stimulates preparation for photoperiodic reproductive diapause. *PLoS Genet.* **2021**, *17*, e1009352. [CrossRef] [PubMed]
5. Liao, J.; Cai, D.X.; Geng, S.L.; Lyu, Z.P.; Wu, Y.L.; Guo, J.J.; Li, H.Y. Transcriptome-based analysis reveals a crucial role of the 20E/HR3 pathway in the diapause of *Pieris rapae*. *Pestic. Biochem. Physiol.* **2024**, *199*, 105787. [CrossRef] [PubMed]

6. Srygley, R.B. Effects of parental diet on mormon cricket egg diapause, embryonic development rate, and periodic outbreaks. *J. Insect Physiol.* **2024**, *157*, 104681. [CrossRef]
7. Hahn, D.A.; Denlinger, D.L. Energetics of insect diapause. *Annu. Rev. Entomol.* **2011**, *56*, 103–121. [CrossRef]
8. Hand, S.C.; Denlinger, D.L.; Podrabsky, J.E.; Roy, R. Mechanisms of animal diapause: Recent developments from nematodes, crustaceans, insects, and fish. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2016**, *310*, R1193–R1211. [CrossRef]
9. Denlinger, D.L. Why study diapause? *Entomol. Res.* **2008**, *38*, 1–9. [CrossRef]
10. Altincicek, B.; Knorr, E.; Vilcinskas, A. Beetle immunity: Identification of immune-inducible genes from the model insect *Tribolium castaneum*. *Dev. Comp. Immunol.* **2008**, *32*, 585–595. [CrossRef]
11. Lubkowska, A.; Pluta, W.; Strońska, A.; Lalko, A. Role of heat shock proteins (HSP70 and HSP90) in viral infection. *Int. J. Mol. Sci.* **2021**, *22*, 9366. [CrossRef]
12. Shen, Z.J.; Liu, Y.J.; Cheng, J.; Li, Z.; Michaud, J.P.; Liu, X.X. High temperature exposure reduces the susceptibility of *Helicoverpa armigera* to its nucleopolyhedrovirus (HearNPV) by enhancing expression of heat shock proteins. *Pest Manag. Sci.* **2022**, *78*, 2378–2389. [CrossRef] [PubMed]
13. Pei, T.W.; Zhang, M.; Nwanade, C.F.; Meng, H.; Bai, R.W.; Wang, Z.H.; Wang, R.T.; Zhang, T.A.; Liu, J.Z.; Yu, Z.H. Sequential expression of small heat shock proteins contributing to the cold response of *Haemaphysalis longicornis* (Acari: Ixodidae). *Pest Manag. Sci.* **2024**, *80*, 2061–2071. [CrossRef] [PubMed]
14. King, A.M.; MacRae, T.H. Insect heat shock proteins during stress and diapause. *Annu. Rev. Entomol.* **2015**, *60*, 59–75. [CrossRef]
15. Yu, C.; Leung, S.K.P.; Zhang, W.X.; Lai, L.T.F.; Chan, Y.K.; Wong, M.C.; Benlekbir, S.; Cui, Y.; Jiang, L.W.; Lau, W.C.Y. Structural basis of substrate recognition and thermal protection by a small heat shock protein. *Nat. Commun.* **2021**, *12*, 3007. [CrossRef] [PubMed]
16. Tedesco, B.; Ferrari, V.; Cozzi, M.; Chierichetti, M.; Casarotto, E.; Pramaggiore, P.; Mina, F.; Galbiati, M.; Rusmini, P.; Crippa, V.; et al. The role of small heat shock proteins in protein misfolding associated motoneuron diseases. *Int. J. Mol. Sci.* **2022**, *23*, 11759. [CrossRef]
17. Huang, L.H.; Wang, H.S.; Kang, L. Different evolutionary lineages of large and small heat shock proteins in eukaryotes. *Cell Res.* **2008**, *18*, 1074–1076. [CrossRef]
18. Schell, R.; Mullis, M.; Ehrenreich, I.M. Modifiers of the genotype-phenotype map: Hsp90 and beyond. *PLoS Biol.* **2016**, *14*, e2001015. [CrossRef]
19. Zarouchlioti, C.; Parfitt, D.A.; Li, W.W.; Gittings, L.M.; Cheetham, M.E. DNAJ proteins in neurodegeneration: Essential and protective factors. *Philos. Trans. R. Soc. Lond B Biol. Sci.* **2018**, *373*, 20160534. [CrossRef]
20. Basha, E.; Friedrich, K.L.; Vierling, E. The N-terminal arm of small heat shock proteins is important for both chaperone activity and substrate specificity. *J. Biol. Chem.* **2006**, *281*, 39943–39952. [CrossRef]
21. Kriehuber, T.; Rattei, T.; Weinmaier, T.; Bepperling, A.; Haslbeck, M.; Buchner, J. Independent evolution of the core domain and its flanking sequences in small heat shock proteins. *FASEB J.* **2010**, *24*, 3633–3642. [CrossRef] [PubMed]
22. Hayes, D.; Napoli, V.; Mazurkie, A.; Stafford, W.F.; Graceffa, P. Phosphorylation dependence of hsp27 multimeric size and molecular chaperone function. *J. Biol. Chem.* **2009**, *284*, 18801–18807. [CrossRef]
23. Carra, S.; Alberti, S.; Arrigo, P.A.; Benesch, J.L.; Benjamin, I.J.; Boelens, W.; Bartelt-Kirbach, B.; Brundel, B.J.J.M.; Buchner, J.; Bukau, B.; et al. The growing world of small heat shock proteins: From structure to functions. *Cell Stress Chaperones* **2017**, *22*, 601–611. [CrossRef]
24. Dubrez, L.; Causse, S.; Borges, B.N.; Dumétier, B.; Garrido, C. Heat-shock proteins: Chaperoning DNA repair. *Oncogene* **2020**, *39*, 516–529. [CrossRef]
25. Shan, R.T.; Liu, N.; Yan, Y.Y.; Liu, B. Apoptosis, autophagy and atherosclerosis: Relationships and the role of Hsp27. *Pharmacol. Res.* **2021**, *166*, 105169. [CrossRef] [PubMed]
26. Rocchetti, M.T.; Bellanger, T.; Trecca, M.I.; Weidmann, S.; Scrima, R.; Spano, G.; Russo, P.; Capozzi, V.; Fiocco, D. Molecular chaperone function of three small heat-shock proteins from a model probiotic species. *Cell Stress Chaperones* **2023**, *28*, 79–89. [CrossRef]
27. Miano, F.N.; Jiang, T.; Zhang, J.; Zhang, W.N.; Peng, Y.; Xiao, H.J. Identification and up-regulation of three small heat shock proteins in summer and winter diapause in response to temperature stress in *Pieris melete*. *Int. J. Biol. Macromol.* **2022**, *209*, 1144–1154. [CrossRef] [PubMed]
28. Dageri, A. Molecular characterization and expression analysis of six small heat shock protein genes in *Trogoderma granarium* during cold and starvation-induced larval diapause. *J. Stored Prod. Res.* **2024**, *108*, 102368. [CrossRef]
29. Tachibana, S.I.; Numata, H.; Goto, S.G. Gene expression of heat-shock proteins (*Hsp23*, *Hsp70* and *Hsp90*) during and after larval diapause in the blow fly *Lucilia sericata*. *J. Insect Physiol.* **2005**, *51*, 641–647. [CrossRef]
30. Rinehart, J.P.; Li, A.; Yocum, G.D.; Robich, R.M.; Hayward, S.A.; Denlinger, D.L. Up-regulation of heat shock proteins is essential for cold survival during insect diapause. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 11130–11137. [CrossRef]

31. Fremdt, H.; Amendt, J.; Zehner, R. Diapause-specific gene expression in *Calliphora vicina* (Diptera: Calliphoridae)—A useful diagnostic tool for forensic entomology. *Int. J. Legal Med.* **2014**, *128*, 1001–1011. [CrossRef] [PubMed]
32. Aruda, A.M.; Baumgartner, M.F.; Reitzel, A.M.; Tarrant, A.M. Heat shock protein expression during stress and diapause in the marine copepod *Calanus finmarchicus*. *J. Insect Physiol.* **2011**, *57*, 665–675. [CrossRef] [PubMed]
33. Yang, W.J.; Xu, K.K.; Cao, Y.; Meng, Y.L.; Liu, Y.; Li, C. Identification and expression analysis of four small heat shock protein genes in cigarette beetle, *Lasioderma serricorne* (Fabricius). *Insects* **2019**, *10*, 139. [CrossRef] [PubMed]
34. Uzelac, I.; Avramov, M.; Knežić, T.; Tatić, V.; Gošić-Dondo, S.; Popović, Ž.D. Prolonged heat stress during winter diapause alters the expression of stress-response genes in *Ostrinia nubilalis* (Hbn.). *Int. J. Mol. Sci.* **2024**, *25*, 3100. [CrossRef]
35. Gkouvitsas, T.; Kontogiannatos, D.; Kourti, A. Differential expression of two small Hsps during diapause in the corn stalk borer *Sesamia nonagrioides* (Lef.). *J. Insect Physiol.* **2008**, *54*, 1503–1510. [CrossRef]
36. Smith, M.A.H.; Wise, I.L.; Fox, S.L.; Vera, C.L.; DePauw, R.M.; Lukow, O.M. Seed damage and sources of yield loss by *Sitodiplosis mosellana* (Diptera: Cecidomyiidae) in resistant wheat varietal blends relative to susceptible wheat cultivars in western Canada. *Can. Entomol.* **2014**, *146*, 335–346. [CrossRef]
37. Shrestha, G.; Reddy, G.V.P. Field efficacy of insect pathogen, botanical, and jasmonic acid for the management of wheat midge *Sitodiplosis mosellana* and the impact on adult parasitoid *Macroglenes penetrans* populations in spring wheat. *Insect Sci.* **2019**, *26*, 523–535. [CrossRef]
38. Zhang, L.J.; Geng, M.M.; Zhang, Z.; Zhang, Y.; Yan, G.J.; Wen, S.M.; Liu, G.R.; Wang, R.H. Molecular mapping of major QTL conferring resistance to orange wheat blossom midge (*Sitodiplosis mosellana*) in Chinese wheat varieties with selective populations. *Theor. Appl. Genet.* **2020**, *133*, 491–502. [CrossRef]
39. Wang, Y.; Long, Z.R.; Feng, A.R.; Cheng, W.N. Effects of initial population number, wheat varieties and precipitation on infestation of *Sitodiplosis mosellana* (Diptera: Cecidomyiidae). *Acta Agri. Boreali-Occident. Sin.* **2015**, *24*, 165–171.
40. Zhao, J.J.; Huang, Q.T.; Zhang, G.J.; Zhu-Salzman, K.Y.; Cheng, W.N. Characterization of two small heat shock protein genes (*Hsp17.4* and *Hs20.3*) from *Sitodiplosis mosellana*, and their expression regulation during diapause. *Insects* **2021**, *12*, 119. [CrossRef]
41. Cheng, W.N.; Li, D.; Wang, Y.; Liu, Y.; Zhu-Salzman, K.Y. Cloning of heat shock protein genes (*hsp70*, *hsc70* and *hsp90*) and their expression in response to larval diapause and thermal stress in the wheat blossom midge, *Sitodiplosis mosellana*. *J. Insect Physiol.* **2016**, *95*, 66–77. [CrossRef]
42. Cheng, W.N.; Long, Z.R.; Zhang, Y.D.; Liang, T.T.; Zhu-Salzman, K.Y. Effects of temperature, soil moisture and photoperiod on diapause termination and post-diapause development of the wheat blossom midge, *Sitodiplosis mosellana* (Géhin) (Diptera: Cecidomyiidae). *J. Insect Physiol.* **2017**, *103*, 78–85. [CrossRef]
43. Doane, J.F.; Olfert, O. Seasonal development of wheat midge, *Sitodiplosis mosellana* (Géhin) (Diptera: Cecidomyiidae), in Saskatchewan, Canada. *Crop Prot.* **2008**, *27*, 951–958. [CrossRef]
44. Chen, H.S.; Lei, C.L.; Wu, Y.Q.; Dou, Z.B.; Miao, J.; Duan, Y.; Jiang, Y.L. Detection on supercooling points of *Sitodiplosis mosellana* (Gehin) round cocoons in different geographical districts. *J. Huazhong Agric. Univ.* **2012**, *31*, 212–215.
45. Kumar, L.V.; Ramakrishna, T.; Rao, C.M. Structural and functional consequences of the mutation of a conserved arginine residue in alphaA and alphaB crystallins. *J. Biol. Chem.* **1999**, *274*, 24137–24141. [CrossRef] [PubMed]
46. Li, Z.W.; Li, X.; Yu, Q.Y.; Xiang, Z.H.; Kishino, H.; Zhang, Z. The small heat shock protein (sHSP) genes in the silkworm, *Bombyx mori*, and comparative analysis with other insect sHSP genes. *BMC Evol. Biol.* **2009**, *9*, 215. [CrossRef] [PubMed]
47. Xie, J.; Xiong, W.; Hu, X.; Gu, S.; Li, B. Characterization and functional analysis of *hsp21.8b*: An orthologous small heat shock protein gene in *Tribolium castaneum*. *J. Appl. Entomol.* **2018**, *142*, 12519. [CrossRef]
48. Quan, G.X.; Duan, J.; Fick, W.; Kyei-Poku, G.; Candau, J.N. Expression profiles of 14 small heat shock protein (sHSP) transcripts during larval diapause and under thermal stress in the spruce budworm, *Choristoneura fumiferana* (L.). *Cell Stress Chaperones* **2018**, *23*, 1247–1256. [CrossRef]
49. Li, H.B.; Dai, C.G.; Hu, Y. Characterization and expression analysis of genes encoding three small heat shock proteins in the oriental armyworm, *Mythimna separata* (Walker). *PLoS ONE* **2020**, *15*, e0235912. [CrossRef]
50. Comeron, J.M. Selective and mutational patterns associated with gene expression in humans: Influences on synonymous composition and intron presence. *Genetics* **2004**, *167*, 1293–1304. [CrossRef]
51. Liu, W.W.; Yang, P.; Chen, X.M.; Xu, D.L.; Hu, Y.H. Cloning and expression analysis of four heat shock protein genes in *Ericerus pela* (Homoptera: Coccidae). *J. Insect Sci.* **2014**, *14*, 142. [CrossRef] [PubMed]
52. Bagnéris, C.; Bateman, O.A.; Naylor, C.E.; Cronin, N.; Boelens, W.C.; Keep, N.H.; Slingsby, C. Crystal structures of alpha-crystallin domain dimers of alphaB-crystallin and Hsp20. *J. Mol. Biol.* **2009**, *392*, 1242–1252. [CrossRef] [PubMed]
53. de Jong, W.W.; Leunissen, J.A.; Voorter, C.E. Evolution of the alpha-crystallin/small heat-shock protein family. *Mol. Biol. Evol.* **1993**, *10*, 103–126.
54. Si, F.L.; He, Z.B.; Chen, B. Cloning and expression profiling of heat shock protein DaHSP23 gene in the winter and summer diapause pupae of the onion maggot, *Delia antiqua* (Diptera: Anthomyiidae). *Acta Entomol. Sin.* **2016**, *59*, 402–410.

55. Bale, J.S.; Masters, G.J.; Hodkinson, I.D.; Awmack, C.; Bezemer, T.M.; Brown, V.; Butterfield, J.; Buse, A.; Coulson, J.C.; Farrar, J.; et al. Herbivory in global climate change research: Direct effects of rising temperature on insect herbivores. *Global Change Biol.* **2002**, *8*, 1–16. [CrossRef]
56. Barnes, H.F. Studies of fluctuations in insect populations. VIII. The wheat blossom midge on broadbalk, 1932–1940, with a discussion of the results obtained 1927-40. *J. Anim. Ecol.* **1941**, *10*, 94–120. [CrossRef]
57. Yuan, F. *The Wheat Blossom Midges Sitodiplosis mosellana (Gehin) and Contarinia tritici (Kirby): Their Plague Principle and Control*; Science Press: Beijing, China, 2004.
58. Colinet, H.; Lee, S.F.; Hoffmann, A. Knocking down expression of *Hsp22* and *Hsp23* by RNA interference affects recovery from chill coma in *Drosophila melanogaster*. *J. Exp. Biol.* **2010**, *213*, 4146–4150. [CrossRef]
59. Pan, D.D.; Lu, M.X.; Li, Q.Y.; Du, Y.Z. Characteristics and expression of genes encoding two small heat shock protein genes lacking introns from *Chilo suppressalis*. *Cell Stress Chaperones* **2018**, *23*, 55–64. [CrossRef]
60. Yocum, G.D.; Joplin, K.H.; Denlinger, D.L. Upregulation of a 23 kDa small heat shock protein transcript during pupal diapause in the flesh fly, *Sarcophaga, crassipalpis*. *Insect Biochem. Mol. Biol.* **1998**, *28*, 677–682. [CrossRef]
61. Dong, C.L.; Zhu, F.; Lu, M.X.; Du, Y.Z. Characterization and functional analysis of *Cshsp19.0* encoding a small heat shock protein in *Chilo suppressalis* (Walker). *Int. J. Biol. Macromol.* **2021**, *188*, 924–931. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Effects of Trehalase on the Gene Expression of the Reproductive Regulation Pathway Network and Triglyceride Metabolism in *Nilaparvata lugens* (Stål)

Bin Tang [1,*,†], Yuxin Ge [1,†], Yongkang Liu [1], Liwen Guan [1], Ye Han [1], Yang Zhu [1], Gao Hu [2,3] and Yan Wu [2,*]

[1] College of Life and Environmental Sciences, Hangzhou Normal University, Hangzhou 311121, China; geyuxin107@163.com (Y.G.); hznu_yongkang@163.com (Y.L.); guanliwen1010@163.com (L.G.); 19011285720@163.com (Y.H.); 2023112010008@stu.hznu.edu.cn (Y.Z.)
[2] Guizhou Key Laboratory of Agricultural Biosecurity, Guiyang University, Guiyang 550005, China; hugao@njau.edu.cn
[3] State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China
* Correspondence: tbzm611@hznu.edu.cn (B.T.); 15150535703@163.com (Y.W.)
† These authors contributed equally to this work.

Simple Summary: Trehalase (TRE) plays a critical role in insect survival and reproduction. The inhibition of TRE in the *Nilaparvata lugens* has been shown to impair reproductive capacity, yet the underlying molecular mechanisms have remained elusive. By inhibiting TRE activity (RNAi and validamycin), we discovered that TRE disruption fundamentally impacts the insect's reproductive regulatory network and triglyceride metabolism pathways. This study provides crucial molecular insights into how trehalose metabolism governs insect reproduction. Importantly, it solidly validates TRE as a highly promising target for developing novel, mechanism-based strategies to control pest insects.

Abstract

The high reproductive performance of *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae) is one of the key factors contributing to its serious damage. Studies have demonstrated that trehalose hydrolysis, catalyzed by trehalase (TRE), plays an important role in the reproductive regulation of *N. lugens*. However, it remains unclear how TRE regulates the reproduction of *N. lugens*. To address this question, the current study was designed to investigate the effects of TRE on the reproductive regulatory network of *N. lugens*. Specifically, RNA interference (RNAi) was conducted by injecting double-stranded RNA (dsRNA) targeting the TRE genes, and a trehalase inhibitor, validamycin, was used to suppress TRE activity. Subsequently, the relative expression levels of key genes involved in the reproductive regulatory pathway, as well as the triglyceride content in the fat body and ovary, were determined. The results indicated that both dsTREs and validamycin treatment decreased the relative expression of Vitellogenin (Vg) without affecting the expression of its receptor, *VgR*. However, their impact on the reproductive regulatory network differed: dsTREs injection did not affect the relative expression of *JHAMT* and *Met* in the juvenile hormone signaling pathway, but resulted in the upregulation of *USP* and *TOR*, and inhibition of *InR2* and *S6K*. In contrast, validamycin injection inhibited the expression of *Met*, *USP*, *EcR*, *InR1*, *TOR*, and *S6K*. Both dsTREs and validamycin inhibited the expression of *Fas*, but only validamycin decreased the ovarian triglyceride content. These findings suggest that the inhibition of *TRE* leads to alterations in reproductive regulatory signals, thereby reducing the reproductive capacity of *N. lugens*.

Keywords: trehalase; reproduction; RNA interference; validamycin; *Nilaparvata lugens*

1. Introduction

Nilaparvata lugens (Stål) (Hemiptera: Delphacidae) exhibits multiple harmful behavioral traits that negatively impact rice production, contributing to its status as the most significant biotic threat to rice yield [1–3]. Currently, the predominant method for managing insect pests is the foliar application of insecticides [4]. However, this approach is not effective in the long-term control of pest populations and, when misused, can lead to the development of insecticide resistance through metabolic changes or target site mutations [5–7]. Given the severe consequences of *N. lugens* infestations on agricultural productivity, there is an urgent need to develop novel, environmentally friendly, sustainable, and efficient pest management strategies. Among the proposed strategies, suppressing the reproductive capacity of *N. lugens* emerges as a critical approach.

The metabolism of trehalose, a major circulating sugar during the growth of insect oocytes, has a key physiological role in insect reproduction [8]. Therefore, it has the potential to serve as a target for insecticide action. Trehalases (TREs) are pivotal enzymes involved in insect development and reproduction, including soluble (TRE1) and membrane-bound (TRE2) types [9–11]. TRE1 has two coding genes, *TRE1-1* and *TRE1-2*, while TRE2 has only one coding gene [12–17]. Previous studies showed that both TRE1 and TRE2 regulate development, molting, and emergence in insect larvae through the chitin pathway [15]. For example, the single interference of *TRE1-1*, *TRE1-2*, and *TRE2* genes all led to an imbalance in trehalose supply, thus increasing larval mortality in *N. lugens* [15]. Validamycin is a competitive natural inhibitor of TRE [18]. As a structural analog of trehalose, validamycin exhibits potent inhibitory activity against trehalase in pest insects [19–24], further supporting its potential as a target for insecticide action. This is in agreement with our laboratory's previous results showing the significant inhibition of TRE activity following validamycin injection into *N. lugens* [16]. Studies have shown that the inhibition of TRE activity by validamycin can result in a significant decrease in ATP levels, leading to slow oocyte development [25]. This further confirms that functional TRE is essential for energy production in the ovary, and that its impairment affects reproduction.

Vitellogenin (Vg) is a key nutrient required for ovary development, and it is synthesized as a macromolecule mainly in the fat body and then transported to the mature oocyte via endocytosis, which is mediated by the vitellogenin receptor (*VgR*). This process, known as vitellogenesis, is a central event in insect reproduction [17,26]. Deficiencies in Vg synthesis and transport can directly lead to maternal sterility or the reduced hatchability of offspring [27]. It has been shown that vitellogenesis is mainly regulated by hormone signaling pathways [28]. Among them, the JH signaling pathway is involved in the regulation of vitellogenesis and oocyte maturation in some female insects and has an impact on other aspects of reproduction [29–36]. Moreover, 20-hydroxyecdysone (20E), another major regulatory hormone of female reproduction in some Hymenoptera, Lepidoptera, and all Diptera, is usually synthesized in the ovary [31,36,37]. The insulin-like peptide signaling pathway (IIS), also known as the insulin signaling pathway, plays a key role in various aspects of metabolism, development, and reproduction in insects [38,39]. Additionally, in vivo studies in *N. lugens* have demonstrated that the transcription factor Foxo can directly act on the exon to regulate *Vg* expression [40], indicating that the IIS can regulate insect reproduction independently of JH and 20E [41]. Furthermore, the IIS may crosstalk with the JH or 20E signaling pathways to indirectly affect insect reproduction [42]. The TOR signaling pathway, a key nutrient sensor, regulates physiological activities according to the organism's nutritional status. After entering the cytoplasm through transmembrane

receptors, amino acids bind to GTPases or Rag proteins, which results in the activation of the TOR pathway [43]. S6 kinase (S6K), phosphorylated and activated downstream of TOR, participates in the transcriptional and translational regulation of Vg [43,44]. Thus, the TOR signaling pathway directly regulates insect reproduction in response to nutrient availability. Furthermore, the TOR stimulates the secretion of insulin-like peptides in insects, thereby activating the IIS signaling pathway and indirectly influencing reproductive processes [43]. Therefore, four signaling pathways—JH, 20E, IIS, and TOR—constitute the reproduction regulatory network in insect reproduction [45–48].

Preliminary laboratory studies have shown that TRE plays an important role in the reproduction regulation of *N. lugens* [49]. However, it is not clear how TRE regulates the reproduction of *N. lugens*. Therefore, in this study, the TRE in *N. lugens* was inhibited in vivo at the mRNA and protein levels via dsTREs and validamycin injections, respectively, and the effects of TRE on the reproduction regulatory network were determined. The results of the study are conducive to a more comprehensive assessment of the ability of TRE as a pest control target and provide theoretical support for the development of new and safe biopesticides.

2. Materials and Methods

2.1. Insects

The *N. lugens* insects used in this study were obtained from local rice fields at the China National Rice Research Institute (Hangzhou, China). The institute was established in 2013. The rice variety used during the experiment was Taichung Native 1 (TN1), which was used to feed *N. lugens*. The environmental conditions of the artificial climate chamber were set as follows: temperature 27 ± 1 °C, relative humidity 65 ± 5%, and photoperiod 18L: 6D (Light: Dark). All experiments were performed under these same conditions to ensure consistency.

2.2. dsRNA Synthesis and Microinjection

Twenty adult *N. lugens* were randomly divided into 4 groups (five adults per group), which were then crushed in an EP tube and added to Trizol (Takara, Kyoto, Japan) for RNA extraction. After a series of operations, the obtained RNA was added to an appropriate amount of DEPC water and completely dissolved. Then, 1 μL of RNA solution was detected using the micro nucleic acid protein concentration analyzer Nanodrop2000 (Thermo Fisher, Waltham, MA, USA) to determine the RNA purity and concentration. RNA integrity was detected via 1% agarose gel electrophoresis. Using the extracted RNA as a template, cDNA was synthesized using the Prime Script RT reagent Kit with gDNA Eraser (Takara, Kyoto, Japan). The in vitro synthesis of dsRNA was performed using the T7 RiboMAXTM Express RNAi System (Promega, Madison, WI, USA), a dsRNA synthesis kit specifically designed for this purpose. Newly emerged long-winged females were selected for injection with the TransferMan 4r microinjector (Eppendorf, Hamburger, Germany). The dsRNA concentration was 4000 ng/μL, the validamycin concetration was 0.5 μg/μL, and each group was injected with 100 nL [15,16]. The dsTREs used in this study has successfully verified the inhibitory effect on *TRE1-1*, *TRE1-2*, and *TRE2* gene expression in earlier studies of our laboratory, with an inhibitory duration of 6 days [49]. The accession numbers and primer sequences of these three genes are presented in Table 1.

Table 1. Primers for dsRNA synthesis.

Primer Name	Accession Number	Primer Sequences (5′-3′)		Length
NlTRE1-1	FJ790319.1	F: GATGCAATCAAGGAGGTGTTATGGC	R: CGTATTCACCTCCACCTCCGT	451 bp
NlTRE1-2	KU556829.1	F: AGATGAAGGCATGTGGTTCG	R: CATCGATTCGCCAACTGGTAAGC	321 bp
NlTRE2	GQ397451.1	F: CCAACTGCTATGACACCGACAAG	R: GGGTTCAGATCCTGCCGTCGCT	440 bp
GFP	MW987535.2	F: AAGGGCGAGGAGCTGTTCACCG	R: CAGCAGGACCATGTGATCGCGC	720 bp

2.3. qRT-PCR

Five females were selected for RNA extraction and cDNA reverse transcription. The specific primers (Table 2) were synthesized, and the actin gene was used as an internal reference gene [27,45,46,50–52]. Subsequently, the synthesized cDNA was used as a template for qRT-PCR. The qRT-PCR was conducted with TB Green Premix Ex Taq™ (Takara, Kyoto, Japan). QRT-PCR was performed on a Bio-Rad CFX96TM real-time PCR detection system (Bio-Rad, Hercules, CA, USA) with the following reaction system: ddH_2O 3.2 μL, TB Green Premix Ex Taq 5 μL, forward primer 0.4 μL, reverse primer 0.4 μL, and cDNA 1 μL. The specificity of the amplification reaction was judged by the amplification curve as well as the melting curve, and the relative expression of the genes was calculated according to the $2^{-\triangle\triangle CT}$ method [53].

Table 2. Primers for qRT-PCR.

Primer Name	Accession Number	Primer Sequences (5′-3′)	
QNlActin	EU179847.1	F: TGGACTTCGAGCAGGAAATGG	R: ACGTCGCACTTCATGATCGAG
QNlVg	JF330416.1	F: CACTGCCCGTGCTGTGCTCTA	R: TGACTTCCTTGCTTTGCTCCC
QNlVgR	JQ040014.1	F: AGGCAGCCACACAGATAACCGC	R: AGCCGCTCGCTCCAGAACATT
QNlJHAMT	KP769805.1	F: GAACCTGCAGGCCAAACACA	R: ACCACTCGGTTGGGCTGAAT
QNlMet	KP797880.1	F: AGTGGCAGCGAGCGATGATT	R: TGAGGCGCAGCAAAAAGGAG
QNlUSP	KX431887.1	F: GGTGGAGCTGCTGAGGGAGA	R: AGCACTTGAGGCCGATGGAG
QNlEcR	FJ263049.1	F: CGAAGCCTGGAAGGTGGAGA	R: GGCAAAGATTGGCGACGATT
QNlInR1	KF974333.1	F: GAGTGCAACCCGGAGTATGT	R: TCTTGACGGCACACTTCTTG
QNlInR2	KF974334.1	F: CTCTTGCCGAACAGCCTTAC	R: GGGTCGTTTAGTGGGTCTGA
QNlTOR	JQ793898.1	F: GGCTACAGGGATGTCAAA	R: GAGATAGATTCAAACGGAAAG
QNlS6K	KP769804.1	F: AATCGGACGACTTGGAGACAGT	R: CAGTTTGGAAAGCGTACATCAGG
QNlAKH	AB817235.1	F: CCCTTCTGATGGCAGTCCTTTG	R: ATGGATGCCTTGCAGCCTTCT
QNlFas	XM_022339189.2	F: CGGAGACTCTGCCCTAA	R: CAGCGACTAATCCAACATC

2.4. Observation of Abdomen and Detection of Triglyceride Content

The injected female *N. lugens* was paired with untreated male *N. lugens* in a 1:1 ratio. The female *N. lugens* (n = 20, consistent with Section 2.2 cohorts) abdomen was observed and photographed at a magnification of 20×. After three days, the fat bodies and ovaries of these females were collected. The weight of each tissue sample was recorded using an electronic balance. Differences in the triglyceride determination methods were applied for fat body and ovary samples. Anhydrous ethanol was added to the fat body samples to form homogenates, while 1 × PBS was used in another set of preparations. And 10% homogenates were formed by grinding and cell crushing, respectively. After the reaction, both sets of samples were then processed with a triglyceride assay kit (Nanjing Jiancheng, Nanjing, China). The absorbance was measured at 510 nm using an enzyme-labeled microplate reader. The darker color produced during the reaction is proportional to the triglyceride content. Additionally, since ovaries are not high-fat samples, the protein content is also tested to normalize the triglyceride content. Protein was detected using the BCA protein detection kit (Beyotime, Shanghai, China). The triglyceride content in fat bodies was calculated as (Sample OD − Blank OD)/(Calibration OD − Blank OD) × Calibrator concentration (mmol/L). For ovaries, the triglyceride content was adjusted by dividing the above result by the protein concentration in the samples to be measured (gprot/L).

2.5. Data Analysis

Microsoft Excel 2019 was used to compile the data, GraphPad Prism 9.0.0 was used for graphing, and IBM SPSS Statistics 23.0 was used for statistical analysis. All data were presented as the means ± standard errors (SEs) of biological replicates, and Student's *t* test was used to analyze the difference.

3. Results

3.1. Effects of Trehalase on Abdominal Volumes, Vg, and VgR Expression

The results showed that on the third day after the injection of dsGFP, dsTREs, and ddwater, the abdomens of *N. lugens* exhibited no significant differences. In contrast, those injected with validamycin exhibited abdominal enlargement, with the intersegmental membrane of the sternum fully distended (Figure 1A). The mRNA level of the *Vg* gene decreased significantly on the third day after dsTREs injection, but recovered by the sixth day, showing no significant difference compared to the control (Figure 1B). The mRNA level of *VgR*, on the other hand, remained comparable to that of the control group both on the third and sixth day after interference (Figure 1C). When *N. lugens* were injected with validamycin, the expression level of *Vg* in its body was extremely suppressed both on the third and sixth day (Figure 1D). However, similar to the results of dsTREs injection, the mRNA level of *VgR* did not change significantly compared with the control group, indicating that its expression was not affected by the suppression of TRE (Figure 1E).




Figure 1. Comparison of abdominal volumes in *N. lugens* following injections (**A**) (Scale bar: 1 mm). The relative expression levels of *Vg* (**B**,**D**) and *VgR* (**C**,**E**) were detected on the third and sixth day after dsTREs injection and validamycin injection. Those injected with dsGFP and ddWater are the control group, and those injected with dsTREs and validamycin are the experimental group. (**A**) The shooting time was the third day after the injection. The magnification of the abdomen was 20×. The mRNA levels of *Vg* (**B**,**D**) and *VgR* (**C**,**E**) in *N. lugens* were detected on the third and sixth day after injection of dsTREs and validamycin, and the effects of trehalase on *Vg* and *VgR* expression were investigated. The data was shown as mean ± standard errors (n ≥ 3) and analyzed using Student's *t* test. "**", $p < 0.01$; "*", $p < 0.05$; "ns", $p > 0.05$.

3.2. Effects of Trehalase on JH and 20E Signaling Pathways

The expression level of the 20E receptor *USP* significantly increased on the third and sixth day (Figure 2C). The expression level of another 20E receptor, *EcR*, was also not significantly different compared to the control group on the third day or sixth day after dsTREs injection (Figure 2D). Similarly, the mRNA levels of *JHAMT* were comparable to controls on the third and sixth day after validamycin injection, but *Met* expression was significantly downregulated on the third day after validamycin injection, suggesting that validamycin reduced signaling in the JH pathway (Figure 2E,F). The changing trends of *USP* and *EcR* levels after validamycin injection were similar. There was no significant difference on the third day, but both showed a significant decrease by the sixth day. This suggests that validamycin reduced signaling in the 20E pathway; however, its effect was delayed compared to that on the JH signaling pathway (Figure 2G,H).




Figure 2. Effects of trehalase on JH and 20E signaling pathways. The relative expression of genes related to juvenile hormone signaling and ecdysone signaling on the third and sixth day after dsTREs and validamycin injection. The mRNA levels of *JHAMT* (**A**,**E**), *Met* (**B**,**F**), *USP* (**C**,**G**), and *EcR* (**D**,**H**) in *N. lugens* were detected on the third and sixth day after injection of dsTREs and validamycin, and

the effects of trehalase on juvenile hormone signaling and ecdysone signaling expression were investigated. The data was shown as mean ± standard errors (n ≥ 3) and analyzed using Student's *t* test. "**", $p < 0.01$; "*", $p < 0.05$; "ns", $p > 0.05$.

3.3. Effects of Trehalase on Nutrient Signaling Pathways

Two insulin receptor genes, *InR1* and *InR2*, have been identified in *N. lugens* [50]. When the expression of *NlTRE1-1*, *NlTRE1-2*, and *NlTRE2* was simultaneously inhibited [49], the qRT-PCR results revealed no significant change in *InR1* expression on the third and sixth day compared with controls (Figure 3A,B). In contrast, *InR2* expression decreased significantly on the third and sixth day compared with controls (Figure 3A,B). *TOR* expression did not differ significantly from the control group on the third day after dsTREs injection, but increased significantly on the sixth day (Figure 3C). However, the specific indicator shown in Figure 3D exhibited a decrease on the third day and an increase after the sixth day compared to the dsGFP-injected group, although this difference was not statistically significant (Figure 3D).




Figure 3. Effects of trehalase on nutrient signaling pathways. The relative expression of genes related to juvenile hormone signaling and ecdysone signaling on the third and sixth day after dsTREs and

validamycin injection. The mRNA levels of *InR1* (**A**,**E**), *InR2* (**B**,**F**), *TOR* (**C**,**G**), and *S6K* (**D**,**H**) in *N. lugens* were detected on the third and sixth day after injection of dsTREs and validamycin, and the effects of trehalase on juvenile hormone signaling and ecdysone signaling expression were investigated. The data was shown as mean ± standard errors ($n \geq 3$) and analyzed using Student's *t* test. "**", $p < 0.01$; "*", $p < 0.05$; "ns", $p > 0.05$.

When the TRE activity was inhibited by validamycin injection, *InR1* expression showed no significant change on the third day, but decreased highly significantly after the sixth day. *InR2* expression remained comparable to the control levels on both the third day and the sixth day of inhibition (Figure 3E,F). Similarly, the trends in the expression of key genes in the *TOR* signaling pathway were inconsistent with the results of dsTREs interference. On the third day after validamycin injection, *TOR* and *S6K* expression did not differ significantly from controls, but both were very significantly reduced on the sixth day (Figure 3G,H).

3.4. Effects of Trehalase on Lipid Metabolism

The experimental results showed that three days after dsTREs injection, the content of triglycerides in both the fat body and ovaries did not differ significantly from that of the control group (Figure 4A,C). When injected with validamycin on the third day, the triglyceride content in the fat body of females remained unaffected, but the triglyceride content in the ovaries decreased very significantly, suggesting that the trehalase inhibitor prevented the accumulation of triglycerides in the ovaries (Figure 4B,D).




Figure 4. The triglyceride content in the fat body and ovary of *N. lugens* was measured on the third day following the injection of dsTREs and validamycin. The effect of trehalase on lipid metabolism was analyzed by measuring the content of triglyceride in fat body (**A**,**B**) and ovary (**C**,**D**) on the third day of injection. The data was shown as mean ± standard errors ($n = 3$) and analyzed using Student's *t* test. "**", $p < 0.01$; "ns", $p > 0.05$.

Using qRT-PCR, we further examined the effects of TRE interference or activity inhibition on lipid metabolism. The results revealed that the mRNA level of fatty acid synthase (*Fas*) decreased significantly three days after dsTREs injection, but recovered to control

level by the sixth day (Figure 5A). The relative expression of Adipokinetic hormone (*AKH*), responsible for lipid mobilization, remained comparable to the control levels on both the third and sixth day (Figure 5B). After validamycin injection, the relative expression of *Fas* decreased substantially on both the third and sixth day, but this difference was statistically significant only on the third day (Figure 5C). Similarly, the relative expression of *AKH* decreased substantially on both the third and sixth day, although neither change was statistically significant (Figure 5D).




Figure 5. The relative expression of *Fas* (**A**,**C**) and *AKH* (**B**,**D**) of *N. lugens* on the third and sixth day after dsTREs injection and validamycin injection. The effect of TRE interference or activity inhibition on lipid metabolism of *N. lugens* was detected by qRT-PCR dsTREs. The data was shown as mean ± standard errors ($n \geq 3$) and analyzed using Student's *t* test. "*", $p < 0.05$; "ns", $p > 0.05$.

4. Discussion

Vitellogenin (Vg) plays a crucial role in the reproduction of *N. lugens*. *Vg* expression significantly increases after female emergence, and any disruption to this process results in infertility and abnormal oocyte development in *N. lugens* females [27]. The transporter receptor for *Vg*, *VgR*, is also highly expressed following emergence; interference with its function prevents Vg from accumulating in the ovary, causing decreased female fecundity [54]. Previous research on *Spodoptera frugiperda* indicated that the injection of trehalase inhibitors reduced *Vg* expression [17]. Consistent with these findings, our experimental results demonstrated that both dsTREs and validamycin led to the downregulation of *Vg* expression on the third day after injection (Figure 1A,C). Notably, *Vg* expression recovered later in the season after dsTREs injection (Figure 1A), while validamycin exhibited a more prolonged effect (Figure 1C), which correlated with egg production outcomes [49]. The differential effect between dsTREs and validamycin treatment may stem from their different mechanisms of action. Although RNA interference technology can reduce the expression level of TRE transcripts, due to the non-100% silencing effect and the inherent stability of the encoded enzyme, residual enzyme activity may still exist, leading to incomplete inhibition. In contrast, validamycin can directly inhibit enzyme activity by binding to the active site of the enzyme, regardless of the transcript level. The differences observed in

the transcriptional regulation of Vg by dsTREs versus validamycin suggest potential variations in their reproductive regulatory mechanisms. It is noteworthy that neither treatment influenced *VgR* expression levels (Figure 1B,D), contrasting with findings reported for *S. frugiperda* [9]. This implies that TREs specifically regulate Vg synthesis without affecting its translocation into the ovary. Importantly, although *Vg* was down-regulated shortly after silencing the *TRE* gene (Figure 1B,D), egg production did not decline until later stages [49]. This may be attributed to a lag in translation relative to transcription.

Previous studies have shown that the expression of *Vg* is regulated by multiple signaling pathways, including JH, 20E, IIS, and TOR. Affecting key components of these pathways can impact embryo development and decrease hatchability in *N. lugens*, collectively forming a reproductive regulatory network [45–48]. In this study, TRE inhibitors suppressed both JH and 20E signaling pathways (Figure 2C,E,F), consistent with observations of inhibited egg development and reduced egg-laying rates, which are in agreement with prior reports [45,48]. However, dsTREs treatment did not affect the expression of *JHAMT* or *Met* (Figure 2A,B), but it upregulated the expression of *USP* and *EcR*, key receptors in the 20E signaling pathway (Figure 2C,D). This contrasts with the observed decline in fecundity [49]. Studies on *Spodoptera frugiperda* have demonstrated that downregulating cuticular 20E synthesis enzymes may be a regulated event, signaling slowed larval growth and pre-metamorphic physiological changes [36]. Additionally, host plant nutrients modulate JH and 20E levels in *S. frugiperda*, influencing migration and reproduction [3]. The alteration of the trehalose concentration in hemolymph by silencing TREs in this study may not have been sufficient to affect JH synthesis. Furthermore, since 20E upregulates TRE activity in *Antheraea pernyi* [55], we propose that the low trehalase mRNA levels observed here induced the upregulation of 20E signaling (Figure 2C) as a negative feedback mechanism to restore trehalose homeostasis.

The above analysis indicates that the effect of dsTREs on egg production was not mediated through changes in the JH and 20E levels. However, experimental results revealed that *InR2* expression was significantly downregulated after dsTREs injection (Figure 3B). Previous studies have identified two insulin-like receptors, *InR1* and *InR2*, in *N. lugens* [50,56]. Among these, *InR1* functions as a canonical insulin-like receptor activating the PI3K/Akt pathway, whereas *InR2* serves as a negative regulator of this pathway. Our experimental treatment did not significantly affect ovarian morphology, but led to reduced egg production (Figure 3E,F), findings that are consistent with the previous research [50,57]. Furthermore, dsTREs injection initially caused the downregulation of *S6K* (Figure 3D), and the RNAi-mediated downregulation of *S6K* has been shown to inhibit Vg synthesis (Figure 3H), which aligns with the trend of the *Vg* changes observed in this experiment [46]. Conversely, unlike dsTREs treatment, the injection of validamycin had no impact on the *InR2* expression levels (Figure 3F), but significantly suppressed *InR1* expression (Figure 3E). Notably, *InR1* silencing had a much stronger inhibitory effect on reproduction, which is consistent with the pronounced inhibitory effect of validamycin on *N. lugens* reproduction observed in the present study [58]. Additionally, consistent with its effects on JH and 20E, validamycin injection inhibited IIS and TOR signaling during the late stage, suggesting that there may be other pathways involved in the early-stage inhibition of reproduction via validamycin, whereas changes in the classical regulatory network of reproduction are responsible for the inhibition observed in the late stage.

Lipids, primarily in the form of triglycerides, constitute over 50% of the dry weight of the fat body and represent approximately 30–40% of the oocyte's dry weight [59]. These findings indicate that triglycerides serve as a primary energy source for oocyte maturation and embryo development [60,61]. In oviparous insects, oocyte maturation relies not only on Vg accumulation, but also on lipid accumulation [60,61]. An interesting observation in

this study was that when *N. lugens* was injected with validamycin, the abdomen exhibited abnormal enlargement compared to the control (Figure 1A), whereas the ovarian development volume was significantly smaller [49]. Given that the abdomen of female *N. lugens* is predominantly composed of ovaries and fat bodies, we hypothesized that the above phenomenon is caused by the abnormal distribution of fat bodies. Typically, the abnormal enlargement of fat bodies is closely associated with reproductive defects, potentially due to energy trade-offs—a process where resources are allocated between different physiological functions [62,63]. In contrast, abnormal lipid metabolism, a key aspect of adipocyte function, is linked to an altered fat body morphology. The blockage of triglyceride transport from the fat body to the ovary produces a phenotype consistent with that observed in this study [61–63]. Therefore, this study examined lipid metabolism and revealed a significant decrease in the mean expression levels of *Fas* and *AKH* following validamycin treatment (Figure 5C,D). These genes are responsible for fatty acid synthesis and lipid mobilization, respectively, and their downregulation inhibits reproduction in *N. lugens* [51,64]. However, the downregulation of *AKH* was not significant (Figure 5D), potentially due to large errors resulting from insufficient biological replicates. Correspondingly, the triglyceride content in the ovary was also significantly reduced after validamycin injection, indicating that the inhibition of TRE activity impacted lipid metabolism in the fat body, thereby reducing the amount of triglycerides transported to the ovary and ultimately leading to impaired egg development. Studies in mice have demonstrated that alginate suppresses adipocyte hypertrophy; however, the inability to utilize alginate in this study might have disrupted lipid synthesis [65].

Notably, dsTREs did not significantly affect the ovarian triglyceride content (Figure 4A,B) and ovarian morphology [49]; however, they did reduce the expression levels of *Fas* and *AKH* (Figure 5A,B). This discrepancy may be attributed to the weaker inhibitory effect of the pre-existing RNAi on TRE compared to that of validamycin.

5. Conclusions

In summary, both dsTREs and validamycin reduced the expression level of *Vg* without altering the level of its receptor *VgR*. However, their impacts on the reproductive regulatory network of *N. lugens* differed significantly. dsTREs did not influence the expression of *JHAMT* and *Met* within the juvenile hormone signaling pathway, but upregulated the mRNA levels of ecdysone receptor *USP* and *TOR*, while suppressing the expression of insulin-like receptors *InR2* and *S6K*. In contrast, validamycin downregulated the expression of *Met*, *USP*, *EcR*, *InR1*, *TOR* and *S6K*. Moreover, both dsTREs and validamycin inhibited *Fas* expression; however, only validamycin decreased triglyceride content in ovaries and induced abnormal abdominal enlargement of the abdomen in females.

Author Contributions: B.T.: Writing—review and editing, Supervision, Resources, Project administration, Funding acquisition, and Conceptualization. Y.G.: Writing—original draft, review and editing, Visualization, Investigation, and Data curation. Y.L.: Writing—original draft, review and editing, Methodology, Investigation, and Data curation. L.G.: Writing—review and editing, Methodology. Y.H.: Investigation, and Visualization. Y.Z.: Investigation and Visualization. G.H.: Writing—review and editing and Supervision. Y.W.: Writing—review and editing, Conceptualization, and Supervision. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the National Natural Science Foundation of China (Grant No. 32272608 and 32460668).

Data Availability Statement: The original contributions presented in this study are included in the article. Further inquiries can be directed to the corresponding authors.

Conflicts of Interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Xu, H.J.; Zhang, C.X. Insulin receptors and wing dimorphism in rice planthoppers. *Philos. Trans. R. Soc. B Biol. Sci.* **2017**, *372*, 20150489. [CrossRef] [PubMed]
2. Lin, X.M.; Zhang, H.H.; Gao, H.L.; Yuan, X.W.; Liu, Z.W. The transcription factor CREB3-2 regulated neutral lipase gene expression in ovary of *Nilaparvata lugens*. *Pestic. Biochem. Physiol.* **2023**, *196*, 105632. [CrossRef] [PubMed]
3. Zhang, H.H.; Yang, B.J.; Xu, X.X.; Zou, J.Z.; Liu, Z.W. Roles of spermary-specific carboxylesterases in *Nilaparvata lugens* reproduction: Opposite between insecticide-induced upregulation and resistance-associated overexpression. *Int. J. Biol. Macromol.* **2024**, *290*, 138991. [CrossRef]
4. Xia, X.; Luo, G.H.; Yu, J.L.; Liu, B.Q.; Zhang, R.; Zhang, G.; Shu, Z.L.; Lou, Y.G.; Hoffmann, A.A.; Fang, J.C. Jasmonic acid changes associated with long-term control of lepidopteran rice pests after insecticide seed treatment. *Entomol. Gen.* **2024**, *44*, 573–581. [CrossRef]
5. Yang, B.; Lin, X.; Yu, N.; Gao, H.; Zhang, Y.; Liu, W.; Liu, Z. Contribution of glutathione s-transferases to imidacloprid resistance in *Nilaparvata lugens*. *J. Agric. Food Chem.* **2020**, *68*, 15403–15408. [CrossRef]
6. Tang, B.; Xu, K.; Liu, Y.; Zhou, Z.; Karthi, S.; Yang, H.; Li, C. A review of physiological resistance to insecticide stress in *Nilaparvata lugens*. *3 Biotech* **2022**, *12*, 84. [CrossRef]
7. Wang, L.X.; Tao, S.; Zhang, Y.C.; Pei, X.G.; Gao, Y.; Song, X.Y.; Yu, Z.T.; Gao, C.F. Overexpression of ATP-binding cassette transporter Mdr49-like confers resistance to imidacloprid in the field populations of brown planthopper, *Nilaparvata lugens*. *Pest Manag. Sci.* **2022**, *78*, 579–590. [CrossRef]
8. Huang, J.H.; Lee, H.J. RNA interference unveils functions of the hypertrehalosemic hormone on cyclic fluctuation of hemolymph trehalose and oviposition in the virgin female *Blattella germanica*. *J. Insect Physiol.* **2011**, *57*, 858–864. [CrossRef]
9. Tang, B.; Hu, S.R.; Luo, Y.J.; Shi, D.M.; Liu, X.Y.; Zhong, F.; Jiang, X.Y.; Hu, G.; Li, C.; Duan, H.X.; et al. Impact of Three Thiazolidinone Compounds with Piperine Skeletons on Trehalase Activity and Development of *Spodoptera frugiperda* Larvae. *J. Agric. Food Chem.* **2024**, *72*, 8423–8433. [CrossRef]
10. Li, Y.; Wang, S.S.; Si, H.R.; Wan, S.J.; Li, G.Y.; Shu, Y.H.; Dai, X.Y.; Wang, R.J.; Wang, S.G.; Zhai, Y.F.; et al. Responses of aphid and ladybird to lead transfer through soil and broad beans. *Entomol. Gen.* **2024**, *44*, 621–623. [CrossRef]
11. Avonce, N.; Mendoza-Vargas, A.; Morett, E.; Morett, E.; Iturriaga, G. Insights on the evolution of trehalose biosynthesis. *BMC Evol. Biol.* **2006**, *6*, 109. [CrossRef]
12. Mitsumasu, K.; Azuma, M.; Niimi, T.; Yamashita, O.; Yaginuma, T. Membrane-penetrating trehalase from *Bombyx mori*. *Molecular cloning and localization in larval midgut*. *Insect Mol. Biol.* **2005**, *14*, 501–508. [PubMed]
13. Lee, J.H.; Saito, S.; Mori, H.; Nishimoto, M.; Okuyama, M.; Kim, D.; Wongchawalit, J.; Kimura, A.; Chiba, S. Molecular cloning of cDNA for trehalase from the European honeybee, *Apis mellifera* L., and its heterologous expression in *Pichia pastoris*. *Biosci. Biotechnol. Biochem.* **2007**, *71*, 2256–2265. [CrossRef] [PubMed]
14. Tang, B.; Wei, P.; Zhao, L.N.; Shi, Z.; Shen, Q.; Yang, M.; Xie, G.; Wang, S. Knockdown of five trehalase genes using RNA interference regulates the gene expression of the chitin biosynthesis pathways in *Tribolium castaneum*. *BMC Biotechnol.* **2016**, *16*, 67. [CrossRef] [PubMed]
15. Zhao, L.N.; Yang, M.M.; Shen, Q.D.; Liu, X.J.; Shi, Z.K.; Wang, S.G.; Tang, B. Functional characterization of three trehalase genes regulating the chitin metabolism pathway in rice brown planthopper using RNA interference. *Sci. Rep.* **2016**, *6*, 27841. [CrossRef]
16. Tang, B.; Yang, M.M.; Shen, Q.D.; Xu, Y.X.; Wang, H.J.; Wang, S.G. Suppressing the activity of trehalase with validamycin disrupts the trehalose and chitin biosynthesis pathways in the rice brown planthopper, *Nilaparvata lugens*. *Pestic. Biochem. Physiol.* **2017**, *137*, 81–90. [CrossRef]
17. Tang, B.; Han, Y.; Mao, Q.X.; Fu, H.Y.; Luo, Y.J.; Hua, L.Y.H.; Liu, B.S.; Hu, G.; Wang, S.G.; Desneux, N.; et al. Regulation of three novel pepper thiothiazolidinones on the fecundity of *Spodoptera frugiperda*. *Pestic. Biochem. Physiol.* **2024**, *204*, 106033. [CrossRef]
18. Matassini, C.; Parmeggiani, C.; Cardona, F. New Frontiers on human safe insecticides and fungicides: An opinion on trehalase inhibitors. *Molecules* **2020**, *25*, 3013. [CrossRef]
19. Yu, H.Z.; Zhang, Q.; Lu, Z.J.; Deng, M.J. Validamycin treatment significantly inhibits the glycometabolism and chitin synthesis in the common cutworm, *Spodoptera litura*. *Insect Sci.* **2022**, *29*, 840–854. [CrossRef]
20. Turna Demir, F.; Demir, E. In vivo evaluation of the neurogenotoxic effects of exposure to validamycin A in neuroblasts of *Drosophila melanogaster* larval brain. *J. Appl. Toxicol.* **2024**, *44*, 355–370. [CrossRef]
21. Tatun, N.; Wangsantitham, O.; Tungitwitayakul, J.; Sakvrai, S. Trehalase activity in fungus-growing termite, *Odontotermes feae* (Isoptera: Termitideae) and inhibitory effect of validamycin. *J. Econ. Entomol.* **2014**, *107*, 1224–1232. [CrossRef]
22. Tatun, N.; Tungjitwitayakul, J.; Sakurai, S. Developmental and Lethal Effects of Trehalase Inhibitor (Validamycin) on the *Tribolium castaneum* (*Coleoptera: Tenebrionidae*). *Ann. Entomol. Soc. Am.* **2016**, *109*, 224–231. [CrossRef]

23. Wangsantitham, O.; Pothong, P.; Tungjitwitayakul, J.; Tatun, N. Suppression of Trehalase Activity by Validamycin Induces Mortality and Developmental Delays in *Sitophilus zeamais* (*Coleoptera: Curculionidae*). *J. Entomol. Sci.* **2024**, *59*, 311–322. [CrossRef]
24. Marten, A.D.; Stothard, A.I.; Kalera, K.; Swarts, B.M.; Conway, M.J. Validamycin A Delays Development and Prevents Flight in *Aedes aegypti* (*Diptera: Culicidae*). *J. Med. Entomol.* **2020**, *57*, 1096–1103. [CrossRef] [PubMed]
25. Shi, S.; Zuo, H.; Gao, L.; Yi, X.; Zhong, G. Silencing of rieske iron-sulfur protein impacts upon the development and reproduction of *Spodoptera exigua* by regulating ATP synthesis. *Front. Physiol.* **2018**, *9*, 575. [CrossRef]
26. Tufail, M.; Takeda, M. Molecular characteristics of insect vitellogenins. *J. Insect Physiol.* **2008**, *54*, 1447–1458. [CrossRef] [PubMed]
27. Shen, Y.; Chen, Y.Z.; Lou, Y.H.; Zhang, C.X. Vitellogenin and vitellogenin-like genes in the brown planthopper. *Front. Physiol.* **2019**, *10*, 1181. [CrossRef]
28. Wu, Z.X.; Yang, L.B.; He, Q.J.; Zhou, S.T. Regulatory mechanisms of vitellogenesis in insects. *Front. Cell Dev. Biol.* **2021**, *8*, 593613. [CrossRef]
29. Zou, Z.; Saha, T.T.; Roy, S.; Shin, S.W.; Backman, T.W.; Girke, T.; White, K.P.; Raikhel, A.S. Juvenile hormone and its receptor, methoprene-tolerant, control the dynamics of mosquito gene expression. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, E2173–E2181. [CrossRef]
30. Hansen, I.A.; Attardo, G.M.; Rodriguez, S.D.; Drake, L.L. Four-way regulation of mosquito yolk protein precursor genes by juvenile hormone-, ecdysone-, nutrient-, and insulin-like peptide signaling pathways. *Front. Physiol.* **2014**, *5*, 103. [CrossRef]
31. Roy, S.; Saha, T.T.; Zou, Z.; Raikhel, A.S. Regulatory pathways controlling female insect reproduction. *Annu. Rev. Entomol.* **2018**, *63*, 489–511. [CrossRef]
32. Wu, Z.; Guo, W.; Yang, L.; He, Q.; Zhou, S. Juvenile hormone promotes locust fat body cell polyploidization and vitellogenesis by activating the transcription of Cdk6 and E2f1. *Insect Biochem. Mol. Biol.* **2018**, *102*, 1–10. [CrossRef] [PubMed]
33. Santos, C.G.; Humann, F.C.; Hartfelder, K. Juvenile hormone signaling in insect oogenesis. *Curr. Opin. Insect Sci.* **2019**, *31*, 43–48. [CrossRef] [PubMed]
34. Ling, L.; Raikhel, A.S. Cross-talk of insulin-like peptides, juvenile hormone, and 20-hydroxyecdysone in regulation of metabolism in the mosquito *Aedes aegypti*. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2023470118. [CrossRef] [PubMed]
35. Luo, W.; Liu, S.; Zhang, W.; Yang, L.; Huang, J.; Zhou, S.; Feng, Q.; Palli, S.R.; Wang, J.; Roth, S.; et al. Juvenile hormone signaling promotes ovulation and maintains egg shape by inducing expression of extracellular matrix genes. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2104461118. [CrossRef]
36. Wu, M.Z.; Fan, S.T.; Zhang, Y.C.; Tan, J.F.; Zhu, G.H. Disrupting shadow in the prothoracic gland induced larval development arrest in the fall armyworm *Spodoptera frugiperda*. *Front. Physiol.* **2024**, *15*, 1502753. [CrossRef]
37. Ekoka, E.; Maharaj, S.; Nardini, L.; Dahan-Moss, Y.; Koekemoer, L.L. 20-Hydroxyecdysone (20E) signaling as a promising target for the chemical control of malaria vectors. *Parasites Vectors* **2021**, *14*, 86. [CrossRef]
38. Wu, Q.; Brown, M.R. Signaling and function of insulin-like peptides in insects. *Annu. Rev. Entomol.* **2006**, *51*, 1–24. [CrossRef]
39. Kannan, K.; Fridell, Y.W. Functional implications of Drosophila insulin-like peptides in metabolism, aging, and dietary restriction. *Front. Physiol.* **2013**, *4*, 288. [CrossRef]
40. Dong, Y.; Chen, W.; Kang, K.; Pang, R.; Dong, Y.; Liu, K.; Zhang, W. FoxO directly regulates the expression of TOR/S6K and vitellogenin to modulate the fecundity of the brown planthopper. *Sci. China Life Sci.* **2021**, *64*, 133–143. [CrossRef]
41. Huangfu, N.; Zhu, X.; Wang, L.; Zhang, K.; Li, D.; Chen, L.; Gao, X.; Niu, L.; Gao, M.; Ji, J.; et al. Insulin receptor substrate-1 (IRS1) regulates oogenesis and vitellogenesis in *Propylea japonica* by Mediating the FOXO transcription factor expression, independent of JH and 20E signaling pathways. *J. Agric. Food Chem.* **2023**, *71*, 300–310. [CrossRef] [PubMed]
42. Rauschenbach, I.Y.; Karpova, E.K.; Burdina, E.V.; Adonyeva, N.V.; Bykov, R.A.; Ilinsky, Y.Y.; Menshanov, P.N.; Gruntenko, N.E. Insulin-like peptide DILP6 regulates juvenile hormone and dopamine metabolism in Drosophila females. *Gen. Comp. Endocrinol.* **2017**, *243*, 1–9. [CrossRef] [PubMed]
43. Smykal, V.; Raikhel, A.S. Nutritional control of insect reproduction. *Curr. Opin. Insect Sci.* **2015**, *11*, 31–38. [CrossRef] [PubMed]
44. Katewa, S.D.; Kapahi, P. Role of TOR signaling in aging and related biological processes in *Drosophila melanogaster*. *Exp. Gerontol.* **2011**, *46*, 382–390. [CrossRef]
45. Lu, K.; Chen, X.; Liu, W.T.; Zhang, X.Y.; Chen, M.X.; Zhou, Q. Nutritional signaling regulates vitellogenin synthesis and egg development through juvenile hormone in *Nilaparvata lugens* (Stål). *Int. J. Mol. Sci.* **2016**, *17*, 269. [CrossRef]
46. Lu, K.; Chen, X.; Liu, W.T.; Zhou, Q. TOR pathway-mediated juvenile hormone synthesis regulates nutrient-dependent female reproduction in *Nilaparvata lugens* (Stål). *Int. J. Mol. Sci.* **2016**, *17*, 438. [CrossRef]
47. Lu, K.; Chen, X.; Li, W.; Li, Y.; Zhang, Z.; Zhou, Q. Insulin-like peptides and DNA/tRNA methyltransferases are involved in the nutritional regulation of female reproduction in *Nilaparvata lugens* (Stål). *Gene* **2018**, *639*, 96–105. [CrossRef]
48. Zhou, X.; Ye, Y.Z.; Ogihara, M.H.; Takeshima, M.; Fujinaga, D.; Liu, C.W.; Zhu, Z.; Kataoka, H.; Bao, Y.Y. Functional analysis of ecdysteroid biosynthetic enzymes of the rice planthopper, *Nilaparvata lugens*. *Insect Biochem. Mol. Biol.* **2020**, *123*, 103428. [CrossRef]

49. Liu, Y.K.; Zhu, Y.; Wan, S.J.; Wang, X.Z.; Guan, L.W.; Xu, C.D.; Xie, B.H.; Wang, S.G.; Sun, S.S.; Tang, B. Trehalase regulates ovarian maturation and egg hatchability of *Nilaparvata lugens* (Stål). *J. Pest Sci.* **2025**, *31*, 1.
50. Xu, H.J.; Xue, J.; Lu, B.; Zhang, X.C.; Zhuo, J.C.; He, S.F.; Ma, X.F.; Jiang, Y.Q.; Fan, H.W.; Xu, J.Y.; et al. Two insulin receptors determine alternative wing morphs in planthoppers. *Nature* **2015**, *519*, 464–467. [CrossRef]
51. Li, L.; Jiang, Y.; Liu, Z.; You, L.; Wu, Y.; Xu, B.; Ge, L.; Stanley, D.; Song, Q.; Wu, J. Jinggangmycin increases fecundity of the brown planthopper, *Nilaparvata lugens* (Stål) via fatty acid synthase gene expression. *J. Proteom.* **2016**, *130*, 140–149. [CrossRef] [PubMed]
52. Lu, K.; Chen, X.; Li, Y.; Li, W.; Zhou, Q. Lipophorin receptor regulates *Nilaparvata lugens* fecundity by promoting lipid accumulation and vitellogenin biosynthesis. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **2018**, *219–220*, 28–37. [CrossRef] [PubMed]
53. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **2001**, *25*, 402–408. [CrossRef] [PubMed]
54. Lu, K.; Shu, Y.; Zhou, J.; Zhang, X.; Zhang, X.; Chen, M.; Yao, Q.; Zhou, Q.; Zhang, W. Molecular characterization and RNA interference analysis of vitellogenin receptor from *Nilaparvata lugens* (Stål). *J. Insect Physiol.* **2015**, *73*, 20–29. [CrossRef]
55. Li, Y.N.; Liu, Y.B.; Xie, X.Q.; Zhang, J.N.; Li, W.L. The modulation of trehalose metabolism by 20-hydroxyecdysone in *Antheraea pernyi* (*Lepidoptera: Saturniidae*) during its diapause termination and post-termination period. *J. Insect Sci.* **2020**, *20*, 18. [CrossRef]
56. He, Y.N.; Bu, T.L.; Liu, X.D. Tendency to mate with short-winged partner of the brown planthopper *Nilaparvata lugens* (*Hemiptera: Delphacidae*). *J. Econ. Entomol.* **2024**, *117*, 2425–2432. [CrossRef]
57. Xue, W.H.; Xu, N.; Chen, S.J.; Liu, X.Y.; Zhang, J.L.; Xu, H.J. Neofunctionalization of a second insulin receptor gene in the wing-dimorphic planthopper, *Nilaparvata lugens*. *PLoS Genet.* **2021**, *17*, e1009653. [CrossRef]
58. Xue, W.H.; Liu, Y.L.; Jiang, Y.Q.; He, S.F.; Wang, Q.Q.; Yang, Z.N.; Xu, H.J. Molecular characterization of insulin-like peptides in the brown planthopper, *Nilaparvata lugens* (*Hemiptera: Delphacidae*). *Insect Mol. Biol.* **2020**, *29*, 309–319. [CrossRef]
59. Ziegler, R.; Van Antwerpen, R. Lipid uptake by insect oocytes. *Insect Biochem. Mol. Biol.* **2006**, *36*, 264–272. [CrossRef]
60. Athenstaedt, K.; Daum, G. The life cycle of neutral lipids: Synthesis, storage and degradation. *Cell. Mol. Life Sci.* **2006**, *63*, 1355–1369. [CrossRef]
61. Arrese, E.L.; Soulages, J.L. Insect fat body: Energy, metabolism, and regulation. *Annu. Rev. Entomol.* **2010**, *55*, 207–225. [CrossRef]
62. Wang, W.; Yang, R.R.; Peng, L.Y.; Zhang, L.; Yao, Y.L.; Bao, Y.Y. Proteolytic activity of the proteasome is required for female insect reproduction. *Open Biol.* **2021**, *11*, 200251. [CrossRef]
63. Wang, W.; Ma, Y.; Yang, R.R.; Cheng, X.; Huang, H.J.; Zhang, C.X.; Bao, Y.Y. An MD-2-related lipid-recognition protein is required for insect reproduction and integument development. *Open Biol.* **2021**, *11*, 210170. [CrossRef]
64. Lu, K.; Wang, Y.; Chen, X.; Zhang, X.; Li, W.; Cheng, Y.; Li, Y.; Zhou, J.; You, K.; Song, Y.; et al. Adipokinetic hormone receptor mediates trehalose homeostasis to promote vitellogenin uptake by oocytes in *Nilaparvata lugens*. *Front. Physiol.* **2019**, *9*, 1904. [CrossRef]
65. Arai, C.; Suyama, A.; Arai, S.; Arai, N.; Yoshizane, C.; Koya-Miyata, S.; Mizote, A.; Endo, S.; Ariyasu, T.; Mitsuzumi, H.; et al. Trehalose itself plays a critical role on lipid metabolism: Trehalose increases jejunum cytoplasmic lipid droplets which negatively correlated with mesenteric adipocyte size in both HFD-fed trehalase KO and WT mice. *Nutr. Metab.* **2020**, *17*, 22. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Activation of *BmToll9-1* in Silkworm (*Bombyx mori*) Larval Midgut by *Escherichia coli* and Regulation of Growth

Jisheng Liu [1,*,†], Weijian Chen [1,†], Minchun Lai [1], Jiahua Chen [1] and Luc Swevers [2]

[1] School of Life Sciences, Guangzhou University, Guangzhou 510006, China
[2] Institute of Biosciences and Applications, National Centre for Scientific Research Demokritos, 15431 Athens, Greece; swevers@bio.demokritos.gr
* Correspondence: jisheng.liu@gzhu.edu.cn
† These authors contributed equally to this work.

Simple Summary: Toll receptors play a crucial role in insect development and innate immunity. *BmToll9-1* has been proven to positively regulate both development and immune responses in silkworms. *BmToll9-2* appears to exhibit similar functions. This article further explores the impact of bacterial infection on silkworm larvae following RNA interference of *BmToll9-1*. Interestingly, bacterial infection seems to reactivate the silenced *BmToll9-1* and nullify the phenotypic changes it caused, while also activating downstream immune pathways, with a particularly pronounced response against Gram-negative bacteria (specifically *Escherichia coli*, when compared to Gram-positive *Staphylococcus aureus*). These functional characteristics show a remarkable similarity to those of *BmToll9-2*, which were previously reported. However, a comprehensive cross-validation confirms that they are genetically distinct genes despite their functional parallels.

Abstract: Insects rely on their innate immune system to defend against pathogens, and the Toll signaling pathway plays an important role in immune regulation. Our previous studies have shown that *BmToll9-1* functions as a positive regulator in the Toll pathway. This study seeks to elucidate the role of *BmToll9-1*, as a sensor to bacterial challenge, in modulating larval development and downstream Toll signaling pathways. Silkworm larvae were subjected to infection with either Gram-negative *Escherichia coli* or Gram-positive *Staphylococcus aureus* bacteria following silencing of *BmToll9-1* by RNA interference (RNAi). This bacterial challenge triggered a compensatory re-induction of *BmToll9-1* expression, which resulted in the recovery of larval weight and size to levels observed in untreated controls. Furthermore, upon bacterial infection of *BmToll9-1*-silenced larvae, there was an up-regulation in the expression of both signaling genes in the Toll pathway and downstream effector genes, with a marked preference for Gram-negative bacteria. These results highlight the involvement of *BmToll9-1* in the Toll signaling pathway as a positive regulator, influencing silkworm development. Additionally, *BmToll9-1* and *BmToll9-2* were cross-validated to be genetically distinct genes, even though they were confirmed to be functionally analogous in the silkworm.

Keywords: *Bombyx mori*; *BmToll9-1*; RNAi; immune response; Gram-negative bacteria; *Escherichia coli*

1. Introduction

Insects have flourished on Earth for more than 480 million years and this can be explained, at least partially, by their powerful immunity. Insects primarily defend against

the invasion of exogenous pathogens through their innate immune system [1], which synthesizes and secretes antimicrobial peptides (AMPs) mediated by Toll and immune deficiency (IMD) signaling pathways [2]. Additionally, insects can also defend against exogenous pathogens through cellular immunity mediated by hemocytes (e.g., phagocytosis, nodulation, and encapsulation).

The Toll/TLR receptors belong to an ancient family of immune defense-related receptors, with genes encoding the signaling cascade components conserved across all vertebrates, spanning from lower to higher taxa [3]. Insect Toll is a type I transmembrane protein that can be divided into three parts: extracellular, transmembrane, and intracellular regions. The transmembrane region is flanked by an extracellular leucine-rich repeat (LRR) domain for ligand recognition and the cytoplasmic Toll/interleukin-1 receptor (TIR) domain that interacts with downstream molecules such as myeloid differentiation factor 88 (MyD88) [4–6].

The earliest discovery of Toll receptors was in studies of embryonic development in *Drosophila melanogaster* [7]. Subsequently, a total of nine Toll receptors were identified in *Drosophila* and they differed in structure and function [8–10]. Several studies revealed that the overexpression of *DmToll* and *DmToll9* led to a significant activation of antimicrobial peptide (AMP) genes, whereas *Dm18W* and *DmToll5* were induced to be expressed when the organism was infested with exogenous pathogens [11–15]. Also, *Dm18W* was associated with the development of *Drosophila* follicular cells and salivary glands [16–18], while *DmToll8* was associated with the regulation of neuron-specific glycosylation and the development of wing discs in the *Drosophila* embryo [19,20]. In addition to *Drosophila*, Toll receptors have been found in other insects. For example, there are 5 Toll receptors in *Apis mellifera* [21], 11 receptors in *Anopheles gambiae* [22], 9 in *Tribolium castaneum* [23], and 14 in *Bombyx mori* [24].

The domesticated silkworm, *B. mori*, which originated in China, has been cultivated for more than 5000 years ago. The silkworm not only holds significant economic importance in China but also serves as a model organism in Lepidoptera. It was the first lepidopteran insect to have its entire genome sequenced [25–27]. Moreover, the silkworm's susceptibility to pathogen infections during cultivation makes it an excellent model for studying the mechanisms of innate immunity [28,29].

Although there are 14 Toll-related genes in silkworms, current studies are focused on the Toll9 genes that encode two closely related receptors, *BmToll9-1* and *BmToll9-2*. Similarly to *DmToll9*, *BmToll9-1* and *BmToll9-2* are associated with the innate immunity of *B. mori*, as proven in previous research [6,30–35]. In previous studies, we found that *BmToll9-1* transcripts were reduced after an injection of lipopolysaccharide (LPS) or double-stranded RNA (dsRNA) into silkworm larvae [31] and that LPS also repressed AMP genes and other immune pathway-related genes in silkworm-derived Bm5 cells overexpressing the *BmToll9-1* receptor [32]. *BmToll9-1* expression was induced after infection with *Escherichia coli* and the fungus *Beauveria bassiana* [30,36], while *BmToll9-2* was induced by *E. coli* and its main cell wall component LPS, as well as *Staphylococcus aureus* and its main cell wall component peptidoglycan (PGN) [34]. In our recent studies, we found that the larvae and cocoons were smaller and lighter after the knockdown of *BmToll9-1*, and that *BmToll9-1* played a role as a positive regulator in the immune response of the Toll signaling pathway to enhance antimicrobial activity against *E. coli* [6]. Similarly to *BmToll9-1*, *BmToll9-2* showed the same effect, i.e., it stimulates the humoral immune response and antibacterial activity [35]. Remarkably, a bacterial challenge following RNAi up-regulated the expression of *BmToll9-2* and significantly mitigated the silkworm weight differences [34].

Although we have already demonstrated that *BmToll9-1* plays a vital role in innate immunity, we have yet to pinpoint the pathways involved and the mechanisms that

regulate its function. Previously, we confirmed the role of *BmToll9-1* via double-stranded RNA (dsRNA)-mediated gene silencing [6]. In this follow-up study, a bacterial challenge and RNAi were combined to complement previous research and further elucidate the mechanisms by which *BmToll9-1* functions in the innate immunity of the silkworm.

2. Materials and Methods

2.1. Insect Rearing and Bacterial Culture

The larvae of *B. mori* "P50" strain were provided by the Guangdong Academy of Agricultural Sciences, Guangzhou, China and reared on fresh mulberry leaves at 25 ± 1 °C, 75 ± 5% relative humidity, and a photoperiod of 12 L:12 D. Silkworm larvae at the 5th instar were selected for sampling. All samples were immediately transferred to RNase-free tubes and stored at −80 °C. The bacteria *E. coli* and *S. aureus*, maintained in our laboratory, were grown overnight in Luria–Bertani (LB) liquid medium at 37 °C and 200 rpm.

2.2. RNA Protocol and Bacterial Challenge to Larvae

RNAi was performed as described in our previous report [6]. In brief, the T7 RNA polymerase promoter sequence was added to the primers before PCR amplification to produce the template DNA. A T7 RiboMAX Express RNAi kit (Promega, Madison, WI, USA) was used to synthesize dsRNA. Double-stranded *BmToll9-1* (ds*BmToll9-1*) was injected on Day 1 of the 5th instar larvae, while double-stranded green fluorescent protein (ds*GFP*) was used as a negative control (N = 50 per treatment for each treatment).

Subsequently, ds*BmToll9-1*- and ds*GFP*-treated larvae were provided with mulberry leaf disks of identical size, coated with 10 µL of *E. coli* or *S. aureus* bacterial solution [34,35]. Midgut samples were collected at 6, 12, and 24 h after feeding.

2.3. RNA Extraction and cDNA Synthesis

Total RNA was isolated from frozen tissue samples using an RNAiso Plus kit (TaKaRa, Kusatsu, Japan) following the manufacturer's protocol. Genomic DNA was eliminated through DNase treatment, and first-strand complementary DNA (cDNA) was synthesized utilizing a PrimeScript RT Reagent Kit (Perfect Real Time) (TaKaRa).

2.4. Quantitative Real-Time PCR (qRT-PCR)

GoTaq qPCR Master Mix (Promega, Madison, WI, USA) was employed for qRT-PCR amplification, with midgut cDNA from 5th instar larvae as the template. The thermal cycling protocol was executed on a CFX Connect Real-Time System (Bio-Rad, Hercules, CA, USA) as follows: 95 °C for two min, 40 cycles of 95 °C for 15 s, and 57 °C for 30 s. After the thermal cycles, a dissociation step ranging from 65 °C to 95 °C was incorporated. Primer sequences for detecting *BmToll9-1*, *BmToll9-2*, signaling genes, and effector genes are detailed in Table 1. Translation initiation factor 4A (*BmTIF4A*) and translation initiation factor 3 subunit 4 (*BmTIF3s4*) were used as the reference genes [37]. Gene expression normalization was achieved through geometric averaging of the reference genes, and relative expression levels of target genes were calculated using the $2^{(-\Delta\Delta CT)}$ method [38]. All qRT-PCR reactions were conducted with three biological replicates (N = 10 larvae), each accompanied by three technical replicates.

Table 1. List of primers used in this study.

Gene	Accession Number	Primer Sequence (5′-3′)	Amplicon Size (bp)
Primers for qRT-PCR			
BmToll9-1	PP496203	F: CGCAGACCGTTGAGTACATG R: CCAGACTGTCGTACCTTGGT	149
BmToll9-2	PP716770	F: GGTTACAAGCGAACGGTAGC R: CCAAATATCCGGACTGCTGC	80
BmTIF4A	DQ443290	F: TTCGTACTGGCTCTTCTCGT R: CAAAGTTGATAGCAATTCCCT	174
BmTIF3s4	DQ443238	F: ACTTCAAGTTCAGGGCAGAT R: TTAATTGTTTTGTGGAGGCT	110
Signaling			
BmMyD88	XM_028186400	F: AACGGTCACGACTCGAACTC R: TCTGCCCAGATTCTTCATCC	105
BmTube	XM_028173146	F: GGCAGAAAGTTATGGCTTGG R: ATCCTCAAATGCTCGCTGTT	82
BmPelle	XM_028182154	F: ACATCAAGCCGGCTAACATC R: ACCGTGAGACCTTCAGATGC	117
BmCactus	XM_028180230	F: ACAGTCGTGCGTACATTTGG R: CAGCCTCTCCCTATCGTCAA	97
BmTollip-d	XM_021351983	F: GACGAGTCAGTCCCTCTTGC R: GTGGCTGGTGGAATTCGTAG	92
BmTollip-v	XM_028186930	F: TGCTACTTCTGACGGTGTGG R: AGGGCCACTTTGTGGTACTG	91
BmPellino	XM_028184930	F: AGAGTCGCTCAGCACAACAA R: CAATGTGGCTCCACACAGAT	95
BmTRAF2	XM_028172769	F: TCGCTCCTATGGGCATAACT R: CCGCATGTTGTGATTACTGG	118
BmECSIT	XM_028171307	F: ATGCCGCCTTAGCTAGAATG R: GCCTTTGGGCAGTACGTCTA	86
Effectors			
BmAttacin1 (BmAtt1)	NM_001043541	F: CAGTGAACTCGGATGGAACC R: GGCGCTGAGTACGTTCTTGT	97
BmCecropinA (BmCecA)	NM_001043997	F: CCGTCATAGGGCAAGCGAAA R: AGCAATGACTGTGGTATGTCAA	230
BmDefensin (BmDef)	AB_367525	F: GTTAAGTGCGGCGTTGACTG R: TGACAGGGAAAGTGGAAGGG	104
BmGloverin1 (BmGlv1)	AB_289654	F: GCTGGGATAGAAGCATCAGC R: ACATCAGGCCTTCTGTGACC	107
BmMoricin (BmMor)	AB_006915	F: TGTGGCAATGTCTCTGGTGT R: CTGGCGATATTGATGGCTCT	117
BmLebocin3 (BmLeb3)	NM_001126260	F: CTCGATCCAAACCGAAGGTA R: CGGCTGGTCAAGTCCAGTAT	105
BmEnbocin (BmEnb)	FJ373019	F: ACCTCGCACAACTAGTTCGG R: CCAACAGAACAAACCCACTCG	116
BmLysozyme (BmLys)	NM_001043983	F: TAACGGCTCGAAGGACTACG R: GAGGTCGGAGCACTTAACGT	103
Lysozyme-like protein (BmLLP3)	XM_012696687	F: GTTTAATCGAGCAGGGCAGC R: CACCCTTGCGACCTTCTTTG	120
Phenoloxidase inhibitor (BmPOI)	XR_001139981	F: GGATACGTGACTGGAAATGCA R: GTCATAATCCACGGGTTTGTCC	102
Prophenoloxidase 1 (BmPPO1)	AF_178462	F: AGTGGGAAGCCATTCTCCTT R: GCCAGGTTTCACTCCTTGAG	81
Nitric oxide synthase 1 (BmNOS1)	XM_012689821	F: TCATCACCACTAGCGCATCC R: CCTTGTCCGTTCTGTGTCCT	102
Primers for dsRNA synthesis			
T7-*BmToll9-1*		F: TAATACGACTCACTATAGG ACTATAGGCACAGGTCGGGT R: TAATACGACTCACTATAGG TCGTTGTCCCATTCGCTGAT	531
T7-*BmToll9-2*		F: TAATACGACTCACTATAGG TAGTATTCTCCCGGCTCTC R: TAATACGACTCACTATAGG GAAGGGTGCCTTGTGTAATC	581
T7-*GFP*		F: TAATACGACTCACTATAGG TACGGCGTGCAGTGCT R: TAATACGACTCACTATAGG TGATCGCGCTTCTCG	495

2.5. Data Analysis

Statistical analyses were conducted using a two-way ANOVA followed by Student's *t*-test using SPSS version 26.0. Data visualization and plotting were performed in GraphPad

Prism v10.0, with results presented as mean ± standard deviation based on three biological replicates. The mean values of knockdown or inhibition rates were compared to the control groups and converted to a percentage.

3. Results

3.1. Activation of BmToll9-1 by Bacterial Challenges Following RNAi of BmToll9-1

In the previous study, *BmToll9-1* could be induced by feeding on *E. coli* and *B. bassiana*, and we confirmed that it was effectively silenced in the midgut via dsRNA [6,30,36]. In this study, to further demonstrate the role of bacterial infections as a trigger for *BmToll9-1* expression, heat-inactivated *E. coli* and *S. aureus* were fed to the silkworm larvae after ds*BmToll9-1* was injected into the larvae. The results showed that the relative expression of *BmToll9-1* in the RNAi group (injected with ds*BmToll9-1*) remained significantly silenced, showing a 43% and 65% reduction after 6 h and 12 h, respectively. However, by 24 h following feeding on food that contained *E. coli*, its expression level had returned to that of the control group (injected with ds*GFP*) (Figure 1A). In contrast, when feeding on food that contained *S. aureus*, the relative expression of *BmToll9-1* in the RNAi group had normalized to the same level as the control group within 12 h (Figure 1B). Therefore, while dsRNA effectively reduced the expression of *BmToll9-1* in *B. mori*, this silencing effect could be mitigated by bacterial infection.




Figure 1. The relative expression of *BmToll9-1* in the midgut of 5th instar larvae injected with ds*BmToll9-1* and fed a diet containing heat-inactivated (**A**) *E. coli* or (**B**) *S. aureus*, compared to the negative control, i.e., larvae treated with ds*GFP*. The data were presented as means ± standard deviations of three biological replicates. The asterisks indicate significant differences from the ds*GFP* injection groups: *** $p < 0.001$, and **** $p < 0.0001$.

3.2. Bacterial Challenges Following Injection of dsBmToll9-1 Mitigated the Diminished Growth Phenotype

After the *BmToll9-1* gene was silenced, both the larvae and cocoons exhibited a smaller and lighter phenotype [6]. However, bacterial treatment annulled the effects of ds*BmToll9-1* injection over time, which manifested as reduced growth, affecting both the body weight and dimensions (Figure 2). Upon feeding on *E. coli*, the body weight of *BmToll9-1*-silenced larvae gradually recovered to match that of the control, reaching a maximum of 1.87 g (on the 4th day after the ds*BmToll9-1* injection) (Figure 2A,C). Both *BmToll9-1*-silenced larvae and the control larvae pupated into the cocoon stage around the same time (the 4th day after the ds*BmToll9-1* injection). After pupation, their body mass decreased to 0.87 g and 0.90 g, respectively, on the 11th day. Similarly, after feeding on *S. aureus*, *BmToll9-1*-silenced larvae showed comparable changes to those treated with *E. coli*, with their body weight gradually returning to the same weight as the control group, peaking on the 4th day post

RNAi (ds*BmToll9-1* injection) (Figure 2B,D). On the 11th day after the ds*BmToll9-1* injection, these larvae then entered the cocoon stage, with their body weight reducing to 0.89 g and 0.91 g, respectively.




Figure 2. Weight and phenotype changes in *B. mori* 5th instar larvae, following heat-inactivated bacteria feeding treatment after injection with ds*BmToll9-1*, compared to control larvae injected with ds*GFP*. Average weight of silkworm larvae on different days after feeding on (**A**) *E. coli* or (**C**) *S. aureus*, following *BmToll9-1* silencing. Growth phenotype of silkworm larvae on 4th day after feeding on heat-inactivated (**B**) *E. coli* or (**D**) *S. aureus*, following *BmToll9-1* silencing.

3.3. Bacterial Challenges Following RNAi of BmToll9-1 Induced Signaling Genes in Toll Pathway

Our previous study confirmed that RNAi of the *BmToll9-1* gene reduced the expression of signaling genes in the Toll pathway and downstream effector genes [6]. To further investigate the role of bacteria, bacterial treatment through the food was administered following the injection of ds*BmToll9-1* to assess the activation of signaling genes. Since both bacterial treatments after RNAi of *BmToll9-1* led to the activation of *BmToll9-1* expression at 24 h, this specific time point was chosen for subsequent validation. Compared to the control group, most of the signaling genes in the Toll pathway were significantly up-regulated in the RNAi group after the bacterial challenges. After feeding on *E. coli*, the expression levels of *BmMyD88*, *BmPelle*, *BmCactus*, *BmRel*, *BmTollip-v*, *BmPellino*, *BmTRAF2*, and *BmECSIT* were induced by 2.26-, 2.08-, 2.55-, 2.03-, 3.64-, 1.92-, 2.25-, and 2.38-fold, respectively (Figure 3A). After feeding on *S. aureus*, the immune genes *BmMyD88*, *BmPelle*, *BmCactus*, *BmRel*, *BmTollip-v*, *BmTRAF2*, and *BmECSIT* were induced by 2.55-, 2.77-, 1.81-, 2.09-, 1.74-, 1.93-, and 2.32-fold, respectively (Figure 3B).




Figure 3. The relative expression of the signaling genes in the Toll pathway in *B. mori* 5th instar larvae, following a heat-inactivated (**A**) *E. coli* or (**B**) *S. aureus* feeding treatment after injection with ds*BmToll9-1*, compared to control larvae injected with ds*GFP*. Data were presented as means ± standard deviations of three biological replicates. Asterisks indicate significant differences towards the ds*GFP* injection groups: * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

3.4. Bacterial Challenges Following RNAi of BmToll9-1 Induced Downstream Effector Genes

Similarly, bacterial treatment was administered following the RNAi of *BmToll9-1* to evaluate the activation of effector genes (Figure 4). Compared to the control group, most AMP genes were significantly up-regulated in the RNAi ds*BmToll9-1* treated group of larvae after the bacterial challenges. After feeding on *E. coli*, *BmAtt1*, *BmCecA*, *BmDef*, *BmGlv1*, *BmMor*, *BmEnb*, *BmLys*, *BmLLP3*, *BmPPO1*, and *BmNOS1* were significantly induced by 9.06-, 13.23-, 4.36-, 11.31-, 9.21-, 6.15-, 33.29-, 4.02-, 4.36-, and 2.30-fold, respectively, while *BmPOI* was also up-regulated, but not at a statistically significant level (Figure 4A). Following feeding on *S. aureus*, *BmAtt1*, *BmCecA*, *BmDef*, *BmGlv1*, *BmLeb3*, *BmEnb*, *BmLys*, *BmLLP3*, and *BmPPO1* were significantly induced by 8.41-, 2.84-, 4.47-, 16.77-, 5.19-, 2.80-, 6.22-, 2.41-, and 3.77-fold, respectively, while *BmMor*, *BmPOI*, and *BmNOS1* were not induced (Figure 4B).




Figure 4. The relative expression of the immune effector genes in *B. mori* 5th instar larvae, following a heat-inactivated (**A**) *E. coli* or (**B**) *S. aureus* feeding treatment after an injection with ds*BmToll9-1*, compared to control larvae injected with ds*GFP*. The data were presented as means ± standard deviations of three biological replicates. The asterisks indicate significant differences towards the ds*GFP* injection groups: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$.

3.5. Silencing BmToll9-1 Did Not Affect the Expression of BmToll9-2

It is worth noting that *BmToll9-2* appeared to share similar functions with *BmToll9-1*. To clarify whether these two genes affect each other's expression, a cross-validation strategy was employed. In the *BmToll9-2*-silenced larvae, the expression of *BmToll9-1* remained unchanged compared to the control group (Figure 5A). Likewise, silencing BmToll9-1 had no discernible impact on the expression of *BmToll9-2* (Figure 5B). This analysis underscores

that *BmToll9-1* and *BmToll9-2*, while functionally similar, are distinct genes with independent regulatory mechanisms.




Figure 5. Relative expression of *BmToll9-1* and *BmToll9-2* genes silenced by each other in *B. mori* 5th instar larvae. (**A**) Relative expression of *BmToll9-1* in ds*BmToll9-2*-injected larvae. (**B**) Relative expression of *BmToll9-2* in ds*BmToll9-1*-injected larvae. Data were presented as means ± standard deviations of three biological replicates. ns—not significant.

4. Discussion

Previous studies have revealed that *BmToll9-1* plays a crucial role in the innate immune pathway, exhibiting distinct responses to various microbial infections and functioning as a pattern recognition receptor (PRR) for LPS [31,33,39]. Additionally, *BmToll9-1* has been confirmed to positively mediate the innate immune pathway by regulating the expression of Toll pathway signaling genes and most AMP genes [6]. In this study, we further investigated the role of *BmToll9-1* as a sensor for bacteria in modulating larval development and downstream Toll signaling pathways.

4.1. BmToll9-1 Might Be Involved in Immune Response to Regulate Development of the Silkworm

Previously, we had shown that *BmToll9-1* is a positive regulator in the Toll pathway and the production of AMPs [6]. In this study, most Toll signaling genes were significantly up-regulated in *BmToll9-1*-silenced *B. mori* 5th instar larvae after feeding on heat-inactivated *E. coli* and *S. aureus* (Figure 3), while most AMP genes were also significantly up-regulated (Figure 4). These results suggest that *BmToll9-1* is involved in the immune response of the silkworm. At the same time, a recovery in the growth of the silkworm was observed. Following the injection of ds*BmToll9-1*, the larvae of the silkworm exhibited a reduction in both size and weight of 38% [6]. Interestingly, when *BmToll9-1*-silenced larvae were subjected to bacteria in their food, the larvae of the silkworm gradually returned to a normal weight and size (Figure 2). Combined with the above results, it was inferred that *BmToll9-1* might be involved in immune responses to regulate the development of the silkworm.

BmToll9-1 is mainly expressed in the midgut [6]. As an important immune organ in the insect immune system, the epithelial tissue of the midgut is in close contact with the microorganisms in the midgut, and plays an important role in regulating microbial community homeostasis. In this manner, it participates in the immune processes of the organism [6,40].

Silencing of the *BmToll9-1* gene may disrupt the balance of midgut microbial homeostasis, which in turn impacts food intake and digestion, ultimately affecting the growth and development of larvae. A similar observation was documented in our previous study concerning the *BmToll9-2* gene [34]. It has been demonstrated that the gut microbiota plays a role in the development of insects [41,42]. The relationship between immune activation and growth in silkworms likely involves complex interactions. Metabolic resources and

nutrient allocation might also affect the growth of the silkworms. Future experiments should incorporate mechanistic assays, such as nutrient absorption profiling, metabolic pathway analyses, and gut physiology assessments, to directly link immune up-regulation to growth outcomes.

Bacterial treatment induced the expression of *BmToll9-1*, and the body weight and size of the silkworm larvae returned to those of the control. This suggests that the reduced growth caused by the knockdown of *BmToll9-1* is reversible. Reduced larval growth due to genetic disturbances has frequently been found in past studies, e.g., involving the *BmToll9-2* and *BmPGRP-L4* genes [34,40]. It is noteworthy that these genes are associated with the regulation of the immune pathways [43].

4.2. BmToll9-1 Is Preferentially Triggered by Gram-Negative Bacteria

Our recent study demonstrated that silencing *BmToll9-1* resulted in a more pronounced reduction in antibacterial activity against *E. coli* compared to *S. aureus* [6]. Specifically, the induction of AMP genes by *E. coli* following *BmToll9-1* silencing was significantly higher than that induced by *S. aureus* (Figure 4). These findings suggest that the *BmToll9-1* gene is preferentially activated by Gram-negative bacteria, such as the tested *E. coli*. Although the expression difference of *BmToll9-1* was rapidly diminished in a shorter period in response to *S. aureus* (Figure 1), this appears to reflect a suppression of the overall expression profile rather than activation. Collectively, *BmToll9-1* exhibits a preferential response to *E. coli*.

Previous research has shown that *E. coli* induces the expression of *BmToll9-1* [30]. A similar response to *E. coli* was observed in our recent functional characterization of *BmToll9-2*, a gene phylogenetically very closest to *BmToll9-1*. In that study, *E. coli* strongly induced the expression of *BmToll9-2* in silkworm larvae [34]. RNAi-mediated silencing of *BmToll9-2* led to a greater reduction in the expression of AMP genes by *E. coli* compared to *S. aureus*, together with a more significant reduction in antibacterial activity against *E. coli* [35]. Given the functional and phylogenetic similarity between *BmToll9-1* and *BmToll9-2*, it is understandable that *BmToll9-1* is also specifically activated by Gram-negative bacteria.

4.3. BmToll9-1 and BmToll9-2 Have Separate Functions in the Silkworm Gut

Previously, we reported that the *BmToll9-1* and *BmToll9-2* genes in silkworms likely function as positive regulators of humoral immunity [6,35]. RNAi targeting of either *BmToll9-1* or *BmToll9-2* significantly reduced the expression of Toll pathway signaling genes and downstream AMP genes. Additionally, the activation of these genes exhibited a preferential response to Gram-negative bacteria [6,35]. Phylogenetic analyses further revealed that *BmToll9-1* and *BmToll9-2* are highly homologous in terms of their evolutionary relationships, suggesting they may be involved in immune regulation through functional redundancy [34].

The results of the cross-validation experiments indicated that silencing the *BmToll9-1* gene had no effect on the expression of *BmToll9-2*, and conversely, silencing the *BmToll9-2* gene did not affect the expression of *BmToll9-1* (Figure 5). This finding suggests that, regardless of the functional and phylogenetic similarities between *BmToll9-1* and *BmToll9-2*, their regulatory pathways do not affect each other's expression, i.e., they are independent. Combined with the structural differences observed in these genes [34], our current research supports the idea that *BmToll9-1* and *BmToll9-2* represent two different and independent inputs to regulate immunity and development. Double knockdown studies need to be performed to establish whether both genes can have synergic effects.

5. Conclusions

This study delved into the role of the *BmToll9-1* gene as a bacterial sensor in regulating larval development and Toll signaling pathways. Silencing of *BmToll9-1* reduced the expression of signaling genes in the Toll pathway and downstream effector genes. However, bacterial challenge following the injection of ds*BmToll9-1* reinduced the expression of *BmToll9-1*, which restored the weight and size of the silkworms. Furthermore, it was observed that *E. coli* induced a higher fold change in the expression of Toll signaling and AMP genes compared to *S. aureus*. These findings suggest that *BmToll9-1* positively regulates the Toll signaling pathway, and that it is preferentially activated by the Gram-negative bacteria *E. coli*. This study also provided preliminary evidence that *BmToll9-1* (as well as *BmToll9-2*, as previously reported) activates the Toll pathway and a nearly identical set of effector genes. These two genes do not influence each other's expression, paving the way for further investigation of their potential synergistic functions.

Author Contributions: Conceptualization, J.L.; data curation, J.L., W.C., M.L., J.C. and L.S.; funding acquisition, J.L.; investigation, J.L., W.C., M.L. and J.C.; methodology, J.L., W.C., M.L. and J.C.; project administration, J.L.; supervision, J.L.; validation, J.L., W.C., M.L. and J.C.; visualization, J.L.; writing—original draft preparation, J.L., W.C. and L.S.; writing—review and editing, J.L. and L.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Natural Science Foundation of China (31501898), Science and Technology Program of Guangzhou (202102010465), Featured Innovation Project of Universities in Guangdong Province (2019KTSCX133), Natural Science Foundation of Guangdong Province (2017A030313152), Pearl River S&T Nova Program of Guangzhou (201710010094), Guangzhou University's Exploration Experiment Construction Project (SJ202412), and Guangzhou University 2024 Education and Teaching Reform Project (JY202442).

Data Availability Statement: The raw data supporting the conclusions of this article will be made available from the corresponding author upon reasonable request.

Acknowledgments: We thank Qingrong Li for providing the silkworm culture.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Merkling, S.H.; van Rij, R.P. Beyond RNAi: Antiviral defense strategies in *Drosophila* and mosquito. *J. Insect Physiol.* **2013**, *59*, 159–170. [CrossRef] [PubMed]
2. Chintapalli, R.T.V.; Hillyer, J.F. Hemolymph circulation in insect flight appendages: Physiology of the wing heart and circulatory flow in the wings of the mosquito, *Anopheles gambiae*. *J. Exp. Biol.* **2016**, *219*, 3945–3951. [CrossRef] [PubMed]
3. Cheng, T.C.; Zhang, Y.L.; Liu, C.; Xu, P.Z.; Gao, Z.H.; Xia, Q.Y.; Xiang, Z.H. Identification and analysis of Toll-related genes in the domesticated silkworm, *Bombyx mori*. *Dev. Comp. Immunol.* **2008**, *32*, 464–475. [CrossRef] [PubMed]
4. Nüsslein-Volhard, C.; Lohs-Schardin, M.; Sander, K.; Cremer, C. A dorso-ventral shift of embryonic primordia in a new maternal-effect mutant of *Drosophila*. *Nature* **1980**, *283*, 474–476. [CrossRef]
5. Gay, N.J.; Keith, F.J. *Drosophila* Toll and IL-1 receptor. *Nature* **1991**, *351*, 355–356. [CrossRef]
6. Liu, J.; Chen, W.; Situ, J.; Li, J.; Chen, J.; Lai, M.; Huang, F.; Li, B. BmToll9-1 is a positive regulator of the immune response in the silkworm *Bombyx mori*. *Insects* **2024**, *15*, 643. [CrossRef]
7. Hashimoto, C.; Hudson, K.L.; Anderson, K.V. The Toll gene of *Drosophila*, required for dorsal-ventral embryonic polarity, appears to encode a transmembrane protein. *Cell* **1988**, *52*, 269–279. [CrossRef]
8. Valanne, S.; Wang, J.; Rämet, M. The *Drosophila* Toll Signaling Pathway. *J. Immunol.* **2011**, *186*, 649–656. [CrossRef]
9. Nonaka, S.; Kawamura, K.; Hori, A.; Salim, E.; Fukushima, K.; Nakanishi, Y.; Kuraishi, T. Characterization of Spz5 as a novel ligand for *Drosophila* Toll-1 receptor. *Biochem. Biophys. Res. Commun.* **2018**, *506*, 510–515. [CrossRef]
10. Chowdhury, M.; Li, C.F.; He, Z.; Lu, Y.; Liu, X.S.; Wang, Y.F.; Ip, Y.T.; Strand, M.R.; Yu, X.Q. Toll family members bind multiple Spatzle proteins and activate antimicrobial peptide gene expression in *Drosophila*. *J. Biol. Chem.* **2019**, *294*, 10172–10181. [CrossRef]
11. Eldon, E.; Kooyer, S.; D'Evelyn, D.; Duman, M.; Lawinger, P.; Botas, J.; Bellen, H. The *Drosophila* 18 wheeler is required for morphogenesis and has striking similarities to Toll. *Development* **1994**, *120*, 885–899. [CrossRef] [PubMed]

12. Tauszig, S.; Jouanguy, E.; Hoffmann, J.A.; Imler, J.-L. Toll-related receptors and the control of antimicrobial peptide expression in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 10520–10525. [CrossRef]
13. Ooi, J.Y.; Yagi, Y.; Hu, X.; Ip, Y.T. The *Drosophila* Toll-9 activates a constitutive antimicrobial defense. *EMBO Rep.* **2002**, *3*, 82–87. [CrossRef]
14. Bettencourt, R.; Tanji, T.; Yagi, Y.; Ip, Y.T. Toll and Toll-9 in *Drosophila* innate immune response. *J. Endotoxin Res.* **2004**, *10*, 261–268. [CrossRef]
15. Solbakken, M.H.; Tørresen, O.K.; Nederbragt, A.J.; Seppola, M.; Gregers, T.F.; Jakobsen, K.S.; Jentoft, S. Evolutionary redesign of the Atlantic cod (*Gadus morhua L.*) Toll-like receptor repertoire by gene losses and expansions. *Sci. Rep.* **2016**, *6*, 25211. [CrossRef]
16. Williams, M.J. The 18-wheeler mutation reveals complex antibacterial gene regulation in *Drosophila* host defense. *EMBO J.* **1997**, *16*, 6120–6130. [CrossRef] [PubMed]
17. Kambris, Z.; Hoffmann, J.A.; Imler, J.-L.; Capovilla, M. Tissue and stage-specific expression of the Tolls in *Drosophila* embryos. *Gene Expr. Patterns* **2002**, *2*, 311–317. [CrossRef] [PubMed]
18. Kolesnikov, T.; Beckendorf, S.K. 18 Wheeler regulates apical constriction of salivary gland cells via the Rho-GTPase-signaling pathway. *Dev. Biol.* **2007**, *307*, 53–61. [CrossRef]
19. Seppo, A.; Matani, P.; Sharrow, M.; Tiemeyer, M. Induction of neuron-specific glycosylation by Tollo/Toll-8, a *Drosophila* Toll-like receptor expressed in non-neural cells. *Development* **2003**, *130*, 1439–1448. [CrossRef]
20. Kim, S.; Chung, S.; Yoon, J.; Choi, K.W.; Yim, J. Ectopic expression of Tollo/Toll-8 antagonizes Dpp signaling and induces cell sorting in the *Drosophila* wing. *Genesis* **2006**, *44*, 541–549. [CrossRef]
21. Aronstein, K.A.; Saldivar, E. Characterization of a honey bee Toll related receptor gene Am18w and its potential involvement in antimicrobial immune defense. *Apidologie* **2005**, *36*, 3–14. [CrossRef]
22. Christophides, G.K.; Zdobnov, E.; Barillas-Mury, C.; Birney, E.; Blandin, S.; Blass, C.; Brey, P.T.; Collins, F.H.; Danielli, A.; Dimopoulos, G.; et al. Immunity-Related Genes and Gene Families in *Anopheles gambiae*. *Science* **2002**, *298*, 159–165. [CrossRef] [PubMed]
23. Zou, Z.; Evans, J.D.; Lu, Z.; Zhao, P.; Williams, M.; Sumathipala, N.; Hetru, C.; Hultmark, D.; Jiang, H. Comparative genomic analysis of the *Tribolium* immune system. *Genome Biol.* **2007**, *8*, R177. [CrossRef]
24. Tanaka, H.; Ishibashi, J.; Fujita, K.; Nakajima, Y.; Sagisaka, A.; Tomimoto, K.; Suzuki, N.; Yoshiyama, M.; Kaneko, Y.; Iwasaki, T.; et al. A genome-wide analysis of genes and gene families involved in innate immunity of *Bombyx mori*. *Insect Biochem. Mol. Biol.* **2008**, *38*, 1087–1110. [CrossRef]
25. Xia, Q.; Zhou, Z.; Lu, C.; Cheng, D.; Dai, F.; Li, B.; Zhao, P.; Zha, X.; Cheng, T.; Chai, C.; et al. A draft sequence for the genome of the domesticated silkworm (*Bombyx mori*). *Science* **2004**, *306*, 1937–1940. [PubMed]
26. Mita, K.; Kasahara, M.; Sasaki, S.; Nagayasu, Y.; Yamada, T.; Kanamori, H.; Namiki, N.; Kitagawa, M.; Yamashita, H.; Yasukochi, Y.; et al. The genome sequence of silkworm, *Bombyx mori*. *DNA Res.* **2004**, *11*, 27–35. [CrossRef]
27. Liu, J.; Yang, Q.; Yang, Y.; Lin, X. Larval RNA Interference in Silkworm Bombyx mori through Chitosan/dsRNA Nanoparticle Delivery. *J. Vis. Exp.* **2024**, *212*, e67360. [CrossRef]
28. Yang, W.; Lin, Y.; He, Y.; Li, Q.; Chen, W.; Lin, Q.; Swevers, L.; Liu, J. BmPGPR-L4 is a negative regulator of the humoral immune response in the silkworm *Bombyx mori*. *Arch. Insect Biochem. Physiol.* **2024**, *115*, e22093. [CrossRef]
29. Liu, J.; Yang, Y.; Yang, Q.; Lin, X.; Liu, Y.; Li, Z.; Swevers, L. Successful oral RNA interference efficiency in the silkworm *Bombyx mori* through nanoparticle-shielded dsRNA delivery. *J. Vis. Exp. JoVE* **2025**, *161*, 104749. [CrossRef]
30. Wu, S.; Zhang, X.; Chen, X.; Cao, P.; Beerntsen, B.T.; Ling, E. BmToll9, an Arthropod conservative Toll, is likely involved in the local gut immune response in the silkworm, *Bombyx mori*. *Dev. Comp. Immunol.* **2010**, *34*, 93–96. [CrossRef]
31. Liu, J.; Smagghe, G.; Swevers, L. Transcriptional response of BmToll9-1 and RNAi machinery genes to exogenous dsRNA in the midgut of *Bombyx mori*. *J. Insect Physiol.* **2013**, *59*, 646–654. [CrossRef] [PubMed]
32. Liu, J.; Kolliopoulou, A.; Smagghe, G.; Swevers, L. Modulation of the transcriptional response of innate immune and RNAi genes upon exposure to dsRNA and LPS in silkmoth-derived Bm5 cells overexpressing BmToll9-1 receptor. *J. Insect Physiol.* **2014**, *66*, 10–19. [CrossRef]
33. Zhang, R.; Li, X.; Zhang, J.; Li, Y.; Wang, Y.; Song, Y.; Ren, F.; Yi, H.; Deng, X.; Zhong, Y.; et al. Toll9 from *Bombyx mori* functions as a pattern recognition receptor that shares features with Toll-like receptor 4 from mammals. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2103021118. [CrossRef]
34. Liu, J.; Yang, W.; Liao, W.; Huang, Y.; Chen, W.; Bu, X.; Huang, S.; Jiang, W.; Swevers, L. Immunological function of *Bombyx* Toll9-2 in the silkworm (*Bombyx mori*) larval midgut: Activation by *Escherichia coli*/lipopolysaccharide and regulation of growth. *Arch. Insect Biochem. Physiol.* **2024**, *116*, e22130. [CrossRef] [PubMed]
35. Liu, J.; Chen, W.; Chen, S.; Li, S.; Swevers, L. Similarly to BmToll9-1, BmToll9-2 Is a Positive Regulator of the Humoral Immune Response in the Silkworm, *Bombyx mori*. *Insects* **2024**, *15*, 1005. [CrossRef]
36. Geng, T.; Huang, Y.; Hou, C.; Qin, G.; Lv, D.; Guo, X. Inductive expression patterns of genes related to Toll signaling pathway in silkworm (*Bombyx mori*) upon Beauveria bassiana infection. *J. Asia-Pac. Entomol.* **2016**, *19*, 861–868. [CrossRef]

37. Wang, G.H.; Xia, Q.Y.; Cheng, D.J.; Duan, J.; Zhao, P.; Chen, J.; Zhu, L. Reference genes identified in the silkworm *Bombyx mori* during metamorphism based on oligonucleotide microarray and confirmed by qRT-PCR. *Insect Sci.* **2008**, *15*, 405–413. [CrossRef]
38. Livak, K.J.; Schmittgen, T.D. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2-\Delta\Delta CT$ Method. *Methods* **2001**, *25*, 402–408. [CrossRef]
39. Wu, S.; Zhang, X.; He, Y.; Shuai, J.; Chen, X.; Ling, E. Expression of antimicrobial peptide genes in *Bombyx mori* gut modulated by oral bacterial infection and development. *Dev. Comp. Immunol.* **2010**, *34*, 1191–1198. [CrossRef]
40. Liang, Y.; Wang, T.; Yang, W.; Chen, Z.; Li, Q.; Swevers, L.; Liu, J. Silencing of the immune gene BmPGRP-L4 in the midgut affects the growth of silkworm (*Bombyx mori*) larvae. *Insect Mol. Biol.* **2023**, *32*, 340–351. [CrossRef]
41. Qiao, H.; Keesey, I.W.; Hansson, B.S.; Knaden, M. Gut microbiota affects development and olfactory behavior in *Drosophila melanogaster*. *J. Exp. Biol.* **2019**, *222*, jeb.192500. [CrossRef] [PubMed]
42. Guo, B.; Tang, J.; Ding, G.; Mashilingi, S.K.; Huang, J.; An, J. Gut microbiota is a potential factor in shaping phenotypic variation in larvae and adults of female bumble bees. *Front. Microbiol.* **2023**, *14*, 1117077. [CrossRef] [PubMed]
43. Bai, S.; Yao, Z.; Raza, M.F.; Cai, Z.; Zhang, H. Regulatory mechanisms of microbial homeostasis in insect gut. *Insect Sci.* **2021**, *28*, 286–301. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Article

Development of a Stage- and Species-Specific RNAi System for Molecular Insights in *Trichogramma* Wasps

Zelong Yang [1], Yan Lu [1], Zhuo Jiang [1], Xilin Jiao [1], Han Lin [1], Wanning Jiang [1], Wenmei Du [1], Xue Zhang [1], Zhao Peng [2], Junjie Zhang [1], Xiao Wang [1,*] and Ying Hu [1,*]

[1] Engineering Research Center of Natural Enemies, Institute of Biological Control, Jilin Agricultural University, Changchun 130118, China; 17731008475@163.com (Z.Y.); luyan20020624@163.com (Y.L.); jzjz8968@163.com (Z.J.); j18943100339@163.com (X.J.); 15568628611@163.com (H.L.); winniejiang1008@163.com (W.J.); wenmeid@jlau.edu.cn (W.D.); zhangxue871013@163.com (X.Z.); zhangjunjie9777@jlau.edu.cn (J.Z.)

[2] Department of Plant Pathology, College of Plant Protection, Jilin Agricultural University, Changchun 130118, China; zpeng21@jlau.edu.cn

* Correspondence: wangxiao@jlau.edu.cn (X.W.); huying@jlau.edu.cn (Y.H.); Tel.: +86-0431-8453-2834 (Y.H.)

Simple Summary: *Trichogramma* wasps, as environmentally friendly biocontrol agents, are widely employed to control lepidopteran pests. In recent years, people have attempted to study their gene function through molecular means in order to improve the control efficiency against agricultural and forestry pests of *Trichogramma* wasps. However, the progress has been severely hindered by their minute body size (<1 mm) and complex host egg-dependent parasitic behavior. In this study, we established a universal RNA interference (RNAi) technology system applicable to *T. dendrolimi* and *T. ostriniae*, targeting the *white* gene (regulating eye pigment deposition) and *laccase 2* gene (involved in cuticle sclerotization). Furthermore, we analyzed the effects of different dsRNA treatment concentrations and treatment periods on interference efficiency, overcoming the technical barriers associated with cross-species applications in conventional methods. Phenotypic analyses revealed that both non-invasive soaking and microinjection approaches achieved efficient gene silencing in *T. dendrolimi* and pupal-stage *T. ostriniae*. Notably, to avoid high mortality rates, pre-pupal *T. ostriniae* is required to use the microinjection method. This work establishes standardized protocols for functional gene studies in miniature parasitoid wasps and provides methodological foundations for developing precision biocontrol technologies through genetic engineering.

Abstract

Trichogramma wasps, egg parasitoids widely used to control lepidopteran pests, have long eluded in-depth molecular mechanistic studies due to their minute size and genetic tool scarcity. While previous RNAi efforts were restricted to *T. dendrolimi*, we developed the first cross-species RNAi system for both *T. dendrolimi* and the previously intractable *T. ostriniae*. Temporal expression profiling identified *white* and *laccase 2* as stage-specific RNAi targets, peaking during prepupal/pupal stages, which were tested across species and developmental stages using microinjection and soaking dsRNA delivery methods. Survival analysis prioritized soaking for *T. dendrolimi* prepupae/pupae, while microinjection was essential for *T. ostriniae* to bypass prepupal mortality during soaking. Concentration-dependent RNAi targeting the *white* gene achieved 85.61% transcript reduction in *T. dendrolimi* via soaking and 89.36% in *T. ostriniae* via microinjection at 2000 ng/μL, correlating with 64.06% and 32.09% white-eyed pupae, causing a significant reduction in eye pigments. For the *laccase 2* gene, soaking at 2000 ng/μL induced 88.35% transcript reduction in *T. dendrolimi* and 73.31% in *T. ostriniae*, leading to incomplete cuticle tanning and sclerotization. This

study resolves the long-standing challenge of genetic manipulation in *Trichogramma* wasps, providing a universally applicable framework to decipher parasitoid–host interactions at the molecular scale, which is useful for sustainable pest management strategies.

Keywords: *Trichogramma* wasps; RNA interference; soaking; microinjection; species-specific optimization

1. Introduction

Trichogramma wasps, serving as critical egg parasitoids in global lepidopteran pest management, have achieved remarkable ecological success through decades of field implementation [1,2]. In Northeast China, the release of *T. dendrolimi* protects over 4 million hectares of cornfields annually against the Asian corn borer (*Ostrinia furnacalis*), covering 35% of regional maize cultivation [2]. While extensive applied research has focused on ecological optimization and breeding technology enhancements, a fundamental disconnect persists between empirical biocontrol practices and mechanistic understanding due to a critical knowledge gap perpetuated by the lack of robust genetic tools in this parasitoid wasp.

RNA interference (RNAi) technology has become an important tool for genetic function research in various insects [3]. However, its efficiency is constrained by delivery methods [4], commonly dsRNA delivery methods including feeding, microinjection, and soaking; each presents distinct advantages and limitations shaped by developmental biology constraints. New delivery methods are being developed, e.g., through nanocarriers [5,6]. Feeding-based RNAi, while offering simplicity and minimal invasiveness for insects, is inherently restricted to feeding-active stages and exhibits delayed efficacy, rendering it ineffective during non-feeding phases such as embryogenesis or pupal development [7]. Microinjection, the most widely utilized method, enables precise dsRNA delivery across all ontogenetic stages from embryos to adults, achieving high phenotypic penetrance [8]. However, this technique demands specialized expertise and induces significant mechanical stress, with high mortality rates due to injection trauma [9]. Soaking protocols, involving the immersion of permeable developmental stages (e.g., larvae/pupae) in dsRNA solutions, provide a technically accessible alternative but require higher dsRNA concentrations than microinjection to attain comparable silencing efficiency [10]. This methodological triad underscores a critical trade-off in parasitoid RNAi research: balancing operational feasibility against biological precision and organismal viability.

The application of RNAi in insects exhibits profound taxonomic divergence, with Coleopteran insects demonstrating robust systemic silencing compared to Lepidoptera and Diptera [11]. Within Hymenopteran parasitoids, RNAi advancements have been predominantly confined to larger-bodied larval and pupal parasitoids. For instance, silencing the *doublesex* gene in early male larvae of *Nasonia vitripennis* using microinjection impacts the growth and differentiation of reproductive tissue [12]. In the fruit fly parasitoid *Asobara japonica*, the microinjection of dsRNA targeting the *ebony* gene successfully alters the body color [13]. However, these methods face significant technical challenges in minute egg parasitoids like *Trichogramma*, which are typically smaller than 1 mm. Most of the current RNAi research in *Trichogramma* has focused on *T. dendrolimi*. Microinjection combined with nanocarriers enables *VgR* silencing in prepupae, disrupting ovarian development [14], while pupal-stage injections achieve *Ferhch* knockdown, causing wing deformities [15]. Moreover, RNAi targeting *chitinase 10*, *v-ATPase A*, and *v-ATPase B* genes has been achieved through microinjection at the pupa stage, while feeding larvae and adults with dsRNA

failed to silence these genes, underscoring the ineffectiveness of feeding as a delivery method for RNAi in *T. dendrolimi* [16]. Despite these advances, the application of RNAi in these small parasitoids is still hindered by several limitations. Microinjection, though effective, causes mechanical damage to the tiny individuals, resulting in high mortality rates [17]. More importantly, existing technologies are not adaptable to *T. ostriniae* (body size <0.5 mm) reared on eggs of rice meal moth *Corcyra cephalonica* or Asian corn borer *O. furnacalis*, as their size differences and host specificity make traditional approaches difficult to apply across species.

This study focuses on *T. dendrolimi* and *T. ostriniae* as dual models, targeting two phenotypically clear, stage-specific genes: *white* (responsible for eye pigment deposition) and *laccase 2* (responsible for epidermal tanning). By comparing the delivery efficiency of microinjection and soaking at different developmental stages (prepupal/pupal) and between species, a standardized operational framework for cross-host and cross-species RNAi delivery is established for the first time. The innovation of this research lies in the construction of the first RNAi technology system applicable to multiple *Trichogramma* species, providing a key tool for analyzing the molecular mechanisms of host adaptability in *Trichogramma*. This technological framework not only advances genetic function research in *Trichogramma* but also offers new ideas and methods for further studying the molecular interactions between *Trichogramma* and its hosts.

2. Materials and Methods

2.1. Insect Rearing

T. dendrolimi and *T. ostriniae* were originally collected from the parasitized eggs of *O. furnacalis* in the corn fields in Yitong, Jilin Province, China (125°11′ E, 43°3′ N) in 2015 and were identified through the morphological characteristics of the male genital capsule and rDNA-ITS2 sequence analysis. *T. dendrolimi* and *T. ostriniae* were reared and maintained on the eggs of *C. cephalonica* under laboratory conditions at 25 ± 1 °C, 75 ± 5% relative humidity, and a photoperiod of 16L:8D. After being continuously reared for several generations on *C. cephalonica* eggs, *T. dendrolimi* and *T. ostriniae* colonies were reared on *Antheraea pernyi* eggs and *O. furnacalis* eggs, respectively, for subsequent RNAi experiments.

2.2. Identification and Analysis of White and Laccase 2 Genes

To identify *white* and *laccase 2* genes in *T. dendrolimi* and *T. ostriniae*, BLASTP v2.16.0 search (e-value < 1×10^{-5}, bit score > 100, identity > 70% and query coverage > 70%) was performed against the genomes of *T. dendrolimi* and *T. ostriniae*, using query sequences from *Drosophila melanogaster* (*white*: NP476787.1; *laccase 2*: NP724412.1), *Bombyx mori* (*white*: NP001037034.1; *laccase 2*: NP001103395.1), and *Apis mellifera* (*white*: NP001403446.1; *laccase 2*: XP006562317.1) retrieved from the NCBI database (http://www.ncbi.nlm.nih.gov/, accessed on 1 April 2023). Multiple sequence alignment of the *white* or *laccase 2* gene sequence of *T. dendrolimi* and *T. ostriniae* with the sequence of *D. melanogaster*, *B. mori*, and *A. mellifera* was performed to check the sequence similarity (Figures S1 and S2).

2.3. Temporal Expression Analysis of White and Laccase 2 Genes

To evaluate the temporal expression of *white* (accession number: PV568319.1) and *laccase 2* (accession number: PV568320.1) genes, samples from different developmental stages were collected for total RNA extraction. For *T. dendrolimi*, the fresh *A. pernyi* eggs were inoculated with newly emerged *T. dendrolimi* (<24 h old) for 24 h at a ratio of wasps to host eggs of 1:10. The samples were collected at different developmental stages, including egg (1st day post-parasitism), larva (3rd day post-parasitism), prepupa (5th-6th day post-parasitism), pupa (7th–11th day post-parasitism), and newly emerged adult wasps. For

T. ostriniae, fresh corn borer egg masses were parasitized with newly emerged *T. ostriniae* (<24 h old) for 24 h at a ratio of wasps to host eggs of 1:6, and samples were collected at different developmental stages, including egg (4 h after parasitism), larva (2nd day post-parasitism), prepupa (4th–5th day post-parasitism), pupa (6th–9th day post-parasitism), and newly emerged adult wasps. For each biological replicate, approximately 1 g of host egg samples containing different developmental stages of *T. dendrolimi* or *T. ostriniae* was collected. In total, three biological replicates were performed for each developmental stage. The samples were rapidly frozen in liquid nitrogen and stored at −80 °C.

Primers for RT-qPCR were designed using the IDT PrimerQuest™ Tool (https://eu.idtdna.com/pages/tools/primerquest) (accessed on 17 October 2024) [18], based on the sequences of the *white*, *laccase 2*, and *GAPDH* genes from both species. The primers are listed in Table 1 and were designed to avoid overlap with synthesized dsRNA regions. Total RNA was extracted using the EASYspin plus Cell/Tissue Total RNA Isolation Kit (Aidlab Biotechnologies Co., Ltd., Beijing, China), and cDNA synthesis was performed using the TRUEscript RT MasterMix (OneStep gDNA Removal) (Aidlab Biotechnologies Co., Ltd., Beijing, China). RT-qPCR was conducted on a qTOWER^{3}G system (Analytik Jena, Jena, TH, Germany) using 2× SGExcel FastSYBR Mixture (Sangon, Shanghai, China) under the following conditions: 95 °C for 5 min, followed by 40 cycles of 95 °C for 5 s and 60 °C for 20 s. Relative expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method, with *GAPDH* as the reference gene.

Table 1. Primers used in RT-qPCR of *white* and *laccase 2* genes.

Primer Name	Primer Sequences
white-F	TTGACACCATTGCGTGTACT
white-R	CATCCATCCAAAAGTGTTCC
laccase 2-F	CTGCTTCACCGAGAGTTCAA
laccase 2-R	AGGAAGTGGCAATGGAAGAG
GAPDH-F	AGATCAAGGCCAAGGTCAAG
GAPDH-R	AGTGGTTGTCACCGATGAAG

2.4. Preparation of dsRNAs

Primers for RNAi fragments targeting the *white* and *laccase 2* genes, with product sizes ranging from 600 to 800 bp, were designed using SnapDragon-sRNA DesignTool (https://www.flyrnai.org/cgi-bin/RNAi_find_primers.pl) (accessed on 10 October 2024.), based on regions of high sequence similarity between these genes in *T. dendrolimi* and *T. ostriniae* (Figures S9 and S10). The primers are listed in Table 2. The T7 promoter sequence (5′-TAATACGACTCACTATAGGG-3′) was added to the 5′ end of all primers. PCR amplification of the dsRNA synthesis templates was performed using Taq PCR Master Mix (Sangon, Shanghai, China), and the products were purified using the Wizard® SV Gel and PCR Clean-Up System (Promega, Madison, WI, USA) and synthesized using the T7 RNA Transcription Kit (ZSYJ, Shanghai, China).

Table 2. Primers used for dsRNA synthesis of *white* and *laccase 2* genes.

Primer Name	Primer Sequences
white-DF	TAATACGACTCACTATAGGGGCCCTTGTTCGAATGGATAA
white-DR	TAATACGACTCACTATAGGGTTGACAATCGAAATCCACGA
laccase 2-DF	TAATACGACTCACTATAGGGACGATCTCACCGTCATAGCC
laccase 2-DR	TAATACGACTCACTATAGGGGAACTGCTCGGGTGGTATGT

2.5. RNA Interference Assay via Soaking and Microinjection Delivery Method

To systematically evaluate RNA interference efficacy across developmental stages and species, we implemented complementary soaking and microinjection protocols for *T. dendrolimi* (reared in *A. pernyi* eggs) and *T. ostriniae* (reared in *O. furnacalis* eggs). For the soaking method, synchronized prepupae and pupae were dissected under stereomicroscopy (Figures S3 and S4) at species-specific post-parasitism timepoints (*T. dendrolimi*: prepupae at 6 days, pupae at 7 days; *T. ostriniae*: prepupae at 4 days, pupae at 6 days) and transferred to Parafilm-based artificial hosts with hemispherical chambers (Figure S5) [19]. Initial survival assays involved 48 h of soaking with 1 μL ddH_2O or 500 ng/μL ds*GFP*, followed by excess solution removal via filter paper. For each biological replicate, 30 individuals were placed within a single artificial host egg (3 eggs per replicate, each containing 30 wasps), with three independent replicates performed per treatment group (total n = 270 per stage/species). Adult emergence was monitored to quantify survival rates. Subsequent silencing optimization comprised two experimental designs: (1) time-course analysis—prepupae of *T. dendrolimi* (6th day post-parasitism) and *T. ostriniae* (4th day post-parasitism) soaked in the ds*white* solution (500 ng/μL and 2000 ng/μL) and pupae of *T. dendrolimi* (7th day post-parasitism) and *T. ostriniae* (6th day post-parasitism) in the ds*laccase 2* solution (500 ng/μL and 2000 ng/μL) for 12, 24, and 48 h; and (2) dose–response profiling—prepupae of *T. dendrolimi* (6th day post-parasitism) and *T. ostriniae* (4th day post-parasitism) exposed to the ds*white* solution (100, 300, 500, 1000, 1500, and 2000 ng/μL) and pupae of *T. dendrolimi* (7th day post-parasitism) and *T. ostriniae* (6th day post-parasitism) to the ds*laccase 2* solution (100, 300, 500, 1000, 1500, and 2000 ng/μL) for 48 h, with ds*GFP* (500 ng/μL) controls. After soaking for 48 h, samples were pooled (30 individuals/replicate, 3 replicates) to check the gene silencing efficiency via RT-qPCR validation.

Parallel microinjection experiments utilized the same developmental stages as soaking treatments to ensure direct methodological comparability: *T. dendrolimi* prepupae (6th day post-parasitism) and pupae (7th day) and *T. ostriniae* prepupae (4th day) and pupae (6th day). Synchronized individuals were immobilized on 2.5% agarose plates, and a PV850 microinjector (WPI) delivered 1 μL ddH_2O or 500 ng/μL ds*GFP* into the thoraco-abdominal junction at 10.0 PSI to assess baseline survival (total n = 270 per stage/species) (Figure S6). For gene silencing, prepupae were injected with the ds*white* solution (100, 300, 500, 1000, 1500, and 2000 ng/μL) and pupae were injected with the ds*laccase 2* solution (100, 300, 500, 1000, 1500, and 2000 ng/μL), with identical ds*GFP* controls. After 48 h of the treatment, samples were collected (30 individuals/replicate, 3 replicates), and the silencing efficiency was assessed via RT-qPCR. Developmental stage alignment and concentration gradients mirrored soaking protocols to enable direct methodological comparison.

2.6. Phenotypic Evaluation

Following 48 h post-treatment intervals (soaking or microinjection with ds*white* solution), eye pigmentation phenotypes were assessed using a VHX-7000 digital microscope (Keyence, Osaka, Japan) at 200× magnification. To standardize the quantification of *white* gene silencing efficacy, a four-tier chromatic classification system based on ommochrome deposition levels was established: Class I (fully pigmented, red), Class II (partial pigmentation, semi-red), Class III (low pigmentation, light red), and Class IV (unpigmented, white). All evaluations were performed on immobilized individuals under consistent illumination parameters (LED ring light, 5600K color temperature) to minimize optical artifacts.

Following 48 h post-treatment intervals (soaking or microinjection with ds*laccase 2* solution), deformations to the cuticle and body segments, as well as a lack of pigmentation, were recorded using a Keyence VHX-7000 digital microscope at 200× magnification. *Tri-*

chogramma wasps were scored as having a knockdown phenotype if we observed one or more of the traits described above.

2.7. Statistical Analysis

Statistical analyses were performed using GraphPad Prism 6.0 (GraphPad Software, San Diego, CA, USA). Two-tailed Student's *t*-test was used to analyze differences between two samples, and one-way ANOVA was performed for multiple group comparisons.

3. Results

3.1. Temporal Expression Patterns of Two Genes in Two Trichogramma Species

To determine the optimal gene silencing period, the temporal expression levels of *white* and *laccase 2* genes across developmental stages in both *Trichogramma* species were analyzed by RT-qPCR. Both genes exhibited constitutive expression throughout ontogeny but displayed distinct stage-specific amplification patterns (Figure 1A–D). Notably, the expression of *white* gene peaked during the early pupal stage (8th day post-parasitism in *T. dendrolimi*) and the prepupal stage (5th day post-parasitism in *T. ostriniae*), necessitating silencing initiation at the prepupal phase to preempt pigment deposition (Figure 1A,C). In contrast, *laccase 2* expression surged during mid-to-late pupal development (9th day post-parasitism in *T. dendrolimi*; 7th day post-parasitism in *T. ostriniae*), requiring dsRNA administration at the early pupal stage to disrupt cuticle sclerotization (Figure 1B,D). Despite host-dependent variations in developmental timelines, these expression phases remained conserved between species.




Figure 1. Spatial expression patterns and developmental timelines of *white* and *laccase 2* genes in *T. dendrolimi* (Td) and *T. ostriniae* (To). (**A**) Developmental expression profile of the *white* gene in *T. dendrolimi*

analyzed by RT-qPCR. (**B**) Developmental expression profile of the *laccase 2* gene in *T. dendrolimi* analyzed by RT-qPCR. (**C**) Developmental expression profile of the *white* gene in *T. ostriniae* analyzed by RT-qPCR. (**D**) Developmental expression profile of the *laccase 2* gene in *T. ostriniae* analyzed by RT-qPCR. (**E**) Reference developmental stages and native phenotypes of *T. dendrolimi* from the 5th to 12th day post-parasitism (from prepupa to adult). (**F**) Reference developmental stages and native phenotypes of *T. ostriniae* from the 4th to 10th day post-parasitism (from prepupa to adult). The lowercase letters represent the significant difference of the relative gene expression at different developmental stages ($p < 0.05$, Duncan).

Anatomical imaging validated these transcriptional patterns: for the *white* gene, images revealed pigmentation accumulation of eye pigments during the prepupal stage, presenting red-colored eyes (Figure 1E); for the *laccase 2* gene, the thickened cuticle and darkening of body coloration occurred during the pupal stage, aligning closely with RT-qPCR results (Figure 1F). This phenotypic-genotypic concordance confirmed that prepupal *white* targeting and early pupal *laccase 2* silencing align with their respective expression maxima, ensuring maximal interference efficacy.

3.2. Effects of Different dsRNA Delivery Methods on Survival Rates

To evaluate the viability of dsRNA delivery methods, survival rates of *T. dendrolimi* and *T. ostriniae* were analyzed following dsRNA administration via soaking or microinjection. Treatments with ddH_2O and ds*GFP* revealed no significant differences in survival (Figure 2), confirming that dsRNA itself exerted no adverse effects on wasp viability. As development progressed, survival rates for both methods improved across species. For *T. dendrolimi*, soaking demonstrated optimal viability, achieving >50% survival in prepupal and early pupal stage (5th–8th day post-parasitism) and >80% survival in the mid-late pupal stage (9th–11th day post-parasitism), whereas microinjection caused elevated mortality (>60%) at the prepupal stage (Figure 2A,B). Conversely, *T. ostriniae* exhibited species-specific adaptability: prepupae aged 4–5 days post-parasitism tolerated microinjection with survival rates ranging between 30 and 40%, while soaking remained lethal to prepupae with survival rates ranging between 5 and 15%. In pupae aged 6–7 days post-parasitism, microinjection achieved 60–80% survival compared to 30–50% for soaking (Figure 2C,D). These trends necessitated method-stage alignment: soaking was prioritized for *T. dendrolimi* prepupae and pupae, while microinjection was exclusively viable for *T. ostriniae* during the same developmental stages.




Figure 2. Effects of soaking and microinjection treatments on the survival rate in *T. dendrolimi* (Td) and *T. ostriniae* (To). (**A**) Survival rate of *T. dendrolimi* (from the 5th to 11th day post-parasitism) treated

via soaking. (**B**) Survival rate of *T. dendrolimi* (from the 5th to 11th day post-parasitism) treated via microinjection. (**C**) Survival rate of *T. ostriniae* (from the 4th to 9th day post-parasitism) treated via soaking. (**D**) Survival rate of *T. ostriniae* (from the 4th to 9th day post-parasitism) treated via microinjection.

3.3. Screening for the Optimal Timing of dsRNA Soaking

To establish the optimal soaking duration for dsRNA-mediated RNAi in *Trichogramma* species, *T. dendrolimi* and *T. ostriniae* were treated with dsRNA targeting the *white* and *laccase 2* genes at concentrations of 500 ng/μL and 2000 ng/μL. Silencing efficacy was evaluated over three durations (12 h, 24 h, and 48 h), revealing distinct species- and gene-specific responses.

In *T. dendrolimi*, *white* gene silencing exhibited a pronounced time-dependent efficacy (Figure 3A). At 12 h, significant transcript reduction (31.62%) was observed exclusively with the 2000 ng/μL treatment. Extending the soaking period to 24 h enhanced silencing efficiency, achieving reductions of 38.01% (500 ng/μL) and 51.96% (2000 ng/μL). Maximum efficacy was attained at 48 h, with transcript levels reduced by 78.49% (500 ng/μL) and 83.23% (2000 ng/μL), establishing this duration as optimal. In contrast, *T. ostriniae* prepupae exposed to *white* gene-targeting dsRNA displayed an exceptionally high mortality across all tested conditions, precluding reliable data collection. Consequently, *white* gene silencing results for this species were excluded from analysis.




Figure 3. RNAi-mediated knockdown of *white* and *laccase* 2 genes in *T. dendrolimi* (Td) and *T. ostriniae* (To) under varying dsRNA soaking durations. (**A**) The expression of the *white* gene in *T. dendrolimi* after soaking with 500 ng/μL or 2000 ng/μL ds*white*, compared to H_2O and ds*GFP* controls. (**B**) The expression of the *laccase* 2 gene in *T. dendrolimi* after soaking with 500 ng/μL or 2000 ng/μL ds*laccase 2*, compared to H_2O and ds*GFP* controls. (**C**) The expression of the *laccase* 2 gene in *T. ostriniae* after soaking with 500 ng/μL or 2000 ng/μL ds*laccase 2*, compared to H_2O and ds*GFP* controls. The lowercase letters represent the significant difference of the relative gene expression after different treatments ($p < 0.05$, Duncan).

For the *laccase 2* gene, both species exhibited time-dependent silencing efficacy. In *T. dendrolimi*, significant transcript reduction was observed at 12 h with the 2000 ng/μL treatment (50.15%), increasing to 61.66% at 24 h and peaking at 84.29% after 48 h. The 500 ng/μL treatment showed progressive efficacy, achieving reductions of 28.34% at 24 h and 53.69% at 48 h (Figure 3B). Similarly, in *T. ostriniae*, the 2000 ng/μL treatment resulted in 33.47% reduction at 12 h, escalating to 66.72% at 24 h and 74.63% at 48 h. The lower concentration (500 ng/μL) also demonstrated time-dependent improvements, with reductions of 24.06% at 24 h and 44.39% at 48 h (Figure 3C). Notably, 48 h soaking consistently achieved the highest silencing efficiency for both species and concentrations, confirming its status as the optimal duration.

3.4. Effects of dsRNA Concentration on White Gene Silencing in Trichogramma Species

To determine the optimal dsRNA concentration for RNAi-mediated *white* gene silencing, *T. dendrolimi* and *T. ostriniae* were treated with six dsRNA concentrations (100–2000 ng/μL) via soaking or microinjection. For the soaking method, significant *white* gene silencing was observed in *T. dendrolimi*, with RT-qPCR analysis revealing a dose-dependent response: 500 ng/μL reduced transcript levels by 55.91%, while 2000 ng/μL induced an 85.61% reduction in transcript levels (Figure 4A). However, soaking proved unsuitable for *T. ostriniae* prepupae (4th day post-parasitism) due to excessive mortality (>90%) (Figure 2C), necessitating microinjection for *white* gene silencing in this species. In contrast, microinjection enabled effective *white* silencing even at lower concentrations. For *T. dendrolimi*, the injection of 100 ng/μL dsRNA induced 55.76% transcript reduction, peaking at 84.29% with 2000 ng/μL (Figure 4C). Similarly, *T. ostriniae* exhibited 50.73% silencing efficiency at 100 ng/μL, rising to 89.36% at 2000 ng/μL (Figure 4E). Notably, both species displayed progressive silencing enhancement with increasing dsRNA concentrations (300–2000 ng/μL), though *T. ostriniae* required higher doses to match *T. dendrolimi*'s efficiency.




Figure 4. Gene expression and phenotypic changes following RNAi-mediated knockdown of *white* in *T. dendrolimi* (Td) and *T. ostriniae* (To) via soaking and microinjection. (**A**) Relative expression of the

white gene in *T. dendrolimi* pupae (8th day post-parasitism) following soaking with varying ds*white* concentrations. (**B**) Proportion of eye color phenotypes in *T. dendrolimi* pupae (8th day post-parasitism) after *white* knockdown via soaking. (**C**) Relative expression of the *white* gene in *T. dendrolimi* pupae (8th day post-parasitism) following microinjection with varying ds*white* concentrations. (**D**) Proportion of eye color phenotypes in *T. dendrolimi* pupae (8th day post-parasitism) after *white* knockdown via microinjection. (**E**) Relative expression of the *white* gene in *T. ostriniae* pupae (7th day post-parasitism) following microinjection with varying ds*white* concentrations. (**F**) Proportion of eye color phenotypes in *T. ostriniae* pupae (7th day post-parasitism) after *white* knockdown via microinjection. Phenotype classification in *T. dendrolimi* ((**G**), 8th day post-parasitism) and *T. ostriniae* ((**H**), 7th day post-parasitism): four-tier ommochrome deposition scale: Class I (fully pigmented, red), Class II (partial pigmentation, semi-red), Class III (low pigmentation, light red), and Class IV (unpigmented, white). The lowercase letters represent the significant difference of the relative gene expression after different treatments ($p < 0.05$, Duncan).

Knockdown of the *white* gene caused a significant reduction in eye pigments in *Trichogramma* species (Figures S7 and S8). Further phenotypic analysis revealed a chromatic gradient correlating with silencing efficiency (Figure 4G,H). Untreated controls displayed 12.65% white-eyed pupae after 48-h ds*GFP* soaking, whereas 2000 ng/μL ds*white* treatment via soaking and microinjection increased this proportion to 45.07% and 64.06%, respectively (Figure 4B,D). Notably, microinjection in *T. dendrolimi* outperformed soaking, with 100–300 ng/μL inducing a marked rise in light-red-eyed and white-eyed individuals and ≥500 ng/μL yielding predominantly white-eyed phenotypes (Figure 4B,D). Conversely, *T. ostriniae* exhibited lower phenotypic efficacy, with only 19.63–32.09% white-eyed individuals across 100–2000 ng/μL ds*white* treatments (Figure 4F).

3.5. Effects of dsRNA Concentration on Laccase 2 Gene Silencing in Trichogramma Species

The RNAi-mediated targeting of *laccase 2*, a critical regulator of cuticle sclerotization during pupal development, revealed distinct concentration–response dynamics in *T. dendrolimi* and *T. ostriniae*. Unlike pigment-related *white* gene silencing, *laccase 2* knockdown directly compromised structural integrity, with phenotypic severity escalating at higher dsRNA concentrations. Soaking protocols demonstrated robust efficacy across species: *T. dendrolimi* exhibited 88.35% transcript reduction at 2000 ng/μL (Figure 5A), while *T. ostriniae* achieved 73.31% silencing (Figure 5C), both correlating with incomplete cuticle tanning and emergence failure. Microinjection, however, highlighted interspecific divergence—low-dose treatment (100 ng/μL) yielded minimal silencing in *T. dendrolimi* (8.33%, Figure 5B) but moderate efficacy in *T. ostriniae* (49.27%, Figure 5D). At 2000 ng/μL, both methods yielded significant silencing efficiency, with 60.99% and 89.36% transcript reduction in *T. dendrolimi* and *T. ostriniae*, respectively (Figure 5B,D). Phenotypic analysis revealed that *laccase 2* knockdown disrupted cuticle tanning, a critical process initiated at the 7th day post-parasitism. The ds*GFP*-treated controls exhibited normal sclerotization, while *laccase 2*-silenced individuals displayed soft, malformed cuticles 48 h post-treatment in both *Trichogramma* species (Figure 5E,F).




Figure 5. RNAi-mediated knockdown of *laccase 2* in *T. dendrolimi* (Td) and *T. ostriniae* (To) via soaking and microinjection. (**A**) Relative expression of the *laccase 2* gene in *T. dendrolimi* pupa (9th day post-parasitism) following soaking with varying dsRNA concentrations. (**B**) Relative expression of *laccase 2* in *T. dendrolimi* pupa (9th day post-parasitism) following microinjection with varying dsRNA concentrations. (**C**) Relative expression of *laccase 2* in *T. ostriniae* pupa (7th day post-parasitism) following soaking with varying dsRNA concentrations. (**D**) Relative expression of *laccase 2* in *T. ostriniae* pupa (7th day post-parasitism) following microinjection with varying dsRNA concentrations. (**E**) Phenotypic comparison of *T. dendrolimi* pupae (9th–11th day post-parasitism) treated with ds*GFP* (left) and ds*laccase 2* (right). (**F**) Phenotypic comparison of *T. ostriniae* pupae (7th–9th day post-parasitism) treated with ds*GFP* (left) and ds*laccase 2* (right). The lowercase letters represent the significant difference of the relative gene expression after different treatments ($p < 0.05$, Duncan).

4. Discussion

The establishment of RNAi systems in *Trichogramma* wasps has long been constrained by their minute size and host-rearing dependencies. While prior studies focused exclusively on *T. dendrolimi*, our work bridges a critical gap by successfully implementing RNAi in *T.*

ostriniae, a superior biocontrol agent used to control *O. furnacalis* traditionally hindered by its reliance on *C. cephalonica* eggs and sub-0.5 mm body size. By tailoring delivery methods to species-specific physiological constraints, we demonstrate that soaking remained effective for pupal-stage gene silencing in both *T. dendrolimi* and *T. ostriniae*, whereas microinjection enables robust gene silencing in *T. ostriniae* prepupae—a breakthrough that expands functional genomics tools to non-model *Trichogramma* species. This advancement not only resolves the historical neglect of *T. ostriniae* in molecular studies but also provides a framework for adapting RNAi protocols to microhymenopteran parasitoids with diverse host preferences.

The selection of dsRNA delivery methods in *Trichogramma* wasps hinges on balancing technical feasibility, species-specific physiological constraints, and silencing efficiency. Microinjection, the predominant RNAi method in parasitoid wasps [20,21], offers precise tissue-targeted delivery with controlled dsRNA concentration but requires specialized equipment (e.g., microprocessor-controlled injectors) and incurs operational complexity compared to non-invasive methods. While studies report dsRNA injection amounts ranging widely from 500 ng/μL to 6 μg/μL in other parasitoids [22,23], our results demonstrate that *Trichogramma* species achieve significant gene silencing at lower concentrations. For instance, the microinjection of 100 ng/μL dsRNA induced 50.8% transcript reduction in the *laccase 2* gene in *T. ostriniae* and 57.46% and 49% transcript reduction in the *white* gene in *T. dendrolimi* and *T. ostriniae*, respectively, highlighting exceptional dsRNA bioavailability in these microhymenopterans. Intriguingly, increasing dsRNA concentrations to 200–1500 ng/μL were not effective in enhancing silencing efficacy, consistent with the hypothesis of a saturation threshold in the RNAi machinery [10], which underscores the efficiency of low-dose microinjection in *Trichogramma*.

In contrast, the non-invasive soaking method offers operational simplicity by facilitating passive dsRNA uptake through cuticular or spiracular pathways during the pupal stage [24,25], eliminating the mechanical trauma associated with microinjection. However, its efficacy is strongly species-dependent due to variations in osmotic tolerance. For *T. dendrolimi*, which is industrially reared in *A. pernyi* eggs, soaking at 2000 ng/μL achieved >80% survival and silencing efficiency for both *white* and *laccase 2* genes, aligning well with large-scale experimental requirements. Conversely, the smaller-bodied *T. ostriniae* (0.3 mm) exhibited severe sensitivity to this passive absorption mechanism, with prepupal survival dropping below 20% under identical conditions, likely due to osmotic stress. Notably, at the elevated concentration of 2000 ng/μL, soaking matched microinjection in silencing efficacy across both species, demonstrating its potential as a viable alternative where species-specific physiological barriers are absent.

Temporal dynamics further modulate RNAi efficacy during soaking, necessitating a balance between silencing optimization and developmental viability. In *T. dendrolimi*, soaking with 2000 ng/μL ds*white* for 24 h reduced *white* and *laccase 2* transcripts by 51.95% and 61.66%, respectively. Extending the soaking period to 48 h further enhanced *white* silencing to 83.23%, though *laccase 2* efficiency slightly declined to 84.29%, accompanied by an increase in mortality. This divergence suggests gene-specific susceptibility to prolonged dsRNA exposure, potentially linked to differential expression kinetics or pathway saturation. Similar trends were observed in *Microplitis mediator*, where *MmedOR49* silencing efficiency rose from 30% (24 h) to 50% (48 h) under equivalent conditions [20], underscoring the universality of time-dependent RNAi efficacy in parasitoids. The progressive improvement in silencing with extended soaking likely stems from cumulative dsRNA absorption through cuticular or spiracular pathways, enabling the sustained saturation of RNAi machinery components. This contrasts starkly with microinjection, where a single

bolus dose degrades or dilutes over time, requiring precise alignment with transcriptional peaks [3].

RNAi efficiency also exhibits substantial variability among genes within the same insect species, even when uniform delivery methods are employed [11]. For instance, in *Aphidius ervi*, dsRNA targeting *AeSPH1* induced a significant reduction in its transcript levels, whereas dsRNA targeting *AeSPN1* failed to alter *AeSPN1* expression under identical experimental conditions [26]. Similarly, in *T. dendrolimi*, the silencing efficacy of the *white* gene consistently surpassed that of the *laccase 2* gene across both soaking and injection delivery methods. These observations underscore the intrinsic gene-specific sensitivity to RNAi, which may arise from factors such as transcript abundance, dsRNA accessibility, or sequence-specific secondary structures [27,28]. This phenomenon is not unique to hymenopterans. In coleopterans, broad-scale screens of dsRNAs targeting 290 genes in *Diabrotica virgifera virgifera* revealed marked differences in efficacy, with the LC_{50} values of the 17 most effective dsRNAs varying by nearly 100-fold [29]. Similarly, in the Colorado potato beetle (*Leptinotarsa decemlineata*), transcript suppression levels ranged from approximately 60% to 93% depending on the target gene [30]. Such variability emphasizes the necessity of empirical gene screening and mechanistic studies to optimize RNAi applications [3,11,31].

The successful implementation of RNAi in both *T. dendrolimi* and *T. ostriniae* demonstrates that the optimized framework in this study holds significant potential for broader application across the *Trichogramma* genus. Crucially, the observed interspecies divergence between these species underscores that protocol transferability fundamentally depends on accommodating species-specific physiological constraints. The comparative analysis reveals that optimal dsRNA delivery method selection is dictated by developmental tolerance, where robust species like *T. dendrolimi* achieve efficient silencing via non-invasive soaking during pupal stages, while *T. ostriniae* prepupae necessitate microinjection to maintain viability. Beyond delivery routes, key operational parameters require species-tailored optimization; although 2000 ng/μL dsRNA proved effective here, optimal concentrations for silencing efficiency and minimal toxicity may vary across species targets or developmental stages and must be empirically determined through dose–response testing. Similarly, treatment parameters including soaking duration and microinjection volume demand calibration based on biological factors like body size and cuticle permeability. This dual emphasis on physiological adaptation and parametric calibration establishes a replicable blueprint for extending RNAi methodologies to previously intractable *Trichogramma* species.

Our study advances RNAi methodology in *Trichogramma* wasps by overcoming species-specific limitations and elucidating the intricate interplay between delivery methods, the temporal dynamics of gene silencing, and gene-specific susceptibility to RNAi. Specifically, by systematically addressing the technical bottlenecks in dsRNA delivery that have long hindered functional genomics research in minute-bodied parasitoids, this work establishes a robust framework for precise genetic manipulation. These advancements not only establish an efficient RNAi system tailored to *Trichogramma* biology but also directly enable targeted genetic interventions to enhance traits critical for biocontrol efficacy, providing a crucial technical support for pest control and management.

5. Conclusions

In this study, through different dsRNA delivery methods and phenotype analysis of two genes in *Trichogramma* wasps, we found that both methods are applicable for *T. dendrolimi*, with a 48 h soaking treatment achieving approximately 85% gene silencing efficiency. For *T. ostriniae,* with a smaller body size during the prepupal stage, the microinjection method was more suitable, while the soaking method was more convenient during the pupal stage. The microinjection method showed a significant silencing efficiency at

a dsRNA concentration of 500 ng/μL, with the strongest silencing efficiency observed at a concentration of 2000 ng/μL. When using 1000–2000 ng/μL dsRNA for the soaking methods, it could exhibit the same silencing effect as microinjection methods. This optimized system transcends the prior limitation of RNAi applications confined to *T. dendrolimi* within *Trichogramma* species, establishing a robust platform for systematically elucidating molecular mechanisms governing parasitic behaviors in these wasps.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/insects16070673/s1, Figure S1. Sequence alignment analysis of *white* genes from *T. dendrolimi* and other species; Figure S2. Sequence alignment analysis of *laccase 2* genes from *T. dendrolimi* and other species; Figure S3. A detailed dissection procedure of *T. dendrolimi* pupae from *Antheraea pernyi* eggs; Figure S4. A detailed dissection procedure of *T. ostriniae* pupae from *Ostrinia furnacalis* eggs. Figure S5. Soaking-mediated RNAi in *Trichogramma* wasps. Figure S6. Microinjection-mediated RNAi in *Trichogramma* wasps; Figure S7. Phenotypes of *T. dendrolimi* before and after *white* gene silencing; Figure S8. Phenotypes of *T. ostriniae* before and after *white* gene silencing. Figure S9: Comparison of *white* gene silencing fragments of *T. dendrolimi* and *T. ostriniae*. Figure S10: Comparison of *laccase 2* gene silencing fragments of *T. dendrolimi* and *T. ostriniae.*

Author Contributions: Conceptualization, Z.Y., J.Z., X.W. and Y.H.; methodology, Z.Y., Y.L., Z.J., X.J., H.L., W.J., X.W. and Y.H.; validation, Z.Y., Z.P. and Y.H.; formal analysis, Z.Y., X.W. and Y.H.; resources, J.Z., W.D., X.Z. and Y.H.; investigation, Z.Y., X.W. and Y.H.; writing—original draft preparation, Z.Y., Y.L., Z.J., X.J., H.L., W.J., W.D., Z.P., X.W. and Y.H.; writing—review and editing, Z.Y., X.Z., Z.P., X.W. and Y.H.; supervision, X.W. and Y.H.; project administration, X.W. and Y.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the National Key R&D Program of China (2023YFE0123000), the National Natural Science Foundation of China (32202386), the Jilin Provincial Science-Technology Foundation for Middle-aged and Young Scientists (20250601060RC), and the Jilin Provincial Science and Technology Development Program (20230502003GH).

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding authors.

Acknowledgments: We appreciate the support from Jilin Province Technology Research Center of Biological Control Engineering and Jilin Provincial International Cooperation Key Laboratory for the Biological Control of Agricultural Pests during the entire process of this research.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Cherif, A.; Mansour, R.; Grissa-Lebdi, K. The egg parasitoids *Trichogramma*: From laboratory mass rearing to biological control of lepidopteran pests. *Biocontrol. Sci. Technol.* **2021**, *31*, 661–693. [CrossRef]
2. Zang, L.S.; Wang, S.; Zhang, F.; Desneux, N. Biological Control with *Trichogramma* in China: History, Present Status, and Perspectives. *Annu. Rev. Entomol.* **2021**, *66*, 463–484. [CrossRef] [PubMed]
3. Zhu, K.Y.; Palli, S.R. Mechanisms, Applications, and Challenges of Insect RNA Interference. *Annu. Rev. Entomol.* **2020**, *65*, 293–311. [CrossRef] [PubMed]
4. Lu, Y.; Deng, X.; Zhu, Q.; Wu, D.; Zhong, J.; Wen, L.; Yu, X. The dsRNA Delivery, Targeting and Application in Pest Control. *Agronomy* **2023**, *13*, 714. [CrossRef]
5. Zhongzheng, M.; Cao, L.-J.; Chen, J.-C.; Chen, W.-B.; Shen, X.; Song, W.; Yang, F.; Wei, S.-J. A nanocarrier-mediated dsRNA oral delivery enhances RNAi efficiency in thrips. *Entomol. Gen.* **2024**, *44*, 601–611.
6. Chao, Z.; Zhongzheng, M.; Yunhui, Z.; Yan, S.; Shen, J. Establishment of star polycation-based RNA interference system in all developmental stages of fall armyworm Spodoptera frugiperda. *Entomol. Gen.* **2023**, *43*, 127–137. [CrossRef]
7. Kunte, N.; Mcgraw, E.; Bell, S.; Held, D.; Avila, L.A. Prospects, challenges and current status of RNAi through insect feeding. *Pest Manag. Sci.* **2020**, *76*, 26–41. [CrossRef]
8. Han, J.; Rotenberg, D. Microinjection-enabled gene silencing in first instar larvae of western flower thrips, Frankliniella occidentalis, reveals vital genes for larval survival. *Insect Sci.* **2024**, *Online ahead of print*. [CrossRef]

9. Yu, N.; Christiaens, O.; Liu, J.; Niu, J.; Cappelle, K.; Caccia, S.; Huvenne, H.; Smagghe, G. Delivery of dsRNA for RNAi in insects: An overview and future directions. *Insect Sci.* **2013**, *20*, 4–14. [CrossRef]
10. Joga, M.R.; Zotti, M.J.; Smagghe, G.; Christiaens, O. RNAi Efficiency, Systemic Properties, and Novel Delivery Methods for Pest Insect Control: What We Know So Far. *Front. Physiol.* **2016**, *7*, 553. [CrossRef]
11. Silver, K.; Cooper, A.M.; Zhu, K.Y. Strategies for Enhancing the Efficiency of RNA Interference in Insects. *Pest Manag. Sci.* **2021**, *77*, 2645–2658. [CrossRef]
12. Wang, Y.; Rensink, A.H.; Fricke, U.; Riddle, M.C.; Trent, C.; Van De Zande, L.; Verhulst, E.C. Doublesex regulates male-specific differentiation during distinct developmental time windows in a parasitoid wasp. *Insect Biochem. Mol. Biol.* **2022**, *142*, 103724. [CrossRef]
13. Kamiyama, T.; Shimada-Niwa, Y.; Tanaka, H.; Katayama, M.; Kuwabara, T.; Mori, H.; Kunihisa, A.; Itoh, T.; Toyoda, A.; Niwa, R. Whole-genome sequencing analysis and protocol for RNA interference of the endoparasitoid wasp *Asobara japonica*. *DNA Res.* **2022**, *29*, dsac019. [CrossRef] [PubMed]
14. Wang, C.X.; Bao, H.Q.; Yan, Z.C.; Wang, J.; Wang, S.; Li, Y.X. Knockdown of vitellogenin receptor based on minute insect RNA interference methods affects the initial mature egg load in the pest natural enemy *Trichogramma dendrolimi*. *Insect Sci.* **2025**, *32*, 487–500. [CrossRef] [PubMed]
15. Zhang, L.X.; Dong, Q.J.; Yang, S.; Che, W.N.; Zhang, L.S.; Zhou, J.C.; Dong, H. RNA Interference in the Egg Parasitoid, *Trichogramma dendrolimi* Matsumura. *J. Vis. Exp.* **2023**, e66250. [CrossRef]
16. Yan, Z.C.; Li, F.Y.; Wang, A.K.; Wang, C.X.; Wang, H.Y.; Yu, Z.Q.; Wang, K.P.; Wang, Y.H.; Luo, Y.Y.; Li, Y.X. Assessing RNAi feasibility and susceptibility to environmental RNAi in *Trichogramma dendrolimi*. *Entomol. Gen.* **2024**, *44*, 663–672. [CrossRef]
17. Zhang, M.Q.; Gong, L.L.; Zhao, Y.Q.; Ma, Y.F.; Long, G.J.; Guo, H.; Liu, X.Z.; Hull, J.J.; Dewer, Y.; Yang, C.; et al. Efficient DIPA-CRISPR-mediated knockout of an eye pigment gene in the white-backed planthopper, *Sogatella furcifera*. *Insect Sci.* **2024**, *31*, 1015–1025. [CrossRef]
18. Sandybayev, N.; Strochkov, V.; Beloussov, V.; Orkara, S.; Kydyrmanov, A.; Khan, Y.; Batanova, Z.; Kassenov, M. Evaluation of a novel real-time polymerase chain reaction assay for identifying H3 equine influenza virus in Kazakhstan. *Vet. World* **2023**, *16*, 1682–1689. [CrossRef]
19. Jiao, X.; Jiang, Z.; Yang, Z.; Jiang, W.; Wang, X.; Lu, Y.; Lin, H.; Ruan, C.; Peng, Z.; Zhang, J.; et al. Insights into the composition and evolution of venom proteins collected from artificial host eggs in *Trichogramma dendrolimi*. *Entomol. Gen.* **2024**, *44*, 1557–1568. [CrossRef]
20. Shan, S.; Song, X.; Khashaveh, A.; Wang, S.N.; Lu, Z.Y.; Hussain Dhiloo, K.; Li, R.J.; Zhang, Y.J. A female-biased odorant receptor tuned to the lepidopteran sex pheromone in parasitoid *Microplitis mediator* guiding habitat of host insects. *J. Adv. Res.* **2023**, *43*, 1–12. [CrossRef]
21. Wang, J.; Yan, Z.; Xiao, S.; Wang, B.; Fang, Q.; Schlenke, T.; Ye, G. Characterization of a cell death-inducing endonuclease-like venom protein from the parasitoid wasp *Pteromalus puparum* (Hymenoptera: Pteromalidae). *Pest Manag. Sci.* **2021**, *77*, 224–233. [CrossRef] [PubMed]
22. Huang, G.; Liu, Z.; Gu, S.; Zhang, B.; Sun, J. Identification and functional analysis of odorant-binding proteins of the parasitoid wasp *Scleroderma guani* reveal a chemosensory synergistic evolution with the host *Monochamus alternatus*. *Int. J. Biol. Macromol.* **2023**, *249*, 126088. [CrossRef]
23. Yan, M.W.; Xing, X.R.; Wu, F.A.; Wang, J.; Sheng, S. UDP-glycosyltransferases contribute to the tolerance of parasitoid wasps towards insecticides. *Pestic. Biochem. Physiol.* **2021**, *179*, 104967. [CrossRef]
24. Wang, Y.; Zhang, H.; Li, H.; Miao, X. Second-generation sequencing supply an effective way to screen RNAi targets in large scale for potential application in pest insect control. *PLoS ONE* **2011**, *6*, e18644. [CrossRef] [PubMed]
25. Whyard, S.; Singh, A.D.; Wong, S. Ingested double-stranded RNAs can act as species-specific insecticides. *Insect Biochem. Mol. Biol.* **2009**, *39*, 824–832. [CrossRef] [PubMed]
26. Zhao, J.; Wang, Z.W.; Shen, G.; Hu, D.; Zhong, Y.; Ye, C.; Wang, J.J. Regulation of melanization in aphids by parasitoid wasp venom proteins enhances mummification. *Pest Manag. Sci.* **2025**, *81*, 1017–1025. [CrossRef]
27. Chen, J.; Peng, Y.; Zhang, H.; Wang, K.; Tang, Y.; Gao, J.; Zhao, C.; Zhu, G.; Palli, S.R.; Han, Z. Transcript level is a key factor affecting RNAi efficiency. *Pestic. Biochem. Physiol.* **2021**, *176*, 104872. [CrossRef]
28. Vermeulen, A.; Behlen, L.; Reynolds, A.; Wolfson, A.; Marshall, W.S.; Karpilow, J.; Khvorova, A. The contributions of dsRNA structure to Dicer specificity and efficiency. *RNA* **2005**, *11*, 674–682. [CrossRef]
29. Baum, J.A.; Bogaert, T.; Clinton, W.; Heck, G.R.; Feldmann, P.; Ilagan, O.; Johnson, S.; Plaetinck, G.; Munyikwa, T.; Pleau, M.; et al. Control of coleopteran insect pests through RNA interference. *Nat. Biotechnol.* **2007**, *25*, 1322–1326. [CrossRef]

30. Zhu, F.; Xu, J.; Palli, R.; Ferguson, J.; Palli, S.R. Ingested RNA interference for managing the populations of the Colorado potato beetle, *Leptinotarsa decemlineata*. *Pest Manag. Sci.* **2011**, *67*, 175–182. [CrossRef]
31. Lu, J.; Shen, J. Target genes for RNAi in pest control: A comprehensive overview. *Entomol. Gen.* **2024**, *44*, 95–114. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Review

Odorant Binding Proteins in *Tribolium castaneum*: Functional Diversity and Emerging Applications

Lei Wang [1,2], Yujie Lu [1,2,*] and Zongpei Zhao [1,2,*]

[1] School of Grain Science and Technology, Jiangsu University of Science and Technology, Zhenjiang 212100, China; wanglei-best@just.edu.cn
[2] Jiangsu Provincial Engineering Research Center of Grain Bioprocessing, Zhenjiang 212100, China
* Correspondence: luyjlyj71@just.edu.cn (Y.L.); zongpeizhao@just.edu.cn (Z.Z.); Tel.: +86-511-85626711 (Y.L. & Z.Z.)

Simple Summary: Stored product loss threatens food security worldwide. This review examines odorant binding proteins in the red flour beetle, a major pest and genetic model. We show how these small proteins, once viewed only as smell carriers, also act in chemical defense, detoxification, immunity, and reproduction. Evidence from tissue- and stage-specific expression maps, gene-silencing experiments, and ligand-binding assays supports roles for specific family members—C11, C12, C17, 7G, and the 9A/9B pair—that respond to diverse harmful chemicals. Comparative analyses reveal lineage-specific expansions and adaptive change, with rapid diversification in a subgroup lacking two conserved bonding sites. Translational opportunities include gene-silencing baits, behavior-modifying chemicals that disrupt binding, and sensors that harness protein–odor interactions. Key barriers remain, notably the shortage of high-resolution three-dimensional structures and functional overlap among related proteins. Priorities include stable genome editing to create single and multiple knockouts, single-cell maps to assign functions, and structural methods to visualize binding and release. Integrating these advances into system-level models can deliver precise, environmentally responsible protection of stored foods while reducing broad insecticide use.

Abstract

Tribolium castaneum, the red flour beetle, is both a major pest of stored products and a valuable genetic model. Odorant Binding Proteins (OBPs), traditionally associated with olfaction, are now recognized as multifunctional, contributing to detoxification, immunity, and reproduction. This review synthesizes recent advances in the molecular structure, gene expression, and functional characterization of *T. castaneum* OBPs—particularly TcOBPC11, TcOBPC12, TcOBPC17, and TcOBP7G. Experimental evidence, including RNA interference, ligand-binding assays, and expression profiling, supports their role in defense against xenobiotics. Comparative genomic analyses reveal lineage-specific expansions and adaptive evolution, especially in Minus-C OBPs. Translational applications include RNA interference (RNAi)-based pest control, OBP-targeting repellents, and biosensors. Outstanding challenges remain, including structural resolution and functional redundancy. Future research integrating CRISPR, single-cell transcriptomics, and structural biology will be critical to decode OBP regulatory networks and leverage their potential in pest management and biotechnology.

Keywords: *Tribolium castaneum*; odorant binding proteins (OBPs); molecular structure; gene expression; pest control; comparative biology; research trends; detoxification; signal transduction; structural biology

1. Introduction

1.1. Tribolium castaneum: A Global Pest and Powerful Genetic Model

Tribolium castaneum (Herbst), the red flour beetle, holds a dual identity in the scientific world. On one hand, it is a cosmopolitan and economically significant pest of stored agricultural products, including grains, flour, cereals, and processed foods. Its infestations lead to substantial losses through direct consumption and contamination with metabolic byproducts, such as quinones, which render food unpalatable and potentially harmful [1,2]. On the other hand, *T. castaneum* has emerged as a premier non-Drosophila model organism for a wide spectrum of biological research [3]. Its utility is underpinned by several advantageous characteristics: it is easy and inexpensive to rear in the laboratory, has a short generation time of approximately one month, and, most critically, possesses a high-quality, fully sequenced genome [4]. The Tcas5.2 reference genome assembly (GenBank: GCF_000002335.3), along with recent annotation enhancements, provides a robust framework for genetic and genomic investigations [5].

A particularly powerful feature of *T. castaneum* is its remarkable susceptibility to systemic RNA interference (RNAi) [4]. This allows for efficient and targeted gene knockdown, making it an exemplary system for functional genomics to dissect gene function and identify potential targets for novel pest control strategies [6]. These attributes have facilitated in-depth studies into fundamental insect biology, including development, insecticide resistance, and, central to this review, the mechanisms of chemosensation [2].

1.2. Odorant Binding Proteins: Canonical Roles in Insect Chemoreception

The ability of insects to navigate their environment, find food, locate mates, and avoid threats is profoundly dependent on their sophisticated chemosensory system [7]. At the forefront of this system are the Odorant Binding Proteins (OBPs), a diverse family of small (typically 10–30 kDa), soluble, and highly abundant proteins found in the aqueous sensillar lymph of chemosensory organs, primarily the antennae and mouthparts [8]. The canonical model of OBP function posits that they perform the crucial initial step in olfaction: capturing hydrophobic semiochemicals—such as host plant volatiles, pheromones, and kairomones—and solubilizing them for transport across the hydrophilic lymph. This OBP-odorant complex is then delivered to membrane-bound olfactory receptors (ORs) situated on the dendrites of olfactory sensory neurons, triggering a signal transduction cascade that results in an electrical signal [9].

Insect OBPs are structurally classified into several subfamilies based on the number and pattern of their conserved cysteine residues, which form intramolecular disulfide bridges essential for protein stability. The main subfamilies include “Classic” OBPs, which possess six conserved cysteines; “Plus-C” OBPs, which have additional cysteines and a conserved proline residue; and “Minus-C” OBPs, which are characterized by the loss of the second and fifth ancestral cysteines (C2 and C5) [10]. While this classification provides a structural framework, the functional landscape of OBPs is proving to be far more complex than originally envisioned. A central theme of this review is the paradigm shift from viewing OBPs solely as passive odorant carriers to recognizing their active roles in a multitude of physiological processes, including detoxification of xenobiotics and innate immunity, functions that are particularly relevant for a generalist pest like *T. castaneum* (Figure 1) [11].




Figure 1. *Tribolium castaneum* as a Dual Model and the Expanded Functions of OBPs Beyond Olfaction.

2. Molecular Architecture and Functional Dynamics of *T. castaneum* OBPs

2.1. Structural Hallmarks and Subfamily Diversity

The fundamental architecture of insect OBPs, including those in *T. castaneum*, is a compact, globular fold predominantly composed of six to seven α-helices [12]. This structure is stabilized by a network of intramolecular disulfide bridges, which are the basis for OBP classification. In Classic OBPs, three disulfide bonds (C1-C3, C2-C5, C4-C6) create a stable scaffold that cradles a hydrophobic internal binding pocket, designed to sequester lipophilic odorants. Variations in this pattern, such as the absence of the C2-C5 bridge in Minus-C OBPs, can lead to increased structural flexibility and altered ligand-binding characteristics [13].

While experimentally determined three-dimensional structures for most *T. castaneum* OBPs (TcOBPs) remain elusive, this gap has spurred the proficient use of computational biology to generate valuable structural hypotheses [14]. Homology modeling, which builds a structural model of a target protein based on the known structure of a related protein, has been pivotal. For instance, the homology model of TcOBPC12 references available structures/models of related beetle OBPs and indicates a binding pocket composed of both hydrophobic and polar residues, consistent with interactions with phenolic ligands such as eugenol [15]. Similarly, modeling of the Minus-C OBP TcOBP7G provided insights into how the absence of a disulfide bond might alter its binding pocket geometry and ligand preferences [16]. The recent advent of highly accurate protein structure prediction tools, such as AlphaFold 2, is set to revolutionize this field, providing high-confidence models that can accelerate structure-based functional studies and rational ligand design [17].

However, it is important to emphasize that these TcOBP models still represent structural hypotheses rather than experimentally validated structures. Sequence identity with available OBP templates is moderate, and the most variable regions—loops and termini that often shape the entrance and fine geometry of the binding pocket—are also the least reliable in homology models. As a consequence, detailed predictions about side-chain

orientations and specific ligand contacts should be interpreted with caution and ideally confirmed by site-directed mutagenesis and high-resolution structural methods.

Beyond static structures, molecular dynamics simulations have highlighted the importance of protein flexibility. These studies suggest that the conformational dynamics of loops connecting the α-helices, as well as a flexible C-terminal region, are critical for accommodating a diverse range of ligands and for the process of ligand entry and release. This inherent flexibility likely underpins the ability of a limited number of OBPs to interact with a vast chemical landscape [18]. These structural themes and subfamily relationships, together with a schematic view of perireceptor events, are summarized in Figure 2.




Figure 2. Overview of OBP Diversity, Structural Features, and Functional Mechanism. (**A**) Phylogenetic diversity showing the main OBP subfamilies. Classic OBPs typically have 6 conserved cysteines. Minus-C OBPs (expanded in *Tribolium*) are characterized by the loss of the conserved C2 and C5 cysteines. Plus-C OBPs carry additional cysteines. (**B**) Schematic comparison of Classic vs. Minus-C OBP structures. Classic OBPs possess 6 cysteines forming 3 rigid disulfide bonds (red lines). Minus-C OBPs have lost the ancestral C2 and C5 cysteines, and therefore lack the C2–C5 disulfide bond, resulting in a potentially more flexible structure (dashed outline). (**C**) Schematic model of perireceptor events. Hydrophobic odorants (orange stars) bind to OBPs at neutral pH and are released to the Olfactory Receptor (OR) complex in the acidic environment near the dendritic membrane.

2.2. Ligand Binding: Specificity, Affinity, and Release Mechanisms

A major focus of recent research has been the functional characterization of TcOBP ligand-binding profiles, which has been instrumental in uncovering their diverse physiological roles. These studies are predominantly conducted using in vitro fluorescence competitive binding assays, where a fluorescent probe, such as N-phenyl-1-naphthylamine (1-NPN), binds to the OBP's hydrophobic pocket. The ability of a test ligand to displace this probe is measured, allowing for the calculation of binding affinity (dissociation constant, Kd, or inhibition constant, Ki) [19].

These assays have revealed distinct binding specificities that align with the beetle's ecology. Several TcOBPs are clearly implicated in defense and detoxification. TcOBPC12 binds the plant-derived compound eugenol; reported affinities vary across assays, with estimates in the low micromolar range [14]. Its role in responding to various chemical agents was further supported by subsequent studies. Likewise, TcOBPC11 and TcOBPC17 bind components of plant essential oils, such as those from *Artemisia vulgaris* [20,21]. More recently, TcOBPC02 has been suggested to participate in the beetle's defense against eucalyptol, although direct experimental validation remains limited [22]. In contrast, other OBPs have more canonical olfactory roles. The antennally expressed TcOBP9A and TcOBP9B are

implicated in sensing broad food-related volatiles, including green leaf volatiles; current evidence is primarily from expression patterns and functional inference, suggesting a role in general food-source recognition rather than specific pheromone detection [23].

A critical and still-debated aspect of OBP function is the mechanism of ligand release at the surface of the neuronal dendrite. The leading hypothesis is a pH-dependent conformational change [24]. The sensillar lymph is maintained at a generally neutral pH, while the microenvironment near the negatively charged dendritic membrane is thought to be more acidic. This pH drop is proposed to induce a conformational change in the OBP, particularly in the C-terminal region, which "unplugs" the binding pocket and lowers the affinity for the ligand, facilitating its release to the receptor [25]. This model is supported by biophysical evidence for TcOBPs. Circular dichroism (CD) spectroscopy has demonstrated that both TcOBPC12 and TcOBP9A undergo significant conformational changes, with a decrease in α-helical content, as the pH drops from 7.4 to 5.0. This change is accompanied by a marked reduction in ligand-binding affinity, providing strong evidence for this release mechanism in *T. castaneum* [14,21,23]. In addition to pH-driven conformational changes, several alternative or complementary mechanisms have been proposed based on work in other insects. These include allosteric gating triggered by direct interactions with olfactory receptors or membrane lipids, OBP multimerization, and the participation of auxiliary proteins such as sensory neuron membrane proteins (SNMPs). Although such mechanisms have not yet been dissected in detail in *T. castaneum*, they highlight that pH-dependent release is unlikely to be the only solution and that OBP–receptor coupling probably involves a broader set of dynamic molecular events.

At the same time, fluorescence competitive binding assays using 1-NPN have inherent limitations. They report relative affinity under a defined set of in vitro conditions rather than absolute specificity under physiological conditions, and are sensitive to factors such as inner-filter effects, probe solubility, and pH-dependent changes in protein conformation. Consequently, high apparent affinity in these assays does not automatically translate into in vivo functional relevance, and binding data are best interpreted in combination with expression profiles and genetic evidence such as RNAi phenotypes.

2.3. The OBP-Receptor Interface and Signal Transduction

The final step in the canonical OBP pathway involves the delivery of the odorant to a membrane-bound receptor complex, which converts the chemical signal into an electrical one. In insects, this is typically a heteromeric complex formed by a specific, ligand-binding Odorant Receptor (OR) and a highly conserved coreceptor known as Orco [26,27]. In addition to OR–Orco complexes, insects also deploy ionotropic receptors (IRs) and gustatory receptors (GRs) for chemosensory detection, and it is conceivable that soluble binding proteins contribute to ligand delivery or modulation across these receptor classes as well, although such roles remain largely unexplored in *T. castaneum*.

T. castaneum possesses a remarkably large repertoire of over 250 predicted OR genes, reflecting the chemical complexity of its environment [28]. The Orco protein is essential for the proper trafficking, membrane localization, stability, and ion channel function of the entire OR-Orco complex [27,29,30]. Studies in other beetles have shown that silencing Orco via RNAi effectively disables odorant sensing, underscoring its central role [31].

While direct evidence for specific OBP-OR interactions in vivo remains limited, it has been hypothesized that general odorant binders like TcOBP9A may interact with certain Ors, such as TcOR16 and TcOR18, based on co-expression and response profiles [23,32,33]. In addition, preliminary evidence suggests that TcOBPC12 may colocalize with TcOrco in sensory neurons under chemical stress conditions, although further validation is needed to confirm a functional OBP-receptor coupling [21,22]. The olfactory system is further compli-

cated by the presence of other protein families, such as SNMPs, which may also participate in the receptor complex, particularly for pheromone detection in some insects [34].

3. Genomic Organization, Expression, and Regulation

3.1. A Spatiotemporal Expression Atlas of T. castaneum OBPs

The functional diversity of the OBP family is mirrored by the remarkable specificity of their gene expression patterns across different tissues, developmental stages, and sexes. This spatiotemporal regulation provides strong clues about the physiological roles of individual OBPs, as summarized in Table 1. A key finding is that many TcOBPs are not confined to the antennae. For instance, TcOBPC11, TcOBPC17, and the recently characterized TcOBPC02 are all highly expressed in tissues central to metabolism and detoxification, such as the fat body, Malpighian tubules, epidermis, and hemolymph [35]. This expression pattern strongly supports their proposed roles in defense against xenobiotics, which are processed in these organs. In contrast, other OBPs like TcOBP9A and TcOBP9B show classic antennal-specific expression, consistent with a primary role in olfaction [23].

Table 1. Functionally Characterized Odorant Binding Proteins in *Tribolium castaneum*.

OBP	GenBank Accession	Subfamily	Primary Expression Tissues	Key Ligands/Inducers	Established Function(s)	References
TcOBPC11	XM_962706	Classic	Fat body, Malpighian tubules, Epidermis, Antennae	*Artemisia vulgaris* essential oil	Detoxification, Defense, Immunity	[35]
TcOBPC12	NP_001107842.1	Classic	Epidermis, Fat body, Antennae	Eugenol, Various chemical agents, Insecticides	Detoxification, Defense, Immunity	[21]
TcOBPC17	XP_008194483.1	Classic	Head, Fat body, Epidermis, Hemolymph	*Artemisia vulgaris* essential oil	Detoxification, Defense	[20]
TcOBPC02	-	Classic	Head, Epidermis, Hemolymph	Eucalyptol	Phytochemical Defense	[22]
TcasOBP9A	NP_001107850.1	Classic	Antennae	General odorants (e.g., green leaf volatiles)	Olfaction (General Volatile Detection)	[23]
TcasOBP9B	NP_001107851.1	Classic	Antennae	General odorants (e.g., green leaf volatiles)	Olfaction (General Volatile Detection)	[23]
TcasOBP7G	-	Minus-C	Reproductive tissues, Antennae	Juvenile Hormone III, Phytochemicals	Reproduction, Defense, Immunity	[16]

Expression levels are also dynamically regulated throughout the beetle's life cycle. For example, TcOBPC11 expression peaks during the late larval and adult stages, while TcOBPC02 is most abundant in early larvae and pupae [22]. This temporal orchestration likely reflects the different chemical challenges and physiological requirements of each developmental stage. Furthermore, sexually dimorphic expression points to roles in reproduction.

TcOBP10 is reported to show male-biased expression in the antennae, suggesting a possible role in detecting sex pheromones. TcOBP7G has been detected in female reproductive tissues and may be associated with oviposition, but direct functional validation is still lacking [16].

Our ability to map these expression patterns has been greatly enhanced by methodological advances. While quantitative real-time PCR (qRT-PCR) and RNA sequencing (RNA-Seq) provide quantitative data, techniques like fluorescent in situ hybridization (FISH) and, more recently, single-cell RNA sequencing (scRNA-seq) offer unprecedented

spatial resolution. For example, FISH revealed that the closely related and genomic-linked TcOBP9A and TcOBP9B are, remarkably, expressed in mutually exclusive, non-overlapping populations of sensory neurons within the antenna [23]. Although scRNA-seq datasets for *T. castaneum* remain limited, preliminary studies suggest distinct clusters of sensory neurons may express unique combinations of OBPs and ORs [36].

3.2. The Regulatory Network: Transcriptional, Hormonal, and Environmental Control

The specific expression patterns of OBP genes are orchestrated by a complex regulatory network that integrates diverse internal and external signals. A dominant theme in *T. castaneum* research is the induction of OBP expression by chemical cues. Exposure to plant secondary metabolites, such as components of essential oils like eugenol and eucalyptol, triggers a significant upregulation of defense-related OBPs, including TcOBPC11, TcOBPC12, TcOBPC17, and TcOBPC02 [20,21,35]. This response is not limited to natural compounds; sublethal doses of synthetic insecticides also induce these same genes, providing a direct link between OBP expression and insecticide resistance mechanisms [9].

The molecular basis for this induction is beginning to be understood. Although functional data are limited, promoter analysis of the TcOBPC12 gene has identified binding sites for well-known stress-responsive transcription factors, including Nuclear Factor Erythroid 2-Related Factor 2 (*Nrf2*) and Activator Protein-1 (*AP-1*) [37]. These binding motifs suggest potential regulation by these factors during xenobiotic challenge [9].

Internal physiological states also exert tight control. The major insect developmental hormones, juvenile hormone (JH) and ecdysone, are critical for orchestrating the stage-specific expression profiles of OBPs like TcOBPC11 and TcOBP7G, ensuring their functions are aligned with developmental transitions such as metamorphosis [35]. Even the beetle's nutritional status plays a role; starvation has been shown to modulate the expression of several antennal OBPs. This response may involve insulin signaling pathways, as reported in Drosophila, although experimental evidence in *T. castaneum* is still lacking [38,39]. Physical environmental factors, including temperature and humidity, also fine-tune OBP expression, allowing the beetle to adapt its sensory capabilities to prevailing abiotic conditions [40]. This intricate regulation underscores that OBP genes are not merely simple olfactory components but are sophisticated regulatory hubs. They integrate a wide array of signals—developmental, physiological, and environmental—to produce a tailored protein output that optimizes the insect's fitness in its immediate context. This plasticity is a key adaptive trait that contributes to the success of this generalist pest [7,11].

3.3. Emerging Layers of Regulation: Epigenetic Modifications and miRNA Regulations

Beyond direct transcriptional control, research is uncovering more subtle layers of regulation that fine-tune OBP gene expression. These epigenetic and post-transcriptional mechanisms add significant complexity and responsiveness to the system. While this is a frontier area for TcOBPs specifically, strong parallels exist with other insect systems [41].

Epigenetic modifications, which influence gene expression without altering the DNA sequence, are increasingly implicated in insect adaptation. In *T. castaneum*, genome annotations suggest a simplified DNA methylation machinery, consistent with observations in other beetles. Although DNA methylation typically plays a limited role, histone modifications such as acetylation and methylation are hypothesized to contribute to transcriptional regulation. While no direct chromatin immunoprecipitation sequencing (ChIP-seq) data are currently available in *T. castaneum*, analogous evidence from other insects suggests that chromatin remodeling could influence OBPs expression in response to essential oils and xenobiotics [42].

Non-coding RNA regulation is increasingly recognized as a key layer of gene expression control in insects. In *T. castaneum*, microRNAs (miRNAs) have been shown to modulate physiological processes such as diapause and reproduction. For example, *tca-miR-277-3p* regulates reproductive diapause by targeting transcription factors like *Foxo* in a mating-dependent manner. Although no direct miRNA–OBP interaction has been experimentally confirmed, it remains a plausible regulatory mechanism [43]. Additionally, alternative splicing of OBP genes may generate isoforms with distinct C-terminal regions, offering functional diversification that merits further investigation [44].

4. The Expanding Functional Repertoire of *T. castaneum* OBPs

4.1. Central Roles in Detoxification and Defense Against Xenobiotics

One of the most significant paradigm shifts in OBP biology, powerfully illustrated by research in *T. castaneum*, is the recognition of their central role in detoxification and defense. This function moves far beyond the canonical model of olfaction and repositions many OBPs as key players in the insect's chemical defense arsenal. The evidence for this is multifaceted and compelling. A suite of TcOBPs, including TcOBPC11, TcOBPC12, TcOBPC17, and TcOBPC02, is robustly induced upon exposure to a wide range of xenobiotics, from plant-derived toxins like essential oils and their components (e.g., eugenol, eucalyptol) to synthetic insecticides [20–22].

Crucially, this correlation is backed by direct functional evidence. RNAi-mediated knockdown of these specific OBP genes often renders the beetle more susceptible to the corresponding toxins, supporting the view that these proteins are not just bystanders but likely contribute substantially to defense [22,35]. Taken together, TcOBPC11, TcOBPC12, TcOBPC17, and TcOBPC02 therefore represent the best-supported examples of detoxification-related OBPs in *T. castaneum*, where induction profiles, ligand-binding assays, and RNAi phenotypes converge. At the same time, RNAi-based evidence has its own limitations: knockdown efficiency can vary among tissues and developmental stages, and residual transcript levels may partially mask gene function. Thus, while these OBPs are strongly implicated in xenobiotic defense, definitive dissection of redundancy within the family will ultimately require stable multi-gene knockouts. This defensive role is not limited to natural compounds. Constitutively higher expression of these same OBPs is observed in insecticide-resistant strains of *T. castaneum*, and the proteins have been shown to bind insecticides directly, implicating them in resistance mechanisms [14,35]. This phenomenon is not unique to *T. castaneum*; similar roles for OBPs and the related Chemosensory Proteins (CSPs) in insecticide resistance are being uncovered across a wide range of insect pests [45].

The proposed mechanism for this defensive function is sequestration. By binding to toxic molecules in the hemolymph, gut, or other tissues, these soluble OBPs effectively act as molecular sponges, reducing the free concentration of the toxin and preventing it from reaching its physiological target sites [7,9,46]. This action can be considered a "Phase 0" detoxification step, where the OBP captures and transports the xenobiotic, possibly to dedicated metabolic enzymes (like cytochrome P450s) for subsequent degradation and excretion. Given the constant chemical assault faced by a generalist pest in stored products—from plant allelochemicals, fungal mycotoxins, and pesticides—it is plausible that for some OBP lineages in *T. castaneum*, broad-spectrum defensive sequestration represents a co-opted role, whereas olfactory transport is likely ancestral; definitive evolutionary assignments remain to be tested. The olfactory role may have been a later, more specialized co-option of this versatile and robust binding scaffold [20,22,35].

4.2. Contributions to Innate Immunity and Other Physiological Processes

The functional diversification of TcOBPs extends even beyond detoxification. There is growing correlative evidence for their involvement in the insect's innate immune system. Exposure to entomopathogenic fungi like *Beauveria bassiana* and bacteria such as *Bacillus thuringiensis* leads to the upregulation of OBPs like TcOBPC11 and TcOBPC12 in immune-competent tissues like the fat body and cuticle [9,20,35]. This suggests that OBPs may participate in the immune response, perhaps by binding to pathogen-associated molecular patterns (PAMPs) or by detecting pathogen-released volatiles that signal danger [46,47]. However, in contrast to detoxification, direct genetic evidence for immune functions of TcOBPs is still sparse. To our knowledge, there are currently no published studies in *T. castaneum* that combine OBP knockdown with quantitative measurements of infection outcome (e.g., pathogen load or host survival). This gap makes it difficult to distinguish between OBPs acting as bona fide immune effectors and OBPs that are simply co-regulated with other stress-responsive genes. Highlighting this limitation is important to avoid over-interpretation and to motivate future work that explicitly tests OBP-mediated innate immunity.

Reproduction is another domain where OBPs play a role. The Minus-C OBP TcOBP7G is implicated in reproductive processes in *T. castaneum*, potentially by binding and transporting signaling molecules like Juvenile Hormone III (JH III) [7,16]. In other insect species, OBPs have been identified as components of the seminal fluid that are transferred from males to females during mating, where they can modulate female post-mating physiology and behavior, such as oviposition [48]. While not yet fully explored in *T. castaneum*, this represents a probable and important function. Finally, research in other insects has hinted at OBP involvement in non-olfactory sensory modalities, such as hygrosensation (humidity detection) and thermosensation [7]. These emerging, non-canonical functions highlight the remarkable evolutionary plasticity of the OBP family, which has been adapted to serve a wide array of physiological needs.

5. Translational Prospects: Leveraging OBP Biology for Innovation

5.1. Targeting OBPs for Novel Pest Management Strategies

The fundamental knowledge gained about TcOBP function is directly fueling the development of innovative and potentially more sustainable pest management strategies. Because OBPs are a critical interface between the insect and its chemical ligands, disrupting their function offers a powerful means of controlling pest behavior and survival [22].

5.1.1. OBP-Modulating Compounds as Repellents and Insecticides

The prominent role of OBPs like TcOBPC12 in detoxification makes them excellent targets for chemical intervention. Compounds designed to bind tightly to the OBP's active site could act as potent synergists, blocking the protein's ability to sequester a toxin and thereby increasing the efficacy of existing insecticides or natural pesticides [14,21]. Taking this a step further, structure-based virtual screening, guided by homology models of TcOBPC12, has suggested potential candidates that may not only bind with high affinity but also exhibit putative insecticidal activity and disrupt the beetle's orientation behavior [14,21].

Conversely, understanding the ligands for OBPs involved in attraction can lead to behavior-modifying strategies. Identifying the specific host volatiles that bind to generalist OBPs like TcOBP9A and TcOBP9B can inform the creation of more effective attractant blends for monitoring and mass-trapping systems [23]. In parallel, screening for compounds that act as antagonists—binding to the OBP but failing to elicit an attractive response—can yield novel repellents or confusants. For instance, *T. castaneum* actively avoids volatiles such as linalool and geraniol, which are emitted by its predator *Xylocoris flavipes*, making these compounds promising candidates for development as natural repellents [49,50].

5.1.2. RNA Interference (RNAi) as a Tool for OBP-Targeted Control

The potent and systemic RNAi response in *T. castaneum* makes it an ideal organism for exploring gene silencing as a pest control technology. The proof-of-concept for targeting OBPs is robust: laboratory studies have repeatedly shown that delivering double-stranded RNA (dsRNA) to silence defense-related OBPs like TcOBPC11, TcOBPC12, and TcOBPC17 significantly increases the beetle's mortality when subsequently exposed to the corresponding toxins [4,21,35].

The primary obstacle for RNAi-based pest control has always been the effective and stable delivery of dsRNA in a field setting [51]. However, significant progress is being made on this front. Nanoparticle carriers, such as chitosan, liposomes, and star polycations, as well as emerging platforms like carbon-based nanomaterials and silica nanoparticles, offer a promising solution [52]. As highlighted in a recent review, these diverse nanocarriers unveil new frontiers in pest control by acting as protective shields against nucleases and facilitating efficient transmembrane delivery, thereby significantly enhancing dsRNA stability and cellular uptake [52]. For example, using chitosan nanoparticles to encapsulate dsRNA has been shown to protect the fragile molecules from environmental degradation and nucleases in the insect gut, leading to more efficient gene silencing and higher mortality in model insects [53,54].

Another avenue is the self-assembled RNA nanostructures (SARNs) platform, a scalable RNA delivery system enabling targeted gene silencing in diverse insect species. SARNs improve RNA stability and delivery in plants and in model pests with chewing mouthparts (*T. castaneum*) and piercing-sucking mouthparts (*Nilaparvata lugens*). SARNs surpass traditional dsRNA systems in stability and translocation, offering a cost-effective, field-deployable solution for RNAi-based pest control.

From a broader perspective, *T. castaneum* illustrates both the strengths and the practical limitations of RNAi-based strategies. Its high systemic RNAi sensitivity makes it an ideal model to demonstrate proof-of-concept for OBP-targeted gene silencing, yet the very efficient responses observed under controlled laboratory conditions do not automatically translate into field-ready products. Environmental nucleases, UV exposure, and formulation costs all act against stable delivery of dsRNA in real storage environments. Nanocarriers and RNA nanostructures clearly improve stability and uptake, but their large-scale production, regulatory acceptance, and long-term environmental impact are still open questions. For these reasons, we focus here on how OBP biology can be harnessed by RNAi, while referring readers to dedicated RNAi reviews for a more exhaustive treatment of RNAi mechanisms across insects. Key OBP-targeted pest management strategies, including behavioral modulators, synergists, and RNAi-based approaches, are summarized in Table 2.

Table 2. Emerging Pest Management Strategies Targeting *T. castaneum* OBPs.

Strategy	Target OBP(s)	Approach/Method	Key Finding	References
RNAi-mediated lethality enhancement	TcOBPC11, TcOBPC12, TcOBPC17	Larval dsRNA injection or feeding	Knockdown significantly increases mortality when larvae are subsequently exposed to eugenol or *Artemisia vulgaris* essential oil (mortality significantly higher than control).	[20,21,35]
	TcOBPC02	Larval dsRNA injection	Knockdown increases the susceptibility of larvae to eucalyptol.	[22]
Proof-of-concept synergist	TcOBPC12	In silico docking and fluorescence competition binding assays	Docking and fluorescence competition identified several ligands with strong micromolar affinity to TcOBPC12; proposed as candidate synergists against eugenol, but no in vivo insecticidal tests have been performed.	[14,21]
Antennal sensitivity enhancement	TcOBP9A, TcOBP9B	Electroantennography (EAG) + RNAi	Knockdown of TcOBP9A or TcOBP9B significantly reduces antennal responses to several food-related volatiles (e.g., 2-hexanone, (E)-2-heptenal, 6-methyl-5-hepten-2-one), indicating that these OBPs enhance odor detection at the antennal level; behavioral attraction assays were not performed.	[23]
Behavioral repellence (predator-derived)	Not directly via OBP blockade	Exposure to *Xylocoris flavipes* volatiles	Volatiles emitted by the predator *X. flavipes*, particularly linalool and geraniol, significantly reduce the orientation of *T. castaneum* towards food sources and act as potent spatial repellents in laboratory olfactometer assays; no OBP-based mechanism has been validated.	[49,50]

5.2. OBPs in Biotechnology: From Biosensors to Bioremediation

5.2.1. OBP-Based Biosensors

The ability of OBPs to bind specific volatile compounds with high affinity and selectivity makes them ideal biorecognition elements for the development of highly sensitive biosensors. These "bio-electronic noses" have potential applications in diverse fields, including the detection of pheromones for pest monitoring, identification of food contaminants and spoilage markers, and sensing of environmental pollutants or medical biomarkers [55,56].

Recent advances have focused on integrating OBPs with nanotechnology platforms to create these devices. In other insect models, OBPs have been immobilized on nanomaterials such as graphene and gold nanoparticles, generating electrical signals upon odorant binding. Similar strategies are being explored using recombinant *TcOBP9A* and *TcOBP9B*, with promising preliminary results [56,57]. Other approaches have successfully coupled TcOBP9A to gold nanoparticles to create an electrochemical sensor for citrus volatiles and linked TcOBPC12 with fluorescent quantum dots to fabricate an optical nanosensor for phenolic compounds [58].

5.2.2. Other Potential Applications

The fundamental ability of OBPs to bind and transport small hydrophobic molecules is being explored for other innovative applications. In environmental science, detoxification-related OBPs like TcOBPC11 and TcOBPC12 have been immobilized onto biopolymer matrices to create novel bio-filters capable of sequestering and removing phenolic pollutants from contaminated water. In the pharmaceutical field, theoretical applications of OBPs in drug delivery are being explored, based on their ability to transport hydrophobic molecules [59,60]. However, functional validation in pharmaceutical contexts is still lacking. These emerging applications demonstrate the remarkable versatility of the OBP scaffold and the potential for translating fundamental insect biology into diverse technological solutions. The following table (Table 3) summarizes the current and emerging biotechnological and environmental applications of TcOBPs, highlighting their functional versatility and translational potential.

Table 3. Biotechnological and Environmental Applications of *T. castaneum* OBPs.

Application Field	Insect OBP(s)	Platform/Technique	Actual Reported Outcome	References
Graphene-based bioelectronic nose prototype	TcOBP9A, TcOBP9B (preliminary)	Graphene/reduced Graphene Oxide (rGO) field-effect transistor (FET)	Recombinant TcOBP9A/B immobilized on graphene-based FET devices show reproducible electrical responses to model volatiles such as sulcatone and (S)-(+)-3-octanol in buffer; proposed as a proof-of-concept olfactory biosensor, but no food-matrix or field applications have been reported.	[56–58]
Fluorescence binding assay/future nanosensor	TcOBPC12 (preliminary)	Solution fluorescence quenching	Intrinsic fluorescence quenching assays demonstrate binding of eugenol and related phenolic/terpenoid compounds to TcOBPC12 in vitro; the protein has been suggested as a candidate recognition element for future nanosensors, but no immobilized sensor device has yet been reported.	[21]
Environmental pollutant removal (non-*T. castaneum*)	Porcine and lepidopteran OBPs	OBPs immobilized on biopolymers	Laboratory-scale studies with porcine and lepidopteran OBPs demonstrate up to ~90% removal of phenolic pollutants in aqueous systems; no experimental bioremediation has been performed with *T. castaneum* OBPs.	[59]

6. Evolutionary and Comparative Perspectives

6.1. Evolution of the OBP Gene Family in Coleoptera

Placing the *T. castaneum* OBP family into a broader evolutionary context reveals key patterns of adaptation within the order Coleoptera, the beetles. The *T. castaneum* genome contains approximately 50 functional OBP genes, a moderately sized repertoire compared to other insects [23,35]. Comparative genomic analyses across multiple beetle species have shown that the OBP gene family has been shaped by lineage-specific events. For example, the Colorado potato beetle, *Leptinotarsa decemlineata*, a folivorous pest with a well-annotated genome, also exhibits lineage-specific expansions in certain OBP subfamilies, suggesting that ecological specialization and host-plant chemistry can independently drive OBP diversification in different beetle lineages. Notably, the Minus-C OBP subfamily appears to have undergone a specific expansion in the *T. castaneum* lineage [16]. Given that Minus-C OBPs are frequently implicated in non-olfactory physiological roles, this expansion may represent an adaptation to the unique challenges of the stored-product environment, such as hygrosensation and thermosensation modalities [7].

While precise dating remains to be confirmed, phylogenetic analysis suggests that the gene duplication giving rise to TcOBP9A and TcOBP9B occurred early in beetle evolution, possibly within the Cucujiformia infraorder. This indicates an ancient origin for this pair, followed by a long period of conserved function or subtle divergence. The evolutionary pressures acting on these genes are not uniform. Analyses of selection have found strong evidence of positive selection—where mutations that change the protein sequence are favored—acting on several detoxification-related OBPs, including TcOBPC11 and TcOBPC12 [23,61]. This suggests an ongoing evolutionary arms race, where the beetle's defensive proteins are rapidly evolving to cope with a changing landscape of chemical threats from plants, microorganisms, and pesticides. In contrast, core olfactory OBPs involved in general odorant perception tend to be under stronger purifying selection, indicating that their fundamental functions are more conserved across beetle lineages. These comparative studies, which frequently include OBP sequences from a wide range of other beetle species, are essential for distinguishing between ancient, conserved functions and more recent, lineage-specific adaptations [61].

6.2. Functional Divergence and Conservation Across Insecta

Expanding the comparison to other major insect orders, such as Diptera (flies) and Lepidoptera (moths), highlights both universal principles and fascinating divergences in OBP evolution. The overall size of the OBP gene repertoire varies dramatically between species, from as few as eight in the whitefly to over 100 in the German cockroach, likely reflecting vast differences in their ecological niches and reliance on chemical communication [62,63]. The primary engine driving this diversity is a process known as birth-and-death evolution, where new genes are created through duplication and subsequently either lost or retained and modified for new functions.

This process results in a remarkable degree of functional plasticity within the OBP family. The same basic protein scaffold can be repurposed to meet different ecological needs. A commonly cited example involves TcOBP9A/B and the Drosophila protein Lush (DmOBP76a), which are hypothesized to share a common ancestor. However, while Lush is specialized for pheromone detection and courtship, TcOBP9A/B are believed to facilitate detection of general food volatiles, suggesting functional divergence despite possible homology [23,62]. This demonstrates how function can become decoupled from ancestry, allowing orthologous genes to be co-opted for entirely different roles in different lineages.

Conversely, the phenomenon of functional convergence is also apparent. The role of TcOBPC12 in binding and detoxifying xenobiotics shows striking functional parallels with certain insecticide-resistance-associated OBPs in mosquitoes [62]. Although there is little sequence homology, these proteins appear to have evolved similar detoxification capabilities, likely under analogous selective pressures. The widespread occurrence of non-olfactory OBP functions, particularly in detoxification and immunity, across diverse insect orders suggests that this is a common evolutionary trajectory. It may reflect repeated, independent co-option of this versatile binding scaffold in response to chemical stress, or alternatively, indicate an ancient but now broadly diversified ancestral role [8,9].

7. Synthesis and Future Outlook

7.1. Summary of Key Advances and Emerging Paradigms

Research has matured from foundational gene identification to the detailed functional characterization of several key players, including TcOBPC11, TcOBPC12, TcOBPC17, TcOBP7G, and the TcOBP9A/B pair. A central and recurring theme has been the paradigm shift away from an exclusively olfaction-centric view of OBP function. Compelling evidence now firmly establishes that many TcOBPs are integral components of the beetle's defense and detoxification systems, responding to and protecting against a wide array of xenobiotics. This has been corroborated by detailed gene expression studies mapping OBP activity to specific tissues and developmental stages, with RNAi providing definitive functional validation. Comparative genomics has successfully placed TcOBPs within the broader evolutionary tapestry of insect chemosensation, revealing a dynamic history of gene duplication, functional divergence, and adaptive selection. This fundamental knowledge is now being translated into tangible applications, with the targeting of OBPs for novel pest control strategies—particularly through RNAi and behavior-modifying compounds—emerging as a highly promising avenue.

7.2. Current Gaps, Challenges, and Future Research Frontiers

Despite this significant progress, substantial knowledge gaps and challenges persist. First, the precise biological roles of the majority of the ~50 TcOBPs remain unresolved, and ligands are known with confidence for only a small subset. Second, for non-olfactory OBPs that are highly expressed in tissues such as fat body, hemolymph, and gut, their exact subcellular localization and in vivo binding partners are largely unknown. Third,

interactions between TcOBPs and canonical detoxification systems—cytochrome P450s, glutathione S-transferases (GSTs), and esterases—have barely been explored, even though such crosstalk is likely central to xenobiotic handling. For nearly all TcOBPs, high-resolution 3D structures are lacking, which severely hinders a deep mechanistic understanding of ligand binding and prevents true structure-based rational design of inhibitors. The intricate choreography of OBP-receptor interactions and the full complexity of OBP gene regulatory networks are still being mapped. From an applied perspective, translating the success of laboratory-based RNAi into field-stable, cost-effective, and environmentally safe pest control products presents formidable logistical and regulatory hurdles [19,64]. A key biological challenge is the issue of functional redundancy; with a large family of related proteins, the knockout of a single OBP may be compensated for by another, potentially masking its true function and complicating functional characterization.

Addressing these gaps will require the deployment of a new generation of research tools and a shift towards more integrative research paradigms. The future of OBP research lies not in any single technology, but in the synergistic integration of multiple cutting-edge approaches. A truly comprehensive understanding will emerge from combining systems-level discovery, precise genetic manipulation, high-resolution molecular analysis, and sophisticated functional assays. This integrated, multi-scale approach represents the next frontier. Key future research directions include:

- CRISPR/Cas9 Gene Editing: Moving beyond the transient effects of RNAi, CRISPR/Cas9 technology allows for the creation of stable, heritable knockout lines for precise functional analysis. Crucially, it enables the generation of multi-gene knockouts, which will be essential for overcoming the challenge of functional redundancy and dissecting the roles of clustered or closely related OBP genes.
- Single-Cell Transcriptomics (scRNA-seq): This revolutionary technology provides the ability to resolve OBP expression profiles at the ultimate level of resolution: single cells. Applying scRNA-seq to the beetle's antennae and other tissues will create a high-resolution atlas of the chemosensory and defensive systems, identifying the precise cellular context in which each OBP functions and revealing novel, rare cell types that may have been missed by bulk analyses.
- Artificial Intelligence and Machine Learning: The growing volume of sequence, structure, and functional data for OBPs is ripe for the application of AI. Machine learning models can be trained to predict ligand-binding properties with increasing accuracy, to perform virtual screening of immense chemical libraries for novel OBP inhibitors or modulators, and to mine large-scale 'omics datasets to generate new, data-driven hypotheses about OBP function.
- Structural Biology: A concerted effort to solve the experimental 3D structures of key TcOBPs—both in their unbound (apo) form and in complex with ecologically relevant ligands—is a critical priority. Techniques like X-ray crystallography, Nuclear Magnetic Resonance (NMR) spectroscopy, and cryo-electron microscopy (Cryo-EM) will provide the atomic-level detail necessary to understand the dynamics of ligand binding and release, which is the essential foundation for the rational design of next-generation pest control molecules.
- Systems Biology: Ultimately, the goal is to move beyond studying individual proteins in isolation. The future lies in integrating these multilevel data streams—genomic, transcriptomic, proteomic, structural, and functional—to build systems-level models of the beetle's entire chemosensory and defensive apparatus. This will allow researchers to understand how the system as a whole responds to complex chemical environments and to predict the consequences of targeted interventions. From an applied standpoint, it is also important to recognize practical constraints. Field deployment of

OBP-targeted RNAi will have to contend with dsRNA instability, formulation costs, and regulatory hurdles, while OBP-based biosensors must be engineered into robust, user-friendly devices before they can move beyond laboratory prototypes. Likewise, turning high-affinity OBP ligands into viable repellents, attractants, or synergists involves extensive medicinal chemistry and safety evaluation. Explicitly acknowledging these constraints provides a more realistic context for the promising translational opportunities outlined above.

In conclusion, the study of Odorant Binding Proteins in *Tribolium castaneum* has evolved into a dynamic and interdisciplinary field. These small, versatile proteins, once viewed as passive transporters, are now understood to be central players in the beetle's complex interactions with its chemical world, from olfaction and reproduction to detoxification and immunity. As research continues to leverage cutting-edge technologies and integrative approaches, TcOBPs will undoubtedly yield further fundamental insights into insect biology and provide innovative solutions for pest management and biotechnology.

Author Contributions: Conceptualization, Z.Z. and Y.L.; methodology, L.W. and Z.Z.; writing—original draft preparation, L.W. and Z.Z.; writing—review and editing, Z.Z.; supervision, Z.Z. and Y.L.; project administration, Y.L.; funding acquisition, Y.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by grants from the National Key R&D Program of China (No. 2023YFC2604903) and the Natural Science Foundation of China (No.31871975).

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors have declared no conflicts of interest.

References

1. Agarwal, A.; Agashe, D. The Red Flour Beetle *Tribolium castaneum*: A Model for Host-Microbiome Interactions. *PLoS ONE* **2020**, *15*, e0239051. [CrossRef]
2. Campbell, J.F.; Athanassiou, C.G.; Hagstrum, D.W.; Zhu, K.Y. *Tribolium castaneum*: A Model Insect for Fundamental and Applied Research. *Annu. Rev. Entomol.* **2022**, *67*, 347–365. [CrossRef]
3. Klingler, M.; Bucher, G. The Red Flour Beetle *T. castaneum*: Elaborate Genetic Toolkit and Unbiased Large Scale RNAi Screening to Study Insect Biology and Evolution. *EvoDevo* **2022**, *13*, 14. [CrossRef]
4. Rösner, J.; Wellmeyer, B.; Merzendorfer, H. *Tribolium castaneum*: A Model for Investigating the Mode of Action of Insecticides and Mechanisms of Resistance. *Curr. Pharm. Des.* **2020**, *26*, 3554–3568. [CrossRef]
5. Herndon, N.; Shelton, J.; Gerischer, L.; Ioannidis, P.; Ninova, M.; Dönitz, J.; Waterhouse, R.M.; Liang, C.; Damm, C.; Siemanowski, J. Enhanced Genome Assembly and a New Official Gene Set for *Tribolium castaneum*. *BMC Genom.* **2020**, *21*, 47. [CrossRef] [PubMed]
6. Schmitt-Engel, C.; Schultheis, D.; Schwirz, J.; Ströhlein, N.; Troelenberg, N.; Majumdar, U.; Dao, V.A.; Grossmann, D.; Richter, T.; Tech, M. The iBeetle Large-Scale RNAi Screen Reveals Gene Functions for Insect Development and Physiology. *Nat. Commun.* **2015**, *6*, 7822. [CrossRef] [PubMed]
7. Rihani, K.; Ferveur, J.-F.; Briand, L. The 40-Year Mystery of Insect Odorant-Binding Proteins. *Biomolecules* **2021**, *11*, 509. [CrossRef]
8. Sun, J.S.; Xiao, S.; Carlson, J.R. The Diverse Small Proteins Called Odorant-Binding Proteins. *R. Soc. Open Biol.* **2018**, *8*, 180208. [CrossRef] [PubMed]
9. Abendroth, J.A.; Moural, T.W.; Wei, H.; Zhu, F. Roles of Insect Odorant Binding Proteins in Communication and Xenobiotic Adaptation. *Front. Insect Sci.* **2023**, *3*, 1274197. [CrossRef]
10. Mam, B.; Karpe, S.D.; Sowdhamini, R. Minus-C Subfamily Has Diverged from Classic Odorant-Binding Proteins in Honeybees. *Apidologie* **2023**, *54*, 12. [CrossRef]
11. Rana, A.; Sharma, D.; Choudhary, K.; Kumari, P.; Ruchika, K.; Yangchan, J.; Kumar, S. Insight into Insect Odorant Binding Proteins: An Alternative Approach for Pest Management. *J. Nat. Pestic. Res.* **2024**, *8*, 100069. [CrossRef]
12. Leal, W.S. Odorant Reception in Insects: Roles of Receptors, Binding Proteins, and Degrading Enzymes. *Annu. Rev. Entomol.* **2013**, *58*, 373–391. [CrossRef] [PubMed]

13. Zheng, Z.-C.; Li, D.-Z.; Zhou, A.; Yi, S.-C.; Liu, H.; Wang, M.-Q. Predicted Structure of a Minus-C OBP from Batocera Horsfieldi (Hope) Suggests an Intermediate Structure in Evolution of OBPs. *Sci. Rep.* **2016**, *6*, 33981. [CrossRef]
14. Wang, L.; Hou, M.; Liang, C.; Xu, Q.; Lu, Y.; Zhao, Z. Role of Odorant Binding Protein C12 in the Response of *Tribolium castaneum* to Chemical Agents. *Pestic. Biochem. Physiol.* **2024**, *201*, 105861. [CrossRef] [PubMed]
15. Qin, J.-H.; Wang, C.-Q.; Li, K.-B.; Cao, Y.-Z.; Peng, Y.; Feng, H.-L.; Yin, J. Molecular Characterization of Sex Pheromone Binding Proteins from *Holotrichia Oblita* (Coleoptera: Scarabaeida). *Int. J. Biol. Macromol.* **2021**, *193*, 8–18. [CrossRef]
16. Gao, S.; Zhang, X.; Zhou, C.; Zhang, G.; Zhang, Y.; Li, R.; Zhang, K. Minus-C Odorant Binding Protein TcasOBP7G Contributes to Reproduction and Defense against Phytochemical in the Red Flour Beetle, *Tribolium castaneum*. *J. Asia-Pac. Entomol.* **2023**, *26*, 102151. [CrossRef]
17. Wang, L.; Wen, Z.; Liu, S.-W.; Zhang, L.; Finley, C.; Lee, H.-J.; Fan, H.-J.S. Overview of AlphaFold2 and Breakthroughs in Overcoming Its Limitations. *Comput. Biol. Med.* **2024**, *176*, 108620. [CrossRef]
18. Li, D.; Li, C.; Liu, D. Analyses of Structural Dynamics Revealed Flexible Binding Mechanism for the *Agrilus Mali* Odorant Binding Protein 8 towards Plant Volatiles. *Pest Manag. Sci.* **2021**, *77*, 1642–1653. [CrossRef]
19. Shukla, S.; Nakano-Baker, O.; Godin, D.; MacKenzie, D.; Sarikaya, M. iOBPdb a Database for Experimentally Determined Functional Characterization of Insect Odorant Binding Proteins. *Sci. Data* **2023**, *10*, 295. [CrossRef]
20. Gao, S.-S.; Li, R.-M.; Xue, S.; Zhang, Y.-C.; Zhang, Y.-L.; Wang, J.-S.; Zhang, K.-P. Odorant Binding Protein C17 Contributes to the Response to Artemisia Vulgaris Oil in *Tribolium castaneum*. *Front. Toxicol.* **2021**, *3*, 627470. [CrossRef]
21. Gao, S.; Lu, R.; Zhang, Y.; Sun, H.; Li, S.; Zhang, K.; Li, R. Odorant Binding Protein C12 Is Involved in the Defense against Eugenol in *Tribolium castaneum*. *Pestic. Biochem. Physiol.* **2021**, *179*, 104968. [CrossRef] [PubMed]
22. Gao, S.; Guo, M.; Yu, W.; Miao, W.; Ya, H.; Liu, D.; Li, R.; Zhang, K. Odorant Binding Protein TcOBPC02 Contributes to Phytochemical Defense in the Red Flour Beetle, *Tribolium castaneum*. *Entomol. Res.* **2024**, *54*, e70009. [CrossRef]
23. Montino, A.; Balakrishnan, K.; Dippel, S.; Trebels, B.; Neumann, P.; Wimmer, E.A. Mutually Exclusive Expression of Closely Related Odorant-Binding Proteins 9A and 9B in the Antenna of the Red Flour Beetle *Tribolium castaneum*. *Biomolecules* **2021**, *11*, 1502. [CrossRef] [PubMed]
24. Manoharan, M.; Fuchs, P.F.; Sowdhamini, R.; Offmann, B. Insights on pH-Dependent Conformational Changes of Mosquito Odorant Binding Proteins by Molecular Dynamics Simulations. *J. Biomol. Struct. Dyn.* **2014**, *32*, 1742–1751. [CrossRef]
25. Pelosi, P.; Iovinella, I.; Zhu, J.; Wang, G.; Dani, F.R. Beyond Chemoreception: Diverse Tasks of Soluble Olfactory Proteins in Insects. *Biol. Rev.* **2017**, *93*, 184–200. [CrossRef]
26. Zhao, J.; Chen, A.Q.; Ryu, J.; Del Mármol, J. Structural Basis of Odor Sensing by Insect Heteromeric Odorant Receptors. *Science* **2024**, *384*, 1460–1467. [CrossRef]
27. Wang, Y.; Qiu, L.; Wang, B.; Guan, Z.; Dong, Z.; Zhang, J.; Cao, S.; Yang, L.; Wang, B.; Gong, Z. Structural Basis for Odorant Recognition of the Insect Odorant Receptor OR-Orco Heterocomplex. *Science* **2024**, *384*, 1453–1460. [CrossRef]
28. Engsontia, P.; Sanderson, A.P.; Cobb, M.; Walden, K.K.; Robertson, H.M.; Brown, S. The Red Flour Beetle's Large Nose: An Expanded Odorant Receptor Gene Family in *Tribolium castaneum*. *Insect Biochem. Mol. Biol.* **2008**, *38*, 387–397. [CrossRef]
29. Butterwick, J.A.; Del Mármol, J.; Kim, K.H.; Kahlson, M.A.; Rogow, J.A.; Walz, T.; Ruta, V. Cryo-EM Structure of the Insect Olfactory Receptor Orco. *Nature* **2018**, *560*, 447–452. [CrossRef]
30. Yan, H.; Jafari, S.; Pask, G.; Zhou, X.; Reinberg, D.; Desplan, C. Evolution, Developmental Expression and Function of Odorant Receptors in Insects. *J. Exp. Biol.* **2020**, *223*, jeb208215. [CrossRef]
31. He, P.; Ma, Y.-F.; Wang, M.-M.; Wang, H.; Dewer, Y.; Abd El-Ghany, N.M.; Chen, G.-L.; Yang, G.-Q.; Zhang, F.; He, M. Silencing the Odorant Coreceptor (Orco) Disrupts Sex Pheromonal Communication and Feeding Responses in *Blattella germanica*: Toward an Alternative Target for Controlling Insect-transmitted Human Diseases. *Pest Manag. Sci.* **2021**, *77*, 1674–1682. [CrossRef]
32. Abdel-Latief, M. A Family of Chemoreceptors in *Tribolium castaneum* (Tenebrionidae: Coleoptera). *PLoS ONE* **2007**, *2*, e1319. [CrossRef]
33. Miyazaki, H.; Otake, J.; Mitsuno, H.; Ozaki, K.; Kanzaki, R.; Chieng, A.C.-T.; Hee, A.K.-W.; Nishida, R.; Ono, H. Functional Characterization of Olfactory Receptors in the Oriental Fruit Fly *Bactrocera dorsalis* That Respond to Plant Volatiles. *Insect Biochem. Mol. Biol.* **2018**, *101*, 32–46. [CrossRef]
34. Cassau, S.; Krieger, J. The Role of SNMPs in Insect Olfaction. *Cell Tissue Res.* **2021**, *383*, 21–33. [CrossRef] [PubMed]
35. Zhang, Y.; Gao, S.; Xue, S.; Zhang, K.; Wang, J.; Li, B. Odorant-Binding Proteins Contribute to the Defense of the Red Flour Beetle, *Tribolium castaneum*, against Essential Oil of Artemisia Vulgaris. *Front. Physiol.* **2020**, *11*, 819. [CrossRef] [PubMed]
36. Zhu, X.; Xu, B.; Qin, Z.; Kader, A.; Song, B.; Chen, H.; Liu, Y.; Liu, W. Identification of Candidate Olfactory Genes in *Scolytus Schevyrewi* Based on Transcriptomic Analysis. *Front. Physiol.* **2021**, *12*, 717698. [CrossRef] [PubMed]
37. He, F.; Ru, X.; Wen, T. NRF2, a Transcription Factor for Stress Response and Beyond. *Int. J. Mol. Sci.* **2020**, *21*, 4777. [CrossRef]
38. Chen, Y.; Hong, B.; Zhang, Y.; Chen, X.; Zhang, T.; Zhong, G.; Yi, X. FoxO Directly Regulates the Expression of Odorant Receptor Genes to Govern Olfactory Plasticity upon Starvation in *Bactrocera dorsalis*. *Insect Biochem. Mol. Biol.* **2023**, *153*, 103907. [CrossRef]

39. Rohner, P.T.; Casasa, S.; Moczek, A.P. Assessing the Evolutionary Lability of Insulin Signalling in the Regulation of Nutritional Plasticity across Traits and Species of Horned Dung Beetles. *J. Evol. Biol.* **2023**, *36*, 1641–1648. [CrossRef]
40. Stočes, D.; Šipoš, J. Multilevel Analysis of Ground Beetle Responses to Forest Management: Integrating Species Composition, Morphological Traits and Developmental Instability. *Ecol. Evol.* **2025**, *15*, e70793. [CrossRef]
41. Mika, K.; Benton, R. Olfactory Receptor Gene Regulation in Insects: Multiple Mechanisms for Singular Expression. *Front. Neurosci.* **2021**, *15*, 738088. [CrossRef] [PubMed]
42. Länger, Z.M.; Israel, E.; Engelhardt, J.; Kalita, A.I.; Keller Valsecchi, C.I.; Kurtz, J.; Prohaska, S.J. Multiomics Reveal Associations between CpG Methylation, Histone Modifications and Transcription in a Species That Has Lost DNMT3, the Colorado Potato Beetle. *J. Exp. Zool. Part B Mol. Dev. Evol.* **2025**, *344*, 454–469. [CrossRef]
43. Güney, G.; Schmitt, K.; Zicola, J.; Toprak, U.; Rostás, M.; Scholten, S.; Cedden, D. The MicroRNA Pathway Regulates Obligatory Aestivation in a Flea Beetle. *bioRxiv* **2025**. [CrossRef]
44. Ma, H.; Liu, Y.; Tian, X.; Chen, Y.; Gao, S. Transcriptome-Wide Evaluation Characterization of microRNAs and Assessment of Their Functional Roles as Regulators of Diapause in *Ostrinia Furnacalis* Larvae (Lepidoptera: Crambidae). *Insects* **2024**, *15*, 702. [CrossRef] [PubMed]
45. Tsouri, A.; Douris, V. The Role of Chemosensory Proteins in Insecticide Resistance: A Review. *Insects* **2025**, *16*, 496. [CrossRef]
46. Zhang, W.; Xie, M.; Eleftherianos, I.; Mohamed, A.; Cao, Y.; Song, B.; Zang, L.-S.; Jia, C.; Bian, J.; Keyhani, N.O. An Odorant Binding Protein Is Involved in Counteracting Detection-Avoidance and Toll-Pathway Innate Immunity. *J. Adv. Res.* **2023**, *48*, 1–16. [CrossRef] [PubMed]
47. Bianchi, F.; Flisi, S.; Careri, M.; Riboni, N.; Resimini, S.; Sala, A.; Conti, V.; Mattarozzi, M.; Taddei, S.; Spadini, C. Vertebrate Odorant Binding Proteins as Antimicrobial Humoral Components of Innate Immunity for Pathogenic Microorganisms. *PLoS ONE* **2019**, *14*, e0213545. [CrossRef]
48. Brown, N.C.; Gordon, B.; McDonough-Goldstein, C.E.; Misra, S.; Findlay, G.D.; Clark, A.G.; Wolfner, M.F. The Seminal Odorant Binding Protein Obp56g Is Required for Mating Plug Formation and Male Fertility in *Drosophila Melanogaster*. *Elife* **2023**, *12*, e86409. [CrossRef]
49. Lu, S.; Yang, L.; Wu, Z.; Chen, M.; Lu, Y. Volatiles of the Predator *Xylocoris Flavipes* Recognized by Its Prey Tribolium Castaneum (Herbst) and *Oryzaephilus Surinamensis* (Linne) as Escape Signals. *Insects* **2024**, *16*, 31. [CrossRef]
50. Pajaro-Castro, N.; Caballero-Gallardo, K.; Olivero-Verbel, J. Neurotoxic Effects of Linalool and β-Pinene on *Tribolium castaneum* Herbst. *Molecules* **2017**, *22*, 2052. [CrossRef]
51. Lu, Y.; Deng, X.; Zhu, Q.; Wu, D.; Zhong, J.; Wen, L.; Yu, X. The dsRNA Delivery, Targeting and Application in Pest Control. *Agronomy* **2023**, *13*, 714. [CrossRef]
52. Liu, J.; He, Q.; Lin, X.; Smagghe, G. Recent Progress in Nanoparticle-Mediated RNA Interference in Insects: Unveiling New Frontiers in Pest Control. *J. Insect Physiol.* **2025**, *167*, 104884. [CrossRef]
53. Kolge, H.; Kadam, K.; Ghormade, V. Chitosan Nanocarriers Mediated dsRNA Delivery in Gene Silencing for *Helicoverpa Armigera* Biocontrol. *Pestic. Biochem. Physiol.* **2023**, *189*, 105292. [CrossRef]
54. Liu, Y.; Zhang, J.; Li, S.; Chai, L.; Chang, B.H.; Malak, M.; El Wakil, A.; Moussian, B.; Zhao, Z.; Zeng, Z. Chitosan Nanoparticle-mediated Delivery of dsRNA for Enhancing RNAi Efficiency in *Locusta migratoria*. *Pest Manag. Sci.* **2025**, *87*, 5260–5269. [CrossRef]
55. Hurot, C.; Brenet, S.; Buhot, A.; Barou, E.; Belloir, C.; Briand, L.; Hou, Y. Highly Sensitive Olfactory Biosensors for the Detection of Volatile Organic Compounds by Surface Plasmon Resonance Imaging. *Biosens. Bioelectron.* **2019**, *123*, 230–236. [CrossRef] [PubMed]
56. Teixeira, G.D.; Esteves, C.; Moro, A.J.; Lima, J.C.; Barbosa, A.J.; Roque, A.C.A. An Odorant-Binding Protein Based Electrical Sensor to Detect Volatile Organic Compounds. *Sens. Actuators B Chem.* **2024**, *411*, 135726. [CrossRef]
57. Ha, T.S.; Smith, D.P. Recent Insights into Insect Olfactory Receptors and Odorant-Binding Proteins. *Insects* **2022**, *13*, 926. [CrossRef]
58. Bensana, A.; Achi, F. Analytical Performance of Functional Nanostructured Biointerfaces for Sensing Phenolic Compounds. *Colloids Surf. B Biointerfaces* **2020**, *196*, 111344. [CrossRef] [PubMed]
59. Liu, J.; Li, Y.; Wang, P.; Zhang, Y.; Tian, Z. High-Efficiency Removal of Pyrethroids Using a Redesigned Odorant Binding Protein. *J. Hazard. Mater.* **2024**, *463*, 132856. [CrossRef] [PubMed]
60. Capo, A.; Cozzolino, S.; Cavallari, A.; Bruno, U.; Calabrese, A.; Pennacchio, A.; Camarca, A.; Staiano, M.; D'Auria, S.; Varriale, A. The Porcine Odorant-Binding Protein as a Probe for an Impedenziometric-Based Detection of Benzene in the Environment. *Int. J. Mol. Sci.* **2022**, *23*, 4039. [CrossRef]
61. Soffan, A.; Subandiyah, S.; Makino, H.; Watanabe, T.; Horiike, T. Evolutionary Analysis of the Highly Conserved Insect Odorant Coreceptor (Orco) Revealed a Positive Selection Mode, Implying Functional Flexibility. *J. Insect Sci.* **2018**, *18*, 18. [CrossRef] [PubMed]
62. Rondón, J.J.; Moreyra, N.N.; Pisarenco, V.A.; Rozas, J.; Hurtado, J.; Hasson, E. Evolution of the Odorant-Binding Protein Gene Family in Drosophila. *Front. Ecol. Evol.* **2022**, *10*, 957247. [CrossRef]

63. Robertson, H.M.; Baits, R.L.; Walden, K.K.; Wada-Katsumata, A.; Schal, C. Enormous Expansion of the Chemosensory Gene Repertoire in the Omnivorous German Cockroach *Blattella germanica*. *J. Exp. Zool. Part B Mol. Dev. Evol.* **2018**, *330*, 265–278. [CrossRef] [PubMed]
64. Venthur, H.; Zhou, J.-J. Odorant Receptors and Odorant-Binding Proteins as Insect Pest Control Targets: A Comparative Analysis. *Front. Physiol.* **2018**, *9*, 1163. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

insects

MDPI

Review

Perspectives of RNAi, CUADb and CRISPR/Cas as Innovative Antisense Technologies for Insect Pest Control: From Discovery to Practice

Hemant Kumar [1], Nikita Gal'chinsky [2,*], Verma Sweta [3], Nikita Negi [4], Roman Filatov [2], Anamika Chandel [5], Jamin Ali [6], Vol Oberemok [2] and Kate Laikova [2]

[1] Department of Entomology, Agriculture College, Garhwa, Birsa Agricultural University, Ranchi 834006, India; hksahu.iari@gmail.com
[2] Department of General Biology and Genetics, V.I. Vernadsky Crimean Federal University, 295007 Simferopol, Russia; variant_022@mail.ru (R.F.); voloberemok@gmail.com (V.O.); botan_icus@mail.ru (K.L.)
[3] Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India; drsvbiotech@gmail.com
[4] Division of Genomic Resources, National Bureau of Agricultural Insect Resources, Bengaluru 560024, India; nikinegi10@gmail.com
[5] Division of Vegetable Sciences, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India; anamikathakuriari@gmail.com
[6] College of Plant Protection, Jilin Agricultural University, Changchun 130118, China; j.alirana@yahoo.com
* Correspondence: pcr.product@gmail.com

Simple Summary: RNA interference (RNAi), contact unmodified antisense DNA biotechnology (CUADb), and clustered regularly interspaced short palindromic repeats (CRISPR) along with CRISPR-associated proteins (CRISPR/Cas) represent three innovative cost-effective antisense technologies for insect pest control. Each method operates through the formation of specific unmodified nucleic acid complexes, RNAi involves guide RNA binding to messenger RNA (mRNA); CUADb: employs guide DNA interacting with ribosomal RNA (rRNA); and CRISPR/Cas: utilizes guide RNA targeting genomic DNA. These complexes function via nucleic acid-guided nucleases: Argonaute in RNAi; rRNase in CUADb, and CRISPR-associated proteins in CRISPR/Cas systems.

Abstract

Pest management is undergoing a transformative shift with the development of the cutting-edge antisense technologies: RNA interference (RNAi), contact unmodified antisense DNA biotechnology (CUADb), and the CRISPR-associated proteins (CRISPR/Cas). These approaches function by facilitating sequence-specific pairing of nucleic acids followed by nuclease-mediated cleavage, offering exceptional precision for targeted pest control. While RNA-guided mechanisms such as RNAi and CRISPR/Cas were initially characterized in non-insect systems, primarily as innate defenses against viral infections, the DNA-guided CUADb pathway was first identified in insect pests as a functional pest control strategy. Its broader role in ribosomal RNA (rRNA) biogenesis was recognized later. Together, these discoveries have revealed an entirely new dimension of gene regulation, with profound implications for sustainable pest management. Despite sharing a common principle of sequence-specific targeting RNAi, CUADb, and CRISPR/Cas differ in several key aspects, including their mechanisms of action, target specificity, and applicability. Rather than serving as universal solutions, each technology is likely to be optimally effective against specific pest groups. Moreover, these technologies allow for rapid adaptation of control strategies to overcome target-site resistance, ensuring long-term efficacy. This review summarizes the

core functional characteristics, potential applications, and current limitations of each antisense technology, emphasizing their complementary roles in advancing environmentally sustainable pest control. By integrating foundational biological discoveries with applied innovations, this work provides a new perspectives on incorporating antisense-based strategies into next-generation integrated pest management systems.

Keywords: RNAi; CUADb; CRISPR/Cas; antisense technologies; insect pest control

1. Introduction

Nucleic acids, DNA and RNA, orchestrate cellular processes through precise complementary interactions [1]. The fundamental principles of Watson–Crick base pairing, combined with the action of specific enzymes, govern essential biological mechanisms such as replication, transcription, translation, and regulation of gene expression [2]. The specificity and fidelity of these processes arise from the unique combinations of nitrogenous bases, which constitute the molecular foundation of genetic control. Three innovative antisense technologies, RNA interference (RNAi) [3], contact unmodified antisense DNA biotechnology (CUADb) [4–6], and clustered regularly interspaced short palindromic repeats (CRISPR) along with CRISPR-associated proteins (CRISPR/Cas) [7–9], have harnessed these nucleic acid interactions to induce targeted genetic effects. These technologies operate through the formation of sequence-specific duplexes: RNAi (guide RNA–messenger RNA (mRNA)) [10], CUADb (guide DNA-ribosomal RNA (rRNA)) [5,11–14], and CRISPR/Cas (guide RNA-genomic DNA) [9,15,16], which subsequently recruit specialized nucleases such as Argonaute (Ago) [17], rRNase [5,6], and CRISPR-associated proteins [9].

The emergence of these technologies was made possible by significant advances in bioinformatics, in vitro nucleic acid synthesis, and the development of modern molecular tools. Typically, a pivotal insight or an experimental breakthrough reveals a pattern that can be reliably applied in practice. For RNAi, the key finding was the use of an antisense fragment within double-stranded RNA (dsRNA); for CUADb, it was the understanding that antisense oligodeoxyribonucleotides can be used as contact insecticides; and for CRISPR/Cas, it was the identification that genomic DNA could serve as the target for antisense-guided effects. These innovations have set new standards in molecular genetics and are now being widely explored for their applications in insect pest control. Each of these technologies emerged from fundamental research, evolving from uncertain beginnings into powerful tools with transformative potential. Since their core mechanism relies on the complementary binding of antisense molecule, either DNA or RNA, to a target nucleic acid, they are collectively referred to as antisense technologies (Figure 1). While RNA- and DNA-guided nucleases have been extensively studied, a targeted DNA-cleaving mechanism utilizing guide DNA to cut pest genomic DNA via a specific nuclease has not yet been developed, representing a potential direction for future research.

Despite the role of chemical insecticides is pest management [18], several factors drive the search for alternative control strategies. The most important of these is high economic cost of insect pest damage to agriculture and growing prevalence of insecticide resistance, which has increased dramatically and persistently since the mid-20th century [18–20]. Insecticide resistance generally arises through natural selection, whereby the frequency of specific alleles increases due to random mutations within pest populations [21]. Antisense technologies (RNAi, CUADb, and CRISPR/Cas) are able to counteract insecticide resistance by targeting conserved genes or conserved gene regions and by facilitating the rapid development of effective pest control agents when resistance to existing insec-

ticides emerges. While CUADb [6] and RNAi [3] demonstrate promising potential as next-generation bioinsecticides, due to their fast biodegradability, selectivity, and low carbon footprint [12], CRISPR/Cas is primarily used to genetically attenuate insect pest populations through genetic engineering [7]. Nevertheless, these innovative antisense approaches, and their potential combinations, offer a broad and flexible toolkit for insect pest control. The primary challenge lies in selecting the most suitable strategy for each specific pest management scenario.




Figure 1. Antisense technologies based on unmodified nucleic acids and used for insect pest control: guide nucleic acid (RNA or DNA) forms duplex with target nucleic acid (RNA or DNA) and the target nucleic acid is cleaved by specific nuclease.

The development of antisense technologies was predicted on the discovery of the DNA double helix, followed by a formative period marked by pioneering studies, such as the site-specific modification of valine tRNA by Grineva and colleagues [22] and the work of Zamecnik and Stephenson using modified DNA against the *Rous sarcoma* virus [23]. However, it became critically important to develop methods employing unmodified (natural) nucleic acids and to transform them into cost-effective, environmentally safe antisense technologies. While the RNAi was discovered by a research group in the USA in 1998 [3], CUADb was introduced in 2008 and subsequently developed further in Crimea [4,6], the development of the CRISPR/Cas9 was established in 2012 through the contributions of several research groups, primarily from Lithuania, Sweden and the United States [7,24,25]. These efforts collectively lead to the establishment of the three main antisense technologies for pest control at the turn of the 21st century. Although CUADb offers a relatively straightforward framework for pesticide design, RNAi and CRISPR/Cas are still progressing toward streamlined, reliable strategies for creating selective and effective pest control agents. The objective of this review is to provide a concise historical overview of the emergence of antisense technologies and to assess their potential in insect pest control. Overall, we provide an evaluation of the current capabilities and limitations of antisense technologies in this field, and propose directions for future development.

2. RNAi

2.1. History of Discovery of RNAi

The discovery of RNAi was inspired by the pioneering work of Zamecnik and Stephenson (1978), who demonstrated that a short antisense sequence of modified nucleic acid could inhibit the replication of the *Rous sarcoma* virus [23]. In 1998, Mello and Fire investigated the effect of both antisense and sense RNA fragments on the development of the nematode *Caenorhabditis elegans*. Their aim was to elucidate the mechanisms underlying the efficacy of sense RNA fragments synthesized by bacteriophage RNA polymerase (specifically, DNA-dependent RNA polymerases), using dsRNA fragments as experimental controls. Bacteriophage polymerases, although highly specific, occasionally produce ectopic transcripts, and DNA transgene arrays, also known as extrachromosomal arrays, are known to generate aberrant RNA products. Mello and Fire hypothesized that the interfering RNA population might include molecules with double-stranded features. Unexpectedly, it was the dsRNA fragment that triggered a potent silencing effect on the endogenous *mex-3* mRNA transcript, which is highly expressed in the gonads and early embryos of the nematode. The groundbreaking discovery, published in *Nature* in 1998 [3], established the foundational concept of RNAi: dsRNA molecules can initiate gene silencing by targeting homologous transcripts [26,27]. Subsequent research revealed that short antisense RNA fragments (21–23 nucleotides in length) are produced during RNAi and mediate the degradation of target mRNAs [28] (Figure 2).




Figure 2. Main route of RNAi used for creation of dsRNA insecticides (exogenous siRNA pathway): 1–dsRNA cleavage by Dicer; 2–silencing complex formation and silencing complex activation; 3–target mRNA degradation by Ago.

2.2. How RNAi Works in Insect Pests

RNAi has been extensively studied for its potential in agricultural pest management, particularly through the silencing of essential insect genes to suppress pest populations.

The small interfering RNA (siRNA) pathway, the most thoroughly characterized RNAi mechanism, consists of two branches: the exogenous siRNA (exo-siRNA) pathway, which mediates antiviral defense, and the endogenous siRNA (endo-siRNA) pathway, responsible for transposon suppression within the genome [29]. The exo-siRNA pathway is commonly utilized in experimental and applied contexts by introducing dsRNA to silence specific target genes [30,31]. For effective gene silencing, long dsRNA fragments (>200 bp) are typically used to trigger the production of short antisense siRNAs [26,27]. The RNAi process begins with cellular uptake of dsRNA, followed by cleavage into siRNAs (21–25 nucleotides) by the enzyme Dicer. These siRNAs are then incorporated into the RNA-induced silencing complex (RISC), where Argonaute (Ago) proteins serve as the catalytic core [32]. One strand of the siRNA duplex, known as the guide strand, directs RISC to the complementary mRNA target, leading to transcript degradation or translational repression [33].

Although much of the research has focused on the siRNA machinery, the RNAi pathway in insects is more complex and includes three sub-pathways: siRNA, micro (miRNA), and piwi-interacting RNA (piRNAs). miRNAs (18–24 nucleotides) are involved in diverse regulatory processes and interact with Ago proteins, while piRNAs (26–31 nucleotides) are primarily responsible for silencing transposable elements, particularly in germline cells, through interactions with PIWI proteins. Artificially synthesized dsRNA may influence not only the siRNA pathway but also genes associated with the miRNA pathway, suggesting a broader, interconnected regulatory network [34].

RNAi is particularly robust and systemic in Coleopteran insects [29]. However, systemic RNAi, defined as the environmental uptake of dsRNA followed by intercellular transport, is limited in Diptera, Lepidoptera [35–37], and sap-feeding Hemiptera [38,39], thereby constraining its utility in some agricultural contexts. Nonetheless, RNAi has demonstrated considerable potential for control of holometabolous and hemimetabolous insect pests, by targeting key genes involved in development, detoxification, and reproduction (Table 1).

Table 1. List of pest species successfully targeted by RNAi.

Sl. No.	Names of Model Insects	Targeted Gene(s)	Affected Processes	References
1	Beet armyworm, *Spodoptera exigua* (Lepidoptera)	*Chitin synthase gene A*	Chitin synthesis	[40]
2	Brown planthopper, *Nilaparvata lugens* (Hemiptera)	*NlHT1*, *Nlcar*, *Nltry* *NlTPS*	Digestive system Enzymatic activity	[41,42]
3	African sweet potato weevil, *Cylas puncticollis* (Coleoptera)	*Snf7*	Digestive system	[43]
4	Tomato pinworm, *Tuta absoluta* (Lepidoptera)	*Vacuolar ATPase-A and Arginine kinase*	High mortality	[44]
5	Oriental fruit fly, *Bactrocera dorsalis* (Diptera)	*α-Spectrin*	Oviposition and ovary size	[45]
6	Cotton mealybug, *Phenacoccus solenopsis* (Hemiptera)	*Krüppel homologue-1*, *ADP-ATP/Translocase*, *IDGF-1* *Bursicon*, *V-ATPase*	Not specified Cuticle hardening and V-ATPases act as proton pumps	[46,47]
7	Diamondback moth, *Plutella xylostella* (Lepidoptera)	*PxCht*	Chitin synthesis	[48]

Table 1. *Cont.*

Sl. No.	Names of Model Insects	Targeted Gene(s)	Affected Processes	References
8	Fall armyworm, *Spodoptera frugiperda* (Lepidoptera)	*Met*, *EcR*, *USP* *COPIα*, *COPIβ*, *GSTU1*	Reproductive system, fertility Insect reproduction	[49,50]
9	White-backed planthopper, *Sogatella furcifera* (Hemiptera)	*hsc70-3*, *PP-α*	Insect metamorphosis	[51]
10	Soybean aphid, *Aphis glycines* (Hemiptera)	*Cytochrome P450 monooxygenases (CYP450s)*	Insect resistance	[52]
11	Asian citrus psyllid, *Diaphorina citri* (Hemiptera)	*CHC*, *vATPase-A*, *Snf7*	Transmembrane system	[53]
12	*Trichogramma dendrolimi* (Lepidoptera)	*Vitellogenin receptor (VgR)*	Female reproductive system	[54]
13	Domestic silk moth, *Bombyx mori* (Lepidoptera)	*BmToll9-1*, *BmToll9-2*, *PGRP-L4*	Toll and immune deficiency signaling pathways	[55–58]
14	Silverleaf whitefly, *Bemisia tabaci* (Hemiptera)	*Cysteine protease*	Digestive system	[59]
15	Cowpea weevil, *Callosobruchus maculatus* (Coleoptera)	*Olfactory receptor coreceptor (Cmac\Orco)*	Insect sensory system	[60]
16	White-backed planthopper, *S. furcifera* (Hemiptera)	*β-N-acetylhexosaminidase genes*	Insect metamorphosis	[61]
17	Desert locust, *Schistocerca gregaria* (Orthoptera)	*Cytochrome P450*	Ecdysteroid pathway	[62]
18	Red flour beetle, *Tribolium castaneum* (Coleoptera)	*CPAPs*	Cuticular proteins	[63]
19	Chinese white pine beetle, *Dendroctonus armandi* (Coleoptera)	*Aquaporin*	Osmoregulation	[64]
20	Kissing bug, *Rhodnius prolixus* (Hemiptera)	*Nitrophorin 2 (NP2)*	Anticoagulant and apyrase activities in saliva	[65]
21	Citrus aphid, *Toxoptera citricida* (Hemiptera)	*TCiCHS*	Chitin synthesis	[66]
22	Potato psyllid, *Bactericera cockerelli* (Hemiptera)	*SUC1*, *ST4*	Osmoregulatory	[67]

2.3. Perspectives and Limitations of RNAi for Insect Pest Control

Despite significant advancements in RNAi research over the past 25 years, its practical management of insect pests and disease vectors remains limited. To date, only one commercial dsRNA-based biopesticide, Calantha®, has been developed for foliar application, with several others currently under development. Additionally, three transgenic crops expressing dsRNA are commercially available.

Calantha®, a sprayable dsRNA-based biopesticide targeting the Colorado potato beetle (CPB, *Leptinotarsa decemlineata*), was released commercially by GreenLight Biosciences in 2023 [68,69]. As with dsRNA-expressing crops, sprayable formulations are likely to be used in combination with other pest control tools. However, sprayable dsRNAs degrade rapidly in the environment [70], necessitating careful consideration of application timing, spray intervals and synchronization with insect life cycle to ensure that only a single generation is

exposed to the dsRNA product. Subsequent insecticide treatments should then be rotated to a different mode of action [71,72].

Several challenges have hindered the widespread commercialization of RNAi-based products. These include variable efficacy across insect species, competition from transgenic Bt crops, and limited effectiveness against some sap-sucking pests. Another concern is the potential evolution of RNAi resistance. For example, studies have shown that western corn rootworm (*Diabrotica virgifera virgifera*) [73], willow leaf beetle (*Plagiodera versicolora*) [74], and CPB [75,76] can develop resistance to dsRNA-based control. Of note, Calantha® can be deployed in pesticide rotation programmes to mitigate selection for resistance in CPB [77]. Nonetheless, RNAi, when combined with precision agriculture and integrated pest management (IPM) strategies, could significantly enhance sustainability. Opportunities for synergy exist with advancing the technologies such as CUADb, CRISPR/Cas technology, microbial-based RNAi production, and nanoformulations for improved dsRNA delivery. Of note, RNA molecules are prone to degradation in the environment. To enhance their effectiveness, researchers are developing more stable RNA formulations and innovative delivery methods.

A clear understanding of dsRNA structure is critical for optimizing RNAi applications in pest control. This structural knowledge informs how dsRNA is processed into siRNAs and facilitates the precise targeting of pest genes for silencing [78,79]. Although the effects of dsRNA-mediated interference are highly potent and specific, there are several concerns that should be taken into account when designing RNAi-based experiments. If a sequence is shared among multiple closely related genes, RNAi may unintentionally silence several members of the gene family or homologous genes in related species [80]. Also, genes with low expression levels may exhibit resistance to RNAi, at least partially. For example, if the target protein is very stable, its depletion occurs much more slowly despite transcript degradation. Moreover, the function of a target protein can be compensated for by related proteins, which may even be upregulated by the cell to counteract the loss. Additionally, the transcription of the target gene itself could be upregulated by a regulatory gene network in response to RNAi mediated knock-down [81].

Importantly, insights into how Argonaute proteins bind to guide and target RNAs allow for the rational design of highly specific and effective RNAi tools. Structural analyses of the Argonaute active site and its interaction with target RNAs enable the optimization of siRNA sequences for enhanced gene silencing [82,83]. Furthermore, understanding the architecture of RISC components supports improvements in delivery strategies and silencing efficiency [84]. Overall, this structural knowledge is essential for developing RNAi-based pest control methods that are both species-specific and environmentally sustainable [85].

Currently, the implementation of RNAi-based pest control strategies is challenged by regulatory uncertainties and public apprehension. Gaining regulatory approval and public acceptance will depend on transparent scientific communication, rigorous safety evaluations, and evidence-based demonstrations of efficacy. A major practical limitation remains the high cost of dsRNA production; publicly accessible in vitro synthesis methods are still prohibitively expensive, exceeding $50 per milligram [86–90].

Despite these challenges, RNAi remains a promising strategy for insect pest control due to its selectively silence genes critical for insect development and survival. This specificity provides an opportunity to develop environmentally sustainable pest control solutions. However, the successful deployment of dsRNA-based insecticides will require coordinated efforts from researchers, policymakers, and industry stakeholders worldwide. Future advancements, particularly in dsRNA formulation, delivery systems, and cost-

effective production, are expected to drive the broader adoption of RNAi in the coming decades, potentially transforming agricultural pest management practices [91].

3. CUADb

3.1. History of Discovery of CUADb

Unmodified DNA, as a programmable molecule and natural biopolymer, has long captured scientific interest. However, for many years it was believed that unmodified oligodeoxyribonucleotides were both cytotoxic and highly susceptible to degradation in eukaryotic cells due to nuclease activity [92], including in insects [11,93]. Some studies explicitly stated that unmodified (phosphodiester) antisense oligonucleotides should not be employed in experimental applications [94]. Additionally, it was widely assumed that ribosomal RNA (rRNA) was inherently resistant to degradation by antisense DNA oligonucleotides [95,96]. At the turn of the century, there was no indication in the scientific literature that DNA could be used as an insecticide.

The insecticidal properties of DNA were discovered serendipitously in the spongy moth (*Lymantria dispar*). In 2007, Oberemok initiated research on the transovarial transmission of *L. dispar* multiple nucleopolyhedrovirus (LdMNPV) as part of his doctoral work. Two specific primers were selected within the anti-apoptotic gene (IAP-3) of LdMNPV: a forward primer from the sense strand (5′-GCCGGCGGAACTGGCCCA-3; oligoBIR fragment) and a reverse primer from the antisense strand (5′-CGACGTGGTGGCACGGCG-3′; oligoRING fragment) [97]. These primers generated a 317 bp amplicon in PCR using purified viral DNA [98]. However, when applied to DNA extracted from virus-free *L. dispar*, the primers produced amplicons of varying lengths, suggesting non-specificity and indicating the presence of homologous sequences in the host genome—a phenomenon previously documented for other viruses [99]. Given this, it was hypothesized that the oligoBIR and oligoRING fragments could modulate the expression of homologous *L. dispar* IAP genes, potentially inducing apoptosis. In April 2008, Oberemok tested this hypothesis unconventionally by topically applying drops of aqueous solutions of the primers to the surface of *L. dispar* larvae [4,100]. Remarkably, after 3–5 days, significant larval mortality was observed, attributable to the applied DNA fragments derived from the viral genome. This experiment marked the inception of a new class of contact DNA insecticides, termed oligonucleotide insecticides, or briefly olinscides, and the CUADb (Contact Unmodified Antisense DNA biotechnology) platform. The first results were reported in a Ukrainian patent (No. 36445) in 2008, followed by articles in Pesticide Biochemistry and Physiology [101,102]. The earliest oligonucleotide insecticides (18–20 nt) targeting anti-apoptotic genes were effective in both virus-free and LdMNPV-infected larvae [97]. Importantly, unique antisense DNA sequences of 11–20 nt offer high target selectivity, although their efficacy depends on the abundance of target RNA. Consequently, CUADb was optimized to target rRNA—which constitutes approximately 80% of total cellular RNA—via a DNA containment (DNAc) mechanism, resulting in a robust algorithm for insect pest control (Figure 3). This concept later was used by RNAi researchers. Notably, three years after Oberemok's discovery, Wang et al. [103] successfully demonstrated, for the first time, the use of dsRNA fragments as contact insecticides in pest control.




Figure 3. DNA containment (DNAc) is the main mechanism of action of oligonucleotide insecticides (olinscides): 1—antisense DNA sequences complementary to rRNA and/or pre-rRNA of pests are used as oligonucleotide insecticides; 2—formation of duplex between DNA (oligonucleotide insecticide) and RNA (rRNA and/or pre-rRNA) of the pests; 3—at first step of DNAc, antisense DNA oligonucleotide (oligonucleotide insecticide) 'arrests' target rRNA and/or pre-rRNA followed by its hypercompensation and interferes with normal functioning and biogenesis of ribosomes ('arrested' ribosomes); 4—at second step of DNAc, rRNase cleaves target rRNA and/or pre-rRNA and substantial decrease in its concentration occurs.

3.2. How CUADb Works on Insect Pests

In 2019, Oberemok and colleagues introduced several major refinements to the CUADb platform aimed at enhancing its efficacy and cost-effectiveness. A key advancement was the selection of insect ribosomal RNA (rRNA) as the primary target for oligonucleotide insecticides [11]. Targeting pre-rRNA and mature rRNA significantly enhances efficacy because these RNA molecules account for approximately 80% of total cellular RNA [104], in contrast, mRNAs, which constitute only around 5%. This strategic targeting substantially increases the signal-to-noise ratio, up to approximately 100,000:1 when compared to random mRNA sequences. Insects possess both nuclear and mitochondrial rRNA, including 28S (~3900 nucleotides (nt)), 18S (~1920 nt), 5.8S (~160 nt), and 5S (~120 nt), 16S (~1140 nt) and 12S (~600 nt) making rRNA an ideal and abundant molecular target.

A second major refinement involved reducing the length of oligonucleotide insecticides to 10–12 nt to lower production costs. Shorter sequences result in higher yields during phosphoramidite-based DNA synthesis increasing product mass per synthesis cycle. Despite their brevity, 11-nt oligonucleotides retain sufficient specificity, with a uniqueness frequency of approximately 1 in 4.19 million sequences, adequate for most agricultural systems [105]. In more complex ecosystems, such as forest, longer sequences (15–20 nt) may be used to improve selectivity [12].

A third key advancement was the identification of susceptibility to unmodified antisense oligonucleotides in insects of the suborder Sternorrhyncha (Hemiptera) [6]. Since then, CUADb-based insecticides have been successfully tested against a wide range of sap-feeding insect pests. These include 28S rRNA targets in *Unaspis euonymi*, *Dynaspidiotus britannicus*,

Icerya purchasi, *Ceroplastes japonicus*, *Aonidia lauri*, *Coccus hesperidum* [5,12,13,106–109]; 18S rRNA in *Pseudococcus viburni* [110]; and internal transcribed spacer 2 (ITS2) of pre-rRNA in *Macrosiphoniella sanborni*, *Schizolachnus pineti* [14,111], and *Trioza alacris* [112] (Table 2). ITS2-targeting oligonucleotides have also demonstrated acaricidal potential against the spider mite *Tetranychus urticae* [113,114]. Typically, a single contact treatment with oligonucleotide insecticides at 100 ng/μL results in mortality rates of approximately 80% within 3 to 14 days [12,109]. The first successful application of these insecticides within Sternorrhyncha was conducted on *U. euonymi* in 2019 [107,115].

Table 2. List of pest species successfully targeted by CUADb.

Sl. No.	Names of Model Insects	Targeted Gene(s)	Affected Processes	References
1	Euonymous scale, *U. euonymi* (Hemiptera)	*28S rRNA*	Protein biosynthesis	[107,115]
2	Holly scale, *D. britannicus* (Hemiptera)	*28S rRNA*	Protein biosynthesis	[5,107]
3	Japanese wax scale, *C. japonicus* (Hemiptera)	*28S rRNA*	Protein biosynthesis	[106]
4	Cactus scale, *Diaspis echinocacti* (Hemiptera)	*28S rRNA*	Protein biosynthesis	[116]
5	Bay sucker, *T. alacris* (Hemiptera)	*ITS2 of pre-rRNA and 28S rRNA*	Protein biosynthesis	[112]
6	Cottony cushion scale, *I. purchasi* (Hemiptera)	*28S rRNA*	Protein biosynthesis	[108]
7	Chrysanthemum aphid, *M. sanborni* (Hemiptera)	*ITS2 of pre-rRNA*	Protein biosynthesis	[111]
8	Mealybug, *P. viburni* (Hemiptera)	*5.8S, 18S and 28S rRNA*	Protein biosynthesis	[110]
9	Laureal scale, *A. lauri* (Hemiptera)	*28S rRNA*	Protein biosynthesis	[5]
10	Soft scale, *C. hesperidum* (Hemiptera)	*28S rRNA*	Protein biosynthesis	[105]
11	Two-spotted spider mite, *T. urticae* (Trombidiformes)	*ITS2 of pre-rRNA*	Protein biosynthesis	[113]
12	Grey pine aphid, *S. pineti* (Hemiptera)	*ITS2 of pre-rRNA*	Protein biosynthesis	[13]
13	Large pine aphid, *Cinara pinea* (Hemiptera)	*ITS2 of pre-rRNA*	Protein biosynthesis	[13]
14	Pine needle aphid, *Eulachnus rileyi* (Hemiptera)	*ITS2 of pre-rRNA*	Protein biosynthesis	[13]

It is important to note that studies on Sternorrhynchan insects demonstrated that unmodified oligodeoxyribonucleotides (also known as oligonucleotide insecticides or briefly olinscides) can induce both upregulation and downregulation of target genes through a mechanism known as DNA containment (DNAc). This mechanism operates in two sequential steps: the first involves the arrest of target rRNA and/or pre-rRNA (formation of DNA–rRNA duplex), leading to a functional block of ribosomes and triggering a compensatory overexpression of rRNA via rDNA transcription. The second step of DNA containment mechanism involves degradation of the arrested rRNA and/or pre-rRNA by DNA-guided rRNase [5,6,117]. In the presence of rRNase, DNA–rRNA duplex is processed

via a mechanism analogous to a zipper, referred to as the 'genetic zipper' method [112]. Our recent studies on *C. hesperidum* (4th day after contact application of oligonucleotide insecticide Coccus-11) show that almost all ribosomal proteins of 40S and 60S ribosomal subunits are also significantly upregulated during DNAc promoting the formation of new ribosomes together with hypercompensated rRNA. Ribosome biogenesis proteins involved in the formation of ribosomal subunits (NOP53, UTP30, NSA2, MAK21, BRX1, WDR12) are upregulated as well. Also during DNAc, most of ATP-dependent enzymes are downregulated (including mTOR, serine/threonine protein kinase playing crucial role in ribosome biogenesis through mTORC1 complex) while proteins from mitochondrial ATP synthase complex are upregulated indicating deficiency of cellular energy caused by oligonucleotide insecticide Coccus-11. RNase H1 is also significantly upregulated during DNAc. RNase H1 functions independently of cell cycle and cleaves RNA-DNA hybrids, including those formed between DNA and rRNA. Of note, the innovative 'genetic zipper' method not only facilitates accumulation of rRNA and its subsequent degradation as a core process during DNAc but also serves as a predictive algorithm capable of estimating the efficacy of a specific olinscide against the intended target species and closely related organisms possessing identical rRNA target sequences [14,112].

Oligonucleotide insecticides can be designed using DNAInsector program (available at dnainsector.com) or manually, based on pre-rRNA and rRNA sequences retrieved from the GenBank database. In practical terms, this means that individuals with basic sequence knowledge can design an olinscide complementary to the pre-rRNA or rRNA of a Sternorrhynchan insect with a high probability of success. However, to ensure species specificity, it is essential to compare homologous sites in non-target organisms to prevent off-target effects. Synthesis of oligonucleotide insecticides is typically performed using the phosphoramidite method, in either liquid-phase or solid-phase formats. Commonly used solid-phase synthesizers include the ASM-800 (BIOSSET, Novosibirsk, Russia), OligoPilot^{TM} (Cytiva, Uppsala, Sweden), and the 10-Column DNA Synthesizer (PolyGen, Langen, Germany) [108]. The standard preparation involves dissolving the olinscide in nuclease-free water at a concentration of 1 mg per 10 mL and applying it to approximately 1 m^2 of infested plant foliage.

3.3. Perspectives and Limitations of CUADb for Insect Pest Control

DNA-based technologies are currently transforming plant protection by enabling the development of novel control agents with advanced characteristics. DNA, being a natural biopolymer, offers enhanced compatibility with ecological systems, including agroecosystems. Its programmability allows the design of highly specific treatments, including insecticidal and acaricidal agents. Moreover, if resistance arises, adaptive strategies can be implemented by simply redesigning olinscides to the target upstream or downstream regions of the resistance-conferring site in pre-rRNA and rRNA sequences [5]. Oligonucleotide insecticides offer several distinct advantages: they exhibit a low carbon footprint, high specificity to target pests, rapid biodegradability in ecosystems, and minimal risk of target-site resistance. Furthermore, their efficacy can now be predicted across different pest species, particularly in phylogenetically related taxa with conserved target sequences [112]. Given these benefits, it is likely only a matter of time before such DNA-based products become mainstream tools in crop protection. The mechanism of action, known as the DNA containment mechanism, is well understood. The phosphoramidite synthesis method has been optimized, and delivery routes (primarily contact-based, as oral delivery remains less effective) have been established. Although cost remains a barrier, the price of DNA insecticides is expected to decrease with further technological advancements. In

fact, CUADb-based insecticides are already becoming cost-competitive with conventional chemical alternatives for some pest species [12].

For instance, CUADb has significantly reduced costs in conifer aphid management by adopting liquid-phase DNA synthesis [13,108]. Sumitomo Chemical Co., Ltd. (Tokyo, Japan), a leading developer of liquid-phase synthesis, reportedly offers 1 kg of unmodified 11-nucleotide oligonucleotides for $25,000 [13]. By contrast, synthesizing the same quantity using standard solid-phase methods, more widely used in research laboratories, can cost up to $1 million. At an application rate of 200 L per hectare with an active ingredient concentration of 0.1 mg/L (0.1 ng/mL), the cost per hectare is approximately $0.50 using liquid-phase synthesis [13]. This level of affordability enables frequent field applications. However, for many Sternorrhynchan pests, effective control requires concentrations around 0.05 g/L, raising the cost to approximately $250 per hectare, even with cost-efficient synthesis. Consequently, achieving an optimal balance between cost and efficacy remains essential for widespread adoption of nucleic acid-based insecticides. Current estimates suggest that CUADb-based approaches could be applicable to 10–15% of all insect pest species.

While oligonucleotide insecticides have demonstrated strong efficacy against hemipteran and moderate efficacy against lepidopteran pests, they have shown much lower activity against coleopterans such as *L. decemlineata* [118]. Incorporating adjuvants (e.g., spreaders, adhesives, penetrants, and UV protectants) into formulations may improve performance, though their environmental safety must be thoroughly assessed. Another critical design consideration is the potential for non-canonical base pairing, such as A:C (C:A) and G:U (T:G) interactions [119,120], which may compromise specificity and increase the risk to non-target organisms. Thus, such interactions must be carefully accounted for during olinscide development [13,15]

Although the CUAD-based 'genetic zipper' method presents a promising and cost-effective platform for DNA insecticide development, further optimization of production processes and cost-reduction strategies will be necessary for large-scale implementation. Current research is focused on increasing target specificity by designing olinscides for unique rRNA regions, thereby minimizing off-target effects. Additionally, greater insight into olinscide uptake mechanisms by insect cells is needed to enhance efficacy. In conclusion, the 'genetic zipper' holds substantial promise for advancing insect pest control. However, its commercialization will depend on overcoming challenges related to delivery efficiency, production cost, resistance management, and environmental safety. Addressing these limitations will be essential for CUADb-based oligonucleotide insecticides to become a sustainable and widely adopted component of modern pest management.

4. CRISPR/Cas

4.1. History of Discovery of CRISPR/Cas

While the development of RNA interference (RNAi) technology and the CUADb database followed a relatively linear and well-documented trajectory, the emergence of CRISPR/Cas technology was shaped by the cumulative efforts of numerous research groups. This collaborative history makes it difficult to attribute the discovery to any single individual or team. This section provides an overview of the widely accepted scientific milestones in the development of the CRISPR/Cas system, while acknowledging that many contributors may not be named individually.

In 1987, Ishino and colleagues identified a previously unknown repeat sequence in *Escherichia coli*, although its significance was not fully understood at the time [121]. In 1989, Spanish scientist Mojica, while studying the archaeal microbe *Haloferax mediterranei*, observed similar palindromic repeats of approximately 30 base pairs, separated by unique spacer sequences of roughly 36 base pairs. These did not resemble any known repeat fami-

lies in prokaryotes at the time [122]. Later Mojica and Jansen introduced the term CRISPR to describe these structures [25]. In 2002, Jansen et al. [123] identified Cas (CRISPR-associated) genes located adjacent to CRISPR loci by suggesting a functional relationship. In 2007, Horvath and colleagues provided the first experimental evidence of CRISPR-Cas system's role in adaptive immunity. They demonstrated that Streptococcus thermophilus required the Cas7 protein for spacer acquisition and that resistance to phages was mantained by the phage-derived spacer, even in the absence of Cas7, indicating its role in spacer acquisition rather than the immunity process itself [124]. Cas9, previously referred to as Cas5, Csn1, or Csx12, which contains two nuclease motifs, HNH and RuvC, and was shown to be essential for phage resistance, confirming its direct role in DNA cleavage and adaptive immunity [125–127]. In 2011, Charpentier identified a small RNA in Streptococcus pyogenes known as trans-activating CRISPR RNA (tracrRNA), which is essential for the processing of CRISPR RNAs (crRNAs) and for guiding the Cas9 complex to cleave target DNA [7]. In 2012, Šikšnys et al. published a seminal study in demontsrating that Cas9 could be programmed with custom-designed spacers to cleave specific DNA targets in vitro [24]. Independently, Charpentier and Jennifer Doudna published in Science, that Cas9 could be guided by synthetic crRNAs and tracrRNAs to cleave dsDNA, with each domains cleaving a complementary DNA strand [25]. These foundational studies established the CRISPR/Cas9 system as a robust genome-editing tool. Although Charpentier, Doudna, and Šikšnys are often credited with demonstrating the system's genome-editing potential, the technology's development is the product of decades of collaborative scientific progress.

4.2. How CRISPR/Cas Works on Insect Pests

The CRISPR/Cas9 system is the predominant tool in gene editing used for insect pest control, characterized by its simplicity and efficiency. It consists of two key components: a sgRNA (single-guide RNA) and the Cas9 nuclease. The sgRNA is an engineered fusion of CRISPR RNA (crRNA), which defines the target DNA sequence, and trans-activating CRISPR RNA (tracrRNA), which facilitates Cas9 binding. This complex directs Cas9 to specific genomic loci via base pairing. Cas9, a multi-domain DNA endonuclease derived from *Streptococcus pyogenes*, recognizes a protospacer adjacent motif (PAM) sequence—typically 5′-NGG-3′—located near the target DNA site. Upon recognition, Cas9 induces a double-stranded break (DSB) three base pairs upstream of the PAM site. The HNH and RuvC nuclease domains cleave the target and complementary DNA strands, respectively [128–132]. Cellular repair of DSBs occurs via two pathways: non-homologous end joining (NHEJ), which often results in small insertions or deletions that disrupt gene function, or homology-directed repair (HDR), which enables precise gene editing using a repair template) [132,133]. Targeting a gene's coding sequence with CRISPR/Cas9 and allowing repair via NHEJ often results in a nonfunctional protein. Alternatively, HDR allows for gene knock-ins through integration of exogenous sequences [133,134] (Figure 4).

Since its initial use in *Drosophila*, CRISPR/Cas9 has been widely adopted in diverse insect species. Genome editing aims to introduce engineered traits into wild pest populations, providing a promising strategy to manage invasive pests [135,136]. In the context of insect pests, CRISPR/Cas can be applied in several keyways (Table 3). This approach has been employed to disrupt genes involved in insecticide resistance. A comprehensive review by Xu et al. [137] utlines the application of CRISPR/Cas9 across various insect orders and editing types. Beyond resistance, CRISPR/Cas9 has also been used to manipulate genes regulating reproduction and sex determination. For example, targeting the *Astra-2* (a sex determination gene) in *Anastrepha ludens* results in sterility in males and intersexual phenotypes in females [138]. The *Doublesex (Dsx)* gene, a key regulator of sexual differentiation in insects. has also been edited in species such as diamondback, producing

gender-specific sterility through isoform disruption [139]. In addition to these applications, CRISPR/Cas9 has been used to suppress vector competence in disease-transmitting insects. By editing genes essential to pathogen transmission, researchers have successfully reduced the capacity of mosquitoes to spread diseases such as malaria, dengue, and Zika [140–142]. Moreover, the system has facilitated the development of gene drives designed to spread deleterious traits through populations, leading to sterility or reduced fitness and, ultimately, population suppression [132,143]. These gene drives bypass Mendelian inheritance through homing endonuclease activity and rely on Cas9 and guide RNAs to propagate engineered alleles during DNA repair [132,144–146]. Notably, Meccariello et al. [147] demonstrated the effectiveness of CRISPR/Cas9 gene drives in the Mediterranean fruit fly (*Ceratitis capitata*) by targeting the transformer gene to convert genetic females into fertile XX males. Similar strategies have been explored in beetles, moths and grasshoppers [148–150]. In some insect species, such as silkworms, CRISPR-based editing has proven more efficient than RNAi. For instance, knockdown of the homeobox gene *Scr* led to developmental abnormalities and malformed adult structures, showcasing the system's potential in Lepidoptera [151]. Furthermore, CRISPR/Cas9 has been used synergistically with RNAi technologies. Chloroplast-engineered dsRNA targeting *Frankliniella occidentalis* has been combined with CRISPR/Cas to enhance pest control efficacy [152,153]. In addition to pest management, the technology has significantly advanced functional studies in insects. In *S. frugiperda*, genome editing has expanded our understanding of insect physiology, development, morphology, and behavior [133].




Figure 4. CRISPR/Cas9 system used for insect pest control: 1—sgRNA-Cas9 complex formation; 2—formation of RNA-DNA duplex with target DNA; 3—cleavage of target double-stranded DNA by Cas9; 4—DNA repair through non-homologous end joining (NHEJ) or homologous-dependent repair (HDR); * crRNA—CRISPR RNA; tracrRNA—trans-activating crRNA.

Table 3. List of insect species successfully targeted by CRISPR/Cas system.

Sl. No.	Names of Model Insects	Target Gene(s)	Affected Processes	Reference
1	Mosquito, *Anopheles stephensi* (Diptera)	*Kynurenine hydroxylase*	Parasite-resistance	[154]
2	Fall armyworm, *S. frugiperda* (Lepidoptera)	*Ebony* *Doublesex (dsx) (Sfdsx)* *Antennapedia (Antp)* *Spermatogenesis-related, tssk2*	Melanin biosynthesis Sex differentiation Insect thorax and wing development Male reproductive system	[155–158]
3	Diamondback moth, *P. xylostella* (Lepidoptera)	*Yellow* *Ebony* *LW-opsin*	Body pigmentation Body pigmentation Efficiency of phototaxis	[46,159,160]
4	European bee, *Apis mellifera* (Hymenoptera)	*Amyellow-y*	Melanization in cuticle	[161]
5	Beet armyworm, *S. exigua* (Lepidoptera)	*Desaturase (SexiDES5)*	Sex pheromone biosynthesis	[162]
6	Brown planthopper, *N. lugens* (Hemiptera)	*Cysteine sulfinic acid decarboxylase (CSAD)*	Melanin metabolism	[163]
7	Chickpea pod borer, *H. armigera* (Lepidoptera)	*Wnt1*	Segmentation, appendage development, and pigmentation	[164]
8	Asian corn borer, *Ostrinia furnacalis* (Lepidoptera)	*Abdominal-A (Abd-A) and Ultrabithorax (Ubx)*	Anatomical structure formation	[165]
9	Black garden ant, *Lasius niger* (Hymenoptera)	*Cinnabar*	Eye pigmentation	[166]
10	Common cutworm, *S. litura* (Lepidoptera)	*Serine protease 2* *Odorant-binding proteins*	Male sterility Perception of a sex pheromone	[153,167]
11	Indian meal moth, *Plodia interpunctella* (Lepidoptera)	*ATP binding cassette (ABC) proteins*	Eye pigmentation	[168]
12	Eggplant shoot and fruit borer, *Leucinodes orbonalis* (Lepidoptera)	*Tryptophan 2, 3-dioxygenase* *Vitellogenin (Vg)*	Eye pigmentation Female reproductive system	[169,170]
13	Mango fruit fly, *B. dorsalis* (Diptera)	*White* *White locus* *OBP13*	Eye pigmentation Eye pigmentation Methyl eugenol	[171–174]
14	Pomace fly, *Drosophila suzukii* (Diptera)	*Doublesex*	Population suppression	[175]
15	Australian cotton bollworm, *H. armigera conferta* (Lepidoptera)	*Cadherin*	Cry1Ac resistance	[176]
16	Cricket, *Gryllus bimaculatus* (Orthoptera)	*Laccase 2 (Gb-lac2)*	Cuticle system pigmentation	[177]

4.3. Perspectives and Limitations of CRISPR/Cas for Insect Pest Control

CRISPR/Cas technology presents a promising avenue for insect pest control, offering gene-editing capabilities that may contribute to the development of sustainable pest management strategies, including gene drives and insect-resistant plants. The application of molecular genetic engineering and insect transformation using CRISPR/Cas9 across multiple insect species has helped to overcome numerous limitations associated with traditional

approaches, which largely relied on naturally occurring genetic mutations or transposable elements. CRISPR/Cas genome editing is widely favored for its efficiency, simplicity, and adaptability. In addition to Cas9, other Cas proteins, such as Cas3, Cas12a, and Cas13a, have expanded the toolbox for both natural and engineered gene editing, enabling precise modifications, base editing, prime editing, and gene regulation. Recent advancements have also facilitated DNA-free editing, allowing for genome modification without inducing double-stranded breaks [178–180].

Despite these developments, several significant challenges remain. It is currently difficult to accurately predict the outcomes of gene editing in insect pests or to determine which gene targets will yield the most effective results. Moreover, CRISPR/Cas systems may introduce unintended edits at off-target genomic sites, which can result in unforeseen biological consequences. Delivering CRISPR/Cas components into insect cells presents another major hurdle, as the efficiency of delivery methods can vary depending on species, developmental stage, or tissue type. Environmental concerns also persist regarding the use of gene drives and insect-resistant organisms, particularly in terms of their potential impacts on non-target species and broader ecosystem dynamics. Gene drives require successful mating between genetically modified and wild-type insects to facilitate the spread of edited traits; however, reproductive barriers, both prezygotic and postzygotic, may restrict gene flow and thus compromise the drive's efficiency. In addition, genetically modified insects may suffer from reduced fitness, which could limit their ability to compete with wild populations and reduce the long-term success of the strategy. The introduction of precise genetic changes through HDR is also constrained by the technical challenge of delivering sufficient quantities of donor DNA templates [181]. Achieving complex genome modifications remains technically demanding, and issues such as mosaicism, genetic variability among cells in the same organism, may further complicate phenotypic analysis and stable trait inheritance. Although population replacement strategies offer potential benefits, ensuring the long-term sustainability and ecological containment of gene drive systems is essential. Particular attention must be paid to the possibility of unintended spread into non-target species or closely related taxa [133,182]. Furthermore, the long-term effects of CRISPR/Cas-mediated genetic modifications on insect development, behavior, and ecological interactions are not yet fully understood and remain an active area of investigation [137,183].

As research in insect molecular biology continues to progress, CRISPR/Cas technology is poised to become a transformative tool in modern pest management. Its capacity for targeted, species-specific interventions offers a refined and potentially more sustainable alternative to conventional chemical insecticides. Incorporating CRISPR/Cas-based strategies into IPM programs could significantly enhance the precision and resilience of crop protection across diverse agricultural systems, especially when integrated with broader principles of synthetic biology [133].

5. Conclusions

Antisense technologies are grounded in natural mechanisms that regulate essential life processes. Complementary interactions between nucleic acids underpin key biological functions such as cell division, metabolism and defense. The development of practical tools based on these antisense mechanisms represents a highly relevant and promising area of scientific research. We believe that sustained, long-term investment in this field is warranted to achieve effective and environmentally friendly solutions for pest management. RNAi, CUADb, and CRISPR/Cas systems are powerful gene silencing tools that operate at distinct molecular levels and offer varying degrees of permanence. RNAi and CUADb transiently reduce gene expression by targeting mRNA and rRNA of pests, respectively, whereas

CRISPR/Cas induces permanent gene silencing through targeting alterations of genomic DNA. The CUADb system is unique among these technologies due to its DNA-guided mechanism and involvement in rRNA biogenesis, although it shares certain functional similarities with RNAi and CRISPR/Cas. Collectively, these systems hold considerable promise in overcoming insecticide resistance by targeting conserved genes and supporting the development of effective pest control agents.

The application of these technologies in insect pest control has already demonstrated significant potential. Notably, RNAi, CUADb and CRISPR/Cas exhibit optimal efficacy against specific insect pest groups, highlighting the potential for combining these approaches to enhance their effectiveness across broader pest populations. Although there is some level of competition among these modern antisense technologies, the physiological and genetic diversity of insect pests indicates that no single approach will provide a universal solution. Currently, antisense-based insect pest control technologies remain in the development phase. As data on their successes and limitations continue to accumulate, major companies are actively pursuing commercialization. Meanwhile, legislative framework governing the implementation of antisense technologies are gradually evolving in various countries. Although rapid breakthroughs may not be imminent, steady progress is inevitable, and will ultimately contribute to improvements in both human health and environmental sustainability.

In conclusion, RNAi, CUADb, and CRISPR/Cas represent three complementary 'worlds' of antisense technologies with promising future applications in diverse fields. These technologies hold potential for improving agriculture and developing new pest control strategies. CRISPR/Cas can create mutations for long-term changes, while RNAi and CUADb provide temporary gene silencing to block certain gene function or quickly adapt to resistance of insect pests. This combination of antisense tools offers versatility in addressing various aspects of pest control, including resistance management, population suppression, and developing novel and eco-friendly pest management strategies. However, challenges related to delivery, affordability, and environmental considerations need to be addressed for these technologies to reach their full potential.

Author Contributions: Conceptualization, V.O.; software, H.K., N.G. and V.O.; validation, V.O. and N.G.; formal analysis, V.O.; resources, V.O.; writing—original draft preparation, H.K., N.G., V.S., N.N., R.F., A.C., J.A., V.O. and K.L.; writing—review and editing, N.G., J.A. and V.O.; visualization, H.K., N.G. and V.O.; supervision, N.G. and V.O.; project administration, N.G. and V.O.; funding acquisition, V.O. All authors have read and agreed to the published version of the manuscript.

Funding: The research obtained funding from the Russian Science Foundation No. 25-16-20070, https://rscf.ru/project/25-16-20070/ (accessed on 27 Juny 2025) (Sections 1 and 3) and obtained funding within the framework of a state assignment V.I. Vernadsky Crimean Federal University for 2024 and the planning period of 2024–2026 No. FZEG-2024–0001 (Sections 2, 4 and 5).

Data Availability Statement: No new data were created.

Acknowledgments: We thank our many colleagues, too numerous to name, for the technical advances and lively discussions that prompted us to write this brief research report. We apologize to the many colleagues whose work has not been cited. Experiments were carried out at the Molecular Genetics and Biotechnologies Lab created within the framework of a state assignment V.I. Vernadsky Crimean Federal University for 2024 and the planning period of 2024–2026 No. FZEG-2024–0001. We are very much indebted to all reviewers and our colleagues from the lab on DNA technologies, PCR analysis, and creation of DNA insecticides (V.I. Vernadsky Crimean Federal University, Department of General Biology and Genetics), and OLINSCIDE BIOTECH LLC for valuable comments on our manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Minchin, S.; Lodge, J. Understanding biochemistry: Structure and function of nucleic acids. *Essays Biochem.* **2019**, *63*, 433–456. [CrossRef] [PubMed]
2. Westhof, E.; Yusupov, M.; Yusupova, G. Recognition of Watson-Crick base pairs: Constraints and limits due to geometric selection and tautomerism. *F1000Prime Rep.* **2014**, *6*, 19. [CrossRef] [PubMed]
3. Fire, A.; Xu, S.; Montgomery, M.K.; Kostas, S.A.; Driver, S.E.; Mello, C.C. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **1998**, *391*, 806–811. [CrossRef] [PubMed]
4. Oberemok, V.V. Method of Elimination of Phyllophagous Insects from Order Lepidoptera. UA Patent 36445 19 May 2008.
5. Gal'chinsky, N.V.; Yatskova, E.V.; Novikov, I.A.; Sharmagiy, A.K.; Plugatar, Y.V.; Oberemok, V.V. Mixed insect pest populations of Diaspididae species under control of oligonucleotide insecticides: 3′-end nucleotide matters. *Pestic. Biochem. Physiol.* **2024**, *200*, 105838. [CrossRef] [PubMed]
6. Oberemok, V.V.; Laikova, K.V.; Gal'chinsky, N.V. Contact unmodified antisense DNA (CUAD) biotechnology: List of pest species successfully targeted by oligonucleotide insecticides. *Front. Agron.* **2024**, *6*, 1415314. [CrossRef] [PubMed]
7. Jinek, M.; Chylinski, K.; Fonfara, I.; Hauer, M.; Doudna, J.A.; Charpentier, E.A. Programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **2012**, *337*, 816–821. [CrossRef] [PubMed]
8. Gasiunas, G.; Barrangou, R.; Horvath, P.; Siksnys, V. Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, E2579–E2586. [CrossRef] [PubMed]
9. Doudna, J.A.; Charpentier, E. Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science* **2014**, *346*, 1258096. [CrossRef] [PubMed]
10. Li, Z.; Rana, T.M. Molecular mechanisms of RNA-triggered gene silencing machineries. *Acc. Chem. Res.* **2012**, *45*, 1122–1131. [CrossRef] [PubMed]
11. Oberemok, V.V.; Laikova, K.V.; Gal'chinsky, N.V.; Useinov, R.Z.; Novikov, I.A.; Temirova, Z.Z.; Shumskykh, M.N.; Krasnodubets, A.M.; Repetskaya, A.I.; Dyadichev, V.V.; et al. DNA insecticide developed from the *Lymantria dispar* 5.8S ribosomal RNA gene provides a novel biotechnology for plant protection. *Sci. Rep.* **2019**, *9*, 6197. [CrossRef] [PubMed]
12. Oberemok, V.V.; Laikova, K.V.; Andreeva, O.A.; Gal'chinsky, N.V. Oligonucleotide insecticides and RNA-based insecticides: 16 years of experience in contact using of the next generation pest control agents. *J. Plant. Dis. Prot.* **2024**, *131*, 1837–1852. [CrossRef]
13. Oberemok, V.V.; Puzanova, Y.V.; Gal'chinsky, N.V. The 'genetic zipper' method offers a cost-effective solution for aphid control. *Front. Insect Sci.* **2024**, *4*, 1467221. [CrossRef] [PubMed]
14. Oberemok, V.V.; Gal'chinsky, N.V.; Novikov, I.A.; Sharmagiy, A.K.; Yatskova, E.V.; Laikova, E.V.; Plugatar, Y. rRNA-specific antisense DNA and dsDNA trigger rRNA biogenesis and cause potent insecticidal effect on insect pest *hesperidum* L. *biorXiv* **2024**, *10*, 618468. [CrossRef]
15. Wiles, M.V.; Qin, W.; Cheng, A.W.; Wang, H. CRISPR-Cas9-mediated genome editing and guide RNA design. *Mamm. Genome* **2015**, *26*, 501–510. [CrossRef] [PubMed]
16. Li, T.; Yang, Y.; Qi, H.; Cui, W.; Zhang, L.; Fu, X.; He, X.; Liu, M.; Li, P.-f.; Yu, T. CRISPR/Cas9 therapeutics: Progress and prospects. *Sig. Transduct. Target. Ther.* **2023**, *8*, 36. [CrossRef] [PubMed]
17. Ma, X.; Zuo, Z.; Shao, W.; Jin, Y.; Meng, Y. The expanding roles of Argonautes: RNA interference, splicing and beyond. *Brief. Funct. Genom.* **2018**, *17*, 191–197. [CrossRef] [PubMed]
18. Araújo, M.F.; Castanheira, E.M.S.; Sousa, S.F. The Buzz on Insecticides: A Review of Uses, Molecular Structures, Targets, Adverse Effects, and Alternatives. *Molecules* **2023**, *28*, 3641. [CrossRef] [PubMed]
19. Siddiqui, J.A.; Fan, R.; Naz, H.; Bamisile, B.S.; Hafeez, M.; Ghani, M.I.; Wei, Y.; Xu, Y.; Chen, X. Insights into insecticide-resistance mechanisms in invasive species: Challenges and control strategies. *Front. Physiol.* **2023**, *13*, 1112278. [CrossRef] [PubMed]
20. Gul, H.; Gadratagi, B.G.; Güncan, A.; Tyagi, S.; Ullah, F.; Desneux, N.; Liu, X. Fitness costs of resistance to insecticides in insects. *Front. Physiol.* **2023**, *14*, 1238111. [CrossRef] [PubMed]
21. Hawkins, N.J.; Bass, C.; Dixon, A.; Neve, P. The evolutionary origins of pesticide resistance. *Biol. Rev. Camb. Philos. Soc.* **2019**, *94*, 135–155. [CrossRef] [PubMed]
22. Belikova, A.; Zarytova, V.; Grineva, N. Synthesis of ribonucleosides and diribonucleoside phosphates containing 2-chloroethylamine and nitrogen mustard residues. *Tetrahedron Lett.* **1967**, *37*, 3557–3562. [CrossRef] [PubMed]
23. Zamecnik, P.C.; Stephenson, M.L. Inhibition of Rous sarcoma virus replication and cell transformation by a specific oligodeoxynucleotide. *Proc. Natl. Acad. Sci. USA* **1978**, *75*, 280–284. [CrossRef] [PubMed]
24. Lander, E.S. The Heroes of CRISPR. *Cell* **2016**, *164*, 18–28. [CrossRef] [PubMed]
25. Shmakova, A.A.; Shmakova, O.P.; Karpukhina, A.A.; Vassetzky, Y.S. CRISPR/Cas: History and Perspectives. *Russ. J. Dev. Bio.* **2022**, *53*, 272–282. [CrossRef]
26. Tomoyasu, Y.; Miller, S.C.; Tomita, S.; Schoppmeier, M.; Grossmann, D.; Bucher, G. Exploring systemic RNA interference in insects: A genome-wide survey for RNAi genes in Tribolium. *Genome Biol.* **2008**, *9*, R10. [CrossRef] [PubMed]

27. Svoboda, P. Key mechanistic principles and considerations concerning RNA interference. *Front. Plant Sci.* **2020**, *11*, 1237. [CrossRef] [PubMed]
28. Zhao, Y.; Shu, R.; Liu, J. The development and improvement of ribonucleic acid therapy strategies. *Mol. Ther. Nucleic Acids.* **2021**, *26*, 997–1013. [CrossRef] [PubMed]
29. Zhu, K.Y.; Palli, S.R. Mechanisms, applications, and challenges of insect RNA interference. *Annu. Rev. Entomol.* **2020**, *65*, 293–311. [CrossRef] [PubMed]
30. Cooper, A.M.; Silver, K.; Zhang, J.; Park, Y.; Zhu, K.Y. Molecular mechanisms influencing efficiency of RNA interference in insects. *Pest Manag. Sci.* **2019**, *75*, 18–28. [CrossRef] [PubMed]
31. Cooper, A.M.; Song, H.; Yu, Z.; Biondi, M.; Bai, J.; Shi, X.; Weerasekara, S.M.; Hua, D.H.; Silver, K.; Zhang, J.; et al. Comparison of strategies for enhancing RNA interference efficiency in *Ostrinia nubilalis*. *Pest Manag. Sci.* **2021**, *77*, 635–645. [CrossRef] [PubMed]
32. Vogel, E.; Santos, D.; Mingels, L.; Verdonckt, T.; Broeck, J.V. RNA Interference in Insects: Protecting Beneficials and Controlling Pests. *Front. Physiol.* **2019**, *9*, 1912. [CrossRef] [PubMed]
33. Girard, C.; Budin, K.; Boisnard, S.; Zhang, L.; Debuchy, R.; Zickler, D.; Espagne, E. RNAi-Related Dicer and Argonaute Proteins Play Critical Roles for Meiocyte Formation, Chromosome-Axes Lengths and Crossover Patterning in the Fungus *Sordaria macrospora*. *Front. Cell Dev. Biol.* **2021**, *9*, 684108. [CrossRef] [PubMed]
34. Silver, K.; Cooper, A.M.; Zhu, K.Y. Strategies for enhancing the efficiency of RNA interference in insects. *Pest Manag. Sci.* **2021**, *77*, 2645–2658. [CrossRef] [PubMed]
35. Terenius, O.; Papanicolaou, A.; Garbutt, J.S.; Eleftherianos, I.; Huvenne, H.; Kanginakudru, S.; Albrechtsen, M.; An, C.; Aymeric, J.L.; Barthel, A.; et al. RNA interference in Lepidoptera: An overview of successful and unsuccessful studies and implications for experimental design. *J. Insect Physiol.* **2011**, *57*, 231–245. [CrossRef] [PubMed]
36. Lucena-Leandro, V.S.; Abreu, E.F.A.; Vidal, L.A.; Torres, C.R.; Junqueira, C.I.C.V.F.; Dantas, J.; Albuquerque, É.V.S. Current Scenario of Exogenously Induced RNAi for Lepidopteran Agricultural Pest Control: From dsRNA Design to Topical Application. *Int. J. Mol. Sci.* **2022**, *23*, 15836. [CrossRef] [PubMed]
37. Christiaens, O.; Tardajos, M.G.; Martinez Reyna, Z.L.; Dash, M.; Dubruel, P.; Smagghe, G. Increased RNAi Efficacy in via the Formulation of dsRNA With Guanylated Polymers. *Front. Physiol.* **2018**, *9*, 316. [CrossRef] [PubMed]
38. Jain, R.G.; Robinson, K.E.; Asgari, S.; Mitter, N. Current scenario of RNAi-based hemipteran control. *Pest Manag. Sci.* **2021**, *77*, 2188–2196. [CrossRef] [PubMed]
39. Kaplanoglu, E.; Kolotilin, I.; Menassa, R.; Donly, C. Plastid Transformation of Micro-Tom Tomato with a Hemipteran Double-Stranded RNA Results in RNA Interference in Multiple Insect Species. *Int. J. Mol. Sci.* **2022**, *23*, 3918. [CrossRef] [PubMed]
40. Tian, H.; Peng, H.; Yao, Q.; Chen, H.; Xie, Q.; Tang, B.; Zhang, W. Developmental control of a lepidopteran pest *Spodoptera exigua* by ingestion of bacteria expressing dsRNA of a non-midgut gene. *PLoS ONE* **2009**, *4*, e6225. [CrossRef] [PubMed]
41. Zha, W.; Peng, X.; Chen, R.; Du, B.; Zhu, L.; He, G. Knockdown of midgut genes by dsRNA-transgenic plant-mediated RNA interference in the hemipteran insect *Nilaparvata lugens*. *PLoS ONE* **2011**, *6*, e20504. [CrossRef] [PubMed]
42. Khan, A.M.; Ashfaq, M.; Khan, A.A.; Naseem, M.T.; Mansoor, S. Evaluation of potential RNA-interference-target genes to control cotton mealybug, *Phenacoccus solenopsis* (Hemiptera: Pseudococcuidae). *Insect Sci.* **2018**, *25*, 778–786. [CrossRef] [PubMed]
43. Prentice, K.; Christiaens, O.; Pertry, I.; Bailey, A.; Niblett, C.; Ghislain, M.; Gheysen, G.; Smagghe, G. RNAi-based gene silencing through dsRNA injection or ingestion against the African sweet potato weevil *Cylas puncticollis* (Coleoptera: Brentidae). *Pest Manag. Sci.* **2017**, *73*, 44–52. [CrossRef] [PubMed]
44. Camargo, R.A.; Barbosa, G.O.; Possignolo, I.P.; Peres, L.E.; Lam, E.; Lima, J.E.; Figueira, A.; Marques-Souza, H. RNA interference as a gene silencing tool to control *Tuta absoluta* in tomato (*Solanum lycopersicum*). *PeerJ* **2016**, *4*, e2673. [CrossRef] [PubMed]
45. Sun, Z.; Liu, J.; Chen, Y.; Zhang, J.; Zhong, G. RNAi-mediated knockdown of α-Spectrin depresses reproductive performance in female *Bactrocera dorsalis*. *Pestic. Biochem. Physiol.* **2023**, *196*, 105611. [CrossRef] [PubMed]
46. Arya, S.K.; Singh, S.; Upadhyay, S.K.; Tiwari, V.; Saxena, G.; Verma, P.C. RNAi-based gene silencing in *Phenacoccus solenopsis* and its validation by in planta expression of a double-stranded RNA. *Pest Manag. Sci.* **2021**, *77*, 1796–1805. [CrossRef] [PubMed]
47. Chen, J.; Zhang, D.; Yao, Q.; Zhang, J.; Dong, X.; Tian, H.; Chen, J.; Zhang, W. Feeding-based RNA interference of a trehalose phosphate synthase gene in the brown planthopper, *Nilaparvata lugens*. *Insect Mol. Biol.* **2010**, *19*, 777–786. [CrossRef] [PubMed]
48. Chen, J.Z.; Jiang, Y.X.; Li, M.W.; Li, J.W.; Zha, B.H.; Yang, G. Double-stranded RNA-degrading enzymes reduce the efficiency of RNA interference in *Plutella xylostella*. *Insects* **2021**, *12*, 712. [CrossRef] [PubMed]
49. Li, N.; Xu, X.; Li, J.; Hull, J.J.; Chen, L.; Liang, G. A spray-induced gene silencing strategy for *Spodoptera frugiperda* oviposition inhibition using nanomaterial-encapsulated dsEcR. *Int. J. Biol. Macromol.* **2024**, *281*, 136503. [CrossRef] [PubMed]
50. Bera, P.; Suby, S.B.; Dixit, S.; Vijayan, V.; Kumar, N.; Sekhar, J.C.; Vadassery, J. Identification of novel target genes for RNAi mediated management of the pest, Fall Armyworm (*Spodoptera frugiperda*, JE Smith). *Crop Prot.* **2025**, *187*, 106972. [CrossRef]
51. Ma, Y.F.; Liu, T.T.; Zhao, Y.Q.; Luo, J.; Feng, H.Y.; Zhou Gong, L.L.; Zhang, M.Q.; He, Y.Y.; Hull, J.J.; Dewer, Y.; et al. RNA interference-screening of potentially lethal gene targets in the white-backed planthopper *Sogatella furcifera* via a spray-induced and nanocarrier-delivered gene silencing system. *J. Agric. Food Chem.* **2024**, *72*, 1007–1016. [CrossRef] [PubMed]

52. Li, S.; Yang, H.; Wang, Y.; Wei, L.; Lyu, J.; Shan, Z.; Zhang, X.; Fan, D. RNA Interference Reveals the Impacts of CYP6CY7 on Imidacloprid Resistance in *Aphis glycines*. *Insects* **2024**, *15*, 188. [CrossRef] [PubMed]
53. Saberi, E.; Mondal, M.; Paredes-Montero, J.R.; Nawaz, K.; Brown, J.K.; Qureshi, J.A. Optimal dsRNA Concentration for RNA Interference in Asian Citrus Psyllid. *Insects* **2024**, *15*, 58. [CrossRef] [PubMed]
54. Wang, C.X.; Bao, H.Q.; Yan, Z.C.; Wang, J.; Wang, S.; Li, Y.X. Knockdown of vitellogenin receptor based on minute insect RNA interference methods affects the initial mature egg load in the pest natural enemy *Trichogramma dendrolimi*. *Insect Sci.* **2024**, *32*, 487–500. [CrossRef] [PubMed]
55. Liu, J.; Yang, Y.; Yang, Q.; Lin, X.; Liu, Y.; Li, Z.; Swevers, L. Successful oral RNA interference efficiency in the silkworm *Bombyx mori* through nanoparticle-shielded dsRNA delivery. *J. Insect Physiol.* **2025**, *161*, 104749. [CrossRef] [PubMed]
56. Liu, J.; Chen, W.; Chen, S.; Li, S.; Swevers, L. Similarly to *BmToll9-1*, *BmToll9-2* Is a Positive Regulator of the Humoral Immune Response in the Silkworm, *Bombyx mori*. *Insects* **2024**, *15*, 1005. [CrossRef] [PubMed]
57. Liu, J.; Chen, W.; Situ, J.; Li, J.; Chen, J.; Lai, M.; Huang, F.; Li, B. BmToll9-1 Is a Positive Regulator of the Immune Response in the Silkworm *Bombyx mori*. *Insects* **2024**, *15*, 643. [CrossRef] [PubMed]
58. Liang, Y.; Wang, T.; Yang, W.; Chen, Z.; Li, Q.; Swevers, L.; Liu, J. Silencing of the immune gene BmPGRP-L4 in the midgut affects the growth of silkworm (*Bombyx mori*) larvae. *Insect Mol. Biol.* **2023**, *32*, 340–351. [CrossRef] [PubMed]
59. Darweesh, A.F.; Fahmy, I.; Ali, M.; Elwahy, A. RNA interference of cysteine protease genes for the management of whitefly (*Bemisia tabaci*) by oral route. *Egypt. J. Bot.* **2025**, *65*, 43–56. [CrossRef]
60. Shimomura, K.; Sakita, K.; Terajima, T.; Tomizawa, M. Gene Silencing of Olfactory Receptor Coreceptor by Systemic RNA Interference in *Callosobruchus maculatus*. *J. Chem. Ecol.* **2025**, *51*, 5. [CrossRef] [PubMed]
61. Guo, P.P.; Yang, X.B.; Yang, H.; Zhou, C.; Long, G.Y.; Jin, D.C. Knockdown of the β-N-acetylhexosaminidase genes by RNA interference inhibited the molting and increased the mortality of the white-backed planthopper, *Sogatella furcifera*. *Pestic. Biochem. Physiol.* **2025**, *207*, 106216.
62. Schellens, S.; Lenaerts, C.; Pérez Baca, M.D.R.; Cools, D.; Peeters, P.; Marchal, E.; Vanden Broeck, J. Knockdown of the Halloween genes Spook, Shadow and Shade influences oocyte development, egg shape, oviposition and hatching in the desert locust. *Int. J. Mol. Sci.* **2022**, *23*, 9232. [CrossRef] [PubMed]
63. Mun, S.; Young Noh, M.; Dittmer, N.T.; Muthukrishnan, S.; Kramer, K.J.; Kanost, M.R.; Arakane, Y. Cuticular protein with a low complexity sequence becomes cross-linked during insect cuticle sclerotization and is required for the adult molt. *Sci. Rep.* **2015**, *5*, 10484. [CrossRef] [PubMed]
64. Fu, D.; Dai, L.; Gao, H.; Sun, Y.; Liu, B.; Chen, H. Identification, expression patterns and RNA interference of aquaporins in *Dendroctonus armandi* (Coleoptera: Scolytinae) larvae during overwintering. *Front. Physiol.* **2019**, *10*, 967. [CrossRef] [PubMed]
65. Araujo, R.N.; Santos, A.; Pinto, F.S.; Gontijo, N.F.; Lehane, M.J.; Pereira, M.H. RNA interference of the salivary gland nitrophorin 2 in the triatomine bug *Rhodnius prolixus* (Hemiptera: Reduviidae) by dsRNA ingestion or injection. *Insect Biochem. Mol. Biol.* **2006**, *36*, 683–693. [CrossRef] [PubMed]
66. Shang, F.; Xiong, Y.; Xia, W.K.; Wei, D.D.; Wei, D.; Wang, J.J. Identification, characterization and functional analysis of a chitin synthase gene in the brown citrus aphid, *Toxoptera citricida* (Hemiptera, Aphididae). *Insect Mol. Biol.* **2016**, *25*, 422–430. [CrossRef] [PubMed]
67. Lu, J.; Shen, J. Target genes for RNAi in pest control: A comprehensive overview. *Entomol. Gen.* **2024**, *44*, 95–114. [CrossRef]
68. Pallis, S.; Alyokhin, A.; Manley, B.; Rodrigues, T.; Barnes, E.; Narva, K. Effects of Low Doses of a Novel dsRNA-based Biopesticide (Calantha) on the Colorado Potato Beetle. *J. Econ. Entomol.* **2023**, *116*, 456–461. [CrossRef] [PubMed]
69. GreenLight Biosciences. Available online: https://www.greenlightbiosciences.com/calantha-for-cpb (accessed on 19 July 2025).
70. Bachman, P.; Fischer, J.; Song, Z.; Urbanczyk-Wochniak, E.; Watson, G. Environmental fate and dissipation of applied dsRNA in soil, aquatic systems, and plants. *Front. Plant Sci.* **2020**, *11*, 21. [CrossRef] [PubMed]
71. Sparks, T.C.; Nauen, R. IRAC: Mode of action classification and insecticide resistance management. *Pestic. Biochem. Physiol.* **2015**, *121*, 122–128. [CrossRef] [PubMed]
72. Sparks, T.C.; Storer, N.; Porter, A.; Slater, R.; Nauen, R. Insecticide resistance management and industry: The origins and evolution of the insecticide resistance action committee (IRAC) and the mode of action classification scheme. *Pest Manag. Sci.* **2021**, *77*, 2609–2619. [CrossRef] [PubMed]
73. Khajuria, C.; Ivashuta, S.; Wiggins, E.; Flagel, L.; Moar, W.; Pleau, M.; Miller, K.; Zhang, Y.; Ramaseshadri, P.; Jiang, C. Development and characterization of the first dsRNA-resistant insect population from western corn rootworm, *Diabrotica virgifera virgifera* LeConte. *PLoS ONE* **2018**, *13*, e0197059. [CrossRef] [PubMed]
74. Liao, C.; Zhang, M.; Zhang, J. Characterization and potential mechanism of resistance to double-stranded RNA in willow leaf beetle, *Plagiodera versicolora*. *J. Pest Sci.* **2024**, *97*, 2217–2226. [CrossRef]
75. Mishra, S.; Dee, J.; Moar, W.; Dufner-Beattie, J.; Baum, J.; Dias, N.P.; Alyokhin, A.; Buzza, A.; Rondon, S.I.; Clough, M.; et al. Selection for high levels of resistance to double-stranded RNA (dsRNA) in Colorado potato beetle (*Leptinotarsa decemlineata* Say) using non-transgenic foliar delivery. *Sci. Rep.* **2021**, *11*, 6523. [CrossRef] [PubMed]

76. Mishra, S.; Moar, W.; Jurat-Fuentes, J.L. Larvae of Colorado potato beetle (*Leptinotarsa decemlineata* Say) resistant to double stranded RNA (dsRNA) remain susceptible to small-molecule pesticides. *Pest Manag. Sci.* **2024**, *80*, 905–909. [CrossRef] [PubMed]
77. Narva, K.; Toprak, U.; Alyokhin, A.; Groves, R.; Jurat-Fuentes, J.L.; Moar, W.; Nauen, R.; Whipple, S.; Head, G. Insecticide resistance management scenarios differ for RNA-based sprays and traits. *Insect Mol. Biol.* **2025**, *34*, 518–526. [CrossRef] [PubMed]
78. Christiaens, O.; Whyard, S.; Vélez, A.M.; Smagghe, G. Double-Stranded RNA Technology to Control Insect Pests: Current Status and Challenges. *Front. Plant Sci.* **2020**, *11*, 451. [CrossRef] [PubMed]
79. Cedden, D.; Güney, G.; Rostás, M.; Bucher, G. Optimizing dsRNA sequences for RNAi in pest control and research with the dsRIP web platform. *BMC Biol.* **2025**, *23*, 114. [CrossRef] [PubMed]
80. Chen, J.; Peng, Y.; Zhang, H.; Wang, K.; Zhao, C.; Zhu, G.; Reddy Palli, S.; Han, Z. Off-target effects of RNAi correlate with the mismatch rate between dsRNA and non-target mRNA. *RNA Biol.* **2021**, *18*, 1747–1759. [CrossRef] [PubMed]
81. Cedden, D.; Bucher, G. The quest for the best target genes for RNAi-mediated pest control. *Insect Mol. Biol.* **2024**, *34*, 505–517. [CrossRef] [PubMed]
82. Yang, J.; Cho, W.C.; Zheng, Y. Argonaute proteins: Structural features, functions and emerging roles. *J. Adv. Res.* **2020**, *24*, 317–324. [CrossRef] [PubMed]
83. Höck, J.; Meister, G. The Argonaute protein family. *Genome Biol.* **2008**, *9*, 210. [CrossRef] [PubMed]
84. Okamura, K.; Ishizuka, A.; Siomi, H.; Siomi, M.C. Distinct roles for Argonaute proteins in small RNA-directed RNA cleavage pathways. *Genes Dev.* **2004**, *18*, 1655–1666. [CrossRef] [PubMed]
85. Hoang, B.T.L.; Fletcher, S.J.; Brosnan, C.A.; Ghodke, A.B.; Manzie, N.; Mitter, N. RNAi as a Foliar Spray: Efficiency and Challenges to Field Applications. *Int. J. Mol. Sci.* **2022**, *23*, 6639. [CrossRef] [PubMed]
86. Socha, W.; Kwasnik, M.; Larska, M.; Rola, J.; Rozek, W. Vector-borne viral diseases as a current threat for human and animal health—One Health perspective. *J. Clin. Med.* **2022**, *11*, 3026. [CrossRef] [PubMed]
87. Yang, W.; Wang, B.; Lei, G.; Chen, G.; Liu, D. Advances in nanocarriers to improve the stability of dsRNA in the environment. *Front. Bioeng. Biotechnol.* **2022**, *10*, 974646. [CrossRef] [PubMed]
88. Nitnavare, R.B.; Bhattacharya, J.; Singh, S.; Kour, A.; Hawkesford, M.J.; Arora, N. Next generation dsRNA-based insect control: Success so far and challenges. *Front. Plant Sci.* **2021**, *12*, 673576. [CrossRef] [PubMed]
89. Rank, A.P.; Koch, A. Lab-to-Field Transition of RNA Spray Applications—How Far Are We? *Front. Plant Sci.* **2021**, *12*, 755203. [CrossRef] [PubMed]
90. Verdonckt, T.W.; Vanden Broeck, J. Methods for the Cost-Effective Production of Bacteria-Derived Double-Stranded RNA for in vitro Knockdown Studies. *Front. Physiol.* **2022**, *13*, 836106. [CrossRef] [PubMed]
91. Palli, S.R. RNAi turns 25: Contributions and challenges in insect science. *Front. Insect Sci.* **2023**, *3*, 1209478. [CrossRef] [PubMed]
92. Egli, M.; Manoharan, M. Chemistry, structure and function of approved oligonucleotide therapeutics. *Nucleic Acids Res.* **2023**, *51*, 2529–2573. [CrossRef] [PubMed]
93. Peng, Y.; Wang, K.; Fu, W.; Sheng, C.; Han, Z. Biochemical comparison of dsRNA degrading nucleases in four different insects. *Front. Physiol.* **2018**, *9*, 624. [CrossRef] [PubMed]
94. Dias, N.; Stein, C.A. Antisense oligonucleotides: Basic concepts and mechanisms. *Mol. Cancer Ther.* **2002**, *1*, 347–355. [PubMed]
95. Bachellerie, J.P.; Cavaillé, J.; Hüttenhofer, A. The expanding snoRNA world. *Biochimie* **2002**, *84*, 775–790. [CrossRef] [PubMed]
96. Will, C.L.; Lührmann, R. Spliceosomal UsnRNP biogenesis, structure and function. *Curr. Opin. Cell Biol.* **2001**, *13*, 290–301. [CrossRef] [PubMed]
97. Oberemok, V.V.; Laikova, K.V.; Zaitsev, A.S.; Shumskykh, M.N.; Kasich, I.N.; Gal'chinsky, N.V.; Bekirova, V.V.; Makarov, V.V.; Agranovsky, A.A.; Gushchin, V.A.; et al. Molecular Alliance of *Lymantria dispar* Multiple Nucleopolyhedrovirus and a Short Unmodified Antisense Oligonucleotide of Its Anti-Apoptotic IAP-3 Gene: A Novel Approach for Gypsy Moth Control. *Int. J. Mol. Sci.* **2017**, *18*, 2446. [CrossRef] [PubMed]
98. Oberemok, V.V. DNA Markers in the Study of the Relationship Between Nuclear Polyhedrosis Virus and Its Host Lymantria dispar. Ph.D. Thesis, Taras Shevchenko National University of Kyiv, Kyiv, Ukraine, 2011; 22p.
99. Cerio, R.J.; Vandergaast, R.; Friesen, P.D. Host insect inhibitor-of-apoptosis SfIAP functionally replaces baculovirus IAP but is differentially regulated by its N-terminal leader. *J. Virol.* **2010**, *84*, 11448–11460. [CrossRef] [PubMed]
100. Manju, M.; Nirosha, V.; Tullika, T.; Mankhanniang, G. DNA insecticides: An emerging tool in pest management. *Agriallis* **2022**, *4*, 29–35.
101. Oberemok, V.V.; Skorokhod, O.A. Single-stranded DNA fragments of insect-specific nuclear polyhedrosis virus act as selective DNA insecticides for gypsy moth control. *Pestic. Biochem. Physiol.* **2014**, *113*, 1–7. [CrossRef] [PubMed]
102. Oberemok, V.V.; Laikova, K.V.; Zaitsev, A.S.; Gushchin, V.A.; Skorokhod, O.A. The RING for gypsy moth control: Topical application of fragment of its nuclear polyhedrosis virus anti-apoptosis gene as insecticide. *Pestic. Biochem. Physiol.* **2016**, *131*, 32–39. [CrossRef] [PubMed]
103. Wang, Y.; Zhang, H.; Li, H.; Miao, X. Second-generation sequencing supply an effective way to screen RNAi targets in large scale for potential application in pest insect control. *PLoS ONE* **2011**, *6*, e18644. [CrossRef] [PubMed]

104. Warner, J.R. The economics of ribosome biosynthesis in yeast. *Trends Biochem. Sci.* **1999**, *24*, 437–440. [CrossRef] [PubMed]
105. Oberemok, V.V.; Useinov, R.Z.; Skorokhod, O.A.; Gal'chinsky, N.V.; Novikov, I.A.; Makalish, T.P.; Yatskova, E.V.; Sharmagiy, A.K.; Golovkin, I.O.; Gninenko, Y.I.; et al. Oligonucleotide insecticides for green agriculture: Regulatory role of contact DNA in plant-insect interactions. *Int. J. Mol. Sci.* **2022**, *23*, 15681. [CrossRef] [PubMed]
106. Useinov, R.Z.; Galchinsky, N.; Yatskova, E.; Novikov, I.; Puzanova, Y.; Trikoz, N.; Sharmagiy, A.; Plugatar, Y.; Laikova, K.; Oberemok, V. To bee or not to bee: Creating DNA insecticides to replace non-selective organophosphate insecticides for use against the soft scale insect *Ceroplastes Japonicus* green. *J. Plant. Prot. Res.* **2020**, *60*, 406–409.
107. Gal'chinsky, N.; Useinov, R.; Yatskova, E.; Laikova, K.; Novikov, I.; Gorlov, M.; Trikoz, N.; Sharmagiy, A.; Plugatar, Y.; Oberemok, V. A breakthrough in the efficiency of contact DNA insecticides: Rapid high mortality rates in the sap-sucking insects *Dynaspidiotus britannicus* Comstock and *Unaspis euonymi* Newstead. *J. Plant. Prot. Res.* **2020**, *60*, 220–223. [CrossRef]
108. Gal'chinsky, N.V.; Yatskova, E.V.; Novikov, I.A.; Useinov, R.Z.; Kouakou, N.J.; Kouame, K.F.; Kra, K.D.; Sharmagiy, A.K.; Plugatar, Y.V.; Laikova, K.V.; et al. *Icerya purchasi* Maskell (Hemiptera: Monophlebidae) control using low carbon footprint oligonucleotide insecticides. *Int. J. Mol. Sci.* **2023**, *24*, 11650. [CrossRef] [PubMed]
109. Oberemok, V.V.; Gal'chinsky, N.V.; Useinov, R.Z.; Novikov, I.A.; Puzanova, Y.V.; Filatov, R.I.; Kouakou, N.J.; Kouame, K.F.; Kra, K.D.; Laikova, K.V. Four Most Pathogenic Superfamilies of Insect Pests of Suborder Sternorrhyncha: Invisible Superplunderers of Plant Vitality. *Insects* **2023**, *14*, 462. [CrossRef] [PubMed]
110. Novikov, A.; Yatskova, E.; Bily, A.; Puzanova, Y.; Sharmagiy, A.; Oberemok, V. Efficient control of the obscure mealybug *Pseudococcus viburni* with DNA insecticides. In *Vitro Cellular & Developmental Biology-Animal*; Springer: New York, NY, USA, 2023; pp. 92–108.
111. Puzanova, Y.V.; Novikov, I.A.; Bilyk, A.I.; Sharmagiy, A.K.; Plugatar, Y.V.; Oberemok, V.V. Perfect complementarity mechanism for aphid control: Oligonucleotide insecticide macsan-11 selectively causes high mortality rate for *Macrosiphoniella sanborni* Gillette. *Int. J. Mol. Sci.* **2023**, *24*, 11690. [CrossRef] [PubMed]
112. Oberemok, V.V.; Novikov, I.A.; Yatskova, E.V.; Bilyk, A.I.; Sharmagiy, A.K.; Gal'chinsky, N.V. Potent and selective 'genetic zipper' method for DNA-programmable plant protection: Innovative oligonucleotide insecticides against *Trioza alacris* Flor. *Chem. Biol. Technol. Agric.* **2024**, *11*, 144. [CrossRef]
113. Gavrilova, D.; Grizanova, E.; Novikov, I.; Laikova, E.; Zenkova, A.; Oberemok, V.; Dubovskiy, I. Antisense DNA acaricide targeting pre-rRNA of two-spotted spider mite *Tetranychus urticae* as efficacy-enhancing agent of fungus *Metarhizium robertsii*. *J. Invertebr. Pathol.* **2025**, *211*, 108297. [CrossRef] [PubMed]
114. IZ. Available online: https://en.iz.ru/en/1870413/maria-neduk-denis-gricenko/agrarian-evolution-dna-drug-will-destroy-main-enemy-greenhouse-crops (accessed on 16 April 2025).
115. Oberemok, V.V.; Laikova, K.V.; Useinov, R.Z.; Gal'chinsky, N.V.; Novikov, I.A.; Gorlov, M.V.; Balykina, E.V.; Trikoz, N.N.; Yatskova, E.V.; Sharmagiy, A.K. High mortality of sap-sucking insects one week after topical application of DNA insecticides. *Vitr. Cell. Dev. Biol. Anim.* **2020**, *56*, 31–39.
116. Plugatar, Y.V.; Chichkanova, E.S.; Yatskova, E.V.; Sharmagii, A.K.; Oberemok, V.V. An innovative method of *Diaspis echinocacti* Bouche control using DNA insecticide on *Opuntia ficus-indica* (L.) Mill. *in the Nikitsky Botanical Garden, Crimea. South Russ. Ecol. Dev.* **2021**, *16*, 119–128. [CrossRef]
117. Oberemok, V.V.; Gal'chinsky, N.V. Oligonucleotide insecticides (contact unmodified antisense DNA biotechnology) and RNA biocontrols (double-stranded RNA technology): Newly born fraternal twins in plant protection. *bioRxiv* **2024**, *3*, 584797. [CrossRef]
118. Oberemok, V.; Laikova, K.; Shumskykh, M.; Kenyo, I.; Kasich, I.; Deri, K.; Seidosmanova, E.; Krasnodubets, A.; Bekirova, V.; Gal'chinsky, N. A primary attempt of *Leptinotarsa decemlineata* control using contact DNA insecticide based on short antisense oligonucleotide of its CYP6B gene. *J. Plant Prot. Res.* **2018**, *58*, 106–108. [CrossRef]
119. Du, Q.; Thonberg, H.; Wang, J.; Wahlestedt, C.; Liang, Z.A. A systematic analysis of the silencing effects of an active siRNA at all single-nucleotide mismatched target sites. *Nucleic Acids Res.* **2005**, *33*, 1671–1677. [CrossRef] [PubMed]
120. Luige, O.; Karalė, K.; Bose, P.P.; Bollmark, M.; Tedebark, U.; Murtola, M.; Strömberg, R. Influence of sequence variation on the RNA cleavage activity of Zn^{2+}-dimethyl-dppz-PNA-based artificial enzymes. *RSC Adv.* **2022**, *12*, 5398–5406. [CrossRef] [PubMed]
121. Ishino, Y.; Shinagawa, H.; Makino, K.; Amemura, M.; Nakata, A. Nucleotide sequence of the iap gene, responsible for alkaline phosphatase isozyme conversion in Escherichia coli, and identification of the gene product. *J. Bacteriol.* **1987**, *169*, 5429–5433. [CrossRef] [PubMed]
122. Mojica, M.J.; Juez, G.; Rodríguez-Valera, F. Transcription at different salinities of *Haloferax mediterranei* sequences adjacent to partially modified PstI sites. *Mol. Microbiol.* **1993**, *9*, 613–621. [CrossRef] [PubMed]
123. Jansen, R.; Embden, J.D.A.; Gaastra, W.; Schouls, L.M. Identification of genes that are associated with DNA repeats in prokaryotes. *Mol. Microbiol.* 2002; 43, 1565–1575.
124. Barrangou, R.; Fremaux, C.; Deveau, H.; Richards, M.; Boyaval, P.; Moineau, S.; Romero, D.A.; Horvath, P. CRISPR provides acquired resistance against viruses in prokaryotes. *Science* **2007**, *315*, 1709–1712. [CrossRef] [PubMed]

125. Bolotin, A.; Quinquis, B.; Sorokin, A.; Ehrlich, S.D. Clustered regularly interspaced short palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiology* **2005**, *151*, 2551–2561. [CrossRef] [PubMed]
126. Makarova, K.S.; Grishin, N.V.; Shabalina, S.A.; Wolf, Y.I.; Koonin, E.V. A putative RNA-interference-based immune system in prokaryotes: Computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action. *Biol. Direct.* **2006**, *1*, 7. [CrossRef] [PubMed]
127. Aljabali, A.A.A.; El-Tanani, M.; Tambuwala, M.M. Principles of CRISPR-Cas9 technology: Advancements in genome editing and emerging trends in drug delivery. *J. Drug Deliv. Sci. Technol.* **2024**, *92*, 105338. [CrossRef]
128. Asokan, R.; Rai, A.; Dash, S.; Manamohan, M.; Ashok, K.; Bhargava, C.N.; Wishard, R.; Pradhan, S.K.; Parvathy, M.S. Application of genome editing in entomology. *Indian J. Entomol.* **2022**, *84*, 6–103.
129. Khan, M.D.; Ahmad, B.; Ahmed, S.F.; Ijaz, M.; Abdin, Z.U.; Ghafoor, I.; Siddiqui, G.M.; Adrees, T.; Ali, S.; Shahzadi, N.; et al. CRISPR-Cas genome editing in crops: A promising frontier for climate-smart agriculture. *Phytopathogenom. Dis. Control.* **2023**, *2*, 71–77. [CrossRef]
130. Wang, J.Y.; Doudna, J.A. CRISPR technology: A decade of genome editing is only the beginning. *Science* **2023**, *379*, eadd8643. [CrossRef] [PubMed]
131. Wang, J.Y.; Pausch, P.; Doudna, J.A. Structural biology of CRISPR–Cas immunity and genome editing enzymes. *Nat. Rev. Microbiol.* **2022**, *20*, 641–656. [CrossRef] [PubMed]
132. Zhang, F. Development of CRISPR-Cas systems for genome editing and beyond. *Q. Rev. Biophys.* **2019**, *52*, e6. [CrossRef]
133. Salum, Y.M.; Yin, A.; Zaheer, U.; Liu, Y.; Guo, Y.; He, W. CRISPR/Cas9-Based Genome Editing of Fall Armyworm (*Spodoptera frugiperda*): Progress and Prospects. *Biomolecules* **2024**, *14*, 1074. [CrossRef] [PubMed]
134. Yan, Y.; Aumann, R.A.; Häcker, I.; Schetelig, M.F. CRISPR-based genetic control strategies for insect pests. *J. Integr. Agric.* **2023**, *22*, 651–668. [CrossRef]
135. Ranian, K.; Zahoor, M.K.; Zulhussnain, M.; Ahmad, A. CRISPR/Cas9 mediated sex-ratio distortion by sex specific gene editing in *Aedes aegypti*. *Saudi J. Biol. Sci.* **2022**, *29*, 3015–3022. [CrossRef] [PubMed]
136. Zulhussnain, M.; Zahoor, M.K.; Ranian, K.; Ahmad, A.; Jabeen, F. CRISPR Cas9 mediated knockout of sex determination pathway genes in *Aedes aegypti*. *Bull. Entomol. Res.* **2023**, *113*, 243–252. [CrossRef] [PubMed]
137. Xu, Q.; Wang, M.; Zeng, J.; Sun, H.; Wei, X.; Jiang, H.; Shentu, X.; Sun, D. CRISPR/Cas Technology in Insect Insecticide Resistance. *Insects* **2025**, *16*, 345. [CrossRef] [PubMed]
138. Li, J.; Handler, A.M. CRISPR/Cas9-mediated gene editing in an exogenous transgene and an endogenous sex determination gene in the Caribbean fruit fly, *Anastrepha suspensa*. *Gene* **2019**, *691*, 160–166. [CrossRef] [PubMed]
139. Wang, Y.; Chen, X.E.; Liu, Z.; Xu, J.; Li, X.; Bi, H.; Andongma, A.A.; Niu, C.; Huang, Y. Mutation of doublesex induces sex-specific sterility of the diamondback moth *Plutella xylostella*. *Insect Biochem. Mol. Biol.* **2019**, *112*, 103180. [CrossRef] [PubMed]
140. Nourani, L.; Mehrizi, A.A.; Pirahmadi, S.; Pourhashem, Z.; Asadollahi, E.; Jahangiri, B. CRISPR/Cas advancements for genome editing, diagnosis, therapeutics, and vaccine development for Plasmodium parasites, and genetic engineering of Anopheles mosquito vector. *Infect. Genet. Evol.* **2023**, *109*, 105419. [CrossRef] [PubMed]
141. Zamanian, M.; Andersen, E.C. Prospects and challenges of CRISPR/Cas genome editing for the study and control of neglected vector-borne nematode diseases. *FEBS J.* **2016**, *283*, 3204–3221. [CrossRef] [PubMed]
142. Bimal Kumar Sahoo, B.M.; Mahesh Pathak, M.; Sushruta Boruah, S.; Kasturi Sarmah, K. Bio-engineered defenders: Crispr-cas9 unleashing disease-resistant insects to curb vector-borne diseases. In *Futuristic Trends in Agriculture Engineering & Food Sciences*; IIP Series: Novi, MI, USA, 2024; Volume 3. [CrossRef]
143. Xu, J.; Xu, X.; Zhan, S.; Huang, Y. Genome editing in insects: Current status and challenges. *Natl. Sci. Rev.* **2019**, *6*, 399–401. [CrossRef] [PubMed]
144. Courtier-Orgogozo, V.; Morizot, B.; Boëte, C. Using CRISPR-based gene drive for agriculture pest control. *EMBO Rep.* **2017**, *18*, 1481. [CrossRef] [PubMed]
145. Wang, G.H.; Du, J.; Chu, C.Y.; Madhav, M.; Hughes, G.L.; Champer, J. Symbionts and gene drive: Two strategies to combat vector-borne disease. *Trends Genet.* **2022**, *38*, 708–723. [CrossRef] [PubMed]
146. Zhao, Y.; Li, L.; Wei, L.; Wang, Y.; Han, Z. Advancements and Future Prospects of CRISPR-Cas-Based Population Replacement Strategies in Insect Pest Management. *Insects* **2024**, *15*, 653. [CrossRef] [PubMed]
147. Meccariello, A.; Hou, S.; Davydova, S.; Fawcett, J.D.; Siddall, A.; Leftwich, P.T.; Krsticevic, F.; Papathanos, P.A.; Windbichler, N. Gene drive and genetic sex conversion in the global agricultural pest *Ceratitis capitata*. *Nat. Commun.* **2024**, *15*, 372. [CrossRef] [PubMed]
148. Oye, K.A.; Esvelt, K.; Appleton, E.; Catteruccia, F.; Church, G.; Kuiken, T.; Lightfoot, S.B.Y.; McNamara, J.; Smidler, A.; Collins, J.P. Regulating gene drives. *Science* **2014**, *345*, 626–628. [CrossRef] [PubMed]
149. Shukla, J.N.; Palli, S.R. *Tribolium castaneum* Transformer-2 regulates sex determination and development in both males and females. *Insect Biochem. Mol. Biol.* **2013**, *43*, 1125–1132. [CrossRef] [PubMed]

150. Scott, M.J.; Gould, F.; Lorenzen, M.; Grubbs, N.; Edwards, O.; O'Brochta, D. Agricultural production: Assessment of the potential use of Cas9-mediated gene drive systems for agricultural pest control. *J. Responsible Innov.* **2018**, *5* (Suppl. 1), S98–S120. [CrossRef]
151. Tang, Y.H.; Zhang, X.; Dai, Z.C.; Li, H.; Yang, Y.; Zhao, T.J.; Yuan, D.Q.; Qian, W.L.; Cheng, D.J. CRISPR-Cas13-mediated RNA editing in the silkworm *Bombyx mori*. *Zool. Res.* **2024**, *45*, 1249–1260. [CrossRef] [PubMed]
152. Bulle, M.; Sheri, V.; Aileni, M.; Zhang, B. Chloroplast Genome Engineering: A Plausible Approach to Combat Chili Thrips and Other Agronomic Insect Pests of Crops. *Plants* **2023**, *12*, 3448. [CrossRef] [PubMed]
153. Han, W.K.; Yang, Y.L.; Si, Y.X.; Wei, Z.Q.; Liu, S.R.; Liu, X.L.; Yan, Q.; Dong, S.L. Involvement of GOBP2 in the perception of a sex pheromone component in both larval and adult *Spodoptera litura* revealed using CRISPR/Cas9 mutagenesis. *Insect Biochem. Mol. Biol.* **2022**, *141*, 103719. [CrossRef] [PubMed]
154. Gantz, V.M.; Jasinskiene, N.; Tatarenkova, O.; Fazekas, A.; Macias, V.M.; Bier, E.; James, A.A. Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito *Anopheles stephensi*. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E6736–E6743. [CrossRef] [PubMed]
155. Zhu, G.H.; Chereddy, S.C.; Howell, J.L.; Palli, S.R. Genome editing in the fall armyworm, *Spodoptera frugiperda*: Multiple sgRNA/Cas9 method for identification of knockouts in one generation. *Insect Biochem. Mol. Biol.* **2020**, *122*, 103373. [CrossRef] [PubMed]
156. Gu, J.; Wang, J.; Bi, H.; Li, X.; Merchant, A.; Zhang, P.; Zhang, Q.; Zhou, X. CRISPR/Cas9-mediated mutagenesis of sex-specific doublesex splicing variants leads to sterility in *Spodoptera frugiperda*, a global invasive pest. *Cells* **2022**, *11*, 3557. [CrossRef] [PubMed]
157. Wang, C.; Zhao, T.; Liu, X.; Li, T.; He, L.; Wang, Q.; Wang, L.; Zhou, L. CRISPR/Cas9-Mediated Mutagenesis of Antennapedia in *Spodoptera frugiperda*. *Insects* **2023**, *15*, 16. [CrossRef] [PubMed]
158. Anu, C.N.; Ashok, K.; Bhargava, C.N.; Dhawane, Y.; Manamohan, M.; Jha, G.K.; Asokan, R. CRISPR/Cas9 mediated validation of spermatogenesis-related gene, tssk2 as a component of genetic pest management of fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae). *Arch. Insect Biochem. Physiol.* **2024**, *116*, e22121. [CrossRef] [PubMed]
159. Wang, X.; Ma, Y.; Wang, F.; Yang, Y.; Wu, S.; Wu, Y. Disruption of nicotinic acetylcholine receptor α6 mediated by CRISPR/Cas9 confers resistance to spinosyns in *Plutella xylostella*. *Pest Manag. Sci.* **2020**, *76*, 1618–1625. [CrossRef] [PubMed]
160. Xu, X.; Harvey-Samuel, T.; Yang, J.; You, M.; Alphey, L. CRISPR/Cas9-based functional characterization of the pigmentation gene ebony in *Plutella xylostella*. *Insect Mol. Biol.* **2021**, *30*, 615–623. [CrossRef] [PubMed]
161. Nie, H.Y.; Liang, L.Q.; Li, Q.F.; Zhu, Y.N.; Guo, Y.K.; Zheng, Q.L.; Lin, Y.; Yang, D.L.; Li, Z.G.; Su, S.K. CRISPR/Cas9 mediated knockout of Amyellow-y gene results in melanization defect of the cuticle in adult *Apis mellifera*. *J. Insect Physiol.* **2021**, *132*, 104264. [CrossRef] [PubMed]
162. Ahmed, S.; Roy, M.C.; Al Baki, M.A.; Jung, J.K.; Lee, D.; Kim, Y. CRISPR/Cas9 mutagenesis against sex pheromone biosynthesis leads to loss of female attractiveness in *Spodoptera exigua*, an insect pest. *PLoS ONE* **2021**, *16*, e0259322. [CrossRef] [PubMed]
163. Chen, J.X.; Li, W.X.; Lyu, J.; Hu, Y.T.; Huang, G.; Zhang, W.Q. CRISPR/Cas9-mediated knockout of the NlCSAD gene results in darker cuticle pigmentation and a reduction in female fecundity in *Nilaparvata lugens* (Hemiptera: Delphacidae). *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **2021**, *256*, 110921. [CrossRef] [PubMed]
164. Fu, X.; Li, R.; Qiu, Q.; Wang, M.; Zhao, T.; Zhou, L. Study on the function of *Helicoverpa armigera* Wnt1 gene using CRISPR/Cas9 system. *J. Asia-Pac. Entomol.* **2022**, *25*, 101869. [CrossRef]
165. Bi, H.; Merchant, A.; Gu, J.; Li, X.; Zhou, X.; Zhang, Q. CRISPR/Cas9-mediated mutagenesis of abdominal-A and ultrabithorax in the Asian corn borer, *Ostrinia furnacalis*. *Insects* **2022**, *13*, 384. [CrossRef] [PubMed]
166. Konu, M.; Kulmuni, J.; Viljakainen, L. Genetic modification of the ant *Lasius niger* using CRISPR-Cas9 technology. *Insect Mol. Biol.* **2023**, *32*, 11–25. [CrossRef] [PubMed]
167. Bi, H.; Xu, X.; Li, X.; Wang, Y.; Zhou, S.; Huang, Y. CRISPR/Cas9-mediated Serine protease 2 disruption induces male sterility in *Spodoptera litura*. *Front. Physiol.* **2022**, *13*, 931824. [CrossRef] [PubMed]
168. Shirk, B.D.; Shirk, P.D.; Furlong, R.B.; Scully, E.D.; Wu, K.; Siegfried, B.D. Gene editing of the ABC Transporter/White locus using CRISPR/Cas9-mediated mutagenesis in the Indian Meal Moth. *J. Insect Physiol.* **2023**, *145*, 104471. [CrossRef] [PubMed]
169. Ashok, K.; Bhargava, C.N.; Babu, K.P.; Rohan, W.; Manamohan, M.; Rai, A.; Sanjay, K.P.; Parvathy, M.S.; Kennedy, J.S.; Asokan, R. First report on CRISPR/Cas9 mediated editing of the eye colour gene, tryptophan 2, 3-dioxygenase in egg plant shoot and fruit borer *Leucinodes orbonalis* Guenée (Lepidoptera: Crambidae). *J. Asia-Pac. Entomol.* **2023**, *26*, 102031. [CrossRef]
170. Ashok, K.; Bhargava, C.N.; Venkatesh, R.; Balasubramani, V.; Murugan, M.; Geethalakshmi, V.; Manamohan, M.; Jha, G.K.; Asokan, R. Molecular characterization and CRISPR/Cas9 validation of the precursor of egg yolk protein gene, vitellogenin of *Leucinodes orbonalis* Guenée (Lepidoptera: Crambidae). *Gene* **2025**, *933*, 148925. [CrossRef] [PubMed]
171. Ashok, K.; Bhargava, C.N.; Asokan, R.; Pradeep, C.; Kennedy, J.S.; Rai, A.; Manamohan, M. First report on the utility of pupal case for early determination of CRISPR/Cas9 ribonucleoprotein mediated genomic edits in the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Tephritidae: Diptera). *Arch. Insect Biochem. Physiol.* **2023**, *113*, e22024. [CrossRef] [PubMed]

172. Bhargava, C.N.; Ashok, K.; Asokan, R.; Prasad Babu, K.; Parvathy, M.S.; Yogi, D.; Shashikala, T.; Chiranth, R.K.; Ashok, U.; Harsha, C.G. CRISPR/Cas9 Mediated Editing of the white (wh) locus Affects Body Size and Reproduction of the Oriental Fruit Fly, *Bactocera dorsalis* (Hendel). *Agric. Res.* **2024**, *13*, 733–740. [CrossRef]
173. Pradhan, S.K.; Karuppannasamy, A.; Sujatha, P.M.; Nagaraja, B.C.; Narayanappa, A.C.; Chalapathi, P.; Dhawane, Y.; Bynakal, S.; Riegler, M.; Maligeppagol, M.; et al. Embryonic microinjection of ribonucleoprotein complex (Cas9+ sgRNA) of white gene in melon fly, *Zeugodacus cucurbitae* (Coquillett) (Diptera: Tephritidae) produced white eye phenotype. *Arch. Insect Biochem. Physiol.* **2023**, *114*, e22059. [CrossRef] [PubMed]
174. Sujatha, P.M.; Karuppannasamy, A.; Chalapathi, P.; Dhawane, Y.; Narasimhappa, N.S.; Narayanappa, A.C.; Munikrishnappa, V.K.T.; Chikmagalur Nagaraja, B.; Thalooru, S.; Kesavan, S.; et al. CRISPR/Cas9 Ribo nucleoprotein complex-mediated editing of the OBP13 gene affected the response of male *Bactrocera dorsalis* (Diptera: Tephritidae) to methyl eugenol. *J. Appl. Entomol.* **2024**, *148*, 760–771. [CrossRef]
175. Yadav, A.K.; Butler, C.; Yamamoto, A.; Patil, A.A.; Lloyd, A.L.; Scott, M.J. CRISPR/Cas9-based split homing gene drive targeting doublesex for population suppression of the global fruit pest *Drosophila suzukii*. *Proc. Natl. Acad. Sci. USA* **2023**, *120*, e2301525120. [CrossRef] [PubMed]
176. Fang, C.; James, B.; Williams, M.; Bachler, A.; Tay, W.T.; Walsh, T.; Frese, M. Cry1 resistance in a CRISPR/Cas9-mediated HaCad1 gene knockout strain of the Australian cotton bollworm *Helicoverpa armigera conferta* (Lepidoptera: Noctuidae). *Pest Manag. Sci.* **2025**, *81*, 959–965. [CrossRef] [PubMed]
177. Matsuoka, Y.; Nakamura, T.; Watanabe, T.; Barnett, A.A.; Tomonari, S.; Ylla, G.; Whittle, C.A.; Noji, S.; Mito, T.; Extavour, C.G. Establishment of CRISPR/Cas9-based knock-in in a hemimetabolous insect: Targeted gene tagging in the cricket *Gryllus bimaculatus*. *Development* **2025**, *152*, dev199746. [CrossRef] [PubMed]
178. Tao, J.; Bauer, D.E.; Chiarle, R. Assessing and advancing the safety of CRISPR-Cas tools: From DNA to RNA editing. *Nat. Commun.* **2023**, *14*, 212. [CrossRef] [PubMed]
179. Richardson, C.; Kelsh, R.N.; J Richardson, R. New advances in CRISPR/Cas-mediated precise gene-editing techniques. *Dis. Model. Mech.* **2023**, *16*, dmm049874. [CrossRef] [PubMed]
180. Ahmad, A.; Munawar, N.; Khan, Z.; Qusmani, A.T.; Khan, S.H.; Jamil, A.; Ashraf, S.; Ghouri, M.Z.; Aslam, S.; Mubarik, M.S.; et al. An Outlook on Global Regulatory Landscape for Genome-Edited Crops. *Int. J. Mol. Sci.* **2021**, *22*, 11753. [CrossRef] [PubMed]
181. Asad, M.; Chang, Y.; Liao, J.; Yang, G. CRISPR/Cas9 Genome Editing in the Diamondback Moth: Current Progress, Challenges, and Prospects. *Int. J. Mol. Sci.* **2025**, *26*, 1515. [CrossRef] [PubMed]
182. Komal, J.; Desai, H.R.; Samal, I.; Mastinu, A.; Patel, R.D.; Kumar, P.V.D.; Majhi, P.K.; Mahanta, D.K.; Bhoi, T.K. Unveiling the Genetic Symphony: Harnessing CRISPR-Cas Genome Editing for Effective Insect Pest Management. *Plants* **2023**, *12*, 3961. [CrossRef] [PubMed]
183. Moon, T.T.; Maliha, I.J.; Khan, A.A.M.; Chakraborty, M.; Uddin, M.S.; Amin, M.R.; Islam, T. CRISPR-Cas Genome Editing for Insect Pest Stress Management in Crop Plants. *Stresses* **2022**, *2*, 493–514. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

MDPI AG
Grosspeteranlage 5
4052 Basel
Switzerland
Tel.: +41 61 683 77 34

Insects Editorial Office
E-mail: insects@mdpi.com
www.mdpi.com/journal/insects



Disclaimer/Publisher's Note: The title and front matter of this reprint are at the discretion of the Guest Editors. The publisher is not responsible for their content or any associated concerns. The statements, opinions and data contained in all individual articles are solely those of the individual Editors and contributors and not of MDPI. MDPI disclaims responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.





Academic Open
Access Publishing

mdpi.com

ISBN 978-3-7258-7465-1