

Article

Existence of Solutions for Nonlinear Fractional Differential Equations and Inclusions Depending on Lower-Order Fractional Derivatives

Subramanian Muthaiah ¹  and Dumitru Baleanu ^{2,3*} 

¹ Department of Mathematics, Sri Ramakrishna Mission Vidyalaya College of Arts and Science, Coimbatore 641020, India; subramanianmcbe@gmail.com

² Department of Mathematics, Cankaya University, Balgat 06530, Ankara, Turkey;

³ Institute of Space Science, 077125 Magurele-Bucharest, Romania

* Correspondence: dimitru@cankaya.edu.tr

Received: 23 March 2020; Accepted: 13 April 2020; Published: 25 April 2020



Abstract: This article deals with the solutions of the existence and uniqueness for a new class of boundary value problems (BVPs) involving nonlinear fractional differential equations (FDEs), inclusions, and boundary conditions involving the generalized fractional integral. The nonlinearity relies on the unknown function and its fractional derivatives in the lower order. We use fixed-point theorems with single-valued and multi-valued maps to obtain the desired results, through the support of illustrations, the main results are well explained. We also address some variants of the problem.

Keywords: single-valued map; multi-valued map; Caputo derivative; generalized Riemann–Liouville integral

1. Introduction

The subject of the fractional boundary value problem (BVP) has been intensively discussed in recent years by several researchers and in the literature, for example [1–19] and the references cited therein, where a variety of findings relevant to both the theoretical and implementation aspects of the topic can be found. It has improved the classic modeling of many significant materials and processes with the use of fractional calculus tools as a fractional-order operator can take the history of the phenomena involved into account. The extensive applications of fractional calculus can easily be seen in many engineering and technical sciences such as biology, environmental problems, aerodynamics, electron-analytic chemistry, etc. We direct the viewer to the article [20–28] and the references listed in it for examples and information. Recently, some authors analyzed the problems of fractional differential equations and inclusions. Ahmad et al. discussed in [29] the fractional differential equations (FDEs) and inclusions with nonlocal Erdelyi–Kober integral conditions:

$$\begin{aligned} {}^C\mathcal{D}^q x(t) &= f(t, x(t)), \quad \tau \in [0, T] := \mathcal{K}, \\ x(0) &= g(x), \quad x(T) = \alpha \mathcal{J}_\eta^{\gamma, \delta} x(\xi). \end{aligned}$$

and:

$$\begin{aligned} {}^C\mathcal{D}^q x(t) &\in f(t, x(t)), \quad \tau \in [0, T] := \mathcal{K}, \\ x(0) &= g(x), \quad x(T) = \alpha \mathcal{J}_\eta^{\gamma, \delta} x(\xi). \end{aligned}$$

Ntouyas et al. [30] investigated the existence of solutions for fractional differential inclusion. Salem et al. [31] studied the FDEs and inclusions, under integral-multipoint conditions.

In this article, we examine a new BVP of FDEs and inclusions:

$${}^C\mathcal{D}^\zeta y(\tau) = g(\tau, y(\tau), {}^C\mathcal{D}^\rho y(\tau), {}^C\mathcal{D}^{\rho+1} y(\tau)), \tau \in [0, T] := \mathcal{K}, \tag{1}$$

$${}^C\mathcal{D}^\zeta y(\tau) \in \mathcal{G}(\tau, y(\tau), {}^C\mathcal{D}^\rho y(\tau), {}^C\mathcal{D}^{\rho+1} y(\tau)), \tau \in [0, T] := \mathcal{K}, \tag{2}$$

augmented with the boundary conditions given by:

$$y(0) = 0, y'(0) = 0, \int_0^T y(\sigma) d\sigma = \zeta^\rho \mathcal{J}^\omega y(\zeta), y'(T) = 0, \tag{3}$$

where ${}^C\mathcal{D}^{(\cdot)}$ denotes the Caputo fractional derivatives (CFDs) of order (\cdot) , $3 < \zeta \leq 4$, $0 < \rho \leq 1$, ${}^\rho \mathcal{J}^\omega$ denote the generalized Riemann–Liouville fractional integral (GRLFI) of order $0 < \omega < 1$, $\rho > 0$, $g : \mathcal{K} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a continuous function, $\mathcal{G} : \mathcal{K} \times \mathbb{R}^3 \rightarrow \mathcal{T}(\mathbb{R})$ is a multivalued map, $\mathcal{T}(\mathbb{R})$ is all nonempty subsets of \mathbb{R} , and $0 < \zeta < T$, ζ is a real constant.

For $0 < \rho \leq 1$, let $\mathcal{Y} = \{y : y, {}^C\mathcal{D}^\rho y, {}^C\mathcal{D}^{\rho+1} y \in C(\mathcal{K}, \mathbb{R})\}$ denote the $[0, T] \rightarrow \mathbb{R}$ continuous function space of Banach endowed with the $\|y\|_* = \|y\| + \|{}^C\mathcal{D}^\rho y\| + \|{}^C\mathcal{D}^{\rho+1} y\| = \sup_{\tau \in \mathcal{K}} \{|y(\tau)| + |{}^C\mathcal{D}^\rho y(\tau)| + |{}^C\mathcal{D}^{\rho+1} y(\tau)|\}$ norm. For a normed space $(\mathcal{Y}, \|\cdot\|)$, let $\mathcal{T}_{cl}(\mathcal{Y}) = \{\mathcal{Z} \in \mathcal{T}(\mathcal{Y}) : \mathcal{Y} \text{ is closed}\}$, $\mathcal{T}_{bd}(\mathcal{Y}) = \{\mathcal{Z} \in \mathcal{T}(\mathcal{Y}) : \mathcal{Y} \text{ is bounded}\}$, $\mathcal{T}_{cpt}(\mathcal{Y}) = \{\mathcal{Z} \in \mathcal{T}(\mathcal{Y}) : \mathcal{Y} \text{ is compact}\}$, and $\mathcal{T}_{cpt, cx}(\mathcal{Y}) = \{\mathcal{Z} \in \mathcal{T}(\mathcal{Y}) : \mathcal{Y} \text{ is compact and convex}\}$. Define the set of choices \mathcal{G} by each $C(\mathcal{K}, \mathbb{R})$,

$$\mathcal{W}_{\mathcal{G}, y} = \{\phi : \mathcal{L}^1(\mathcal{K}, \mathbb{R}) : \phi(\tau) \in \mathcal{G}(\tau, y(\tau), {}^C\mathcal{D}^\rho y(\tau), {}^C\mathcal{D}^{\rho+1} y(\tau)) \text{ for a.e. } \tau \in \mathcal{K}\}.$$

The remaining part of the article is structured accordingly. We recall some definitions in Section 2 and establish a lemma regarding the linear problem variant (1)–(3). Sections 3 and 4 include the consequences of existence. We emphasize that the techniques used in these sections in fixed-point theory are the standard. Finally, we glance at a new problem similar to (1)–(3) and discuss the approach to solving them.

2. Preliminaries

Here, we are reminded of some basic concepts in the fractional calculus [23,32,33] and of the results that we need to accomplish during the upcoming analysis.

Definition 1. A continuous function $g : (0, \infty) \rightarrow \mathbb{R}$ is defined by the RLFI of order $\zeta > 0$:

$$\mathcal{I}^\zeta g(\tau) = \frac{1}{\Gamma(\zeta)} \int_0^\tau (\tau - \sigma)^{\zeta-1} g(\sigma) d\sigma,$$

provided the right-hand side (RHS) is point-wise defined on $(0, \infty)$.

Definition 2. The CFD of order ζ for a function $g : [0, \infty) \rightarrow \mathbb{R}$ can be written as:

$${}^C\mathcal{D}^\zeta g(\tau) = \frac{1}{\Gamma(n - \zeta)} (\tau - \sigma)^{n-\zeta-1} g^n(\sigma) d\sigma, \quad n - 1 < \zeta < n, \quad n = [\zeta] + 1,$$

where $[\zeta]$ denotes the integer part of the real number $[\zeta]$.

Definition 3. The GRLFI of order $\zeta > 0$ and $\rho > 0$, of a function $g(\tau)$, $\forall 0 < \tau < \infty$, is defined as:

$${}^\rho \mathcal{I}^\zeta g(\tau) = \frac{\rho^{1-\zeta}}{\Gamma(\zeta)} \int_0^\tau \frac{\sigma^{\rho-1}}{(\tau^\rho - \sigma^\rho)^{1-\zeta}} d\sigma,$$

provided the RHS is point-wise defined on $(0, \infty)$.

Definition 4. A multi-valued map $\mathcal{G} : \mathcal{K} \times \mathbb{R}^3 \rightarrow \mathcal{T}(\mathbb{R})$ is Caratheodory if:

- (i) $\tau \mapsto \mathcal{G}(\tau, y, z, w)$ is measurable for each $y, z, w \in \mathbb{R}$;
- (ii) $(y, z, w) \mapsto \mathcal{G}(\tau, y, z, w)$ is upper semicontinuous (USC) $\forall \tau \in \mathcal{K}$; In addition, a \mathcal{G} feature of Caratheodory is called \mathcal{L}^1 -Caratheodory, if:
- (iii) for each $\alpha > 0$, there exists $\Lambda_\alpha \in \mathcal{L}^1(\mathcal{K}, \mathbb{R}^+)$ such that $\|\mathcal{G}(\tau, y, z, w)\| = \sup\{|\phi| : \phi \in \mathcal{G}(\tau, y, z, w)\} \leq \Lambda_\alpha(\tau) \forall \|y\|, \|z\|, \|w\| \leq \alpha$ and for almost everywhere $\tau \in \mathcal{K}$.

Lemma 1. Let $\hat{g} \in C[0, T]$. Then, the unique solution of the linear FDE:

$${}^C\mathcal{D}^\varsigma y(\tau) = \hat{g}(\tau), \quad \tau \in \mathcal{K}, \tag{4}$$

subject to the boundary condition (3) is given by:

$$y(\tau) = \mathcal{I}^\varsigma \hat{g}(\tau) + \kappa_1(\tau) \left[\xi^\rho \mathcal{J}^\omega \mathcal{I}^\varsigma \hat{g}(\zeta) - \int_0^T \mathcal{I}^\varsigma \hat{g}(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\varsigma-1} \hat{g}(\tau), \tag{5}$$

where:

$$\kappa_1(\tau) = \frac{\tau^2 3T^2 - 2T\tau^3}{\vartheta}, \quad \kappa_2(\tau) = \frac{\tau^2 v_2 - v_1 \tau^3}{\vartheta}, \quad \vartheta = 3T^2 v_1 - 2T v_2, \tag{6}$$

$$v_1 = \frac{T^3}{3} - \frac{\xi \zeta^{\rho\omega+2}}{\rho^\omega} \frac{\Gamma(\frac{2}{\rho} + 1)}{\Gamma(\frac{2}{\rho} + \omega + 1)}, \quad v_2 = \frac{T^4}{4} - \frac{\xi \zeta^{\rho\omega+3}}{\rho^\omega} \frac{\Gamma(\frac{3}{\rho} + 1)}{\Gamma(\frac{3}{\rho} + \omega + 1)}. \tag{7}$$

Definition 5. A multi-valued operator $\mathcal{U} : \mathcal{Y} \rightarrow \mathcal{T}_{cl}(\mathcal{Y})$ is called:

- (a) ι -Lipschitz iff there exists $\iota > 0$ such that $\mathcal{A}_d(\mathcal{U}(y), \mathcal{U}(z)) \leq \iota d(x, y)$ for each $y, z \in \mathcal{Y}$ and
- (b) a contraction iff it is ι -Lipschitz with $\iota > 1$.

Definition 6. A function $y \in C(\mathcal{K}, \mathbb{R})$ is said to be a solution of the BVP $y(0) = 0, y'(0) = 0, \int_0^T y(\sigma) d\sigma = \xi^\rho \mathcal{J}^\omega y(\zeta), y'(T) = 0$, and there exists a function $\phi \in \mathcal{W}_{\mathcal{G}, y}$ such that $\phi(\tau) \in \mathcal{G}(\tau, y(\tau), {}^C\mathcal{D}^\rho y(\tau), {}^C\mathcal{D}^{\rho+1} y(\tau))$, and:

$$y(\tau) = \mathcal{I}^\varsigma \phi(\sigma)(\tau) + \kappa_1(\tau) \left[\xi^\rho \mathcal{J}^\omega \mathcal{I}^\varsigma \phi(\sigma)(\zeta) - \int_0^T \mathcal{I}^\varsigma \phi(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\varsigma-1} \phi(\sigma)(\tau).$$

3. Single-Valued Maps for the Problem (1) and (3)

With respect to Lemma 1, the problem (1) and (3) is turned into a fixed point problem equivalent to:

$$y = \Upsilon y, \tag{8}$$

where $\Upsilon : \mathcal{Y} \rightarrow \mathcal{Y}$ is defined by:

$$\begin{aligned} (\Upsilon y)(\tau) &= \mathcal{I}^\varsigma g(\sigma, y(\sigma), {}^C\mathcal{D}^\rho y(\sigma), {}^C\mathcal{D}^{\rho+1} y(\sigma))(\tau) \\ &\quad + \kappa_1(\tau) \left[\xi^\rho \mathcal{J}^\omega \mathcal{I}^\varsigma g(\sigma, y(\sigma), {}^C\mathcal{D}^\rho y(\sigma), {}^C\mathcal{D}^{\rho+1} y(\sigma))(\zeta) \right. \\ &\quad \left. - \int_0^T \mathcal{I}^\varsigma g(\sigma, y(\sigma), {}^C\mathcal{D}^\rho y(\sigma), {}^C\mathcal{D}^{\rho+1} y(\sigma)) d\sigma \right] \\ &\quad + \kappa_2(\tau) \mathcal{I}^{\varsigma-1} g(\sigma, y(\sigma), {}^C\mathcal{D}^\rho y(\sigma), {}^C\mathcal{D}^{\rho+1} y(\sigma))(\tau). \end{aligned} \tag{9}$$

We represent it as suitable for computing:

$$\psi_1 = \frac{T^\zeta}{\Gamma(\zeta + 1)} + \frac{\tilde{\kappa}_2 T^{\zeta-1}}{\Gamma(\zeta)} + \tilde{\kappa}_1 \left(\frac{\xi \zeta^{\zeta+\rho\omega}}{\rho^\omega \Gamma(\zeta + 1)} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\Gamma(\zeta + 2)} \right), \tag{10}$$

$$\psi_2 = \frac{T^{\zeta-\varrho}}{\Gamma(\zeta - \varrho + 1)} + \frac{\varphi_1}{\Gamma(\zeta + 1)} \left(\frac{\xi \zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) + \frac{\varphi_2 T^{\zeta-1}}{\Gamma(\zeta)}, \tag{11}$$

$$\psi_3 = \frac{T^{\zeta-\varrho-1}}{\Gamma(\zeta - \varrho)} + \frac{\delta_1}{\Gamma(\zeta + 1)} \left(\frac{\xi \zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) + \frac{\delta_2 T^{\zeta-1}}{\Gamma(\zeta)}, \tag{12}$$

$$\hat{\psi}_1 = \psi_1 - \frac{T^\zeta}{\Gamma(\zeta + 1)}, \hat{\psi}_2 = \psi_2 - \frac{T^{\zeta-\varrho}}{\Gamma(\zeta - \varrho + 1)}, \hat{\psi}_3 = \psi_3 - \frac{T^{\zeta-\varrho-1}}{\Gamma(\zeta - \varrho)}. \tag{13}$$

Theorem 1. Assume that there exists $\lambda \in C(\mathcal{K}, \mathbb{R}^+)$ such that $|g(\tau, y(\tau), {}^C\mathcal{D}^\varrho y(\tau), {}^C\mathcal{D}^{\varrho+1} y(\tau))| \leq \lambda(\tau)$ for $\tau \in \mathcal{K}$ with $\max_{\tau \in \mathcal{K}} |\lambda(\tau)| = \|\lambda\|$. The problem (1) and (3) has at least one solution on \mathcal{K} .

Proof. First, we demonstrate that operator Y is completely continuous. Let $\mathcal{H} \subset \mathcal{Y}$ be a bounded set. Then, use the premise $|g(\tau, y(\tau), {}^C\mathcal{D}^\varrho y(\tau), {}^C\mathcal{D}^{\varrho+1} y(\tau))| \leq \lambda(\tau), \forall y \in \mathcal{H}$; we get:

$$|(Yy)(\tau)| \leq \mathcal{I}^\zeta \lambda(\sigma)(\tau) + \kappa_1(\tau) \left[\xi^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \lambda(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \lambda(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \lambda(\sigma)(\tau),$$

which yields when taking the norm for $\tau \in \mathcal{K}$,

$$\|Yy\| \leq \frac{\|\lambda\|}{\Gamma(\zeta + 1)} \left[(\tilde{\kappa}_2 \zeta T^{\zeta-1} + T^\zeta) + \tilde{\kappa}_1 \left(\frac{\xi \zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) \right] = \mathcal{P}_1,$$

where $\max_{\tau \in \mathcal{K}} |\kappa_i(\tau)| = \tilde{\kappa}_i, i = 1, 2$ κ_i 's are given by (6). Similarly, we can obtain:

$$\|{}^C\mathcal{D}^\varrho Yy\| \leq \|\lambda\| \left[\frac{T^{\zeta-\varrho}}{\Gamma(\zeta - \varrho + 1)} + \frac{\varphi_1}{\Gamma(\zeta + 1)} \left(\frac{\xi \zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) + \frac{\varphi_2 T^{\zeta-1}}{\Gamma(\zeta)} \right] = \mathcal{P}_2,$$

where $\varphi_i = \max_{\tau \in \mathcal{K}} |\varphi_i(\tau), i = 1, 2$ and:

$$\varphi_1(\tau) = {}^C\mathcal{D}^\varrho \kappa_1(\tau) = \frac{6T^2\tau^{2-\varrho}}{\vartheta\Gamma(3-\varrho)} - \frac{12T\tau^{3-\varrho}}{\vartheta\Gamma(4-\varrho)}, \varphi_2(\tau) = {}^C\mathcal{D}^\varrho \kappa_2(\tau) = \frac{2\nu_2\tau^{2-\varrho}}{\vartheta\Gamma(3-\varrho)} - \frac{6\nu_1\tau^{3-\varrho}}{\vartheta\Gamma(4-\varrho)}.$$

Likewise, we can obtain:

$$\|{}^C\mathcal{D}^{\varrho+1} Yy\| \leq \|\lambda\| \left[\frac{T^{\zeta-\varrho-1}}{\Gamma(\zeta - \varrho)} + \frac{\delta_1}{\Gamma(\zeta + 1)} \left(\frac{\xi \zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) + \frac{\delta_2 T^{\zeta-1}}{\Gamma(\zeta)} \right] = \mathcal{P}_3,$$

where $\delta_i = \max_{\tau \in \mathcal{K}} |\delta_i(\tau), i = 1, 2$ and:

$$\delta_1(\tau) = {}^C\mathcal{D}^{\varrho+1} \kappa_1(\tau) = \frac{6T^2\tau^{1-\varrho}}{\vartheta\Gamma(2-\varrho)} - \frac{12T\tau^{2-\varrho}}{\vartheta\Gamma(3-\varrho)}, \delta_2(\tau) = {}^C\mathcal{D}^{\varrho+1} \kappa_2(\tau) = \frac{2\nu_2\tau^{1-\varrho}}{\vartheta\Gamma(2-\varrho)} - \frac{6\nu_1\tau^{2-\varrho}}{\vartheta\Gamma(3-\varrho)}.$$

For $0 < \tau_1 < \tau_2 < T$ and $\forall y \in \mathcal{H}$, we have:

$$\begin{aligned}
 |(Yy)(\tau_2) - (Yy)(\tau_1)| \leq & \|\lambda\| \left\{ \left| \frac{|\tau_2^\zeta - \tau_1^\zeta| + 2(\tau_2 - \tau_1)^\zeta}{\Gamma(\zeta + 1)} \right| + \left| \left(\frac{(\tau_2^2 - \tau_1^2)v_2 - v_1(\tau_2^3 - \tau_1^3)}{\vartheta} \right) \frac{T^{\zeta-1}}{\Gamma(\zeta)} \right| \right. \\
 & + \left| \frac{(\tau_2^2 - \tau_1^2)3T^2 - 2T(\tau_2^3 - \tau_1^3)}{\vartheta\Gamma(\zeta + 1)} \right. \\
 & \left. \times \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) \right\}. \tag{14}
 \end{aligned}$$

Similarly, we can accomplish:

$$\begin{aligned}
 |({}^C\mathcal{D}^\varrho Yy)(\tau_2) - ({}^C\mathcal{D}^\varrho Yy)(\tau_1)| \leq & \|\lambda\| \left\{ \left| \frac{|\tau_2^{\zeta-\varrho} - \tau_1^{\zeta-\varrho}| + 2(\tau_2 - \tau_1)^{\zeta-\varrho}}{\Gamma(\zeta - \varrho + 1)} \right| \right. \\
 & + \left| \left(\frac{2v_2(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3 - \varrho)} - \frac{6v_1(\tau_2^{3-\varrho} - \tau_1^{3-\varrho})}{\vartheta\Gamma(4 - \varrho)} \right) \frac{T^{\zeta-1}}{\Gamma(\zeta)} \right| \\
 & + \left| \frac{6T^2(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3 - \varrho)} - \frac{12T(\tau_2^{3-\varrho} - \tau_1^{3-\varrho})}{\vartheta\Gamma(4 - \varrho)} \right. \\
 & \left. \times \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega\Gamma(\zeta + 1)} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\Gamma(\zeta + 2)} \right) \right\}. \tag{15}
 \end{aligned}$$

Likewise, we obtain:

$$\begin{aligned}
 |({}^C\mathcal{D}^{\varrho+1} Yy)(\tau_2) - ({}^C\mathcal{D}^{\varrho+1} Yy)(\tau_1)| \leq & \|\lambda\| \left\{ \left| \frac{|\tau_2^{\zeta-\varrho-1} - \tau_1^{\zeta-\varrho-1}| + 2(\tau_2 - \tau_1)^{\zeta-\varrho-1}}{\Gamma(\zeta - \varrho)} \right| \right. \\
 & + \left| \left(\frac{2v_2(\tau_2^{1-\varrho} - \tau_1^{1-\varrho})}{\vartheta\Gamma(2 - \varrho)} - \frac{6v_1(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3 - \varrho)} \right) \frac{T^{\zeta-1}}{\Gamma(\zeta)} \right| \\
 & + \left| \frac{6T^2(\tau_2^{1-\varrho} - \tau_1^{1-\varrho})}{\vartheta\Gamma(2 - \varrho)} - \frac{12T(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3 - \varrho)} \right. \\
 & \left. \times \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega\Gamma(\zeta + 1)} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\Gamma(\zeta + 2)} \right) \right\}. \tag{16}
 \end{aligned}$$

The RHS of the inequalities (14)–(16) tends to zero as $\tau_2 - \tau_1 \rightarrow 0$ independently of y . Thus, Y is equicontinuous. Therefore, by the lemma (see Lemma 1.2 [21]), $\{Yy : y \in \mathcal{H}\}$, $\{{}^C\mathcal{D}^\varrho Yy : y \in \mathcal{H}\}$, and $\{{}^C\mathcal{D}^{\varrho+1} Yy : y \in \mathcal{H}\}$ are relatively compact in $C(\mathcal{K})$. Hence, $Y(\mathcal{H})$ is a relatively compact subset of \mathcal{Y} . Next, we take the set $\mathcal{V} = \{y \in \mathcal{Y} | y = \mu Yy, 0 < \mu < 1\}$, into consideration and prove it is bounded. Let $y \in \mathcal{V}$. Then, $y = \mu Yy, 0 < \mu < 1$. For any $\tau \in \mathcal{K}$, it follows from $y(\tau) = \mu |Yy(\tau)|$ that:

$$\|Yy\| \leq \frac{\|\lambda\|}{\Gamma(\zeta + 1)} \left[(\tilde{\kappa}_2\zeta T^{\zeta-1} + T^\zeta) + \tilde{\kappa}_1 \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) \right]$$

This indicates that the set \mathcal{V} is bounded. Thus, operator Y has at least one fixed point by Theorem (see Theorem [20]) The problem (1) and (3) has at least one solution on \mathcal{K} . □

Theorem 2. Let $g : \mathcal{K} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ be a continuous function that holds the following conditions:

$$(\mathcal{G}_1) |g(\tau, y_1, y_2, y_3) - g(\tau, z_1, z_2, z_3)| \leq \mathcal{P}(|y_1 - z_1| + |y_2 - z_2| + |y_3 - z_3|), \quad \forall \tau \in \mathcal{K}, y_1, z_1, y_2, z_2, y_3, z_3 \in \mathbb{R}, \mathcal{P} > 0.$$

- (G₂) $|g(\tau, y(\tau), {}^C\mathcal{D}^\varrho y(\tau), {}^C\mathcal{D}^{\varrho+1} y(\tau))| \leq \Lambda(\tau)$ for $\tau \in \mathcal{K}$ and $\Lambda \in C(\mathcal{K}, \mathbb{R}^+)$ with $\max_{\tau \in \mathcal{K}} |\Lambda(\tau)| = \|\Lambda\|$.
 (G₃) $\mathcal{P}\hat{\psi} < 1$, where $\hat{\psi} = \max\{\hat{\psi}_1, \hat{\psi}_2, \hat{\psi}_3\}$ and $\hat{\psi}_1, \hat{\psi}_2$ and $\hat{\psi}_3$ are given by (13). The problem (1) and (3) has at least one solution on \mathcal{K} .

Proof. Define $\mathcal{B}_\epsilon = \{y \in \mathcal{Y} : \|y\| \leq \epsilon\}$, where $\epsilon \geq \|\Lambda\|\psi$ with:

$$\psi = \max\{\psi_1, \psi_2, \psi_3\}, \tag{17}$$

where ψ_1, ψ_2 , and ψ_3 are defined by (10)–(12), respectively. In order to demonstrate the premise of Theorem (see Theorem 4.4.1 [20]), the operator Y provided by (9) is divided into $Y = Y_1 + Y_2$ by \mathcal{B}_ϵ .

$$\begin{aligned} (Y_1y)(\tau) &= \mathcal{I}^\zeta g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau) \\ (Y_2y)(\tau) &= |\kappa_1(\tau)| \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\zeta) \right. \\ &\quad \left. - \int_0^T \mathcal{I}^\zeta g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma)) d\sigma \right] \\ &\quad + |\kappa_2(\tau)| \mathcal{I}^{\zeta-1} g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau). \end{aligned}$$

It can be easily shown that for $y, z \in \mathcal{B}_\epsilon$, and using (17), $\|(Y_1y) + (Y_2z)\| \leq \|\Lambda\|\psi \leq \epsilon$, $\|({}^C\mathcal{D}^\varrho Y_1y) + ({}^C\mathcal{D}^\varrho Y_2z)\| \leq \|\Lambda\|\psi \leq \epsilon$ and $\|({}^C\mathcal{D}^{\varrho+1} Y_1y) + ({}^C\mathcal{D}^{\varrho+1} Y_2z)\| \leq \|\Lambda\|\psi \leq \epsilon$, that means $Y_1y + Y_2z \in \mathcal{B}_\epsilon$. Next, we are going to show that Y_2 is a contraction. Let $y, z \in \mathbb{R}$, $\tau \in \mathcal{K}$. Then, we use the statement (G₁),

$$\begin{aligned} \|Y_2y - Y_2z\| &\leq \mathcal{P}\|y - z\| \sup_{\tau \in \mathcal{K}} \left[|\kappa_1(\tau)| \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta(1)(\zeta) + \int_0^T \mathcal{I}^\zeta(1) d\sigma \right] + |\kappa_2(\tau)| \mathcal{I}^{\zeta-1}(1)(\tau) \right] \\ &\leq \mathcal{P}\hat{\psi}_1 \|y - z\| \leq \mathcal{P}\hat{\psi} \|y - z\|. \end{aligned}$$

Similarly,

$$\|{}^C\mathcal{D}^\varrho Y_2y - {}^C\mathcal{D}^\varrho Y_2z\| \leq \mathcal{P}\hat{\psi}_2 \|y - z\| \leq \mathcal{P}\hat{\psi} \|y - z\|.$$

Likewise,

$$\|{}^C\mathcal{D}^{\varrho+1} Y_2y - {}^C\mathcal{D}^{\varrho+1} Y_2z\| \leq \mathcal{P}\hat{\psi}_3 \|y - z\| \leq \mathcal{P}\hat{\psi} \|y - z\|.$$

This follows from the observation (G₃) that the Y_2 operator is a contraction. Next we are going to show the Y_1 is compact and continuous. The continuity of g means operator Y_1 is continuous. Y_1 is uniformly bounded on \mathcal{B}_ϵ as $\|Y_1y\| \leq \frac{\|\Lambda\|T^\zeta}{\Gamma(\zeta+1)}$, $\|{}^C\mathcal{D}^\varrho Y_1y\| \leq \frac{\|\Lambda\|T^{\zeta-\varrho}}{\Gamma(\zeta-\varrho+1)}$ and $\|{}^C\mathcal{D}^{\varrho+1} Y_1y\| \leq \frac{\|\Lambda\|T^{\zeta-\varrho-1}}{\Gamma(\zeta-\varrho)}$. Furthermore, with $\sup_{(\tau, y, z, w) \in \mathcal{K} \times \mathcal{B}_\epsilon} |g(\tau, y, z, w)| = \hat{g} < \infty$ and $\tau_2 > \tau_1$, we have:

$$|(Y_1y)(\tau_2) - (Y_1y)(\tau_1)| \leq \left[\frac{\hat{g}|\tau_2^\zeta - \tau_1^\zeta| + 2(\tau_2 - \tau_1)^\zeta}{\Gamma(\zeta + 1)} \right]. \tag{18}$$

Similarly, we can obtain:

$$|({}^C\mathcal{D}^\varrho Y_1y)(\tau_2) - ({}^C\mathcal{D}^\varrho Y_1y)(\tau_1)| \leq \left[\frac{\hat{g}|\tau_2^{\zeta-\varrho} - \tau_1^{\zeta-\varrho}| + 2(\tau_2 - \tau_1)^{\zeta-\varrho}}{\Gamma(\zeta - \varrho + 1)} \right]. \tag{19}$$

Likewise, we obtain:

$$|({}^C\mathcal{D}^{\varrho+1}Y_1y)(\tau_2) - ({}^C\mathcal{D}^{\varrho+1}Y_1y)(\tau_1)| \leq \left[\frac{\widehat{g}|\tau_2^{\zeta-\varrho-1} - \tau_1^{\zeta-\varrho-1}| + 2(\tau_2 - \tau_1)^{\zeta-\varrho-1}}{\Gamma(\zeta - \varrho)} \right]. \tag{20}$$

The RHS of the inequalities (18)–(20) tend to zero as $\tau_2 - \tau_1 \rightarrow 0$ independently of y . Thus, Y_1 is relatively compact on \mathcal{B}_ϵ . Hence, by the lemma (see Lemma 1.2 [21]), Y_1 is compact on \mathcal{B}_ϵ . Therefore, all the claims of the theorem (see Theorem 4.4.1 [20]) are fulfilled. Therefore, at least one solution exists (1) and (3) for the problem on \mathcal{K} . \square

Theorem 3. Assume that $g : \mathcal{K} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a continuous function satisfying condition $|g(\tau, y_1, y_2, y_3) - g(\tau, z_1, z_2, z_3)| \leq \mathcal{Q}(|y_1 - z_1| + |y_2 - z_2| + |y_3 - z_3|), \forall \tau \in \mathcal{K}, y_1, z_1, y_2, z_2, y_3, z_3 \in \mathbb{R}$, with $\mathcal{Q} < \frac{1}{\psi}$, where $\psi = \max\{\psi_1, \psi_2, \psi_3\}$, and ψ_1, ψ_2 and ψ_3 are respectively given by (10)–(12). Therefore, a unique solution exists (1) and (3) for the problem on \mathcal{K} .

Proof. We demonstrate that $Y\mathcal{B}_\epsilon \subset \mathcal{B}_\epsilon$, where Y is described by (9), $\mathcal{B}_\epsilon = \{y \in \mathcal{Y} : \|y\| \leq \epsilon\}$ with $\epsilon \geq \frac{\mathcal{M}\psi}{1-\mathcal{Q}\psi}, \mathcal{M} = \sup_{\tau \in \mathcal{K}} |g(\tau, 0, 0, 0)|$. For $y \in \mathcal{B}_\epsilon, \tau \in \mathcal{K}$, we have that:

$$\begin{aligned} |g(\tau, y, z, w)| &= |g(\tau, y, z, w) - g(\tau, 0, 0, 0) + g(\tau, 0, 0, 0)| \\ &\leq |g(\tau, y, z, w) - g(\tau, 0, 0, 0)| + |g(\tau, 0, 0, 0)| \leq \mathcal{Q}(|y| + |z| + |w|) + \mathcal{M} \\ &\leq \mathcal{Q}\|y\| + \mathcal{M} \leq \mathcal{Q}\epsilon + \mathcal{M}, \end{aligned}$$

which yields along with the given conditions:

$$\begin{aligned} \|Yy\| &\leq \left[\frac{T^\zeta}{\Gamma(\zeta + 1)} + \frac{\tilde{\kappa}_2 T^{\zeta-1}}{\Gamma(\zeta)} + \frac{\tilde{\kappa}_1}{\Gamma(\zeta + 1)} \left(\frac{\tilde{\zeta} \zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) \right] \\ &\leq (\mathcal{Q}\epsilon + \mathcal{M})\psi_1 \leq (\mathcal{Q}\epsilon + \mathcal{M})\psi \leq \epsilon. \end{aligned}$$

Similarly, we can obtain:

$$\|{}^C\mathcal{D}^\varrho Yy\| \leq (\mathcal{Q}\epsilon + \mathcal{M})\psi_2 \leq (\mathcal{Q}\epsilon + \mathcal{M})\psi \leq \epsilon.$$

Likewise, we obtain:

$$\|{}^C\mathcal{D}^{\varrho+1}Yy\| \leq (\mathcal{Q}\epsilon + \mathcal{M})\psi_3 \leq (\mathcal{Q}\epsilon + \mathcal{M})\psi \leq \epsilon.$$

Therefore, we get $Yy \in \mathcal{B}_\epsilon$, which means $Y\mathcal{B}_\epsilon \subset \mathcal{B}_\epsilon$. Then, for $y, z \in \mathcal{Y}$, for each $\tau \in \mathcal{K}$, we have:

$$\begin{aligned} \|Yy - Yz\| &\leq \mathcal{Q}\|y - z\| \sup_{\tau \in \mathcal{K}} \left\{ \mathcal{I}^\zeta(1)(\tau) + \kappa_1(\tau) \left[\tilde{\zeta}^\rho \mathcal{J}^\omega \mathcal{I}^\zeta(1)(\zeta) - \int_0^T \mathcal{I}^\zeta(1)d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1}(1)(\tau) \right\} \\ &\leq \mathcal{Q}\psi_1\|y - z\| \leq \mathcal{Q}\psi\|y - z\|. \end{aligned}$$

Similarly,

$$\|{}^C\mathcal{D}^\varrho Y_2y - {}^C\mathcal{D}^\varrho Y_2z\| \leq \mathcal{Q}\psi_2\|y - z\| \leq \mathcal{Q}\psi\|y - z\|.$$

Likewise,

$$\|{}^C\mathcal{D}^{\varrho+1}Y_2y - {}^C\mathcal{D}^{\varrho+1}Y_2z\| \leq \mathcal{Q}\psi_3\|y - z\| \leq \mathcal{Q}\psi\|y - z\|.$$

Therefore, the operator Y is a contraction in the light of condition $Q < \frac{1}{\psi}$. Thus, Theorem (see Theorem 1.2.2 [20]) follows that the problem (1) and (3) has a unique solution on \mathcal{K} . \square

4. Multi-Valued Maps for the Problem (2) and (3)

Theorem 4. Assume that:

- (\mathcal{F}_1) $\mathcal{G} : \mathcal{K} \times \mathbb{R}^3 \rightarrow \mathcal{T}(\mathbb{R})$ is \mathcal{L}^1 -Caratheodory and has nonempty compact and convex values;
- (\mathcal{F}_2) there exists a function $\gamma \in C(\mathcal{K}, \mathbb{R})$, and a nondecreasing, sub-homogeneous function $\Delta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that (i.e., $\Delta(\beta y) \leq \beta \Delta(y) \forall \beta \geq 1$ and $y \in \mathbb{R}^+$):

$$\|\mathcal{G}(\tau, y)\|_{\mathcal{T}} := \sup\{|q| : q \in \mathcal{G}(\tau, y, z, w)\} \leq \gamma(\tau)\Delta(\|y\| + \|z\| + \|w\|)$$

for each $(\tau, y, z, w) \in \mathcal{K} \times \mathbb{R}^3$;

- (\mathcal{F}_3) there exists a constant $\mathcal{E} > 0$ such that:

$$\frac{\mathcal{E}}{\|\gamma\|\psi\Delta(\mathcal{E})} > 1,$$

Then, there is at least one solution for the BVP (2) and (3) on \mathcal{K} .

Proof. Define an operator $\Delta_{\mathcal{G}} : C(\mathcal{K}, \mathbb{R}) \rightarrow \mathcal{T}(C(\mathcal{K}, \mathbb{R}))$ by $\Delta_{\mathcal{G}}(y) = \{u \in C(\mathcal{K}, \mathbb{R}) : u(\tau) = \mathcal{S}(y)(\tau)\}$ where:

$$\mathcal{S}(y)(\tau) = \left\{ \mathcal{I}^{\varsigma} \phi(\sigma)(\tau) + \kappa_1(\tau) \left[\xi^{\rho} \mathcal{J}^{\omega} \mathcal{I}^{\varsigma} \phi(\sigma)(\zeta) - \int_0^T \mathcal{I}^{\varsigma} \phi(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\varsigma-1} \phi(\sigma)(\tau), \phi \in \mathcal{W}_{\mathcal{G}, y} \right\}$$

We will prove that $\Delta_{\mathcal{G}}$ follows the theorem’s assumptions (see 8. Theorem 8.5 [22]). The proof requires multiple measures. First, we demonstrate that for every $C(\mathcal{K}, \mathbb{R})$, $\Delta_{\mathcal{G}}$ is convex. This phase is evident as $\mathcal{W}_{\mathcal{G}, y}$ is convex, so we skip the proof. Next, we show that $\Delta_{\mathcal{G}}$ maps in $C(\mathcal{K}, \mathbb{R})$ bounded sets to bound sets. Let $\mathcal{B}_{\epsilon} = \{y \in C(\mathcal{K}, \mathbb{R}) : \|y\| \leq \epsilon\}$ be a bounded ball in $C(\mathcal{K}, \mathbb{R})$ for a positive number ϵ . Then, for each $u \in \Delta_{\mathcal{G}}(y)$, $y \in \mathcal{B}_{\epsilon}$, there exists $\phi \in \mathcal{W}_{\mathcal{G}, y}$ such that:

$$\mathcal{S}(y)(\tau) = \mathcal{I}^{\varsigma} \phi(\sigma)(\tau) + \kappa_1(\tau) \left[\xi^{\rho} \mathcal{J}^{\omega} \mathcal{I}^{\varsigma} \phi(\sigma)(\zeta) - \int_0^T \mathcal{I}^{\varsigma} \phi(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\varsigma-1} \phi(\sigma)(\tau), \phi \in \mathcal{W}_{\mathcal{G}, y}.$$

Then, we have for $\tau \in \mathcal{K}$:

$$\begin{aligned} |u(\tau)| &\leq \frac{1}{\Gamma(\varsigma+1)} \left[(\tilde{\kappa}_2 \xi T^{\varsigma-1} + T^{\varsigma}) + \tilde{\kappa}_1 \left(\frac{\xi \zeta^{\varsigma+\rho\omega}}{\rho^{\omega}} \frac{\Gamma(\frac{\varsigma}{\rho} + 1)}{\Gamma(\frac{\varsigma}{\rho} + \omega + 1)} + \frac{T^{\varsigma+1}}{\varsigma + 1} \right) \right] \|\gamma\| \Delta(\|y\|_{\mathcal{Y}}) \\ &\leq \psi_1 \|\gamma\| \Delta(\|y\|_{\mathcal{Y}}), \end{aligned}$$

which yields on the norm for $\tau \in \mathcal{K}$,

$$\|u\| \leq \psi_1 \|\gamma\| \Delta(\|y\|_{\mathcal{Y}}) \leq \psi_1 \|\gamma\| \Delta(\epsilon).$$

Similarly, we have:

$$\begin{aligned} \|{}^C \mathcal{D}^{\varrho} u(\tau)\| &\leq \left[\frac{T^{\varsigma-\varrho}}{\Gamma(\varsigma-\varrho+1)} + \frac{\varphi_1}{\Gamma(\varsigma+1)} \left(\frac{\xi \zeta^{\varsigma+\rho\omega}}{\rho^{\omega}} \frac{\Gamma(\frac{\varsigma}{\rho} + 1)}{\Gamma(\frac{\varsigma}{\rho} + \omega + 1)} + \frac{T^{\varsigma+1}}{\varsigma + 1} \right) + \frac{\varphi_2 T^{\varsigma-1}}{\Gamma(\varsigma)} \right] \|\gamma\| \Delta(\|y\|_{\mathcal{Y}}) \\ &\leq \psi_2 \|\gamma\| \Delta(\|y\|_{\mathcal{Y}}), \end{aligned}$$

Likewise, we have:

$$\begin{aligned} \|{}^C\mathcal{D}^{\varrho+1}u(\tau)\| &\leq \|\lambda\| \left[\frac{T^{\zeta-\varrho-1}}{\Gamma(\zeta-\varrho)} + \frac{\delta_1}{\Gamma(\zeta+1)} \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho}+1)}{\Gamma(\frac{\zeta}{\rho}+\omega+1)} + \frac{T^{\zeta+1}}{\zeta+1} \right) + \frac{\delta_2 T^{\zeta-1}}{\Gamma(\zeta)} \right] \|\gamma\|\Delta(\|y\|_Y) \\ &\leq \psi_3 \|\gamma\|\Delta(\|y\|_Y). \end{aligned}$$

As $u \in \Delta_{\mathcal{G}}(y)$, $y \in \mathcal{B}_\epsilon$ is arbitrary, so we have:

$$\begin{aligned} \|\Delta_{\mathcal{G}}(y)\|_Y &= \|\Delta_{\mathcal{G}}(y)\| + \|{}^C\mathcal{D}^{\varrho}\Delta_{\mathcal{G}}(y)\| + \|{}^C\mathcal{D}^{\varrho+1}\Delta_{\mathcal{G}}(y)\| \\ &\leq \|\gamma\|\Delta(\epsilon)(\psi). \end{aligned}$$

Next, we demonstrate that $\Delta_{\mathcal{G}}$ maps bounded into equicontinuous sets of $C(\mathcal{K}, \mathbb{R})$. Let $\tau_1, \tau_2 \in \mathcal{K}$ with $\tau_2 > \tau_1$ and $y \in \mathcal{B}_\epsilon$. For each $u \in \Delta_{\mathcal{G}}(y)$, we obtain:

$$\begin{aligned} |u(\tau_2) - u(\tau_1)| &\leq \left\{ \left[\frac{|\tau_2^\zeta - \tau_1^\zeta| + 2(\tau_2 - \tau_1)^\zeta}{\Gamma(\zeta+1)} \right] + \left| \left(\frac{(\tau_2^2 - \tau_1^2)v_2 - v_1(\tau_2^3 - \tau_1^3)}{\vartheta} \right) \frac{T^{\zeta-1}}{\Gamma(\zeta)} \right| \right. \\ &\quad \left. + \left| \frac{(\tau_2^2 - \tau_1^2)3T^2 - 2T(\tau_2^3 - \tau_1^3)}{\vartheta\Gamma(\zeta+1)} \right| \right. \\ &\quad \left. \times \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho}+1)}{\Gamma(\frac{\zeta}{\rho}+\omega+1)} + \frac{T^{\zeta+1}}{\zeta+1} \right) \right\} \|\gamma\|\Delta(\epsilon). \end{aligned}$$

Similarly, we can obtain:

$$\begin{aligned} |{}^C\mathcal{D}^{\varrho}u(\tau_2) - {}^C\mathcal{D}^{\varrho}u(\tau_1)| &\leq \left\{ \left[\frac{|\tau_2^{\zeta-\varrho} - \tau_1^{\zeta-\varrho}| + 2(\tau_2 - \tau_1)^{\zeta-\varrho}}{\Gamma(\zeta-\varrho+1)} \right] \right. \\ &\quad \left. + \left| \left(\frac{2v_2(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3-\varrho)} - \frac{6v_1(\tau_2^{3-\varrho} - \tau_1^{3-\varrho})}{\vartheta\Gamma(4-\varrho)} \right) \frac{T^{\zeta-1}}{\Gamma(\zeta)} \right| \right. \\ &\quad \left. + \left| \frac{6T^2(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3-\varrho)} - \frac{12T(\tau_2^{3-\varrho} - \tau_1^{3-\varrho})}{\vartheta\Gamma(4-\varrho)} \right| \right. \\ &\quad \left. \times \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega\Gamma(\zeta+1)} \frac{\Gamma(\frac{\zeta}{\rho}+1)}{\Gamma(\frac{\zeta}{\rho}+\omega+1)} + \frac{T^{\zeta+1}}{\Gamma(\zeta+2)} \right) \right\} \|\gamma\|\Delta(\epsilon). \end{aligned}$$

Likewise, we obtain:

$$\begin{aligned} |{}^C\mathcal{D}^{\varrho+1}u(\tau_2) - {}^C\mathcal{D}^{\varrho+1}u(\tau_1)| &\leq \left\{ \left[\frac{|\tau_2^{\zeta-\varrho-1} - \tau_1^{\zeta-\varrho-1}| + 2(\tau_2 - \tau_1)^{\zeta-\varrho-1}}{\Gamma(\zeta-\varrho)} \right] \right. \\ &\quad \left. + \left| \left(\frac{2v_2(\tau_2^{1-\varrho} - \tau_1^{1-\varrho})}{\vartheta\Gamma(2-\varrho)} - \frac{6v_1(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3-\varrho)} \right) \frac{T^{\zeta-1}}{\Gamma(\zeta)} \right| \right. \\ &\quad \left. + \left| \frac{6T^2(\tau_2^{1-\varrho} - \tau_1^{1-\varrho})}{\vartheta\Gamma(2-\varrho)} - \frac{12T(\tau_2^{2-\varrho} - \tau_1^{2-\varrho})}{\vartheta\Gamma(3-\varrho)} \right| \right. \\ &\quad \left. \times \left(\frac{\xi\zeta^{\zeta+\rho\omega}}{\rho^\omega\Gamma(\zeta+1)} \frac{\Gamma(\frac{\zeta}{\rho}+1)}{\Gamma(\frac{\zeta}{\rho}+\omega+1)} + \frac{T^{\zeta+1}}{\Gamma(\zeta+2)} \right) \right\} \|\gamma\|\Delta(\epsilon). \end{aligned}$$

Clearly, the RHS of the above-mentioned inequalities tends to be zero as $\tau_2 - \tau_1 \rightarrow 0$. As $\Delta_{\mathcal{G}}$ fulfills the premises, the lemma (see Lemma 1.2 [21]) follows that $\Delta_{\mathcal{G}} : C(\mathcal{K}, \mathbb{R}) \rightarrow \mathcal{T}(C(\mathcal{K}, \mathbb{R}))$ is completely continuous. We demonstrate that the $\Delta_{\mathcal{G}}$ is USC at the end. It is enough to prove that $\Delta_{\mathcal{G}}$ has a closed

graph in the lemma (see Proposition 1.2 [32]). Let $y_n \rightarrow y_*$, $u_n \in \Delta_G(y_n)$, and $u_n \rightarrow u_*$. Then, we have to prove that $u_* \in \Delta_G(y_*)$; there exists $\phi_n \in \mathcal{W}_{G,y_n}$ such that for each $\tau \in \mathcal{K}$,

$$u_n(\tau) = \mathcal{I}^\zeta \phi_n(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi_n(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi_n(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi_n(\sigma)(\tau).$$

Therefore, it is enough to prove that $\phi_* \in \mathcal{W}_{G,y_*}$ exists so that for each $\tau \in \mathcal{K}$,

$$u_*(\tau) = \mathcal{I}^\zeta \phi_*(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi_*(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi_*(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi_*(\sigma)(\tau).$$

Consider the linear operator $\Psi : \mathcal{L}^1(\mathcal{K}, \mathbb{R}) \rightarrow C(\mathcal{K}, \mathbb{R})$ provided by:

$$\phi \mapsto \Psi(\phi)(\tau) = \mathcal{I}^\zeta \phi(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi(\sigma)(\tau).$$

Observe that:

$$\begin{aligned} \|u_n(\tau) - u_n(\tau)\| &= \|\mathcal{I}^\zeta(\phi_n - \phi_*)(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta(\phi_n - \phi_*)(\sigma)(\zeta) \right. \\ &\quad \left. - \int_0^T \mathcal{I}^\zeta(\phi_n - \phi_*)(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1}(\phi_n - \phi_*)(\sigma)(\tau)\| \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

The lemma (see Lemma [34]) follows that $\Psi \circ \mathcal{W}_G$ is a closed graph operator. We also have $u_n(\tau) \in \Psi(\mathcal{W}_{G,y_n})$. Therefore, since $y_n \rightarrow y_*$, we have:

$$\begin{aligned} u_*(\tau) &= \mathcal{I}^\zeta \phi_*(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi_*(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi_*(\sigma) d\sigma \right] \\ &\quad + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi_*(\sigma)(\tau), \text{ for some } \phi_* \in \mathcal{W}_{G,y_*}. \end{aligned}$$

Next, we demonstrate that an open set exists $\mathcal{V} \subseteq C(\mathcal{K}, \mathbb{R})$ with $y \notin \Delta_G(y)$ for any $\iota \in (0, T)$ and all $y \in \partial \mathcal{V}$. Let $\iota \in (0, T)$ and $y \in \iota \Delta_G(y)$. Then, there exists $\phi \in \mathcal{L}^1(\mathcal{K}, \mathbb{R})$ with $\phi \in \mathcal{W}_{G,y}$ such that for $\tau \in \mathcal{K}$, we can obtain:

$$\|y\|_Y = \|y\| + \|\mathcal{C}\mathcal{D}^\rho y\| + \|\mathcal{C}\mathcal{D}^{\rho+1} y\| \leq \|\gamma\| \Delta(\|y\|_Y)(\psi).$$

This ensures that $\frac{\|y\|_Y}{\|\gamma\| \Delta(\|y\|_Y)(\psi)} \leq 1$. With regard to (C3), there exists \mathcal{E} such that $\|y\| \neq \mathcal{E}$. Let us set $\mathcal{V} = \{y \in C(\mathcal{K}, \mathbb{R}) : \|y\| < \mathcal{E}\}$. Remember that operator $\Delta_G : \bar{\mathcal{V}} \rightarrow \mathcal{T}(C(\mathcal{K}, \mathbb{R}))$ is USC and completely continuous. There is no $y \in \partial \mathcal{V}$ of the choice of \mathcal{V} such that $y \in \iota \Delta_G(y)$ for some $\tau \in (0, T)$. Therefore, we deduce from Theorem (see 8.Theorem 8.5 [22]) that Δ_G has a fixed point \mathcal{V} to the problem (2) and (3). \square

Theorem 5. Assume that:

(F₄) $\mathcal{G} : \mathcal{K} \times \mathbb{R}^3 \rightarrow \mathcal{T}_{cp}(\mathbb{R})$ is such that $\mathcal{G}(\cdot, y(\tau), \mathcal{C}\mathcal{D}^\rho y(\tau), \mathcal{C}\mathcal{D}^{\rho+1} y(\tau)) : \mathcal{K} \rightarrow \mathcal{T}_{cp}(\mathbb{R})$ is measurable for each $y \in \mathcal{R}$;

(F₅) $\mathcal{A}_d(\mathcal{G}(\tau, y, z, w), \mathcal{G}(\tau, \hat{y}, \hat{z}, \hat{w})) \leq s(\tau)[|y - \hat{y}| + |z - \hat{z}| + |w - \hat{w}|] \forall \tau \in \mathcal{K}$ and $y, z, w, \hat{y}, \hat{z}, \hat{w} \in \mathbb{R}$ with $s \in C(\mathcal{K}, \mathbb{R}^+)$ and $d(0, \mathcal{G}(\tau, 0, 0, 0)) \leq s(\tau) \forall \tau \in \mathcal{K}$. Then, the problem (2) and (3) has at least one solution for \mathcal{K} if:

$$\|s\| \psi < 1. \tag{21}$$

Proof. Consider operator $\Delta_G : C(\mathcal{K}, \mathbb{R}) \rightarrow \mathcal{T}(C(\mathcal{K}, \mathbb{R}))$ defined in Theorem 4 at the beginning of the proof. Remember that for each $C(\mathcal{K}, \mathbb{R})$, set $\mathcal{W}_{G,y}$ is not empty by Hypothesis (F₄), so \mathcal{G} has a measurable range (see Theorem III.6 [35]). Now, we prove that the operator Δ_G fulfills the lemma's

assumptions (see Lemma [36]). To show that $\Delta_{\mathcal{G}}(y) \in \mathcal{T}_{cl}(C(\mathcal{K}, \mathbb{R}))$ for each $y \in C(\mathcal{K}, \mathbb{R})$, let $p_{n_n \geq 0} \in \Delta_{\mathcal{G}}(y)$ be such that $p_n \rightarrow p$ ($n \rightarrow \infty$) in $C(\mathcal{K}, \mathbb{R})$. Then, $p \in C(\mathcal{K}, \mathbb{R})$, and there exists $\phi_n \in \mathcal{W}_{\mathcal{G}, y}$ such that, for each $\tau \in \mathcal{K}$,

$$p_n(\tau) = \mathcal{I}^\zeta \phi_n(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi_n(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi_n(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi_n(\sigma)(\tau).$$

Since \mathcal{G} has compact values, we move a subsequence (if required) to get p_n converging to p in $\mathcal{L}^1(\mathcal{K}, \mathbb{R})$. Therefore, $\phi \in \mathcal{W}_{\mathcal{G}, y}$, and for each $\tau \in \mathcal{K}$, we have:

$$p_n(\tau) \rightarrow p(\tau) = \mathcal{I}^\zeta \phi(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi(\sigma)(\tau).$$

Thus, $p \in \Delta_{\mathcal{G}}(y)$. Now, we demonstrate that there exists $\varepsilon := \|s\| \psi < 1$ such that $\mathcal{A}_d(\Delta_{\mathcal{G}}(y), \Delta_{\mathcal{G}}(\hat{y})) \leq \varepsilon \|y - \hat{y}\|_Y$ for each $y, \hat{y} \in C(\mathcal{K}, \mathbb{R})$. Let $y, \hat{y} \in C(\mathcal{K}, \mathbb{R})$ and $u_1 \in \Delta_{\mathcal{G}}(y)$. Then, there exists $\phi_1(\tau) \in \mathcal{G}(\tau, y(\tau), {}^C\mathcal{D}^\rho y(\tau), {}^C\mathcal{D}^{\rho+1} y(\tau))$ such that, for each $\tau \in \mathcal{K}$,

$$u_1(\tau) = \mathcal{I}^\zeta \phi_1(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi_1(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi_1(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi_1(\sigma)(\tau).$$

By \mathcal{F}_2 , we have:

$$\mathcal{A}_d(\mathcal{G}(\tau, y, z, w), \mathcal{G}(\tau, \hat{y}, \hat{z}, \hat{w})) \leq s(\tau) [|y - \hat{y}| + |z - \hat{z}| + |w - \hat{w}|].$$

Therefore, there exists $l \in \mathcal{G}(\tau, \hat{y}, \hat{z}, \hat{w})$ such that:

$$|\phi_1(\tau) - l| \leq s(\tau) [|y(\tau) - \hat{y}(\tau)| + |z(\tau) - \hat{z}(\tau)| + |w(\tau) - \hat{w}(\tau)|], \quad \tau \in \mathcal{K}.$$

Define $\mathcal{V} : \mathcal{K} \rightarrow \mathcal{T}(\mathbb{R})$ by:

$$\mathcal{V}(\tau) = \{l \in \mathbb{R} : |\phi_1(\tau) - l| \leq s(\tau) [|y(\tau) - \hat{y}(\tau)| + |z(\tau) - \hat{z}(\tau)| + |w(\tau) - \hat{w}(\tau)|]\}.$$

As the $\mathcal{V}(\tau) \cap \mathcal{G}(\tau, \hat{y}, \hat{z}, \hat{w})$ operator can be measurable (Proposition III.4 [35]), a $\phi_2(\tau)$ function exists, which is a selection measurable for $\mathcal{V}(\tau) \cap \mathcal{G}(\tau, \hat{y}, \hat{z}, \hat{w})$. Therefore, $\phi_2(\tau) \in \mathcal{G}(\tau, \hat{y}, \hat{z}, \hat{w})$, and for each $\tau \in \mathcal{K}$, we have $|\phi_1(\tau) - \phi_2(\tau)| \leq s(\tau) [|y(\tau) - \hat{y}(\tau)| + |z(\tau) - \hat{z}(\tau)| + |w(\tau) - \hat{w}(\tau)|]$. For each $\tau \in \mathcal{K}$, define:

$$u_2(\tau) = \mathcal{I}^\zeta \phi_2(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta \phi_2(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta \phi_2(\sigma) d\sigma \right] + \kappa_2(\tau) \mathcal{I}^{\zeta-1} \phi_2(\sigma)(\tau).$$

Thus,

$$\begin{aligned} |u_1(\tau) - u_2(\tau)| &= \mathcal{I}^\zeta |\phi_1 - \phi_2|(\sigma)(\tau) + \kappa_1(\tau) \left[\zeta^\rho \mathcal{J}^\omega \mathcal{I}^\zeta |\phi_1 - \phi_2|(\sigma)(\zeta) - \int_0^T \mathcal{I}^\zeta |\phi_1 - \phi_2|(\sigma) d\sigma \right] \\ &\quad + \kappa_2(\tau) \mathcal{I}^{\zeta-1} |\phi_1 - \phi_2|(\sigma)(\tau) \\ &\leq \frac{\|s\|}{\Gamma(\zeta + 1)} \left[(\tilde{\kappa}_2 \zeta T^{\zeta-1} + T^\zeta) + \tilde{\kappa}_1 \left(\frac{\zeta \zeta^{\rho+\omega}}{\rho^\omega} \frac{\Gamma(\frac{\zeta}{\rho} + 1)}{\Gamma(\frac{\zeta}{\rho} + \omega + 1)} + \frac{T^{\zeta+1}}{\zeta + 1} \right) \right] \|y - \hat{y}\|_Y, \end{aligned}$$

which yields $\|u_1 - u_2\| \leq \|s\| \psi \|y - \hat{y}\|_Y$. Furthermore, we have:

$$\|{}^C\mathcal{D}^\rho u_1(\tau) - {}^C\mathcal{D}^\rho u_2(\tau)\| \leq \|s\| \psi \|y - \hat{y}\|_Y.$$

In a similar manner, we have:

$$\|{}^C\mathcal{D}^{\varrho+1}u_1(\tau) - {}^C\mathcal{D}^{\varrho+1}u_2(\tau)\| \leq \|s\|\psi_3\|y - \hat{y}\|_Y.$$

As a result, we get $\|u_1 - u_2\| \leq \|s\|\psi\|y - \hat{y}\|_Y$. Likewise, swap the positions of y and \hat{y} ; we can get $\mathcal{A}_d(\Delta_G(y), \Delta_G(\hat{y})) \leq \|s\|\psi\|y - \hat{y}\|_Y$. Since Δ_G is a contraction by (21), it follows that it has a fixed point y by the lemma (see Lemma [36]), which is a solution of the problem (2) and (3). \square

5. Examples

Example 1. Consider a fractional BVP given by:

$${}^C\mathcal{D}^{\frac{94}{25}}y(\tau) = g(\tau, y(\tau), {}^C\mathcal{D}^{\frac{17}{50}}y(\tau), {}^C\mathcal{D}^{\frac{67}{50}}y(\tau)), \quad \tau \in [0, 1], \tag{22}$$

$$y(0) = 0, \quad y'(0) = 0, \quad \int_0^1 y(\sigma) d\sigma = \frac{11}{20} \frac{9}{25} \mathcal{J}^{\frac{23}{50}} y\left(\frac{31}{50}\right), \quad y'(1) = 0. \tag{23}$$

Here, $\varsigma = \frac{94}{25}$, $\varrho = \frac{17}{50}$, $\xi = \frac{11}{20}$, $\zeta = \frac{31}{50}$, $T = 1$, $\rho = \frac{9}{25}$, $\omega = \frac{23}{50}$. We have that with the data $\kappa_1 = 20.988457646948103$, $\kappa_2 = 1.5906878444726738$, $\nu_1 = 0.1992226279453819$, $\nu_2 = 0.17972085404600993$, $\vartheta = 0.2382261757441258$. Now, we demonstrate the outcomes by selecting different $\frac{17}{50}$ $\frac{67}{50}$ $g(\tau, y(\tau), {}^C\mathcal{D}^{\frac{17}{50}}y(\tau), {}^C\mathcal{D}^{\frac{67}{50}}y(\tau))$ values.

- (i) Consider:

$$g(\tau, y(\tau), {}^C\mathcal{D}^{\frac{17}{50}}y(\tau), {}^C\mathcal{D}^{\frac{67}{50}}y(\tau)) = \frac{1}{\sqrt{\tau^2 + 64}} \left(\frac{|y(\tau)|}{|y(\tau)| + 1} + \cos^2({}^C\mathcal{D}^{\frac{17}{50}}y(\tau)) + \sin({}^C\mathcal{D}^{\frac{67}{50}}y(\tau)) \right)$$

Clearly, $g(\tau, y(\tau), {}^C\mathcal{D}^{\frac{17}{50}}y(\tau), {}^C\mathcal{D}^{\frac{67}{50}}y(\tau)) = \frac{3}{\sqrt{\tau^2 + 64}} = \lambda(\tau)$. Therefore, the assumption of Theorem 1 holds. Hence, in Theorem 1, at least one solution has been found for the problem (22)–(23) on $[0, 1]$.

- (ii) To prove that Theorem 2 is valid, nonlinear function g is taken from the form:

$$g(\tau, y(\tau), {}^C\mathcal{D}^{\frac{17}{50}}y(\tau), {}^C\mathcal{D}^{\frac{67}{50}}y(\tau)) = \frac{1}{7(3+\tau^2)} \left(y(\tau) + \frac{|{}^C\mathcal{D}^{\frac{17}{50}}y(\tau)|}{17} + \cos({}^C\mathcal{D}^{\frac{67}{50}}y(\tau)) \right).$$

With the data given, we get $\kappa_1 = 20.988457646948103$, $\kappa_2 = 1.5906878444726738$, $\nu_1 = 0.1992226279453819$, $\nu_2 = 0.17972085404600993$, $\vartheta = 0.2382261757441258$, $\hat{\psi}_1 = 0.6727808331171499$, $\hat{\psi}_2 = 0.9525720580682466$, $\hat{\psi}_3 = 2.054165331280194$, and $\hat{\psi} = \max\{\hat{\psi}_1, \hat{\psi}_2, \hat{\psi}_3\} = 2.054165331280194$. By the resulting inequalities

$$\begin{aligned} |g(\tau, y(\tau), {}^C\mathcal{D}^{\frac{17}{50}}y(\tau), {}^C\mathcal{D}^{\frac{67}{50}}y(\tau)) - g(\tau, z(\tau), {}^C\mathcal{D}^{\frac{17}{50}}z(\tau), {}^C\mathcal{D}^{\frac{67}{50}}z(\tau))| &= \frac{1}{7(3+\tau^2)} (|y - z| + \\ |{}^C\mathcal{D}^{\frac{17}{50}}y(\tau) - {}^C\mathcal{D}^{\frac{17}{50}}z(\tau)| + |{}^C\mathcal{D}^{\frac{67}{50}}y(\tau) - {}^C\mathcal{D}^{\frac{67}{50}}z(\tau)|) &\leq \frac{1}{21} \|y - z\|, \end{aligned}$$

we have $\mathcal{P} = \frac{1}{21}$. Therefore, the hypothesis of Theorem 2 is fulfilled. Consequently, Theorem 2's assumption applies, and the problem (22)–(23) has at least one solution on $[0, 1]$.

- (iii) Let us consider:

$$g(\tau, y(\tau), {}^C D_{50}^{17} y(\tau), {}^C D_{50}^{67} y(\tau)) = \frac{1}{4\sqrt{\tau^2 + 16}} \left(y(\tau) + ({}^C D_{50}^{17} y(\tau)) + \frac{{}^C D_{50}^{67} y(\tau)}{|{}^C D_{50}^{67} y(\tau)| + 1} + 1 \right).$$

By the resulting inequalities,

$$|g(\tau, y(\tau), {}^C D_{50}^{17} y(\tau), {}^C D_{50}^{67} y(\tau)) - g(\tau, z(\tau), {}^C D_{50}^{17} z(\tau), {}^C D_{50}^{67} z(\tau))| = \frac{1}{4\sqrt{\tau^2 + 16}} (|y - z| + |{}^C D_{50}^{17} y(\tau) - {}^C D_{50}^{17} z(\tau)| + |{}^C D_{50}^{67} y(\tau) - {}^C D_{50}^{67} z(\tau)|) \leq \frac{1}{16} \|y - z\|, \text{ we have } Q = \frac{1}{16}. \text{ With the data given, we get } \kappa_1 = 20.988457646948103, \kappa_2 = 1.5906878444726738, \nu_1 = 0.1992226279453819, \nu_2 = 0.17972085404600993, \vartheta = 0.2382261757441258, \hat{\psi}_1 = 0.7322037594741672, \hat{\psi}_2 = 1.0485700387520187, \hat{\psi}_3 = 2.3824784252186952, \text{ and } \hat{\psi} = \max\{\hat{\psi}_1, \hat{\psi}_2, \hat{\psi}_3\} = 2.3824784252186952. \text{ Therefore, the hypothesis of Theorem 3 is fulfilled. Consequently, Theorem 3's assumption applies, and the problem (22)–(23) has a unique solution on } [0, 1].$$

Example 2. Consider the following problem of inclusions:

$${}^C D_{25}^{94} y(\tau) \in \mathcal{G}(\tau, y(\tau), {}^C D_{50}^{17} y(\tau), {}^C D_{50}^{67} y(\tau)), \tau \in [0, 1], \tag{24}$$

$$y(0) = 0, y'(0) = 0, \int_0^1 y(\sigma) d\sigma = \frac{11}{20} {}^J_{25}^{23} y\left(\frac{31}{50}\right), y'(1) = 0. \tag{25}$$

- (i) To show the illustration of Theorem 4, we take \mathcal{G} under consideration.

$$\mathcal{G}(\tau, y(\tau), {}^C D_{50}^{17} y(\tau), {}^C D_{50}^{67} y(\tau)) = \frac{1}{3\sqrt{625 + \tau^2}} \left(y(\tau) + \sin({}^C D_{50}^{17} y(\tau)) + \left(\frac{|{}^C D_{50}^{67} y(\tau)|}{67} + 1 \right) \right). \tag{26}$$

Apparently, $\gamma = \frac{1}{75}, \Delta(\|y\|_{\mathcal{Y}}) = 1 + \|y\|_{\mathcal{Y}}$, and state (\mathcal{F}_3) with $\mathcal{E} > \mathcal{E}_1$ is satisfied. Therefore, all criteria of Theorem 4 have been fulfilled, and at least one solution exists to the problem (24)–(25) with \mathcal{G} given in (26) on $[0, 1]$.

- (ii) Let us choose \mathcal{G} for the example of Theorem 5.

$$\mathcal{G}(\tau, y(\tau)) = \left[0, \frac{1}{8(4 + \tau^2)} \left(y(\tau) + \frac{|{}^C D_{50}^{17} y(\tau)|}{17} + \cos({}^C D_{50}^{67} y(\tau)) + \frac{1}{\tau + 1} \right) \right]. \tag{27}$$

It is clear that:

$$\mathcal{A}_d(\mathcal{G}(\tau, y), \mathcal{G}(\tau, \hat{y})) \leq \frac{1}{8(\tau^2 + 4)} \|y - \hat{y}\|_{\mathcal{Y}}.$$

Allowing $s(\tau) = \frac{1}{8(\tau^2+4)}$, we can check easily that $d(0, \mathcal{G}(\tau, 0)) \leq s(\tau)$ holds $\forall \tau \in [0, 1]$ and that $\|s\| \psi < 1$. Since we are satisfied with the assumptions of Theorem 5, we conclude that the problem (24)–(25) with \mathcal{G} as indicated by (27) has at least one solution for $[0, 1]$.

6. Discussion

We discussed the solutions of the existence and uniqueness for FDEs and inclusions supplemented by GRLFI boundary conditions. We used fixed point theorems for single-valued and multi-valued maps to evaluate the desired results. When we fixed the parameters involved in the problem (ρ, ω, ζ) (1)–(3), our results corresponded to certain specific problems. Suppose that taking $\rho = 1$ in the results provided, we are given the problems (1) with the form:

$$y(0) = y'(0) = 0, \int_0^T y(\sigma) d\sigma = \zeta \mathcal{I}^\omega y(\zeta), y'(T) = 0, \tag{28}$$

while the results are:

$$y(0) = y'(0) = 0, \int_0^T y(\sigma) d\sigma = 0, y'(T) = 0, \tag{29}$$

followed by $\zeta = 0$. When $\rho = \omega = 1$, we can obtain:

$$y(0) = y'(0) = 0, \int_0^T y(\sigma) d\sigma = \zeta \int_0^\zeta y(\sigma) d\sigma, y'(T) = 0. \tag{30}$$

Concerning the problem (1) with (28) instead of (3), we obtained the operator $\hat{Y} : \mathcal{Y} \rightarrow \mathcal{Y}$ defined by:

$$\begin{aligned} (\hat{Y}y)(\tau) = & \kappa_1(\tau) \left[\zeta \mathcal{I}^{\zeta+\omega} g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\zeta) \right. \\ & \left. - \int_0^T \mathcal{I}^\zeta g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma)) d\sigma \right] \\ & + \kappa_2(\tau) \mathcal{I}^{\zeta-1} g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau) \\ & + \mathcal{I}^\zeta g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau), \end{aligned}$$

where:

$$\begin{aligned} \kappa_1(\tau) = \frac{\tau^2 3T^2 - 2T\tau^3}{\vartheta}, \quad \kappa_2(\tau) = \frac{\tau^2 \nu_2 - \nu_1 \tau^3}{\vartheta}, \quad \vartheta = 3T^2 \nu_1 - 2T\nu_2, \\ \nu_1 = \frac{T^3}{3} - \frac{2\zeta \zeta^{\omega+2}}{\Gamma(3+\omega)}, \quad \nu_2 = \frac{T^4}{4} - \frac{6\zeta \zeta^{\omega+3}}{\Gamma(4+\omega)}. \end{aligned}$$

Similarly, the problem (1) related to operator $\bar{Y} : \mathcal{Y} \rightarrow \mathcal{Y}$ with conditions (29) rather than (3) is:

$$\begin{aligned} (\bar{Y}y)(\tau) = & \kappa_2(\tau) \mathcal{I}^{\zeta-1} g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau) \\ & - \kappa_1(\tau) \left[\int_0^T \mathcal{I}^\zeta g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma)) d\sigma \right] \\ & + \mathcal{I}^\zeta g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau), \end{aligned}$$

where:

$$\kappa_1(\tau) = \frac{\tau^2 3T^2 - 2T\tau^3}{\vartheta}, \quad \kappa_2(\tau) = \frac{\tau^2 \nu_2 - \nu_1 \tau^3}{\vartheta}, \quad \vartheta = 3T^2 \nu_1 - 2T\nu_2, \quad \nu_1 = \frac{T^3}{3}, \quad \nu_2 = \frac{T^4}{4}.$$

Likewise, the problem (1) related to operator $\tilde{Y} : \mathcal{Y} \rightarrow \mathcal{Y}$ with conditions (30) rather than (3) is:

$$\begin{aligned}
 (\tilde{Y}y)(\tau) = & \kappa_1(\tau) \left[\xi \int_0^\xi \mathcal{I}^\varsigma g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau) d\sigma \right. \\
 & \left. - \int_0^T \mathcal{I}^\varsigma g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma)) d\sigma \right] \\
 & + \kappa_2(\tau) \mathcal{I}^{\varsigma-1} g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau) \\
 & + \mathcal{I}^\varsigma g(\sigma, y(\sigma), {}^C\mathcal{D}^\varrho y(\sigma), {}^C\mathcal{D}^{\varrho+1} y(\sigma))(\tau),
 \end{aligned}$$

where:

$$\begin{aligned}
 \kappa_1(\tau) = \frac{\tau^2 3T^2 - 2T\tau^3}{\vartheta}, \quad \kappa_2(\tau) = \frac{\tau^2 v_2 - v_1 \tau^3}{\vartheta}, \quad \vartheta = 3T^2 v_1 - 2T v_2, \\
 v_1 = \frac{T^3 - \xi \zeta^3}{3}, \quad v_2 = \frac{T^4 - \xi \zeta^4}{4}.
 \end{aligned}$$

The existence and uniqueness of solutions for the new problems can be defined by the \hat{Y} , \bar{Y} , and \tilde{Y} operators similar to those obtained for (1)–(3).

Author Contributions: All authors contributed equally to this paper. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflicts of Interest.

References

1. Subramanian, M.; Kumar, A.; Gopal, T.N. Analysis of fractional boundary value problem with non-local integral strip boundary conditions. *Nonlinear Stud.* **2019**, *26*, 445–454.
2. Subramanian, M.; Kumar, A.R.V.; Gopal, T.N. Analysis of fractional boundary value problem with non local flux multi-point conditions on a Caputo fractional differential equation. *Mathematica* **2019**, 511. [CrossRef]
3. Duraisamy, P.; Kumar, A.V.; Gopal, T.N.; Subramanian, M. Influence of nonlocal discrete and integral boundary conditions involving Caputo derivative in boundary value problem. *J. Phys. Conf. Ser.* **2018**, *1139*, 012014. [CrossRef]
4. Kumar, A.V.; Duraisamy, P.; Gopal, T.N.; Subramanian, M. Analysis of fractional differential equation involving Caputo derivative with nonlocal discrete and multi-strip type boundary conditions. *J. Phys. Conf. Ser.* **2018**, *1139*, 012020. [CrossRef]
5. Muthaiah, S.; Murugesan, M.; Thangaraj, N.G. Existence of Solutions for Nonlocal Boundary Value Problem of Hadamard Fractional Differential Equations. *Adv. Theory Nonlinear Anal. Appl.* **2019**, *3*, 162–173. [CrossRef]
6. Ahmad, B.; Luca, R. Existence of solutions for sequential fractional integro-differential equations and inclusions with nonlocal boundary conditions. *Appl. Math. Comput.* **2018**, *339*, 516–534. [CrossRef]
7. Ahmad, B.; Alghanmi, M.; Ntouyas, S.K.; Alsaedi, A. A study of fractional differential equations and inclusions involving generalized Caputo-type derivative equipped with generalized fractional integral boundary conditions. *AIMS Math.* **2019**, *4*, 26. [CrossRef]
8. Agarwal, R.P.; Alsaedi, A.; Alghamdi, N.; Ntouyas, S.K.; Ahmad, B. Existence results for multi-term fractional differential equations with nonlocal multi-point and multi-strip boundary conditions. *Adv. Differ. Equ.* **2018**, *2018*, 342. [CrossRef]
9. Ahmad, B.; Alghanmi, M.; Ntouyas, S.K.; Alsaedi, A. Fractional differential equations involving generalized derivative with Stieltjes and fractional integral boundary conditions. *Appl. Math. Lett.* **2018**, *84*, 111–117. [CrossRef]
10. YILDIZ, T.A.; Khodabakhshi, N.; Baleanu, D. Analysis of mixed-order Caputo fractional system with nonlocal integral boundary condition. *Turk. J. Math.* **2018**, *42*, 1328–1337.

11. Uğurlu, E.; Baleanu, D.; Taş, K. On the solutions of a fractional boundary value problem. *Turk. J. Math.* **2018**, *42*, 1307–1311.
12. Ali, Z.; Zada, A.; Shah, K. Existence and stability analysis of three point boundary value problem. *Int. J. Appl. Comput. Math.* **2017**, *3*, 651–664. [[CrossRef](#)]
13. Zada, A.; Rizwan, R.; Xu, J.; Fu, Z. On implicit impulsive Langevin equation involving mixed order derivatives. *Adv. Differ. Equ.* **2019**, *2019*, 489. [[CrossRef](#)]
14. Ali, A.; Shah, K.; Jarad, F.; Gupta, V.; Abdeljawad, T. Existence and stability analysis to a coupled system of implicit type impulsive boundary value problems of fractional-order differential equations. *Adv. Differ. Equ.* **2019**, *2019*, 101. [[CrossRef](#)]
15. Shah, K.; Ali, A.; Bushnaq, S. Hyers-Ulam stability analysis to implicit Cauchy problem of fractional differential equations with impulsive conditions. *Math. Methods Appl. Sci.* **2018**, *41*, 8329–8343. [[CrossRef](#)]
16. Subramanian, M.; Kumar, A.V.; Gopal, T.N. Influence of coupled nonlocal slit-strip conditions involving Caputo derivative in fractional boundary value problem. *Discontin. Nonlinearity Complex.* **2019**, *8*, 429–445.
17. Subramanian, M.; Kumar, A.V.; Gopal, T.N. A fundamental approach on non-integer order differential equation *Discontin. Nonlinearity Complex.* **2019**, *8*, 189–199.
18. Subramanian, M.; Kumar, A.V.; Gopal, T.N. A writ large analysis of complex order coupled differential equations in the course of coupled non-local multi-point boundary conditions. *Adv. Stud. Contemp. Math.* **2019**, *29*, 505–520.
19. Subramanian, M.; Kumar, A.V.; Gopal, T.N. A strategic view on the consequences of classical integral sub-strips and coupled nonlocal multi-point boundary conditions on a combined Caputo fractional differential equation. *Proc. Jangjeon Math. Soc.* **2019**, *22*, 437–453.
20. Smart, D.R. *Fixed Point Theorems*; CUP Archive: Cambridge, UK, 1980; Volume 66.
21. Yong, Z.; Jinrong, W.; Lu, Z. *Basic Theory of Fractional Differential Equations*; World Scientific: Singapore, 2016.
22. Granas, A.; Dugundji, J. *Fixed Point Theory*; Springer Science & Business Media: Berlin/Heidelberg, Germany, 2013.
23. Kilbas, A.A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of fRactional Differential Equations*; Elsevier Science Limited: Amsterdam, The Netherlands, 2006; Volume 204.
24. Machado, J.T.; Kiryakova, V.; Mainardi, F. Recent history of fractional calculus. *Commun. Nonlinear Sci. Numer. Simul.* **2011**, *16*, 1140–1153. [[CrossRef](#)]
25. Valério, D.; Machado, J.T.; Kiryakova, V. Some pioneers of the applications of fractional calculus. *Fract. Calc. Appl. Anal.* **2014**, *17*, 552–578. [[CrossRef](#)]
26. Jiang, C.; Zada, A.; Şenel, M.T.; Li, T. Synchronization of bidirectional N-coupled fractional-order chaotic systems with ring connection based on antisymmetric structure. *Adv. Differ. Equ.* **2019**, *2019*, 1–16. [[CrossRef](#)]
27. Jiang, C.; Zhang, F.; Li, T. Synchronization and antisynchronization of N-coupled fractional-order complex chaotic systems with ring connection. *Math. Methods Appl. Sci.* **2018**, *41*, 2625–2638. [[CrossRef](#)]
28. Wang, P.; Li, C.; Zhang, J.; Li, T. Quasilinearization method for first-order impulsive integro-differential equations. *Electron. J. Differ. Equ* **2019**, *46*, 2019.
29. Ahmad, B.; Ntouyas, S.K.; Zhou, Y.; Alsaedi, A. A study of fractional differential equations and inclusions with nonlocal Erdélyi–Kober type integral boundary conditions. *Bull. Iran. Math. Soc.* **2018**, *44*, 1315–1328. [[CrossRef](#)]
30. Ntouyas, S.K.; Etemad, S.; Tariboon, J. Existence of solutions for fractional differential inclusions with integral boundary conditions. *Bound. Value Probl.* **2015**, *2015*, 1–14. [[CrossRef](#)]
31. Salem, S.; Alsaedi, A.; Ahmad, B.; Al-Yami, M. Arbitrary order fractional differential equations and inclusions with new integro-multipoint boundary conditions. *Adv. Differ. Equ.* **2018**, *2018*, 1–19. [[CrossRef](#)]
32. Deimling, K. *Multivalued Differential Equations*; Walter de Gruyter: Berlin, Germany, 2011; Volume 1.
33. Hu, S.; Papageorgiou, N. *Handbook of Multivalued Analysis (Theory)*; Kluwer Academic Publishers: Dordrecht, The Netherlands; Boston, MA, USA; London, UK, 1997.
34. Lasota, A.; Opial, Z. An application of Kakutani-ky fan theorem in theory of ordinary differential equations. *Bull. Acad. Pol. Sci. Ser. Sci. Math. Astron. Phys.* **1965**, *13*, 781.

35. Castaing, C.; Valadier, M. *Convex Analysis and Measurable Multifunctions*; Springer: Berlin/Heidelberg, Germany, 2006; Volume 580.
36. Covitz, H.; Nadler, S. Multi-valued contraction mappings in generalized metric spaces. *Isr. J. Math.* **1970**, *8*, 5–11. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).