



## Review

# The Role and Application of MAdCAM-1/ $\alpha 4\beta 7$ -Induced Lymphocyte Migration in Inflammatory Enterohepatic Diseases

Chuchu Yu <sup>1</sup>, Yuqing Pan <sup>1</sup>, Aojie Mao <sup>1</sup>, Yu Zhao <sup>1,2</sup>, Qiaohong Liu <sup>4,\*</sup> and Yiyang Hu <sup>1,2,3,4,\*</sup><sup>1</sup> Institute of Liver Diseases, Shuguang Hospital Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China<sup>2</sup> Key Laboratory of Liver and Kidney Diseases (Ministry of Education), Shanghai 201203, China<sup>3</sup> Shanghai Clinical Key Laboratory of Traditional Chinese Medicine, Shanghai 201203, China<sup>4</sup> Clinical Research Center, Shuguang Hospital Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

\* Correspondence: sg4080@shutcm.edu.cn (Q.L.); huyiyang@shutcm.edu.cn (Y.H.)

## Abstract

Inflammation drives the development of multiple inflammatory enterohepatic diseases. The recruitment of immune cells to inflammatory tissues is essential for maintaining immune homeostasis, mediating immune responses and regulating inflammation. MAdCAM-1/ $\alpha 4\beta 7$  is a pair of homing ligand and receptor that plays important roles in lymphocyte migration. Their binding induces lymphocytes to cross endothelial structures into corresponding lymphoid tissues, contributing to the inflammatory response. Aberrant lymphocyte migration due to excessive binding is closely related to the occurrence and development of inflammatory bowel disease and liver inflammation. In this review, we focus on the activation of  $\alpha 4\beta 7$  and binding to MAdCAM-1 how to induce the migration of multiple kinds of lymphocytes. Additionally, we describe the intestinal microbiota and its metabolites associated with MAdCAM-1/ $\alpha 4\beta 7$  in inflammatory enterohepatic diseases. We also discuss the current status of the development of monoclonal antibodies and small molecule drugs targeting MAdCAM-1/ $\alpha 4\beta 7$  for the remission and treatment of inflammatory enterohepatic disease. Future research should focus on enhancing hepatic specificity and conducting well-designed clinical trials for inflammatory liver diseases to confirm therapeutic efficacy.

**Keywords:** cell adhesion molecules; mucosal address cell adhesion molecule 1;  $\alpha 4\beta 7$ ; lymphocyte migration; inflammatory enterohepatic diseases



Academic Editor: Giuseppe Murdaca

Received: 3 September 2025

Revised: 14 October 2025

Accepted: 21 October 2025

Published: 29 October 2025

**Citation:** Yu, C.; Pan, Y.; Mao, A.; Zhao, Y.; Liu, Q.; Hu, Y. The Role and Application of MAdCAM-1/ $\alpha 4\beta 7$ -Induced Lymphocyte Migration in Inflammatory Enterohepatic Diseases. *Biomedicines* **2025**, *13*, 2659. <https://doi.org/10.3390/biomedicines13112659>

**Copyright:** © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Inflammation drives the development of multiple inflammatory gut-liver conditions. In fatty liver disease, primary sclerosing cholangitis, liver cirrhosis, and other liver diseases, the inflammatory process is an integral part of the progression of these diseases, in addition to their respective etiologies of fat accumulation, cholestasis, and fibrosis formation [1,2]. While the pathogenesis of inflammatory bowel disease (IBD) involves multiple factors including genetics, environment, immune responses, and gut microbiota [3], the central role of inflammation in driving the progression of both ulcerative colitis (UC) and Crohn's disease (CD) is well-established. So, suppressing inflammation and controlling its progression is important in the treatment of these enterohepatic diseases.

Dysregulation of lymphocyte transport plays an important role in the inflammatory process in these diseases [4]. The ability of lymphocytes to selectively enter specific lymph

phoid tissues is known as “homing”, and integrins and their ligands play a key role among the various molecules responsible for regulating lymphocyte homing, a complex process of specific migration [5]. A central pathway is mediated by integrin  $\alpha 4\beta 7$  following its activation and binding to MAdCAM-1, which directs the migration of various lymphocyte populations, including T cells, whereas overexpression of MAdCAM-1 or excess lymphocyte homing induced by MAdCAM-1/ $\alpha 4\beta 7$  also drives the pathogenesis of intestinal inflammatory disorders, including IBD. Through gut-liver axis communication, these lymphocytes migrate to hepatic tissue, further inducing inflammatory responses that contribute to the pathogenesis of inflammatory liver diseases such as primary sclerosing cholangitis (PSC) and non-alcoholic steatohepatitis (NASH). This continuum of pathological events further confirms the intimate connection between the gut and the liver within the inflammatory immune mechanisms of inflammatory enterohepatic diseases. Given the pivotal role of the MAdCAM-1/ $\alpha 4\beta 7$  pathway in sustaining this gut-liver inflammatory axis, monoclonal antibodies targeting this pathway, such as Vedolizumab, have emerged as a key therapeutic strategy for IBD [6].

This paper is for reviewing and emphasizing the role of MAdCAM-1/ $\alpha 4\beta 7$  in lymphocyte migration and inflammatory enterohepatic diseases. In addition, we will highlight specific intestinal microbiota and their metabolites associated with the expression of MAdCAM-1/ $\alpha 4\beta 7$ , as well as monoclonal antibody drugs and small molecule drugs targeting MAdCAM-1/ $\alpha 4\beta 7$ . Understanding MAdCAM-1/ $\alpha 4\beta 7$ -regulated lymphocyte migration could open new paths for the treatment of gut-liver inflammatory diseases.

## 2. Structure and Expression of MAdCAM-1/ $\alpha 4\beta 7$

Integrins are transmembrane heterodimers that act as major cell adhesion transmembrane receptors and 24 integrin heterodimers have been reported, consisting of 18  $\alpha$ -subunits and 8  $\beta$ -subunits [7], each of which is a surface glycoprotein greater than 1600 amino acids in length, and consists of a short intracellular region, a helical transmembrane region, and a large extracellular region. The extracellular structural domain of each subunit serves as the ligand-binding site for the heterodimer [8]; the extracellular structural domain of  $\alpha 4$  (180 kDa) contains: a hepta-bladed  $\beta$ -propeller, a thigh and two calf structural domains. It is worth noting that unlike other  $\alpha$ -chains in the integrin family,  $\alpha 4$  does not have an  $\alpha I$  structural domain, and forms a binding pocket directly with the  $\beta I$  structural domain in the  $\beta$ -chain via the  $\beta$ -propeller structure. The extracellular structural domain of  $\beta 7$  (130 kDa) contains an insertion of heterodimeric  $\beta I$  structural domain, plexin-semaphorin- integrin (PSI), four cysteine-rich epidermal growth factor modules, and a  $\beta$ -tail structural domain (Figure 1) [9]. Integrins can be categorized into four types based on cell type or ligand specificity: leukocyte adhesion integrins, RGD (Arginine-Glycine-Aspartic Acid) -binding integrins, collagen-binding integrins, and laminin-binding integrins, of which leukocyte adhesion integrins are mainly involved in the regulation of inflammation [10].  $\alpha 4\beta 7$  belongs to leukocyte adhesion integrins, which are expressed on a wide range of leukocytes, such as naïve and memory CD4<sup>+</sup>, CD8<sup>+</sup> T cells, B cells, eosinophils, NK-cells, and a small number of monocytes [11]. The expression and binding affinity of  $\alpha 4\beta 7$  are enhanced by chemokines, including CXCL12, CCL12, and CCL25, with CXCL12 demonstrating the most pronounced effect [12].




**Figure 1. Schematic structure and affinity regulation of  $\alpha 4\beta 7$ .**  $\alpha 4\beta 7$  is bent in a quiescent state and the ligand-binding pocket is hidden from MAdCAM-1 binding. The “inside-out signaling” triggered by the binding of chemokines and their receptors induces conformational changes in the integrin structure, extending the domains and exposing the hidden ligand-binding pocket, which increases the affinity of  $\alpha 4\beta 7$  and begins to bind to MAdCAM-1. This ligand binding leads to integrin-mediated “outside-in signaling”, affecting cytoskeletal rearrangement, further aggregation of  $\alpha 4\beta 7$ , and a change in ‘valency,’ which enhances adhesion strength [13].

The interaction of integrins with the Immunoglobulin superfamily (IgSF) is particularly important during lymphocyte homing [14]. Mucosal address cell adhesion molecule 1 (MAdCAM-1), the ligand for  $\alpha 4\beta 7$ , is a member of the IgSF family. Other members include intercellular adhesion molecule (ICAM), vascular cell adhesion molecule-1 (VCAM-1), and platelet/endothelial cell adhesion molecule-1 (PECAM-1), etc. [15,16]. MAdCAM-1 contains two distal immunoglobulin-like structural domains, a mucin-like region, a transmembrane structural domain, and a cytoplasmic structural domain. It binds to integrin  $\alpha 4\beta 7$  via its immunoglobulin-like structural domain and to L-selectin through its mucin-like region when properly O-glycosylated (Figure 2) [15]. MAdCAM-1 is expressed on high endothelial venule (HEV) endothelial cells in gut-associated lymphoid tissues (GALT), as well as in the small intestine and colon of both healthy individuals and patients with IBD. Its expression is upregulated during active inflammation. While absent in normal liver tissues, MAdCAM-1 is expressed on portal vein endothelial cells and hepatic sinusoidal endothelial cells in inflammatory liver disease [12]. In humans and mice, its expression is induced by TNF- $\alpha$ , which is mediated through the NF- $\kappa$ B pathway [17,18].




**Figure 2. Schematic structure of MAdCAM-1.** MAdCAM-1 contains two distal immunoglobulin-like structural domains, D1 and D2, which constitute the key structural elements for  $\alpha 4\beta 7$  binding; a mucin-like region which is the site of O-linked glycosylation and the L-selectin-binding domain; an  $\alpha$ -helix transmembrane domain and a shorter cytoplasmic domain.

### 3. Activation of $\alpha 4\beta 7$ and Binding to MAdCAM-1 Induces Lymphocyte Migration

#### 3.1. T Cell

Naïve T cells released from the thymus need to enter secondary lymphoid organs to encounter their cognate antigens, at which point the naïve T cells express CD62L (L-selectin), C-C-chemokine receptor (CCR7), and low levels of  $\alpha 4\beta 7$  integrins [19]. However, because integrins are inactive (bent conformation) on resting circulating immune cells and do not readily interact with endothelial cells [20], the adhesion cascade begins with “rolling” mediated by “tethering”, where cells need to be pre-recruited to the canalicular wall and slowed down, which is mediated by selectins on lymphocytes and selectin ligands on the endothelium. In addition, the low-affinity linkage of  $\alpha 4\beta 7$  to MAdCAM-1 is involved in rolling. These interactions establish loose contacts between cells, which are repeatedly separated and re-established, resulting in leukocytes being able to have ample opportunity to sense chemokines while slowly rolling along the vessel wall [21]. These chemokines and G protein-coupled chemokine receptors form a rapid intracellular “inside-out” signaling [22] that induces conformational changes in integrins. These changes involve: the extension of the integrin’s extracellular region, the segregation of its subunits and transmembrane domains, and a rearrangement of the  $\alpha$ - $\beta$  chains in the area that binds ligands. Consequently, the integrin shifts from a compact, curved shape to an open one. This open conformation results in a higher binding affinity. When  $\alpha 4\beta 7$  binds to MAdCAM-1, cellular “outside-in” signaling occurs, resulting in a change in “valence” at the contact site, an increase in the number of receptor-ligand bonds [13], and a further strengthening of cellular adhesion, which allows T cells to enter into the intestinal lymphoid tissue via HEVs [23]. After entering the gut-associated lymphoid tissue and meeting with antigen-presenting dendritic cells, the dendritic cells convert retinol to retinoic acid by producing retinaldehyde dehydrogenase. The retinoic acid then activates the nuclear receptor, retinoic acid receptor (RAR), which forms a heterodimer with the retinoid X receptor (RXR) [24]. This complex then binds to retinoic acid response elements (RAREs) in the promoter regions of genes encoding  $\alpha 4\beta 7$  and CCR9, inducing their transcription in T lymphocytes and

promoting the differentiation of naïve T cells into effector T cells with high  $\alpha 4\beta 7$  expression. These effector T cells subsequently return to the circulation via efferent lymphatics, and when once again following the circulation to the GALT, at which point they can be directly implanted in the lamina propria through the high affinity of  $\alpha 4\beta 7$  for MAdCAM-1 and the binding of CCR9 and CCL25 [12]. In addition, it has been reported that CCL25 induces a more extended active conformation of  $\alpha 4\beta 7$  by binding to CCR9, resulting in an increased affinity of  $\alpha 4\beta 7$  for MAdCAM-1 [25,26]. This process is summarized in Figure 3.




**Figure 3. Activation of  $\alpha 4\beta 7$  on naïve T cells and migration of naïve and mature T cells mediated by binding of  $\alpha 4\beta 7$  to MAdCAM-1.** ① After leaving the thymus, Naïve T lymphocytes enter the circulatory system and begin to slowly roll along endothelial cells through the “tethering” mainly mediated by L-selectin when passing through HEV. ② After activation of  $\alpha 4\beta 7$  mediated by CCR7, CCL19 and CCL21, the binding of  $\alpha 4\beta 7$  and MAdCAM-1 induces T cells to enter the Peyer’s patches (PPs) and mesenteric lymph nodes (MLN). ③ T cells encounter antigen-presenting dendritic cells in PPs and MLN that encode  $\alpha 4\beta 7$  expression in the presence of retinoic acid, differentiate into effector T cells highly expressing  $\alpha 4\beta 7$ , and ④ subsequently return to the circulatory system via efferent lymphatics. ⑤ At this point, effector T cells that are highly expressing  $\alpha 4\beta 7$  can migrate into the lamina propria directly in binding to MAdCAM-1. ⑥ Effector T cells highly expressing  $\alpha 4\beta 7$  enter the liver parenchyma via hepatic sinusoidal endothelial cells in response to the inflammatory response after MAdCAM-1/ $\alpha 4\beta 7$  binding.

T cells isolated from the livers of mice with chronic DSS (Dextran Sulfate Sodium Salt) colitis, including both CD4+ and CD8+ T cells, expressed integrin  $\alpha 4\beta 7$  at higher levels than in control mice [27]. The transfer of intestinal pathogens beyond the mucosal barrier to the portal circulation and the liver allows MAdCAM-1, which is not expressed in the normal liver, to be expressed in the inflammatory environment. Lymphocyte recruitment requires

not only MAdCAM-1/ $\alpha 4\beta 7$  interactions, but also specific chemokine signaling provided by CCL25, whose expression is usually restricted to the gut and thymus. However, it has been reported that CCR9+ (receptor for CCL25) cells make up to 20% of the lymphocytes that infiltrate the liver in PSC [28]. In summary, long-lived mucosal T cells are recruited to the liver in response to the abnormal expression of endothelial adhesion molecules and chemokines, which are typically restricted to the intestine [29]. Another evidence that  $\alpha 4\beta 7$ + T lymphocytes in the liver are initially activated in the intestine is that dendritic cells isolated from the liver or hepatic lymph nodes of patients with PSC fail to induce  $\alpha 4\beta 7$  and CCR9 expression in the absence of exogenous retinoic acid [30]. In contrast to the homing of lymphocytes in the intestines, although integrin  $\alpha 4\beta 7$  was detected on intestinal-activated CD8+ T cells in iFABP-OVA mice expressing ovalbumin (OVA) in small intestinal enterocytes and on hepatic-activated CD8+ T cells in TF-OVA mice expressing ovalbumin in hepatocytes, the co-expression of  $\alpha 4\beta 7$  and CCR9 was restricted to intestinal-activated CD8+ T cells, which exhibited a stronger migration potential. And this T-cell migration appeared to be unidirectional: gut-activated T cells were present in both the liver and intestine, whereas liver-activated T cells aggregated only in the liver [31]. Unlike homing in the gut, the low blood flow through the narrow vessels in the hepatic sinusoids means that T cells already have the opportunity to continuously interact with the endothelium, rendering the selectins that are important in intestinal homing redundant. However, the currently undefined process of “tethering” may still be necessary [32]. Furthermore, upregulation of the extracellular enzyme vascular adhesion protein-1 (VAP-1) during chronic inflammation induces MAdCAM-1 upregulation on the sinusoidal endothelium through deamination of methylamine (MA) in addition to directly supporting lymphocyte adhesion [33,34], as shown in Figure 3.

### 3.2. Migration of Lymphocytes Other than T Cells

#### 3.2.1. B Cell

B-lymphocyte homing appears to be controlled in a manner similar to that of T cells: naïve and memory B cells are recruited to GALT through the interaction of CD62L, CCR7, and  $\alpha 4\beta 7$  with CCL21 and MAdCAM-1 expressed on HEVs. Upregulation of intestinal homing markers, such as CCR9 and  $\alpha 4\beta 7$  integrins, on B-lymphocytes is mediated by dendritic cells that secrete retinoic acid (RA) [23]. Activated plasmablasts exhibited upregulated CCR9, CCR10, and  $\alpha 4\beta 7$ , with  $\alpha 4\beta 7$  being critical for their intestinal homing. In  $\beta 7^{-/-}$  mice, the number of IgA-secreting plasma cells in the lamina propria was reduced, and these mice were incapable of generating an IgA response, which impairs mucosal immunity [35]. Furthermore, the serum and intestinal levels of IgA following stimulation with T cell-dependent antigens were significantly lower in  $\beta 7^{-/-}$  mice compared to wild-type mice [35,36]. Similarly, MAdCAM-1 $^{-/-}$  mice show a failure to generate an IgA response [37]. Unlike T-cell migration, the homing efficiency of B cells is lower than that of T cells because the level of L-selectin expression in B cells is more than 50% lower than that of T cells, despite the fact that B cells can be activated in HEV by two different chemokines and corresponding receptors (CCR7 and CCL21, CXCR4 and CXL12) [19,38]. Additionally, B cells, while capable of homing to the PPs to support mucosal immunity, are less efficient at returning to peripheral lymph nodes compared to T cells. Instead, B cells preferentially migrate to the spleen, whereas T cells primarily localize to the lymph nodes [39]. CD22 is more important for B-cell migration, but its absence does not affect T-cell migration [40].

#### 3.2.2. Eosinophil

Most studies on eosinophils have focused on rolling and adhesion mediated by the interaction of  $\alpha 4\beta 1$  with VCAM1 and fibronectin, but an important role of  $\alpha 4\beta 7$  for eosinophil

recruitment into the gut under inflammatory conditions has also been reported: experimental gastrointestinal anergy induction in mice revealed that, eosinophil levels were significantly reduced in the colon of mice lacking  $\beta 7$  compared to wild-type mice [41]. In a clinical observation, vedolizumab treatment for one year led to improved global assessment scores and reduced eosinophil density in three of four eosinophilic gastrointestinal diseases (EGID) patients [42].

### 3.2.3. Treg17 Cell

Treg17 cells, a unique subset of regulatory T cells producing IL-17, typically coexpress the canonical Treg marker Foxp3 alongside the Th17-characteristic cytokine IL-17A and transcription factor ROR $\gamma$ t. The development of both Treg17 cells and their effector Th17 counterparts depends on the same lineage-specific master transcription factor, STAT3, reflecting a delicate balance between immune tolerance and inflammation [43]. Current research has moved beyond the simplistic definition of Foxp3+ IL-17+ cells and now aims to identify more specific surface marker combinations—such as the p-mTOR<sup>high</sup> Treg17 population defined by Natarajan et al., which plays a protective role during mucosal infection [44]. Notably, administration of antibiotics resulted in down-regulation of MAdCAM-1 expression levels in the ileum, PPs and MLN, while Treg17 cells migrated out of the ileum to extra-intestinal tumor as well as to tumor-draining lymph nodes. Ectopic expression of MAdCAM-1 in mouse liver reduces the accumulation of  $\alpha 4\beta 7$ + Treg17 cells at tumor, thereby improving the therapeutic efficacy of immunotherapy [45].

### 3.2.4. Non-Classical Monocytes

They express high levels of the CX3C chemokine receptor 1 (CX3CR1), which competitively inhibits CX3C chemokine ligand 1 (CX3CL1) and CCL3 from activating other, more destructive inflammatory cells, thereby protecting the endothelium and barrier system [46]. In dynamic adhesion assays, increased adhesion of peripheral blood monocytes from IBD patients to MAdCAM-1 was observed. Treatment with vedolizumab reduced monocyte adhesion to MAdCAM-1 to background levels. Further analysis revealed that the dynamic adhesion rate of non-classical monocytes to MAdCAM-1 was significantly higher than that of classical monocytes. Vedolizumab markedly reduced the adhesion rate of non-classical monocytes but had no significant effect on classical monocytes. Similarly, after transferring fluorescence-labeled classical and non-classical monocytes into the ileocolic artery of recipient mice, results showed that significantly more non-classical monocytes homed to the intestine compared to classical monocytes. Anti- $\alpha 4\beta 7$  integrin treatment significantly reduced the accumulation of non-classical monocytes in the colon, with no notable effect on classical monocytes. These findings collectively indicate that  $\alpha 4\beta 7$  expression on non-classical monocytes plays a crucial role in mediating dynamic adhesion and gut homing in vivo [47].

### 3.2.5. Innate Lymphoid Cell

Innate lymphoid cells (ILCs) do not express antigen-specific receptors; instead, they function primarily through the secretion of specific cytokines in a manner analogous to T cells, playing critical roles in host defense against pathogens [48]. Innate lymphoid cells (ILCs) types 1 and 3 may upregulate  $\alpha 4\beta 7$  via retinoic acid-mediated gut-specific imprinting thereby accomplishing the “homing receptor switch”. In contrast, expression of  $\alpha 4\beta 7$  on ILCs2 is acquired in a RA-independent manner in the bone marrow and ILCs2 can migrate directly to the intestine [49].

#### 4. MAdCAM-1/ $\alpha$ 4 $\beta$ 7-Induced Lymphocyte Homing Excess in Inflammatory Enterohepatic Disease

In addition to the anatomical homology of the liver and intestines originating from the endoderm, the theory of ‘intestine-liver axis’ proposed by Marshall in 1998 elaborated the mutual regulation and influence between the liver and intestine through the portal vein system, which was even more proved to have metabolic interactions and immune correlations. For example, impaired intestinal barrier function due to multiple causes allows bacterial metabolites to migrate and reach the liver through the portal vein, thereby exacerbating hepatic inflammation and leading to NASH, cirrhosis, and other diseases [50]. Whereas liver insufficiency can cause dysfunction in intestinal secretion, absorption, motility, barrier, and other aspects. For example, cirrhosis can lead to the formation of portal hypertension, which in turn can cause alterations in intestinal dynamics and dysbiosis of intestinal microbiota [51]. In addition to the gut-liver interconnection, activated immune cells disrupt tight junction proteins between epithelial cells, leading to increased intestinal permeability, and interfere with normal metabolic functions of hepatocytes in the liver, such as promoting lipid synthesis and inhibiting fatty acid  $\beta$ -oxidation. Factors released by inflammatory cells and damaged hepatocytes, including TGF- $\beta$ , activate hepatic stellate cells, further aggravating disease progression. Thus, the synergistic interactions between immune and non-immune cells also play a critical role in the initiation and progression of inflammation [52,53].

##### 4.1. Intestinal Diseases

In 1997, in attempting to determine whether MAdCAM-1 or its receptor  $\alpha$ 4 $\beta$ 7 was a suitable target for therapeutic intervention in gut-associated inflammation, D’Picarella et al. found that  $\beta$ 7- and MAdCAM-1-specific antibodies blocked lymphocyte recruitment to the colon in a colitis model, significantly reducing disease severity. This suggests that the interaction between MAdCAM-1 and  $\alpha$ 4 $\beta$ 7 is closely related to the recruitment of lymphocytes to the intestine in chronic inflammatory diseases, and blocking their binding could be a relevant therapeutic target for inflammatory bowel disease patients [54]. In 2006, Christopher Bachmann et al. developed a novel intravascular ultrasound contrast agent targeting MAdCAM-1 to specifically detect and quantify intestinal inflammation in experimental ileitis. This agent provides a reliable, noninvasive method for diagnosing and monitoring intestinal inflammation [55]. Over the years, an increasing number of studies on inflammatory bowel disease have focused on this pathway. For example, angiotensin II type 1 receptor blockade can modulate TNF- $\alpha$ -induced MAdCAM-1 expression by inhibiting NF- $\kappa$ B nuclear translocation, thereby ameliorating colitis [17]. It was not until 2014 that the Food and Drug Administration (FDA) approved Vedolizumab for the treatment of adult patients with moderate to severe inflammatory bowel disease who have had an inadequate response, loss of response, or intolerance to one or more of the standard therapies (corticosteroids, immunomodulators, or tumor necrosis factor-alpha inhibitors) [56]. Moreover, a vaccination study in healthy volunteers showed that vedolizumab attenuated the response to enteral antigen challenge (oral cholera vaccine) without affecting the response to parenteral antigen challenge (intramuscular hepatitis B vaccine), indicating a high degree of gut-specific anti-inflammatory activity for Vedolizumab [57]. In addition to inflammatory bowel disease, a review of Vedolizumab for the treatment of non-inflammatory bowel disease-associated enteropathy demonstrated that Vedolizumab has clinical, endoscopic, and histologic improvements in patients with non-IBD-associated inflammatory bowel disease (e.g., microscopic colitis) as well as inflammatory disorders of the small bowel, including autoimmune enteropathy and common variable immunodeficiency-associated enteritis, suggesting that its role in the gut is not limited to inflammatory bowel disease [58].

## 4.2. Liver Diseases

### 4.2.1. Primary Sclerosing Cholangitis

PSC is an inflammation of the intra- and extra-hepatic bile ducts leading to destruction of the bile duct epithelium and end-stage liver disease [59]. Approximately 70% of patients with PSC have or will develop concomitant IBD, with the majority of PSC-IBD patients suffering from ulcerative colitis (72%). Colon resection prior to the diagnosis of PSC reduces the risk of liver transplantation or death [59,60]. The significant association between PSC and IBD suggests that long-lived memory T lymphocytes expressing intestinal homing markers migrate to the liver via aberrantly expressed mucosal adhesion molecules and chemokines. Analysis of common T-cell receptor clonotypes in paired colon, liver, and blood samples revealed a higher degree of overlap between PSC-IBD samples than between normal intestinal and liver samples. This suggests that memory T-cells in the intestines and livers of PSC-IBD patients are reacting to common antigens [61]. Consistent with this, PSC is characterized by a peribiliary infiltrate composed mainly of T lymphocytes, a large proportion of which express the intestinal homing integrin  $\alpha 4\beta 7$  [62]. Several studies have also reported that MAdCAM-1 is aberrantly expressed on PSC hepatic endothelial cells in both large and small portal veins [63–65]. For example, the proportion of endothelial cells staining positively for MAdCAM-1 calculated by H score was significantly higher in liver tissues of patients with long-term PSC (PSC-IBDLT). The proportion of  $\beta 7$ -positive T cells was also significantly higher than in short-term patients and controls, with a positive correlation between disease duration and *Itgb7* expression. The enzymatic activity of VAP-1 was higher in liver and intestinal tissues of PSC patients compared to controls, promoting the adhesion of intestinal-derived lymphocytes to hepatic endothelial cells in a substrate-dependent manner [34]. Seidel, D. et al. also found that activated CD8<sup>+</sup> T cells in mouse GALT induce immune-mediated cholangitis in an antigen (OVA)-dependent manner [27].

### 4.2.2. Other Inflammatory Liver Diseases

Although the role of MAdCAM-1/ $\alpha 4\beta 7$  in PSC has been widely reported, this theory still lacks comparison with other chronic liver diseases (CLD), Jonathon J Graham et al. found that hepatic expression of MAdCAM-1 was up-regulated in all CLD, including PSC, primary biliary cholangitis (PBC), alcoholic liver disease (ASH), NASH, and Viral hepatitis C (HCV), and that the frequency of hepatic T-cells expressing  $\alpha 4\beta 7$  was increased, as compared to normal liver, suggesting that aberrant hepatic recruitment of T-cells of intestinal origin is not unique to PSC exclusively [66].

Increased infiltration of  $\alpha 4\beta 7$ <sup>+</sup> CD4<sup>+</sup> T cells and increased MAdCAM-1 expression in the liver and intestines of NASH mice. Treatment of mice with a highly specific neutralizing monoclonal antibody against  $\alpha 4\beta 7$  inhibited  $\alpha 4\beta 7$ <sup>+</sup> CD4<sup>+</sup> T cell recruitment in the colon and liver of NASH mice, attenuated hepatic and colonic mucosal inflammation, and protected the colonic epithelial barrier function to improve NASH. Similarly, MAdCAM-1, *Itga4*, and *Itgb7* expression was higher in liver tissues of NASH patients than in controls [67]. However, Hannah K. Drescher et al. found that unlike the protective effect of MAdCAM-1 deficiency,  $\beta 7$  integrin deficiency triggered stronger hepatic immune cell infiltration and subsequent hepatic inflammation, promoting NASH and fibrosis progression [68]. This discrepancy can be partially attributed to the fact that complete deficiency of  $\beta 7$  integrin impedes the migration of regulatory T cells to the liver, thereby weakening the suppression of effector T cell proliferation and promoting the secretion of inflammatory cytokines and chemokines. Furthermore, this enhanced inflammatory microenvironment recruits more neutrophils to the liver.

MAdCAM-1 was expressed in tissue sections from 27/28 patients with cirrhosis, mainly localized in the endothelium of the peribiliary blood vessel plexus and in interstitial

regions within lymphoid aggregates, and there was also a significant upregulation of *MAdCAM-1* mRNA levels compared to the control group, which was consistent with immunohistochemical analysis [69]. Infiltration of  $\alpha 4\beta 7+$  T cells in liver tissue sections was significantly higher in 10 NASH-related and 10 alcoholic steatohepatitis-related cirrhosis patients than in control patients. Mice with CCL4-induced cirrhosis also exhibited greater aggregation of  $\alpha 4\beta 7+$  T cells and higher expression of *Itga4*, *Itgb7*, and *MAdCAM-1* [70]. A comprehensive phenotypic and functional characterization of CD8+ T cells in primary biliary cirrhosis concluded that CD45RO<sup>high</sup> CD57+ CD8<sup>high</sup> T cells are a subpopulation of cytotoxic memory cells that play a key role in the destruction of bile duct epithelial cells in PBC and that these cells have a higher  $\alpha 4\beta 7$  expression compared to other CD8<sup>high</sup> T cells [71].

Increased expression of *MAdCAM-1* in liver tissue was shown in mice with immune hepatitis induced by Concanavalin A (ConA). In contrast, *MAdCAM-1*-deficient and  $\beta 7$ -deficient mice exhibited reduced production of pro-inflammatory mediators and showed protection against ConA-induced liver injury. Additionally, lymphocytes isolated from lymph nodes of wild-type and  $\beta 7$ -deficient mice were transferred intravenously into recipient mice lacking mature lymphocytes. The results showed that recipient mice receiving lymphocytes from wild-type mice had more apoptotic hepatocytes and higher expression of *MAdCAM-1*, indicating that  $\beta 7$  is involved in ConA-induced liver injury and plays a key role in *MAdCAM-1* expression [72]. *MAdCAM-1* has been reported as one of the nine groups of susceptibility loci shared by inflammatory bowel disease and autoimmune liver disease [73].

As early as 1999, Hillan KJ et al. found that 20 out of 30 patients recruited with HCV showed *MAdCAM-1* positivity in liver tissue, with a significant correlation between its expression and the hepatic activity index (HAI) classification [64].

4.2.3. Acute Liver Injury

Acute liver failure (ALF) mice exhibited elevated serum *MAdCAM-1* levels that were significantly positively correlated with ALT, an indicator of liver injury. Similarly, serum *MAdCAM-1* levels appeared to be significantly elevated in patients with liver failure and were suggested to be a promising early predictor of liver failure. Hepatic tissue *MAdCAM-1* mRNA levels also appeared to be significantly increased, and there was a significant positive correlation with hepatic tissue pro-inflammatory markers, such as TNF $\alpha$  and IL6. Accordingly, acute liver failure mice showed accumulation of  $\alpha 4\beta 7+$  CD4+ T cells in the liver [74].

Taken together, this suggests that the role of *MAdCAM-1*/ $\alpha 4\beta 7$  in hepatic inflammation is broad and not limited to primary sclerosing cholangitis (Table 1).

Table 1. Role of *MAdCAM-1*/ $\alpha 4\beta 7$  in inflammatory liver disease.

| Diseases | Cohort/Model                                             | Main Effects                                                                                                                                                                                                                     | Ref. |
|----------|----------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| PSC      | 5 patients with PSC                                      | Antibodies against <i>MAdCAM-1</i> (10G3), $\alpha 4\beta 7$ (ACT-1) blocked the adhesion of peripheral blood lymphocytes to <i>MAdCAM-1</i> -transfected CHO cells, to the hepatic portal vein, and the hepatic blood sinusoids | [63] |
|          | 16 patients with PSC who underwent liver transplantation | 14/16 cases with positive immunohistochemical <i>MAdCAM-1</i> staining                                                                                                                                                           | [63] |
|          | Liver biopsy of 9 patients with PSC                      | $\alpha 4\beta 7+$ T lymphocyte infiltration was significantly higher than in patients with non-inflammatory liver disease                                                                                                       | [62] |

Table 1. Cont.

| Diseases  | Cohort/Model                                                                                                 | Main Effects                                                                                                                                                                                                                                                                              | Ref. |
|-----------|--------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| NASH      | 5 patients with PSC who underwent liver transplantation                                                      | 5/5 cases with positive immunohistochemical MAdCAM-1 staining                                                                                                                                                                                                                             | [64] |
|           | Liver biopsy of 7 patients with Long-term PSC                                                                | The H-score of immunohistochemical MAdCAM-1 staining was significantly higher than that of the control. The proportion of $\beta$ 7-positive T cells in the liver was increased and positively correlated with PSC disease duration                                                       | [65] |
|           | Western diet-fed F11r <sup>-/-</sup> mice were administered mAbs targeting $\alpha$ 4 $\beta$ 7 and MAdCAM-1 | Amelioration of hepatic inflammation and fibrosis observed in histopathology. Significant reduction in serum ALT and AST levels, inflammatory factors such as <i>TNF-<math>\alpha</math></i> , and liver fibrosis indicators such as <i>Acta2</i> .                                       | [67] |
|           | Western diet-fed F11r <sup>-/-</sup> mice                                                                    | $\alpha$ 4 $\beta$ 7+ CD4+ T cells exhibited increased accumulation in peripheral blood, Peyer's patches, colonic lamina propria, and liver tissues, accompanied by elevated MAdCAM-1 expression in both colon and liver compared to WT mice                                              | [67] |
|           | Liver biopsy of 5 patients with NASH                                                                         | Immunohistochemical analysis revealed a larger MAdCAM-1-positive area in the livers of NASH patients compared to controls, with significant upregulation of <i>MAdCAM-1</i> , <i>ITGA4</i> , and <i>ITGB7</i> mRNA levels                                                                 | [67] |
| Cirrhosis | MAdCAM-1-deficient mice were fed high-fat diet or methionine-choline deficient diet                          | MAdCAM-1 deficiency ameliorates liver injury and fibrosis by enhancing antioxidant and anti-inflammatory immune response                                                                                                                                                                  | [68] |
|           | CCL4-induced mice were administered mAbs targeting $\alpha$ 4 $\beta$ 7 and MAdCAM-1                         | Protected against liver injury and inhibited the progression of fibrosis, reduced serum ALT, AST levels, the positive staining of Sirius red, and the expression of liver fibrosis indicator <i>Col1a2</i> , and so on                                                                    | [70] |
|           | CCL4-induced mice                                                                                            | $\alpha$ 4 $\beta$ 7+ CD4+ T cells were markedly accumulated in the liver tissues, accompanied by enhanced immunopositivity of MAdCAM-1 and $\alpha$ 4 $\beta$ 7 compared to WT mice. The mRNA levels of <i>MAdCAM-1</i> , <i>ITGA4</i> , and <i>ITGB7</i> were significantly upregulated | [70] |
|           | 10 patients with NASH-related and 10 patients with ASH-related cirrhosis who underwent liver transplantation | Stronger immunopositivity for $\alpha$ 4 $\beta$ 7 compared to controls                                                                                                                                                                                                                   | [70] |
|           | 28 patients with cirrhosis who underwent liver transplantation                                               | 27/28 cases with positive immunohistochemical MAdCAM-1 staining, with significant upregulation of <i>MAdCAM-1</i> mRNA level                                                                                                                                                              | [69] |
|           | 24 patients with PBC                                                                                         | The frequency of CD45RO <sup>high</sup> CD57+ CD8 <sup>high</sup> subpopulations expressing $\alpha$ 4 $\beta$ 7 in the peripheral blood of PBC patients was significantly higher than that of controls                                                                                   | [71] |
|           | Liver biopsy of 21 patients with PBC and 5 patients with PBC who underwent liver transplantation             | 15/21 cases with positive immunohistochemical MAdCAM-1 staining, 5/5 cases with positive immunohistochemical MAdCAM-1 staining                                                                                                                                                            | [64] |

Table 1. Cont.

| Diseases                        | Cohort/Model                                                              | Main Effects                                                                                                                                                                                                                            | Ref. |
|---------------------------------|---------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
|                                 | 11 patients with PBC who underwent liver transplantation                  | 7/11 cases with positive immunohistochemical MAdCAM-1 staining                                                                                                                                                                          | [63] |
| Acute Immune-Mediated Hepatitis | ConA-induced mice                                                         | The area of positive immunofluorescence staining and mRNA level of MAdCAM-1 in liver tissue were significantly higher than those in WT mice                                                                                             | [72] |
|                                 | ConA-induced MAdCAM-1 <sup>-/-</sup> and $\beta 7^{-/-}$ mice             | Reducing the production of pro-inflammatory mediators, such as IFN- $\gamma$ , resulted in liver tissue necrosis, hepatocyte apoptosis, and serum ALT levels, which were significantly lower than those of WT mice after receiving ConA | [72] |
| ALF                             | LPS/D-GalN-induced mice                                                   | MAdCAM-1 mRNA and protein levels in serum and liver tissue were significantly higher than those in WT mice. Higher aggregates of $\alpha 4\beta 7^{+}$ CD4 <sup>+</sup> T cells in liver tissue of ALF mice.                            | [74] |
|                                 | 12 patients with ALF                                                      | Exhibited a significantly higher level of serum MAdCAM1, more area of MAdCAM-1, $\beta 7$ positive staining in liver tissue compared with healthy controls                                                                              | [74] |
| HCV                             | Liver biopsy of 30 patients with HCV                                      | 20/30 cases with positive immunohistochemical MAdCAM-1 staining                                                                                                                                                                         | [64] |
| Autoimmune hepatitis            | 10 patients with autoimmune hepatitis who underwent liver transplantation | 7/10 cases with positive immunohistochemical MAdCAM-1 staining                                                                                                                                                                          | [63] |

#### 4.3. Lymphocyte Migration-Associated Intestinal Microbiota and Metabolites

The role of intestinal microbiota and its metabolites in the gut-liver axis is well established and it has been reported that the expression of MAdCAM-1/ $\alpha 4\beta 7$  is associated with specific intestinal microbiota and metabolites, especially with bile acids. Conrad Rauber and Guido Kroemer et al. of the Gustave Roussy Cancer Center in France found that after receiving antibiotic treatment, the *Enterocloster* spp. colonizes the intestines of mice and down-regulates MAdCAM-1 expression in PPs and MLN through the accumulation of its metabolites lithocholic acid (LCA) and ursodeoxycholic acid (UDCA), which in turn triggers an exodus of immunosuppressive  $\alpha 4\beta 7^{+}$  CD4<sup>+</sup> Treg17 cells from the intestine to tumor-draining lymph nodes, aiding in tumor immune escape and thus diminishing the efficacy of the immune checkpoint inhibitor (ICB). And suggests that serum sMAdCAM-1 is a surrogate marker for intestinal dysbiosis. It is noteworthy that exposure to either *Enterocloster* spp., UDCA, or LCA significantly reduced the fluorescence intensity of GFP under the control of the *Madcam1* promoter in sinusoidal endothelial cells (TSEC) [45]. *Lactobacillus plantarum* treatment ameliorated histological damage and decreased the expression of adhesion molecules MAdCAM-1, ICAM-1, and  $\alpha 4\beta 7$  in IL10<sup>-/-</sup> colitis mice [75]. Recipient mice receiving a high-fat diet or low-fiber dietary microbiota exhibited increased expression of MAdCAM-1 in aortic sinus plaques as well as an increase in  $\beta 7$ -positive cells compared to controls [76]. Bing Han, Xiaodan Lv et al. in addition to demonstrating the ameliorative effect of Vedolizumab in the mouse model of 2,4,6-trinitrobenzenesulfonic acid-induced colitis, found that baseline intestinal microbiota differed at all levels between the Vedolizumab-responsive group and the non-responsive group. And it was further found by antibiotic depletion of intestinal microbiota experiments that the depletion of

microbiota resulted in partial attenuation of the beneficial effects of anti- $\alpha 4\beta 7$  integrins, including weight gain, reduction in DAI scores, and reversal of severe pathologic symptoms. And recipient mice receiving fecal bacteria from Vedolizumab-responsive patients were found to exhibit elevated body weight, improved pathology, and reduced intestinal inflammation through fecal transplantation experiments. In addition, the metabolite clustering of mice in the Vedolizumab-responsive and non-responsive groups was significantly differentiated, with higher levels of bile acids such as chenodeoxycholic acid (CDCA), LCA, and Taurochenodeoxycholic acid (TCDCA) in the responding group. Moreover, CDCA, a recognized FXR agonist, showed significantly higher expression of both FXR and FXR target gene FGF15 in the responding group than in the non-responding group [77]. The association of UDCA with MAdCAM-1 was also validated in in vivo and in vitro experiments. UDCA reduced TNF- $\alpha$ -induced MAdCAM-1 expression in TSEC and brain endothelial cells (bEnd.3) in a dose-dependent manner. Similarly, a significant reduction in MAdCAM-1 expression was seen with the administration of UCDA in MDR2<sup>-/-</sup>/DSS, an animal model mimicking human PSC-IBD [78]. In addition to bile acids, cysteamine is the metabolite with the highest ability to induce functional MAdCAM-1 expression on the hepatic endothelium among VAP-1 substrates [34]. Furthermore, Treg cells were increased in the colon, MLN, and PPs in mice treated with butyric acid, while Treg expressing  $\alpha 4\beta 7$ , CCR9, and GPR15 were increased in pancreatic lymph nodes (PLN) and pancreas suggesting that intestinal-activated Treg cells migrate to the pancreas and have the function of restoring immune tolerance and delaying diabetic onset during Type 1 Diabetes [79].

Other metabolites also have a role in lymphocyte migration. Short-chain fatty acids (SCFA) increase the expression and mRNA levels of *L-selectin* on neutrophil surfaces. Both propionic acid and butyric acid among SCFA promote neutrophil migration in vitro and in vivo [80]. Compared to the control group, the medium-chain fatty acid octanoic acid increased the expression of L-selectin on T lymphocytes, decreased the proportion of free-flowing T cells, and increased the proportion of rolling lymphocytes compared to the control group. However, it had no significant effect on cell adhesion or migration rate [81]. Administration of propionic acid significantly reduced the migration of Th1 cells from the colon to the spleen in experimental autoimmune uveitis (EAU) to stabilize the subclinical intestinal changes that occur in EAU. Propionic acid may induce Th17 retention in the ileum prior to the onset of uveitis, increasing IL-17 production in the ileum and decreasing the number of Th17 cells in the MLN [82]. In addition, the gut microbiota may also regulate ILC migration and localization through local production of metabolites. The GPR109A receptor on ILC3 senses higher levels of butyric acid in ileal PPs compared to jejunal PPs, promoting ILC3 localization in jejunal PPs [83]. In addition to butyric acid, ILC3 migration is also affected by 7 $\alpha$ ,25-dihydroxycholesterol and tryptophan, and deletion of 7 $\alpha$ ,25-OHC prevents ILC3 from localizing to crypt plaques (CPs) in the colon and small bowel, and 7 $\alpha$ ,25-OHC also mediates the localization of ILC3 to the lymphoid follicles (ILF) and interfollicular regions within the MLN. Increased production of 7 $\alpha$ ,25-OHC in inflammatory states promotes GPR183+ ILC3 migration to colonic inflammation sites [84,85]. Tryptophan may enhance ILC3 migration through Ahr regulation of CCR6 [86].

5-Hydroxyindoleacetic acid (5HIAA) and its receptor GPR35 promote neutrophil aggregation to inflamed peritoneum, lymph nodes, skin and other sites. And 5HIAA is required for the induction of neutrophil aggregation [87]. Sodium lactate inhibits CD4+ T cell motility by interfering with glycolysis required for T cell migration, whereas lactate inhibits CD8+ T cells. This effect on different T cell subsets is mediated by different transporter proteins [88]. Migration of sarcosine-treated dendritic cells to lymph nodes and spleen was significantly increased after intradermal injection in the inguinal region of mice. Sarcosine also significantly increased the migration of human and mouse dendritic

cells and the expression of CXCR2, CXCL3, and CXCL1 in in vitro experiments. Sarcosine-induced migration was blocked by CXCR2-neutralizing antibodies [89]. Prostaglandin E2 (PGE2) induces dendritic cells to express the lymphoid homing chemokine receptor CCR7 and promotes their migration. Neutrophils treated with pneumococcal haemolysin or *Pseudomonas aeruginosa* enhance PGE2 production, impeding neutrophil activation and migration, thereby inhibiting an effective immune response [90]. It is concluded in Table 2.

#### 4.4. Others

In addition, other integrins [91] also play important roles in inducing lymphocyte migration in response to inflammatory signals and in the pathogenesis of inflammatory enterohepatic diseases [92]. Examples include  $\alpha$ L $\beta$ 2 (LFA-1) and its ligand ICAM-1. The therapeutic agent efalizumab, which blocks this interaction, improves dermatopathological features of psoriasis by decreasing recruitment of inflammatory cells at psoriatic lesions [93,94]. Lifitegrast, a novel small-molecule antagonist that blocks ICAM-1 binding to  $\alpha$ L $\beta$ 2, reduces corneal inflammation and is approved in the United States for the treatment of dry eye [95]. In terms of intestinal inflammation, Pavlick et al. also found that RAG-1<sup>−/−</sup> recipient mice receiving LFA-1<sup>−/−</sup> donor T cells showed less evidence of colitis than recipient mice receiving wild-type mouse T cells [96]. The use of an antibody that blocks the integrin MAC-1, which is formed from the  $\beta$ 2 subunit and  $\alpha$ M, also inhibits inflammatory cell recruitment, reduces ulcer area and number of ulcers, and ameliorates tissue damage in rats with colitis [97]. In addition to leukocyte adhesion integrins, the  $\alpha$ v subunit-containing  $\alpha$ v $\beta$ 3 and  $\alpha$ v $\beta$ 5 of the RGD-binding integrins bind to bone bridging proteins in the livers of NAFLD mice, which inhibits autophagosome-lysosome fusion and promotes lipid accumulation. Free fatty acids-induced autophagic damage was attenuated and lipid accumulation was reduced in HepG2 cells using  $\alpha$ v $\beta$ 3 and  $\alpha$ v $\beta$ 5 antibodies [98]. Deletion of the gene encoding the  $\alpha$ v integrin subunit effectively targets myofibroblasts in multiple organs and is protective in multiple fibrosis models. Blockade of  $\alpha$ v-containing integrins by the small molecule CWHM 12 attenuated liver and lung fibrosis [99].  $\alpha$ v $\beta$ 3 expressed on stellate cells also has a clear role in fibrosis in NASH, and the use of an antibody that blocks integrin  $\alpha$ v $\beta$ 3 reduces the expression of laminin in high glucose-induced human liver sinusoidal endothelial cells [100]. Expression of  $\alpha$ v $\beta$ 3 detected using the 18F-FPP-RGD2 PET probe in NASH mouse model is strongly associated with histological fibrosis [101,102]. In addition, the use of anti-Itgb1 antibody attenuated pro-inflammatory monocyte infiltration, suppressed hepatic inflammation, and ameliorated hepatic injury and fibrosis in the livers of mice fed a high fat, fructose, and cholesterol diet [103].

**Table 2.** Role of intestinal microbiota or metabolites on lymphocyte migration.

| Intestinal Microbiota/Metabolites | Cells          | Mechanisms                                                                                                                                                       | Refs. |
|-----------------------------------|----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| propanoic acid                    | Th1/Th17 cells | Propionate reduces migration of Th1 cells from the colon to the spleen in experimental EAU and induces Th17 retention in the ileum prior to the onset of uveitis | [82]  |
| SCFA                              | neutrophil     | SCFA promote L-selectin expression on neutrophils to stimulate their migration                                                                                   | [80]  |
| butyric acid                      | Treg cell      | Butyric acid promotes migration of gut-activated Treg cells (expressing $\alpha$ 4 $\beta$ 7, CCR9 and GPR15) to the pancreas and PLN                            | [79]  |

Table 2. Cont.

| Intestinal Microbiota/Metabolites                       | Cells                     | Mechanisms                                                                                                                                                                                                                                                       | Refs.   |
|---------------------------------------------------------|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| butyric acid                                            | ILC3                      | Butyric acid regulates the localization of NKp46+ ILC3 in Peyer's patch                                                                                                                                                                                          | [83]    |
| 7 $\alpha$ ,25-Dihydroxycholesterol(7 $\alpha$ ,25-OHC) | ILC3                      | 7 $\alpha$ ,25-OHC and its receptor GPR183 mediate localization of CCR6+ LTi-like ILC3 to crypts, isolated lymphoid follicles                                                                                                                                    | [84,85] |
| tryptophan                                              | ILC3                      | Tryptophan may be able to enhance ILC3 migration through Ahr regulation of CCR6                                                                                                                                                                                  | [86]    |
| 5-Hydroxyindoleacetic acid (5HIAA)                      | neutrophil                | 5HIAA and its receptor GPR35 promote neutrophil aggregation to inflamed peritoneum, lymph nodes, skin and other sites                                                                                                                                            | [87]    |
| lactic acid/sodium lactate                              | T cell                    | Sodium lactate inhibits CD4+ T cell motility and lactate inhibits CD8+ T cell motility                                                                                                                                                                           | [88]    |
| sarcosine                                               | dendritic cell            | Sarcosine increases migration of mouse and human dendritic cells through the CXC chemokine pathway                                                                                                                                                               | [89]    |
| ProstaglandinE2 (PGE2)                                  | dendritic cell/neutrophil | PGE2 induces dendritic cells expressing CCR7 to migrate to lymph nodes, inhibiting neutrophil activation and migration                                                                                                                                           | [90]    |
| caprylic acid                                           | T cell                    | caprylic acid increases L-selectin expression to stimulate lymphocyte rolling but does not affect lymphocyte adhesion                                                                                                                                            | [81]    |
| <i>Enterocloster</i> LCA, UDCA                          | Treg17 cell               | <i>Enterocloster</i> down-regulates MAdCAM-1 expression in PPs and MLN through accumulation of its metabolites LCA and UDCA, which in turn causes $\alpha$ 4 $\beta$ 7+ CD4+ Treg17 cells to leave the intestine and metastasize into tumor-draining lymph nodes | [45]    |
| <i>Lactobacillus plantarum</i>                          | /                         | <i>Lactobacillus plantarum</i> reduces MAdCAM-1, ICAM-1 and $\alpha$ 4 $\beta$ 7 expression and ameliorates histological damage in IL10 <sup>-/-</sup> colitis mice                                                                                              | [75]    |
| UDCA                                                    | /                         | UDCA inhibits MAdCAM-1 expression in TNF- $\alpha$ -induced TSEC, bEnd.3 cells and MDR2 <sup>-/-</sup> /DSS animal model                                                                                                                                         | [78]    |
| cysteamine                                              | /                         | Deamidation of cysteamine by VAP-1 induces up-regulation of MAdCAM-1 on hepatic endothelium                                                                                                                                                                      | [34]    |

## 5. $\alpha$ 4 $\beta$ 7/MAdCAM-1 as a Drug Target in Inflammatory Enterohepatic Diseases

A number of therapeutic agents targeting the MAdCAM-1/ $\alpha$ 4 $\beta$ 7 interaction (Figure 4) have been approved or are under investigation in various clinical trial phases.




**Figure 4.**  $\alpha 4\beta 7$ /MAdCAM-1 as a Drug Target. Blocking the interaction between MAdCAM-1 and  $\alpha 4\beta 7$  with monoclonal antibodies or small-molecule drugs inhibits lymphocyte migration, thereby ameliorating inflammation.

#### 5.1. Monoclonal Antibody

On May 20, 2014, vedolizumab [Entyvio (US, Europe)], a humanized monoclonal antibody against  $\alpha 4\beta 7$  integrin, received its first global approval in the United States. Vedolizumab is used for the treatment of adult patients with moderately to severely active ulcerative colitis and Crohn's disease who have had an inadequate response, loss of response, or intolerance to TNF- $\alpha$  inhibitors or immunomodulators, or who have had an inadequate response or intolerance to, or have developed dependence on corticosteroids [56]. In ulcerative colitis, Vedolizumab was used to induce and maintain clinical response and clinical remission. No significant difference in adverse event incidence was observed between the Vedolizumab and placebo groups [104]. Although adverse events were more common with Vedolizumab for the treatment of Crohn's disease, patients with active Crohn's disease treated with Vedolizumab were more likely to achieve remission at week 6 compared to those treated with placebo. Patients who responded to induction therapy were more likely to achieve remission at week 52 if they continued to receive Vedolizumab, compared to those who switched to placebo [105]. In addition to their role in IBD, Kate D. Lynch, Roger W. Chapman et al. found no definitive biochemical response to Vedolizumab in a cohort with primary sclerosing cholangitis and inflammatory bowel disease who had received at least three doses. However, a 20% reduction in alkaline phosphatase (ALP) was observed in some patients, and those with cirrhosis and higher baseline ALP levels were more likely to show a reduction in ALP [106]. This result may be due to the indirect therapeutic effect of vedolizumab, as it has been shown to produce only gut-specific anti-inflammatory activity [57].

Natalizumab, a recombinant humanized monoclonal antibody against the  $\alpha 4$  integrin, was the first drug approved for the treatment of Crohn's disease, increasing clinical remission and response rates and improving quality of life [107]. However, Natalizumab has been limited due to the fact that 1 in 1000 treated individuals may develop progressive multifocal leukoencephalopathy [108]. Abrilumab, a fully human monoclonal IgG2 antibody against  $\alpha 4\beta 7$ , demonstrated significantly better remission rates after treatment in both

Phase II studies against moderate to severe CD and UC than in the placebo group [109]. However, no Phase III clinical trial enrollment information is available to date. A phase III, multicenter, double-blind, placebo-controlled study of Etrolizumab, a monoclonal antibody specifically targeting the  $\beta 7$  subunit, in patients with moderately-to-severely active ulcerative colitis who had been previously treated with anti-TNF drugs, showed that the Etrolizumab group was able to achieve a higher rate of remission compared to the control group. However, no significant difference was observed between the groups on the primary maintenance endpoint of remission at week 66 [110]. In HIBISCUS I and HIBISCUS II, two identically designed, multicenter, phase III, randomized, double-blind, placebo-controlled and active-controlled studies, Etrastuzumab given to patients with moderately to severely active ulcerative colitis who had not been treated with an anti-TNF agent, Etrastuzumab was superior to placebo in inducing remission in HIBISCUS I, but its performance was not superior to Adalimumab in both HIBISCUS I and HIBISCUS II trials [111], and it also performed similarly to Infliximab (tumor necrosis factor inhibitor) [112]. Ontamalimab, a fully human immunoglobulin G2 monoclonal antibody against MAdCAM-1, was shown to achieve the primary endpoint of clinical remission in more patients than the placebo group. It also showed a significant advantage in endoscopic improvement with no safety concerns [113].

### 5.2. Small Molecule Drug

MORF-057, an orally administered small molecule inhibitor designed to selectively inhibit integrin  $\alpha 4\beta 7$ , is currently undergoing its evaluation in adult patients with moderately to severely active ulcerative colitis and Crohn's disease [114,115]. AJM300, an oral  $\alpha 4$  integrin antagonist, induced clinical responses in patients with moderately active ulcerative colitis and in patients with inadequate response or intolerance to Mesalazine, and there was no difference in the incidence of adverse events compared to controls [116]. It was first approved for the treatment of patients with moderate ulcerative colitis who were not responding well to treatment with 5-aminosalicylic acid in Japan in 2022 [117]. Although the oral  $\alpha 4\beta 7$  antagonist peptide PTG-100 initially failed to meet the primary endpoint in the phase II study, blinded re-readings of endoscopies from 65 participants evaluated for futility showed that the trial would not reach futility, and colon biopsy results suggested that PTG-100 could improve histology and that the trial could continue as planned [118]. The key clinical trial outcomes of these drugs are summarized in Table 3, while their current developmental status is presented in Table 4.

**Table 3.** Clinical trial results of MAdCAM-1/ $\alpha 4\beta 7$ -targeted therapies.

| Drug Name   | Disease | Efficacy                                                                                                                        | Safety                   | Limitations                                                                                                                       |
|-------------|---------|---------------------------------------------------------------------------------------------------------------------------------|--------------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| Natalizumab | CD      | Primary endpoint (clinical remission at Week 6) was not met; however, positive signals were observed at multiple time points    | Generally well-tolerated | The short-term (12-week) study design precluded assessment of long-term risks, such as progressive multifocal leukoencephalopathy |
| Vedolizumab | UC      | Clinical response significantly higher at Wk6 (47.1% vs. 25.5%); remission at Wk52 significantly higher (41.8%/44.8% vs. 15.9%) | Generally well-tolerated | Optimal induction duration and minimum effective dose not determined                                                              |

Table 3. Cont.

| Drug Name   | Disease | Efficacy                                                                                                                                                             | Safety                                                   | Limitations                                                                                                                        |
|-------------|---------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|
|             | CD      | Significantly higher clinical remission vs. placebo at Wk6 (14.5% vs. 6.8%) and Wk52 (39.0%/36.4% vs. 21.6%) in maintenance responders                               | Higher incidence of serious AEs (24.4% vs. 15.3%)        | Optimal patient population not defined; combination therapy effects not assessed                                                   |
|             | PSC     | Biochemical response in a subset of patients (20.6%); effective for concomitant IBD                                                                                  | Significant liver biochemical worsening in some patients | Retrospective design; lack of control group; UDCA use post-baseline unclear                                                        |
| Etrolizumab | UC      | Significant induction of remission at Wk 14 (18.5% vs. 6.3%)                                                                                                         | Generally well-tolerated                                 | Lack of statistically significant efficacy in the maintenance phase                                                                |
|             | CD      | Significant improvement in clinical remission (35% vs. 24%) and endoscopic improvement (24% vs. 12%) at Week 66 vs. placebo. No significant benefit during induction | Generally well-tolerated                                 | mITT analysis; potentially underestimated placebo response; unusually long study duration; premature closure of induction cohort 3 |
| Abrilumab   | UC      | Unadjusted remission rates at Week 8 were significantly higher with abrilumab (13.3%, 12.7%) than placebo (4.3%)                                                     | Generally well-tolerated                                 | Dosing errors; short-term study; limited subgroup analysis by prior TNF antagonist exposure                                        |
| Ontamalimab | UC      | Significantly higher clinical remission vs. placebo at Wk 12 (29.8%/29.5% vs. 15.8%/12.5%) and Wk 52 (53.5%/40.2% vs. 8.2%/12.8%)                                    | Generally well-tolerated                                 | Clinical development program prematurely discontinued                                                                              |
| AJM300      | UC      | Clinical response rate significantly higher with AJM300 vs. placebo at Week 8 (45% vs. 21%)                                                                          | Generally well-tolerated                                 | Limited sample size/retreatment cycles; severe/refractory UC populations not evaluated; open-label retreatment phase               |
| PTG-100     | UC      | Primary endpoint (remission at Wk12) not met initially                                                                                                               | Generally well-tolerated                                 | limited sample size; initial primary endpoint failure; results interpreted with caution.                                           |
| MORF-057    | UC      | At Week 12, mean reduction from baseline in RHI score was −6.4; 22.9% of participants achieved RHI remission (score ≤ 3)                                             | Generally well-tolerated                                 | Small sample size; high dropout rate; lack of placebo control                                                                      |

**Table 4.** Current development status of MAdCAM-1/ $\alpha$ 4 $\beta$ 7-targeted therapies.

| Drug Name   | Mechanism of Action                                          | Disease | Approval/Phase  | Primary Developer/Company         | Source                                                                                                                                                                                                                                                    |
|-------------|--------------------------------------------------------------|---------|-----------------|-----------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Natalizumab | Humanized IgG4 anti- $\alpha$ 4 integrin subunit mAb         | CD      | Licensed in USA | Biogen Idec                       | <a href="https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&amp;AppNo=125104">https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&amp;AppNo=125104</a> [119]<br>(accessed on 19 February 2025) |
| Vedolizumab | Humanized IgG1 anti- $\alpha$ 4 $\beta$ 7 integrin dimer mAb | CD/UC   | Licensed        | Takeda                            | <a href="https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&amp;AppNo=125476">https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&amp;AppNo=125476</a> [120]<br>(accessed on 19 February 2025) |
| Etrolizumab | Humanized IgG1 anti- $\beta$ 7 integrin subunit mAb          | UC/CD   | Phase III       | Roche                             | UC: NCT02165215<br>NCT02100696<br>CD: NCT02394028                                                                                                                                                                                                         |
| Abrilumab   | Human IgG2 anti- $\alpha$ 4 $\beta$ 7 integrin dimer mAb     | CD/UC   | Phase II        | Amgen                             | CD: NCT01696396<br>UC: NCT01694485                                                                                                                                                                                                                        |
| Ontamalimab | Human IgG2 anti-MAdCAM-1 mAb                                 | UC      | Phase III       | Takeda                            | NCT03290781                                                                                                                                                                                                                                               |
| AJM300      | $\alpha$ 4 integrin subunit antagonist                       | UC      | Phase III       | Meiji Seika Pharma/<br>EA Pharma  | NCT03531892                                                                                                                                                                                                                                               |
| PTG-100     | $\alpha$ 4 $\beta$ 7 integrin-specific peptide               | UC      | Phase IIb       | Protagonist Therapeutics          | NCT02895100                                                                                                                                                                                                                                               |
| MORF-057    | small molecule $\alpha$ 4 $\beta$ 7 integrin inhibitor       | UC/CD   | Phase II        | Morphic Therapeutic/<br>Eli Lilly | UC: NCT05291689<br>NCT05611671<br>CD: NCT06226883                                                                                                                                                                                                         |

## 6. Conclusions and Perspective

The recruitment and migration of lymphocytes are central to the immune response, enabling their precise homing to sites of inflammation. This process plays a critical role in maintaining immune homeostasis, initiating specific immune reactions, and regulating inflammatory progression. Furthermore, it is also highly regulated and requires coordination by adhesion molecules expressed on homing lymphocytes and corresponding ligands expressed by endothelial cells, in which MAdCAM-1/ $\alpha$ 4 $\beta$ 7 has an important role in the regulatory process [121]. Excessive lymphocyte migration, triggered by dysregulated MAdCAM-1/ $\alpha$ 4 $\beta$ 7 expression, not only disrupts intestinal immune balance and drives local inflammation but also mediates the spread of inflammation to the liver through the gut-liver axis. This sequential process constitutes the core pathological feature of inflammatory enterohepatic diseases, highlighting the integral link between gut and liver in inflammatory mechanisms. Therefore, inhibiting aberrant MAdCAM-1/ $\alpha$ 4 $\beta$ 7 binding is essential for curbing this connected series of intestinal and hepatic inflammatory events [122].

Blocking lymphocyte migration to the liver may be a potential mechanism to reduce inflammation and prevent disease progression. Administration of both MAdCAM-1 mAb and  $\alpha 4\beta 7$  mAb to Western diet-induced NASH mice ameliorated their inflammation and liver fibrosis [67]. Similarly, the use of MAdCAM-1 mAb and  $\alpha 4\beta 7$  mAb resulted in reduced hepatic inflammation and blocked the progression of fibrosis in mice with CCL4-induced hepatic fibrosis [70]. The lack of MAdCAM-1 or  $\beta 7$  integrins also reduced pro-inflammatory mediator production and attenuated ConA-induced liver injury [72]. However, Drescher et al. also found that  $\beta 7$  integrin deficiency triggered more intense hepatic immune cell infiltration and subsequent hepatic inflammation. This discrepancy can be partially explained by key methodological differences between the present study and the work by Rai et al. Specifically, Drescher et al. utilized  $\beta 7$  integrin-deficient mice, whereas Rai et al. employed a neutralizing monoclonal antibody against  $\alpha 4\beta 7$  to block its interaction with MAdCAM-1. The complete absence of  $\beta 7$  integrin impaired the migration of regulatory T cells to the liver, thereby attenuating the suppression of effector T cell proliferation and promoting the secretion of inflammatory cytokines and chemokines. This enhanced inflammatory milieu recruited more neutrophils to the liver. Moreover,  $\beta 7^{-/-}$  mice exhibited elevated levels of IL-17 mRNA, a cytokine produced by Th17 cells that is known to promote neutrophil recruitment. More directly, neutrophils do not express integrin  $\alpha 4\beta 7$ , indicating that their migration to inflammatory sites is independent of  $\beta 7$  [68]. And this controversy may require further research. Moreover, in clinical trials, Vedolizumab did not significantly improve the levels of biochemical markers such as hepatic transaminases and bilirubin in patients with PSC, but the 20% reduction in ALP was observed in some patients, and patients with cirrhosis and higher ALP at baseline were more likely to respond with a reduction in ALP [106,123]. This may be related to the fact that vedolizumab, by blocking  $\alpha 4\beta 7$  integrin, exclusively produces gut-specific anti-inflammatory activity. In healthy non-human primates treated with vedolizumab, a significant decrease in the frequency of  $\beta 7^{+}$  lymphocytes in gastrointestinal tissues and a significant increase in the frequency of  $\alpha 4\beta 7^{+}$  memory helper T lymphocytes in peripheral blood were observed. Additionally, Vedolizumab did not suppress systemic adaptive or innate immune responses. These data suggest that Vedolizumab has a gut-specific anti-inflammatory effect [124]. Moreover, a vaccination study in healthy volunteers showed that vedolizumab attenuated the response to enteral antigen challenge (oral cholera vaccine) without affecting the response to parenteral antigen challenge (intramuscular hepatitis B vaccine), indicating a high degree of gut-specific anti-inflammatory activity for Vedolizumab [57]. In a limited cohort, all three patients with comorbid CD and cirrhosis, who had undergone ileal or colonic resection, or who were unresponsive to treatments such as infliximab, were well tolerated by vedolizumab in the present study, and none developed significant infectious complications or decompensated cirrhosis [125].

The current gut selectivity of drugs is likely attributable to the high constitutive expression of MAdCAM-1 in the gut, compared to its minimal expression in the liver under physiological conditions. However, hepatic MAdCAM-1 is upregulated during immune activation. Therefore, future therapeutic strategies for liver inflammatory diseases may focus on blocking the migration of  $\alpha 4\beta 7$ -expressing cells to the liver. One such strategy could involve the construction of GalNAc-modified, ROS-responsive polymeric nanoparticles to encapsulate  $\alpha 4\beta 7$ /MAdCAM-1 antagonists [126,127]. Such a targeted delivery system could enhance liver targeting while reducing systemic immunosuppression. Combination strategies with agents that address other hepatic pathological features, such as disordered lipid metabolism, should also be considered. Furthermore, given that combination therapies using microecological modulators and small-molecule drugs have been previously documented [128], the potential synergistic effect between anti-MAdCAM-1/ $\alpha 4\beta 7$  therapy

and FXR agonists—such as obeticholic acid—is highly promising. Such a combination strategy could simultaneously inhibit the infiltration of aberrant immune cells into inflamed tissues and restore microbiome-metabolite homeostasis, thereby addressing both the inflammatory and metabolic pathological features of the disease. More importantly, to rigorously test this hypothesis in humans, well-designed Phase II/I trials in PSC and NASH should be implemented. These trials ought to incorporate appropriate and stable biomarkers for efficacy evaluation and adopt stratified randomization designs to enable further assessment in specific patient subpopulations. Beyond therapeutic applications, specific molecular imaging probes such as  $^{18}\text{F}$ -FPP-RGD2 enable non-invasive, in vivo, quantitative detection of specific molecular pathway activity underlying diseases. These probes can be used not only for diagnosis and staging but also for real-time treatment guidance and monitoring. Additionally, as highlighted in the study by Marine Fidelle et al., serum MAdCAM-1 may serve as a strong prognostic indicator of treatment response in cancer patients [45], raising the possibility of using specific MAdCAM-1 detection for diagnosing or monitoring treatment of inflammatory enterohepatic diseases. All these aspects warrant further research and exploration.

In addition to the diseases mentioned above, gut-associated lymphoid tissues have been shown to be a key site of human immunodeficiency virus (HIV) viral replication, especially during the acute phase of HIV infection. The doping of host cell proteins by HIV viroosomes has long been recognized, and  $\alpha 4\beta 7$  remains functionally active after its doping, with the ability to bind to MAdCAM-1, and may contribute to early infection of intestinal tissues through binding to MAdCAM-1 [129]. Thus, antibodies that block viral admixture of  $\alpha 4\beta 7$  may ameliorate intestinal damage and slow disease progression during early HIV infection [130]. In addition, drugs targeting  $\alpha 4$  are being experimented and explored in diseases such as multiple sclerosis, asthma and allergic conjunctivitis [131–135]. The role of MAdCAM-1/ $\alpha 4\beta 7$  in inflammatory diseases is widely recognized.

Currently, several emerging drugs targeting MAdCAM-1/ $\alpha 4\beta 7$  to block lymphocyte homing primarily focus on IBD; however, their potential role in liver inflammation remains largely unexplored and warrants investigation. This review summarizes the current state of development, key research findings, and existing challenges, aiming to facilitate the discovery and clinical development of molecules or drugs targeting MAdCAM-1/ $\alpha 4\beta 7$  that could benefit liver inflammation.

**Author Contributions:** Conceptualization, C.Y. and Q.L.; investigation, Y.P. and A.M.; writing—original draft preparation, C.Y. and Q.L.; writing—review and editing, Y.Z. and Y.H.; visualization, C.Y., Y.P. and A.M.; supervision, Y.Z. and Y.H. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the National Natural Science Foundation of China, grant number No. 82204840, No. 82174186, and Yangfan Project of Shanghai Science and Technology Commission, grant number 22YF1449400.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** The figures for this article were generated through the Biorender website, and permission to publish them has been obtained (<https://biorender.com/>).

**Conflicts of Interest:** The authors declare no conflicts of interest.

## Abbreviations

|          |                                             |
|----------|---------------------------------------------|
| IBD      | Inflammatory bowel disease                  |
| UC       | Ulcerative colitis                          |
| CD       | Crohn's disease                             |
| PSC      | Primary sclerosing cholangitis              |
| NASH     | Non-alcoholic steatohepatitis               |
| PSI      | plexin-semaphorin-integrin                  |
| RGD      | Arginine-Glycine-Aspartic Acid              |
| IgSF     | Immunoglobulin superfamily                  |
| MAdCAM-1 | Mucosal address in adhesion molecule 1      |
| ICAM     | Intercellular adhesion molecule             |
| VCAM-1   | Vascular cell adhesion molecule             |
| PECAM-1  | Platelet/endothelial cell adhesion molecule |
| HEV      | High endothelial venule                     |
| GALT     | Gut-associated lymphoid tissues             |
| RAR      | retinoic acid receptor                      |
| RXR      | retinoid X receptor                         |
| RAREs    | retinoic acid response elements             |
| PPs      | Peyer's patche                              |
| MLN      | Mesenteric lymph nodes                      |
| DSS      | Dextran Sulfate Sodium Salt                 |
| OVA      | Ovalbumin                                   |
| VAP1     | Vascular adhesion protein-1                 |
| MA       | Methylamine                                 |
| RA       | Retinoic acid                               |
| EGID     | eosinophilic gastrointestinal diseases      |
| CX3CR1   | CX3C chemokine receptor 1                   |
| CX3CL1   | CX3C chemokine ligand 1                     |
| ILCs     | Innate lymphoid cells                       |
| FDA      | Food and Drug Administration                |
| PBC      | Primary biliary cirrhosis                   |
| CLD      | Chronic liver diseases                      |
| ASH      | Alcoholic steatohepatitis                   |
| ALF      | Acute liver failure                         |
| HCV      | Viral hepatitis C                           |
| HAI      | hepatic activity index                      |
| ConA     | Concanavalin A                              |
| LCA      | Lithocholic acid                            |
| UDCA     | Ursodeoxycholic acid                        |
| ICB      | immune checkpoint inhibitor                 |
| CDCA     | Chenodeoxycholic acid                       |
| TCDCA    | Taurochenodeoxycholic acid                  |
| PLN      | Pancreatic lymph nodes                      |
| SCFA     | Short-chain fatty acids                     |
| EAU      | Experimental autoimmune uveitis             |
| CPs      | Crypt plaques                               |
| ILF      | Lymphoid follicles                          |
| 5HIAA    | 5-Hydroxyindoleacetic acid                  |
| PGE2     | Prostaglandin E2                            |
| ALP      | Alkaline phosphatase                        |
| HIV      | human immunodeficiency virus                |

## References

1. Israelsen, M.; Francque, S.; Tsochatzis, E.A.; Krag, A. Steatotic liver disease. *Lancet* **2024**, *404*, 1761–1778. [\[CrossRef\]](#) [\[PubMed\]](#)
2. Lazaridis, K.N.; LaRusso, N.F. Primary Sclerosing Cholangitis. *N. Engl. J. Med.* **2016**, *375*, 1161–1170. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Xia, X.; Huang, Z.; Xu, C.; Fu, H.; Wang, S.; Tian, J.; Rui, K. Regulation of intestinal tissue-resident memory T cells: A potential target for inflammatory bowel disease. *Cell Commun. Signal.* **2024**, *22*, 610. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Zundler, S.; Gunther, C.; Kremer, A.E.; Zaiss, M.M.; Rothhammer, V.; Neurath, M.F. Gut immune cell trafficking: Inter-organ communication and immune-mediated inflammation. *Nat. Rev. Gastroenterol. Hepatol.* **2023**, *20*, 50–64. [\[CrossRef\]](#)
5. Kobayashi, M.; Hoshino, H.; Suzawa, K.; Sakai, Y.; Nakayama, J.; Fukuda, M. Two distinct lymphocyte homing systems involved in the pathogenesis of chronic inflammatory gastrointestinal diseases. *Semin. Immunopathol.* **2012**, *34*, 401–413. [\[CrossRef\]](#)
6. Neurath, M.F. Current and emerging therapeutic targets for IBD. *Nat. Rev. Gastroenterol. Hepatol.* **2017**, *14*, 269–278, Erratum in *Nat. Rev. Gastroenterol. Hepatol.* **2017**, *14*, 688. [\[CrossRef\]](#)
7. Barczyk, M.; Carracedo, S.; Gullberg, D. Integrins. *Cell Tissue Res.* **2010**, *339*, 269–280. [\[CrossRef\]](#)
8. Takada, Y.; Ye, X.; Simon, S. The integrins. *Genome Biol.* **2007**, *8*, 215. [\[CrossRef\]](#)
9. Luo, B.; Carman, C.V.; Springer, T.A. Structural basis of integrin regulation and signaling. *Annu. Rev. Immunol.* **2007**, *25*, 619–647. [\[CrossRef\]](#)
10. Lamb, C.A.; O’Byrne, S.; Keir, M.E.; Butcher, E.C. Gut-Selective Integrin-Targeted Therapies for Inflammatory Bowel Disease. *J. Crohn’s Colitis* **2018**, *12*, S653–S668. [\[CrossRef\]](#)
11. Soler, D.; Chapman, T.; Yang, L.; Wyant, T.; Egan, R.; Fedyk, E.R. The binding specificity and selective antagonism of vedolizumab, an anti- $\alpha 4\beta 7$  integrin therapeutic antibody in development for inflammatory bowel diseases. *J. Pharmacol. Exp. Ther.* **2009**, *330*, 864–875. [\[CrossRef\]](#)
12. de Krijger, M.; Wildenberg, M.E.; de Jonge, W.J.; Ponsioen, C.Y. Return to sender: Lymphocyte trafficking mechanisms as contributors to primary sclerosing cholangitis. *J. Hepatol.* **2019**, *71*, 603–615. [\[CrossRef\]](#)
13. Carman, C.V.; Springer, T.A. Integrin avidity regulation: Are changes in affinity and conformation underemphasized? *Curr. Opin. Cell Biol.* **2003**, *15*, 547–556. [\[CrossRef\]](#)
14. Hynes, R.O. Integrins: Bidirectional, Allosteric Signaling Machines. *Cell* **2002**, *110*, 673–687. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Tan, K.; Casanovas, J.M.; Liu, J.H.; Briskin, M.J.; Springer, T.A.; Wang, J.H. The structure of immunoglobulin superfamily domains 1 and 2 of MAdCAM-1 reveals novel features important for integrin recognition. *Structure* **1998**, *6*, 793–801. [\[CrossRef\]](#)
16. Newman, P.J. The biology of PECAM-1. *J. Clin. Investig.* **1997**, *99*, 3–8. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Mizushima, T.; Sasaki, M.; Ando, T.; Wada, T.; Tanaka, M.; Okamoto, Y.; Ebi, M.; Hirata, Y.; Murakami, K.; Mizoshita, T.; et al. Blockage of angiotensin II type 1 receptor regulates TNF- $\alpha$ -induced MAdCAM-1 expression via inhibition of NF- $\kappa$ B translocation to the nucleus and ameliorates colitis. *Am. J. Physiol. Gastrointest. Liver Physiol.* **2010**, *298*, G255–G266. [\[CrossRef\]](#) [\[PubMed\]](#)
18. Takeuchi, M.; Baichwal, V.R. Induction of the gene encoding mucosal vascular addressin cell adhesion molecule 1 by tumor necrosis factor  $\alpha$  is mediated by NF- $\kappa$ B proteins. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 3561–3565. [\[CrossRef\]](#)
19. von Andrian, U.H.; Mempel, T.R. Homing and cellular traffic in lymph nodes. *Nat. Rev. Immunol.* **2003**, *3*, 867–878. [\[CrossRef\]](#)
20. Kinashi, T. Intracellular signalling controlling integrin activation in lymphocytes. *Nat. Rev. Immunol.* **2005**, *5*, 546–559. [\[CrossRef\]](#)
21. Cybulsky, M.I.; Nourshargh, S.; Ley, K.; Laudanna, C. Getting to the site of inflammation: The leukocyte adhesion cascade updated. *Nat. Rev. Immunol.* **2007**, *7*, 678–689. [\[CrossRef\]](#)
22. Vinogradova, O.; Velyvis, A.; Velyviene, A.; Hu, B.; Haas, T.; Plow, E.; Qin, J. A structural mechanism of integrin  $\alpha$ (IIb) $\beta$ (3) “inside-out” activation as regulated by its cytoplasmic face. *Cell* **2002**, *110*, 587–597. [\[CrossRef\]](#)
23. Zundler, S.; Becker, E.; Schulze, L.L.; Neurath, M.F. Immune cell trafficking and retention in inflammatory bowel disease: Mechanistic insights and therapeutic advances. *Gut* **2019**, *68*, 1688–1700. [\[CrossRef\]](#)
24. Colucci, M.; Zumerle, S.; Bressan, S.; Gianfanti, F.; Troiani, M.; Valdata, A.; D’Ambrosio, M.; Pasquini, E.; Varesi, A.; Cogo, F.; et al. Retinoic acid receptor activation reprograms senescence response and enhances anti-tumor activity of natural killer cells. *Cancer Cell* **2024**, *42*, 646–661. [\[CrossRef\]](#)
25. Dammes, N.; Goldsmith, M.; Ramishetti, S.; Dearling, J.; Veiga, N.; Packard, A.B.; Peer, D. Conformation-sensitive targeting of lipid nanoparticles for RNA therapeutics. *Nat. Nanotechnol.* **2021**, *16*, 1030–1038. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Wang, S.; Lin, C.; Li, Y.; Liu, Z.; Wang, J.; Zhang, Y.; Yan, Z.; Zhang, Y.; Li, G.; Chen, J. Distinct chemokines selectively induce HIV-1 gp120-integrin  $\alpha 4\beta 7$  binding via triggering conformer-specific activation of  $\alpha 4\beta 7$ . *Signal Transduct. Target. Ther.* **2021**, *6*, 265. [\[CrossRef\]](#)
27. Seidel, D.; Eickmeier, I.; Kuhl, A.A.; Hamann, A.; Loddenkemper, C.; Schott, E. CD8 T cells primed in the gut-associated lymphoid tissue induce immune-mediated cholangitis in mice. *Hepatology* **2014**, *59*, 601–611. [\[CrossRef\]](#)

28. Eksteen, B.; Grant, A.J.; Miles, A.; Curbishley, S.M.; Lalor, P.F.; Hubscher, S.G.; Briskin, M.; Salmon, M.; Adams, D.H. Hepatic endothelial CCL25 mediates the recruitment of CCR9+ gut-homing lymphocytes to the liver in primary sclerosing cholangitis. *J. Exp. Med.* **2004**, *200*, 1511–1517. [[CrossRef](#)] [[PubMed](#)]
29. Adams, D.H.; Eksteen, B. Aberrant homing of mucosal T cells and extra-intestinal manifestations of inflammatory bowel disease. *Nat. Rev. Immunol.* **2006**, *6*, 244–251. [[CrossRef](#)]
30. Eksteen, B.; Mora, J.R.; Haughton, E.L.; Henderson, N.C.; Lee Turner, L.; Villablanca, E.J.; Curbishley, S.M.; Aspinall, A.I.; von Andrian, U.H.; Adams, D.H. Gut Homing Receptors on CD8 T Cells Are Retinoic Acid Dependent and Not Maintained by Liver Dendritic or Stellate Cells. *Gastroenterology* **2009**, *137*, 320–329. [[CrossRef](#)] [[PubMed](#)]
31. Eickmeier, I.; Seidel, D.; Grun, J.R.; Derkow, K.; Lehnardt, S.; Kuhl, A.A.; Kuhl, A.A.; Hamann, A.; Schott, E. Influence of CD8 T cell priming in liver and gut on the enterohepatic circulation. *J. Hepatol.* **2014**, *60*, 1143–1150. [[CrossRef](#)] [[PubMed](#)]
32. Wong, J.; Johnston, B.; Lee, S.S.; Bullard, D.C.; Smith, C.W.; Beaudet, A.L.; Kubes, P. A minimal role for selectins in the recruitment of leukocytes into the inflamed liver microvasculature. *J. Clin. Invest.* **1997**, *99*, 2782–2790. [[CrossRef](#)] [[PubMed](#)]
33. Liaskou, E.; Karikoski, M.; Reynolds, G.M.; Lalor, P.F.; Weston, C.J.; Pullen, N.; Salmi, M.; Jalkanen, S.; Adams, D.H. Regulation of mucosal addressin cell adhesion molecule 1 expression in human and mice by vascular adhesion protein 1 amine oxidase activity. *Hepatology* **2011**, *53*, 661–672. [[CrossRef](#)]
34. Trivedi, P.J.; Tickle, J.; Vesterhus, M.N.; Eddowes, P.J.; Bruns, T.; Vainio, J.; Parker, R.; Smith, D.; Liaskou, E.; Thorbjørnsen, L.W.; et al. Vascular adhesion protein-1 is elevated in primary sclerosing cholangitis, is predictive of clinical outcome and facilitates recruitment of gut-tropic lymphocytes to liver in a substrate-dependent manner. *Gut* **2018**, *67*, 1135–1145. [[CrossRef](#)] [[PubMed](#)]
35. Schippers, A.; Kochut, A.; Pabst, O.; Frischmann, U.; Clahsen, T.; Tenbrock, K.; Muller, W.; Wagner, N. beta7 integrin controls immunogenic and tolerogenic mucosal B cell responses. *Clin. Immunol.* **2012**, *144*, 87–97. [[CrossRef](#)]
36. Kuklin, N.A.; Rott, L.; Feng, N.; Conner, M.E.; Wagner, N.; Muller, W.; Greenberg, H.B. Protective intestinal anti-rotavirus B cell immunity is dependent on alpha 4 beta 7 integrin expression but does not require IgA antibody production. *J. Immunol.* **2001**, *166*, 1894–1902. [[CrossRef](#)]
37. Schippers, A.; Leuker, C.; Pabst, O.; Kochut, A.; Prochnow, B.; Gruber, A.D.; Leung, E.; Krissansen, G.W.; Wagner, N.; Muller, W. Mucosal addressin cell-adhesion molecule-1 controls plasma-cell migration and function in the small intestine of mice. *Gastroenterology* **2009**, *137*, 924–933. [[CrossRef](#)]
38. Tang, M.L.; Steeber, D.A.; Zhang, X.Q.; Tedder, T.F. Intrinsic differences in L-selectin expression levels affect T and B lymphocyte subset-specific recirculation pathways. *J. Immunol.* **1998**, *10*, 5113–5132. [[CrossRef](#)]
39. Stevens, S.K.; Weissman, I.L.; Butcher, E.C. Differences in the migration of B and T lymphocytes: Organ-selective localization in vivo and the role of lymphocyte-endothelial cell recognition. *J. Immunol.* **1982**, *128*, 844–851. [[CrossRef](#)]
40. Lee, M.; Kiefel, H.; LaJevic, M.D.; Macauley, M.S.; Kawashima, H.; O'Hara, E.; Pan, J.; Paulson, J.C.; Butcher, E.C. Transcriptional programs of lymphoid tissue capillary and high endothelium reveal control mechanisms for lymphocyte homing. *Nat. Immunol.* **2014**, *15*, 982–995. Erratum in *Nat. Immunol.* **2015**, *16*, 214. [[CrossRef](#)]
41. Brandt, E.B.; Zimmermann, N.; Muntel, E.E.; Yamada, Y.; Pope, S.M.; Mishra, A.; Hogan, S.P.; Rothenberg, M.E. The alpha4beta7-integrin is dynamically expressed on murine eosinophils and involved in eosinophil trafficking to the intestine. *Clin. Exp. Allergy* **2006**, *36*, 543–553. [[CrossRef](#)]
42. Olbrich, C.L.; Simerly, L.; de Zoeten, E.F.; Furuta, G.T.; Spencer, L.A. Climbing New Mountains: How Antibodies Blocking alpha4beta7 Integrins Tamed Eosinophilic Inflammation of the Intestinal Tract. *Dig. Dis. Sci.* **2019**, *64*, 2068–2071. [[CrossRef](#)] [[PubMed](#)]
43. Kluger, M.A.; Melderis, S.; Nosko, A.; Goerke, B.; Luig, M.; Meyer, M.C.; Turner, J.; Meyer-Schwesinger, C.; Wegscheid, C.; Tiegs, G.; et al. Treg17 cells are programmed by Stat3 to suppress Th17 responses in systemic lupus. *Kidney Int.* **2016**, *89*, 158–166. [[CrossRef](#)]
44. Bhaskaran, N.; Faddoul, F.; Paes Da Silva, A.; Jayaraman, S.; Schneider, E.; Mamileti, P.; Weinberg, A.; Pandiyan, P. IL-1beta-MyD88-mTOR Axis Promotes Immune-Protective IL-17A (+) Foxp3 (+) Cells During Mucosal Infection and Is Dysregulated with Aging. *Front. Immunol.* **2020**, *11*, 595936. [[CrossRef](#)]
45. Fidelle, M.; Rauber, C.; Alves, C.S.C.; Tian, A.L.; Lahmar, I.; de La Varenne, A.M.; Zhao, L.; Thelemaque, C.; Lebhar, I.; Messaoudene, M.; et al. A microbiota-modulated checkpoint directs immunosuppressive intestinal T cells into cancers. *Science* **2023**, *380*, eabo2296. [[CrossRef](#)] [[PubMed](#)]
46. Bai, R.; Li, Z.; Lv, S.; Wang, R.; Hua, W.; Wu, H.; Dai, L. Persistent Inflammation and Non-AIDS Comorbidities During ART: Coming of the Age of Monocytes. *Front. Immunol.* **2022**, *13*, 820480. [[CrossRef](#)] [[PubMed](#)]
47. Schleier, L.; Wiendl, M.; Heidbreder, K.; Binder, M.; Atreya, R.; Rath, T.; Becker, E.; Schulz-Kuhnt, A.; Stahl, A.; Schulze, L.L.; et al. Non-classical monocyte homing to the gut via  $\alpha 4 \beta 7$  integrin mediates macrophage-dependent intestinal wound healing. *Gut* **2020**, *69*, 252–263. [[CrossRef](#)]

48. Li, Q.; Li, Y.; Zhang, B.; Zhai, Y.; Wen, L.; Cheng, G.; Zhao, Z. The role of innate lymphoid cells in kidney disease. *Cytokine Growth Factor Rev.* **2025**, *85*, 146–157. [\[CrossRef\]](#)
49. Kim, M.H.; Taparowsky, E.J.; Kim, C.H. Retinoic Acid Differentially Regulates the Migration of Innate Lymphoid Cell Subsets to the Gut. *Immunity* **2015**, *43*, 107–119. [\[CrossRef\]](#)
50. Albillos, A.; de Gottardi, A.; Rescigno, M. The gut-liver axis in liver disease: Pathophysiological basis for therapy. *J. Hepatol.* **2020**, *72*, 558–577. [\[CrossRef\]](#)
51. Baffy, G. Potential mechanisms linking gut microbiota and portal hypertension. *Liver Int.* **2019**, *39*, 598–609. [\[CrossRef\]](#) [\[PubMed\]](#)
52. Zhou, Y.; Zhang, H.; Yao, Y.; Zhang, X.; Guan, Y.; Zheng, F. CD4 (+) T cell activation and inflammation in NASH-related fibrosis. *Front. Immunol.* **2022**, *13*, 967410. [\[CrossRef\]](#) [\[PubMed\]](#)
53. Bhat, A.A.; Uppada, S.; Achkar, I.W.; Hashem, S.; Yadav, S.K.; Shanmugakonar, M.; Al-Naemi, H.A.; Haris, M.; Uddin, S. Tight Junction Proteins and Signaling Pathways in Cancer and Inflammation: A Functional Crosstalk. *Front. Physiol.* **2018**, *9*, 1942. [\[CrossRef\]](#)
54. Picarella, D.; Hurlbut, P.; Rottman, J.; Shi, X.; Butcher, E.; Ringler, D.J. Monoclonal antibodies specific for beta 7 integrin and mucosal addressin cell adhesion molecule-1 (MAdCAM-1) reduce inflammation in the colon of scid mice reconstituted with CD45RB<sup>high</sup> CD4+ T cells. *J. Immunol.* **1997**, *158*, 2099–2106. [\[CrossRef\]](#)
55. Bachmann, C.; Klibanov, A.L.; Olson, T.S.; Sonnenschein, J.R.; Rivera-Nieves, J.; Cominelli, F.; Ley, K.F.; Lindner, J.R.; Pizarro, T.T. Targeting mucosal addressin cellular adhesion molecule (MAdCAM)-1 to noninvasively image experimental Crohn's disease. *Gastroenterology* **2006**, *130*, 8–16. [\[CrossRef\]](#)
56. Poole, R.M. Vedolizumab: First global approval. *Drugs* **2014**, *74*, 1293–1303. [\[CrossRef\]](#)
57. Wyant, T.; Leach, T.; Sankoh, S.; Wang, Y.; Paolino, J.; Pasetti, M.F.; Feagan, B.G.; Parikh, A. Vedolizumab affects antibody responses to immunisation selectively in the gastrointestinal tract: Randomised controlled trial results. *Gut* **2014**, *64*, 77–83. [\[CrossRef\]](#)
58. Akhtar, H.J.; Nguyen, T.M.; Ma, C.; Jairath, V. Vedolizumab for the Treatment of Noninflammatory Bowel Disease Related Enteropathy. *Clin. Gastroenterol. Hepatol.* **2022**, *20*, e614–e623. [\[CrossRef\]](#)
59. Weismuller, T.J.; Trivedi, P.J.; Bergquist, A.; Imam, M.; Lenzen, H.; Ponsioen, C.Y.; Holm, K.; Gotthardt, D.; Farkkila, M.A.; Marschall, H.U.; et al. Patient Age, Sex, and Inflammatory Bowel Disease Phenotype Associate with Course of Primary Sclerosing Cholangitis. *Gastroenterology* **2017**, *152*, 1975–1984. [\[CrossRef\]](#) [\[PubMed\]](#)
60. Sorensen, J.O.; Nielsen, O.H.; Andersson, M.; Ainsworth, M.A.; Ytting, H.; Belard, E.; Jess, T. Inflammatory bowel disease with primary sclerosing cholangitis: A Danish population-based cohort study 1977–2011. *Liver Int.* **2018**, *38*, 532–541. [\[CrossRef\]](#) [\[PubMed\]](#)
61. Henriksen, E.K.; Jorgensen, K.K.; Kaveh, F.; Holm, K.; Hamm, D.; Olweus, J.; Melum, E.; Chung, B.K.; Eide, T.J.; Lundin, K.E.; et al. Gut and liver T-cells of common clonal origin in primary sclerosing cholangitis-inflammatory bowel disease. *J. Hepatol.* **2017**, *66*, 116–122. [\[CrossRef\]](#) [\[PubMed\]](#)
62. Ponsioen, C.Y.; Kuiper, H.; Ten, K.F.; van Milligen, D.W.M.; van Deventer, S.J.; Tytgat, G.N. Immunohistochemical analysis of inflammation in primary sclerosing cholangitis. *Eur. J. Gastroenterol. Hepatol.* **1999**, *11*, 769–774. [\[CrossRef\]](#) [\[PubMed\]](#)
63. Grant, A.J.; Lalor, P.F.; Hubscher, S.G.; Briskin, M.; Adams, D.H. MAdCAM-1 expressed in chronic inflammatory liver disease supports mucosal lymphocyte adhesion to hepatic endothelium (MAdCAM-1 in chronic inflammatory liver disease). *Hepatology* **2001**, *33*, 1065–1072. [\[CrossRef\]](#)
64. Hillan, K.J.; Hagler, K.E.; MacSween, R.N.M.; Ryan, A.M.; Renz, M.E.; Chiu, H.H.; Ferrier, R.K.; Bird, G.L.; Dhillon, A.P.; Ferrell, L.D.; et al. Expression of the mucosal vascular addressin, MAdCAM-1, in inflammatory liver disease. *Liver* **1999**, *19*, 509–518. [\[CrossRef\]](#)
65. de Krijger, M.; Visseren, T.; Wildenberg, M.E.; Hooijer, G.; Verstegen, M.; van der Laan, L.; de Jonge, W.J.; Verheij, J.; Ponsioen, C.Y. Characterization of gut-homing molecules in non-endstage livers of patients with primary sclerosing cholangitis and inflammatory bowel disease. *J. Transl. Autoimmun.* **2020**, *3*, 100054. [\[CrossRef\]](#)
66. Graham, J.J.; Mukherjee, S.; Yuksel, M.; Sanabria, M.R.; Si, T.; Huang, Z.; Huang, X.; Abu Arqoub, H.; Patel, V.; McPhail, M.; et al. Aberrant hepatic trafficking of gut-derived T cells is not specific to primary sclerosing cholangitis. *Hepatology* **2022**, *75*, 518–530. [\[CrossRef\]](#)
67. Rai, R.P.; Liu, Y.; Iyer, S.S.; Liu, S.; Gupta, B.; Desai, C.; Kumar, P.; Smith, T.; Singhi, A.D.; Nusrat, A.; et al. Blocking integrin  $\alpha 4\beta 7$ -mediated CD4 T cell recruitment to the intestine and liver protects mice from western diet-induced non-alcoholic steatohepatitis. *J. Hepatol.* **2020**, *73*, 1013–1022. [\[CrossRef\]](#) [\[PubMed\]](#)
68. Drescher, H.K.; Schippers, A.; Clahsen, T.; Sahin, H.; Noels, H.; Hornef, M.; Wagner, N.; Trautwein, C.; Streetz, K.L.; Kroy, D.C. beta (7)-Integrin and MAdCAM-1 play opposing roles during the development of non-alcoholic steatohepatitis. *J. Hepatol.* **2017**, *66*, 1251–1264. [\[CrossRef\]](#) [\[PubMed\]](#)

69. Ala, A.; Brown, D.; Khan, K.; Standish, R.; Odin, J.A.; Fiel, M.I.; Schiano, T.D.; Hillan, K.J.; Rahman, S.A.; Hodgson, H.J.F.; et al. Mucosal Addressin Cell Adhesion Molecule (MAdCAM-1) Expression Is Upregulated in the Cirrhotic Liver and Immunolocalises to the Peribiliary Plexus and Lymphoid Aggregates. *Dig. Dis. Sci.* **2013**, *58*, 2528–2541.
70. Gupta, B.; Rai, R.P.; Pal, P.B.; Rossmiller, D.; Chaudhary, S.; Chiaro, A.; Seaman, S.; Singhi, A.D.; Liu, S.; Monga, S.P.; et al. Selective Targeting of alpha (4) beta (7)/MAdCAM-1 Axis Suppresses Fibrosis Progression by Reducing Proinflammatory T Cell Recruitment to the Liver. *Cells* **2024**, *13*, 756. [[CrossRef](#)]
71. Tsuda, M.; Ambrosini, Y.M.; Zhang, W.; Yang, G.X.; Ando, Y.; Rong, G.; Tsuneyama, K.; Sumida, K.; Shimoda, S.; Bowlus, C.L.; et al. Fine phenotypic and functional characterization of effector cluster of differentiation 8 positive T cells in human patients with primary biliary cirrhosis. *Hepatology* **2011**, *54*, 1293–1302. [[CrossRef](#)]
72. Schippers, A.; Hubel, J.; Heymann, F.; Clahsen, T.; Eswaran, S.; Schleputz, S.; Pullen, R.; Gassler, N.; Tenbrock, K.; Tacke, F.; et al. MAdCAM-1/alpha4beta7 Integrin-Mediated Lymphocyte/Endothelium Interactions Exacerbate Acute Immune-Mediated Hepatitis in Mice. *Cell. Mol. Gastroenterol. Hepatol.* **2021**, *11*, 1227–1250. [[CrossRef](#)] [[PubMed](#)]
73. Li, X.; Shen, J.; Ran, Z. Crosstalk between the gut and the liver via susceptibility loci: Novel advances in inflammatory bowel disease and autoimmune liver disease. *Clin. Immunol.* **2017**, *175*, 115–123. [[CrossRef](#)]
74. Fu, S.; Ni, T.; Zhang, M.; Ren, D.; Feng, Y.; Yao, N.; Zhang, X.; Wang, R.; Xu, W.; Yang, N.; et al. Cholinergic Anti-inflammatory Pathway Attenuates Acute Liver Failure Through Inhibiting MAdCAM1/alpha4beta7-mediated Gut-derived Proinflammatory Lymphocytes Accumulation. *Cell. Mol. Gastroenterol. Hepatol.* **2024**, *17*, 199–217. [[CrossRef](#)] [[PubMed](#)]
75. Chu, Z.; Chen, H.; Ma, Y.; Zhou, Y.; Zhang, M.; Zhang, P.; Qin, H. Lactobacillus plantarum Prevents the Upregulation of Adhesion Molecule Expression in an Experimental Colitis Model. *Dig. Dis. Sci.* **2010**, *55*, 2505–2513. [[CrossRef](#)]
76. Laurans, L.; Mouttoulingam, N.; Chajadine, M.; Lavelle, A.; Diedisheim, M.; Bacquer, E.; Creusot, L.; Suffee, N.; Esposito, B.; Melhem, N.J.; et al. An obesogenic diet increases atherosclerosis through promoting microbiota dysbiosis-induced gut lymphocyte trafficking into the periphery. *Cell Rep.* **2023**, *42*, 113350. [[CrossRef](#)]
77. Han, B.; Lv, X.; Liu, G.; Li, S.; Fan, J.; Chen, L.; Huang, Z.; Lin, G.; Xu, X.; Huang, Z.; et al. Gut microbiota-related bile acid metabolism-FXR/TGR5 axis impacts the response to anti-alpha4beta7-integrin therapy in humanized mice with colitis. *Gut Microbes* **2023**, *15*, 2232143. [[CrossRef](#)] [[PubMed](#)]
78. Gao, R.Y.; Shearn, C.T.; Orlicky, D.J.; Battista, K.D.; Alexeev, E.E.; Cartwright, I.M.; Lanis, J.M.; Kostelecky, R.E.; Ju, C.; Colgan, S.P.; et al. Bile acids modulate colonic MAdCAM-1 expression in a murine model of combined cholestasis and colitis. *Mucosal Immunol.* **2021**, *14*, 479–490. [[CrossRef](#)]
79. Jacob, N.; Jaiswal, S.; Maheshwari, D.; Nallabelli, N.; Khatri, N.; Bhatia, A.; Bal, A.; Malik, V.; Verma, S.; Kumar, R.; et al. Butyrate induced Tregs are capable of migration from the GALT to the pancreas to restore immunological tolerance during type-1 diabetes. *Sci. Rep.* **2020**, *10*, 19120. [[CrossRef](#)]
80. Vinolo, M.A.; Rodrigues, H.G.; Hatanaka, E.; Hebeda, C.B.; Farsky, S.H.; Curi, R. Short-chain fatty acids stimulate the migration of neutrophils to inflammatory sites. *Clin. Sci.* **2009**, *117*, 331–338. [[CrossRef](#)]
81. Tsuzuki, Y.; Miura, S.; Kurose, I.; Suematsu, M.; Higuchi, H.; Shigematsu, T.; Kimura, H.; Serizawa, H.; Hokari, R.; Akiba, Y.; et al. Enhanced lymphocyte interaction in postcapillary venules of Peyer's patches during fat absorption in rats. *Gastroenterology* **1997**, *112*, 813–825. [[CrossRef](#)]
82. Nakamura, Y.K.; Janowitz, C.; Metea, C.; Asquith, M.; Karstens, L.; Rosenbaum, J.T.; Lin, P. Short chain fatty acids ameliorate immune-mediated uveitis partially by altering migration of lymphocytes from the intestine. *Sci. Rep.* **2017**, *7*, 11745. [[CrossRef](#)]
83. Kim, S.H.; Cho, B.H.; Kiyono, H.; Jang, Y.S. Microbiota-derived butyrate suppresses group 3 innate lymphoid cells in terminal ileal Peyer's patches. *Sci. Rep.* **2017**, *7*, 3980. [[CrossRef](#)]
84. Emgard, J.; Kammoun, H.; Garcia-Cassani, B.; Chesne, J.; Parigi, S.M.; Jacob, J.M.; Cheng, H.W.; Evren, E.; Das, S.; Czarnewski, P.; et al. Oxysterol Sensing through the Receptor GPR183 Promotes the Lymphoid-Tissue-Inducing Function of Innate Lymphoid Cells and Colonic Inflammation. *Immunity* **2018**, *48*, 120–132. [[CrossRef](#)]
85. Willinger, T. Metabolic Control of Innate Lymphoid Cell Migration. *Front. Immunol.* **2019**, *10*, 2010. [[CrossRef](#)]
86. Qiu, J.; Heller, J.J.; Guo, X.; Chen, Z.M.; Fish, K.; Fu, Y.X.; Zhou, L. The aryl hydrocarbon receptor regulates gut immunity through modulation of innate lymphoid cells. *Immunity* **2012**, *36*, 92–104. [[CrossRef](#)]
87. De Giovanni, M.; Tam, H.; Valet, C.; Xu, Y.; Looney, M.R.; Cyster, J.G. GPR35 promotes neutrophil recruitment in response to serotonin metabolite 5-HIAA. *Cell* **2022**, *185*, 815–830. [[CrossRef](#)] [[PubMed](#)]
88. Haas, R.; Smith, J.; Rocher-Ros, V.; Nadkarni, S.; Montero-Melendez, T.; D'Acquisto, F.; Bland, E.J.; Bombardieri, M.; Pitzalis, C.; Perretti, M.; et al. Lactate Regulates Metabolic and Pro-inflammatory Circuits in Control of T Cell Migration and Effector Functions. *PLoS Biol.* **2015**, *13*, e1002202. [[CrossRef](#)] [[PubMed](#)]
89. Dastmalchi, F.; Karachi, A.; Yang, C.; Azari, H.; Sayour, E.J.; Dechkovskaia, A.; Vlasak, A.L.; Saia, M.E.; Lovatton, R.E.; Mitchell, D.A.; et al. Sarcosine promotes trafficking of dendritic cells and improves efficacy of anti-tumor dendritic cell vaccines via CXC chemokine family signaling. *J. Immunother. Cancer* **2019**, *7*, 321. [[CrossRef](#)] [[PubMed](#)]

90. Agard, M.; Asakrah, S.; Morici, L.A. PGE (2) suppression of innate immunity during mucosal bacterial infection. *Front. Cell. Infect. Microbiol.* **2013**, *3*, 45. [\[CrossRef\]](#)
91. Pang, X.; He, X.; Qiu, Z.; Zhang, H.; Xie, R.; Liu, Z.; Gu, Y.; Zhao, N.; Xiang, Q.; Cui, Y. Targeting integrin pathways: Mechanisms and advances in therapy. *Signal Transduct. Target. Ther.* **2023**, *8*, 1. [\[CrossRef\]](#)
92. Mitroulis, I.; Alexaki, V.I.; Kourtzelis, I.; Ziogas, A.; Hajishengallis, G.; Chavakis, T. Leukocyte integrins: Role in leukocyte recruitment and as therapeutic targets in inflammatory disease. *Pharmacol. Ther.* **2015**, *147*, 123–135. [\[CrossRef\]](#)
93. Leonardi, C.L.; Papp, K.A.; Gordon, K.B.; Menter, A.; Feldman, S.R.; Caro, I.; Walicke, P.A.; Compton, P.G.; Gottlieb, A.B. Extended efalizumab therapy improves chronic plaque psoriasis: Results from a randomized phase III trial. *J. Am. Acad. Dermatol.* **2005**, *52*, 425–433. [\[CrossRef\]](#)
94. Weger, W. Current status and new developments in the treatment of psoriasis and psoriatic arthritis with biological agents. *Br. J. Pharmacol.* **2010**, *160*, 810–820. [\[CrossRef\]](#)
95. Keating, G.M. Lifitegrast Ophthalmic Solution 5%: A Review in Dry Eye Disease. *Drugs* **2017**, *77*, 201–208. [\[CrossRef\]](#)
96. Pavlick, K.P.; Ostanin, D.V.; Furr, K.L.; Laroux, F.S.; Brown, C.M.; Gray, L.; Kevil, C.G.; Grisham, M.B. Role of T-cell-associated lymphocyte function-associated antigen-1 in the pathogenesis of experimental colitis. *Int. Immunol.* **2006**, *18*, 389–398. [\[CrossRef\]](#)
97. Palmen, M.J.; Dijkstra, C.D.; van der Ende, M.B.; Pena, A.S.; van Rees, E.P. Anti-CD11b/CD18 antibodies reduce inflammation in acute colitis in rats. *Clin. Exp. Immunol.* **1995**, *101*, 351–356. [\[CrossRef\]](#)
98. Tang, M.; Jiang, Y.; Jia, H.; Patpur, B.K.; Yang, B.; Li, J.; Yang, C. Osteopontin acts as a negative regulator of autophagy accelerating lipid accumulation during the development of nonalcoholic fatty liver disease. *Artif. Cells Nanomed. Biotechnol.* **2020**, *48*, 159–168. [\[CrossRef\]](#) [\[PubMed\]](#)
99. Henderson, N.C.; Arnold, T.D.; Katamura, Y.; Giacomini, M.M.; Rodriguez, J.D.; McCarty, J.H.; Pellicoro, A.; Raschperger, E.; Betsholtz, C.; Ruminiski, P.G.; et al. Targeting of alphav integrin identifies a core molecular pathway that regulates fibrosis in several organs. *Nat. Med.* **2013**, *19*, 1617–1624. [\[CrossRef\]](#) [\[PubMed\]](#)
100. Liu, J.; Quan, J.; Feng, J.; Zhang, Q.; Xu, Y.; Liu, J.; Huang, W.; Liu, J.; Tian, L. High glucose regulates LN expression in human liver sinusoidal endothelial cells through ROS/integrin alphavbeta3 pathway. *Environ. Toxicol. Pharmacol.* **2016**, *42*, 231–236. [\[CrossRef\]](#) [\[PubMed\]](#)
101. Rokugawa, T.; Konishi, H.; Ito, M.; Iimori, H.; Nagai, R.; Shimosegawa, E.; Shimosegawa, E.; Hatazawa, J.; Abe, K. Evaluation of hepatic integrin alphavbeta3 expression in non-alcoholic steatohepatitis (NASH) model mouse by (18) F-FPP-RGD (2) PET. *EJNMMI Res.* **2018**, *8*, 40. [\[CrossRef\]](#)
102. Hiroyama, S.; Rokugawa, T.; Ito, M.; Iimori, H.; Morita, I.; Maeda, H.; Fujisawa, K.; Matsunaga, K.; Shimosegawa, E.; Abe, K. Quantitative evaluation of hepatic integrin alpha(v)beta(3) expression by positron emission tomography imaging using (18) F-FPP-RGD (2) in rats with non-alcoholic steatohepatitis. *EJNMMI Res.* **2020**, *10*, 118. [\[CrossRef\]](#)
103. Guo, Q.; Furuta, K.; Lucien, F.; Gutierrez Sanchez, L.H.; Hirsova, P.; Krishnan, A.; Kabashima, A.; Pavelko, K.D.; Madden, B.; Alhuwaish, H.; et al. Integrin  $\beta$ 1-enriched extracellular vesicles mediate monocyte adhesion and promote liver inflammation in murine NASH. *J. Hepatol.* **2019**, *71*, 1193–1205. [\[CrossRef\]](#)
104. Feagan, B.G.; Rutgeerts, P.; Sands, B.E.; Hanauer, S.; Colombel, J.; Sandborn, W.J.; Van, A.G.; Axler, J.; Kim, H.-J.; Danese, S.; et al. Vedolizumab as Induction and Maintenance Therapy for Ulcerative Colitis. *N. Engl. J. Med.* **2013**, *369*, 699–710. [\[CrossRef\]](#) [\[PubMed\]](#)
105. Sandborn, W.J.; Feagan, B.G.; Rutgeerts, P.; Hanauer, S.; Colombel, J.; Sands, B.E.; Lukas, M.; Fedorak, R.N.; Lee, S.; Bressler, B.; et al. Vedolizumab as Induction and Maintenance Therapy for Crohn's Disease. *N. Engl. J. Med.* **2013**, *369*, 711–721. [\[CrossRef\]](#) [\[PubMed\]](#)
106. Lynch, K.D.; Chapman, R.W.; Keshav, S.; Montano-Loza, A.J.; Mason, A.L.; Kremer, A.E.; Vetter, M.; de Krijger, M.; Ponsioen, C.Y.; Trivedi, P.; et al. Effects of Vedolizumab in Patients With Primary Sclerosing Cholangitis and Inflammatory Bowel Diseases. *Clin. Gastroenterol. Hepatol.* **2020**, *18*, 179–187. [\[CrossRef\]](#) [\[PubMed\]](#)
107. Ghosh, S.; Goldin, E.; Gordon, F.H.; Malchow, H.A.; Rask-Madsen, J.; Rutgeerts, P.; Vyhnaček, P.; Zádorová, Z.; Palmer, T.; Donoghue, S. Natalizumab for Active Crohn's Disease. *N. Engl. J. Med.* **2003**, *1*, 24–32. [\[CrossRef\]](#)
108. Berger, J.R. Natalizumab and progressive multifocal leucoencephalopathy. *Ann. Rheum. Dis.* **2006**, *65* (Suppl. S3), iii48–iii53. [\[CrossRef\]](#)
109. Sandborn, W.J.; Cyrille, M.; Hansen, M.B.; Feagan, B.G.; Loftus, E.V.; Rogler, G.; Vermeire, S.; Cruz, M.L.; Yang, J.; Boedigheimer, M.J.; et al. Efficacy and Safety of Abrilumab in a Randomized, Placebo-Controlled Trial for Moderate-to-Severe Ulcerative Colitis. *Gastroenterology* **2019**, *156*, 946–957. [\[CrossRef\]](#)
110. Peyrin-Biroulet, L.; Hart, A.; Bossuyt, P.; Long, M.; Allez, M.; Juillerat, P.; Armuzzi, A.; Loftus, E.V., Jr.; Ostad-Saffari, E.; Scalori, A.; et al. Etrolizumab as induction and maintenance therapy for ulcerative colitis in patients previously treated with tumour necrosis factor inhibitors (HICKORY): A phase 3, randomised, controlled trial. *Lancet Gastroenterol. Hepatol.* **2022**, *7*, 128–140. [\[CrossRef\]](#)

111. Rubin, D.T.; Dotan, I.; DuVall, A.; Bouhnik, Y.; Radford-Smith, G.; Higgins, P.; Mishkin, D.S.; Arrisi, P.; Scalori, A.; Oh, Y.S.; et al. Etrolizumab versus adalimumab or placebo as induction therapy for moderately to severely active ulcerative colitis (HIBISCUS): Two phase 3 randomised, controlled trials. *Lancet Gastroenterol. Hepatol.* **2022**, *7*, 17–27. [\[CrossRef\]](#)
112. Danese, S.; Colombel, J.F.; Lukas, M.; Gisbert, J.P.; D’Haens, G.; Hayee, B.; Panaccione, R.; Kim, H.S.; Reinisch, W.; Tyrrell, H.; et al. Etrolizumab versus infliximab for the treatment of moderately to severely active ulcerative colitis (GARDENIA): A randomised, double-blind, double-dummy, phase 3 study. *Lancet Gastroenterol. Hepatol.* **2022**, *7*, 118–127. [\[CrossRef\]](#) [\[PubMed\]](#)
113. Vermeire, S.; Danese, S.; Sandborn, W.J.; Schreiber, S.; Hanauer, S.; D’Haens, G.; Nagy, P.; Thakur, M.; Bliss, C.; Cataldi, F.; et al. Efficacy and Safety of the Anti-Mucosal Addressin Cell Adhesion Molecule-1 Antibody Ontamalimab in Patients with Moderate-to-Severe Ulcerative Colitis or Crohn’s Disease. *J. Crohn’s Colitis* **2024**, *18*, 708–719. [\[CrossRef\]](#)
114. Slack, R.J.; Macdonald, S.; Roper, J.A.; Jenkins, R.G.; Hatley, R. Emerging therapeutic opportunities for integrin inhibitors. *Nat. Rev. Drug Discov.* **2022**, *21*, 60–78. [\[CrossRef\]](#)
115. Sands, B.E.; Schreiber, S.; Danese, S.; Kierkus, J.; Abhyankar, B.; Choi, M.Y.; Soo, C.; Wu, Y.; Sun, F.; Lee, D.; et al. A Phase 2 Study of MORF-057, an Oral alpha4beta7 Integrin Inhibitor in Moderately to Severely Active Ulcerative Colitis. *Clin. Gastroenterol. Hepatol.* **2025**, in press.
116. Matsuoka, K.; Watanabe, M.; Ohmori, T.; Nakajima, K.; Ishida, T.; Ishiguro, Y.; Kanke, K.; Kobayashi, K.; Hirai, F.; Watanabe, K.; et al. AJM300 (carotegrast methyl), an oral antagonist of alpha4-integrin, as induction therapy for patients with moderately active ulcerative colitis: A multicentre, randomised, double-blind, placebo-controlled, phase 3 study. *Lancet Gastroenterol. Hepatol.* **2022**, *7*, 648–657. [\[CrossRef\]](#)
117. Dhillon, S. Carotegrast Methyl: First Approval. *Drugs* **2022**, *82*, 1011–1016. [\[CrossRef\]](#) [\[PubMed\]](#)
118. Sandborn, W.J.; Mattheakis, L.C.; Modi, N.B.; Pugatch, D.; Bressler, B.; Lee, S.; Bhandari, R.; Kanwar, B.; Shames, R.; D’Haens, G.; et al. PTG-100, an Oral alpha4beta7 Antagonist Peptide: Preclinical Development and Phase 1 and 2a Studies in Ulcerative Colitis. *Gastroenterology* **2021**, *161*, 1853–1864. [\[CrossRef\]](#)
119. FDA-Approved Drugs: NATALIZUMAB. Available online: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&ApplNo=125104> (accessed on 19 February 2025).
120. FDA-Approved Drugs: VEDOLIZUMAB. Available online: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&ApplNo=125476> (accessed on 19 February 2025).
121. Habtezion, A.; Nguyen, L.P.; Hadeiba, H.; Butcher, E.C. Leukocyte Trafficking to the Small Intestine and Colon. *Gastroenterology* **2016**, *150*, 340–354. [\[CrossRef\]](#)
122. Takeda, A.; Sasaki, N.; Miyasaka, M. The molecular cues regulating immune cell trafficking. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* **2017**, *93*, 183–195. [\[CrossRef\]](#)
123. Bonfanti, E.; Bracco, C.; Biancheri, P.; Falcetta, A.; Badinella, M.M.; Melchio, R.; Fenoglio, L. Fever During Anti-integrin Therapy: New Immunodeficiency. *Eur. J. Case Rep. Intern. Med.* **2020**, *7*, 1288.
124. Fedyk, E.R.; Wyant, T.; Yang, L.; Csizmadia, V.; Burke, K.; Yang, H.; Kadambi, V.J. Exclusive antagonism of the  $\alpha 4 \beta 7$  integrin by vedolizumab confirms the gut-selectivity of this pathway in primates. *Inflamm. Bowel Dis.* **2012**, *18*, 2107–2119. [\[CrossRef\]](#)
125. Kapila, N.; Flocco, G.; Shen, B.; Esfeh, J.M. The Use of Vedolizumab in Patients with Concomitant Cirrhosis and Crohn’s Disease. *Cureus* **2018**, *10*, e3080. [\[CrossRef\]](#)
126. Huang, X.; Yang, J.; Hu, Y.; Li, S.; Cai, Z.; Li, C.; Qin, W.; Zhang, G. Protective Effects of GalNac-Modified Red Blood Cell-Derived Extracellular Vesicles Against Liver Diseases. *Int. J. Nanomed.* **2025**, *20*, 8993–9017. [\[CrossRef\]](#)
127. Fei, B.; Zhao, Y.; Wang, J.; Wen, P.; Li, J.; Tanaka, M.; Wang, Z.; Li, S. Leveraging adrenergic receptor blockade for enhanced nonalcoholic fatty liver disease treatment via a biomimetic nanoplatfrom. *J. Nanobiotechnol.* **2024**, *22*, 591. [\[CrossRef\]](#) [\[PubMed\]](#)
128. Fei, Y.; Zhang, S.; Han, S.; Qiu, B.; Lu, Y.; Huang, W.; Li, F.; Chen, D.; Berglund, B.; Xiao, H.; et al. The Role of Dihydroresveratrol in Enhancing the Synergistic Effect of Ligilactobacillus salivarius Li01 and Resveratrol in Ameliorating Colitis in Mice. *Research* **2022**, *2022*, 9863845. [\[CrossRef\]](#)
129. Guzzo, C.; Ichikawa, D.; Park, C.; Phillips, D.; Liu, Q.; Zhang, P.; Kwon, A.; Miao, H.; Lu, J.; Rehm, C.; et al. Virion incorporation of integrin alpha4beta7 facilitates HIV-1 infection and intestinal homing. *Sci. Immunol.* **2017**, *2*, eaam7341. [\[CrossRef\]](#)
130. Ansari, A.A.; Reimann, K.A.; Mayne, A.E.; Takahashi, Y.; Stephenson, S.T.; Wang, R.; Wang, X.; Li, J.; Price, A.A.; Little, D.M.; et al. Blocking of alpha4beta7 gut-homing integrin during acute infection leads to decreased plasma and gastrointestinal tissue viral loads in simian immunodeficiency virus-infected rhesus macaques. *J. Immunol.* **2011**, *186*, 1044–1059. [\[CrossRef\]](#) [\[PubMed\]](#)
131. Sneller, M.C.; Clarridge, K.E.; Seamon, C.; Shi, V.; Zorawski, M.D.; Justement, J.S.; Blazkova, J.; Huiting, E.D.; Proshan, M.A.; Mora, J.R.; et al. An open-label phase 1 clinical trial of the anti- $\alpha 4 \beta 7$  monoclonal antibody vedolizumab in HIV-infected individuals. *Sci. Transl. Med.* **2019**, *11*, eaax3447. [\[CrossRef\]](#)
132. Liu, Q.; Lusso, P. Integrin alpha4beta7 in HIV-1 infection: A critical review. *J. Leukoc. Biol.* **2020**, *108*, 627–632. [\[CrossRef\]](#) [\[PubMed\]](#)
133. Grove, R.A.; Shackelford, S.; Sopper, S.; Pirruccello, S.; Horrigan, J.; Havrdova, E.; Gold, M.; Graff, O. Leukocyte counts in cerebrospinal fluid and blood following firtagrast treatment in subjects with relapsing forms of multiple sclerosis. *Eur. J. Neurol.* **2013**, *20*, 1032–1042. [\[CrossRef\]](#) [\[PubMed\]](#)

134. Cortijo, J.; Sanz, M.; Iranzo, A.; Montesinos, J.L.; Nabah, Y.N.A.; Alfón, J.; Gómez, L.A.; Merlos, M.; Morcillo, E.J. A small molecule, orally active,  $\alpha_4\beta_1/\alpha_4\beta_7$  dual antagonist reduces leukocyte infiltration and airway hyper-responsiveness in an experimental model of allergic asthma in Brown Norway rats. *Br. J. Pharmacol.* **2006**, *147*, 661. [[CrossRef](#)] [[PubMed](#)]
135. Dattoli, S.D.; Baiula, M.; De Marco, R.; Bedini, A.; Anselmi, M.; Gentilucci, L.; Spampinato, S. DS-70, a novel and potent  $\alpha_4$  integrin antagonist, is an effective treatment for experimental allergic conjunctivitis in guinea pigs. *Br. J. Pharmacol.* **2018**, *175*, 3891–3910. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.