



Proceeding Paper

# Polyunsaturated Fatty Acid Intake among Young Individuals Attending an English University <sup>†</sup>

Antonio Peña-Fernández <sup>1,2,\*</sup> , Edna Segura <sup>3</sup> , María de los Ángeles Peña <sup>4</sup> , Mark D. Evans <sup>2</sup>  
and Manuel Higuera <sup>3</sup>

<sup>1</sup> Department of Surgery, Medical and Social Sciences, Faculty of Medicine and Health Sciences, University of Alcalá, Ctra. Madrid-Barcelona, Km. 33.600, 28871 Alcalá de Henares, Spain

<sup>2</sup> Leicester School of Allied Health Sciences, De Montfort University, Leicester LE1 9BH, UK; mark.evans@dmu.ac.uk

<sup>3</sup> Scientific Computation & Technological Innovation Center (SCoTIC), Universidad de La Rioja, 26006 Logroño, Spain; edna-viviana.segura@alum.unirioja.es (E.S.); manuel.higuera@unirioja.es (M.H.)

<sup>4</sup> Departamento de Ciencias Biomédicas, Universidad de Alcalá, Ctra. Madrid-Barcelona, Km. 33.600, 28871 Alcalá de Henares, Spain; angeles.pena@uah.es

\* Correspondence: antonio.pena-fernandez@dmu.ac.uk or antonio.penafer@uah.es

<sup>†</sup> Presented at the 3rd International Electronic Conference on Nutrients, 1–15 November 2023; Available online: <https://iecn2023.sciforum.net/>.

**Abstract:** Omega-3 long-chain polyunsaturated fatty acids (PUFAs), such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), are poorly synthesised in the human body and are substantially lower in Western diets compared with their shorter-chain omega-3 essential fatty acid precursors,  $\alpha$ -linolenic acid (18:3n-3; ALA). We assessed their intake among 111 (20.45 years old; 78 females) De Montfort University (DMU, England) students. The dietary intakes of total fat (100.55 vs. 81.72;  $p$ -value = 0.032), PUFA (14.61 vs. 12.91; NS), linoleic acid (LA; 3.893 vs. 2.787;  $p$ -value = 0.0019), ALA (0.925 vs. 0.613;  $p$ -value = 0.00008), arachidonic acid (AA; 0.109 vs. 0.082;  $p$ -value = 0.0303), EPA (0.088 vs. 0.075; NS), DHA (0.153 vs. 0.121; NS), and docosapentaenoic acid (DPA; 0.043 vs. 0.032 all in g/day; NS) were significantly higher or higher in male participants, respectively. The dietary intakes of DHA + EPA in the whole group monitored (0.130 + 0.079 = 0.209 g/day) were lower than the RDI of 0.5 g/day, which considers the intake of one to two portions of fish per week. Our results highlight that some DMU students did not meet the nutritional goals for ALA, EPA, and DHA. DMU students should specifically enhance the intake of oily fish (12.422, 13.406, and 10.054 g/day for the overall, female, and male population, respectively), as these intakes only provide around 0.228, 0.246, and 0.184 g of DHA + EPA/day. Education would be required to increase awareness of the importance of consuming more fish among these young adults. Another option would be to encourage the intake of dietary fish oil supplements or the enrichment of food items largely consumed by young British adults with these long-chain PUFAs.

**Keywords:** essential fatty acids; EPA and DHA intakes; university students; ethnicity; Leicester



**Citation:** Peña-Fernández, A.; Segura, E.; Peña, M.d.l.Á.; Evans, M.D.; Higuera, M. Polyunsaturated Fatty Acid Intake among Young Individuals Attending an English University. *Biol. Life Sci. Forum* **2023**, *29*, 17. <https://doi.org/10.3390/IECN2023-15528>

Academic Editor: Maria-Luz Fernandez

Published: 31 October 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Omega-3 long-chain polyunsaturated fatty acids (PUFAs), such as eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3), are poorly synthesised in the human body and are considered essential components of a healthy and balanced diet [1]. Thus, EPA and DHA have been associated with human homeostasis and healthy ageing and are important for appropriate foetal development and neural, retinal, and immune system function. DHA is the principal omega-3 fatty acid in humans' cortical grey matter, so its intake is crucial in patients with dementia and other neurodegenerative disorders involving the progressive degeneration of the central and peripheral nervous systems [2].

However, the intake of EPA and DHA is substantially lower in Western diets compared with their shorter-chain omega-3 essential fatty acid precursors,  $\alpha$ -linolenic acid (18:3n-3; ALA, C<sub>18:3</sub>), although ALA does not provide the health benefits observed with the intake of EPA and DHA. ALA and the omega-6 fatty acid linoleic acid (LA, C<sub>18:2</sub>) are essential fatty acids that cannot be synthesised by humans but can go through different elongation and desaturation steps to produce long-chain PUFAs in the human body, including EPA and DHA, as well as docosapentaenoic acid (DPA, 22:5n-3) and arachidonic acid (ARA, 20:4n-6), which also have important functions in the human body [3,4]. Thus, AA is as essential for brain development as DHA (Conway et al., 2021).

Moreover, high intake of the essential fatty acids ALA and LA has been associated with the pathogenesis of many inflammatory diseases [5].

Owing to the relevance of long-chain PUFAs, the aim of our work was to assess the intake of total PUFAs, EPA, DHA, DPA, ARA, and essential fatty acids in a young cohort of students attending De Montfort University (DMU), a British university located in the East Midlands, United Kingdom.

## 2. Material and Methods

Comprehensive nutrient intake was collected from 111 (20.45  $\pm$  1.16 yrs-old; 78 female) DMU students between 2015–2016 from three different ethnic backgrounds (41 Asia, 41 Africa, and 27 Europe), using a validated variant of the Nutrition Norfolk Food Frequency Questionnaire (FFQ; version 6, CAMB/PQ/6/1205) [6] with more than 130 food items. More information about the development of this FFQ has been briefly described in Peña-Fernández et al. [7,8].

Questionnaires were processed with Nutritics<sup>®</sup> software (v.5.7 Research Edition, Nutritics Ltd., Dublin, Ireland), as briefly described in Peña-Fernández et al. [8]. This nutritional software has been successfully used in similar studies previously [4,9]. Body mass index (BMI) was calculated as weight in kilograms divided by squared height in meters, using a digital scale (Tanita SC 330-S, London, UK), to identify underweight or obese individuals, depending on their ethnic background [10].

Statistical analyses were performed using the free software R-project, version 4.1.0 [11]. Significance scores were based on Kruskal–Wallis for nonparametric multiple comparisons and one-way analysis of variance for normal multiple comparisons. For normality, the Shapiro–Wilk test was used. Differences were considered statistically significant at *p*-values lower than 0.05.

## 3. Results and Discussion

The dietary intakes of total fat (100.55 vs. 81.72; *p*-value = 0.032), total polyunsaturated fatty acids (PUFA; 14.61 vs. 12.91; NS), linoleic acid (LA; 3.893 vs. 2.787; *p*-value = 0.0019),  $\alpha$ -linolenic acid (ALA; 0.925 vs. 0.613; *p*-value = 0.00008), arachidonic acid (AA; 0.109 vs. 0.082; *p*-value = 0.0303), eicosapentaenoic acid (EPA; 0.088 vs. 0.075; NS), docosahexaenoic acid (DHA; 0.153 vs. 0.121; NS), and docosapentaenoic acid (DPA; 0.043 vs. 0.032 all in g/day; NS) were significantly higher or higher in male participants, respectively.

Our results could be explained by higher intakes of food items rich in long-chain PUFAs [12,13]. Thus, male DMU students were shown to eat higher/significantly higher intakes of food products rich in long-chain PUFAs from aquatic [fish (72.656 vs. 53.907 g/day; *p*-value = 0.826) and seafood dishes (8.995 vs. 0.575 g/day; *p*-value = 0.149)], animals [meat (271.553 vs. 193.063 g/day; *p*-value = 0.016), eggs (17.625 vs. 16.998 g/day; *p*-value = 0.860), and cow milk (228.09 vs. 163.55; *p*-value = 0.037)], and plant sources [vegetable oils (1.094 vs. 0.955 g/day; *p*-value = 0.426)]. However, the consumption of oily fish (13.406 vs. 10.056 g/day; *p*-value = 0.857) and shellfish (12.842 vs. 10.826 g/day; *p*-value = 0.717) was higher in female counterparts, although without showing statistical differences, which in turn might explain that we have not observed differences in the intake of EPA and DHA in the monitored population.

The high intakes of LA and ALA versus EPA and DHA observed would be consistent with those reported in Western countries [13]. However, none of them presented statistical differences according to ethnic background or BMI, except for the intake of total fat ( $p$ -value = 0.0069) and PUFA ( $p$ -value = 0.0013), which were significantly higher in students from Asia and Europe. However, although without significance, the higher intakes of EPA and DHA were seen in Asian [values reported as mean and range (in g/day); 0.086 (0.005–0.498) and 0.144 (0.002–0.818)] and African [0.078 (0.0096–0.512) and 0.125 (0.011–0.723)] students versus European students [0.068 (0.007–0.251) and 0.118 (0.013–0.402)], possibly due to differences in the diet between these individuals based on their different traditions.

### 3.1. Estimated Daily Intakes of EPA + DHA According to Sex and Ethnic Background

The dietary intakes of DHA plus EPA in the whole group monitored (0.130 + 0.079 = 0.209 g/day) met the recommended daily intake (RDI) of 0.2 g/day recommended by the UK's Department of Health but were lower than the RDI of 0.5 g/day that considers the intake of one to two portions of fish per week [14]. The combined intake also did not meet the RDI of EPA + DHA of 0.45 g/day recommended by the Scientific Advisory Committee on Nutrition/Committee on Toxicity [14], and were below the 0.25–2 g/day established by the Food and Agriculture Organization of the United Nations/World Health Organization (FAO/WHO) in 2008 [13].

However, the higher intakes of EPA + DHA in the monitored group than those reported in the 19–24-year-old UK population (97 and 98 mg/day, in males and females, respectively) could be explained by the higher intake of fish observed (59.4) when compared with that reported in the general UK adult population (31 g/day) [15]. Moreover, the intake of total fish was higher in Asian and European students than that reported in African participants (65.895, 63.899 vs. 49.972 g/day), although these differences were not statistically different. Further analysis will evaluate the potential contribution of poultry, milk, and eggs to the intake of EPA and DHA in the monitored group for each ethnic background.

Finally, male and female participants (0.925 vs. 0.613 g/day, respectively) would not cover the adequate intake of ALA (1.6 and 1.1 g/day) established by the US National Institutes of Health [16].

### 3.2. Nutritional Recommendations

In order to meet the RDI of 0.5 g EPA + DHA per day, DMU students should specifically enhance the intake of oily fish, especially male counterparts (12.422, 13.406, and 10.054 g/day for the overall, female, and male population, respectively), as they would provide around 0.228, 0.246, and 0.184 g/day of these two long-chain PUFAs, respectively. These intakes are calculated if we consider the estimation reported by Givens and Gibbs [14], which suggests that 50 g of oily fish per week can provide around 131 mg/day of EPA + DHA. Considering the characteristics of the population monitored, education would be needed to increase awareness of the importance of consuming more fish. This intervention would be more significant if introduced at younger ages in the British educational system, as reported previously [14].

The intake of oily fish ( $r = 0.526$ ;  $p$ -value < 0.001), shellfish ( $r = 0.574$ ;  $p$ -value < 0.001), meat ( $r = 0.459$ ;  $p$ -value < 0.001), fruit ( $r = 0.414$ ;  $p$ -value < 0.001), and grains ( $r = 0.555$ ;  $p$ -value < 0.001) showed a significant and positive correlation with the intake of EPA in the monitored group, while the intake of oily fish ( $r = 0.473$ ;  $p$ -value < 0.001), fish products ( $r = 0.387$ ;  $p$ -value < 0.001), low-fat fish ( $r = 0.414$ ;  $p$ -value < 0.001), shellfish ( $r = 0.450$ ;  $p$ -value < 0.001), and meat ( $r = 0.389$ ;  $p$ -value < 0.001), were positively correlated with the intake of DHA. As a result, an enhancement in the intake of these food products will also contribute to meeting the requirements indicated.

Another option would be to encourage the intake of dietary fish oil supplements in the studied group, although this may be challenging for students that are from low-income backgrounds. Similarly, the enrichment of food items largely consumed by young

British adults with long-chain PUFAs could be another option to ensure that these RDIs are met [14].

#### 4. Conclusions

Our results would highlight that some DMU students monitored did not meet the nutritional goals for ALA, EPA, and DHA. Therefore, it would be advisable to increase the consumption of fish (especially oily fish) and/or foods enriched with these long-chain PUFAs among these university students. Implementation of nutritional supplementation policies for all these components could tackle the low intake observed and aid with the recommended allowances for individuals at risk.

**Author Contributions:** Conceptualisation, A.P.-F.; methodology, A.P.-F., E.S., M.d.l.Á.P., M.D.E. and M.H.; validation, A.P.-F.; formal analysis, A.P.-F., E.S. and M.H.; investigation, A.P.-F., E.S., M.d.l.Á.P., M.D.E. and M.H.; resources, A.P.-F., E.S., M.d.l.Á.P., M.D.E. and M.H.; data curation, A.P.-F., E.S. and M.H.; writing—original draft preparation, A.P.-F.; writing—review and editing, A.P.-F., E.S., M.d.l.Á.P., M.D.E. and M.H.; visualisation, A.P.-F., E.S., M.d.l.Á.P., M.D.E. and M.H.; supervision, A.P.-F.; project administration, A.P.-F. and M.d.l.Á.P.; internal funding acquisition, A.P.-F., M.d.l.Á.P. and M.D.E. All authors have read and agreed to the published version of the manuscript.

**Funding:** This project was funded by the Leicester School of Allied Health Sciences, De Montfort University.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki, and approved by the DMU Research Ethics Committee (Ref. 1674; 11 January 2016), subsequently amended and approved in 2017.

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author. The data are not publicly available due to further processing for a future submission as a manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

1. Tocher, D.R.; Betancor, M.B.; Sprague, M.; Olsen, R.E.; Napier, J.A. Omega-3 long-chain polyunsaturated fatty acids, EPA and DHA: Bridging the gap between supply and demand. *Nutrients* **2019**, *11*, 89. [[CrossRef](#)] [[PubMed](#)]
2. Zárate, R.; el Jaber-Vazdekis, N.; Tejera, N.; Pérez, J.A.; Rodríguez, C. Significance of long chain polyunsaturated fatty acids in human health. *Clin. Transl. Med.* **2017**, *6*, 25. [[CrossRef](#)] [[PubMed](#)]
3. Saini, R.K.; Keum, Y.S. Omega-3 and omega-6 polyunsaturated fatty acids: Dietary sources, metabolism, and significance—A review. *Life Sci.* **2018**, *203*, 255–267. [[CrossRef](#)] [[PubMed](#)]
4. Conway, M.C.; McSorley, E.M.; Mulhern, M.S.; Spence, T.; Wijngaarden, E.V.; Watson, G.E.; Yeates, A.J. The influence of fish consumption on serum n-3 polyunsaturated fatty acid (PUFA) concentrations in women of childbearing age: A randomised controlled trial (the iFish Study). *Eur. J. Nutr.* **2021**, *60*, 1415–1427. [[CrossRef](#)] [[PubMed](#)]
5. Mariamenatu, A.H.; Abdu, E.M. Overconsumption of omega-6 polyunsaturated fatty acids (PUFAs) versus deficiency of omega-3 PUFAs in modern-day diets: The disturbing factor for their “balanced antagonistic metabolic functions” in the human body. *J. Lipids* **2021**, *2021*, 8848161. [[CrossRef](#)] [[PubMed](#)]
6. Bingham, S.; Welch, A.; McTaggart, A.; Mulligan, A.; Runswick, S.; Luben, R.; Day, N. Nutritional methods in the European prospective investigation of cancer in Norfolk. *Public Health Nutr.* **2001**, *4*, 847–858. [[CrossRef](#)] [[PubMed](#)]
7. Peña-Fernández, A.; Ali, N.; Millington, D.; Lobo-Bedmar, M.C.; Haris, P. Human biomonitoring research at De Montfort University: School and university participants’ recruitment experience. In Proceedings of the INTED2018 Proceedings, Valencia Spain, 5–7 March 2018; pp. 6258–6262.
8. Peña-Fernández, A.; Higuera, M.; Segura, E.; Evans, M.D.; Peña, M.A. Inadequacies of iron dietary intake of in normal- and over-weight young university students from Leicester, England. *Biol. Life Sci. Forum*, 2023; *submitted for publication*.
9. Kutbi, H.A. Picky Eating in School-Aged Children: Sociodemographic determinants and the associations with dietary intake. *Nutrients* **2021**, *13*, 2518. [[CrossRef](#)] [[PubMed](#)]
10. Centre for Public Health Excellence at NICE (UK); National Collaborating Centre for Primary Care (UK). *Obesity: The Prevention, Identification, Assessment and Management of Overweight and Obesity in Adults and Children*; National Institute for Health and Clinical Excellence (UK): London, UK, 2006.

11. Core Team R. *A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2021. Available online: <https://www.R-project.org/> (accessed on 11 September 2023).
12. Abedi, E.; Sahari, M.A. Long-chain polyunsaturated fatty acid sources and evaluation of their nutritional and functional properties. *Food Sci. Nutr.* **2014**, *2*, 443–463. [[CrossRef](#)] [[PubMed](#)]
13. Redruello-Requejo, M.; Samaniego-Vaesken, M.d.L.; Puga, A.M.; Montero-Bravo, A.; Ruperto, M.; Rodríguez-Alonso, P.; Partearroyo, T.; Varela-Moreiras, G. Omega-3 and omega-6 polyunsaturated fatty acid intakes, determinants and dietary sources in the spanish population: Findings from the ANIBES study. *Nutrients* **2023**, *15*, 562. [[CrossRef](#)] [[PubMed](#)]
14. Givens, D.I.; Gibbs, R.A. Current intakes of EPA and DHA in European populations and the potential of animal-derived foods to increase them: Symposium on 'How can the n-3 content of the diet be improved?'. *Proc. Nutr. Soc.* **2008**, *67*, 273–280. [[CrossRef](#)] [[PubMed](#)]
15. Gibbs, R.A.; Givens, D.I.; Rymer, C. The effect of age and gender on current and potential intakes of very long chain n-3 polyunsaturated fatty acids from oil-rich fish and animal derived foods in the UK. In Proceedings of the British Society of Animal Science, Southport, UK, 2–4 April 2007; Cambridge University Press: Cambridge, UK, 2007; p. 9.
16. US National Institutes of Health. Omega-3 Fatty Acids. 2023. Available online: <https://ods.od.nih.gov/factsheets/Omega3FattyAcids-HealthProfessional/> (accessed on 11 September 2023).

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.