



Rat Strain-Specific Differences in Alcohol Intake Following Patterned Feeding of a Palatable Diet [†]

Brooke White , Sabrina Pham, John Michael Houeye, Kaiyah Rush and Sunil Sirohi *

Laboratory of Endocrine and Neuropsychiatric Disorders, Division of Basic Pharmaceutical Sciences, College of Pharmacy, Xavier University of Louisiana, 1 Drexel Dr, New Orleans, LA 70125, USA; bwhite17@xula.edu (B.W.); spham6@xula.edu (S.P.); jhoueye@xula.edu (J.M.H.); krush2@xula.edu (K.R.)

* Correspondence: ssirohi@xula.edu; Tel.: +1-504-520-5471

[†] Presented at the 3rd International Electronic Conference on Nutrients, 1–15 November 2023; Available online: <https://iecn2023.sciforum.net/>.

Abstract: A total of 29.5 million people aged 12 and older met the diagnostic criteria for alcohol use disorder (AUD) in the United States in 2021, which presents a significant social and economic burden to modern society. Impaired nutritional status has been frequently documented in patients with AUDs and could contribute to escalated alcohol consumption and behavioral impairments commonly observed in AUD. Interestingly, increased highly palatable food intake during recovery has been reported in patients with AUD, suggesting the importance of understanding the relationship between palatable food and problematic alcohol drinking. We have previously shown that patterned feeding of a palatable diet attenuated alcohol drinking in Long Evans rats. The present study evaluated the impact of patterned feeding on high and low alcohol drinking. Individually housed male high-drinking (P), moderate-drinking (Long Evans), and low-drinking (Wistar) rats received intermittent access (24 h, Tuesdays, and Thursdays) to a nutritionally complete high-fat diet (Int-HFD) or standard chow (controls). Normal chow and water were available ad libitum to all groups of rats. Intermittent HFD access induced a feeding pattern in which the Int-HFD group of rats escalated their caloric intake on Tuesdays and Thursdays. Two weeks of Int-HFD pre-exposure preceded any alcohol access, after which all rats were given unsweetened alcohol (20% v/v) in their home cages via a two-bottle choice paradigm of voluntary alcohol drinking. Alcohol was available for 24 h on chow-only days (Mondays, Wednesdays, and Fridays) while Int-HFD feeding continued. Long Evans rats receiving the Int-HFD displayed a significant ~50% reduction in alcohol drinking when compared to controls. The Int-HFD group of P rats also reduced their alcohol intake significantly ($p < 0.05$) by ~20%, in comparison to respective controls. Interestingly, alcohol drinking in Wistar rats was not affected ($p > 0.05$) by intermittent HFD exposure. These data highlight rat strain-specific differences in alcohol intake following patterned feeding of a palatable diet and identify Long Evans rats as an ideal model to evaluate the impact of palatable diet on alcohol drinking.

Keywords: Wistar; Long Evans; P rat; alcohol use disorder; alcohol drinking; high-fat diet; palatable diet



Citation: White, B.; Pham, S.; Houeye, J.M.; Rush, K.; Sirohi, S. Rat Strain-Specific Differences in Alcohol Intake Following Patterned Feeding of a Palatable Diet. *Biol. Life Sci. Forum* **2023**, *29*, 24. <https://doi.org/10.3390/IECN2023-15821>

Academic Editor: Paula Chen

Published: 3 November 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Alcohol use disorder (AUD) is a debilitating disorder which significantly impacts an individual's health and ability to function and has extensive economic impacts. An estimated 140,000 people die annually of alcohol-related causes [1] and the life expectancy of someone with AUD has been shown to be reduced by as much as 28 years when compared to healthy individuals [2]. Additionally, when measured in disability-adjusted life-years, alcohol misuse contributes significantly to years of life lost due to improper health or disability [3]. In 2010, the cost of alcohol misuse in the United States (US) totaled \$249 billion and \$191.1 billion of this financial burden was attributed to binge drinking [4].

Impaired nutritional status is frequently reported in patients with AUD along with emotional and physiological abnormalities. The cause of nutritional deficiency in this

population is multifaceted; it is impacted by reduced nutritional intake, altered nutrient absorption, and changes in nutrient metabolism [5–7]. Particularly, deficiencies in vitamins and minerals have been widely reported in people with AUD [8]. For example, vitamin B12 and C levels are negatively impacted by excessive alcohol consumption and are associated with cognitive dysfunction [9,10]. Vitamin D is another essential nutrient affected by chronic alcohol intake and these deficiencies have been implicated in increased negative affect [11]. Importantly, unlike other substances of abuse, alcohol contains calories and when the calories from alcohol replace those typically obtained from a healthy diet, nutritional status is negatively affected. In individuals with severe AUD more than 30% of daily caloric intake could be derived from alcohol alone, which negatively affects dietary carbohydrate, fat, and protein intake [12]. Together, a compromised nutritional status as a result of chronic alcohol consumption could impair health and could contribute to AUD and related pathologies.

Interestingly, increased preference for palatable diets (PDs), or food high in sugar [13–15] and carbohydrates [14], have been observed in people with AUD during recovery, suggesting potentially protective effects of increased PD intake in people with AUD [13–15]. Furthermore, Alcoholics Anonymous also suggest sweet-tasting foods consumption to curb alcohol cravings [16]. Several preclinical studies from our laboratory and others have also evaluated the impact of PD consumption on alcohol drinking [17–19]. For example, our lab has previously demonstrated a reduced alcohol deprivation effect following intermittent high-fat diet (Int-HFD) access [17] and attenuated alcohol drinking following two weeks of Int-HFD pre-exposure [18] in Long Evans rats. The objective of the present study was to compare the impact of two weeks of Int-HFD pre-exposure on subsequent alcohol drinking in lower-drinking Wistar rats, moderately drinking Long Evans rats, and higher-drinking P rats.

2. Materials and Methods

2.1. Animals

Male Wistar (RccHan[®]: WIST, Envigo RMS, Inc., Indianapolis, IN, USA), Long Evans (HsdBlue: LE, Envigo RMS, Inc., Indianapolis, IN, USA) and alcohol-preferring (P) rats (Indiana University) were used. The vivarium was controlled for temperature (~70F) and humidity (~60%) with a 12 h reverse light-dark cycle (lights on from 1:00 a.m. to 1:00 p.m.). On arrival, animals were handled before any experimental manipulation or baseline data (body weight, food intake and water intake) were collected.

2.2. Diet and Alcohol

All animals had ad libitum access to standard rodent chow (Tekland–Envigo Diets #2020X, 3.1 kcal/g with 16% calories from fat and 60% calories from carbohydrates) and tap water. The experimental group was given intermittent access to a high-fat diet (HFD; Research Diets #D03082706, 4.5 kcal/g with 40% calories from fat, 46% calories from carbohydrates, and 7.9% calories from sugar) in addition to standard chow. The 190 proof alcohol was purchased from Greenfield Global, MI and the desired 20% *v/v* was prepared at least one day in advance every week. Voluntary alcohol consumption was measured in home cages with a two-bottle choice paradigm and the position of the alcohol and water bottles were switched daily to minimize conditioning effects. Food, alcohol, and water were provided ~2 h into the dark cycle, and intake was measured manually after 24 h of access.

2.3. Procedure

Male rats ($n = 6$) matched for body weight, food intake, and water intake were randomly divided into control and Int-HFD groups. To evaluate the effect of Int-HFD feeding on subsequent alcohol intake, Int-HFD rats were given 24 h, intermittent (Tuesday and Thursday) access to the HFD for two weeks prior to alcohol exposure (see Figure 1A). Following Int-HFD pre-exposure, alcohol consumption was measured on chow-only days (Monday, Wednesday and Friday) while intermittent HFD feeding continued (see Figure 1B).




Figure 1. Schematic representation of patterned feeding and alcohol access schedules. Green boxes represent high-fat diet (HFD) access, yellow boxes represent chow access, and the grey bottles represent alcohol access. **(A)** Intermittent high-fat diet (Int-HFD) rats were given 2 weeks of intermittent 24 h HFD access (Tuesday and Thursday) while controls received additional chow. **(B)** Following HFD pre-exposure, rats received alcohol access on chow-only days (Monday, Wednesday, Friday) simultaneous to Int-HFD feeding.

3. Results and Discussion

Consistent with our previously published data, all groups of Int-HFD rats displayed caloric overconsumption on HFD access days. Long Evans rats displayed a gradual increase in alcohol drinking over time, an effect significantly (50%) reduced in the Int-HFD group of rats. When evaluated under identical conditions, alcohol intake in the Int-HFD P rats was also significantly (~20%) reduced compared to the chow controls, whereas no effect of Int-HFD on alcohol intake was evident in the Wistar rats. Together, intermittent access to a HFD differentially impacted alcohol intake in low-, moderate-, and high-drinking rats with greater effects seen in moderate drinking Long Evans rats. The present study emphasizes strain-specific differences in the effect of an intermittent HFD on subsequent alcohol intake.

It is important to note that several critical factors could impact alcohol drinking following dietary manipulations. For example, significant differences in the behavioral effects of a high-sugar diet and a high-fat diet have been previously identified [20], emphasizing dietary content as a factor requiring further evaluation. Furthermore, the length of PD exposure may have an impact on alcohol drinking. In the present study, all the experimental parameters were kept constant between rat strains and there was no significant change in body weight observed in any Int-HFD rats when compared to respective controls throughout the experiment. Interestingly, a previous study reported attenuated alcohol intake in Wistar rats following 3–4 weeks of junk-food diet (averaging 42% fat, 52% carbohydrates and 6% protein) access. In contrast to the present study, junk-food diet feeding induced an obesity phenotype [19], emphasizing the potential impact of the PD access periods duration.

In alignment with previous published reports, Wistar rats in the current study displayed a low initiation of alcohol drinking and minimal escalation of intake over time. On the other hand, P rats displayed a high initiation of alcohol drinking, maintained throughout the testing period. Interestingly, previous studies comparing blood alcohol levels (BALs) during an intermittent alcohol access paradigm reported higher BALs in Long Evans rats when compared to Wistar and P rats. Even with lower alcohol intake, the amount of ethanol consumption required by P rats and Wistar rats to reach the same pharmacologically relevant blood ethanol content (BEC) was greater than that required by Long Evans rats [21–23]. We have previously reported the involvement of central mechanisms in mediating the effects of an intermittent HFD on alcohol drinking, as selective alterations in the neurotransmitter receptor expression in the brain's reward circuitry were

observed in Int-HFD Long Evans rats when compared to chow controls [18]. Therefore, the low alcohol drinking levels and likely lower BALs of the Wistar rats in the present study could explain the lack of intermittent HFD access effects on alcohol drinking, a contention needing further evaluation.

4. Conclusions

In conclusion, there was no observed effect of intermittent HFD access on alcohol intake in the low-drinking Wistar rats. Int-HFD P rats displayed significantly attenuated alcohol intake (20%) when compared to the chow controls; however, the effect was smaller than that observed in Int-HFD Long Evans rats (50%). These results emphasize rat strain-specific differences in the effect of an Int-HFD on subsequent alcohol intake and warrant future investigation. Nevertheless, this study identifies Long Evans rats as a potentially ideal model for evaluating central mechanisms of diet-induced effects on alcohol intake.

Author Contributions: Conceptualization, S.S.; Methodology, S.S.; Software, S.S.; Validation, S.S.; Formal Analysis, S.S.; Investigation, B.W., J.M.H., K.R. and S.S.; Resources, S.S.; Data curation, S.S.; Writing—original draft, B.W. and S.P.; Writing—review and editing, S.S.; Visualization, S.S.; Supervision, S.S.; Project administration, S.S.; Funding acquisition, S.S. All authors have read and agreed to the published version of the manuscript.

Funding: Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health, project number 5SC3GM127173-04 to S.S. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Institutional Review Board Statement: The animal study protocol was approved by the Institutional Animal Care and Use Committee at the Xavier University of Louisiana on 25 October 2021, with protocol number 100722-003DBPS.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data that support the findings of this study are available from the corresponding author, S.S. upon reasonable request.

Acknowledgments: We would like to thank Sean Graves for his technical support.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Centers for Disease Control and Prevention. Alcohol Related Disease Impact (ARDI) Application Website. 2022. Available online: www.cdc.gov/ARDI (accessed on 6 October 2023).
2. Westman, J.; Wahlbeck, K.; Laursen, T.M.; Gissler, M.; Nordentoft, M.; Hällgren, J.; Arffman, M.; Ösby, U. Mortality and life expectancy of people with alcohol use disorder in Denmark, Finland and Sweden. *Acta Psychiatr. Scand.* **2015**, *131*, 297–306. [[CrossRef](#)] [[PubMed](#)]
3. GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: A systematic analysis for the Global Burden of Disease Study 2019. *Lancet* **2020**, *396*, 1223–1249. [[CrossRef](#)] [[PubMed](#)]
4. Sacks, J.J.; Gonzales, K.R.; Bouchery, E.E.; Tomedi, L.E.; Brewer, R.D. 2010 National and State Costs of Excessive Alcohol Consumption. *Am. J. Prev. Med.* **2015**, *49*, e73–e79. [[CrossRef](#)] [[PubMed](#)]
5. Glória, L.; Cravo, M.; Camilo, M.E.; Resende, M.; Cardoso, J.N.; Oliveira, A.G.; Leitão, C.N.; Mira, F.C. Nutritional deficiencies in chronic alcoholics: Relation to dietary intake and alcohol consumption. *Am. J. Gastroenterol.* **1997**, *92*, 485–489. [[PubMed](#)]
6. Green, P.H. Alcohol, nutrition and malabsorption. *Clin. Gastroenterol.* **1983**, *12*, 563–574. [[CrossRef](#)] [[PubMed](#)]
7. Maillot, F.; Farad, S.; Lamisse, F. Alcool et nutrition [Alcohol and nutrition]. *Pathol.-Biol.* **2001**, *49*, 683–688. [[CrossRef](#)] [[PubMed](#)]
8. Lieber, C.S. Relationships between nutrition, alcohol use, and liver disease. *Alcohol Res. Health J. Natl. Inst. Alcohol Abus. Alcohol.* **2003**, *27*, 220–231.
9. Vedder, L.C.; Hall, J.M.; Jabrouin, K.R.; Savage, L.M. Interactions between chronic ethanol consumption and thiamine deficiency on neural plasticity, spatial memory, and cognitive flexibility. *Alcohol. Clin. Exp. Res.* **2015**, *39*, 2143–2153. [[CrossRef](#)]
10. Clergue-Duval, V.; Azuar, J.; Fonsart, J.; Delage, C.; Rollet, D.; Amami, J.; Frapsauce, A.; Gautron, M.A.; Hispard, E.; Bellivier, F.; et al. Ascorbic Acid Deficiency Prevalence and Associated Cognitive Impairment in Alcohol Detoxification Inpatients: A Pilot Study. *Antioxidants* **2021**, *10*, 1892. [[CrossRef](#)]

11. Neupane, S.P.; Lien, L.; Hilberg, T.; Bramness, J.G. Vitamin D deficiency in alcohol-use disorders and its relationship to comorbid major depression: A cross-sectional study of inpatients in Nepal. *Drug Alcohol Depend.* **2013**, *133*, 480–485. [[CrossRef](#)]
12. Gruchow, H.W.; Sobocinski, K.A.; Barboriak, J.J.; Scheller, J.G. Alcohol consumption, nutrient intake and relative body weight among US adults. *Am. J. Clin. Nutr.* **1985**, *42*, 289–295. [[CrossRef](#)] [[PubMed](#)]
13. Stickel, A.; Rohdemann, M.; Landes, T.; Engel, K.; Banas, R.; Heinz, A.; Müller, C.A. Changes in Nutrition-Related Behaviors in Alcohol-Dependent Patients After Outpatient Detoxification: The Role of Chocolate. *Subst. Use Misuse* **2016**, *51*, 545–552. [[CrossRef](#)] [[PubMed](#)]
14. Yung, L.; Gordis, E.; Holt, J. Dietary choices and likelihood of abstinence among alcoholic patients in an outpatient clinic. *Drug Alcohol Depend.* **1983**, *12*, 355–362. [[CrossRef](#)]
15. Braun, T.D.; Kunicki, Z.J.; Blevins, C.E.; Stein, M.D.; Marsh, E.; Feltus, S.; Miranda, R., Jr.; Thomas, J.G.; Abrantes, A.M. Prospective Associations between Attitudes toward Sweet Foods, Sugar Consumption, and Cravings for Alcohol and Sweets in Early Recovery from Alcohol Use Disorders. *Alcohol. Treat. Q.* **2021**, *39*, 269–281. [[CrossRef](#)] [[PubMed](#)]
16. Alcoholics Anonymous. *Living Sober*; Alcoholics Anonymous World Services, Inc.: New York, NY, USA, 2007; pp. 18–19.
17. Shah, K.; Shaw, C.; Sirohi, S. Reduced alcohol drinking following patterned feeding: Role of palatability and acute contingent availability. *Physiol. Behav.* **2020**, *224*, 113020. [[CrossRef](#)] [[PubMed](#)]
18. Villavasso, S.; Shaw, C.; Skripnikova, E.; Shah, K.; Davis, J.F.; Sirohi, S. Nutritional Contingency Reduces Alcohol Drinking by Altering Central Neurotransmitter Receptor Gene Expression in Rats. *Nutrients* **2019**, *11*, 2731. [[CrossRef](#)]
19. Cook, J.B.; Hendrickson, L.M.; Garwood, G.M.; Toungate, K.M.; Nania, C.V.; Morikawa, H. Junk food diet-induced obesity increases D2 receptor autoinhibition in the ventral tegmental area and reduces ethanol drinking. *PLoS ONE* **2017**, *12*, e0183685. [[CrossRef](#)]
20. Avena, N.M.; Rada, P.; Hoebel, B.G. Sugar and fat bingeing have notable differences in addictive-like behavior. *J. Nutr.* **2009**, *139*, 623–628. [[CrossRef](#)]
21. Carnicella, S.; Ron, D.; Barak, S. Intermittent ethanol access schedule in rats as a preclinical model of alcohol abuse. *Alcohol* **2014**, *48*, 243–252. [[CrossRef](#)]
22. Bell, R.L.; Hauser, S.R.; Liang, T.; Sari, Y.; Maldonado-Devincini, A.; Rodd, Z.A. Rat animal models for screening medications to treat alcohol use disorders. *Neuropharmacology* **2017**, *122*, 201–243. [[CrossRef](#)]
23. Simms, J.A.; Steensland, P.; Medina, B.; Abernathy, K.E.; Chandler, L.J.; Wise, R.; Bartlett, S.E. Intermittent access to 20% ethanol induces high ethanol consumption in Long-Evans and Wistar rats. *Alcohol. Clin. Exp. Res.* **2008**, *32*, 1816–1823. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.