

Agroforestry as a Climate-Smart Strategy: Examining the Factors Affecting Farmers' Adoption [†]

Md. Manik Ali ^{1,*}, Abinash Chandra Pal ¹, Md. Shafiqul Bari ¹, Md. Lutfor Rahman ² and Israt Jahan Sarmin ¹

¹ Department of Agroforestry and Environment, Hajee Mohammad Danesh Science and Technology University, Dinajpur 5200, Bangladesh; btbh.abinash135@gmail.com (A.C.P.); barimshafiqul@gmail.com (M.S.B.); israthstu92@gmail.com (I.J.S.)

² Department of Agricultural Chemistry, Hajee Mohammad Danesh Science and Technology University, Dinajpur 5200, Bangladesh; mdlutfarhstu@gmail.com

* Correspondence: manikali@hstu.ac.bd

[†] Presented at the 2nd International Online Conference on Agriculture, 1–15 November 2023; Available online: <https://iocag2023.sciforum.net/>.

Abstract: Agroforestry production systems have shown growing adoption in Bangladesh, offering ecological and economic benefits in the face of climate change. This study investigates the scale of agroforestry adoption, investment returns, factors influencing uptake, and challenges faced by farmers. Using a multistage random sample of 340 respondents, we find that while 75% of farmers are aware of agroforestry, adoption remains limited. Our analysis focuses on specific tree–crop combinations favored by farmers as agroforestry practices. The results demonstrate that, in cropland agroforestry, Eucalyptus tree with rice (69.05% adoption rate) is predominant, while home-stead/orchard system agroforestry favors mango tree intercropped with potato (73.33%). Financial and investment analyses using Benefit–Cost Ratio (BCR), Net Present Value (NPV), and Internal Rate of Return (IRR) prove that agroforestry is a more favorable alternative for farmers considering adoption, as it provides superior BCR, NPV, and IRR. For example, litchi-based agroforestry systems with vegetables like brinjal (eggplant), potato, and chilies offer higher NPVs (19.00, 19.73, and 18.46, respectively) and IRRs (54.45, 68.00, and 47.19, respectively) compared to monocropping where NPV was 14.38. A binary logistic model reveals that larger farm sizes, younger respondents, higher education levels, training experiences, more frequent extension visits, and improved market access positively influence agroforestry adoption. The study also identifies key challenges for farmers using the Problem Facing Index (PFI). The most significant obstacles include lack of training facilities (PFI-894), shortage of skilled labor (PFI-687), and insufficient technical expertise (PFI-647). Therefore, to promote wider adoption, targeted training programs that address the identified challenges are crucial. It will empower farmers to reap the tangible benefits of agroforestry as a sustainable and climate-smart agricultural practice.

Keywords: agroforestry; CSA; adoption; farmer's income; investment analysis



Citation: Ali, M.M.; Pal, A.C.; Bari, M.S.; Rahman, M.L.; Sarmin, I.J. Agroforestry as a Climate-Smart Strategy: Examining the Factors Affecting Farmers' Adoption. *Biol. Life Sci. Forum* **2024**, *30*, 29. <https://doi.org/10.3390/IOCAG2023-17340>

Academic Editor: Amanullah

Published: 18 April 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Climate change poses a formidable challenge to global agricultural systems, threatening food security, livelihoods, and the overall sustainability of farming practices [1]. As the world confronts the challenges posed by a changing climate, the need for innovative and sustainable solutions is imperative [2,3]. Among the array of climate-smart strategies, agroforestry stands out as a promising approach that integrates trees and shrubs into agricultural landscapes to simultaneously mitigate and adapt to climate change [4]. Agroforestry is recognized as an integrated land-use management approach that has gained attention for its multifaceted benefits [5]. As an integrated land-use management approach, agroforestry has gained attention for its potential to mitigate and adapt to climate change

while promoting sustainable agricultural practices [6,7]. The synergistic relationship between trees, crops, and livestock in agroforestry systems contributes to enhanced resilience, carbon sequestration, soil fertility, and water conservation [8]. These multifunctional landscapes have been recognized for their ability to provide a more robust and diversified livelihood for farmers, reducing vulnerability to climate-induced shocks [9]. While the benefits of agroforestry are evident, the adoption of such practices by farmers is influenced by a complex interplay of factors. These factors range from socio-economic and cultural considerations to institutional support, knowledge dissemination, and policy frameworks [10,11]. Understanding these factors is essential for designing targeted interventions that encourage widespread adoption of agroforestry, thereby harnessing its full potential as a climate-smart strategy. Research in this field has indicated that the adoption of agroforestry practices varies significantly across different regions and farming communities. Factors such as land tenure systems, access to resources, market dynamics, and farmers' perceptions play a crucial role in shaping adoption patterns. Additionally, the effectiveness of extension services, agroforestry training programs, and policy incentives can either facilitate or hinder the uptake of these practices. For instance, studies emphasize the potential of agroforestry as a low-hanging fruit in climate change mitigation [12]. The synergies between agroforestry and climate resilience have been explored, highlighting the robust approach of evergreen agriculture in promoting food security in Africa [13]. These studies provide foundational insights into the benefits and potential of agroforestry, setting the stage for more nuanced examinations of the factors influencing adoption. In this context, this study seeks to delve into the nuanced factors affecting farmers' adoption of agroforestry as a climate-smart strategy. Through a comprehensive approach that includes synthesizing the existing literature, conducting field surveys, and analyzing case studies, the research aims to contribute valuable insights into the intricacies of agroforestry adoption. This knowledge is pivotal for developing context-specific strategies, policies, and extension programs that promote the widespread adoption of agroforestry, fostering sustainable and climate-resilient agricultural systems.

2. Methodologies

2.1. Study Area

The selection of the study area was predicated on the prevalence of agroforestry practices, focusing on the implementation in the Baliadangi Upazila of the Thakurgaon District within the Rangpur Division of Bangladesh. Encompassing an area of 284.12 square kilometers, Baliadangi Upazila is situated between 25°59' and 26°12' north latitudes and 88°10' and 88°22' east longitudes.

2.2. Materials and Methods

In this study, a meticulously designed survey questionnaire was employed to explore farmers' perceptions, attitudes, and adoption behaviors concerning agroforestry practices. This survey will encompass various dimensions, including socio-economic characteristics, land tenure dynamics, resource accessibility, awareness of agroforestry benefits, and factors influencing adoption decisions. A multistage random sampling procedure was employed to collect data. A sample size of 340 households was established using Cochran's formula [14], taking into account a 95% confidence interval and a 5% margin of error. The data collection phase will target a diverse sample of farmers through in-person interviews utilizing both closed-ended questions for quantitative rigor and open-ended questions for qualitative depth. Data quality was enhanced through focus group discussions (FGDs) and key informant interviews, and supplementary data were collected from authoritative sources. Data analysis was performed utilizing the statistical software STATA-14.00. Descriptive statistics and logistic regression analysis (logit model) were employed to discern the factors influencing agroforestry adoption, encompassing socio-economic variables, resource accessibility, and awareness of agroforestry practices. In the context of investment analysis, key financial metrics such as Discounted Net Present Value (NPV), Benefit–Cost

Ratio (BCR), and Internal Rate of Return (IRR) were employed. The Problem Confrontation Index (PCI) was calculated by multiplying the weighted sum of the problems' responses. Each farmer was asked to rate the complexity of each challenge by selecting one of the five options: "Very High," "High," "Medium," "Low," or "Not at all." These replies were given weights of 4, 3, 2, 1, and 0 accordingly [15].

3. Results

3.1. Demographic Profile of the Respondents in the Study Area

The study reveals significant insights into the demographic and economic landscapes of the respondents. The data in Table 1 indicate that the majority of participants are relatively young, with an average age of 42 years, and predominantly lack formal education, as reflected by an average educational status score near to seven. Marital status leans towards being married, suggesting family-centric households. The average household size is substantial at five. Land ownership is notably small, with most respondents relying on external sources for land. The financial situation is challenging, with a low average annual income of BDT 118,735.87, predominantly derived from low-wage agricultural work. Monthly expenditures, averaging around BDT 10,000, further underscore the economic constraints faced by the households. Overall, the findings paint a picture of a community grappling with limited resources, emphasizing the need for targeted interventions to address their economic challenges.

Table 1. Demographic profile of the respondents in the study area.

Variables	Mean	St. Err.	Min	Max
Adoption of agroforestry as a climate-smart strategy	0.582	0.64	0	1
Age of the respondent (years)	42	2.74	19	67
Farm size (hectares)	0.97	3.27	0.19	5.71
Family size (number)	5	2.17	2	7
Education level (years of schooling)	6.81	7.87	0	12
Marital status (binary variable)	0.87	0.31	0	1
Training experience (binary variable)	0.38	0.92	0	1
Number of extension visits (number of visits/month)	0.48	1.94	0	6
Improved market access (binary variable)	0.62	4.11	0	1
Income level (yearly, in BDT)	118,735.87	85,901.27	50,000	375,000
Farming experience (number of years)	12.91	5.48	3	60

Note: binary variable, if yes 1, otherwise 0.

3.2. Major Agroforestry Practices and Tree–Crop Combinations

The investigation into agroforestry practices in the study area reveals a nuanced landscape characterized by varying tree–crop combinations within cropland and homestead/orchard-based agroforestry systems. Notably, in cropland agroforestry, the preeminent practice involves the strategic pairing of Eucalyptus with rice, attaining a substantial adoption rate of 69.05% (Table 2). This is succeeded by Eucalyptus synergies with maize and wheat. Conversely, in homestead/orchard-based agroforestry, mango intercropped with potato emerges as the predominant choice, boasting an impressive adoption rate of 73.33%, underscoring its popularity among local farmers. Additional favored combinations include mango with red amaranth, litchi with red amaranth, and litchi with rice. These rankings not only underscore the diversity in tree–crop amalgamations but also shed light on farmers' discerning preferences, influenced by the intricate interplay of regional agroecological nuances and agricultural practices. This comprehensive insight derived from the table facilitates a deeper comprehension and advocacy for sustainable and diversified farming systems.

Table 2. Major agroforestry practices adopted by the farmers in the study area.

Crop Land Agroforestry			Homestead/Orchard-Based Agroforestry		
Tree–Crop Combination	Practiced by Farmers (%)	Rank Order	Tree–Crop Combination	Practiced by Farmers (%)	Rank Order
Eucalyptus + Maize	57.14	2	Mango + Potato	73.33	1
Eucalyptus + Rice	69.05	1	Mango + Bean	69.05	3
Eucalyptus + Wheat	52.38	3	Mango + Brinjal	64.29	5
Eucalyptus + Mustard	2.38	6	Mango + Onion + Garlic	52.38	7
Mahogany + Rice	42.86	4	Mango + Red Amaranth	73.81	2
Mahogany + Wheat	42.86	4	Mango + Radish	23.81	15
Mahogany + Maize	42.86	4	Mango + Pointed Gourd	28.57	13
Mahogany + Napier	2.38	6	Mango + Tomato	42.86	8
Akashmoni + Rice	2.38	6	Mango + Cauliflower	14.29	16
Mango + Rice	42.86	4	Litchi + Potato	66.67	4
Mango + Wheat	42.86	4	Litchi + Malabar Spinach	40.48	9
Litchi + Rice	52.38	3	Litchi + Onion	35.72	11
Litchi + Mustard	42.86	4	Litchi + Red Amaranth	57.14	6
Mango + Turmeric	9.52	5	Litchi + Sweet Gourd	38.10	10
Akashmoni + Maize	2.38	6	Malta + Potato	2.38	20
			Guava + Cucumber	7.14	19
			Lemon + Corolla	9.52	18

3.3. Investment Analysis of Different Agroforestry Combinations

Table 3 presents a detailed analysis of three different tree species cultivation frameworks: monocrop, agroforestry (agroforestry combined with vegetables), and important economic metrics including Net Present Value (NPV), Internal Rate of Return (IRR), and Benefit–Cost Ratio (BCR). Related crops such as potatoes, brinjal (eggplant), and chilies are included in the agroforestry model. In more detail, litchi performs better than monocrop options in the agroforestry paradigm, with an IRR of 28 and a greater NPV of 14.38. Notably, brinjal, potato, and chili had BCR values of 1.77, 2.00, and 1.46, respectively. Similarly, mango showed higher economic returns with agroforestry than as a monocrop, with an NPV of 18.36 and an IRR of 45. The BCR values obtained from the integration of potato, brinjal, and chili are 1.63, 2.22, and 1.91, respectively. Eucalyptus, Akashmoni, and Mahogany are similar in that they have higher NPV and IRR values in the agroforestry space, and they are paired with vegetable crops that have better BCRs.

Table 3. Investment analysis of the different popular combinations of the agroforestry system.

Tree	Monocrop		Agroforestry (Combined with Vegetables)								
	NPV ¹	IRR	Chili			Brinjal			Potato		
			BCR	NPV ¹	IRR	BCR	NPV ¹	IRR	BCR	NPV ¹	IRR
Litchi	14.38	28	1.77	18.46	47.19	2.00	19.00	54.45	1.46	19.73	68
Mango	18.36	45	1.91	21.18	72.6	2.22	22.40	83.49	1.63	23.00	88
Mahogany	10.95	25	1.46	15.50	50.82	1.41	13.46	33.88	1.21	16.38	57
Eucalyptus	14.38	28	1.69	17.78	43.56	1.96	18.54	48.4	1.46	19.73	68
Akashmoni	7.45	22	1.17	12.22	47.19	1.35	12.79	52.03	1.37	13.08	55

Note: NPV ¹ (Net Present Value at Year 1 in thousand USD).

3.4. Factors Affecting Farmer's Adoption of Agroforestry as a Climate-Smart Strategy

Table 4 shows the result of logit model effect of explanatory variables on the dependent variables. This result of this study exhibits that farm size ($p < 0.01$), age of the respondent ($p < 0.01$), education ($p < 0.05$), training ($p < 0.05$), number of extension visits ($p < 0.01$), and improved market access ($p < 0.10$) have positive significant influence on the adoption of agroforestry as climate smart strategy by the farmers, implying that an increase in these explanatory variables results in the positive increase in the adoption of agroforestry as climate smart strategy.

Table 4. Factors affecting farmer's adoption of agroforestry as a climate-smart strategy.

Variables	Coefficient	St. Err.	t-Value	Sig
Farm size	0.022	0.005	4.13	***
Age of the respondent	3.121	0.389	8.03	***
Education level	1.583	0.71	2.23	**
Training experience	0.004	0.002	2.15	**
Number of extension visits	0.771	0.167	4.63	***
Improved market access	0.401	0.208	1.93	*
Household size	-0.236	0.506	-0.47	
Income level	0.103	0.068	1.51	
Distance from the nearest market	-0.018	0.014	-1.22	
Constant	-2.489	0.902	-2.76	***
Mean of dependent variables	0.517	SD of dependent variables		0.501
Pseudo r-squared	0.395	Number of observations		294
Chi-square	160.944	Prob > chi2		0.000
Akaike criterion (AIC)	266.286	Bayesian criterion (BIC)		303.122

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

3.5. Major Problem Faced by the Farmer for Adopting Agroforestry

The study comprehensively examines challenges faced by farmers in the study area, providing insights into agroforestry complexities. A detailed figure categorizes issues impacting labor dynamics, productivity, environment, and infrastructure, contributing to the Problem Confrontation Index (PCI) (Figure 1). The top concerns are the lack of training facilities (1st, PCI-894) and a shortage of skilled labor (2nd, PCI-687). Insufficient expertise (3rd, PCI-647) and pest-related issues (4th, PCI-625) are also significant. Land availability, marketing infrastructure, and quality inputs follow in PCI rankings. Farmers express concerns about allelopathy, land damage from spreading roots, and complex production worries. Issues like trees falling on crops, limited access to high-quality seedlings, and theft rank are of lesser significance.




Figure 1. Graph showing different problems faced by the farmers for agroforestry adoption. Notes: 1= less productive than a monoculture; 2 = insects and pests are harbored; 3 = allelopathic impact; 4 = land damaged by quickly spreading roots; 5 = trees falling on crops; 6 = absence of skilled labor; 7 = inadequate facilities' training; 8 = lack of land availability; 9 = issues with thieves; 10 = absence of marketing infrastructure; 11 = insufficient expertise and technical assistance; 12 = lack of access to high-quality seedlings; and 13 = absence of quality fungicides, insecticides, and fertilizers.

4. Discussion

The agroforestry practices identified in the study underscore the diverse combinations of tree–crop pairings within cropland and homestead-based systems [16]. Notably, eucalyptus paired with rice and mango intercropped with potato emerge as popular choices, emphasizing the farmers' preferences influenced by local agroecological nuances [17,18]. The investment analysis of different agroforestry combinations adds depth to the understanding of economic viability. Litchi and mango, when integrated with vegetables, demonstrate higher Net Present Value (NPV) and Internal Rates of Return (IRR) compared to monocrop options [19]. This highlights the potential for increased sustainability and profitability through the strategic integration of tree species with diverse crops. The factors influencing farmers' adoption of agroforestry as a climate-smart strategy further enrich the study. Farm size, age of the respondent, education, training, number of extension visits, and improved market access positively influence adoption [20]. This aligns with the existing literature emphasizing the role of knowledge, education, and external support in promoting sustainable agricultural practices. The challenges faced by farmers for adopting agroforestry practices are comprehensively explored. Lack of training facilities and a shortage of skilled labor emerge as the primary concerns. Pest-related issues, insufficient expertise, and land availability also pose significant challenges [15].

5. Conclusions

In conclusion, the research provides a holistic understanding of the socio-economic and agroecological contexts, shedding light on the challenges and opportunities for promoting agroforestry as a climate-smart strategy. The findings have implications for policymakers, extension service providers, and researchers working towards sustainable agricultural development.

Author Contributions: Conceptualization, M.M.A. and M.S.B.; methodology, M.M.A., M.S.B. and A.C.P.; software, A.C.P. and M.L.R.; validation, M.M.A., M.S.B. and I.J.S.; formal analysis, M.M.A., A.C.P. and I.J.S.; investigation, M.M.A., M.S.B. and M.L.R.; resources, M.M.A. and M.S.B.; data curation, M.M.A. and A.C.P.; writing—original draft preparation, M.M.A. and A.C.P.; writing—review and editing, M.S.B. and I.J.S.; visualization, M.L.R. and A.C.P.; supervision, M.M.A. and M.S.B.; project administration, M.M.A. and M.S.B.; All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data will be available on request.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Devendra, C. *Climate Change Threats and Effects: Challenges for Agriculture and Food Security*; Academy of Sciences Malaysia: Kuala Lumpur, Malaysia, 2012.
2. Hoegh-Guldberg, O.; Jacob, D.; Taylor, M.; Bolaños, T.G.; Bindi, M.; Brown, S.; Camilloni, I.A.; Diedhiou, A.; Djalante, R.; Ebi, K.; et al. The human imperative of stabilizing global climate change at 1.5 °C. *Science* **2019**, *365*, eaaw6974. [[CrossRef](#)] [[PubMed](#)]
3. Post, J.E.; Altman, W. Managing the Environmental Change Process: Barriers and Opportunities 1. In *Managing Green Teams*; Routledge: London, UK, 2017; pp. 84–101.
4. Vinodhini, S.M.; Manibharathi, S.; Pavithra, G.; Sakthivel, S. Agroforestry: Integrating Trees into Agricultural Systems. In *Recent Approaches in Agriculture*; Elite Publishing House: Delhi, India, 2023; p. 246.
5. Raj, A.; Jhariya, M.K.; Yadav, D.K.; Banerjee, A.; Meena, R.S. Agroforestry: A holistic approach for agricultural sustainability. In *Sustainable Agriculture, Forest and Environmental Management*; Springer: Berlin/Heidelberg, Germany, 2019; pp. 101–131.
6. McCabe, C. *Agroforestry and Smallholder Farmers: Climate Change Adaptation through Sustainable Land Use*; Capstone Collection: Lutz, FL, USA, 2013.
7. Rao, K.P.C.; Verchot, L.V.; Laarman, J. Adaptation to climate change through sustainable management and development of agroforestry systems. *J. SAT Agric. Res.* **2007**, *4*, 1–30.
8. Magalhaes, C.A.S.; Pedreira, B.C.; Tonini, H.; Neto, A.L.F. Crop, livestock and forestry performance assessment under different production systems in the north of Mato Grosso, Brazil. *Agrofor. Syst.* **2019**, *93*, 2085–2096. [[CrossRef](#)]
9. Shi, P.-L.; Duan, C.; Wang, L.; Wu, N.; Kotru, R.; Gurung, J. Integrated landscape approaches to building resilience and multifunctionality in the Kailash Sacred Landscape, China. *J. Mt. Sci.* **2021**, *18*, 3321–3335. [[CrossRef](#)]
10. Zerihun, M.F. A socioeconomic Analysis of Factors that Affect the Adoption of Agroforestry Technologies in the Eastern Cape Province of South Africa. Ph.D. Thesis, Tshwane University of Technology, Pretoria, South Africa, 2014.
11. Sanou, L.; Savadogo, P.; Ezebilo, E.E.; Thiombiano, A. Drivers of farmers' decisions to adopt agroforestry: Evidence from the Sudanian savanna zone, Burkina Faso. *Renew. Agric. Food Syst.* **2019**, *34*, 116–133. [[CrossRef](#)]
12. Nair, P.K.R. Climate change mitigation: A low-hanging fruit of agroforestry. In *Agroforestry-The Future of Global Land Use*; Springer: Berlin/Heidelberg, Germany, 2012; pp. 31–67.
13. Garrity, D.P.; Festus, K.; Akinnifesi, O.C.; Ajayi, S.G.; Weldesemayat, J.G.; Mowo, A.K.; Larwanou, M.; Bayala, J. Evergreen Agriculture: A robust approach to sustainable food security in Africa. *Food Secur.* **2010**, *2*, 197–214. [[CrossRef](#)]
14. Cochran, W.G. *Sampling Techniques*; John Wiley & Sons: Hoboken, NJ, USA, 1977.
15. Hanif, M.A.; Roy, R.M.; Bari, M.S.; Ray, P.C.; Rahman, M.S.; Hasan, M.F. Livelihood improvements through agroforestry: Evidence from Northern Bangladesh. *Small-Scale For.* **2018**, *17*, 505–522. [[CrossRef](#)]
16. Hong, Y.; Heerink, N.; Jin, S.; Berentsen, P.; Zhang, L.; van der Werf, W. Intercropping and agroforestry in China—current state and trends. *Agric. Ecosyst. Environ.* **2017**, *244*, 52–61. [[CrossRef](#)]
17. Paul, C.; Weber, M.; Knoke, T. Agroforestry versus farm mosaic systems—Comparing land-use efficiency, economic returns and risks under climate change effects. *Sci. Total Environ.* **2017**, *587*, 22–35. [[CrossRef](#)] [[PubMed](#)]
18. Bai, W.; Sun, Z.; Zheng, J.; Du, G.; Feng, L.; Cai, Q.; Yang, N.; Feng, C.; Zhang, Z.; Evers, J.B.; et al. Mixing trees and crops increases land and water use efficiencies in a semi-arid area. *Agric. Water Manag.* **2016**, *178*, 281–290. [[CrossRef](#)]
19. Talukder, M.S.; Miah, M.M.U.; Miah, M.G.; Haque, M.M.; Rahman, M.M.; Islam, M.M. Fruit tree-based agroforestry systems and their carbon sequestration potentials in different ecosystem of Bangladesh. *J. Agrofor. Environ.* **2019**, *13*, 43–48.
20. Jahan, H.; Rahman, M.W.; Islam, M.S.; Rezwan-Al-Ramim, A.; Tuhin, M.M.-U.-J.; Hossain, M.E. Adoption of agroforestry practices in Bangladesh as a climate change mitigation option: Investment, drivers, and SWOT analysis perspectives. *Environ. Chall.* **2022**, *7*, 100509. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.