

Characteristics of PSA Bounce after Radiotherapy for Prostate Cancer: A Meta-Analysis

Narisa Dewi Maulany Darwis ^{1,2}, Takahiro Oike ^{1,3,*}, Nobuteru Kubo ^{1,3}, Soehartati A Gondhowiardjo ² and Tatsuya Ohno ^{1,3}

¹ Department of Radiation Oncology, Gunma University Graduate School of Medicine, 3-39-22, Showa-machi, Maebashi, Gunma 371-8511, Japan; m1920021@gunma-u.ac.jp (N.D.M.D.); kubo@gunma-u.ac.jp (N.K.); tohno@gunma-u.ac.jp (T.O.)

² Department of Radiation Oncology, Faculty of Medicine Universitas Indonesia—Dr. Cipto Mangunkusumo National General Hospital, Jl. Diponegoro No. 71, Jakarta Pusat, DKI Jakarta 10430, Indonesia; gondhow@gmail.com

³ Gunma University Heavy Ion Medical Center, 3-39-22, Showa-Machi, Maebashi, Gunma 371-8511, Japan

* Correspondence: oiketakahiro@gunma-u.ac.jp

Received: 14 July 2020; Accepted: 4 August 2020; Published: 5 August 2020

Abstract: The rate and characteristics of prostate-specific antigen (PSA) bounce post-radiotherapy remain unclear. To address this issue, we performed a meta-analysis. Reports of PSA bounce post-radiotherapy with a cutoff of 0.2 ng/mL were searched by using Medline and Web of Science. The primary endpoint was the occurrence rate, and the secondary endpoints were bounce characteristics such as amplitude, time to occurrence, nadir value, and time to nadir. Radiotherapy modality, age, risk classification, androgen deprivation therapy, and the follow-up period were extracted as clinical variables. Meta-analysis and univariate meta-regression were performed with random-effect modeling. Among 290 search-positive studies, 50 reports including 26,258 patients were identified. The rate of bounce was 31%; amplitude was 1.3 ng/mL; time to occurrence was 18 months; nadir value was 0.5 ng/mL; time to nadir was 33 months. Univariate meta-regression analysis showed that radiotherapy modality (29.7%), age (20.2%), and risk classification (12.2%) were the major causes of heterogeneity in the rate of bounce. This is the first meta-analysis of PSA bounce post-radiotherapy. The results are useful for post-radiotherapy surveillance of prostate cancer patients.

Keywords: prostate cancer; radiotherapy; prostate-specific antigen (PSA); PSA bounce; meta-analysis; meta-regression

1. Introduction

Radiotherapy is a definitive treatment for prostate cancer (PCa). Prostate-specific antigen (PSA) is the biomarker used for post-treatment surveillance of PCa patients [1,2]. In curative cases, PSA levels decrease gradually over a period of more than five years after radiotherapy and reach a nadir. In a subset of patients, however, PSA levels fluctuate and show a temporal increase called the PSA bounce [3]. It is difficult to appropriately diagnose PSA increase post-radiotherapy as the bounce; therefore, the PSA increase post-radiotherapy can be the cause of severe anxiety in both PCa patients and clinicians. Misinterpretation may even endanger patients by leading to unnecessary salvage treatment in cases meeting the definition of biochemical failure. PSA bounce can occur in relation to various radiotherapy modalities, including external beam radiotherapy (EBRT), stereotactic body radiotherapy (SBRT), low dose-rate brachytherapy (LDR-BT), and high dose-rate brachytherapy (HDR-BT) [4,5]. As these radiotherapy modalities use different radiation sources, doses, and

fractionation, as well as delivery techniques, they can exert different biological effects on the tumor and the prostate. However, the characteristics of PSA bounce in relation to different radiotherapy modalities remain unclear. To address this issue, we performed a meta-analysis of the characteristics of PSA bounce.

2. Results

A systematic literature review was performed to identify studies reporting PSA bounce post-radiotherapy (see Materials and Methods for details) (Figure 1). The search identified 50 studies including 26,258 patients, which were included in the analysis (Table 1) [6–55]. The number of studies and patients stratified by modality is summarized in Table S1. Among the 50 studies, eight were prospective observational studies [14,15,17,32,39–41,50] and the others were retrospective observational studies.




Figure 1. Preferred reporting items for systematic reviews and meta-analyses (PRISMA) flow diagram of the literature review for prostate-specific antigen (PSA) bounce after radiotherapy.

Table 1. Papers that report PSA bounce after radiotherapy included in the meta-analysis.

Author	Year	n	Modality	Age	Risk Group	ADT	Follow Up (M)	Bounce (%)	Amplitude (ng/mL)	Time to Bounce (M)	Nadir (ng/mL)	Time to Nadir (M)	Reference
Merrick et al.	2002	218	EBRT+LDR-BT	66 ± 7	L, I	No	46 ± 14	23.9	0.9 (0.3–3.0)	19 ± 9	NA	NA	[6]
Patel et al.	2004	295	LDR-BT	NA	L, I	Yes, partly	38 (24–68)	28.0	0.5 (0.2–4.1)	19 (8–40)	NA	NA	[7]
Coen et al.	2004	101	LDR-BT	NA	L, I	No	54 (38–86)	39.6	0.6 (0.2–7.5)	18 (7–71)	NA	NA	[8]
Zietman et al.	2005	190	EBRT	NA	L, I, H	Yes, all	60 (40–75)	39.0	0.9 (0.5–1.8)	28 (17–42)	NA	NA	[9]
Ciezki et al.	2006	162	LDR-BT	68 (45–83)	L, I, H	Yes, partly	73	46.3	NA	15 (2–57)	NA	NA	[10]
Horwitz et al.	2006	4839	EBRT	NA	L, I, H	No	75	18.6	NA	NA	NA	NA	[11]
		2693	LDR-BT	NA	L, I, H	No	60	17.5	NA	NA	NA	NA	
Toledano et al.	2007	295	LDR-BT	60–65	L, I	Yes, partly	40 (9–66)	49.0	0.8, mean (0.1–4.1)	19, mean (6–58)	NA	NA	[12]
Bostantic et al.	2007	57	LDR-BT	65 ± 6	L	No	62 ± 10	14.0	0.4	18 ± 9	NA	NA	[13]
		46	LDR-BT	63 ± 7	L	No	64 ± 12	45.7	0.4	22 ± 11	NA	NA	
		29	LDR-BT	66 ± 6	L	Yes, all	67 ± 12	20.7	0.4	6 ± 6	NA	NA	
		32	LDR-BT	67 ± 5	L	Yes, all	62 ± 12	28.1	0.4	18 ± 8	NA	NA	
Crook et al.	2007	292	LDR-BT	64 (45–80)	L, I	No	44 (8–81)	40.0	0.7 (0.2–11.7)	15 (3–29)	0.05 (0.01–0.20)	40	[14]
Mitchell et al.	2008	205	LDR-BT	62, mean (43–75)	L, I	No	45 (24–85)	37.0	0.9 (0.2–5.8)	14 (1–40)	NA	NA	[15]
Makarewicz et al.	2009	121	EBRT+HDR-BT	68 (47–78)	L, I	No	81 (60–106)	31.0	0.2, mean (0.2–0.7)	14 (7–26)	0.8 (0.01–2.1)	NA	[16]
King et al.	2009	41	SBRT	66 (48–83)	L	No	33 (6–45)	29.0	0.3 (0.2–2.4)	18 (12–33)	0.3 (0.03–2.6)	NA	[17]
Pinkawa et al.	2010	135	EBRT	71 (52–83)	L, I, H	Yes, partly	67 (9–97)	20.0	NA	NA	NA	NA	[18]
		66	EBRT+HDR-BT	72 (63–81)	L, I, H	Yes, partly	75 (7–98)	23.0	NA	NA	NA	NA	
		94	LDR-BT	69 (49–81)	L, I	Yes, partly	76 (8–101)	42.0	NA	NA	NA	NA	
Caloglu et al.	2011	820	LDR-BT	68 (45–87)	L, I, H	Yes, partly	58 (36–123)	30.1	NA	17 (2–68)	NA	NA	[19]
Zwahlen et al.	2011	194	LDR-BT	61 (47–75)	L	No	60 (23–109)	50.0	0.5 (0.2–8.3)	14 (0–70)	0.1 (0.0–3.5)	NA	[20]
Aaltomaa et al.	2011	535	LDR-BT	64 (42–80)	L, I, H	Yes, partly	69 (15–131)	27.4	NA	NA	NA	NA	[21]
Beriwal et al.	2012	155	EBRT+LDR-BT	65 ± 7	L, I, H	Yes, partly	36 (24–60)	29.7	0.6, mean (0.2–2.3)	12, mean (6–36)	NA	NA	[22]
Hinnen et al.	2012	975	LDR-BT	66 ± 6	L, I, H	Yes, partly	78 (27–215)	32.0	1.7 (1.0–2.8, IQR)	19 (12–24, IQR)	NA	12 (6–15)	[23]

Mazeron et al.	2012	198	LDR-BT	67 (49–80)	L, I	No	63 (36–119)	35.9	1.0 ± 1.0	18 ± 9	NA	NA	[24]
Bolzicco et al.	2013	71	SBRT	72 (52–82)	L, I, H	Yes, partly	36 (6–76)	12.6	NA	23 (18–30)	0.4	36	[25]
Chen et al.	2013	100	SBRT	69 (48–90)	L, I, H	Yes, partly	27 (16–42)	31.0	0.5 (0.2–2.2)	15 (3–21)	0.4 (0.1–1.9)	24	[26]
Katz et al.	2013	304	SBRT	69, mean (45–88)	L, I, H	Yes, partly	60 (8–78)	17.0	0.5	30	0.1	60	[27]
King et al.	2013	1100	SBRT	70 (44–91)	L, I, H	Yes, partly	36	16.0	0.5 (0.2–5.29)	NA	NA	NA	[28]
Mehta et al.	2013	157	HDR-BT	63 (42–90)	L, I	Yes, partly	55	43.0	0.6 (0.2–4.5)	13 (0.6–64)	NA	NA	[29]
Lee et al.	2014	29	SBRT	72 (50–86)	L, I, H	Yes, partly	41 (12–69)	28.0	0.6 (0.3–1.5)	9	0.3 (0.003–1.7)	23	[30]
Nishihara et al.	2014	116	LDR-BT	66 (51–80)	L, I	No	42 (18–77)	40.5	0.4 (0.2–5.6)	17 (8–36)	NA	NA	[31]
Vu et al.	2014	120	SBRT	68 (47–88)	L, I, H	Yes, partly	24 (18–78)	28.0	0.5	9	NA	NA	[32]
Patel et al.	2014	114	EBRT+HDR-BT	68 (48–79)	L, I	No	66 (24–124)	39.0	0.4 (0.2–6.6)	16 (3–76)	0.1 (0.01–1.7)	53 (8–118)	[33]
Waters et al.	2014	74	EBRT, hopo	68 ± 5	L	No	36, min	31.1	0.6	NA	NA	NA	[34]
		58	EBRT	66 ± 5	L	No	36, min	20.7	0.3	NA	NA	NA	
		230	LDR-BT	64 ± 6	L	No	36, min	29.6	0.6	NA	NA	NA	
Kole et al.	2015	175	SBRT	69 (48–85)	L, I, H	No	36	36.2	NA	15 (1–42)	0.3 (0.02–1.8)	30 (3–48)	[35]
Kishan et al.	2015	130	SBRT	69 (44–87)	L, I	No	40 (12–93)	30.8	0.5 (0.2–3.6)	14 (3–43)	NA	NA	[36]
	2015	220	HDR-BT	64 (43–84)	L, I	No	49 (12–94)	39.5	0.6 (0.2–7.1)	10 (3–63)	NA	NA	
	2015	89	EBRT	66 (52–85)	L, I	No	27 (12–90)	21.3	0.5 (0.2–7.6)	13 (3–66)	NA	NA	
Leduc et al.	2015	274	LDR-BT	62 (45–76)	L	Yes, partly	50 (24–126)	31.0	1.0 (0.2–12.4)	12 (6–37)	NA	NA	[37]
Quivirin et al.	2015	66	LDR-BT	64 ± 5	L	No	35 (13–72)	36.4	1.8 ± 1.6	12 ± 6	NA	NA	[38]
Engeler et al.	2015	713	LDR-BT	63 (42–82)	L, I, H	Yes, partly	41 (24–132)	24.3	0.7 (0.2–6.1)	12 (6–33)	NA	NA	[39]
Kim et al.	2016	33	SBRT	67 (56–72)	L, I	No	51 (6–71)	30.3	0.2 (0.2–1.3)	10 (6–12)	0.2	33	[40]
Kim et al.	2016	47	SBRT	64 (52–82)	L, I	No	42 (36–78)	51.0	0.5 (0.2–6.2)	9 (3–36)	NA	36 (11, SD)	[41]
Phak et al.	2016	35	EBRT+SBRT	69, mean (60–78)	L, I	No	52 (14–74)	28.6	0.2 (0.2–0.5)	11 (6–25)	0.2 (0.04–1.4)	32 (12–51)	[42]
		42	EBRT	71, mean (61–79)	L, I	No	52 (14–74)	21.4	0.3 (0.2–1.2)	15 (6–30)	0.3 (0.04–1.8)	25 (9–58)	
Freiberger et al.	2017	94	LDR-BT	69 (49–81)	L, I	Yes, partly	108	42.0	NA	NA	0.05, mean	32, mean	[43]
		66	EBRT+HDR-BT	72 (63–81)	L, I, H	Yes, partly	108	24.0	NA	NA	0.1, mean	31, mean	
		135	EBRT	71 (52–83)	L, I, H	Yes, partly	108	25.0	NA	NA	0.5, mean	19, mean	

Hauck et al.	2017	554	HDR-BT	63 (40–83)	L, I, H	Yes, partly	44 (12–162)	43.2	NA	11, mean	0.2	NA	[44]
Kindts et al.	2017	192	LDR-BT	60 (50–65)	L, I	Yes, partly	66	36.0	0.6, mean	18, mean	NA	NA	[45]
Romesser et al.	2017	776	EBRT	61–72, IQR	L, I, H	Yes, partly	110 (83–134)	15.9	0.3 (0.2–0.7, IQR)	24 (16–38, IQR)	NA	NA	[46]
Park et al.	2018	74	SBRT	69 (47–81)	L, I, H	No	63 (12–109)	35.2	0.5 (0.2–2.6)	11 (2–38)	0.1 (0.01–2.6)	47 (1–85)	[47]
Astrom et al.	2018	623	EBRT+HDR-BT	66 (47–79)	L, I, H	Yes, partly	132 (2–266)	26.0	1.5 (0.3–12.0)	15 (3–103)	NA	NA	[48]
Burchardt et al.	2018	41	LDR-BT	64 ± 7	L, I	Yes, partly	37 ± 8	26.8	0.7 ± 1.1	18 ± 6	0.5 ± 1.1	23 ± 14	[49]
		53	HDR-BT	67 ± 7	L, I	Yes, partly	33 ± 9	22.6	0.8 ± 0.5	10 ± 4	0.2 ± 0.4	19 ± 14	
Kubo et al.	2018	352	EBRT+LDR-BT	69 (49–82)	L, I, H	Yes, partly	82 (12–157)	33.2	NA	20 (3–55)	NA	NA	[50]
Roy et al.	2019	287	SBRT	69 (49–82, IQR)	L, I	Yes, partly	60 (46–106)	31.1	0.6 (0.35–1.61, IQR)	17 (11–25, IQR)	NA	NA	[51]
Jiang et al.	2019	1062	SBRT	68 (63–73, IQR)	L, I	No	66 (36–60, IQR)	26.0	0.5 (0.3–1.0, IQR)	18 (12–31, IQR)	0.2 (0.1–0.3, IQR)	40 (24–66, IQR)	[52]
Darwis et al.	2020	131	Carbon ions	64, mean (48–80)	L, I	No	60 (39–60)	55.7	0.7 ± 1.0	15 ± 11	0.5 ± 0.3	42 (9–60)	[53]
Nakai et al.	2020	256	HDR-BT	67 ± 6	L, I	No	91 ± 23	32.3	NA	19 ± 23	NA	NA	[54]
Slade et al.	2020	4004	LDR-BT	64 ± 6	L, I	No	120	31.8	NA	NA	NA	NA	[55]
		473	EBRT	64 ± 6	L, I	No	120	27.7	NA	NA	NA	NA	

PSA, prostate-specific antigen; EBRT, external beam radiotherapy; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy; NA, not assessable; IQR, interquartile range; L, low risk; I, intermediate risk; H, high risk; ADT, androgen deprivation therapy; M, months. Age, follow-up, and bounce outcomes are shown as mean ± standard deviation or in median (range) unless otherwise stated.

A meta-analysis showed that the rate of PSA bounce for all studies was 31% (95% confidence interval (CI), 28–33%) (Figure 2). The bounce rates according to modality were as follows: 34% (95% CI, 30–37%) for LDR-BT, 36% (95% CI, 29–42%) for HDR-BT, 22% (95% CI, 19–25%) for EBRT, 28% (95% CI, 23–32%) for SBRT, 28% (95% CI, 26–31%) for EBRT followed by boost irradiation, and 56% (95% CI, 47–64%) for carbon-ion radiotherapy (Figure 2). For all studies, bounce amplitude was 1.3 ng/mL (95% CI, 1.1–1.4 ng/mL); time to bounce occurrence was 18 months (95% CI, 17–20 months); nadir value was 0.5 ng/mL (95% CI, 0.4–0.6 ng/mL); and time to nadir was 33 months (95% CI, 22–43 months). The results of the meta-analysis stratified by modality are summarized in Table 2, and the original forest plots are shown in Figures S1–4. Nadir value was higher in bounce-positive patients than in bounce-negative patients for EBRT, SBRT, and CIRT, whereas time to nadir was greater in bounce-positive than in bounce-negative patients regardless of modality (Table 3).




Figure 2. Meta-analysis of the rate of prostate-specific antigen (PSA) bounce after radiotherapy. ES, effect size; CI, confidence interval; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate

brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy.

Table 2. Summary of the results of the meta-analysis of PSA bounce characteristics.

Modality	Rate of Bounce (%)	Amplitude (ng/mL)	Time to Occurrence (M)	Nadir (ng/mL)	Time to Nadir (M)
LDR-BT	34 (30–37)	1.7 (1.3–2.0)	18 (17–20)	0.5 (−0.1–1.1)	23 (19–28)
HDR-BT	36 (29–42)	1.4 (0.7–2.2)	18 (12–25)	0.2 (0.09–0.3)	19 (15–23)
EBRT	22 (19–25)	0.8 (0.4–1.2)	24 (20–29)	0.6 (0.5–0.7)	29 (25–32)
SBRT	28 (23–32)	1.0 (0.7–1.2)	17 (14–20)	0.6 (0.3–0.8)	38 (26–51)
EBRT + boost	28 (26–31)	1.0 (0.7–1.4)	18 (14–22)	0.6 (0.4–0.8)	44 (19–70)
CIRT	56 (47–64)	0.7 (0.5–1.0)	15 (12–17)	0.5 (0.4–0.6)	42 (40–44)
Pooled ES	31 (28–33)	1.3 (1.1–1.4)	18 (17–20)	0.5 (0.4–0.6)	35 (28–42)
I^2 , p values	93.5% ($p < 0.05$)	98.3% ($p < 0.05$)	95.7% ($p < 0.05$)	99.5% ($p < 0.05$)	98.4% ($p < 0.05$)

PSA, prostate-specific antigen; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy; ES, effect size; M, months. Data are means (95% confidence interval).

Table 3. Summary of the results of the meta-analysis of PSA nadir and time to nadir stratified by bounce occurrence.

Modality	Nadir (ng/mL)		Time to Nadir (M)	
	Bounce	No bounce	Bounce	No bounce
LDR-BT	NA	NA	NA	NA
HDR-BT	NA	NA	NA	NA
EBRT	0.7 (0.7–0.8)	0.5 (0.5–0.5)	42 (40–43)	29 (28–29)
SBRT	0.6 (0.5–0.7)	0.3 (0.3–0.4)	NA	NA
EBRT + Boost	0.3 (0.2–0.4)	0.5 (0.4–0.6)	64 (58–70)	54 (48–60)
CIRT	0.6 (0.5–0.7)	0.4 (0.3–0.5)	48 (45–50)	36 (33–40)
Pooled effect size	0.6 (0.3–0.8)	0.5 (0.4–0.6)	50 (42–59)	39 (27–51)
I^2 , p values	98.5% ($p < 0.05$)	95.1% ($p < 0.05$)	96.6% ($p < 0.05$)	97.6% ($p < 0.05$)

PSA, prostate-specific antigen; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy; NA, not assessible; M, months. Data are means (95% confidence interval).

The rate and characteristics of the bounce showed significant heterogeneity among the studies (Table 2). To find the cause of the heterogeneity, we performed univariate meta-regression analysis. Age, radiotherapy modality, use of androgen deprivation therapy (ADT), and risk classification were selected as the covariates for meta-regression based on previous studies reporting that these factors affect the bounce kinetics [4,5]. The heterogeneity in the bounce rate was attributed to modality (29.7%), age (20.2%), and risk classification (12.2%) (Figures 3A,B, Table 3). Regarding bounce amplitude, age was a significant cause of heterogeneity (Figure 3C, Table 3). For time to bounce occurrence, modality was a significant cause of heterogeneity (Table 4).




Figure 3. Univariate meta-regression of heterogeneity in the rate of bounce by age (A) or by risk group (B), and that in bounce amplitude by age (C). L, low risk; IM, intermediate risk; H, high risk.

Table 4. Univariate meta-regression for the proportion and characteristics of bounce.

Covariates	Rate of Bounce (<i>n</i> = 65)			Amplitude (<i>n</i> = 37)			Time to Occurrence (<i>n</i> = 45)			Nadir (<i>n</i> = 15)			Time to Nadir (<i>n</i> = 9)		
	Coefficient	<i>p</i>	R ² (%)	Coefficient	<i>p</i>	R ² (%)	Coefficient	<i>p</i>	R ² (%)	Coefficient	<i>p</i>	R ² (%)	Coefficient	<i>p</i>	R ² (%)
Age	−0.07 (−0.10 to −0.03)	<0.01	20.2	−0.14 (−0.22 to −0.06)	<0.01	28.1	0.30 (−0.27 to 0.87)	0.33	1.1	−0.01 (−0.06 to 0.05)	0.78	0.0	−0.32 (−4.82 to 4.18)	0.87	0.0
Modality															
LDR-BT	−0.08 (−0.46 to 0.30)	0.66	29.7	0.25 (−0.61 to 1.10)	0.56	15.4	−5.57 (−11.32 to 0.18)	0.05	5.0	−0.05 (−0.79 to 0.68)	0.88	0.0	−18.96 (−78.66 to 40.74)	0.38	0.0
HDR-BT	NA	NA		NA	NA		NA	NA		NA	NA		NA	NA	
EBRT	−0.63 (−1.07 to −0.19)	<0.01		−0.53 (−1.56 to 0.50)	0.30		NA	NA		0.11 (−0.79 to 1.01)	0.78		−13.31 (−72.86 to 46.24)	0.52	
SBRT	−0.38 (−0.79 to 0.03)	0.07		−0.48 (−1.36 to 0.40)	0.27		−7.24 (−13.36 to −1.11)	0.02		0.07 (−0.61 to 0.75)	0.82		−3.90 (−52.43 to 44.62)	0.81	
EBRT + boost	−0.32 (−0.75 to 0.11)	0.14		−0.41 (−1.35 to 0.53)	0.38		−5.92 (−12.39 to 0.56)	0.07		0.07 (−0.65 to 0.80)	0.82		2.23 (−49.31 to 53.77)	0.89	
CIRT	0.83 (−0.26 to 1.68)	0.05		−0.70 (−2.24 to 0.83)	0.35		−9.50 (−20.68 to 1.67)	0.09		NA	NA		NA	NA	
ADT	−0.14 (−0.34 to 0.06)	0.17		−0.20 (−0.63 to 0.22)	0.34	0.8	0.32 (−2.18 to 2.82)	0.79	0.0	−0.07 (−0.41 to 0.27)	0.66	0.0	−18.10 (−37.28 to 1.08)	0.06	33.8
Risk group	−0.20 (−0.35 to −0.04)	0.01	12.2	−0.23 (−0.62 to 0.15)	0.22	1.3	1.42 (−0.78 to 3.61)	0.20	0.0	0.02 (−0.21 to 0.24)	0.88	0.0	0.94 (−24.00 to 25.89)	0.93	0.0

LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy; ADT, androgen deprivation therapy. NA, not assessable due to collinearity. Data are means (95% confidence interval).

3. Discussion

The strength of this study is that this is the first meta-analysis to investigate the characteristics of PSA bounce post-radiotherapy. We report the rate, amplitude, nadir, and time course of the bounce for different modalities including brachytherapy, EBRT, SBRT, and CIRT. We also report that the bounce occurs more frequently and with greater amplitude in brachytherapy than in EBRT, and a younger age is associated with a higher incidence and greater amplitude of the bounce. These findings have been extensively reported in mono-institutional studies, e.g., the large-scale study by Romesser [46], which were validated here for the first time by meta-analysis. From this standpoint, the results of the present study are useful for post-radiotherapy surveillance of prostate cancer patients to help oncologists and patients interpret temporal PSA increases post-treatment.

The limitations of this study, on the other hand, are the following. First, the studies analyzed were extremely heterogeneous regarding clinical factors such as dose, fractionation, bounce rate according to ADT usage, and risk classification, which was difficult to control in a meta-analysis design. In particular, the ADT strategy (i.e., the presence or absence of adjuvant or neoadjuvant use) should have affected post-radiotherapy PSA kinetics to a large extent, which was difficult to adjust by study design. Second, we were not able to analyze the PSA kinetics post-radiotherapy stratified by bounce positivity except for nadir and time to nadir. This was because extraction of the corresponding data from the original articles was technically impossible; i.e., the original articles did not contain the PSA kinetics data linked to specific clinical variables (e.g., age and risk) in a form that we can compute in the meta-analysis. Third, we were unable to perform multivariate meta-regression analysis because of the small number of studies. Fourth, most of the studies included had a retrospective design, and no randomized studies were identified. Finally, studies on particle therapy were rarely identified (i.e., one study on CIRT and no studies on proton therapy).

The molecular mechanisms underlying PSA bounce remain to be elucidated. Studies have shown that PSA is released from both tumor tissues and the normal prostate glands after irradiation [48]. Radiation-induced antitumor immunity may contribute to the release of PSA from tumor tissues. For example, Yamamoto et al. reported intra-tumoral infiltration of CD3- and CD8-positive lymphocytes in bounce-positive patients [56]. In the present meta-analysis, the bounce was more prevalent after brachytherapy and SBRT than after EBRT. In addition, the bounce rate for CIRT was strikingly high, although only one study was analyzed. These findings may be explained by the highly concentrated dose delivery by brachytherapy, SBRT, and CIRT compared with that of EBRT. Evidence suggests that a high, single-fractionated dose induces antitumor immunity efficiently [57], partially by promoting DNA damage response signaling [58]. In addition, the properties of carbon ions as high linear energy transfer radiation to efficiently induce antitumor immunity (e.g., induction of HMGB1 [59], OX40L, CD40, ICAM-1, and MHC-1, and suppression of PD-L1 [60]) might contribute to the high bounce rate for CIRT. Another possible explanation for the higher bounce rate associated with brachytherapy, SBRT, and CIRT is that the highly concentrated doses delivered by these modalities destroy the normal prostate glands more efficiently. Kirilova et al. showed an increase in metabolism indicative of inflammation in the normal prostate gland of patients experiencing bounce, which supports this notion [61].

In addition to modality, the meta-regression results indicated that younger age is associated with greater bounce occurrence and amplitude. This is consistent with the findings of the systematic literature review, in which 29 of the 50 papers analyzed identify younger age as a predictor of bounce. Yamamoto et al. suggested that this may be related to the higher immunocompetency in younger patients [56]. Further research is warranted to elucidate immunologic responses of PCa and the prostate glands after radiotherapy.

4. Materials and Methods

4.1. Endpoint Definition

The primary endpoint of this study was the rate of PSA bounce. Secondary endpoints included the characteristics of bounce, i.e., bounce amplitude, time to occurrence, nadir value, and time to nadir. Definitions of these endpoints are listed in Table S2.

4.2. Inclusion and Exclusion Criteria

The inclusion criteria were as follows: (i) an original clinical study reporting on radiotherapy for PCa; (ii) available rate of PSA bounce; and (iii) bounce defined as an increase in PSA over a cutoff of 0.2 ng/mL followed by a spontaneous decrease to or below the pre-bounce nadir [19]. The exclusion criteria were as follows: (i) manuscript written in languages other than English; (ii) full manuscript not available; (iii) subgroup analysis of a given reported cohort; (iv) follow-up shorter than 24 months.

4.3. Study Selection

A systematic literature search based on preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines [62] was performed on 20 March 2020, using two databases, Medline and Web of Science. The search strategy and population-intervention-comparison-outcome metrics [63] are described in Tables S3 and S4, respectively. The search results were combined using the bibliographic management software Mendeley Desktop version 1.19.4 (Mendeley, London, UK), and duplicates were eliminated. Two investigators (N.D.M.D. and T.Oi.) independently reviewed all records in the following three steps. In step 1, the titles of all records were reviewed to detect potentially relevant records. In step 2, the abstracts of all records that passed step 1 were reviewed to detect potentially relevant records. In step 3, the entire manuscripts of all records that passed step 2 were examined if they contained extractable data for the primary endpoint.

4.4. Data Extraction

From the studies identified in Section 4.3, two investigators (N.D.M.D. and T.Oi.) independently extracted the following data: primary and secondary endpoints, radiotherapy modality, age, risk classification [64], the use of ADT, and follow-up period.

4.5. Quality Assessment

Two investigators (N.D.M.D. and T.Oi.) independently confirmed that the methodological quality of the included studies was adequate based on the Quality Assessment Tool for Case Series Studies published by the National Heart, Lung, and Blood Institute-National Institute of Health, U.S. [65]. For Sections 4.3–4.5, decisions were made based on discussion by the two investigators to resolve disagreements on the review results.

4.6. Statistical Analysis

Radiotherapy modalities were classified into six groups as follows: iLDR-BT (^{103}Pd , ^{125}I , or ^{131}Cs), HDR-BT (^{192}Ir), EBRT (three-dimensional conformal radiotherapy or intensity-modulated radiation therapy), SBRT (using CyberKnife or linac), EBRT+boost (using LDR-BT, HDR-BT, or SBRT), and CIRT. Meta-analysis of bounce (binomial data) was performed using *metaprop*, a command of Stata (MP 13, StataCorp, College Station, TX, USA) [66]. Meta-analysis of the characteristics of bounce (continuous variables) was performed using *metan*, a Stata command. For the datasets that lacked the mean and standard deviation to be pooled, these values were estimated from the sample size, median, range, and/or interquartile range, as reported previously [67]. A random-effects model was used considering a high extent of inter-study heterogeneity examined using X^2 and I^2 statistics [68]. Meta-regression was performed to analyze the effect of clinical factors on inter-study heterogeneity

in effect size using *metareg*, a Stata command [69]. To construct the *metareg* command for bounce rate, logit prevalence and its standard error were used [70,71]; for the remaining PSA kinetics outcomes, mean and standard error were used [72]. Results with a *p*-value < 0.05 were interpreted as significant.

5. Conclusions

This is the first study to report the results of meta-analysis and meta-regression of PSA bounce post-radiotherapy. Meta-analysis of 50 studies including 26,258 patients showed that the rate of PSA bounce for all studies was 31% (95% CI, 28–33%); bounce amplitude was 1.3 ng/mL (95% CI, 1.1–1.4 ng/mL); time to bounce occurrence was 18 months (95% CI, 17–20 months); nadir value was 0.5 ng/mL (95% CI, 0.4–0.6 ng/mL); and time to nadir was 33 months (95% CI, 22–43 months). The bounce occurred more frequently and with greater amplitude in brachytherapy than in EBRT. Univariate meta-regression showed that younger age is associated with a higher incidence and greater amplitude of bounce. These data will be useful for post-radiotherapy surveillance of PCa patients to help oncologists and patients interpret temporal PSA increases post-treatment.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1. Table S1: The number of studies and patients according to radiotherapy modality, Table S2: Definition of endpoints, Table S3: Search strategy, Table S4: PICO metrics, Figure S1: Meta-analysis of the amplitude of prostate-specific antigen (PSA) bounce after radiotherapy. ES, effect size; CI, confidence interval; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy. Figure S2: Meta-analysis of the time to occurrence of prostate-specific antigen (PSA) bounce after radiotherapy. ES, effect size; CI, confidence interval; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy. Figure S3: Meta-analysis of prostate-specific antigen (PSA) nadir values after radiotherapy. ES, effect size; CI, confidence interval; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy. Figure S4: Meta-analysis of the time to prostate-specific antigen (PSA) nadir after radiotherapy. ES, effect size; CI, confidence interval; LDR-BT, low dose-rate brachytherapy; HDR-BT, high dose-rate brachytherapy; EBRT, external beam radiotherapy; SBRT, stereotactic body radiotherapy; CIRT, carbon ion radiotherapy.

Author Contributions: Conceptualization, T.O. (Takahiro Oike); formal analysis, N.D.M.D. and T.O. (Takahiro Oike); writing—original draft preparation, N.D.M.D.; writing—review and editing, T.O. (Takahiro Oike); supervision, N.K., S.A.G., and T.O. (Tatsuya Ohno). All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by Gunma University Heavy Ion Medical Center. This work was also supported by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science, and Technology of Japan for programs for Leading Graduate Schools, Cultivating Global Leaders in Heavy Ion Therapeutics and Engineering.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Parker, C.; Gillessen, S.; Heidenreich, A.; Horwich, A. Cancer of the prostate: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* **2015**, *26*, v69–v77, doi:10.1093/annonc/mdv222.
2. Mohler, J.L.; Antonarakis, E.S.; Armstrong, A.J.; D’Amico, A.V.; Davis, B.J.; Dorff, T.; Eastham, J.A.; Enke, C.A.; Farrington, T.A.; Higano, C.S.; et al. Prostate cancer, version 2.2019, NCCN clinical practice guidelines in oncology. *J. Natl. Compr. Canc. Netw.* **2019**, *17*, 479–505, doi:10.6004/jnccn.2019.0023.
3. Wallner, K.; Blasko, J.; Dattoli, M. *Prostate Brachytherapy Made Complicated*; SmartMedicine Press: Seattle, WA, USA, 1997; pp. 14.11–14.15.
4. Caloglu, M.; Ciezki, J. Prostate-specific antigen bounce after prostate brachytherapy: Review of a confusing phenomenon. *Urology* **2009**, *74*, 1183–1190, doi:10.1016/j.urology.2009.01.043.

5. Pickles, T. Prostate-specific antigen (PSA) bounce and other fluctuations: Which biochemical relapse definition is least prone to psa false calls? An analysis of 2030 men treated for prostate cancer with external beam or brachytherapy with or without adjuvant androgen deprivation therapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2006**, *64*, 1355–1359, doi:10.1016/j.ijrobp.2005.10.008.
6. Merrick, G.S.; Butler, W.M.; Wallner, K.E.; Galbreath, R.W.; Anderson, R.L. Prostate-specific antigen spikes after permanent prostate brachytherapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2002**, *54*, 450–456, doi:10.1016/S0360-3016(02)02948-6.
7. Patel, C.; Elshaikh, M.A.; Angermeier, K.; Ulchaker, J. PSA bounce predicts early success in patients with permanent Iodine-125 prostate implant. *Urology* **2004**, *63*, 110–113, doi:10.1016/j.urology.2003.08.025.
8. Coen, J.; Zietman, A.; Grocela, J.; Heney, N.; Babayan, R. Definitions of biochemical control after permanent interstitial brachytherapy as sole treatment for localized prostate cancer: Interpreting the PSA bounce. *Int. J. Radiat. Oncol. Biol. Phys.* **2004**, *60*, S444.
9. Zietman, A.L.; Christodouleas, J.P.; Shipley, W.U. PSA bounces after neoadjuvant androgen deprivation and external beam radiation: Impact on definitions of failure. *Int. J. Radiat. Oncol. Biol. Phys.* **2005**, *62*, 714–718, doi:10.1016/j.ijrobp.2004.11.020.
10. Ciezki, J.P.; Reddy, C.A.; Garcia, J.; Angermeier, K.; Ulchaker, J.; Mahadevan, A.; Chehade, N.; Altman, A.; Klein, E.A. PSA kinetics after prostate brachytherapy: PSA bounce phenomenon and its implications for PSA doubling time. *Int. J. Radiat. Oncol. Biol. Phys.* **2006**, *64*, 512–517, doi:10.1016/j.ijrobp.2005.07.960.
11. Horwitz, E.; Levy, L.; Martinez, A.; Potters, L.; Beyer, D.; Blasko, J.; Sandler, H.; Buskirk, S.; Zietman, A.; Kuban, D. The post-treatment PSA bounce for prostate cancer patients treated with external beam RT or permanent brachytherapy alone is not biochemically or clinically significant: A multi-institutional pooled analysis of more than 7500 patients. *Int. J. Radiat. Oncol.* **2006**, *66*, S205, doi:10.1016/j.ijrobp.2006.07.394.
12. Toledano, A.; Chauveinc, L.; Flam, T.; Thiounn, N.; Solignac, S.; Timbert, M.; Rosenwald, J.C.; Cosset, J.M. PSA bounce after permanent implant prostate brachytherapy may mimic a biochemical failure. *Cancer/Radiother.* **2007**, *11*, 105–110, doi:10.1016/j.canrad.2006.10.005.
13. Bostancic, C.H.; Merrick, G.S.; Butler, W.M.; Wallner, K.E.; Allen, Z.; Glbreath, R.; Lief, J.; Gutman, S.E. Isotope and patient age predict for PSA spikes after permanent prostate brachytherapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2007**, *68*, 1431–1437, doi:10.1016/j.ijrobp.2007.01.066.
14. Crook, J.; Gillian, C.; Yeung, I.; Austen, L.; Mclean, M. PSA kinetics and PSA bounce following permanent seed prostate brachytherapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2007**, *69*, 426–433, doi:10.1016/j.ijrobp.2007.03.031.
15. Mitchell, D.M.; Swindell, R.; Elliott, T.; Wylie, J.P.; Taylor, C.M.; Logue, J.P. Analysis of prostate-specific antigen bounce after I125 permanent seed implant for localised prostate cancer. *Radiother. Oncol.* **2008**, *88*, 102–107, doi:10.1016/j.radonc.2008.04.004.
16. Makarewicz, R.; Lebioda, A.; Terlikiewicz, J.; Biedka, M. PSA bouncing after brachytherapy HDR and external beam radiation therapy: A study of 121 patients with minimum 5-years follow-up. *J. Contemp. Brachyther.* **2009**, *1*, 92–96.
17. King, C.R.; Brooks, J.D.; Gill, H.; Pawlicki, T.; Cotruz, C. Stereotactic body radiotherapy for localized prostate cancer: Interim results of a prospective phase II clinical trial. *Int. J. Radiat. Oncol. Biol. Phys.* **2009**, *73*, 1043–1048.
18. Pinkawa, M.; Piroth, M.D.; Holy, R.; Fishedick, K.; Schaar, S.; Borchers, H.; Heidenreich, A.; Eble, M.J. Prostate-specific antigen kinetics following external-beam radiotherapy and temporary (Ir-192) or permanent (I-125) brachytherapy for prostate cancer. *Radiother. Oncol.* **2010**, *96*, 25–29, doi:10.1016/j.radonc.2010.02.010.
19. Caloglu, M.; Ciezki, J.P.; Reddy, C.A.; Angermeier, K.; Ulchaker, J.; Chehade, N.; Altman, A.; Magi-Galuzzi, C.; Klein, E.A. PSA bounce and biochemical failure after brachytherapy for prostate cancer: A study of 820 patients with a minimum of 3 years of follow-up. *Int. J. Radiat. Oncol. Biol. Phys.* **2011**, *80*, 735–741, doi:10.1016/j.ijrobp.2010.02.021.
20. Zwahlen, D.R.; Smith, R.; Andrianopoulos, N.; Matheson, B.; Royce, P.; Millar, J.L. Prostate-specific antigen bounce after permanent Iodine-125 prostate brachytherapy—An Australian analysis. *Int. J. Radiat. Oncol. Biol. Phys.* **2011**, *79*, 179–187, doi:10.1016/j.ijrobp.2009.10.016.
21. Aaltomaa, S.H.; Kataja, V.V.; Raty, A.; Palmgren, J.-E.; Lahtinen, T. Does the outcome of prostate cancer patients with large prostates differ from small prostate size in permanent seed, low dose-rate brachytherapy? *Scand. J. Urol. Nephrol.* **2011**, *45*, 339–345, doi:10.3109/00365599.2011.590994.

22. Beriwal, S.; Smith, R.P.; Houser, C.; Benoit, R.M. Prostate-specific antigen spikes with ¹³¹Cs brachytherapy: Is there a difference with other radioisotopes? *Brachytherapy* **2012**, *11*, 457–459, doi:10.1016/j.brachy.2012.04.004.
23. Hinnen, K.A.; Monninkhof, E.M.; Battermann, J.J.; Van Roermund, J.G.H.; Frank, S.J.; Van Vulpen, M. Prostate specific antigen bounce is related to overall survival in prostate brachytherapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2012**, *82*, 883–888, doi:10.1016/j.ijrobp.2010.11.049.
24. Mazon, R.; Bajard, A.; Montbarbon, X.; Gassa, F.; Malet, C.; Rocher, F.; Clippe, S.; Bringeon, G.; Desmettre, O.; Pommier, P. Permanent ¹²⁵I-seed prostate brachytherapy: Early prostate specific antigen value as a predictor of PSA bounce occurrence. *Radiat. Oncol.* **2012**, *7*, 46, doi:10.1186/1748-717X-7-46.
25. Bolzicco, G.; Favretto, M.S.; Satariano, N.; Scremin, E.; Tambone, C.; Tasca, A. A single-center study of 100 consecutive patients with localized prostate cancer treated with stereotactic body radiotherapy. *BMC Urol.* **2013**, *13*, 49.
26. Chen, L.N.; Suy, S.; Uhm, S.; Oermann, E.K.; Ju, A.W.; Chen, V.; Hanscom, H.N.; Laing, S.; Kim, J.S.; Batipps, G.P. Stereotactic body radiation therapy (SBRT) for clinically localized prostate cancer: The Georgetown University experience. *Radiat. Oncol.* **2013**, *8*, 1–10.
27. Katz, A.J.; Santoro, M.; Diblasio, F.; Ashley, R. Stereotactic body radiotherapy for localized prostate cancer: Disease control and quality of life at 6 years. *Radiat. Oncol.* **2013**, *8*, 1–8, doi:10.1186/1748-717X-8-118.
28. King, C.R.; Freeman, D.; Kaplan, I.; Fuller, D.; Bolzicco, G.; Collins, S.; Meier, R.; Wang, J.; Kupelian, P.; Steinberg, M.; et al. Stereotactic body radiotherapy for localized prostate cancer: Pooled analysis from a multi-institutional consortium of prospective phase II trials. *Radiother. Oncol.* **2013**, *109*, 217–221, doi:10.1016/j.radonc.2013.08.030.
29. Mehta, N.H.; Kamrava, M.; Wang, P.C.; Steinberg, M.; Demanes, J. Prostate-specific antigen bounce after high-dose-rate monotherapy for prostate cancer. *Int. J. Radiat. Oncol. Biol. Phys.* **2013**, *86*, 729–733, doi:10.1016/j.ijrobp.2013.02.032.
30. Lee, Y.H.; Son, S.H.; Yoon, S.C.; Yu, M.; Choi, B.O.; Kim, Y.S.; Jang, H.S.; Lee, S.N.; Jang, J.S.; Hwang, T.K. Stereotactic body radiotherapy for prostate cancer: A preliminary report. *Asia. Pac. J. Clin. Oncol.* **2014**, *10*, 46–53, doi:10.1111/j.1743-7563.2012.01589.x.
31. Nishihara, K.; Nakiri, M.; Chikui, K.; Suekane, S.; Matsuoka, K.; Hattori, C.; Ogo, E.; Abe, T.; Matsumoto, Y.; Ishitake, T. Relationship between sexual function and prostate-specific antigen bounce after Iodine-125 permanent implant brachytherapy for localized prostate cancer. *Int. J. Urol.* **2014**, *21*, 658–663, doi:10.1111/iju.12411.
32. Vu, C.C.; Haas, J.A.; Katz, A.E.; Witten, M.R. Prostate-specific antigen bounce following stereotactic body radiation therapy for prostate cancer. *Front. Oncol.* **2014**, *4*, 1–4, doi:10.3389/fonc.2014.00008.
33. Patel, N.; Souhami, L.; Mansure, J.J.; Duclos, M.; Aprikian, A.; Faria, S.; David, M.; Cury, F.L. Prostate-specific antigen bounce after high-dose-rate prostate brachytherapy and hypofractionated external beam radiotherapy. *Brachytherapy* **2014**, *13*, 450–455, doi:10.1016/j.brachy.2014.05.005.
34. Waters, A.; Delouya, G.; Donath, D.; Lambert, C.; Larrivée, S.; Zorn, K.C.; Taussky, D. Risk factors for PSA bounce following radiotherapy: Outcomes from a multi-modal therapy analysis. *Can. J. Urol.* **2014**, *21*, 7548–7553.
35. Kole, T.P.; Chen, L.N.; Obayomi-Davies, O.; Kim, J.S.; Lei, S.; Suy, S.; Dritschilo, A.; Collins, S.P. Prostate specific antigen kinetics following robotic stereotactic body radiotherapy for localized prostate cancer. *Acta Oncol.* **2015**, *54*, 832–838, doi:10.3109/0284186X.2014.983656.
36. Kishan, A.U.; Wang, P.; Upadhyaya, S.K.; Hauswald, H.; Demanes, D.J.; Nickols, N.G.; Kamrava, M.; Sadeghi, A.; Kupelian, P.A.; Steinberg, M.L.; et al. SBRT and HDR brachytherapy produce lower PSA nadirs and different PSA decay patterns than conventionally fractionated IMRT in patients with low- or intermediate-risk prostate cancer. *Pract. Radiat. Oncol.* **2015**, *6*, 268–275, doi:10.1016/j.prro.2015.11.002.
37. Leduc, N.; Atallah, V.; Creoff, M.; Rabia, N.; Taouil, T.; Escarmant, P.; Vinh-Hung, V. Prostate-specific antigen bounce after curative brachytherapy for early-stage prostate cancer: A study of 274 african-caribbean patients. *Brachytherapy* **2015**, *14*, 826–833, doi:10.1016/j.brachy.2015.09.005.
38. Quivrin, M.; Loffroy, R.; Cormier, L.; Mazoyer, F.; Bertaut, A.; Chambade, D.; Martin, E.; Maingon, P.; Walker, P.; Créhange, G. Multiparametric MRI and post implant CT-based dosimetry after prostate brachytherapy with iodine seeds: The higher the dose to the dominant index lesion, the lower the PSA bounce. *Radiother. Oncol.* **2015**, *117*, 258–261, doi:10.1016/j.radonc.2015.08.020.

39. Engeler, D.S.; Schwab, C.; Thöni, A.F.; Hochreiter, W.; Prikler, L.; Suter, S.; Stucki, P.; Schiefer, J.; Plasswilm, L.; Schmid, H.-P.; et al. PSA bounce after 125I-brachytherapy for prostate cancer as a favorable prognosticator. *Strahlenther. Onkol.* **2015**, *191*, 787–791, doi:10.1007/s00066-015-0860-0.
40. Kim, H.J.; Phak, J.H.; Kim, W.C. Hypofractionated stereotactic body radiotherapy in low- and intermediate-risk prostate carcinoma. *Radiat. Oncol. J.* **2016**, *34*, 260–264.
41. Kim, D.N.; Straka, C.; Cho, L.C.; Lotan, Y.; Yan, J.; Kavanagh, B.; Raben, D.; Cooley, S.; Brindle, J.; Xie, X.J.; et al. Early and multiple PSA bounces can occur following high-dose prostate stereotactic body radiation therapy: Subset analysis of a phase 1/2 trial. *Pract. Radiat. Oncol.* **2016**, *7*, e43–e49.
42. Phak, J.H.; Kim, H.J.; Kim, W.C. Prostate-specific antigen kinetics following hypofractionated stereotactic body radiotherapy boost as post-external beam radiotherapy versus conventionally fractionated external beam radiotherapy for localized prostate cancer. *Prostate Int.* **2016**, *4*, 25–29, doi:10.1016/j.prnil.2015.12.001.
43. Freiburger, C.; Berneking, V.; Vögeli, T.A.; Kirschner-Hermanns, R.; Eble, M.J.; Pinkawa, M. Long-term prognostic significance of rising PSA levels following radiotherapy for localized prostate cancer—Focus on overall survival. *Radiat. Oncol.* **2017**, *12*, 1–7, doi:10.1186/s13014-017-0837-5.
44. Hauck, C.R.; Ye, H.; Chen, P.Y.; Gustafson, G.S.; Limbacher, A.; Krauss, D.J. Increasing fractional doses increases the probability of benign PSA bounce in patients undergoing definitive HDR brachytherapy for prostate cancer. *Int. J. Radiat. Oncol. Biol. Phys.* **2017**, *98*, 108–114, doi:10.1016/j.ijrobp.2017.01.025.
45. Kindts, I.; Stellamans, K.; Billiet, I.; Pottel, H.; Lambrecht, A. I-125 brachytherapy in younger prostate cancer patients: Outcomes in low- and intermediate-risk disease. *Strahlenther. Onkol.* **2017**, *1–7*, doi:10.1007/s00066-017-1142-9.
46. Romesser, P.B.; Pei, X.; Shi, W.; Zhang, Z.; Kollmeier, M.; McBride, S.M.; Zelefsky, M.J. Prostate-specific antigen (PSA) bounce after dose-escalated external beam radiation therapy is an independent predictor of PSA recurrence, metastasis, and survival in prostate adenocarcinoma patients. *Int. J. Radiat. Oncol.* **2017**, *100*, 59–67, doi:10.1016/j.ijrobp.2017.09.003.
47. Park, Y.; Park, H.J.; Jang, W.I.; Jeong, B.K.; Kim, H.-J.; Chang, A.R. Long-term results and PSA kinetics after robotic SBRT for prostate cancer: Multicenter retrospective study in Korea (Korean Radiation Oncology Group Study 15–01). *Radiat. Oncol.* **2018**, *13*, 1–7, doi:10.1186/s13014-018-1182-z.
48. Åström, L.; Sandin, F.; Holmberg, L. Good prognosis following a PSA bounce after high dose rate brachytherapy and external radiotherapy in prostate cancer. *Radiother. Oncol.* **2018**, *129*, 561–566, doi:10.1016/j.radonc.2018.08.011.
49. Burchardt, W.; Skowronek, J. Time to PSA rise differentiates the PSA bounce after HDR and LDR brachytherapy of prostate cancer. *J. Contemp. Brachyther.* **2018**, *10*, 1–9, doi:10.5114/jcb.2018.73786.
50. Kubo, K.; Wadasaki, K.; Kimura, T.; Murakami, Y.; Kajiwara, M.; Teishima, J.; Matsubara, A.; Nagata, Y. Clinical features of prostate-specific antigen bounce after 125I brachytherapy for prostate cancer. *J. Radiat. Res.* **2018**, *59*, 649–655, doi:10.1093/jrr/rry059.
51. Roy, S.; Loblaw, A.; Cheung, P.; Chu, W.; Chung, H.T.; Vesprini, D.; Ong, A.; Chowdhury, A.; Panjwani, D.; Pang, G.; et al. Prostate-specific antigen bounce after stereotactic body radiotherapy for prostate cancer: A pooled analysis of four prospective trials. *Clin. Oncol.* **2019**, *31*, 621–629, doi:10.1016/j.clon.2019.05.001.
52. Jiang, N.Y.; Dang, A.T.; Yuan, Y.; Chu, F.-I.; Shabsovich, D.; King, C.R.; Collins, S.P.; Aghdam, N.; Suy, S.; Mantz, C.A.; et al. Multi-institutional analysis of prostate-specific antigen kinetics after stereotactic body radiation therapy. *Int. J. Radiat. Oncol.* **2019**, *105*, 628–636, doi:10.1016/j.ijrobp.2019.06.2539.
53. Darwis, N.D.; Oike, T.; Kawamura, H. Kinetics of prostate-specific antigen after carbon ion radiotherapy for prostate cancer. *Cancers* **2020**, *12*, 589, doi:10.3390/cancers12030589.
54. Nakai, Y.; Tanaka, N.; Asakawa, I.; Anai, S.; Miyake, M.; Morizawa, Y.; Hori, S.; Owari, T.; Fujii, T.; Yamaki, K.; et al. Prostate-specific antigen bounce after 125I-brachytherapy for prostate cancer is a favorable prognosticator in patients who are biochemical recurrence-free at 4 years and correlates with testosterone. *Jpn. J. Clin. Oncol.* **2020**, *50*, 58–65, doi:10.1093/jjco/hyz136.
55. Slade, A.N.; Dahman, B.; Chang, M.G. Racial differences in the PSA bounce in predicting prostate cancer outcomes after brachytherapy: Evidence from the Department of Veterans Affairs. *Brachytherapy* **2020**, *19*, 6–12, doi:10.1016/j.brachy.2019.08.008.
56. Yamamoto, Y.; Offord, C.P.; Kimura, G.; Kuribayashi, S.; Takeda, H.; Tsuchiya, S.; Shimojo, H.; Kanno, H.; Bozic, I.; Nowak, M.A.; et al. Tumour and immune cell dynamics explain the PSA bounce after prostate cancer brachytherapy. *Br. J. Cancer* **2016**, *115*, 195–202, doi:10.1038/bjc.2016.171.

57. Helm, A.; Ebner, D.K.; Tinganelli, W.; Simoniello, P.; Bisio, A.; Marchesano, V.; Durante, M.; Yamada, S.; Shimokawa, T. Combining heavy-ion therapy with immunotherapy: An update on recent developments. *Int. J. Part. Ther.* **2019**, *5*, 84–93, doi:10.14338/IJPT-18-00024.1.
58. Vanpouille-Box, C.; Alard, A.; Aryankalayil, M.J.; Sarfraz, Y.; Diamond, J.M.; Schneider, R.J.; Inghirami, G.; Coleman, C.N.; Formenti, S.C.; Demaria, S. DNA exonuclease trex1 regulates radiotherapy-induced tumour immunogenicity. *Nat. Commun.* **2017**, *8*, 15618, doi:10.1038/ncomms15618.
59. Onishi, M.; Okonogi, N.; Oike, T.; Yoshimoto, Y.; Sato, H.; Suzuki, Y.; Kamada, T.; Nakano, T. High linear energy transfer carbon-ion irradiation increases the release of the immune mediator high mobility group box 1 from human cancer cells. *J. Radiat. Res.* **2018**, *59*, 541–546, doi:10.1093/jrr/rry049.
60. Mahadevan, L.S.K.; Sahoo, N.; Aliru, M.L.; Krishnan, S. Dependence of immunogenic modulation of tumor cells by proton radiation on the linear energy transfer. *Int. J. Radiat. Oncol.* **2017**, *99*, E607, doi:10.1016/j.ijrobp.2017.06.2061.
61. Kirilova, A.; Damyanovich, A.; Crook, J.; Jezioranski, J.; Wallace, K.; Pintilie, M. 3D MR-spectroscopic imaging assessment of metabolic activity in the prostate during the PSA “bounce” following 125Iodine brachytherapy. *Int. J. Radiat. Oncol. Biol. Phys.* **2011**, *79*, 371–378, doi:10.1016/j.ijrobp.2009.10.066.
62. Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G. Academia and clinic annals of internal medicine preferred reporting items for systematic reviews and meta-analyses : The PRISMA statement. *Ann. Intern. Med.* **2009**, *151*, 264–269.
63. Roever, L. PICO: Model for clinical questions. *Evid. Based Med. Pr.* **2018**, *3*, 1–2, doi:10.4172/2471-9919.1000115.
64. Amico, A.V.D.; Whittington, R.; Malkowicz, S.B.; Schultz, D.; Blank, K.; Broderick, G.A.; Tomaszewski, J.E.; Renshaw, A.A.; Kaplan, I.; Beard, C.J.; et al. Biochemical outcome after radical prostatectomy, external beam radiation therapy, or interstitial radiation therapy for clinically localized prostate cancer. *JAMA* **1998**, *280*, 969–974.
65. National Institute of Health. Study Quality Assessment Tools for Cases Series Studies. Available online: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools> (accessed on 1 April 2020).
66. Nyaga, V.N.; Arbyn, M.; Aerts, M. Metaprop: A stata command to perform meta-analysis of binomial data. *Arch. Publ. Health* **2014**, *72*, 1–10.
67. Wan, X.; Wang, W.; Liu, J.; Tong, T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med. Res. Methodol.* **2014**, *14*, 135, doi:10.1186/1471-2288-14-135.
68. Deeks, J.J.; Higgins, J.P.; Altman, D.G. Chapter 10: Analysing data and undertaking meta-analyses. In *Cochrane Handbook for Systematic Reviews of Interventions Version 6.0*; Higgins, J.P., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M., Welch, V., Eds.; Cochrane: London, UK, 2019.
69. Borenstein, M.; Hedges, L.V.; Higgins, J.P.T.; Rothstein, H.R. Meta-regression. In *Introduction to Meta-Analysis*; John Wiley & Sons: Chichester, West Sussex, UK, 2009.
70. Sanchez, J.; Dohoo, I.R.; Christensen, J.; Rajic, A. Factors influencing the prevalence of salmonella spp. in swine farms: A meta-analysis approach. *Prev. Vet. Med.* **2007**, *81*, 148–177, doi:10.1016/j.prevetmed.2007.04.005.
71. Keithlin, J.; Sargeant, J.; Thomas, M.K.; Fazil, A. Systematic review and meta-analysis of the proportion of campylobacter cases that develop chronic sequelae. *BMC Publ. Health* **2014**, *14*, 1203, doi:10.1186/1471-2458-14-1203.
72. Harbord, R.M.; Higgins, J.P.T. Meta-regression in stata. *Stata J.* **2008**, *8*, 493–519, doi:10.1177/1536867x0800800403.

