

# Corrigendum: Did the addition of concomitant chemotherapy to radiotherapy improve outcomes in hypopharyngeal cancer? A population-based study

S.F. Hall MSc MD\* and R. Griffiths BSc<sup>†</sup>

## Original citation

*Curr Oncol* 2016;23:266–72. (doi: <http://dx.doi.org/10.3747/co.23.3085>)

## Author Description of the Error and Correction

### **Current Text (p. 3, Results, ¶9)**

The LEDFS was 9.93% before 2000 (all patients treated with RT). When LEDFS was compared for patients in the post-2000 era who were specifically selected for each treatment, the LEDFS rates were similar (21.25% for RT, 25.87% for CCRT,  $p = 0.34$ ) and had improved from the previous era regardless of treatment.

### **Corrected Text**

The LEDFS was 37.8% before 2000 (all patients treated with RT). When LEDFS was compared for patients in the post-2000 era who were specifically selected for each treatment, the LEDFS rates were similar (43.75% for RT, 44.06% for CCRT,  $p = 0.96$ ) and had improved from the previous era regardless of treatment.

## CONFLICT OF INTEREST DISCLOSURES

We have read and understood *Current Oncology's* policy on disclosing conflicts of interest, and we declare that we have none.

## AUTHOR AFFILIATIONS

\*Department of Otolaryngology/Head and Neck Surgery and <sup>†</sup>ICES Queen's, Queen's University, Kingston, ON.