

Update on systemic therapy for advanced soft-tissue sarcoma

A. Smrke MD,* Y. Wang MD,* and C. Simmons MD MSc*

ABSTRACT

Background Soft-tissue sarcoma (STS) represents a rare group of mesenchymal neoplasms comprising more than 50 heterogeneous subtypes. Great efforts have been made to increase the understanding of the treatment of advanced STS (unresectable or metastatic disease). We set out to determine whether outcomes for patients with advanced STS have improved over time and to assess the current evidence for systemic therapy.

Methods In a scoping review, we evaluated the contemporary evidence for systemic treatment of advanced STS in adults (>18 years of age). Phase I, II, and III studies of systemic therapy for advanced STS published in the English language were included. After abstract and full-text review of seventy-seven studies, sixty-two trials met the inclusion criteria.

Results The number of clinical trials conducted and published in advanced STS has increased over the last 30 years. Although median overall survival has increased, attempts at improving first-line therapy through dose intensification, doublet chemotherapy, or alternative backbones have not been successful. The optimal therapy beyond anthracyclines remains a challenge, especially given the heterogeneity that grouping multiple STS subtypes within clinical trials creates. However, increasing numbers of agents are being studied, and several studies had shown isolated benefit in progression-free or overall survival.

Summary First-line systemic therapy with an anthracycline remains the standard of care for advanced STS. However, choice of subsequent therapy beyond anthracyclines remains challenging. Novel systemic therapies, use of molecular diagnostics to direct therapy, subtype-specific trials, and learnings from real-world retrospective data are all important for improving outcomes in patients with advanced STS.

Key Words Soft-tissue sarcoma, advanced; systemic therapy; anthracyclines

Curr Oncol. 2020 February;27(S1):25-33

www.current-oncology.com

INTRODUCTION

Soft-tissue sarcoma (STS) refers to a group of uncommon mesenchymal malignancies with more than 50 different subtypes¹. In Canada, STS represents approximately 1175 new cases per year (0.6% of all cancer diagnoses) in adults². The natural history and response to treatment between and within subtypes is heterogeneous. A large proportion of patients (14.5%–26.5%) present with *de novo* metastatic disease³, and 40%–50% of patients with localized disease will develop metastasis¹. Thus, patients with advanced STS, defined as those with unresectable or metastatic disease, represent a significant proportion of patients affected by STS.

Most published randomized trials in advanced STS include all subtypes of STS—the 5 most common histologic

subtypes being liposarcoma (LPS), leiomyosarcoma (LMS), undifferentiated pleomorphic sarcoma (UPS), fibrosarcoma, and synovial sarcoma⁴. The rarity of sarcoma—and the even further rarity of individual subtypes of STS—has limited the ability to conduct large, histology-specific clinical trials to inform practice. Thus, most of the literature concerning the treatment of specific subtypes has focused on retrospective case series.

Outcomes for patients were very poor before the discovery, in 1973, that doxorubicin is active against STS⁵, and anthracyclines have remained the backbone of standard-care treatment for advanced STS⁶. Since the 1970s, various trials have been conducted in hopes of improving patient outcomes or reducing adverse events through regimen intensification, non-anthracycline regimens, or use of alternative anthracyclines. In the present review, we aimed,

Correspondence to: Christine Simmons, Division of Medical Oncology, BC Cancer–Vancouver, 600 West 10th Avenue, Vancouver, British Columbia V5Z 1V6.
E-mail: christine.simmons@bccancer.bc.ca ■ DOI: <https://doi.org/10.3747/co.27.5475>

through literature appraisal, to determine whether survival for patients with STS has continued to improve since the 1970s and to identify the optimum contemporary systemic treatment pathway for patients with advanced STS.

METHODS

Search Strategy and Selection of Studies

A scoping review was conducted using a literature search in PubMed with the keywords “soft tissue sarcoma,” “metastatic,” “unresectable,” “systemic therapy,” “immunotherapy,” and “targeted agents” for 1987 to 15 May 2019. Specific searches based on drugs used in the treatment of STS were also performed. In addition, a search of ongoing clinical trials with the keywords “soft tissue sarcoma,” “metastatic/advanced,” and “systemic therapy” was performed at <https://ClinicalTrials.gov/>. We included studies published in the English language that involved adults (>18 years of age) diagnosed with advanced STS who were enrolled in prospective phase I, II, or III clinical trials and whose primary treatment was systemic therapy. Retrospective studies were excluded from the scoping review. Given well-established and distinctly different treatments, studies were excluded if the predominant histologic subtype was gastrointestinal stromal tumour, Ewing sarcoma, bone sarcoma (osteosarcoma, giant cell tumour of bone, chondrosarcoma), rhabdomyosarcoma, desmoid fibromatosis, and Kaposi sarcoma.

Data Extraction and Analysis

Titles were reviewed for relevance, and duplicates were removed. Abstract and full-text reviews of 152 studies were undertaken by AS, YW, and CS. Disagreements were resolved by consensus. Title, year published, trial type, number of patients, agents used, median progression-free survival (mPFS), median overall survival (mos), clinical benefit rate (complete response, plus partial response, plus stable disease) and response rate [RR (complete response plus partial response)] were extracted from sixty-two studies. Number of studies per year, trial type, line of therapy, and type of systemic therapy were coded. Of thirty-two active clinical trials found at <https://ClinicalTrials.gov/>, AS reviewed all of them, and four were included in the final review. Descriptive statistics were used to generate figures (Figure 1) in the Excel software application (version 16.16.9; Microsoft Corporation, Redmond WA, U.S.A.).

Treatment Choices

Included studies were grouped by line of therapy (first line vs. beyond first line) to generate a summary of the evidence. When a systemic therapy was compared with doxorubicin, the study was coded based on the non-doxorubicin arm.

RESULTS

Studies Found

One hundred fifty-two trials from 1987–2019 underwent abstract and full text review, with sixty-two trials being included in the scoping review. Most were phase II trials ($n = 35$), with fewer being phase III trials ($n = 15$) or phase I trials ($n = 4$); the remaining trials were of mixed or undefined




FIGURE 1 Consort diagram for the scoping review. STS = soft-tissue sarcoma; ES = Ewing sarcoma; RMS = rhabdomyosarcoma; GIST = gastrointestinal stromal tumor; DF = desmoid fibromatosis.

phase ($n = 6$), a meta-analysis ($n = 1$), or an undefined design ($n = 1$) [Figure 2(A)]. The number of published studies in this field has increased over time [Figure 2(B)], to 4–8 per year during 2012–2017 from 0–3 per year during 1987–2002. The types of systemic therapies studied for advanced STS have been increasing. The most commonly studied therapies included alkylating agents (ifosfamide, temozolomide, trabectedin), tyrosine kinase inhibitors, microtubule inhibitors (taxanes, eribulin), or anthracycline derivatives (epirubicin, liposomal doxorubicin, amrubicin) [Figure 2(C)]. Two studies—both published after 2017—focused on immunotherapy.

Outcomes

First Line

Compared with the mos of 7.7–12.0 months with first-line doxorubicin reported by the Cochrane review in 2003⁶, the mos has steadily improved, with the most recently reported




FIGURE 2 (A) Types of published study included ($n = 62$). (B) Trends in publication per year since 1987. (C) Class of systemic therapy used in the study. TKI = tyrosine kinase inhibitor.

mos being 20.4 months for doxorubicin in a doxorubicin-olaratumab phase III trial^{7,8} (Table I). By contrast, the mPFS has, overall, remained stable over time. Compared with doxorubicin, newer anthracyclines developed with the hopes of reducing toxicity have not demonstrated an improved mos^{18–22}. Furthermore, the RRs for comparisons with doxorubicin were all lower in older single-arm studies of gemcitabine (RR: 4%)²³, paclitaxel (RR: 12%)²⁴, perifosine (RR: 5%)²⁵, and temsirolimus (RR: 5%)²⁶. The RR for single-agent ifosfamide at a dose of 12 g/m² was 17%; however, that regimen was associated with significant grades 3 and 4 toxicities²⁷.

Table I summarizes modern first-line trials comparing doxorubicin with doublet or novel systemic therapy backbones. Compared with doxorubicin alone, dose escalation or doublet chemotherapy regimens that, in large phase III trials, attempted to intensify treatment have not improved the mos^{6,11,13}. Compared with doxorubicin alone, doublet therapy with doxorubicin–ifosfamide was associated with a modestly increased RR at the cost of a higher rate of febrile neutropenia, but no mos benefit^{6,28}. Furthermore, ifosfamide variations developed in the hope of reducing the side effects from metabolites have produced disappointing results^{29,30}.

The lone trial to demonstrate improved overall survival was the phase IB/II trial that compared olaratumab–doxorubicin with doxorubicin alone¹⁶. Unfortunately, that result was not replicated in a large phase III study⁸. With respect to anthracycline-sparing regimens, no difference in mPFS or mos for any subtype of STS was observed in a comparison of first-line gemcitabine–docetaxel with doxorubicin alone; however, quality of life was better for patients treated with doxorubicin alone¹⁷.

Beyond First Line

Upon disease progression, trials have attempted to exploit novel doublets or mechanisms of action (Table II), but mos and mPFS results have both remained modest. Only eribulin^{38,40}, gemcitabine–docetaxel³², and gemcitabine–dacarbazine³³ have been associated with a mos benefit in doxorubicin-treated patients. Specifically, compared with dacarbazine, eribulin^{38,40} demonstrated a benefit in mos, but not in mPFS, for patients with LMS or LPS—a result that was likely driven by the dedifferentiated LPS subgroup⁴⁰. Notably, a number of studies have compared novel agents with a backbone of dacarbazine, which showed activity in STS in a single-arm phase II study^{41,47}.

Interestingly, tyrosine kinase inhibitors seem to be associated with an improvement in mPFS, but not mos. A phase II trial of regorafenib^{37,39} and a phase III trial of pazopanib³⁴, both using a placebo comparator in non-adipocytic sarcomas, and a phase III trial of trabectedin compared with dacarbazine³⁶ in patients with LMS or LPS, all showed significantly prolonged mPFS (in the order of months), but not mos.

Immunotherapy

To date, two studies of immunotherapy have been conducted in patients with STS^{48,49,a}. The SARC028 study explored the use of pembrolizumab in patients with STS and bone sarcoma after up to 3 prior lines of therapy. The overall RR was 18% (7 of 40 patients), but was likely driven by patients having

TABLE I Selected randomized controlled trials for first-line treatment including doxorubicin

Reference	Study phase	Pts (n)	Study arms	RR (%)	CBR (%)	mPFS ^a (months)	mOS (months)
Bramwell <i>et al.</i> , 2003 ⁶	Systematic review	2281	Doxorubicin	16–27	NR	NR	7.7–12
Kalofonos <i>et al.</i> , 2004 ⁹	II	30	Doxorubicin 25 mg/m ² days 1–3 and cisplatin 100 mg/m ² day 1	16.7	70	6	11.5
Demetri <i>et al.</i> , 2012 ¹⁰	I/II	86	Doxorubicin 75 mg/m ² day 1	24	93	6.4	21.6
			Doxorubicin 75 mg/m ² and conatumumab 15 mg/kg day 1	20	72	5.6	18.2
Blay <i>et al.</i> , 2014 ¹¹	III ^b	121	Doxorubicin 75 mg/m ² day 1	45.9	86.5	8.3	27.3
			OR Doxorubicin 60 mg/m ² day 1 and ifosfamide 6–9 g/m ² day 1 (37%)				
			Trabectedin 1.5 mg/m ² day 1	37.3	82.4	18.8	38.9
Gelderblom <i>et al.</i> , 2014 ¹²	II	118	Doxorubicin 75 mg/m ² day 1	22.2	27.8	NR	NR
			Brostallicin 10 mg/m ² day 1	3.9	6.5	NR	NR
Judson <i>et al.</i> , 2014 ¹³	III	228	Doxorubicin 75 mg/m ² day 1	14	59.6	4.6	12.8
			Doxorubicin 75 mg/m ² day 1 and ifosfamide 10 g/m ² over days 1–4	26	76.6	7.4	14.3
Bui-Nguyen <i>et al.</i> , 2015 ¹⁴	IIB	133	Doxorubicin 75 mg/m ² day 1	25.6	62.8	5.5	NR
			Trabectedin 1.3 mg/m ² day 1 (3 h)	14.8	55.3	2.8	NR
			Trabectedin 1.5 mg/m ² day 1 (24 h)	4.7	62.8	3.1	NR
Martin-Broto 2016 ¹⁵	II	115	Doxorubicin 75 mg/m ² day 1	NR	NR	5.5 ^c	NR
			Doxorubicin 75 mg/m ² day 1 and trabectedin 1.1 mg/m ² day 1	NR	NR	5.7	NR
Tap <i>et al.</i> , 2016 ¹⁶	II	148	Doxorubicin 75 mg/m ² day 1	11.9	62.7	4.1	26.5 ^d
			Doxorubicin 75 mg/m ² day 1 and olaratumab 15 mg/kg days 1, 8	18.2	77.3	6.6	14.7 ^d
Seddon <i>et al.</i> , 2017 ¹⁷	III	256	Doxorubicin 75 mg/m ² day 1	19	72	5.4	17.6
			Gemcitabine 675 mg/m ² days 1, 8 and docetaxel 75 mg/m ² day 1	20	75	5.4	15.5 ^e
Tap <i>et al.</i> , 2019 ⁷ Eli Lilly and Company ⁸	III	485	Doxorubicin 75 mg/m ² day 1	NA	NA	6.8	20.4
			Doxorubicin 75 mg/m ² day 1 and olaratumab 15 mg/kg days 1 and 8	NA	NA	5.4	19.7

^a Included studies did not consistently report PFS, and PFS was not included in the final outcomes for the systematic review.

^b Only translocation-related sarcoma, mostly myxoid or round-cell liposarcoma.

^c Stopped for fertility.

^d Statistically significant.

^e Reduced quality of life reported in patients receiving docetaxel–gemcitabine.

Pts = patients; RR = response rate; CBR = clinical benefit rate; mPFS = median progression-free survival; mOS = median overall survival; NR = not reported; NA = not available at time of publication.

the STS subtypes UPS (RR: 40%; 4 of 10) and LPS (RR: 20%; 2 of 10)⁴⁸. To further explore those subtypes, 30 additional patients have been enrolled onto each arm, with results awaited (NCT02301039 at <https://ClinicalTrials.gov/>). Nivolumab as a single agent demonstrated a modest 5% RR⁴⁹. However, the combination of ipilimumab–nivolumab in patients having received at least 1 prior line of therapy demonstrated a 16% RR (6 of 38 patients: 1 uterine LMS, 1 non-uterine LMS, 1 myxofibrosarcoma, 2 UPS, 1 angiosarcoma)⁴⁹.

DISCUSSION

Despite an increase in the number of studies and systemic therapies developed, doxorubicin-based chemotherapy remains the backbone of first-line systemic treatment for advanced STS. Although the mOS is similar with first-line gemcitabine–docetaxel and with doxorubicin alone, patients receiving the combination therapy experience reduced quality of life¹⁷, making gemcitabine–docetaxel a less attractive first-line option. Otherwise, first-line doublet

TABLE II Selected randomized controlled trials beyond first line treatment

Reference	Study phase	Pts (n)	Study arms	RR (%)	CBR (%)	mPFS (months)	mOS (months)
<i>Comparative trials</i>							
Bramwell et al., 1987 ³¹	II	171	Cyclophosphamide 1.5 g/m ²	7.5	41.7	NR	NR
			Ifosfamide 5 g/m ²	17.6	57.3	NR	NR
Maki et al., 2007 ³²	II	122	Gemcitabine 1200 mg/m ² days 1, 8	8	27	3	11.5 ^a
			Gemcitabine 1200 mg/m ² days 1, 8 and docetaxel 100 mg/m ² day 1	17	32	6.2	17.9 ^a
García-del-Muro et al., 2011 ³³	II	113	Dacarbazine 1800 mg/m ² every 3 weeks	4	25	2 ^a	8.2 ^a
			Dacarbazine 500 mg/m ² day 1 and gemcitabine 1800 g/m ² day 1 every 2 weeks	12	49	4.2 ^a	16.8 ^a
Van der Graaf et al., 2012 ³⁴	III	372	Placebo (no crossover)	0	38	1.6 ^a	10.7
			Oral pazopanib 800 mg daily	6	73	4.6 ^a	12.5
Blay et al., 2015 ³⁵	III	355	Cisplatin 75 mg/m ² day 1	1	36	1.41	9.33
			Cisplatin 75 mg/m ² and ombrabulin 25 mg/m ² day 1	4	47	1.54	11.43
Demetri et al., 2016 ³⁶	III	518 ^b	Dacarbazine 1000 mg/m ² day 1	6.9	19	1.5 ^a	12.4
			Trabectedin 1.5 mg/m ² day 1	9.9	34	4.2 ^a	12.9
Mir et al., 2016 ³⁷	II	43 LPS	Placebo (could cross over)	0	43	1.7	8.8
			Oral regorafenib 160 mg daily for 21 days; 7 days off	0	57	1.1	4.7
		56 LMS	Placebo (could cross over)	4	58	1.8	9.1
			Oral regorafenib 160 mg daily for 21 days; 7 days off	0	86	3.7	21
		27 SyS	Placebo (could cross over)	0	22	1.0	6.7
			Oral regorafenib 160 mg daily for 21 days; 7 days off	8	85	5.6	13.4
		56 Other	Placebo (could cross over)	0	34	1.0	9.5
			Oral regorafenib 160 mg daily for 21 days; 7 days off	11	78	2.9	12.1
Schöffski et al., 2016 ³⁸	III	452 ^b	Dacarbazine 850–1200 mg/m ² day 1	5	48	2.6	11.5 ^a
			Eribulin 1.4 mg/m ² days 1, 8	4	46	2.6	13.5 ^a
Berry et al., 2017 ³⁹	II	60 ^c	Placebo (could cross over)	NR	NR	1.0 ^a	9.0
			Oral regorafenib 160 mg daily for 21 days; 7 days off	NR	NR	4.0 ^a	13.4
Demetri et al., 2017 ⁴⁰	III	143 ^d	Dacarbazine 850–1200 mg/m ² day 1	0	44.4	1.7	8.4 ^a
			Eribulin 1.4 mg/m ² days 1, 8	0.4	65.2	2.9	15.6 ^a
<i>Selected single-agent trials</i>							
Buesa et al., 1991 ⁴¹	II	44	Dacarbazine 1200 mg/m ² day 1 every 3 weeks	18	36.3	NR	NR
Nielsen 2000 ²⁷	II	124	Ifosfamide 12 g/m ²	16	48	3.5	12.7
Talbot et al., 2003 ⁴²	II	26	Temozolomide	15.4	26.9	2	13.2
George et al., 2009 ⁴³	II	48	Oral sunitinib 37.5 mg daily	2	14	NR	NR
Maki et al., 2009 ⁴⁴	II	144	Oral sorafenib 400 mg twice daily	6	68	3.2	14.3 ^e
Luo et al., 2015 ⁴⁵	II	26	Gemcitabine 1000 mg/m ² days 1, 8; vincristine 1.4 mg/m ² day 1; and cisplatin 25 mg/m ² days 1–3	23	65.3	4.8	15
Subbiah et al., 2018 ⁴⁶	IB/II	25	Pazopanib 800 mg and oral trametinib 2 mg daily	8	56	2.27	9

^a Statistically significant.^b Liposarcoma or leiomyosarcoma only^c Included only non-adipocytic sarcoma, unplanned analysis.^d Planned subgroup analysis of liposarcoma cohort in Demetri et al., 2016³⁶.^e Mostly driven by the angiosarcoma cohort.

RR = response rate; CBR = clinical benefit rate; mPFS = median progression-free survival; mOS = median overall survival; NR = not reported; LPS = liposarcoma; LMS = leiomyosarcoma; SyS = synovial sarcoma.

chemotherapy has generally been shown to increase the RR, but not the mPFS or MOS⁶ (Table 1). The phase IB/II results for olaratumab–doxorubicin¹⁶ initially suggested an impressive MOS benefit, resulting in accelerated conditional approval; however, the larger phase III study of that combination compared with doxorubicin alone was negative⁷. The difference in MOS might have been attributable to overperformance in the doxorubicin-alone arm or to the proportions of the heterogeneous STS population captured in each study being different. Which, if any, subtype of STS might respond best to the combination remains an open question.

However, the foregoing results do reinforce the trend seen in Table 1 that the MOS for first-line doxorubicin-only treatment in STS is indeed increasing over time, but that the RR and progression-free survival have generally remained stagnant. Those observations might be attributable to additional evidence and to an increase in the agents available for second-line therapy, coupled with a better understanding of subtype-directed therapy⁵⁰. Furthermore, supportive care in oncology has improved over time, which could account for improved outcomes such as a quality of life⁵¹.

Although the present review focuses on systemic therapy, patients with advanced STS receive multidisciplinary care from radiation and surgical oncologists. More frequent use of metastasectomy and radiation (specifically, stereotactic ablative radiation therapy) in carefully selected patients with a long disease-free interval or long-interval disease stability might also explain improved outcomes for patients over time^{52,53}.

Single-agent doxorubicin remains the agent of choice in the first-line treatment of most advanced STS; however, in carefully selected patients (such as those with borderline resectable tumours, rapid tumour growth, and a deteriorating performance status or a tumour near critical anatomic structures), doxorubicin–ifosfamide might be considered as first-line treatment because of its higher RR (Figure 3). In patients with cardiac comorbidities, clinicians could consider using liposomal doxorubicin or gemcitabine–docetaxel in the first line to avoid potential worsening of underlying cardiac dysfunction.

After treatment with anthracyclines, subtype becomes an important consideration. Based on our scoping review, we propose an algorithm (Figure 3). Outside the scope of our review, subtype-directed therapy for rare histologies such as alveolar soft-part sarcoma, perivascular epithelioid-cell neoplasms, and angiosarcoma is well summarized in other published review articles⁵⁰. For patients with LMS (and to a greater extent LPS), eribulin is an active therapy that improves MOS^{38,40}. Although activity of gemcitabine–docetaxel was initially reported in a phase II trial that included only patients with LMS⁵⁴, a preplanned subgroup analysis from the GEDDIS trial¹⁷ showed no evidence of a differential benefit based on subtype, including uterine compared with non-uterine LMS compared with other histologies. Thus, gemcitabine–docetaxel is an established treatment for patients with multiple subtypes of STS after failure of anthracyclines. The smaller phase II study of dacarbazine–gemcitabine in previously treated patients




FIGURE 3 Proposed treatment algorithm for advanced soft-tissue sarcoma (STS). ^aDacarbazine, gemcitabine, pazopanib, trabectedin. NGS = next-generation sequencing; PS = performance status; LMS = leiomyosarcoma; LPS = liposarcoma; MDC = multidisciplinary conference.

with STS showed a significant MOS benefit, but the primary outcome of the study was the progression-free survival rate at 3 months. Given that the larger phase III study of gemcitabine–docetaxel represents a higher level of evidence, we favour use of that gemcitabine doublet unless the patient has a contraindication to taxanes. Although randomized data are limited, dacarbazine is often included as a backbone for systemic therapy beyond anthracyclines^{33,36,38,40} and can be considered an option for patients who are fit enough to receive systemic therapy after doublets and for whom subtype-specific therapy has been exhausted.

Delaying the need to switch therapies is an important outcome, but must be balanced when a switch does not translate into a change in MOS. Pazopanib³⁴, regorafenib³⁹, and trabectedin³⁶ are associated only with improved mPFS, not MOS. Although establishing a large trial of heterogeneous STS is challenging, the question of whether those therapies might benefit a specific subgroup remains open.

A large retrospective analysis from the European Organisation for Research and Treatment of Cancer suggested that, in patients with STS, lack of progression is an important predictor of MOS⁵⁵. However, that hypothesis has not held true in large phase III studies that have shown improvement in mPFS, but not MOS^{34,36}—perhaps because of difficulty in interpreting imaging for patients with STS. The size of tumours is often non-uniform, which makes interpretation according to the Response Evaluation Criteria in Solid Tumours difficult. Similarly, response might be better reflected by qualitative change in imaging rather than by size alone⁵⁶. Furthermore, mPFS is critically affected by the design of follow-up imaging, which can influence results⁵⁷.

Another interesting outcome to consider with pazopanib, regorafenib, and trabectedin might be time to second progression, which might help with the sequencing of available agents and represent a clinically meaningful outcome for patients.

Looking beyond cytotoxic or targeted therapy, novel mechanisms of systemic therapy such as immunotherapy have changed the landscape of treatment for a variety of tumour groups⁵⁸. Although early results suggest that immunotherapy could be effective in certain STS subtypes such as UPS and LPS^{48,49}, larger cohorts with longer follow-up are required to better understand the true clinical benefit of those agents for patients with STS.

The rarity of STS and its subtypes generally precludes large subtype-specific randomized trials. However, within the STS subtypes, heterogeneity can be seen in terms of grade, chemoresponsiveness, and prognosis. That heterogeneity often cannot be adequately teased out in large STS trials that group multiple STS histologies having potentially dissimilar biology⁵⁰. Despite histologic heterogeneity, RRS might vary, potentially contributing to the differing results seen in the phase II compared with the phase III trial of olaratumab–doxorubicin⁸. Inclusion criteria such as disease progression within 3 months of enrolment, grade, subtype, and mandated central pathology review should all be carefully considered in attempting to reduce heterogeneity. Alternatively, subtype-specific trials have been successful in the past^{38,59} and could provide robust evidence to inform practice and remove the inherent heterogeneity when many STS subtypes are included in a trial.

Currently, clinicians might be more comfortable applying the results of large, randomized STS trials to patients with common STS histologies such as LPS or LMS; for uncommon subtypes, uncertainty remains, highlighting the continued importance of real-world data from retrospective cohorts of patients with rare STS subtypes as the best evidence to direct therapy. Active trials are generally focusing on small molecules in selected sarcoma subtypes^{56–58}. The focus on subtype-directed therapy has led to the emergence of platforms such as NanoString fusion assays (NanoString Technologies, Seattle, WA, U.S.A.), which can help to confirm the STS subtype, potentially helping to tailor systemic treatment⁶⁰. For example, *TRK* fusions are present in at least 0.4%⁶¹ of the unselected population with sarcoma, but are pathognomonic for infantile fibrosarcoma⁶². Larotrectinib, a potent pan-*TRK* inhibitor has been associated with an impressive 75% RR in adults and children with *TRK* fusions⁶³.

Currently, sequencing tumours in an attempt to identify driver mutations and to personalize care is a growing trend in all tumour types. Sequencing is being done quite frequently in sarcoma as well, although the somatic mutation burden in STS is low relative to that in other tumours⁶⁴. Next-generation sequencing studies in patients with STS show a 41% actionable mutation rate⁶⁵. Importantly, to better understand whether tumour-directed therapy will translate to improved MOS, a number of basket trials matching known actionable mutations to active systemic therapy are enrolling patients with various tumour types (see NCT03297606 and NCT02534649 at <https://ClinicalTrials.gov/>).

We performed a broad scoping review with the aim to include all appropriate and relevant studies. It is important, however, to recognize that our methods do not represent a complete systematic review of the literature. For example, some studies might have been missed, because we included only studies published in English. However, efforts were undertaken to ensure inclusion of key studies for the treatment of advanced STS. Our review focused on randomized or prospective data collection; results from retrospective cohorts of STS patients or specific subtypes were not included.

SUMMARY

Although the number of published studies has increased over time, the most important systemic treatment for advanced STS remains anthracycline-based therapy. For patients treated with first-line doxorubicin, survival has steadily increased over time. However, the selection of optimal systemic therapy after anthracycline-containing regimens remains a challenge, with few agents showing survival benefit. Ongoing studies in immunotherapy, novel chemotherapy combinations, molecular diagnostics, and targeted agents aim to further improve outcomes.

CONFLICT OF INTEREST DISCLOSURES

We have read and understood *Current Oncology's* policy on disclosing conflicts of interest, and we declare that we have none.

AUTHOR AFFILIATIONS

*BC Cancer–Vancouver Centre, Vancouver, BC.

REFERENCES

- Fletcher CDM, Bridge JA, Hogendoorn P, Mertens F. *WHO Classification of Tumors of Soft Tissue and Bone*. 4th ed. Geneva, Switzerland: World Health Organization; 2013.
- Canadian Cancer Society's Advisory Committee on Cancer Statistics. *Canadian Cancer Statistics 2015*. Toronto, ON: Canadian Cancer Society; 2015.
- Italiano A, Mathoulin-Pelissier S, Cesne AL, *et al.* Trends in survival for patients with metastatic soft-tissue sarcoma. *Cancer* 2011;117:1049–54.
- Brennan MF, Antonescu CR, Maki RG. *Management of Soft Tissue Sarcoma*. New York, NY: Springer; 2013.
- Tan C, Etcubanas E, Wollner N, *et al.* Adriamycin—an anti-tumour antibiotic in the treatment of neoplastic diseases. *Cancer* 1973;32:9–17.
- Bramwell V, Anderson D, Charette ML on behalf of the Sarcoma Disease Site Group. Doxorubicin-based chemotherapy for the palliative treatment of adult patients with locally advanced or metastatic soft tissue sarcoma. *Cochrane Database Syst Rev* 2003;:CD003293.
- Tap WD, Wagner AJ, Papai Z, *et al.* ANNOUNCE: a randomized, placebo (PBO)–controlled, double-blind, phase (Ph) III trial of doxorubicin (dox) + olaratumab versus dox + PBO in patients (pts) with advanced soft tissue sarcomas (STS) [abstract LBA3]. *J Clin Oncol* 2019;37:. [Available online at: https://ascopubs.org/doi/abs/10.1200/JCO.2019.37.18_suppl.LBA3?af=R; cited 13 September 2019]
- Eli Lilly and Company. Lilly reports results of phase 3 soft tissue sarcoma study of LARTRUVO® [news release]. Indianapolis, IL: Eli Lilly and Company; 2019.
- Kalofonos HP, Bafaloukos D, Kourelis TG, *et al.* Adriamycin and cis-platinum as first-line treatment in unresectable locally advanced or metastatic adult soft-tissue sarcomas. *Am J Clin Oncol* 2004;27:307–11.
- Demetri GD, Le Cesne A, Chawla SP, *et al.* First-line treatment of metastatic or locally advanced unresectable soft tissue sarcomas with conatumumab in combination with doxorubicin or doxorubicin alone: a phase I/II open-label and double-blind study. *Eur J Cancer* 2012;48:547–63.
- Blay JY, Leahy MG, Nguyen BB, *et al.* Randomised phase III trial of trabectedin versus doxorubicin-based chemotherapy as first-line therapy in translocation-related sarcomas. *Eur J Cancer* 2014;50:1137–47.
- Gelderblom H, Blay J, Seddon B, *et al.* Brostallicin versus doxorubicin as first-line chemotherapy in patients with advanced or metastatic soft tissue sarcoma: an European Organisation for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group randomised phase II and pharmacogenetic study. *Eur J Cancer* 2014;50:388–96.
- Judson I, Verweij J, Gelderblom H, *et al.* on behalf of the European Organisation and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. Doxorubicin alone versus intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic soft-tissue sarcoma: a randomised controlled phase 3 trial. *Lancet Oncol* 2014;15:415–23.
- Bui-Nguyen B, Butrynski JE, Penel N, *et al.* on behalf of the European Organisation for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group and the Sarcoma Alliance for Research Through Collaboration. A phase IIb multicentre study comparing the efficacy of trabectedin to doxorubicin in patients with advanced or metastatic untreated soft tissue sarcoma: the TRUSTS trial. *Eur J Cancer* 2015;51:1312–20.
- Martin-Broto J, Pousa AL, de las Peñas R, *et al.* Randomized phase II study of trabectedin and doxorubicin compared with doxorubicin alone as first-line treatment in patients with advanced soft tissue sarcomas: a Spanish Group for Research on Sarcoma study. *J Clin Oncol* 2016;34:2294–302.
- Tap WD, Jones RL, Van Tine BA, *et al.* Olaratumab and doxorubicin versus doxorubicin alone for treatment of soft-tissue sarcoma: an open-label phase 1b and randomised phase 2 trial. *Lancet* 2016;388:488–97.
- Seddon B, Strauss SJ, Whelan J, *et al.* Gemcitabine and docetaxel versus doxorubicin as first-line treatment in previously untreated advanced unresectable or metastatic soft-tissue sarcomas (GeDDis): a randomised controlled phase 3 trial. *Lancet Oncol* 2017;18:1397–410.
- Mouridsen H, Bastholt L, Somers R, *et al.* Adriamycin versus epirubicin in advanced soft tissue sarcomas. A randomized phase II/phase III study of the EORTC Soft Tissue and Bone Sarcoma Group. *Eur J Cancer Clin Oncol* 1987;23:1477–83.
- Nielsen OS, Dombernowsky P, Mouridsen H, *et al.* High-dose epirubicin is not an alternative to standard-dose doxorubicin in the treatment of advanced soft tissue sarcomas. A study of the EORTC Soft Tissue and Bone Sarcoma Group. *Br J Cancer* 1998;78:1634–9.
- Gupta S, Gouw L, Wright J, *et al.* Phase II study of amrubicin (SM-5887), a synthetic 9-aminoanthracycline, as first line treatment in patients with metastatic or unresectable soft tissue sarcoma: durable response in myxoid liposarcoma with TLS–CHOP translocation. *Invest New Drugs* 2016;34:243–52.
- Chawla SP, Papai Z, Mukhametshina G, *et al.* First-line aldodoxorubicin vs doxorubicin in metastatic or locally advanced unresectable soft-tissue sarcoma: a phase 2b randomized clinical trial. *JAMA Oncol* 2015;1:1272–80.
- Judson I, Radford JA, Harris M, *et al.* Randomised phase II trial of pegylated liposomal doxorubicin (DOXIL/CAELYX) versus doxorubicin in the treatment of advanced or metastatic soft tissue sarcoma: a study by the EORTC Soft Tissue and Bone Sarcoma Group. *Eur J Cancer* 2001;37:870–7.
- Okuno S, Ryan LM, Edmonson JH, Priebat DA, Blum RH. Phase II trial of gemcitabine in patients with advanced sarcomas (E1797): a trial of the Eastern Cooperative Oncology Group. *Cancer* 2003;97:1969–73.
- Balcerzak SP, Benedetti J, Weiss GR, Natale RB. A phase II trial of paclitaxel in patients with advanced soft tissue sarcomas. A Southwest Oncology Group study. *Cancer* 1995;76:2248–52.
- Bailey HH, Mahoney MR, Ettinger DS, *et al.* Phase II study of daily oral perifosine in patients with advanced soft tissue sarcoma. *Cancer* 2006;107:2462–7.
- Okuno S, Bailey H, Mahoney MR, *et al.* A phase 2 study of temsirolimus (CCI-779) in patients with soft tissue sarcomas: a study of the Mayo Phase 2 Consortium (P2C). *Cancer* 2011;117:3468–75.
- Nielsen OS, Judson I, van Hoesel Q, *et al.* Effect of high-dose ifosfamide in advanced soft tissue sarcomas. A multicentre phase II study of the EORTC Soft Tissue and Bone Sarcoma Group. *Eur J Cancer* 2000;36:61–7.
- Frezza AM, Stacchiotti S, Gronchi A. Systemic treatment in advanced soft tissue sarcoma: what is standard, what is new. *BMC Med* 2017;15:109.
- Ryan CW, Merimsky O, Agulnik M, *et al.* PICASSO III: a phase III, placebo-controlled study of doxorubicin with or without palifosfamide in patients with metastatic soft tissue sarcoma. *J Clin Oncol* 2016;34:3898–905.
- Tap WD, Papai Z, Van Tine BA, *et al.* Doxorubicin plus evofosfamide versus doxorubicin alone in locally advanced, unresectable or metastatic soft-tissue sarcoma (TH CR-406/SARC021): an international, multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2017;18:1089–103.
- Bramwell VH, Mouridsen HT, Santoro A, *et al.* Cyclophosphamide versus ifosfamide: final report of a randomized phase II trial in adult soft tissue sarcomas. *Eur J Cancer Clin Oncol* 1987;23:311–21.

32. Maki RG, Wathen JK, Patel SR, *et al.* Randomized phase II study of gemcitabine and docetaxel compared with gemcitabine alone in patients with metastatic soft tissue sarcomas: results of Sarcoma Alliance for Research Through Collaboration study 002 [corrected]. *J Clin Oncol* 2007;25:2755–63.
33. García-del-Muro X, López-Pousa A, Maurel J, *et al.* on behalf of the Spanish Group for Research on Sarcomas. Randomized phase II study comparing gemcitabine plus dacarbazine versus dacarbazine alone in patients with previously treated soft tissue sarcoma: a Spanish Group for Research on Sarcomas study. *J Clin Oncol* 2011;29:2528–33.
34. Van der Graaf WT, Blay JY, Chawla SP, *et al.* on behalf of the EORTC Soft Tissue and Bone Sarcoma Group and the PALETTE Study Group. Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet* 2012;379:1879–86.
35. Blay JY, Papai Z, Tolcher AW, *et al.* Ombrabulin plus cisplatin versus placebo plus cisplatin in patients with advanced soft-tissue sarcomas after failure of anthracycline and ifosfamide chemotherapy: a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2015;16:531–40.
36. Demetri GD, von Mehren M, Jones RL, *et al.* Efficacy and safety of trabectedin or dacarbazine for metastatic liposarcoma or leiomyosarcoma after failure of conventional chemotherapy: results of a phase III randomized multicenter clinical trial. *J Clin Oncol* 2016;34:786–93.
37. Mir O, Brodowicz T, Italiano A, *et al.* Safety and efficacy of regorafenib in patients with advanced soft tissue sarcoma (REGOSARC): a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet Oncol* 2016;17:1732–42.
38. Schöffski P, Chawla S, Maki RG, *et al.* Eribulin versus dacarbazine in previously treated patients with advanced liposarcoma or leiomyosarcoma: a randomised, open-label, multicentre, phase 3 trial. *Lancet* 2016;387:1629–37.
39. Berry V, Basson L, Bogart E, *et al.* REGOSARC: regorafenib versus placebo in doxorubicin-refractory soft-tissue sarcoma—a quality-adjusted time without symptoms of progression or toxicity analysis. *Cancer* 2017;123:2294–302.
40. Demetri GD, Schöffski P, Grignani G, *et al.* Activity of eribulin in patients with advanced liposarcoma demonstrated in a subgroup analysis from a randomized phase III study of eribulin versus dacarbazine. *J Clin Oncol* 2017;35:3433–9.
41. Buesa J, Mouridsen H, van Oosterom A, *et al.* High-dose DTIC in advanced soft-tissue sarcomas in the adult: a phase II study of the EORTC Soft Tissue and Bone Sarcoma Group. *Ann Oncol* 1991;2:307–9.
42. Talbot SM, Keohan ML, Hesdorffer M, *et al.* A phase II trial of temozolomide in patients with unresectable or metastatic soft tissue sarcoma. *Cancer* 2003;98:1942–6.
43. George S, Merriam P, Maki RG, *et al.* Multicenter phase II trial of sunitinib in the treatment of nongastrointestinal stromal tumor sarcomas. *J Clin Oncol* 2009;27:3154–60.
44. Maki RG, D'Adamo DR, Keohan ML, *et al.* Phase II study of sorafenib in patients with metastatic or recurrent sarcomas. *J Clin Oncol* 2009;27:3133–40.
45. Luo Z, Zhang X, Peng W, *et al.* A phase II study of gemcitabine, vincristine, and cisplatin (GVP) as second-line treatment for patients with advanced soft tissue sarcoma. *Medicine (Baltimore)* 2015;94:e1777.
46. Subbiah V, Meyer C, Zinner R, *et al.* Phase IB/II study of the safety and efficacy of combination therapy with multikinase VEGF inhibitor pazopanib and MEK inhibitor trametinib in advanced soft tissue sarcoma. *Clin Cancer Res* 2017;23:4027–34. [Erratum in: *Clin Cancer Res* 2018;24:500–1]
47. Casali P, Abecassis N, Bauer S, *et al.* on behalf of the ESMO Guidelines Committee and EURACAN. Soft tissue and visceral sarcomas: ESMO–EURACAN clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2018;29(suppl 4):iv51–67. [Erratum in: *Ann Oncol* 2018;29(suppl 4):iv268–9]
48. Tawbi HA, Burgess M, Bolejack V, *et al.* Pembrolizumab in advanced soft-tissue sarcoma and bone sarcoma (SARC028): a multicentre, two-cohort, single-arm, open-label, phase 2 trial. *Lancet Oncol* 2017;18:1493–501.
49. D'Angelo SP, Mahoney MR, Van Tine BA, *et al.* Nivolumab with or without ipilimumab treatment for metastatic sarcoma (Alliance A091401): two open-label, non-comparative, randomised, phase 2 trials. *Lancet Oncol* 2018;19:416–26.
50. Katz D, Palmerini E, Pollack SM. More than 50 subtypes of soft tissue sarcoma: paving the path for histology-driven treatments. *Am Soc Clin Oncol Educ Book* 2018;38:925–38.
51. Temel JS, Greer JA, Muzikansky A, *et al.* Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733–42.
52. Palma DA, Olson R, Harrow S, *et al.* Stereotactic ablative radiotherapy versus standard of care palliative treatment in patients with oligometastatic cancers (SABR-COMET): a randomised, phase 2, open-label trial. *Lancet* 2019;393:2051–8.
53. Chudgar NP, Brennan MF, Munhoz RR, *et al.* Pulmonary metastasectomy with therapeutic intent for soft-tissue sarcoma. *J Thorac Cardiovasc Surg* 2017;154:319–30.e1.
54. Hensley ML, Maki R, Venkatraman E, *et al.* Gemcitabine and docetaxel in patients with unresectable leiomyosarcoma: results of a phase II trial. *J Clin Oncol* 2002;20:2824–31.
55. Grünwald V, Litière S, Young R, *et al.* Absence of progression, not extent of tumour shrinkage, defines prognosis in soft-tissue sarcoma—an analysis of the EORTC 62012 study of the EORTC STBSG. *Eur J Cancer* 2016;64:44–51.
56. Choi H. Role of imaging in response assessment and individualised treatment for sarcomas. *Clin Oncol (R Coll Radiol)* 2017;29:481–8.
57. Wilkerson J, Fojo T. Progression-free survival is simply a measure of a drug's effect while administered and is not a surrogate for overall survival. *Cancer J* 2009;15:379–85.
58. Drake CG, Lipson EJ, Brahmer JR. Breathing new life into immunotherapy: review of melanoma, lung and kidney cancer. *Nat Rev Clin Oncol* 2014;11:24–37.
59. Penel N, Bui BN, Bay JO, *et al.* Phase II trial of weekly paclitaxel for unresectable angiosarcoma: the ANGIOTAX study. *J Clin Oncol* 2008;26:5269–74.
60. Chang KT, Goytain A, Tucker T, *et al.* Development and evaluation of a pan-sarcoma fusion gene detection assay using the NanoString nCounter platform. *J Mol Diagn* 2018;20:63–77.
61. Gatalica Z, Xiu J, Swensen J, Vranic S. Molecular characterization of cancers with *NTRK* gene fusions. *Mod Pathol* 2019;32:147–53.
62. Knezevich SR, McFadden DE, Tao W, Lim JF, Sorensen PH. A novel *ETV6–NTRK3* gene fusion in congenital fibrosarcoma. *Nat Genet* 1998;18:184–7.
63. Drilon A, Laetsch TW, Kummar S, *et al.* Efficacy of larotrectinib in *TRK* fusion–positive cancers in adults and children. *N Engl J Med* 2018;378:731–9.
64. Cancer Genome Atlas Research Network. Comprehensive and integrated genomic characterization of adult soft tissue sarcomas. *Cell* 2017;171:950–65.
65. Lucchesi C, Khalifa E, Laizet Y, *et al.* Targetable alterations in adult patients with soft-tissue sarcomas: insights for personalized therapy. *JAMA Oncol* 2018;4:1398–404.