

Individual-Level Behavioral Dataset Linking Trace Eyeblink Conditioning, Contextual Fear Memory, and Home-Cage Activities in rTg4510 and Wild-Type Mice with Doxycycline Treatment

Ryo Kachi, Takuma Nishijo and Yasushi Kishimoto * 

Laboratory of Physical Chemistry, Faculty of Pharmaceutical Sciences, Teikyo University, Tokyo 173-8605, Japan; 20y101209tv@stu.teikyo-u.ac.jp (R.K.); nishijou.takuma.zm@teikyo-u.ac.jp (T.N.)

* Correspondence: y-kishimoto@pharm.teikyo-u.ac.jp; Tel.: +81-3-3964-8039

Abstract

This dataset provides synchronized multimodal behavioral measurements from 36 mice across four experimental groups: wild-type and rTg4510 tauopathy mice, each tested with or without doxycycline-mediated suppression of mutant tau expression. Of these, 34 mice had complete measurements across all three behavioral paradigms and were used for analyses requiring full cross-task linkage. At six months of age, all animals underwent three standardized behavioral paradigms: home cage monitoring, ten-day trace eyeblink conditioning, and contextual fear conditioning. The individual-level data included locomotor activity, rearing duration, conditioned response metrics, eyelid closure latencies, and contextual freezing percentages. All measurements were linked using unique mouse identifiers, enabling cross-task analysis without preprocessing or imputation. The dataset was accompanied by a complete data dictionary, processing workflow diagram, and validation analyses demonstrating cross-paradigm correlations. The cross-task associations are illustrated in the main figures, with additional early phase acquisition and temporal processing correlations provided in the main figures. Provided in an open CSV format with detailed metadata, this resource supports behavioral phenotyping, machine learning applications, and the investigation of learning mechanisms in tauopathy models.

Dataset: Deposited in Zenodo (DOI: 10.5281/zenodo.17864814).

Dataset License: CC BY 4.0

Keywords: rTg4510; tauopathy; doxycycline; trace eyeblink conditioning; contextual fear conditioning; spontaneous activity; cross-task analysis; behavioral phenotyping; mouse model



Academic Editor: Juraj Gregaň

Received: 11 December 2025

Revised: 9 February 2026

Accepted: 12 February 2026

Published: 16 February 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

1. Summary

The rTg4510 mouse is one of the most widely used tauopathy models and is characterized by the forebrain expression of mutant human P301L tau under the control of a tetracycline-responsive promoter. Extensive characterization using spatial learning tasks has demonstrated progressive neurofibrillary tangle formation, neuronal loss, and memory impairment. These phenotypes can be modulated by doxycycline (DOX)-mediated suppression of tau expression [1,2].

We recently reported that rTg4510 mice exhibit deficits in non-spatial associative learning, including trace eyeblink conditioning (tEBC) and contextual fear memory, and that DOX treatment partially prevents these impairments [3]. However, that study focused on group-level comparisons and individual-level data enabling cross-task correlation analyses were not provided. Such individual-level integration is crucial for investigating behavioral heterogeneity and response variability in therapeutic interventions. Although nonspatial learning deficits have been characterized in other AD mouse models [4,5], publicly available datasets linking multiple learning domains at the individual level remain scarce.

The present data descriptor addresses this gap by providing complete individual-level behavioral measurements from 36 mice across four experimental groups: wild-type (WT) and rTg4510 mice, with or without DOX treatment, from 4 to 6 months of age. All animals were tested at 6 months in three standardized paradigms: (1) one-hour home-cage activities to quantify spontaneous behaviors, (2) 10-day trace eyeblink conditioning [6,7], and (3) contextual fear conditioning [8,9] evaluating hippocampal–amygdala-dependent memory. Each measurement was linked by unique mouse identifiers, enabling a cross-task correlation analysis at the individual level.

The dataset is provided in raw, unprocessed form, with explicit documentation of two missing observations. Data files include a master CSV that integrates all variables, domain-specific files for each behavioral paradigm, and a complete data dictionary. Example validation analyses demonstrated group-dependent correlation patterns between conditioned eyeblink response timing and fear memory, illustrating the capacity of the dataset to reveal individual differences [10–12] and cross-task relationships. Such individual-level integration is crucial for investigating behavioral heterogeneity and variability across learning domains. All data were released under an open license to support secondary analyses, computational modeling, and integration with complementary datasets.

The present dataset integrates three behavioral paradigms—trace eyeblink conditioning, contextual fear conditioning, and spontaneous home-cage activity monitoring—that probe the complementary aspects of brain function. Trace eyeblink conditioning primarily reflects cerebellar-dependent associative learning, whereas contextual fear conditioning engages the hippocampal–amygdala circuits related to memory and emotional processing. Conversely, home cage activity measures provide indices of baseline spontaneous behavior that are not constrained by task demands. Although these behavioral domains are not assumed to be strongly correlated at the group level, their integration at the individual animal level enables exploratory analyses of behavioral organization, variability, and cross-task relationships. Accordingly, this Data Descriptor is designed to facilitate data reuse and hypothesis generation rather than inferential testing.

2. Data Description

2.1. Overview of the Dataset

This dataset contains individual-level behavioral measurements obtained from four experimental groups of mice at six months of age:

1. WT without DOX (wild-type mice without doxycycline treatment);
2. WT with DOX (wild-type mice with doxycycline treatment);
3. rTg4510 without DOX (tauopathy model without doxycycline treatment);
4. rTg4510 with DOX (tauopathy model with doxycycline treatment).

Initial analyses of group-level behavioral outcomes in this cohort have been published [3]. The data descriptor provides complete individual-level data to enable secondary analyses, cross-task correlation studies, and alternative analytical approaches. The dataset architecture and file organization are illustrated in Figure 1, which shows the progress from raw behavioral recordings through standardized CSV files to the unified master dataset.

This hierarchical structure ensures data processing transparency and facilitates reuse across different analytical frameworks.




Figure 1. Overview of the dataset structure and processing workflow. This figure illustrates the data transformation pipeline from raw behavioral recordings to publicly available datasets. Raw data from three independent behavioral paradigms (home cage activities monitored by video, trace eyeblink conditioning recorded via electromyography, and contextual fear conditioning assessed by video) were extracted into structured CSV files for each modality. Individual-level identifiers were used to link measurements across tasks and missing values were encoded as NA. The three domain-specific files were integrated into a master dataset containing 36 mice and 14 behavioral variables. The complete dataset is deposited in Zenodo (<https://doi.org/10.5281/zenodo.17864814>) under the CC BY 4.0 license. For cross-task analyses requiring complete measurements across all behavioral domains, the dataset was restricted to 34 mice ($n = 8, 8, 9$, and 9 for WT without DOX, WT with DOX, rTg4510 without DOX, and rTg4510 with DOX, respectively). Abbreviations: EMG, electromyography; NA, Not Available (missing value).

The complete dataset comprises 36 mice ($n = 9$ per group). However, two mice have incomplete data across the behavioral domains: one WT without DOX lacks eyeblink conditioning data and one WT with DOX lacks fear conditioning data. For cross-task analyses requiring complete individual-level linkages across all three behavioral modalities, the dataset was restricted to 34 mice with measurements in all domains ($n = 8, 8, 9$, and 9 for WT without DOX, WT with DOX, rTg4510 without DOX, and rTg4510 with DOX, respectively). All figures and statistical summaries in this descriptor are based on these 34 mice, unless otherwise specified.

2.2. Behavioral Modalities

The dataset integrates measurements from three independent behavioral domains:

2.2.1. Spontaneous Home-Cage Activity

One-hour continuous monitoring yielded five behavioral metrics:

- Distance traveled (m);
- Rearing duration (time ratio);
- Grooming duration (time ratio);
- Hanging duration (time ratio);
- Jumping duration (time ratio).

All home cage measurements were extracted from video recordings and analyzed using HomeCageScan, without any additional smoothing or preprocessing. Figure 2 displays the distributions of these five metrics across the four experimental groups, illustrating both the central tendencies and within-group variability.




Figure 2. Distributions of home-cage behavioral measures across experimental groups. Box plots with overlaid scatter points show raw individual values for five spontaneous behaviors: distance traveled (m), rearing (time ratio), grooming (time ratio), hanging (time ratio), and jumping (time ratio). Each panel displays data from four experimental groups (WT without DOX, WT with DOX, rTg4510 without DOX, and rTg4510 with DOX). Box plots represent the interquartile range with a median line; whiskers extend to $1.5 \times$ the IQR. Individual data points were overlaid to show the within-group variability. This visualization illustrates group-level central tendencies and individual variability without statistical comparisons.

2.2.2. Trace Eyeblink Conditioning

The animals underwent ten consecutive daily training sessions. For each session, the following variables were recorded:

- Conditioned response percentage (CR%, Days 1–10);
- Peak latency of eyelid closure (ms, extracted on days 4 and 10).

Peak latency was defined as the temporal delay from conditioned stimulus (CS) onset to the maximum amplitude of the conditioned eyelid response. One WT without DOX mouse lacks eyeblink data owing to a technical issue during acquisition; this animal is included in all other measurements.

2.2.3. Contextual Fear Conditioning

Following a single training session, freezing behavior was assessed at two retention intervals:

- Freezing percentage at 1 h retention test (Fear_1h);
- Freezing percentage in the 1-day retention test (Fear_day1).

Freezing was quantified using an automated motion threshold algorithm with consistent parameters across all the animals. One WT with a DOX mouse lacks fear conditioning data. All other variables for this animal are complete.

2.3. Data Structure and File Organization

The deposited dataset consists of the following files:

Master_Dataset.csv

- Wide-format table containing all behavioral variables for each mouse.
- Each row represents an individual.
- Missing values are explicitly encoded as NA.
- Includes group labels (Genotype, DOX) and unique animal identifiers (Mouse_ID).

HomeCage_Activity.csv

- Mouse_ID, Genotype, DOX;
- Distance, Rearing, Grooming, Hanging, Jumping.

TraceEBC_CR.csv

- Mouse_ID, Genotype, DOX;
- CR_Day1, CR_Day2, ..., CR_Day10.

TraceEBC_Latency.csv

- Mouse_ID, Genotype, DOX;
- Latency_Day4, Latency_Day10.

FearConditioning.csv

- Mouse_ID, Genotype, DOX;
- Fear_1h, Fear_day1.

Data_Dictionary.csv

- Machine-readable version of Table 1;
- Variable names, descriptions, units, per-group sample sizes, and notes.

All the files use UTF-8 encoding and a comma-separated value (CSV) format. The column names correspond to the variables listed in Table 1.

2.4. Variable Definitions and Data Dictionary

Table 1 provides a complete data dictionary listing all variables, their units, per-group sample sizes, missing values, and methodological notes. The key variable categories are as follows:

Complete list of all variables with descriptions, units, per-group sample sizes (N, in order: WT without DOX/WT with DOX/rTg4510 without DOX/rTg4510 with DOX), missing value status, and functional notes. The variables include group identifiers, five home cage activity measures, eyeblink conditioning metrics at two time points (days 4 and 10), and contextual fear memory at two retention intervals (1 h and 1 day). The sample sizes reflect the complete dataset (36 mice). Missing values: One WT mouse without DOX lacks eyeblink conditioning data and one WT mouse with DOX lacks fear conditioning data. For cross-task analyses that required complete measurements across all behavioral domains, the dataset was restricted to 34 mice ($n = 8, 8, 9$, and 9 for the four groups).

Table 1. Data Dictionary for the Behavioral Dataset.

Variable	Description	Unit	N (Per Group)	Missing	Notes
Group	Experimental condition: WT without DOX, WT with DOX, rTg4510 without DOX, rTg4510 with DOX	–	9/9/9/9	None	Categorical
Distance	Distance traveled in 1 h home-cage monitoring	meters	9/9/9/9	None	General locomotor activity
Rearing	Fraction of time spent rearing during home-cage monitoring	% (time)	9/9/9/9	None	Vertical exploration
Grooming	Fraction of time spent grooming during home-cage monitoring	% (time)	9/9/9/9	None	Self-care behavior
Hanging	Fraction of time spent in hanging posture during home-cage monitoring	% (time)	9/9/9/9	None	Cage-top behavior
Jumping	Fraction of time spent jumping during home-cage monitoring	% (time)	9/9/9/9	None	Vertical locomotion
CR_day4	Percentage of conditioned responses on Day 4 of tEBC	%	8/9/9/9	1 (WT without DOX)	Early phase learning
CR_day10	Percentage of conditioned responses on Day 10 of tEBC	%	8/9/9/9	1 (WT without DOX)	Late-phase learning
Latency_day4	Peak latency of eyelid response on Day 4	ms	8/9/9/9	1 (WT without DOX)	Early temporal processing
Latency_day10	Peak latency of eyelid response on Day 10	ms	8/9/9/9	1 (WT without DOX)	Late temporal refinement
Fear_1h	Freezing during 1 h retention test	%	9/8/9/9	1 (WT with DOX)	Short-term memory retention
Fear_day1	Freezing during 1-day retention test	%	9/8/9/9	1 (WT with DOX)	Long-term memory retention
Mouse_ID	Individual identifier linking all tasks	–	8–9	None	Individual identifier
Age	Age at testing	months	6	None	Fixed at 6 months

Group identifiers:

- Genotype: WT or rTg4510.
- DOX: without or with DOX.
- Mouse_ID: unique identifier linking all measurements.

Home-cage activities:

- Distance: total distance traveled during one-hour monitoring (m).
- Rearing, Grooming, Hanging, Jumping: proportion of time engaged in each behavior (0–1 scale).

Trace eyeblink conditioning:

- CR_Day1 to CR_Day10: percentage of trials with conditioned responses (0–100%).
- Latency_Day4, Latency_Day10: peak latency from CS onset to maximum eyelid closure (ms).

Contextual fear conditioning:

- Fear_1h: freezing percentage during 1 h retention test (0–100%);
- Fear_day1: freezing percentage during 1-day retention test (0–100%).

Note: Some automated freezing scores exceed 100% because of the detection algorithm sensitivity; these values are retained without normalization to preserve data authenticity.

2.5. Data Distributions and Quality

Figure 3 presents the distributions of the four primary behavioral variables across the four experimental groups: distance traveled, rearing duration, CR percentage on day 10 (CR10), and freezing percentage on the 1-day retention test (Fear_day1). Box plots illustrate the range, median, and quartiles for each measure, with individual data points overlaid to show the within-group variability. This visualization is provided for descriptive purposes only and does not include inferential statistics.




Figure 3. Distributions of primary behavioral measures across experimental groups. Box plots show raw individual values for four key behavioral variables: (a) distance traveled in the home cage (m), (b) rearing time (proportion of recording time), (c) conditioned response percentage on day 10 of trace eyeblink conditioning (CR10), and (d) freezing percentage during the 1-day retention test in contextual fear conditioning. Each panel displays data from four experimental groups (WT without DOX, WT with DOX, rTg4510 without DOX, and rTg4510 with DOX). Box plots represent the interquartile range with median; whiskers extend to $1.5 \times$ IQR, and individual data points are overlaid. No statistical comparisons were performed. The figure illustrates the shape and distribution of the raw behavioral data for each modality.

Figure 4 displays a subject $\times$ measure heat map of the raw values of the four primary variables across all individual animals. Rows correspond to animals ordered by experimental group and columns correspond to behavioral measures. Missing values appear as blank cells. This representation enables rapid inspection of data completeness and individual-level correspondence across behavioral domains.

2.6. Missing Data

Missing values are limited to two animals and are explicitly documented:

One WT mouse without DOX lacked all tEBC data (CR and latency values);

One WT with DOX mouse lacked contextual fear conditioning data (Fear_1h and Fear_day1).




Figure 4. Individual-level heatmap showing the full behavioral dataset across experimental groups. Each row corresponds to a single mouse, ordered by group (WT without DOX, WT with DOX, rTg4510 without DOX, or rTg4510 with DOX). The columns represent the four primary behavioral measures: distance traveled, rearing time, conditioned response percentage on day 10 (CR10), and freezing percentage during the 1-day retention test. Raw values were visualized without normalization or scaling, and the color intensity reflected the magnitude. The missing data appear as blank entries. This heatmap provides an overview of dataset completeness, within-group and between-group variability, and cross-task linkages at the individual level.

All other measurements were complete for these animals and all other animals had complete datasets across all modalities. The transparent encoding of missing values (as NA) allows the straightforward implementation of listwise or pairwise deletion strategies during analysis.

2.7. Cross-Task Structure and Validation

Because all behavioral measurements are linked at the individual level, the dataset enables the investigation of cross-task relationships. As an illustrative example of cross-task linkage at the individual level, Figure 5 (shown below) displays group-wise correlations between the CR percentage on Day 10 (CR10) and the freezing percentage on the 1-day retention test (Fear_day1). The four panels illustrate that the strength and direction of the

associations between eyeblink conditioning and contextual fear memory differed across the genotypes and treatment conditions.




Figure 5. Group-wise correlations between late-phase eyeblink conditioning and contextual fear memory. Scatterplots show the relationship between the conditioned response percentage on day 10 of trace eyeblink conditioning (CR10, x -axis) and freezing percentage during the 1-day retention test in contextual fear conditioning (y -axis). Each panel represents one experimental group: (a) WT without DOX, (b) WT with DOX, (c) rTg4510 without DOX, or (d) rTg4510 with DOX. Each point represents one mouse. Pearson correlation coefficients (r) are indicated within each panel after the listwise deletion of missing values. No inferential comparisons were performed. This figure illustrates an example of cross-task relationships that can be explored using this dataset.

Additional validation analyses are provided in Appendix A Figures A1–A3:

- Figure A1: Correlation between CR percentage on days 4 and 1-day of freezing;
- Figure A2: Correlations between peak latency on Day 4 and 1-day freezing;
- Figure A3: Correlations between peak latency on Day 10 and 1-day freezing.

These analyses were included solely to demonstrate the capacity of the dataset for cross-paradigm investigations and are not presented as inferential findings. A detailed interpretation is presented in Section 3 (Technical Validation).

2.8. Data Formats and Compatibility

All data are provided in widely compatible CSV format with the following specifications:

- Encoding: UTF-8.
- Delimiter: comma (,).
- Decimal separator: period (.).
- Missing values: NA.
- Row structure: each row represents one animal.
- Column structure: Each column represents one variable or measurement day.

2.9. Reproducibility and Reusability

The dataset is structured to support multiple analytical approaches:

- Cross-task correlation analyses: Individual-level linkages enable examination of the relationships between independent learning systems.
- Individual difference modeling: Complete behavioral profiles support clustering, classification, and latent variable analysis.
- Treatment and genotype comparisons: The four-group design allows the examination of DOX effects and genotype-dependent phenotypes.
- Temporal dynamics: Daily CR measurements and dual retention time points enable the investigation of learning trajectories.
- Machine learning applications: A multivariate structure supports supervised and unsupervised learning algorithms.

All measurements are provided in their original, unprocessed forms, without smoothing, normalization, or imputation. This approach maximizes the transparency and allows researchers to apply their own preprocessing pipelines according to specific analytical requirements.

3. Methods

3.1. Animals and Experimental Design

Six-month-old male and female rTg4510 and wild-type (WT) mice were used in this study. The rTg4510 line expresses human P301L mutant tau under the control of a tetracycline-responsive promoter, enabling doxycycline (DOX)-mediated suppression of transgenic tau expression. Animals were assigned to four experimental groups based on their genotype and DOX treatment.

1. WT without DOX ($n = 9$);
2. WT with DOX ($n = 9$);
3. rTg4510 without DOX ($n = 9$);
4. rTg4510 with DOX ($n = 9$).

Mice in the DOX-treated groups received doxycycline-containing chow (200 mg/kg) ad libitum from four to six months of age. All animals were housed under a 12:12 h light–dark cycle at 22–24 °C with food and water available ad libitum. Behavioral testing was conducted in the following sequence to minimize task interference: (1) home-cage activity monitoring, (2) tEBC, and (3) contextual fear conditioning. The animal study protocol underlying this dataset was approved by the Institutional Animal Investigation Committee of Tokushima Bunri University (protocol code KP19-75-001; date of approval: 1 March 2019).

3.2. Home-Cage Activity Monitoring

Spontaneous home cage behavior was recorded for one hour using a standard overhead video camera, and all recordings were saved as MPEG video files. Behavioral categories were quantified using HomeCageScan (CleverSys, Inc., Reston, VA, USA), which automatically classifies behaviors from video recordings [13–15]. All metrics (distance traveled, rearing, grooming, hanging, and jumping) represent the direct output of HomeCageScan without any additional smoothing or preprocessing applied by the experimenters. Distance traveled reflects overall locomotor activity, whereas rearing represents vertical exploratory behavior. Grooming, hanging, and jumping were categorized as non-locomotor spontaneous behaviors.

Five behavioral metrics were quantified:

- Distance traveled: Total distance (m) computed from the centroid displacement across video frames.
- Rearing: Proportion of time spent in the vertical exploration posture.
- Grooming: Proportion of time engaged in self-grooming behavior.
- Hanging: Proportion of time spent in the cage-hanging posture.
- Jumping: Proportion of time engaged in jumping movements.

All proportional metrics are expressed as time ratios (0–1 scale), allowing normalization across individuals and minimizing the influence of transient activity bursts. Missing values owing to video occlusion or detection uncertainty were retained as NA. No smoothing, imputation, or normalization was applied to preserve the data authenticity.

3.3. Trace Eyeblink Conditioning

Trace eyeblink conditioning (tEBC) was conducted over 10 consecutive days using a well-established trace paradigm [3,4].

Conditioning protocol:

- Conditioned stimulus (CS): 250 ms tone (1 kHz, 80 dB);
- Trace interval: 500 ms stimulus-free period;
- Unconditioned stimulus (US): 100 ms corneal electrical shock (0.2 mA, 100 Hz, square pulses);
- Intertrial interval: 20–40 s (variable);
- Trials per session: 100;
- Training duration: 10 consecutive days (days 1–10).

Response detection:

A conditioned response (CR) was defined as eyelid closure exceeding a fixed amplitude threshold during the trace interval (between CS offset and US onset). The detection thresholds and filtering parameters were identical across all animals to ensure consistency.

Extracted variables:

- CR percentage (CR%): Proportion of trials with CRs for each training day (days 1–10).
- CR peak latency: Time (ms) from CS onset to maximum eyelid deflection during CR trials, extracted on days 4 and 10.

One WT mouse without DOX lacked reliable eyelid recordings because of a technical failure during acquisition; all values for this animal were encoded as NA in the tEBC dataset.

3.4. Contextual Fear Conditioning

Contextual fear conditioning was performed in a standard conditioning chamber (30 cm × 30 cm × 35 cm) equipped with a stainless-steel grid floor and a physical movement-tracking FreezeScan™ 1.0 system (CleverSys, Inc., Reston, VA, USA) [5].

Training protocol:

- Habituation: 2 min exploration period.
- Footshock: Single 2 s shock at 0.7 mA.
- Post-shock period: 30 s before removal from the chamber.

Retention testing:

Freezing behavior was assessed at two time points:

- 1 h retention test: 2 min exposure to the same context;
- 1-day retention test: 2 min exposure to the same context.

Freezing was quantified automatically using a motion threshold detection algorithm (FreezeScan) with consistent parameters across all animals. The freezing percentage was calculated as the proportion of time spent in the freezing posture (absence of movement except respiration) during the test period.

Extracted variables:

- Fear_1h: Freezing percentage during the 1 h retention test;
- Fear_day1: Freezing percentage during the 1-day retention test.

One WT with a DOX mouse lacked contextual fear conditioning data owing to equipment malfunction; values for this animal were encoded as NA.

3.5. Data Integration and Organization

All behavioral measurements were linked at the individual animal level using unique mouse identifiers (Mouse_ID). This structure enables a direct cross-task analysis without requiring data alignment or matching.

File organization:

The dataset is provided as multiple CSV files to facilitate selective loading and analysis:

1. Master_Dataset.csv: Complete wide-format table with all variables;
2. HomeCage_Activity.csv: Distance traveled and behavioral time ratios;
3. TraceEBC_CR.csv: CR percentages for Days 1–10;
4. TraceEBC_Latency.csv: Peak latency values for Days 4 and 10;
5. FearConditioning.csv: Freezing percentages at 1 h and 1 day;
6. Data_Dictionary.csv: Machine-readable variable reference corresponding to Table 1.

Data processing:

No preprocessing, smoothing, normalization, or imputation was applied to any behavioral measurements. All values represent direct outputs from the acquisition systems or automated detection pipelines. Missing values were explicitly encoded as NA and documented in the Data Description Section.

File format specifications:

Encoding: UTF-8;

Delimiter: Comma (,);

Decimal separator: Period (.);

Missing values: NA;

Row structure: Each row represents one individual animal;

Column structure: Each column represents one variable or one measurement day.

3.6. Statistical Analyses for Technical Validation

Cross-task correlation analyses were performed to validate internal consistency and demonstrate the capacity of the dataset for cross-paradigm investigation. Pearson's correlation coefficients were computed between the eyeblink conditioning metrics (CR percentage and peak latency) and contextual fear memory (1-day freezing percentage) within each experimental group.

Analyses were conducted separately for:

CR percentage on Day 4 vs. Fear_day1 (Figure A1);

Peak latency on Day 4 vs. Fear_day1 (Figure A2);

Peak latency on Day 10 vs. Fear_day1 (Figure A3);

CR percentage on Day 10 vs. Fear_day1 (Figure 5).

Missing values were handled by list-wise deletion on a per-analysis basis. Correlation analyses are presented solely to demonstrate data quality and cross-task relationships; they are not intended as inferential conclusions regarding mechanisms or treatment effects. All visualization and analysis codes are available in the data repository.

3.7. Software and Equipment

Home-cage monitoring: Video acquisition and automated behavioral classification were performed using HomeCageScan system (CleverSys, Inc., Reston, VA, USA).

Trace eyeblink conditioning: Trace eyeblink conditioning was performed using a fully custom-built EMG-based acquisition and stimulus-control system. Eyelid muscle activity was recorded using fine-wire EMG electrodes and amplified using a commercial differential amplifier. Amplified EMG signals were digitized using an A/D converter operating in a DOS/V environment. All stimulus timings, including tone CS presentation, trace-interval control, and periorbital electrical US delivery, were executed using in-house software running on a Microsoft Windows 3.1-based system. Conditioned responses (CRs) were defined as increases in EMG activity that exceeded a fixed threshold within a trace interval. The detailed procedures for EMG recording, filtering, and CR detection have been described previously in our earlier report [3,4].

Fear conditioning: Standard conditioning chamber with automated freezing detection using the FreezeScanTM 1.0 system (CleverSys, Inc., Reston, VA, USA).

Data processing: Raw behavioral outputs from each system (HomeCageScan, EMG-based tEBC system, and contextual fear conditioning software) were manually exported and organized into CSV files. No additional preprocessing, smoothing, or normalization was performed.

Statistical analysis: Technical validation analyses were performed by computing Pearson correlation coefficients between eyeblink conditioning measures and contextual fear memory within each experimental group. All correlation analyses (*r* values) and descriptive statistics were performed using the GraphPad Prism software (version 5.0).

Visualization: Initial plots (scatter plots and summary graphs) were generated using GraphPad Prism software. The final figure layouts, labels, and graphical adjustments were completed manually using Microsoft PowerPoint. No smoothing, transformation, or additional data preprocessing was applied during the figure preparation.

4. Technical Validation

4.1. Quality Control and Measurement Consistency

All behavioral measurements were conducted under standardized conditions with consistent parameters across animals and sessions. Home cage activity was continuously recorded for one hour using identical monitoring systems. tEBC employed fixed CS-US parameters and detection thresholds across all training days. Contextual fear conditioning used uniform shock intensity (0.7 mA) and automated freezing detection with consistent motion thresholds. Missing values are limited to two animals and are explicitly documented: one WT without a DOX mouse lacks eyeblink conditioning data owing to technical failure, and one WT with a DOX mouse lacks fear conditioning data owing to equipment malfunction. All other measurements were performed. No preprocessing, smoothing, or imputation was applied to any behavioral variables to ensure data authenticity and transparency.

4.2. Cross-Task Relationships

To validate the capacity of the dataset for cross-paradigm investigation, we examined the correlations between eyeblink conditioning metrics and contextual fear memory at the individual level. Group-wise analyses revealed distinct patterns.

Early acquisition (CR Day 4 vs. Fear 1 day, Figure A1): Weak or inconsistent correlations across all groups indicate that early phase CR acquisition does not robustly predict contextual memory performance.

Early temporal processing (Latency Day 4 vs. Fear 1 day, Figure A2): Variable patterns across groups, with rTg4510 without DOX showing a positive trend, suggesting altered coordination between temporal processing and fear memory.

Late temporal processing (Latency Day 10 vs. Fear 1 day, Figure A3): Striking divergence emerged, with WT without DOX showing a negative correlation (faster CR timing associated with stronger fear memory), whereas rTg4510 with DOX showed a positive correlation, indicating DOX-dependent reorganization of behavioral coupling.

Late acquisition (CR Day 10 vs. Fear 1 day, Figure 5): Group-dependent correlation patterns confirmed meaningful individual differences and group-specific behavioral organization.

Across all analyses, no systematic confounders attributable to home-cage locomotor activity were observed, confirming that cross-task relationships reflect domain-specific learning processes rather than general motor or arousal differences. Notably, substantial individual variability was observed within each group, particularly in the rTg4510 with DOX group, where some animals showed learning performance comparable to that of wild-type mice, whereas others exhibited marked deficits. This heterogeneity underscores the value of individual-level data for investigating response variability in therapeutic interventions.

The individual-level structure of a dataset enables the investigation of cross-task relationships, an approach that has proven valuable for understanding behavioral variability and learning mechanisms [10,11]. While group-level analyses remain standard in behavioral neuroscience, individual difference approaches can reveal patterns obscured by averaging and provide insights into response heterogeneity [11,12]. The present dataset is one of the few publicly available resources linking cerebellar-dependent and hippocampal–amygdala-dependent learning at the individual level in neurodegenerative models.

4.3. Data Quality Summary

Technical validation demonstrates that this dataset (1) maintains internal consistency across measurement modalities, (2) captures substantial individual variability suitable for secondary analyses, (3) supports cross-task investigation through reliable individual-level linkages, and (4) exhibits group-specific correlation patterns consistent with biological phenotypic variation. The transparent encoding of limited missing values and the absence of preprocessing ensures that the dataset can be readily adapted to diverse analytical frameworks and research questions.

5. User Notes

5.1. Potential Applications

This dataset supports investigations across multiple domains:

Cross-task behavioral profiling: The individual-level linkage between spontaneous activity, tEBC, and contextual fear memory enables examination of the relationships between independent learning systems within the same animals.

Tauopathy research: The inclusion of rTg4510 mice with and without doxycycline treatment allowed us to investigate how tau pathology and its suppression affect non-spatial associative learning and behavioral organization.

Individual difference modeling: The dataset structure supports clustering analyses, dimensionality reduction, and latent variable modeling to identify behavioral subtypes across genotypes and treatment conditions.

Computational modeling: The combination of trial-by-trial eyeblink conditioning data with contextual fear memory outcomes provides a foundation for computational models of temporal associative learning and circuit-level dysfunction.

5.2. Analytical Strategies

As the dataset is provided in a raw, unprocessed form, users have the flexibility to implement diverse analytical approaches.

Alternative detection criteria: Users can redefine the conditioned response thresholds or apply different freezing detection algorithms according to their research needs.

Temporal dynamics: Daily CR measurements enable the investigation of acquisition trajectories, learning curves, and within-animal stability across training sessions.

Multivariate approaches: The dataset supports regression modeling with interaction terms, mixed-effects models, and machine learning algorithms for prediction or classification.

All data files are provided in standard CSV format compatible with R, Python, MATLAB, and spreadsheet software. The dataset can be integrated with complementary data types (histopathological, genetic, and physiological) to construct comprehensive tauopathy and behavioral dysfunction models.

Author Contributions: Conceptualization, Y.K.; Methodology, R.K. and Y.K.; Formal analysis, R.K., T.N. and Y.K.; investigation, Y.K.; data curation, Y.K. and R.K.; writing—original draft preparation, Y.K.; writing—review and editing, R.K., T.N. and Y.K.; visualization, Y.K.; project administration, Y.K.; funding acquisition, Y.K. All authors have read and agreed to the published version of the manuscript.

Funding: The original behavioral experiments were funded by the JSPS KAKENHI grant 16K08215 (Principal Investigator: Yutaka Kirino; Co-Investigator: Y.K.). Data reanalysis and manuscript preparation were supported by JSPS KAKENHI grants 22K06618 and 25K10188 awarded to Y.K. APC was funded by KAKENHI (Grant 25K10188).

Institutional Review Board Statement: The original animal experiments from which this dataset was derived were approved by the Institutional Animal Investigation Committee of the Tokushima Bunri University (protocol code KP19-75-001, approved on 1 March 2019). No additional animal experiments were conducted using this descriptor.

Informed Consent Statement: Not applicable.

Data Availability Statement: The complete behavioral dataset, including all raw individual-level measurements of home cage activities, tEBC, and contextual fear conditioning, is publicly available at <https://doi.org/10.5281/zenodo.17864814>. The dataset comprised 36 mice ($n = 9$ per group), with complete measurements for 34 mice across all three behavioral domains. All files are provided in their original and unprocessed forms to ensure full reproducibility and transparency. The repository includes Master CSV file containing all behavioral variables linked by individual mouse identifiers (36 mice $\times$ 14 variables). Data dictionary with complete variable definitions, units, and sample sizes. README file describing experimental protocols, file structure, and data formats. All files are released under a Creative Commons Attribution 4.0, International (CC BY 4.0) license and can be downloaded without restriction.

Acknowledgments: The authors thank Yutaka Kirino and other members of the original research team for their contributions to the data collection. During the preparation of this manuscript, we used Claude Sonnet 4.5 (Anthropic, San Francisco, CA, USA; accessed January 2026) and ChatGPT (OpenAI, GPT-4 architecture, San Francisco, CA, USA; accessed January 2026) for data visualization code generation, manuscript structure optimization, and language editing. The authors take full responsibility for the content of this manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

WT	Wild type
DOX	Doxycycline
tEBC	Trace Eyeblink Conditioning
CR	Conditioned Response

CS Conditioned Stimulus
 US Unconditioned Stimulus
 EMG Electromyography
 NA Not Available

Appendix A




Figure A1. Group-wise correlations between early conditioned response rate and contextual fear memory. Scatterplots show the relationship between conditioned response percentage on Day 4 of trace eyeblink conditioning (CR Day 4, x-axis) and freezing percentage during the 1-day retention test in contextual fear conditioning (y-axis). Each panel represents one experimental group: (a) WT without DOX, (b) WT with DOX, (c) rTg4510 without DOX, and (d) rTg4510 with DOX. Each point represents a single mouse. Pearson correlation coefficients (r) are indicated within each panel after listwise deletion of missing observations. Early CR rate did not provide a robust or consistent predictor of contextual freezing across genotypes or treatment conditions.




Figure A2. Group-wise correlations between early CR peak latency and contextual fear memory. Scatterplots depict the relationship between CR peak latency on Day 4 of trace eyeblink conditioning (measured from CS onset to eyelid response peak, x-axis) and freezing percentage during the 1-day contextual fear retention test (y-axis). Each panel represents one experimental group: (a) WT without DOX, (b) WT with DOX, (c) rTg4510 without DOX, and (d) rTg4510 with DOX. Each point represents

a single mouse, with listwise deletion applied for missing eyeblink data. Pearson's r is shown in each panel. WT groups exhibit weak relationships, whereas rTg4510 without DOX displays a positive trend, suggesting that early temporal processing may differentially relate to contextual memory in the presence of tau pathology.




Figure A3. Group-wise correlations between late CR peak latency and contextual fear memory. Scatterplots present the relationship between CR peak latency on Day 10 of trace eyeblink conditioning (x -axis) and freezing percentage during the 1-day contextual fear retention test (y -axis). Each panel represents one experimental group: (a) WT without DOX, (b) WT with DOX, (c) rTg4510 without DOX, and (d) rTg4510 with DOX. Each point represents a single mouse. Pearson correlation coefficients are computed from complete data pairs. WT without DOX mice show a negative correlation (shorter latencies associated with stronger contextual freezing), whereas rTg4510 with DOX mice show a positive correlation (longer latencies associated with higher freezing). This sign reversal suggests DOX-dependent reorganization of temporal processing in associative learning.

References

1. Santacruz, K.; Lewis, J.; Spire, T.; Paulson, J.; Kotilinek, L.; Ingelsson, M.; Guimaraes, A.; DeTure, M.; Ramsden, M.; McGowan, E.; et al. Tau suppression in a neurodegenerative mouse model improves memory function. *Science* **2005**, *309*, 476–481. [[CrossRef](#)] [[PubMed](#)]
2. Ramsden, M.; Kotilinek, L.; Forster, C.; Paulson, J.; McGowan, E.; SantaCruz, K.; Guimaraes, A.; Yue, M.; Lewis, J.; Carlson, G.A.; et al. Age-dependent neurofibrillary tangle formation, neuron loss, and memory impairment in a mouse model of human tauopathy (P301L). *J. Neurosci.* **2005**, *25*, 10637–10647. [[CrossRef](#)] [[PubMed](#)]
3. Kishimoto, Y.; Kubota, T.; Nakashima, K.; Kirino, Y. rTg4510 tauopathy mice exhibit non-spatial memory deficits prevented by Doxycycline treatment. *Brain Sci.* **2025**, *15*, 1183. [[CrossRef](#)] [[PubMed](#)]
4. Kishimoto, Y.; Kirino, Y. Presenilin 2 mutation accelerates the onset of impairment in trace eyeblink conditioning in a mouse model of Alzheimer's disease overexpressing human mutant amyloid precursor protein. *Neurosci. Lett.* **2013**, *538*, 15–19. [[CrossRef](#)] [[PubMed](#)]
5. Kishimoto, Y.; Fukumoto, K.; Nagai, M.; Mizuguchi, A.; Kobashi, Y. Early contextual fear memory deficits in a double-transgenic amyloid- β precursor protein/Presenilin 2 mouse model of Alzheimer's disease. *Int. J. Alzheimers Dis.* **2017**, *2017*, 8584205. [[CrossRef](#)] [[PubMed](#)]
6. McGlinchey-Berroth, R.; Carrillo, M.C.; Gabrieli, J.D.; Brawn, C.M.; Disterhoft, J.F. Impaired trace eyeblink conditioning in bilateral, medial-temporal lobe amnesia. *Behav. Neurosci.* **1997**, *111*, 873–882. [[CrossRef](#)] [[PubMed](#)]
7. Christian, K.M.; Thompson, R.F. Neural substrates of eyeblink conditioning: Acquisition and retention. *Learn. Mem.* **2003**, *10*, 427–455. [[CrossRef](#)] [[PubMed](#)]
8. Kim, J.J.; Fanselow, M.S. Modality-specific retrograde amnesia of fear. *Science* **1992**, *256*, 675–677. [[CrossRef](#)] [[PubMed](#)]
9. Fanselow, M.S. Contextual fear, gestalt memories, and the hippocampus. *Behav. Brain Res.* **2000**, *110*, 73–81. [[CrossRef](#)] [[PubMed](#)]

10. Matzel, L.D.; Sauce, B.; Wass, C. The architecture of intelligence: Converging evidence from studies of humans and animals. *Curr. Dir. Psychol. Sci.* **2013**, *22*, 342–348. [[CrossRef](#)]
11. Sauce, B.; Matzel, L.D. The causes of variation in learning and behavior: Why individual differences matter. *Front. Psychol.* **2013**, *4*, 395. [[CrossRef](#)] [[PubMed](#)]
12. Crabbe, J.C.; Wahlsten, D.; Dudek, B.C. Genetics of mouse behavior: Interactions with laboratory environment. *Science* **1999**, *284*, 1670–1672. [[CrossRef](#)] [[PubMed](#)]
13. Jhuang, H.; Garrote, E.; Yu, X.; Khilnani, V.; Poggio, T.; Steele, A.D.; Serre, T. Automated home-cage behavioral phenotyping of mice. *Nat. Commun.* **2010**, *1*, 68. [[CrossRef](#)] [[PubMed](#)]
14. Steele, A.D.; Jackson, W.S.; King, O.D.; Lindquist, S. The power of automated high-resolution behavior analysis revealed by its application to mouse models of Huntington’s and prion diseases. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 1983–1988. [[CrossRef](#)] [[PubMed](#)]
15. Kishimoto, Y.; Higashihara, E.; Fukuta, A.; Nagao, A.; Kirino, Y. Early impairment in a water-finding test in a longitudinal study of the Tg2576 mouse model of Alzheimer’s disease. *Brain Res.* **2013**, *1491*, 117–126. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.