

Analysis of Rhizosphere Bacteriomes from Different Dominant Plants in the Water-Level Fluctuation Zone of the Three Gorges Reservoir

Lanfang Zhou ^{1,2,3,*}, Yutao Gao ⁴, Shengjun Wu ^{2,3,*} and Maohua Ma ^{2,3}

¹ School of River and Ocean Engineering, Chongqing Jiaotong University, Chongqing 400074, China

² Key Laboratory of Reservoir Aquatic Environment, Chongqing Institute of Green and Intelligent Technology, Chinese Academy of Sciences, Chongqing 400714, China

³ Chongqing School, University of Chinese Academy of Sciences, Chongqing 400714, China

⁴ PowerChina Chengdu Engineering Corporation Limited, Chengdu 610072, China

* Correspondence: lfzhou2022@163.com (L.Z.); wsj@cigit.ac.cn (S.W.)




Figure S1 Rarefaction curves based on Shannon index of rhizosphere and non-rhizosphere bacteriomes of *P. lapathifolia* (L.) Delarbre.




Figure S2 Rarefaction curves based on Shannon index of rhizosphere and non-rhizosphere bacteriomes of *A. annua* L.




Figure S3 The relative abundance of rhizosphere and non-rhizosphere bacteriomes at the order (a), family (b) and genus (c) levels from *P. lapathifolia* (L.) Delarbre.




Figure S4 The relative abundance of rhizosphere and non-rhizosphere bacteriomes at the order (a), family (b) and genus (c) levels from *A. annua* L.