



Jac A. Nickoloff ^{1,*}, Neelam Sharma ¹, Lynn Taylor ¹, Sage J. Allen ¹ and Robert Hromas ²

- ¹ Department of Environmental and Radiological Health Sciences, Colorado State University, Ft. Collins, CO 80523, USA; neelam.sharma@colostate.edu (N.S.); lynn.taylor@colostate.edu (L.T.); sage.allen@rams.colostate.edu (S.J.A.)
- ² Division of Hematology and Medical Oncology, Department of Medicine and the Mays Cancer Center, University of Texas Health Science Center, San Antonio, TX 78229, USA; hromas@uthscsa.edu
- * Correspondence: j.nickoloff@colostate.edu

Abstract: DNA replication stress is a constant threat that cells must manage to proliferate and maintain genome integrity. DNA replication stress responses, a subset of the broader DNA damage response (DDR), operate when the DNA replication machinery (replisome) is blocked or replication forks collapse during S phase. There are many sources of replication stress, such as DNA lesions caused by endogenous and exogenous agents including commonly used cancer therapeutics, and difficult-to-replicate DNA sequences comprising fragile sites, G-quadraplex DNA, hairpins at trinucleotide repeats, and telomeres. Replication stress is also a consequence of conflicts between opposing transcription and replication, and oncogenic stress which dysregulates replication origin firing and fork progression. Cells initially respond to replication stress by protecting blocked replisomes, but if the offending problem (e.g., DNA damage) is not bypassed or resolved in a timely manner, forks may be cleaved by nucleases, inducing a DNA double-strand break (DSB) and providing a means to accurately restart stalled forks via homologous recombination. However, DSBs pose their own risks to genome stability if left unrepaired or misrepaired. Here we focus on replication stress response systems, comprising DDR signaling, fork protection, and fork processing by nucleases that promote fork repair and restart. Replication stress nucleases include MUS81, EEPD1, Metnase, CtIP, MRE11, EXO1, DNA2-BLM, SLX1-SLX4, XPF-ERCC1-SLX4, Artemis, XPG, and FEN1. Replication stress factors are important in cancer etiology as suppressors of genome instability associated with oncogenic mutations, and as potential cancer therapy targets to enhance the efficacy of chemoand radiotherapeutics.

Keywords: DNA damage; replication stress; genome instability; DNA double-strand breaks; structurespecific nucleases; DNA damage response

1. Introduction

Accurate DNA replication and proper chromosome segregation to daughter cells are critical to maintaining genome integrity and preventing cancer. Replication of the 6.3 billion bp of the diploid human genome during a typical eight-hour S phase requires >30,000 active origins, ~5000 of which are active at a time [1,2]. Replication forks travel in a highly processive manner, synthesizing ~3000 bp per min, yet forks frequently encounter obstacles that stall replisomes, causing replication stress and triggering stress responses including the intra-S checkpoint [3], fork protection to prevent replisome dissociation or fork collapse, and repair mechanisms that restart damaged forks. Replication stress is caused by a wide variety of endogenous and exogenous factors. Spontaneous DNA damage is caused by endogenous reactive oxygen species formed during cellular metabolism [4,5], misincorporation of ribonucleotides, and DNA lability [5]. DNA damage is also caused by exogenous genotoxic chemicals, and by ionizing and non-ionizing radiation. The vast majority of DNA lesions block replicative polymerases, necessitating lesion repair by an appropriate repair pathway, lesion bypass (damage tolerance) by translesion DNA synthesis (TLS)



Citation: Nickoloff, J.A.; Sharma, N.; Taylor, L.; Allen, S.J.; Hromas, R. Nucleases and Co-Factors in DNA Replication Stress Responses. *DNA* 2022, 2, 68–85. https://doi.org/ 10.3390/dna2010006

Academic Editor: Justin Courcelle

Received: 1 February 2022 Accepted: 22 February 2022 Published: 1 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). polymerases, repriming, homologous recombination (HR) mediated template switching, or passive rescue from an adjacent fork [6,7]. DNA lesions that block replicative polymerases include nucleotides with broken rings, oxidized bases, and chemical adducts, as well as single- and double-strand breaks (DSBs). DNA polymerase inhibitors and depletion of nucleotide pools with hydroxyurea are exogenous sources that cause global replication stress, slowing or stopping most or all replication forks [8].

Additional endogenous sources of replication stress are difficult to replicate DNA sequences and certain chromatin environments (e.g., G-quadraplex DNA, common fragile sites, telomeric DNA) [9–17]. Replication stress is also caused by stable R-loops which form by hybridization of RNA transcripts to DNA templates, especially in G-rich sequences [18–20], and by collisions between opposing transcription and replication machinery, particularly in highly transcribed ribosomal RNA gene arrays, fragile sites, and telomeres [21–27]. Topoisomerases avert replication stress by preventing DNA overwinding in front of replication forks, a type of intrinsic, topological replication stress. A recent yeast study showed that cohesin, a highly conserved protein with essential roles in sister chromatid cohesion required for proper chromosome segregation, increases replication stress in centromeric and ribosomal DNA by trapping topological stress [28]. Although cells suffer replication stress at random sites throughout the genome due to spontaneous (or induced) DNA damage, the stress associated with difficult to replicate sequences and challenging chromatin environments must be managed at those sites in every S phase.

When replication forks are blocked, the initial response has two aims: (1) protect the replication fork by stabilizing the replisome machinery, and (2) protect the fork from nucleolytic attack [29,30]. If a blocked fork is not restarted in a timely manner, it may be cleaved by structure-specific nucleases yielding a single-ended DSB (seDSB) that is processed by resection nucleases to suppress misrepair by canonical non-homologous endjoining (cNHEJ) and promote accurate fork restart by HR. This is important because cNHEJ is the dominant DSB repair pathway in mammalian cells [31,32] and cNHEJ of seDSBs can cause deletions and translocations that produce acentric and dicentric chromosomes that segregate improperly in mitosis or induce breakage-bridge-fusion cycles that further threaten genome integrity [33]. In this review we begin by discussing DDR signaling in response to replication stress, and then describe the activities of key nucleases and co-factors in replication fork protection, fork cleavage to seDSBs, and fork resection that together promote accurate HR-medicated fork repair and restart. These mechanisms are crucial for maintaining genome stability and thus preventing cancer, and they are important as potential targets in cancer therapy.

2. DDR Signaling in Response to Replication Stress

DNA repair and DNA damage checkpoint systems minimize replication fork encounters with blocking lesions [34–36], but with a steady state of ~10,000 DNA lesions per cell in unstressed cells [5] and ~5000 simultaneously active replisomes during S phase [2], fork encounters with blocking lesions are unavoidable. Acute or chronic exposures to genotoxic chemicals and radiation greatly increase replication stress, as does dysregulated replication associated with oncogenic stress [37–39]. Under normal circumstances, the leading and lagging strand replication machines are coupled, traveling together with the MCM (minichromosome maintenance) replicative helicase. If the leading strand polymerase is blocked, MCM helicase may decouple and unwind DNA ahead of the fork, exposing hundreds of bases of single-stranded DNA (ssDNA) [40,41]. As with ssDNA exposed by 5'-3' resection of broken ends at DSBs by resection nucleases EXO1 and DNA2 (with its cofactor BLM) [42–44], the ssDNA exposed by decoupled MCM helicase is rapidly bound by the abundant, heterotrimeric replication protein A (RPA) (Figure 1A). RPA-bound ss-DNA is recognized by the ATR (ataxia telangiectasia and Rad3-related) cofactor ATRIP (ATR-interacting protein), leading to activation of ATR (Figure 1B), the central signaling kinase of the intra-S checkpoint response [45]. In addition to ssDNA-RPA and ATRIP, ATR activation requires several other factors including TopBP1, RAD17-RFC, and the 9-1-1

complex. Additional, distinct ATR activation mechanisms have been described involving NBS1, a component of the MRE11-RAD50-NBS1 (MRN) complex, and the RPA-binding factor ETAA1 [45–47].



Figure 1. Replication Protein A (RPA) roles in replication stress responses. (**A**) RPA is a heterotrimer with 14, 32, and 70 kDa subunits, each with single strand DNA (ssDNA) binding domains called OB (oligonucleotide binding) folds. (**B**) Diagram of a replication fork in which the leading strand DNA polymerase is blocked and decouples from the MCM helicase, creating ssDNA that is bound by RPA. ATRIP recognition of ssDNA-RPA recruits and activates ATR. (**C**) Crosstalk among phosphatidyl inositol 3' kinase-related kinases (PIKKs) and cyclin dependent kinase (CDK) for phosphorylation of serine and threonine residues in the N-terminus of RPA32. The width of each arrow is proportional to the role that each PIKK plays in phosphorylating specific RPA32 residues. Phosphorylated RPA32 residues prime phosphorylation of other residues, indicated by arrows below.

ATR is a member of the phosphatidyl inositol 3' kinase-related kinase (PIKK) family, which also includes ATM and the catalytic subunit of DNA-dependent protein kinase (DNA-PKcs). PIKKs play central roles in DNA damage responses including DSB repair, checkpoint activation, apoptosis, suppression of transcription, and responses to telomere dysfunction and viral infection [48,49]. Activation of each PIKK involves a specific co-factor. The ATR interacting protein ATRIP recruits ATR to RPA-bound ssDNA, initiating ATR activation [50]. The MRN complex and the Ku70/Ku80 heterodimer bind to frank DSB ends, the NBS1 component of MRN recruits and activates ATM, and Ku70/Ku80

recruits and activates DNA-PKcs [48]. Activated PIKKs are autophosphorylated, and they phosphorylate each other and many other targets, showing various degrees of signaling pathway crosstalk [48].

One feature of PIKK crosstalk is apparent in the phosphorylation of RPA bound to ssDNA. RPA is a trimeric complex of 14 kDa, 32 kDa, and 70 kDa subunits with essential roles in DNA replication, DNA repair, and DDR signaling. The N-terminus of the 32 kDa subunit RPA32 (also called RPA2) is phosphorylated on serine and threonine residues by all three PIKKs and cyclin dependent kinase (CDK) (Figure 1C) [51–53]. Phosphorylation of specific RPA32 residues occurs sequentially [54], and residues modified early can prime phosphorylation of other residues (Figure 1C) [51–53]. Following DNA damage, CDK phosphorylates RPA32 S23 and S29, and phospho-S23 (p-S23) primes subsequent phosphorylation of S29, S33 (by ATR), and S4S8 (both phosphorylated by ATM and DNA-PK). These priming effects are sometimes reciprocal; for example, ATR phosphorylation of RPA32 T21 by DNA-PK primes phosphorylation of S4S8 by ATM and DNA-PK, and vice versa (Figure 1C). Thus, CDK phosphorylation of RPA32 initiates a positive feedback loop that results in hyperphosphorylated RPA32, marked by pS4S8 and pT21, which is required for certain downstream events such as apoptosis [55]. RPA-bound ssDNA has emerged as a critical structural foundation for a variety of DDR signaling responses [54]. In addition to its key role in ATR activation, RPA phosphorylation regulates replication in response to stress, and it inhibits resection [56,57]. Most RPA32 phosphorylation events documented to date reflect early DDR signaling as RPA becomes progressively phosphorylated, but in cells stressed with the Topo I inhibitor camptothecin, S12 phosphorylation occurs one and a half to two days after the drug is removed, and this correlates with resumption of DNA replication, suggesting that this modification deactivates Chk1 to terminate checkpoint arrest [51].

RPA modifications regulate cell responses to stress, at least in part, by modulating RPA interactions with DNA and various protein partners, many of which have important DDR signaling and DNA repair roles. For example, RPA phosphorylation reduces its affinity for undamaged double-stranded DNA (dsDNA), but increases its affinity for damaged dsDNA, and it regulates RPA affinity for ssDNA [58,59]. Phospho-RPA shows reduced interactions with ATM, DNA-PK, MRN, 53BP1, and p53 [54]. Conversely, phospho-RPA shows enhanced affinity for PRP19, an E3 ubiquitin ligase important for ATR-ATRIP association with RPA-bound ssDNA, and represents a positive feedback system for ATR activation [60]. Other RPA modifications, including SUMOylation and ubiquitylation, regulate its interactions with other proteins, including the key HR factors RAD51 and RAD52 [54]. RPA inhibitors designed to interfere with RPA binding to ssDNA or RPA phosphorylation are being investigated as cancer chemotherapeutics, including mitigation of tumor resistance to genotoxic chemotherapy [61,62].

Once ATR is activated in response to replication stress, it phosphorylates and activates Chk1 kinase, which then phosphorylates downstream targets including CDK which regulates cell cycle progression. Activation of ATR and Chk1 are critical for the intra-S checkpoint in response to replication stress. This checkpoint enhances DNA repair, promotes protection of stalled replication forks, slows or stops cell cycle progression in S/G2 phases by preventing late origin firing to minimize replication fork encounters with blocking lesions, and activates dormant origins to rescue under-replicated DNA adjacent to blocked or collapsed forks [45,63–65]. Defects in ATR and other replication stress factors are implicated in many human diseases, including cancers, premature ageing, microcephaly, growth retardation, anemia, neurodegenerative disorders, ataxia, and developmental disorders [15].

Although PIKK signaling pathways display crosstalk, each PIKK has a dominant role in specific types of DSB repair. DNA-PK and ATM coordinate repair of two-ended DSBs by cNHEJ and HR, respectively [31,49], and ATR coordinates the replication stress response, including HR-mediated repair of seDSBs at collapsed replication forks [45,66].

3. Protecting and Rescuing Blocked Replication Forks

DNA replication initiates at origins in a complex, highly regulated process involving assembly of pre-replication complexes and licensing factors that ensure DNA is replicated only once per cell cycle [67]. For this reason, there is a premium on protecting replisomes at stalled replication forks to prevent replisome dissociation and fork collapse. Stalled forks are protected by a plethora of repair and replication factors, including RIF1, which inhibits end resection, the MRN-interacting protein MRNIP, the TLS suppressor USP1 which regulates PCNA via de-ubiquitination, HR proteins (RAD51, BRCA1, BRCA2, FANCD2), and RADX which regulates RAD51 [30,68–73]. Cells with defects in any of these fork protection factors are hypersensitive to replication stress.

Maintaining replisomes to protect stalled forks often involves fork regression to a 'chicken foot' structure that resembles four-way branched Holliday junctions of HR reactions (Figure 2A) [29]. Chicken foot structures have a one-ended DSB that at least initially includes ssDNA to which the HR factors RAD51, BRCA1, BRCA2, and the RAD51 paralogs (RAD51B/C/D and XRCC2/3) are recruited [74], although HR factors appear to play distinct roles in HR and fork protection [75]. Recent evidence indicates that fork reversal proceeds in two phases. Limited reversal is catalyzed by helicase-like chromatin remodeling proteins SMARCAL1, HLTF and PICH, the structure-specific nuclease ZRANB3, and RAD51 [29]. PICH has branch migration activity that helps extend fork reversal, which induces topological strain, thus extensive reversal requires topolsomerase II α (TopoII α) to relieve the strain. TopoII α is SUMOylated by ZATT, and SUMO-TopoII α then recruits PICH which branch migrates the four-way structure to further extend the reversed fork [29,76]. RAD51, BRCA1, and BRCA2 protect reversed forks from nuclease attack by MRE11, EXO1, DNA2, and MUS81 [77-79]. Part of the fork protection response involves histone methylation at stalled replication forks by EZH2, as this chromatin modification regulates MUS81 recruitment and subsequent nucleolytic attack of the protected fork [80]. It was recently shown that the WRN interacting protein WRNIP also protects reversed forks from nucleolytic attack [81]. Presumably seDSBs at protected forks are prevented from engaging in cNHEJ with other DSBs, i.e., seDSBs at other stressed forks or ends of frank, two-ended DSBs, to avert genome rearrangements. In cells with defects in any of these fork protection factors, reversed forks are rapidly degraded, accounting for their hypersensitivity to agents that induce replication stress. It has been hypothesized that extensive fork reversal is important to promote fork restart via HR [29,76]. In this model, extensive fork reversal allows sufficient end resection of seDSB ends to establish a functional RAD51-ssDNA nucleoprotein filament to drive HR restart. Forks reversed to a limited extent, by contrast, are preferentially restarted by RECQ1-mediated branch migration to achieve the same goal (Figure 2A).

There are several other mechanisms that repair and restart blocked replication forks, some of which bypass the blocking lesion. A blocked fork may be passively rescued by replication from an existing fork, or by checkpoint-activation of an adjacent dormant origin [65], replisomes may transiently switch templates to bypass a lesion, lesions may be bypassed by error-prone translesion DNA synthesis (TLS) polymerases, or a new fork may be established downstream of the blocking lesion by repriming by PRIMPOL and PRIM1 [6,64,82–87]. An advantage of this set of fork rescue pathways is that they do not create seDSBs and thus eliminate the threat of large-scale genome rearrangement due to cNHEJ-mediated seDSB misrepair. A disadvantage is that these leave behind unrepaired lesions (template switching) or a segment of under-replicated DNA (repriming), or they induce mutations (TLS). If stressed forks are rescued by an adjacent fork, the associated delay poses the risk that the stalled fork will reconfigure into toxic, branched structures catalyzed by HR factors [88].



Figure 2. Replication fork protection and restart. (**A**) Replication forks blocked by a DNA lesion (red star), or stalled by polymerase inhibitors or hydroxyurea, may reverse to a chicken foot, in two steps as shown. RAD51, BRCA2 and other factors protect the seDSB of the reversed fork from nucleolytic attack. Reversed forks may be restarted by RECQ1-mediated branch migration, or by RAD51-mediated strand invasion. (**B**) MUS81-EME1 cleaves four-way Holiday junctions, 3' flaps, and stalled replication forks, which causes fork collapse to a seDSB. MUS81-EME2 cleaves stalled forks to create seDSBs. (**C**) seDSBs at collapsed replication forks are resected to expose ssDNA which is bound by RAD51 to catalyze HR-mediated fork restart, analogous to break-induced replication (BIR).

The threats to genome integrity due to mutation, under-replicated DNA, and delayed restart can be avoided by engaging another set of fork restart mechanisms initiated when forks are cleaved by structure-specific endonucleases MUS81 or EEPD1. Similar to fork restart involving a chicken foot intermediate, fork cleavage restart mechanisms also create seDSBs. The roles of these fork cleavage nucleases and other nucleases/co-factors that contribute to restarting stressed replication forks via HR are described in the following sections.

4. MUS81: An Ancient Structure-Specific Nuclease Involved in HR and Restart of Stressed Replication Forks

MUS81 is a structure-specific 3' endonuclease in the XPF 3' endonuclease family that cleaves a variety of branched DNA structures including 3' flaps and Holliday junctions. Yeast Mus81 was first discovered in 2000 in a two-hybrid screen for proteins that interacted with the RAD54 HR protein and was named for the sensitivity of Mus81-defective cells to methyl methanesulfonate and UV light [89]. Mus81-defective yeast also have a severe meiotic HR defect that together with its interaction with RAD54 suggested an important role in HR [89]. Indeed, yeast Mus81 and its Eme1 cofactor resolve Holliday junctions and human MUS81 cleaves four-way (Holliday) junctions and 3' flap structures [90,91] (Figure 2B). In human cells, MUS81 with its EME1 cofactor resolves Holliday junctions in HR intermediates [92–95], and reversed forks that resemble Holliday junctions [96]. In contrast, MUS81 with its EME2 cofactor cleaves blocked replication forks, causing fork collapse to a seDSB [97-99] (Figure 2B). The seDSB is apparently resected to allow formation of a RAD51-ssDNA nucleoprotein filament that catalyzes fork restart by a mechanism that resembles break-induced replication (BIR) [100], although the resection nuclease(s) involved in processing MUS81-cleaved forks are not known (Figure 2C). Yeast Mus81 also mediates resolution of structures in G2/M that arise when blocked forks are rescued by converging forks to complete DNA replication [101].

MUS81 is an important DDR factor and a relevant tumor marker. MUS81 defects sensitize cells to various genotoxic chemicals [102,103], and it was recently shown that inhibition of MUS81 sensitizes HR-proficient cancer cells to the PARP1 inhibitor, olaparib [104], an agent commonly used to treat cancers with HR defects, such as BRCA1- and BRCA2-

defective breast cancers [105,106]. This suggests that inhibiting the HR functions of MUS81 is synthetically lethal with PARP inhibition, analogous to the synthetic lethality of PARP inhibitors in BRCA- and other HR-defective cells. This finding also suggests that MUS81 inhibition may be an alternative means to sensitize HR-proficient (i.e., BRCA-wildtype) tumors to PARP inhibitors [104]. MUS81 may also underlie an important cancer diagnostic. MUS81 was found to cleave DNA in prostate cancer cells, inducing cytosolic DNA that serves as a prostate tumor marker and promotes STING-dependent immune recognition to drive host rejection of tumor cells in vivo [107]. Interestingly, MUS81 foci correlated with cytosolic DNA levels that may reflect MUS81 cleavage of stressed replication forks [107] caused by oncogenic stress [39]. As noted above, BRCA1/2 help protect reversed replication forks, but in cells with BRCA2 defects, MUS81 is essential for cellular resistance to replication stress and proper chromosome segregation [108]. Because of their fork protection defect, reversed forks in BRCA2-defective cells are susceptible to nucleolytic attack by MRE11 in a reaction initiated by the CtIP nuclease, causing hypersensitivity to replication stress. In these BRCA2-defective cells, stressed fork rescue requires MUS81 cleavage to effect fork restart by an HR mechanism resembling BIR [78]. These findings suggest a treatment strategy for BRCA-defective cells in which this MUS81 fork restart pathway is blocked to enhance tumor killing by replication-stress inducing chemotherapy [78]; this strategy might provide similar benefits with radiotherapy. The roles of MUS81 in cancer depend on the tumor genetic background. Unlike its protective role in BRCA-defective tumors, MUS81 mediates chromosome shattering and apoptosis in cancer cells with microsatellite instability and a defect in the Werner syndrome helicase WRN [109].

Chk1 inhibitors have been explored as cancer chemotherapeutics, but these agents often cause severe side effects. In a recent study [110], Chk1 inhibition increased underreplicated DNA and mitotic defects, including anaphase bridges and intermediates of mitotic DNA synthesis (termed MiDAS). MiDAS completes replication of regions that fail to fully replicate during S phase as a result of replication stress, i.e., at common fragile sites [111,112]. MUS81-EME1 was shown to cleave nascent DNA generated during mitosis in response to insufficient nucleotide pools to maintain MiDAS, and this promoted chromosome instability but did not affect cell survival. In contrast, MUS81-EME2, which normally promotes fork restart, mediates cell death in Chk1-inhibited cells [110]. It was therefore suggested that caution be exercised with Chk1 inhibitors as such treatments may kill certain tumor cells, but those that survive may display chromosomal instability [110]. Such treatments could therefore promote progression of surviving tumor cells to a more aggressive state or increase the risk of secondary tumors [113,114].

5. EEPD1: A 5' Structure-Specific Endonuclease That Complements the 3' MUS81 Nuclease

EEPD1 (endo- exonuclease phosphatase domain protein 1) has a DNase I-like nuclease domain and a DNA binding domain with two helix-hairpin-helix motifs similar to those in prokaryotic RuvA2. As with MUS81, cells with defective EEPD1 are hypersensitive to a variety of genotoxic chemicals and radiation, and replication stress induces chromosome aberrations and mitotic catastrophe [115,116]. iPOND (isolation of proteins on nascent DNA) is a technique that reveals proteins associated with replication forks, including replisome components and proteins recruited to stressed forks [117,118]. iPOND analysis demonstrated that EEPD1 is recruited to stalled replication forks, and similar to MUS81, EEPD1 cleaves fork structures in vitro, and stalled replication forks in vivo (Figure 3) [115]. Once EEPD1 cleaves stalled replication forks, it promotes EXO1-mediated resection of the resulting seDSB to block cNHEJ and promote HR-mediated fork restart [115,116,119]. Resection defects are seen in EEPD1-defective cells at both stressed replication forks and frank DSBs, and these defects suppress ATR activation and downstream stress responses including induction of γ H2AX and Chk1 activation [115]. Replication stress is associated with rapid cell division (i.e., due to oncogenic stress or during embryonic development), and EEPD1 knockdown causes severe developmental defects during early vertebrate development [120]. Unlike MUS81, which evolved ~1.5 billion years ago, EEPD1 arose much later, appearing in chordates and early vertebrates ~500 million years ago. Interestingly, this corresponds to the period in evolution where genome size underwent two successive doublings [121]. It is tempting to speculate that the original MUS81 stressed fork cleavage system required additional assistance to manage increased replication stress associated with genome expansion, and this selective pressure gave rise to EEPD1. Another advantage to EEPD1 is that it is a 5' nuclease and therefore it cleaves the opposite strand at stalled forks as that cleaved by the MUS81 3' nuclease. Although cleavage of either strand produces seDSBs that can initiate HR-mediated fork restart, these distinct restart mechanisms may have different restart kinetics. As shown in Figure 3, MUS81's 3' endonuclease activity cleaves the template strand for leading-strand synthesis, whereas EEPD1's 5' endonuclease activity cleaves the template strand for lagging-strand synthesis. This polarity difference means that the seDSB end produced by MUS81 is forced to invade the lagging strand duplex, which remains discontinuous until Okazaki fragment maturation is complete. Strand invasion by MUS81 seDSBs may not be successful until they are resected enough to allow invasion into a mature lagging strand duplex. In contrast, EEPD1 fork cleavage allows the resected seDSB end to invade the (continuous) leading strand duplex, which requires less resection and therefore may provide a faster fork restart mechanism. There is evidence that even relatively short delays in fork restart can result in genome instability [115,120,122,123], probably because such delays increase the chance that stalled forks will be remodeled into toxic HR intermediates [6,88]. Hence, EEPD1 may have been selected during evolution because it provided an alternative and potentially faster fork restart mechanism to complement the ancient MUS81 mechanism, and thus help manage increased replication stress associated with larger genomes. Another reason EEPD1 may have provided a selective advantage during vertebrate genome evolution is because EEPD1 interacts with and recruits the EXO1 resection nuclease to seDSBs at collapsed replication forks, thereby promoting accurate fork restart by HR [119].

Inactivating mutations in EEPD1 are not seen in cancers, but EEPD1 is overexpressed in subsets of cancers of the brain, breast, colon, cervix, kidney, skin, lung, prostate, head and neck, and uterus [124]. This pattern of few/no mutations and relatively common overexpression is reminiscent of other important DDR factors, such as RAD51, and may reflect the critical nature of replication stress responses to cancer cell survival. Indeed, cancer cells face greater replication stress than normal cells due to dysregulated replication associated with oncogenic stress, nutrient deprivation, hypoxia, attacks by the immune system, and stress associated with genotoxic cancer treatments [17]. Thus, EEPD1 overexpression probably provides a selective advantage to tumor cells, and it may be an important contributor to tumor resistance to therapies that induce replication stress. As noted above, inhibition of MUS81 and BRCA defects are synthetically lethal with PARP1 inhibitors, [108,125]. BRCA defects are also synthetically lethal with defects in the RAD52 HR protein [126,127]. Importantly, BRCA-RAD52 synthetic lethality is suppressed by defects in EEPD1 [128], thus EEPD1 activities at stressed replication forks, and/or during HR repair of frank DSBs, apparently generate intermediates that require processing by either BRCA1/2 or RAD52 to prevent cell death. These findings suggest that although inhibition of RAD52 may provide benefits to patients with BRCA-defective tumors, therapeutic resistance may develop by downregulation or inactivation of EEPD1.



Figure 3. Distinct mechanisms of fork cleavage by MUS81 and EEPD1. (**Left**) The 3' MUS81 nuclease cleaves the leading template strand, producing a seDSB that must be resected to load RAD51. This strand is forced to invade the lagging strand duplex, but strand invasion and reestablishment of the fork may be obstructed if the invasion occurs in the region of immature Okazaki fragments. Fork restart may be delayed until further resection allows invasion into a region with fully mature Okazaki fragments, and/or by delaying invasion until maturation is complete. (**Right**) By cleaving the lagging strand template, EEPD1 avoids this problem as invasion will always occur in the continuous leading strand duplex.

6. Metnase: A Recently Evolved Nuclease-Protein Methyl Transferase That Promotes Replication Fork Restart

Metnase evolved ~50 million years ago when a *Mariner* transposon integrated downstream of a SET protein methylase, and subsequent genetic changes fused the SET and nuclease domains [129]. Metnase is a structure-specific nuclease with numerous genome stabilization functions including promotion of cNHEJ, chromosome decatenation, and restart of stressed replication forks [130–132]. Although defects in the Metnase nuclease delay replication fork restart [123] and Metnase cleaves replication fork structures in vitro [133], Metnase does not cleave stalled forks in vivo like MUS81 and EEPD1 [116]. These findings suggest that Metnase nuclease functions in a later step in replication fork restart, such as trimming flaps in HR-mediated fork repair intermediates [116]. The Metnase protein methylase targets histone H3 K36 to promote recruitment of cNHEJ factors Ku and NBS1 [134], Metnase automethylation regulates its chromosome decatenation function [135], and its methylase also plays an as yet undefined role in promoting restart of stressed replication forks [122]. Metnase is phosphorylated by Chk1, and this modification promotes cNHEJ, but suppresses replication fork restart [136]. Indeed, Metnase regulates Chk1 stability, suggesting a feedback loop between Metnase and Chk1 that coordinates DNA repair and checkpoint processes [137].

7. Other Nucleases with Known or Potential Roles in Replication Stress Responses: CtIP, MRE11, EXO1, DNA2-BLM, SLX1-SLX4, XPF-ERCC1-SLX4, Artemis, XPG, and FEN1

CtIP is a nuclease with important roles in the regulation and initial resection of frank DSBs in collaboration with MRE11 [43,138]. However, MRE11 must be restrained from degrading reversed replication forks [72,139]. In the context of reversed forks CtIP has a nuclease (and MRE11) independent role that protects reversed forks from nucleolytic attack by DNA2 [140]. This CtIP function is even more important in cells with diminished fork protection due to BRCA1 defects, suggesting CtIP as a novel therapeutic target to augment genotoxic cancer therapy of tumors with BRCA1 defects [140]. In a recent study CtIP was shown to be regulated by SUMOylation, a constitutive modification in S phase cells, and this modification was shown to be important for both CtIP roles in resection and in fork protection [141]. This raises the possibility of targeting the CtIP SUMO modification to augment cancer therapy.

EXO1, and DNA2 with its BLM cofactor, are responsible for extensive resection of frank DSB ends, exposing long ssDNA tracts that are first bound by RPA to trigger checkpoint responses as discussed above, before RPA is replaced by RAD51 for HR-mediated DSB repair. Resection also appears to be important at reversed replication forks to recruit RAD51, BRCA1/2 and other fork protection factors. At stressed forks cleaved by EEPD1, there is direct evidence that EEPD1 recruits EXO1 to ensure resection of the seDSB and accurate, HR-mediated fork restart [119]. Although Metnase doesn't cleave stressed forks, it also recruits EXO1 to promote resection of seDSBs at cleaved forks [133]. There is as yet no direct evidence that MUS81 similarly recruits EXO1 and/or DNA2-BLM to seDSBs; if MUS81 lacks this function, this is another likely selective advantage provided by the late-evolving EEPD1 and Metnase proteins.

The SLX4 scaffold protein interacts with many proteins, including three structurespecific nucleases, MUS81-EME1, XPF-ERCC1, and SLX1. These complexes mediate a broad range of DNA transactions that promote genome stability [142]. The SLX1 structurespecific nuclease with its SLX4 co-factor cleaves a variety of branched DNA structures in vitro, it contributes to genome stability by processing branched intermediates during HR, and it promotes inter-strand crosslink repair and telomere maintenance [142–144]. SLX4 also associates with the XPF-ERCC1 structure-specific nuclease and like SLX1-SLX4, this complex also cleaves a variety of branched DNA structures in vitro [142]. XPF-ERCC1 are involved in nucleotide excision repair and inter-strand crosslink repair, and it was recently shown that XPF-ERCC1 is important for DSB repair by HR when substrates form secondary structures, such as AT-rich and G-quadraplex sequences [145]. Although these types of structures can arise during HR repair of frank DSBs and at stressed replication forks, the roles of SLX1 and XPF-ERCC1 in replication stress responses are poorly understood. A recent study implicated both XPF and Artemis (which has nucleolytic roles in cNHEJ) in rapid cleavage of stressed replication forks, although siRNA knockdown of XPF, Artemis, or both proteins had relatively minor effects on the speed and efficiency of replication fork restart [146].

XPG and FEN1 are flap endonucleases with roles in nucleotide excision repair and HR. In addition to its primary role in suppressing replication stress by repairing bulky lesions, XPG was shown to have a non-catalytic role in promoting HR through interactions with RAD51, BRCA1, BRCA2, and PALB2, and the HR defect in XPG-mutant cells causes genome instability and decreases fork restart after HU-induced replication stress [147]. FEN1, named for its flap endonuclease activity, also has 5' exonuclease and gap endonuclease activities, and is involved in Okazaki fragment maturation, base excision repair, HR, and processing of stalled replication forks [148,149].

8. Perspectives

Despite the major advances in molecular characterization of tumors that inform targeted cancer therapies, the majority of cancer patients still receive non-targeted, genotoxic chemo- or radiotherapy, and these genotoxins universally cause replication stress. This has stimulated drug development efforts to augment chemo- and radiotherapy with agents that block general DDR signaling, such as inhibitors of ATM and ATR [150–152], as well as HR factors and replication stress nucleases (Table 1). Characterizing expression levels of MUS81, EEPD1, and Metnase may help illuminate tumor resistance to traditional therapeutics and inform personalized treatments, such as higher genotoxin doses to counteract the enhanced replication stress resistance associated with overexpression of these nucleases. Because of the central nature of DNA replication in cell division, replication stress responses also provide a rich environment for the development of targeted, synthetic lethal treatment strategies [34]. Increasing our understanding of DDR factors that specifically regulate replication responses, including the structure-specific nucleases discussed here, is likely to drive new approaches that exploit tumor dependence on specific replication stress response factors.

Table 1. Functions and inhibitors of key replication stress nucleases and co-factors.

Protein	Biochemical Activities	Biological Functions	Inhibitor References
RPA	Binds ssDNA, ATRIP, and itself	DNA replication and repair; activates ATR through ATRIP binding to RPA-bound ssDNA; replaced by RAD51 on ssDNA during HR	[61,62,153–155]
MRE11	DSB end binding, 3'-5' exonuclease, endonuclease	Early DSB sensor, ATM activation, promotes cNHEJ, initiates resection for HR	[156–158]
CtIP	Endonuclease	Promotes limited resection by MRE11	[159,160]
EXO1	5'-3' exonuclease	Extensive end resection	[161] *
DNA2	5'-3' exonuclease	Extensive end resection	[162,163]
BLM	3'-5' helicase	Unwinds DNA structures during HR, promotes resection by DNA2	[164–167]
RAD51	Strand invasion (recombinase)	Binds dsDNA, ssDNA and itself, catalyzes HR	[168–173]
MUS81-EME2	3' structure specific endonuclease	Cleaves stalled forks, promotes fork restart	[174]
EEPD1	5' structure specific endonuclease	Cleaves stalled forks, promotes fork restart and fork resection by EXO1	None ⁺
Metnase	5' structure specific endonuclease, protein methylase	Cleaves stalled forks, promotes fork restart and fork resection by EXO1	[175]
SLX1-SLX4	5' structure specific endonuclease	Cleaves branched structures, promotes HR, crosslink repair, and telomere maintenance	None ⁺
XPF-ERCC1	5' structure specific endonuclease	Nucleotide excision repair, inter-strand crosslink repair, HR (replication stress?)	[176,177]

* EXO1 activity inhibited indirectly by diallyl disulfide through reduced protein levels. [†] These proteins have been inhibited by using siRNA knockdown.

Author Contributions: Conceptualization, J.A.N. and R.H.; writing—original draft preparation, J.A.N.; writing—review and editing, J.A.N., N.S., L.T., S.J.A. and R.H.; funding acquisition, J.A.N. and R.H. All authors have read and agreed to the published version of the manuscript.

Funding: Research in the Nickoloff lab was supported by NIH General Medical Sciences grant number R01 GM084020 and American Lung Association grant number LCD-686552. Research in the Hromas lab was supported by NIH National Cancer Institute grant number R01 CA139429.

Acknowledgments: We thank our colleagues in the Colorado State University College of Veterinary Medicine and Biomedical Science, and the University of Texas Health Sciences Center, San Antonio for many helpful discussions.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Löb, D.; Lengert, N.; Chagin, V.O.; Reinhart, M.; Casas-Delucchi, C.S.; Cardoso, M.C.; Drossel, B. 3D replicon distributions arise from stochastic initiation and domino-like DNA replication progression. *Nat. Commun.* **2016**, *7*, 11207. [CrossRef] [PubMed]
- Chagin, V.O.; Casas-Delucchi, C.S.; Reinhart, M.; Schermelleh, L.; Markaki, Y.; Maiser, A.; Bolius, J.J.; Bensimon, A.; Fillies, M.; Domaing, P.; et al. 4D Visualization of replication foci in mammalian cells corresponding to individual replicons. *Nat. Commun.* 2016, 7, 11231. [CrossRef] [PubMed]
- 3. IIyer, D.R.; Rhind, N. The Intra-S Checkpoint Responses to DNA Damage. Genes 2017, 8, 74. [CrossRef] [PubMed]
- Sarmiento-Salinas, F.L.; Perez-Gonzalez, A.; Acosta-Casique, A.; Ix-Ballote, A.; Diaz, A.; Trevino, S.; Rosas-Murrieta, N.H.; Millán-Perez-Peña, L.; Maycotte., P. Reactive Oxygen Species: Role in Carcinogenesis, Cancer Cell Signaling and Tumor Progression. *Life Sci.* 2021, 284, 119942. [CrossRef] [PubMed]
- Tubbs, A.; Nussenzweig, A. Endogenous DNA Damage as a Source of Genomic Instability in Cancer. *Cell* 2017, 168, 644–656. [CrossRef]
- Nickoloff, J.A.; Sharma, N.; Taylor, L.; Allen, S.J.; Hromas, R. The Safe Path at the Fork: Ensuring Replication-Associated DNA Double-Strand Breaks are Repaired by Homologous Recombination. *Front. Genet.* 2021, 12. [CrossRef] [PubMed]
- Conti, B.A.; Smogorzewska, A. Mechanisms of direct replication restart at stressed replisomes. DNA Repair 2020, 95, 102947. [CrossRef]
- Lyu, X.; Chastain, M.; Chai, W. Genome-Wide Mapping and Profiling of Gammah2ax Binding Hotspots in Response to Different Replication Stress Inducers. BMC Genom. 2019, 20, 579. [CrossRef]
- 9. Poggi, L.; Richard, G.F. Alternative DNA Structures in Vivo: Molecular Evidence and Remaining Questions. *Microbiol. Mol. Biol. Rev.* 2020, *85*, e00110–e00120. [CrossRef]
- 10. Spiegel, J.; Adhikari, S.; Balasubramanian, S. The Structure and Function of DNA G-Quadruplexes. *Trends Chem.* **2019**, *2*, 123–136. [CrossRef] [PubMed]
- Barlow, J.H.; Faryabi, R.B.; Callen, E.; Wong, N.; Malhowski, A.; Chen, H.T.; Gutierrez-Cruz, G.; Sun, H.W.; McKinnon, P.; Wright, G.; et al. Identification of Early Replicating Fragile Sites That Contribute to Genome Instability. *Cell* 2013, 152, 620–632. [CrossRef] [PubMed]
- 12. Kaushal, S.; Freudenreich, C.H. The role of fork stalling and DNA structures in causing chromosome fragility. *Genes Chromosom. Cancer* **2018**, *58*, 270–283. [CrossRef] [PubMed]
- 13. Bochman, M.L.; Paeschke, K.; Zakian, V.A. DNA secondary structures: Stability and function of G-quadruplex structures. *Nat. Rev. Genet.* **2012**, *13*, 770–780. [CrossRef]
- 14. Gadaleta, M.C.; Noguchi, E. Regulation of DNA Replication through Natural Impediments in the Eukaryotic Genome. *Genes* 2017, *8*, 98. [CrossRef] [PubMed]
- 15. Zeman, M.K.; Cimprich, K.A. Causes and consequences of replication stress. Nat. Cell Biol. 2013, 16, 2–9. [CrossRef] [PubMed]
- Cicconi, A.; Chang, S. Shelterin and the replisome: At the intersection of telomere repair and replication. *Curr. Opin. Genet. Dev.* 2020, 60, 77–84. [CrossRef] [PubMed]
- 17. Gaillard, H.; Garcia-Muse, T.; Aguilera, A. Replication stress and cancer. Nat. Cancer 2015, 15, 276–289. [CrossRef]
- Santos-Pereira, J.M.; Aguilera, A. R loops: New modulators of genome dynamics and function. *Nat. Rev. Genet.* 2015, 16, 583–597. [CrossRef]
- 19. Hamperl, S.; Bocek, M.J.; Saldivar, J.C.; Swigut, T.; Cimprich, K.A. Transcription-Replication Conflict Orientation Modulates R-Loop Levels and Activates Distinct DNA Damage Responses. *Cell* **2017**, *170*, 774–786. [CrossRef]
- 20. Freudenreich, C.H. R-loops: Targets for nuclease cleavage and repeat instability. Curr. Genet. 2018, 64, 789–794. [CrossRef]
- 21. Ivessa, A.; Lenzmeier, B.A.; Bessler, J.B.; Goudsouzian, L.K.; Schnakenberg, S.L.; Zakian, V.A. The *Saccharomyces cerevisiae* Helicase Rrm3p Facilitates Replication Past Nonhistone Protein-DNA Complexes. *Mol. Cell* **2003**, *12*, 1525–1536. [CrossRef]
- 22. Billard, P.; A Poncet, D. Replication Stress at Telomeric and Mitochondrial DNA: Common Origins and Consequences on Ageing. *Int. J. Mol. Sci.* 2019, 20, 4959. [CrossRef]
- 23. Kotsantis, P.; Silva, L.M.; Irmscher, S.; Jones, R.; Folkes, L.; Gromak, N.; Petermann, E. Increased global transcription activity as a mechanism of replication stress in cancer. *Nat. Commun.* **2016**, *7*, 13087. [CrossRef] [PubMed]
- 24. Bermejo, R.; Lai, M.S.; Foiani, M. Preventing Replication Stress to Maintain Genome Stability: Resolving Conflicts between Replication and Transcription. *Mol. Cell* **2012**, *45*, 710–718. [CrossRef] [PubMed]
- Gómez-González, B.; Aguilera, A. Transcription-mediated replication hindrance: A major driver of genome instability. *Genes Dev.* 2019, 33, 1008–1026. [CrossRef]
- 26. Hamperl, S.; Cimprich, K.A. Conflict Resolution in the Genome: How Transcription and Replication Make It Work. *Cell* **2016**, *167*, 1455–1467. [CrossRef]
- 27. Garcia-Muse, T.; Aguilera, A. Transcription–replication conflicts: How they occur and how they are resolved. *Nat. Rev. Mol. Cell Biol.* **2016**, *17*, 553–563. [CrossRef]
- Minchell, N.E.; Keszthelyi, A.; Baxter, J. Cohesin Causes Replicative DNA Damage by Trapping DNA Topological Stress. *Mol. Cell* 2020, 78, 739–751.e8. [CrossRef]
- 29. Qiu, S.; Jiang, G.; Cao, L.; Huang, J. Replication Fork Reversal and Protection. Front. Cell Dev. Biol. 2021, 9, 670392. [CrossRef]
- 30. Rickman, K.; Smogorzewska, A. Advances in understanding DNA processing and protection at stalled replication forks. *J. Cell Biol.* **2019**, *218*, 1096–1107. [CrossRef]

- Chang, H.H.Y.; Pannunzio, N.R.; Adachi, N.; Lieber, M.R. Non-homologous DNA end joining and alternative pathways to double-strand break repair. *Nat. Rev. Mol. Cell Biol.* 2017, 18, 495–506. [CrossRef]
- Scully, R.; Panday, A.; Elango, R.; Willis, N.A. DNA double-strand break repair-pathway choice in somatic mammalian cells. *Nat. Rev. Mol. Cell Biol.* 2019, 20, 698–714. [CrossRef]
- 33. Toledo, F. Mechanisms Generating Cancer Genome Complexity: Back to the Future. Cancers 2020, 12, 3783. [CrossRef]
- 34. Nickoloff, J.A.; Jones, D.; Lee, S.-H.; Williamson, E.A.; Hromas, R. Drugging the Cancers Addicted to DNA Repair. *JNCI J. Natl. Cancer Inst.* 2017, *109*, djx059. [CrossRef]
- Chatterjee, N.; Walker, G.C. Mechanisms of DNA Damage, Repair, and Mutagenesis. *Environ. Mol. Mutagen.* 2017, 58, 235–263. [CrossRef] [PubMed]
- 36. Jossen, R.; Bermejo, R. The DNA damage checkpoint response to replication stress: A Game of Forks. *Front. Genet.* **2013**, *4*, 26. [CrossRef]
- Hills, S.A.; Diffley, J.F. DNA Replication and Oncogene-Induced Replicative Stress. Curr. Biol. 2014, 24, R435–R444. [CrossRef] [PubMed]
- 38. Primo, L.M.F.; Teixeira, L.K. DNA replication stress: Oncogenes in the spotlight. Genet. Mol. Biol. 2020, 43, e20190138. [CrossRef]
- Kotsantis, P.; Petermann, E.; Boulton, S.J. Mechanisms of Oncogene-Induced Replication Stress: Jigsaw Falling into Place. *Cancer Discov.* 2018, *8*, 537–555. [CrossRef] [PubMed]
- 40. Cortez, D. Unwind and slow down: Checkpoint activation by helicase and polymerase uncoupling: Figure 1. *Genes Dev.* **2005**, *19*, 1007–1012. [CrossRef]
- Berti, M.; Cortez, D.; Lopes, M. The plasticity of DNA replication forks in response to clinically relevant genotoxic stress. *Nat. Rev. Mol. Cell Biol.* 2020, 21, 633–651. [CrossRef] [PubMed]
- 42. Daley, J.M.; Niu, H.; Miller, A.S.; Sung, P. Biochemical mechanism of DSB end resection and its regulation. *DNA Repair* 2015, 32, 66–74. [CrossRef] [PubMed]
- Symington, L.S. Mechanism and regulation of DNA end resection in eukaryotes. *Crit. Rev. Biochem. Mol. Biol.* 2016, 51, 195–212. [CrossRef] [PubMed]
- 44. Zhao, F.; Kim, W.; Kloeber, J.A.; Lou, Z. DNA end resection and its role in DNA replication and DSB repair choice in mammalian cells. *Exp. Mol. Med.* **2020**, *52*, 1705–1714. [CrossRef] [PubMed]
- Yazinski, S.A.; Zou, L. Functions, Regulation, and Therapeutic Implications of the ATR Checkpoint Pathway. *Annu. Rev. Genet.* 2016, 50, 155–173. [CrossRef]
- 46. Shiotani, B.; Nguyen, H.D.; Håkansson, P.; Maréchal, A.; Tse, A.; Tahara, H.; Zou, L. Two Distinct Modes of ATR Activation Orchestrated by Rad17 and Nbs1. *Cell Rep.* **2013**, *3*, 1651–1662. [CrossRef]
- 47. Haahr, P.; Hoffmann, S.; Tollenaere, M.; Ho, T.; Toledo, L.; Mann, M.; Bekker-Jensen, S.; Räschle, M.; Mailand, N. Activation of the ATR kinase by the RPA-binding protein ETAA1. *Nat. Cell Biol.* **2016**, *18*, 1196–1207. [CrossRef]
- Blackford, A.N.; Jackson, S.P. ATM, ATR, and DNA-PK: The Trinity at the Heart of the DNA Damage Response. *Mol. Cell* 2017, 66, 801–817. [CrossRef] [PubMed]
- 49. Shibata, A.; Jeggo, P.A. ATM's Role in the Repair of DNA Double-Strand Breaks. Genes 2021, 12, 1370. [CrossRef]
- 50. Saldivar, J.C.; Cortez, D.; Cimprich, K.A. The Essential Kinase ATR: Ensuring Faithful Duplication of a Challenging Genome. *Nat. Rev. Mol. Cell Biol.* **2017**, *18*, 622–636. [CrossRef]
- 51. Liu, S.; Opiyo, S.O.; Manthey, K.; Glanzer, J.G.; Ashley, A.K.; Amerin, C.; Troksa, K.; Shrivastav, M.; Nickoloff, J.A.; Oakley, G.G. Distinct roles for DNA-PK, ATM and ATR in RPA phosphorylation and checkpoint activation in response to replication stress. *Nucleic Acids Res.* 2012, 40, 10780–10794. [CrossRef] [PubMed]
- 52. Anantha, R.W.; Vassin, V.M.; Borowiec, J.A. Sequential and Synergistic Modification of Human RPA Stimulates Chromosomal DNA Repair. *J. Biol. Chem.* 2007, *282*, 35910–35923. [CrossRef] [PubMed]
- Olson, E.; Nievera, C.J.; Klimovich, V.; Fanning, E.; Wu, X. RPA2 Is a Direct Downstream Target for ATR to Regulate the S-phase Checkpoint. J. Biol. Chem. 2006, 281, 39517–39533. [CrossRef] [PubMed]
- 54. Maréchal, A.; Zou, L. RPA-coated single-stranded DNA as a platform for post-translational modifications in the DNA damage response. *Cell Res.* **2014**, *25*, 9–23. [CrossRef]
- 55. Zuazua-Villar, P.; Ganesh, A.; Phear, G.; Gagou, M.E.; Meuth, M. Extensive RPA2 hyperphosphorylation promotes apoptosis in response to DNA replication stress in CHK1 inhibited cells. *Nucleic Acids Res.* **2015**, *43*, 9776–9787. [CrossRef]
- 56. Vassin, V.M.; Anantha, R.W.; Sokolova, E.; Kanner, S.; Borowiec, J.A. Human RPA phosphorylation by ATR stimulates DNA synthesis and prevents ssDNA accumulation during DNA-replication stress. *J. Cell Sci.* 2009, 122, 4070–4080. [CrossRef]
- 57. Soniat, M.M.; Myler, L.R.; Kuo, H.-C.; Paull, T.T.; Finkelstein, I.J. RPA Phosphorylation Inhibits DNA Resection. *Mol. Cell* 2019, 75, 145–153.e5. [CrossRef]
- Patrick, S.M.; Oakley, G.G.; Dixon, K.; Turchi, J.J. DNA Damage Induced Hyperphosphorylation of Replication Protein A. 2. Characterization of DNA Binding Activity, Protein Interactions, and Activity in DNA Replication and Repair. *Biochemistry* 2005, 44, 8438–8448. [CrossRef]
- 59. Oakley, G.G.; Patrick, S.M.; Yao, J.; Carty, M.P.; Turchi, J.J.; Dixon, K. RPA Phosphorylation in Mitosis Alters DNA Binding and Protein–Protein Interactions. *Biochemistry* **2003**, *42*, 3255–3264. [CrossRef]

- Marechal, A.; Li, J.M.; Ji, X.Y.; Wu, C.S.; Yazinski, S.A.; Nguyen, H.D.; Liu, S.; Jimenez, A.E.; Jin, J.; Zou, L. PRP19 Transforms into a Sensor of RPA-ssDNA after DNA Damage and Drives ATR Activation Via a Ubiquitin-Mediated Circuitry. *Mol. Cell* 2014, 53, 235–246. [CrossRef]
- Mishra, A.K.; Dormi, S.S.; Turchi, A.M.; Woods, D.S.; Turchi, J.J. Chemical inhibitor targeting the replication protein A–DNA interaction increases the efficacy of Pt-based chemotherapy in lung and ovarian cancer. *Biochem. Pharmacol.* 2014, 93, 25–33. [CrossRef] [PubMed]
- 62. Glanzer, J.G.; Liu, S.; Wang, L.; Mosel, A.; Peng, A.; Oakley, G.G. RPA Inhibition Increases Replication Stress and Suppresses Tumor Growth. *Cancer Res.* 2014, 74, 5165–5172. [CrossRef] [PubMed]
- 63. Sadoughi, F.; Hallajzadeh, J.; Asemi, Z.; Mansournia, M.A.; Alemi, F.; Yousefi, B. Signaling pathways involved in cell cycle arrest during the DNA breaks. *DNA Repair* **2021**, *98*, 103047. [CrossRef] [PubMed]
- 64. Brambati, A.; Zardoni, L.; Achar, Y.J.; Piccini, D.; Galanti, L.; Colosio, A.; Foiani, M.; Liberi, G. Dormant origins and fork protection mechanisms rescue sister forks arrested by transcription. *Nucleic Acids Res.* **2017**, *46*, 1227–1239. [CrossRef]
- 65. Courtot, L.; Hoffmann, J.-S.; Bergoglio, V. The Protective Role of Dormant Origins in Response to Replicative Stress. *Int. J. Mol. Sci.* **2018**, *19*, 3569. [CrossRef]
- Williams, R.M.; Zhang, X. Roles of ATM and ATR in DNA double strand breaks and replication stress. *Prog. Biophys. Mol. Biol.* 2021, 163, 109–119. [CrossRef] [PubMed]
- 67. Parker, M.W.; Botchan, M.R.; Berger, J.M. Mechanisms and regulation of DNA replication initiation in eukaryotes. *Crit. Rev. Biochem. Mol. Biol.* 2017, 52, 107–144. [CrossRef]
- Mukherjee, C.; Tripathi, V.; Manolika, E.M.; Heijink, A.M.; Ricci, G.; Merzouk, S.; De Boer, H.R.; Demmers, J.; Van Vugt, M.A.T.M.; Chaudhuri, A.R. RIF1 promotes replication fork protection and efficient restart to maintain genome stability. *Nat. Commun.* 2019, 10, 3287. [CrossRef]
- 69. Bennett, L.G.; Wilkie, A.M.; Antonopoulou, E.; Ceppi, I.; Sanchez, A.; Vernon, E.G.; Gamble, A.; Myers, K.N.; Collis, S.J.; Cejka, P.; et al. MRNIP is a replication fork protection factor. *Sci. Adv.* **2020**, *6*, eaba5974. [CrossRef]
- Lim, K.S.; Li, H.; Roberts, E.; Gaudiano, E.F.; Clairmont, C.; Sambel, L.A.; Ponnienselvan, K.; Liu, J.C.; Yang, C.; Kozono, D.; et al. USP1 Is Required for Replication Fork Protection in BRCA1-Deficient Tumors. *Mol. Cell* 2018, 72, 925–941.e4. [CrossRef]
- Rickman, K.A.; Noonan, R.J.; Lach, F.; Sridhar, S.; Wang, A.; Abhyankar, A.; Huang, A.; Kelly, M.; Auerbach, A.D.; Smogorzewska, A. Distinct roles of BRCA2 in replication fork protection in response to hydroxyurea and DNA interstrand cross-links. *Genes Dev.* 2020, 34, 832–846. [CrossRef] [PubMed]
- Schlacher, K.; Wu, H.; Jasin, M. A Distinct Replication Fork Protection Pathway Connects Fanconi Anemia Tumor Suppressors to RAD51-BRCA1/2. *Cancer Cell* 2012, 22, 106–116. [CrossRef]
- 73. Bhat, K.; Krishnamoorthy, A.; Dungrawala, H.; Garcin, E.B.; Modesti, M.; Cortez, D. RADX Modulates RAD51 Activity to Control Replication Fork Protection. *Cell Rep.* **2018**, *24*, 538–545. [CrossRef] [PubMed]
- 74. Berti, M.; Teloni, F.; Mijic, S.; Ursich, S.; Fuchs, J.; Palumbieri, M.D.; Krietsch, J.; Schmid, J.; Garcin, E.B.; Gon, S.; et al. Sequential role of RAD51 paralog complexes in replication fork remodeling and restart. *Nat. Commun.* **2020**, *11*, 1–12. [CrossRef] [PubMed]
- 75. Tye, S.; Ronson, G.E.; Morris, J.R. A fork in the road: Where homologous recombination and stalled replication fork protection part ways. *Semin. Cell Dev. Biol.* **2020**, *113*, 14–26. [CrossRef]
- 76. Tian, T.; Bu, M.; Chen, X.; Ding, L.; Yang, Y.; Han, J.; Feng, X.-H.; Xu, P.; Liu, T.; Ying, S.; et al. The ZATT-TOP2A-PICH Axis Drives Extensive Replication Fork Reversal to Promote Genome Stability. *Mol. Cell* **2020**, *81*, 198–211.e6. [CrossRef]
- 77. Thangavel, S.; Berti, M.; Levikova, M.; Pinto, C.; Gomathinayagam, S.; Vujanovic, M.; Zellweger, R.; Moore, H.; Lee, E.H.; Hendrickson, E.A.; et al. DNA2 Drives Processing and Restart of Reversed Replication Forks in Human Cells. *J. Cell Biol.* 2015, 208, 545–562. [CrossRef]
- Lemaçon, D.; Jackson, J.; Quinet, A.; Brickner, J.R.; Li, S.; Yazinski, S.; You, Z.; Ira, G.; Zou, L.; Mosammaparast, N.; et al. MRE11 and EXO1 nucleases degrade reversed forks and elicit MUS81-dependent fork rescue in BRCA2-deficient cells. *Nat. Commun.* 2017, *8*, 1–12. [CrossRef]
- 79. Mijic, S.; Zellweger, R.; Chappidi, N.; Berti, M.; Jacobs, K.; Mutreja, K.; Ursich, S.; Chaudhuri, A.R.; Nussenzweig, A.; Janscak, P.; et al. Replication fork reversal triggers fork degradation in BRCA2-defective cells. *Nat. Commun.* **2017**, *8*, 859. [CrossRef]
- Rondinelli, B.; Gogola, E.; Yücel, H.; Duarte, A.A.; Van De Ven, M.; Van Der Sluijs, R.; Konstantinopoulos, P.A.; Jonkers, J.; Ceccaldi, R.; Rottenberg, S.; et al. EZH2 promotes degradation of stalled replication forks by recruiting MUS81 through histone H3 trimethylation. *Nat. Cell Biol.* 2017, *19*, 1371–1378. [CrossRef]
- 81. Porebski, B.; Wild, S.; Kummer, S.; Scaglione, S.; Gaillard, P.-H.L.; Gari, K. WRNIP1 Protects Reversed DNA Replication Forks from SLX4-Dependent Nucleolytic Cleavage. *iScience* 2019, 21, 31–41. [CrossRef] [PubMed]
- Quinet, A.; Tirman, S.; Cybulla, E.; Meroni, A.; Vindigni, A. To skip or not to skip: Choosing repriming to tolerate DNA damage. *Mol. Cell* 2021, *81*, 649–658. [CrossRef] [PubMed]
- 83. Goodman, M.F.; Woodgate, R. Translesion DNA Polymerases. *Cold Spring Harb. Perspect. Biol.* 2013, 5, a010363. [CrossRef] [PubMed]
- Ma, X.; Tang, T.; Guo, C. Regulation of translesion DNA synthesis in mammalian cells. *Environ. Mol. Mutagen.* 2020, 61, 680–692. [CrossRef] [PubMed]
- Yekezare, M.; Gomez-Gonzalez, B.; Diffley, J.F. Controlling DNA Replication Origins in Response to DNA Damage—Inhibit Globally, Activate Locally. J. Cell Sci. 2013, 126, 1297–1306. [CrossRef]

- Lehmann, C.P.; Jiménez-Martín, A.; Branzei, D.; Tercero, J.A. Prevention of unwanted recombination at damaged replication forks. *Curr. Genet.* 2020, *66*, 1045–1051. [CrossRef]
- Vaisman, A.; Woodgate, R. Translesion DNA Polymerases in Eukaryotes: What Makes Them Tick? *Crit. Rev. Biochem. Mol. Biol.* 2017, 52, 274–303. [CrossRef]
- Fabre, F.; Chan, A.; Heyer, W.D.; Gangloff, S. Alternate Pathways Involving Sgs1/Top3, Mus81/Mms4, and Srs2 Prevent Formation of Toxic Recombination Intermediates from Single-Stranded Gaps Created by DNA Replication. *Proc. Natl. Acad. Sci.* USA 2002, 99, 16887–16892. [CrossRef]
- 89. Interthal, H.; Heyer, W.-D. MUS81 encodes a novel Helix-hairpin-Helix protein involved in the response to UV- and methylationinduced DNA damage in Saccharomyces cerevisiae. *J. Cell Biol.* 2000, 263, 812–827. [CrossRef]
- 90. Boddy, M.N.; Gaillard, P.H.L.; McDonald, W.H.; Shanahan, P.; Yates, J.R., 3rd; Russell., P. Mus81-Eme1 Are Essential Components of a Holliday Junction Resolvase. *Cell* 2001, 107, 537–548. [CrossRef]
- Chen, X.-B.; Melchionna, R.; Aliouat-Denis, C.-M.; Gaillard, P.-H.; Blasina, A.; Van de Weyer, I.; Boddy, M.N.; Russell, P.; Vialard, J.; McGowan, C.H. Human Mus81-Associated Endonuclease Cleaves Holliday Junctions In Vitro. *Mol. Cell* 2001, *8*, 1117–1127. [CrossRef]
- Dehé, P.M.; Coulon, S.; Scaglione, S.; Shanahan, P.; Takedachi, A.; Wohlschlegel, J.A.; Yates, J.R., 3rd; Llorente, B.; Russell, P.; Gaillard, P.H. Regulation of Mus81-Eme1 Holliday Junction Resolvase in Response to DNA Damage. *Nat. Struct. Mol. Biol.* 2013, 20, 598–603. [CrossRef] [PubMed]
- 93. Naim, V.; Wilhelm, T.; Debatisse, M.; Rosselli, F. Ercc1 and Mus81-Eme1 Promote Sister Chromatid Separation by Processing Late Replication Intermediates at Common Fragile Sites During Mitosis. *Nat. Cell Biol.* **2013**, *15*, 1008–1015. [CrossRef]
- 94. Wyatt, H.D.; Sarbajna, S.; Matos, J.; West, S.C. Coordinated Actions of SLX1-SLX4 and MUS81-EME1 for Holliday Junction Resolution in Human Cells. *Mol. Cell* **2013**, *52*, 234–247. [CrossRef] [PubMed]
- Sarbajna, S.; Davies, D.; West, S.C. Roles of SLX1–SLX4, MUS81–EME1, and GEN1 in avoiding genome instability and mitotic catastrophe. *Genes Dev.* 2014, 28, 1124–1136. [CrossRef]
- 96. Amangyeld, T.; Shin, Y.-K.; Lee, M.; Kwon, B.; Seo, Y.-S. Human MUS81-EME2 can cleave a variety of DNA structures including intact Holliday junction and nicked duplex. *Nucleic Acids Res.* **2014**, *42*, 5846–5862. [CrossRef]
- 97. Pepe, A.; West, S.C. MUS81-EME2 Promotes Replication Fork Restart. Cell Rep. 2014, 7, 1048–1055. [CrossRef]
- 98. Pepe, A.; West, S.C. Substrate Specificity of the Mus81-Eme2 Structure Selective Endonuclease. *Nucleic Acids Res.* 2014, 42, 3833–3845. [CrossRef]
- Gao, H.; Chen, X.-B.; McGowan, C.H. Mus81 Endonuclease Localizes to Nucleoli and to Regions of DNA Damage in Human S-phase Cells. *Mol. Biol. Cell* 2003, 14, 4826–4834. [CrossRef]
- 100. Kramara, J.; Osia, B.; Malkova, A. Break-Induced Replication: The Where, the Why, and the How. *Trends Genet.* **2018**, *34*, 518–531. [CrossRef]
- 101. Pardo, B.; Moriel-Carretero, M.; Vicat, T.; Aguilera, A.; Pasero, P. Homologous recombination and Mus81 promote replication completion in response to replication fork blockage. *EMBO Rep.* 2020, 21, e49367. [CrossRef] [PubMed]
- Dendouga, N.; Gao, H.; Moechars, D.; Janicot, M.; Vialard, J.; McGowan, C.H. Disruption of Murine Mus81 Increases Genomic Instability and DNA Damage Sensitivity but Does Not Promote Tumorigenesis. *Mol. Cell. Biol.* 2005, 25, 7569–7579. [CrossRef] [PubMed]
- Xie, S.; Zheng, H.; Wen, X.; Sun, J.; Wang, Y.; Gao, X.; Guo, L.; Lu, R. MUS81 is associated with cell proliferation and cisplatin sensitivity in serous ovarian cancer. *Biophys. Res. Commun.* 2016, 476, 493–500. [CrossRef] [PubMed]
- Zhong, A.; Zhang, H.; Xie, S.; Deng, M.; Zheng, H.; Wang, Y.; Chen, M.; Lu, R.; Guo, L. Inhibition of MUS81 improves the chemical sensitivity of olaparib by regulating MCM2 in epithelial ovarian cancer. *Oncol. Rep.* 2018, 39, 1747–1756. [CrossRef]
- 105. D'Andrea, A.D. Mechanisms of PARP inhibitor sensitivity and resistance. DNA Repair 2018, 71, 172–176. [CrossRef]
- 106. Pommier, Y.; O'Connor, M.J.; de Bono, J. Laying a Trap to Kill Cancer Cells: Parp Inhibitors and Their Mechanisms of Action. *Sci. Transl. Med.* **2016**, *8*, 362ps17. [CrossRef]
- 107. Ho, S.S.W.; Zhang, W.Y.L.; Tan, N.Y.J.; Khatoo, M.; Suter, M.A.; Tripathi, S.; Cheung, F.S.G.; Lim, W.K.; Tan, P.H.; Ngeow, J.; et al. The DNA Structure-Specific Endonuclease MUS81 Mediates DNA Sensor STING-Dependent Host Rejection of Prostate Cancer Cells. *Immunity* 2016, 44, 1177–1189. [CrossRef]
- 108. Lai, X.; Broderick, R.; Bergoglio, V.; Zimmer, J.; Badie, S.; Niedzwiedz, W.; Hoffmann, J.S.; Tarsounas., M. Mus81 Nuclease Activity Is Essential for Replication Stress Tolerance and Chromosome Segregation in Brca2-Deficient Cells. *Nat. Commun.* 2017, *8*, 15983. [CrossRef]
- 109. van Wietmarschen, N.; Sridharan, S.; Nathan, W.J.; Tubbs, A.; Chan, E.M.; Callen, E.; Wu, W.; Belinky, F.; Tripathi, V.; Wong, N.; et al. Repeat expansions confer WRN dependence in microsatellite-unstable cancers. *Nature* **2020**, *586*, 292–298. [CrossRef]
- Calzetta, N.L.; Besteiro, M.A.G.; Gottifredi, V. Mus81-Eme1–dependent aberrant processing of DNA replication intermediates in mitosis impairs genome integrity. Sci. Adv. 2020, 6, eabc8257. [CrossRef]
- 111. Epum, E.A.; Haber, J.E. DNA replication: The recombination connection. Trends Cell Biol. 2021, 32, 45–57. [CrossRef]
- 112. Özer, Ö.; Hickson, I.D. Pathways for maintenance of telomeres and common fragile sites during DNA replication stress. *Open Biol.* **2018**, *8*, 180018. [CrossRef] [PubMed]
- Brower, V. Tracking Chemotherapy's Effects on Secondary Cancers. JNCI J. Natl. Cancer Inst. 2013, 105, 1421–1422. [CrossRef] [PubMed]

- Kamran, S.; De Gonzalez, A.B.; Ng, A.; Haas-Kogan, D.; Viswanathan, A.N. Therapeutic radiation and the potential risk of second malignancies. *Cancer* 2016, 122, 1809–1821. [CrossRef]
- 115. Wu, Y.; Lee, S.H.; Williamson, E.A.; Reinert, B.L.; Cho, J.H.; Xia, F.; Jaiswal, A.S.; Srinivasan, G.; Patel, B.; Brantley, A.; et al. Eepd1 Rescues Stressed Replication Forks and Maintains Genome Stability by Promoting End Resection and Homologous Recombination Repair. *PLoS Genet.* 2015, *11*, e1005675. [CrossRef]
- 116. Sharma, N.; Speed, M.C.; Allen, C.P.; Maranon, D.G.; Williamson, E.; Singh, S.; Hromas, R.; Nickoloff, J.A. Distinct roles of structure-specific endonucleases EEPD1 and Metnase in replication stress responses. *NAR Cancer* 2020, 2, zcaa008. [CrossRef] [PubMed]
- 117. Cortez, D. Proteomic Analyses of the Eukaryotic Replication Machinery. Methods Enzymol. 2017, 591, 33–53. [CrossRef]
- 118. Sirbu, B.M.; Couch, F.B.; Feigerle, J.T.; Bhaskara, S.; Hiebert, S.W.; Cortez., D. Analysis of Protein Dynamics at Active, Stalled, and Collapsed Replication Forks. *Genes. Dev.* 2011, 25, 1320–1327. [CrossRef]
- 119. Kim, H.-S.; Nickoloff, J.A.; Wu, Y.; Williamson, E.A.; Sidhu, G.S.; Reinert, B.L.; Jaiswal, A.S.; Srinivasan, G.; Patel, B.; Kong, K.; et al. Endonuclease EEPD1 Is a Gatekeeper for Repair of Stressed Replication Forks. *J. Biol. Chem.* 2017, 292, 2795–2804. [CrossRef]
- Chun, C.; Wu, Y.; Lee, S.-H.; Williamson, E.A.; Reinert, B.L.; Jaiswal, A.S.; Nickoloff, J.A.; Hromas, R.A. The homologous recombination component EEPD1 is required for genome stability in response to developmental stress of vertebrate embryogenesis. *Cell Cycle* 2016, 15, 957–962. [CrossRef]
- 121. Panopoulou, G.; Hennig, S.; Groth, D.; Krause, A.; Poustka, A.J.; Herwig, R.; Vingron, M.; Lehrach, H. New Evidence for Genome-Wide Duplications at the Origin of Vertebrates Using an Amphioxus Gene Set and Completed Animal Genomes. *Genome Res.* 2003, 13, 1056–1066. [CrossRef]
- 122. Kim, H.-S.; Kim, S.-K.; Hromas, R.; Lee, S.-H. The Set Domain Is Essential for Metnase Functions in Replication Restart and the 5' End of Ss-Overhang Cleavage. *PLoS ONE* **2015**, *10*, e0139418. [CrossRef] [PubMed]
- 123. Kim, H.-S.; Chen, Q.; Kim, S.-K.; Nickoloff, J.A.; Hromas, R.; Georgiadis, M.M.; Lee, S.-K. The Ddn Catalytic Motif Is Required for Metnase Functions in Nhej Repair and Replication Restart. J. Biol. Chem. 2014, 289, 10930–10938. [CrossRef] [PubMed]
- 124. Park, S.-J.; Yoon, B.-H.; Kim, S.-Y. GENT2: An updated gene expression database for normal and tumor tissues. *BMC Med Genom.* **2019**, *12*, 1–8. [CrossRef] [PubMed]
- 125. Puigvert, J.C.; Sanjiv, K.; Helleday, T. Targeting DNA repair, DNA metabolism and replication stress as anti-cancer strategies. *FEBS J.* **2015**, *283*, 232–245. [CrossRef]
- 126. Feng, Z.; Scott, S.P.; Bussen, W.; Sharma, G.G.; Guo, G.; Pandita, T.K.; Powell, S.N. Rad52 Inactivation Is Synthetically Lethal with Brca2 Deficiency. *Proc. Natl. Acad. Sci. USA* 2011, *108*, 686–691. [CrossRef] [PubMed]
- 127. Lok, B.; Carley, A.C.; Tchang, B.; Powell, S.N. RAD52 inactivation is synthetically lethal with deficiencies in BRCA1 and PALB2 in addition to BRCA2 through RAD51-mediated homologous recombination. *Oncogene* **2012**, *32*, 3552–3558. [CrossRef] [PubMed]
- Hromas, R.; Kim, H.-S.; Sidhu, G.; Williamson, E.; Jaiswal, A.; Totterdale, T.A.; Nole, J.; Lee, S.-H.; Nickoloff, J.A.; Kong, K.Y. The endonuclease EEPD1 mediates synthetic lethality in RAD52-depleted BRCA1 mutant breast cancer cells. *Breast Cancer Res.* 2017, 19, 122. [CrossRef] [PubMed]
- 129. Cordaux, R.; Udit, S.; Batzer, M.A.; Feschotte, C. Birth of a Chimeric Primate Gene by Capture of the Transposase Gene from a Mobile Element. *Proc. Natl. Acad. Sci. USA* 2006, *103*, 8101–8106. [CrossRef]
- 130. De Haro, L.P.; Wray, J.; Williamson, E.A.; Durant, S.T.; Corwin, L.; Gentry, A.C.; Osheroff, N.; Lee, S.-H.; Hromas, R.; Nickoloff, J.A. Metnase promotes restart and repair of stalled and collapsed replication forks. *Nucleic Acids Res.* 2010, *38*, 5681–5691. [CrossRef]
- Lee, S.-H.; Oshige, M.; Durant, S.T.; Rasila, K.K.; Williamson, E.A.; Ramsey, H.; Kwan, L.; Nickoloff, J.A.; Hromas, R. The SET domain protein Metnase mediates foreign DNA integration and links integration to nonhomologous end-joining repair. *Proc. Natl. Acad. Sci. USA* 2005, 102, 18075–18080. [CrossRef] [PubMed]
- 132. Nickoloff, J.A.; Sharma, N.; Taylor, L.; Allen, S.J.; Lee, S.-H.; Hromas, R. Metnase and EEPD1: DNA Repair Functions and Potential Targets in Cancer Therapy. *Front. Oncol.* **2022**, *12*. [CrossRef] [PubMed]
- Kim, H.-S.; Williamson, E.A.; Nickoloff, J.A.; Hromas, R.A.; Lee, S.-H. Metnase Mediates Loading of Exonuclease 1 onto Single Strand Overhang DNA for End Resection at Stalled Replication Forks. J. Biol. Chem. 2017, 292, 1414–1425. [CrossRef] [PubMed]
- 134. Fnu, S.; Williamson, E.A.; De Haro, L.P.; Brenneman, M.; Wray, J.; Shaheen, M.; Radhakrishnan, K.; Lee, S.-H.; Nickoloff, J.A.; Hromas, R. Methylation of histone H3 lysine 36 enhances DNA repair by nonhomologous end-joining. *Proc. Natl. Acad. Sci. USA* 2010, 108, 540–545. [CrossRef] [PubMed]
- 135. Williamson, E.A.; Rasila, K.K.; Corwin, L.K.; Wray, J.; Beck, B.D.; Severns, V.; Mobarak, C.; Lee, S.-H.; Nickoloff, J.A.; Hromas, R. The SET and transposase domain protein Metnase enhances chromosome decatenation: Regulation by automethylation. *Nucleic Acids Res.* 2008, *36*, 5822–5831. [CrossRef]
- 136. Hromas, R.; Williamson, E.A.; Fnu, S.; Lee, Y.-J.; Park, S.-J.; Beck, B.D.; You, J.-S.; Laitao, A.; Nickoloff, J.A.; Lee, S.-H. Chk1 Phosphorylation of Metnase Enhances DNA Repair but Inhibits Replication Fork Restart. *Oncogene* **2012**, *31*, 4245–4254. [CrossRef]
- 137. Williamson, E.A.; Wu, Y.; Singh, S.; Byrne, M.; Wray, J.; Lee, S.-H.; A Nickoloff, J.; Hromas, R. The DNA repair component Metnase regulates Chk1 stability. *Cell Div.* **2014**, *9*, 1. [CrossRef]
- 138. Nimonkar, A.V.; Genschel, J.; Kinoshita, E.; Polaczek, P.; Campbell, J.L.; Wyman, C.; Modrich, P.; Kowalczykowski, S.C. Blm-Dna2-Rpa-Mrn and Exo1-Blm-Rpa-Mrn Constitute Two DNA End Resection Machineries for Human DNA Break Repair. *Genes Dev.* 2011, 25, 350–362. [CrossRef]

- 139. Lyu, X.; Lei, K.; Sang, P.B.; Shiva, O.; Chastain, M.; Chi, P.; Chai, W. Human CST complex protects stalled replication forks by directly blocking MRE11 degradation of nascent-strand DNA. *EMBO J.* **2020**, *40*, e103654. [CrossRef]
- Przetocka, S.; Porro, A.; Bolck, H.A.; Walker, C.; Lezaja, A.; Trenner, A.; von Aesch, C.; Himmels, S.F.; D'Andrea, A.D.; Ceccaldi, R.; et al. Ctip-Mediated Fork Protection Synergizes with Brca1 to Suppress Genomic Instability Upon DNA Replication Stress. *Mol. Cell* 2018, 72, 568–582.e6. [CrossRef]
- Locke, A.J.; Hossain, L.; McCrostie, G.; A Ronato, D.; Fitieh, A.; Rafique, T.A.; Mashayekhi, F.; Motamedi, M.; Masson, J.-Y.; Ismail, I.H. SUMOylation mediates CtIP's functions in DNA end resection and replication fork protection. *Nucleic Acids Res.* 2021, 49, 928–953. [CrossRef] [PubMed]
- 142. Young, S.J.; West, S.C. Coordinated roles of SLX4 and MutSβ in DNA repair and the maintenance of genome stability. *Crit. Rev. Biochem. Mol. Biol.* **2021**, *56*, 157–177. [CrossRef] [PubMed]
- 143. Payliss, B.J.; Patel, A.; Sheppard, A.C.; Wyatt, H.D.M. Exploring the Structures and Functions of Macromolecular SLX4-Nuclease Complexes in Genome Stability. *Front. Genet.* **2021**, *12*. [CrossRef] [PubMed]
- 144. Guervilly, J.-H.; Gaillard, P.-H.L. SLX4: Multitasking to maintain genome stability. *Crit. Rev. Biochem. Mol. Biol.* 2018, 53, 475–514. [CrossRef] [PubMed]
- 145. Li, S.; Lu, H.; Wang, Z.; Hu, Q.; Wang, H.; Xiang, R.; Chiba, T.; Wu, X. Ercc1/Xpf Is Important for Repair of DNA Double-Strand Breaks Containing Secondary Structures. *iScience* 2019, *16*, 63–78. [CrossRef] [PubMed]
- 146. Bétous, R.; De Rugy, T.G.; Pelegrini, A.L.; Queille, S.; De Villartay, J.-P.; Hoffmann, J.-S. DNA replication stress triggers rapid DNA replication fork breakage by Artemis and XPF. *PLoS Genet.* **2018**, *14*, e1007541. [CrossRef]
- 147. Trego, K.S.; Groesser, T.; Davalos, A.R.; Parplys, A.C.; Zhao, W.; Nelson, M.; Hlaing, A.; Shih, B.; Rydberg, B.; Pluth, J.M.; et al. Non-catalytic Roles for XPG with BRCA1 and BRCA2 in Homologous Recombination and Genome Stability. *Mol. Cell* 2016, 61, 535–546. [CrossRef]
- 148. Zheng, L.; Zhou, M.; Chai, Q.; Parrish, J.; Xue, D.; Patrick, S.M.; Turchi, J.J.; Yannone, S.M.; Chen, D.; Shen, B. Novel function of the flap endonuclease 1 complex in processing stalled DNA replication forks. *EMBO Rep.* **2005**, *6*, 83–89. [CrossRef]
- 149. Zheng, L.; Jia, J.; Finger, L.D.; Guo, Z.; Zer, C.; Shen, B. Functional regulation of FEN1 nuclease and its link to cancer. *Nucleic Acids Res.* 2010, *39*, 781–794. [CrossRef]
- 150. Nickoloff, J.A.; Taylor, L.; Sharma, N.; Kato, T.A. Exploiting DNA Repair Pathways for Tumor Sensitization, Mitigation of Resistance, and Normal Tissue Protection in Radiotherapy. *Cancer Drug Resist.* **2014**, *4*, 244–263. [CrossRef]
- 151. Carrassa, L.; Damia, G. DNA damage response inhibitors: Mechanisms and potential applications in cancer therapy. *Cancer Treat. Rev.* **2017**, *60*, 139–151. [CrossRef] [PubMed]
- 152. Jackson, S.P.; Helleday, T. DNA REPAIR. Drugging DNA Repair. Science 2016, 352, 1178–1179. [CrossRef] [PubMed]
- 153. Gavande, N.S.; VanderVere-Carozza, P.S.; Pawelczak, K.S.; Vernon, T.L.; Jordan, M.R.; Turchi, J.J. Structure-Guided Optimization of Replication Protein a (Rpa)-DNA Interaction Inhibitors. ACS Med. Chem. Lett. 2020, 11, 1118–1124. [CrossRef] [PubMed]
- 154. Patrone, J.D.; Waterson, A.G.; Fesik, S.W. Recent advancements in the discovery of protein–protein interaction inhibitors of replication protein A. *MedChemComm* **2016**, *8*, 259–267. [CrossRef] [PubMed]
- Neher, T.M.; Bodenmiller, D.; Fitch, R.; Jalal, S.I.; Turchi, J.J. Novel Irreversible Small Molecule Inhibitors of Replication Protein A Display Single-Agent Activity and Synergize with Cisplatin. *Mol. Cancer Ther.* 2011, 10, 1796–1806. [CrossRef]
- 156. Dupre, A.; Boyer-Chatenet, L.; Sattler, R.M.; Modi, A.P.; Lee, J.H.; Nicolette, M.L.; Kopelovich, L.; Jasin, M.; Baer, R.; Paull, T.T.; et al. A Forward Chemical Genetic Screen Reveals an Inhibitor of the Mre11-Rad50-Nbs1 Complex. *Nat. Chem. Biol.* 2008, *4*, 119–125. [CrossRef]
- 157. Chow, T.Y.-K.; A Alaoui-Jamali, M.; Yeh, C.; Yuen, L.; Griller, D. The DNA Double-Stranded Break Repair Protein Endo-Exonuclease as a Therapeutic Target for Cancer. *Mol. Cancer Ther.* **2004**, *3*, 911–919.
- 158. Wang, Y.-Y.; Hung, A.C.; Lo, S.; Hsieh, Y.-C.; Yuan, S.-S.F. MRE11 as a molecular signature and therapeutic target for cancer treatment with radiotherapy. *Cancer Lett.* **2021**, *514*, 1–11. [CrossRef]
- 159. Lin, Z.P.; Ratner, E.S.; Whicker, M.E.; Lee, Y.; Sartorelli, A.C. Triapine Disrupts Ctip-Mediated Homologous Recombination Repair and Sensitizes Ovarian Cancer Cells to Parp and Topoisomerase Inhibitors. *Mol. Cancer Res.* **2014**, *12*, 381–393. [CrossRef]
- 160. Kuster, A.; Mozaffari, N.L.; Wilkinson, O.J.; Wojtaszek, J.L.; Zurfluh, C.; Przetocka, S.; Zyla, D.; von Aesch, C.; Dillingham, M.S.; Williams, R.S.; et al. A stapled peptide mimetic of the CtIP tetramerization motif interferes with double-strand break repair and replication fork protection. *Sci. Adv.* 2021, 7, eabc6381. [CrossRef]
- Kuo, C.-H.; Leu, Y.-L.; Wang, T.-H.; Tseng, W.-C.; Feng, C.-H.; Wang, S.-H.; Chen, C.-C. A novel DNA repair inhibitor, diallyl disulfide (DADS), impairs DNA resection during DNA double-strand break repair by reducing Sae2 and Exo1 levels. *DNA Repair* 2019, *82*, 102690. [CrossRef] [PubMed]
- 162. Liu, W.; Zhou, M.; Li, Z.; Li, H.; Polaczek, P.; Dai, H.; Wu, Q.; Liu, C.; Karanja, K.K.; Popuri, V.; et al. A Selective Small, Olecule Dna2 Inhibitor for Sensitization of Human Cancer Cells to Chemotherapy. *EBioMedicine* **2016**, *6*, 73–86. [CrossRef] [PubMed]
- 163. Kumar, S.; Peng, X.; Daley, J.; Yang, L.; Shen, J.; Nguyen, N.; Bae, G.; Niu, H.; Peng, Y.; Hsieh, H.-J.; et al. Inhibition of DNA2 nuclease as a therapeutic strategy targeting replication stress in cancer cells. *Oncogenesis* 2017, 6, e319. [CrossRef] [PubMed]
- 164. Chen, X.; Ali, I.Y.; EL Fisher, C.; Arribas-Bosacoma, R.; Rajasekaran, M.B.; Williams, G.; Walker, S.; Booth, J.R.; Hudson, J.J.; Roe, S.M.; et al. Uncovering an allosteric mode of action for a selective inhibitor of human Bloom syndrome protein. *eLife* 2021, 10. [CrossRef] [PubMed]

- 165. Nguyen, G.H.; Dexheimer, T.S.; Rosenthal, A.S.; Chu, W.K.; Singh, D.K.; Mosedale, G.; Bachrati, C.Z.; Schultz, L.; Sakurai, M.; Savitsky, P.; et al. A Small Molecule Inhibitor of the BLM Helicase Modulates Chromosome Stability in Human Cells. *Chem. Biol.* 2013, 20, 55–62. [CrossRef]
- 166. Yin, Q.-K.; Wang, C.-X.; Wang, Y.-Q.; Guo, Q.-L.; Zhang, Z.-L.; Ou, T.-M.; Huang, S.-L.; Li, D.; Wang, H.-G.; Tan, J.-H.; et al. Huang. Discovery of Isaindigotone Derivatives as Novel Bloom's Syndrome Protein (BLM) Helicase Inhibitors That Disrupt the BLM/DNA Interactions and Regulate the Homologous Recombination Repair. J. Med. Chem. 2019, 62, 3147–3162. [CrossRef]
- Datta, A.; Brosh, R.M., Jr. New Insights Into DNA Helicases as Druggable Targets for Cancer Therapy. *Front. Mol. Biosci.* 2018, 5, 59. [CrossRef]
- Budke, B.; Lv, W.; Kozikowski, A.P.; Connell, P.P. Recent Developments Using Small Molecules to Target RAD51: How to Best Modulate RAD51 for Anticancer Therapy? *ChemMedChem* 2016, 11, 2468–2473. [CrossRef]
- 169. King, H.O.; Brend, T.; Payne, H.L.; Wright, A.; Ward, T.; Patel, K.; Egnuni, T.; Stead, L.F.; Patel, A.; Wurdak, H.; et al. RAD51 Is a Selective DNA Repair Target to Radiosensitize Glioma Stem Cells. *Stem Cell Rep.* 2017, *8*, 125–139. [CrossRef]
- 170. Balbous, A.; Cortes, U.; Guilloteau, K.; Rivet, P.; Pinel, B.; Duchesne, M.; Godet, J.; Boissonnade, O.; Wager, M.; Bensadoun, R.J.; et al. A radiosensitizing effect of RAD51 inhibition in glioblastoma stem-like cells. *BMC Cancer* **2016**, *16*, 1–13. [CrossRef]
- Pastushok, L.; Fu, Y.; Lin, L.; Luo, Y.; DeCoteau, J.F.; Lee, K.; Geyer, C.R. A Novel Cell-Penetrating Antibody Fragment Inhibits the DNA Repair Protein RAD51. Sci. Rep. 2019, 9, 11227. [CrossRef] [PubMed]
- 172. Turchick, A.; Hegan, D.C.; Jensen, R.B.; Glazer, P.M. A cell-penetrating antibody inhibits human RAD51 via direct binding. *Nucleic Acids Res.* 2017, 45, 11782–11799. [CrossRef] [PubMed]
- 173. Turchick, A.; Liu, Y.; Zhao, W.; Cohen, I.; Glazer, P.M. Synthetic lethality of a cell-penetrating anti-RAD51 antibody in PTENdeficient melanoma and glioma cells. *Oncotarget* 2019, *10*, 1272–1283. [CrossRef] [PubMed]
- 174. Ngo, S.T.; Van Vu, V.; Phung, H.T.T. Computational investigation of possible inhibitors of the winged-helix domain of MUS81. *J. Mol. Graph. Model.* **2020**, *103*, 107771. [CrossRef] [PubMed]
- 175. Williamson, E.A.; Damiani, L.; Leitao, A.; Hu, C.; Hathaway, H.; Oprea, T.; Sklar, L.; Shaheen, M.; Bauman, J.; Wang, W.; et al. Targeting the Transposase Domain of the DNA Repair Component Metnase to Enhance Chemotherapy. *Cancer Res.* **2012**, *72*, 6200–6208. [CrossRef]
- 176. Arora, S.; Heyza, J.; Zhang, H.; Kalman-Maltese, V.; Tillison, K.; Floyd, A.M.; Chalfin, E.M.; Bepler, G.; Patrick, S.M. Identification of small molecule inhibitors of ERCC1-XPF that inhibit DNA repair and potentiate cisplatin efficacy in cancer cells. *Oncotarget* 2016, 7, 75104–75117. [CrossRef]
- McNeil, E.M.; Astell, K.R.; Ritchie, A.-M.; Shave, S.; Houston, D.R.; Bakrania, P.; Jones, H.M.; Khurana, P.; Wallace, C.; Chapman, T.; et al. Inhibition of the ERCC1–XPF structure-specific endonuclease to overcome cancer chemoresistance. *DNA Repair* 2015, *31*, 19–28. [CrossRef]