

Article

Effects of Dietary Black Soldier Fly (*Hermetia illucens*) Oil Supplementation on Flesh Quality of Largemouth Bass (*Micropterus salmoides*)

Zichuan Wang^{1,2}, Yidan Cao^{1,2}, Wei Yang^{1,2}, Zeting Wang^{1,2}, Yang Kuang^{1,2}, Ping Wu^{1,2,*} , Chunfang Cai^{1,2} and Yuantu Ye^{1,2}

¹ School of Life Sciences, Soochow University, Suzhou 215123, China; wangzc1999@163.com (Z.W.); caoyidan2002@163.com (Y.C.); weiyang3216@163.com (W.Y.); wzt200110@163.com (Z.W.); ky17370516398@163.com (Y.K.); caicf@suda.edu.cn (C.C.); yeyt@suda.edu.cn (Y.Y.)

² Key Laboratory of Aquatic Animal Nutrition, Soochow University, Suzhou 215123, China

* Correspondence: wup@suda.edu.cn; Tel.: +86-13912605792

Abstract

Black soldier fly (*Hermetia illucens*) larvae are a promising source of insect lipids, characterized by rapid fatty acid accumulation and a high lauric acid content. This study investigated the effects of dietary black soldier fly oil (BSFO) on muscle quality in largemouth bass (*Micropterus salmoides*). Experimental diets were formulated to be isonitrogenous, isolipidic, and isophosphoric, with 1.0% and 2.0% BSFO partially replacing soybean oil. A control group received 2.3% soybean oil without BSFO or glycerol monolaurate (GML), while positive controls were supplemented with 0.35% and 0.7% GML. Fish (initial weight: 25.08 ± 0.12 g) were cultured in pond cages for 56 days, and three replicates were established for each treatment group. Muscle quality and nutritional traits were evaluated, including proximate composition, fatty acid profiles, texture properties, fiber diameter, hydroxyproline content, antioxidant capacity, and expression of genes related to muscle development, atrophy, apoptosis, and mTOR signaling. Compared with the control, the 2.0% BSFO group showed a significant increase in muscle hydroxyproline content ($p < 0.05$), while GML supplementation led to a significant decrease ($p < 0.05$). In the 1.0% BSFO group, muscle saturated fatty acid (SFA) and monounsaturated fatty acid (MUFA) contents were unchanged ($p > 0.05$), but n-3/n-6 polyunsaturated fatty acid (PUFA) ratios and highly unsaturated fatty acid (HUFA) levels were significantly elevated ($p < 0.05$). The dietary supplementation of BSFO enhanced the levels of high-quality fatty acids in the muscle tissue. Antioxidant capacity was also significantly enhanced in the 1.0% BSFO group ($p < 0.05$) but reduced in the GML groups ($p < 0.05$). Texture analysis showed that BSFO significantly improved muscle hardness, elasticity, chewiness, and gumminess ($p < 0.05$). Gene expression analysis revealed no significant effects of BSFO on genes related to myogenesis (*myod* and *myog*) and muscle atrophy (*mstn* and *murf1*), or apoptosis-related genes (*caspase8*, *caspase9*, and *caspase3*) ($p > 0.05$); mTOR signaling pathway-related genes (*s6k1* and *akt1*) were significantly upregulated in the 2.0% BSFO group ($p < 0.05$). In contrast, 0.7% GML significantly upregulated genes related to myogenesis (*myod*, *myf5*, and *myog*), muscle atrophy (*mstn*, *fbxo32*, and *murf1*), and apoptosis (*caspase8*, *caspase9*, and *caspase3*) ($p < 0.05$). In summary, dietary supplementation with 2.0% BSFO effectively enhances muscle quality in largemouth bass without negatively impacting muscle development.

Keywords: insect lipids; flesh quality; texture property; antioxidant capacity; muscle development



Academic Editor: Héctor G. Nolasco-Soria

Received: 29 September 2025

Revised: 23 October 2025

Accepted: 25 October 2025

Published: 29 October 2025

Citation: Wang, Z.; Cao, Y.; Yang, W.; Wang, Z.; Kuang, Y.; Wu, P.; Cai, C.; Ye, Y. Effects of Dietary Black Soldier Fly (*Hermetia illucens*) Oil Supplementation on Flesh Quality of Largemouth Bass (*Micropterus salmoides*). *Fishes* **2025**, *10*, 548. <https://doi.org/10.3390/fishes10110548>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Key Contribution: Our study aimed to assess the impact of substituting soybean oil with black soldier fly (*Hermetia illucens*) oil on various aspects of largemouth bass (*Micropterus salmoides*), including overall body composition, muscle characteristics, fatty acid profiles, textural properties, muscle fiber diameter, muscle antioxidant capacity, and the expression levels of genes associated with muscle development and apoptosis. Notably, dietary supplementation with BSFO effectively enhances muscle quality in largemouth bass.

1. Introduction

Lipids are indispensable in fish nutrition, serving as a vital source of energy and essential nutrients that are fundamental to the growth and health of aquatic species. Traditionally, soybean oil has been a preferred lipid source in commercial fish feed due to its high quality and nutritional value [1]. However, the extensive use of plant-based fats, such as soybean oil, has raised concerns over its competition with human land resources and the associated environmental implications, including deforestation and the excessive application of fertilizers and pesticides. For the sustainable development of aquaculture, it has become important to find new sources of high-quality fats with less impact on environment.

Insects have garnered extensive attention from researchers as a potential feed source for some aquatic animals in recent years [2]. Insects have advantages such as high fertility, fast growth rates, renewable resources, and the ability to utilize organic waste. The black soldier fly (*Hermetia illucens*) is one of the resource insects designated by the Food and Agriculture Organization (FAO) of the United Nations due to its nutritional comprehensiveness, being well-tolerated, and ease of factory farming [3]. Black soldier fly larvae are rich in medium-chain fatty acids, with lauric acid constituting 21.4–49.3% of total fatty acids [4]. Initially considered a by-product, black soldier fly oil (BSFO) has now attracted attention as a suitable lipid source due to its high lipid content and unique fatty acid profile [5].

Currently, black soldier fly oil has been reported as a fat source for aquatic animal feed. Fawole et al. [6] found that feeding BSFO did not adversely affect the normal performance of juvenile rainbow trout (*Oncorhynchus mykiss*) or fatty acid deposition in muscle. At the same time, adding BSFO to feed could promote growth, effectively prevent the development of enteritis, and improve the immunity of rainbow trout [5]. Sudha et al. [7] found that BSFO significantly enhanced the expression levels of myod and myog in the white muscle of juvenile striped catfish (*Pangasianodon hypophthalmus*), thereby promoting muscle development. Furthermore, adding BSFO to feed affects the fatty acid composition of fish because of the high content of lauric acid in BSFO [8–10]. So far, BSFO has been less studied in largemouth bass (*Micropterus salmoides*). Xia et al. [11] found that replacing fish oil with BSFO in largemouth bass feed had no negative effects on growth and decreased liver glycogen content. Conversely, adding BSFO to feed was beneficial for the immunity promotion of largemouth bass.

Largemouth bass, known for its excellent growth performance, strong resilience, and low disease incidence, has become a key economic freshwater fish species [12]. FAO statistics reveal that global annual production of largemouth bass increased from 621 kilotonnes to 804 kilotonnes between 2020 and 2022 [13]. With the expansion of largemouth bass farming, there is a growing imperative to develop new feed lipid sources. Moreover, as living standards improve, consumer demand for high-quality aquatic products is increasing, making the flesh quality of fish a critical consideration when seeking lipid substitutes. After lipid substitution, the flesh quality of fish should not deteriorate and, ideally, should be enhanced.

In this study, we investigated the effects of partially replacing soybean oil with BSFO in feed on the body composition and muscle texture of largemouth bass. Our findings aim to provide data supporting the substitution of soybean oil with BSFO in feed and contribute to the development of new, low-carbon feed options.

2. Materials and Methods

2.1. Experimental Feed and Groups

Five isonitrogenous and isolipidic feeds with 53% crude protein and 8.6% crude lipids were formulated using fishmeal and soybean meal as the main protein sources and soybean oil, BSFO, and glyceryl monolaurate as the main lipid sources. A BSFO-free diet was used as the control. The experimental diets were formulated to be isonitrogenous, isolipidic, and isophosphoric, with 1.0% and 2.0% BSFO partially replacing soybean oil. The control diet contained 2.3% soybean oil with neither BSFO nor glycerol monolaurate (GML), whereas the positive control diets were supplemented with 0.35% and 0.7% GML. Feed ingredients were thoroughly pulverized to pass through a 60-mesh sieve; under full mixing, the oil was mixed and the right amount of water was added to make pellets. The pellet feed was dried at 40 °C and stored at −20 °C until use. The experimental diet formulation and proximate composition are shown in Table 1.

Table 1. Largemouth bass experimental diet composition (g/kg) and nutrient composition.

Items	Control	BSFO1	BSFO2	GML0.35	GML0.7
Tapioca	100	100	100	100	100
Full-fat rice bran	116.5	116.5	116.5	116.5	116.5
Cotton meal	80	80	80	80	80
Corn gluten meal	35	35	35	35	35
Fish meal	500	500	500	500	500
Hemoglobin powder	30	30	30	30	30
Chicken meal	90	90	90	90	90
Black soldier fly oil	0	10	20	0	0
Soybean oil	23	13	3	19.5	16
Glyceryl monolaurate	0	0	0	3.5	7
Ca(H ₂ PO ₄) ₂	15	15	15	15	15
Premix ¹	10	10	10	10	10
Ruiantai ²	0.5	0.5	0.5	0.5	0.5
Proximate analysis (% DM)					
Moisture	10.67	6.56	5.51	11.23	10.09
Crude protein	53.30	53.40	53.12	53.37	53.22
Crude lipids	8.66	8.64	8.77	8.66	8.70
Ash	13.51	13.33	13.43	13.39	13.40

¹ Premix contained the following per kilogram of feed: vitamin A, 8 mg; vitamin B1, 18 mg; vitamin B2, 8 mg; vitamin B6, 12 mg; vitamin B12, 0.02 mg; vitamin C, 300 mg; vitamin D3, 3 mg; vitamin K3, 5 mg; folic acid, 5 mg; pantothenic acid, 25 mg; niacin, 25 mg; inositol, 100 mg; Mg, 96 mg; Fe, 64 mg; Zn, 19 mg; Mn, 13 mg; Cu, 2.5 mg; I, 0.021 mg; Se, 0.07 mg; Co, 0.016 mg; K, 0.05 mg. ² Ruiantai is a natural plant-based additive.

The black soldier fly larvae were rinsed three times with tap water and ultrapure water, and then dried using vacuum freeze-drying. The dried BSFL were subjected to mechanical pressing to extract oil. The extracted oil was centrifuged at 13,000 rpm for 10 min at 40 °C, and the clear supernatant was collected as the BSFO used in this study.

2.2. Experimental Fish and the Feeding Trial

The experimental fish were fed in net cages (1.8 m × 1.5 m × 1.8 m) in a pond covering an area of 10,000 m². The experimental fish were fed with the diet of the control group for two weeks for acclimation. A total of 600 healthy largemouth bass were randomly divided into 15 net cages (1.8 m × 1.5 m × 1.8 m), with 40 fish per cage (stocking density: 8.23 fish/m³), which had an initial average body weight of 25.08 ± 0.12 g. During the eight-week culture trial, the fish were fed twice daily (06:30 and 17:00) with 3% of their body

weight. During farming, the water temperature was maintained at 24–35 °C, dissolved oxygen > 6 mg/L, pH = 7.6 ± 0.5 , and $\text{NH}_4^+-\text{N} < 0.01$ mg/L.

2.3. Sample Collection

After the 8-week feeding trial, the fish were fasted for 24 h and anesthetized with an appropriate amount of MS-222 (0.1 mL/L). Three fish were taken from each net cage and stored at -20 °C for body component analysis. Nine fish (three fish for the determination of muscle antioxidant capacity, three fish for the measurement of glycogen and hydroxyproline content in muscle, and three fish for gene expression analysis) in each net cage were randomly selected for dissection, and muscles above the left lateral line were collected, frozen in liquid nitrogen, and stored at -80 °C for biochemical analysis and gene expression analysis. Another 3 fish were dissected, and the muscles above the left lateral line were stored in a 4% paraformaldehyde solution at 4 °C for histologic section preparation. Three fish were taken from each net cage to remove scales and skin, and the muscle below the dorsal fin and above the left lateral line was cut into $1\text{ cm} \times 1\text{ cm} \times 1\text{ cm}$ square muscle blocks for texture analysis.

2.4. Proximate Analysis of Body and Muscle Composition

The moisture, crude protein, crude fat, and crude ash contents were calculated according to the National Standards GB5009.3-2016 [14], GB6432-2018 [15], GB/T6433-2006 [16], and GB5009.4-2016 [17], respectively. The fatty acid composition was determined with a fatty acid profile analyzer (6980N, Agilent Technologies, Santa Clara, CA, USA), and the glycogen and hydroxyproline contents of the muscle were determined using the A043-1-1 and A030-2-1 commercial assay kits (Nanjing Jiancheng Bioengineering Institute, Nanjing, China).

2.5. Muscle Antioxidant Capacity

Muscle samples were homogenized using a high-throughput tissue grinder (JXFSTPRP-24; Shanghai Jingxin, Shanghai, China). Total superoxide dismutase (T-SOD) and total antioxidant capacity (T-AOC) in muscle were measured by the colorimetric method using the A001-1-2 and A015-1-2 commercial assay kits (Nanjing Jiancheng Bioengineering Institute, Nanjing, China).

2.6. Texture Properties of Muscle

Six textural parameters, including hardness, elasticity, chewiness, cohesiveness, gumminess, and resilience, were measured using a texture analyzer (Rapid TA+; Shanghai Tengba Instrument Technology, Shanghai, China). The analyzer was operated with the following parameters: P/36R probe diameter of 3.6 mm, test speeds of 5 mm/s, downward pressure distance of 30% of the sample thickness at 4 s intervals, and a trigger force setting of 5 g.

2.7. Histology of Muscle

Muscle samples were dehydrated in ethanol, infiltrated in xylene, embedded in paraffin, cut serially (5 μm thick), and then stained with hematoxylin and eosin. Sections were observed using a light microscope (DM500; Leica, Wetzlar, German) and photographed. The diameters of the muscle fibers were expressed as the longest axis length using imaging software (ImageJ 2025, USA).

2.8. RT-qPCR Analysis

Total RNA was extracted from muscle using Trizol reagent. RNA concentration was determined using a micro-UV spectrophotometer (NanoDrop 2000; Thermo, Waltham, MA,

USA). The total RNA was reverse-transcribed into cDNA with the PrimerScript™ RT reagent kit with the gDNA Eraser (TaKaRa, Dalian, China) and then stored at -20°C . The relative expression levels of *gapdh* were used as an internal reference, and the relative expression levels of the target genes were calculated using the $2^{-\Delta\Delta\text{CT}}$ method [18]. Primers designed by Primer Premier 5.0 and the sequences are shown in Table 2.

Table 2. The sequences of gene primers used for RT-qPCR.

Gene	Forward Primer (5'-3')	Reverse Primer (5'-3')
<i>myf5</i>	GAGGAGGACGAGCATGTCAG	GAGGTGCAACGTCTCAAAGC
<i>myod</i>	GCGACTAAGCAAGGTGAACG	CTGCAGGGACTCGATGTAGC
<i>myog</i>	GAGTTGGGGTGACAGGAACA	TCTGGTTTGGGTTCATCAGG
<i>mstn</i>	TGATTGCTTTGGGTCCAGTC	CCTTCATTCCGAGTTTGCTC
<i>fbxo32</i>	GACATGGCTGCCAAGAAGAG	TGAAGGCCTCTCCCAGTGTA
<i>MuRF1</i>	TGGACGACACGTGTAAGCTG	GTGGACACCTTCTGCTGCTC
<i>caspase7</i>	CCATGGTGAAGAGGGAATGA	CGGACCCGAATCTGTCTGTA
<i>caspase8</i>	GAGGGGACAAAGAGGTGGAG	GAGCCTGTGGAAGTGTTTCG
<i>caspase9</i>	TCGGGCCTTTCCATTATTTC	AACAACCTGAGTGGGTGCTG
<i>akt1</i>	AGGACGCTACTACGCCATGA	CCGCTCCTCTGAGAACACAC
<i>s6k1</i>	GGCTCATCCCTTCTTTCGAC	GGCTTGATTCCGACTCTCAC
<i>4ebp1</i>	GACTGCCAGAAGACCACTGC	CAGCAGGAACCTTCGGTCAT
<i>gapdh</i>	AAGGGTGGTGCCAAGAGAGT	AGTCTTCTGAGTGGCGGTGA

2.9. Statistical Analysis

SPSS 26 was used for statistical analysis. One-way ANOVA was carried out with Duncan's multiple-range test at a significance level of 0.05. All the results are presented as means $\pm$ standard errors (SEMs). Bar graphs were produced using Origin 2021 software.

3. Results

3.1. The Effect of Black Soldier Fly Oil Supplementation on Body and Muscle Composition

Largemouth bass fed with different levels of BSFO exhibited no significant impact on the crude lipid, crude protein, or ash contents of their whole bodies or muscles ($p > 0.05$) (Table 3).

Table 3. Effects of black soldier fly oil supplementation on body and muscle composition of largemouth bass (dry weight, %).

Items	Control	BSFO1	BSFO2
Whole fish body			
Crude lipids (%)	20.71 $\pm$ 0.98	23.06 $\pm$ 1.67	24.59 $\pm$ 1.84
Crude protein (%)	61.12 $\pm$ 1.12	59.96 $\pm$ 2.52	58.68 $\pm$ 2.24
Ash (%)	15.29 $\pm$ 0.72	13.95 $\pm$ 0.46	13.95 $\pm$ 0.12
Muscle			
Crude lipids (%)	16.51 $\pm$ 0.83	17.04 $\pm$ 1.06	19.80 $\pm$ 1.75
Crude protein (%)	80.09 $\pm$ 0.91	79.49 $\pm$ 2.72	79.09 $\pm$ 1.94
Ash (%)	5.87 $\pm$ 0.27	5.90 $\pm$ 0.12	6.27 $\pm$ 0.12

Control, BSFO1, and BSFO2 represent diets with BSFO inclusion levels of 0%, 1%, and 2%, respectively. All data are expressed as means $\pm$ SEMs ($n = 3$).

3.2. The Effect of Black Soldier Fly Oil and Glyceryl Monolaurate Supplementation on the Glycogen and Hydroxyproline Contents of Muscle

There were no significant effects on glycogen contents in the BSFO and GML groups compared with the control group ($p > 0.05$) (Table 4). Compared with the control group and BSFO1, hydroxyproline content was significantly increased in the BSFO2 group, whereas it was decreased in both the GML0.35 and GML0.7 groups ($p < 0.05$) (Table 4).

Table 4. Effect of black soldier fly oil and glyceryl monolaurate supplementation on the glycogen and hydroxyproline content of muscle in largemouth bass.

Items	Control	BSFO1	BSFO2	GML0.35	GML0.7
Glycogen (mg/g)	1.06 ± 0.08	1.11 ± 0.08	1.1 ± 0.16	1.09 ± 0.02	1.06 ± 0.04
Hydroxyproline (µg/mL)	0.56 ± 0.01 ^b	0.6 ± 0.02 ^b	0.76 ± 0.05 ^a	0.41 ± 0.02 ^c	0.41 ± 0.02 ^c

Control, BSFO1, and BSFO2 represent diets with BSFO inclusion levels of 0%, 1%, and 2%, respectively. GML0.35 and GML0.7 represent diets with GML inclusion levels of 0.35% and 0.7%, respectively. All data are expressed as means ± SEMs (n = 3). Means in the same row with different superscripts are significantly different ($p < 0.05$, one-way ANOVA followed by Duncan's multiple-range test).

3.3. The Effect of Black Soldier Fly Oil Supplementation on the Fatty Acid Composition of Body and Muscle

Table 5 shows the effects on the fatty acid composition of the bodies and muscles of largemouth bass exerted by black soldier fly oil supplementation. C12:0 (lauric acid), C14:0, and Σ SFA contents in the body were significantly higher in the BSFO supplementation groups compared with the control group ($p < 0.05$), and C20:5n-3 (EPA), C22:6n-3 (DHA), Σ n-3 PUFA, and Σ HUFA contents were significantly lower in the BSFO groups compared with the control group ($p < 0.05$). The n-3/n-6 PUFA ratios were not significantly different across all groups ($p > 0.05$).

Table 5. Effect of black soldier fly oil supplementation on the fatty acid composition of whole fish and muscle in largemouth bass (% of total fatty acids).

Items	Body			Muscle		
Fatty acid	Control	BSFO1	BSFO2	Control	BSFO1	BSFO2
C12:0	0.18 ± 0.06 ^c	4.47 ± 0.1 ^b	6.23 ± 0.46 ^a	0.14 ± 0 ^c	2.93 ± 0.22 ^b	4.84 ± 0.33 ^a
C13:0	0.09 ± 0 ^a	0.09 ± 0 ^a	0.06 ± 0 ^b	0.04 ± 0 ^a	0.03 ± 0 ^b	0.02 ± 0 ^b
C14:0	5.43 ± 0.11 ^b	6.59 ± 0.14 ^a	6.36 ± 0.19 ^a	3.24 ± 0.01 ^c	4.07 ± 0.08 ^b	4.42 ± 0.18 ^a
C15:0	0.59 ± 0.02 ^a	0.53 ± 0.03 ^b	0.42 ± 0.03 ^c	0.34 ± 0 ^a	0.3 ± 0.01 ^b	0.29 ± 0 ^b
C16:0	35.51 ± 0.75 ^a	34 ± 0.5 ^{ab}	32.64 ± 0.86 ^b	25.71 ± 0.04 ^a	23.72 ± 0.63 ^b	23.62 ± 0.05 ^b
C17:0	0.59 ± 0.02	0.53 ± 0.05	0.49 ± 0.05	0.35 ± 0 ^a	0.29 ± 0.01 ^c	0.32 ± 0 ^b
C18:0	7.03 ± 0.02 ^a	6.38 ± 0.27 ^b	6.04 ± 0.28 ^b	5.36 ± 0.02 ^a	4.12 ± 0.04 ^b	4.23 ± 0.18 ^b
C21:0	0.14 ± 0.03 ^a	0.1 ± 0.02 ^{ab}	0.07 ± 0.01 ^b	0.48 ± 0.05 ⁵	0.41 ± 0.04	0.43 ± 0.01
C23:0	0.43 ± 0.03 ^a	0.3 ± 0.04 ^b	0.24 ± 0.04 ^b	0.4 ± 0.06 ^a	0.1 ± 0.01 ^b	0.11 ± 0.01 ^b
Σ SFA	50.01 ± 0.73 ^b	53.05 ± 0.22 ^a	52.61 ± 0.77 ^a	36.43 ± 0.03 ^b	36.37 ± 0.43 ^b	38.64 ± 0.34 ^a
C16:1n-7	6.9 ± 0.12	6.96 ± 0.31	6.67 ± 0.26	5.37 ± 0 ^c	5.85 ± 0.24 ^a	5.45 ± 0.17 ^{ab}
C18:1n-9	33.97 ± 0.17 ^{ab}	32.63 ± 1 ^b	34.51 ± 0.72 ^a	29.03 ± 0.1 ^a	26.44 ± 0.92 ^b	27.09 ± 0.2 ^b
C20:1n-9	0.39 ± 0.02 ^a	0.31 ± 0.01 ^b	0.25 ± 0.01 ^c	0.28 ± 0.04 ^a	0.19 ± 0.01 ^b	0.18 ± 0 ^b
C24:1n-9	0.18 ± 0.03	0.2 ± 0.1	0.17 ± 0.03	5.45 ± 0.03 ^b	7.67 ± 0.25 ^a	7.88 ± 0.29 ^a
Σ MUFA	41.5 ± 0.1	40.19 ± 1.03	41.7 ± 0.46	40.3 ± 0.14	40.34 ± 0.79	40.8 ± 0.31
C18:2n-6	6.18 ± 0.72 ^a	4.95 ± 0.82 ^{ab}	4.08 ± 0.39 ^b	18.38 ± 0.02 ^a	16.91 ± 0.72 ^b	14.93 ± 0.02 ^c
C18:3n-6	0.15 ± 0.02	0.11 ± 0.04	0.08 ± 0.02	1.33 ± 0.03 ^a	1.45 ± 0.1 ^a	1.16 ± 0.01 ^b
C20:2n-6	0.06 ± 0.01 ^a	0.04 ± 0 ^{ab}	0.03 ± 0 ^b	0.04 ± 0.01 ^a	0.04 ± 0 ^{ab}	0.03 ± 0 ^b
C20:3n-6	0.04 ± 0	0.03 ± 0.01	0.02 ± 0	0.2 ± 0.01 ^b	0.35 ± 0.02 ^a	0.44 ± 0.07 ^a
C20:4n-6	0.21 ± 0.01 ^a	0.14 ± 0.02 ^b	0.09 ± 0.02 ^c	0.14 ± 0.02	0.14 ± 0.01	0.14 ± 0.01
Σ n-6 PUFA	6.64 ± 0.75	5.28 ± 0.88	4.3 ± 0.4	20.1 ± 0.01 ^a	18.89 ± 0.83 ^b	16.7 ± 0.04 ^c
C18:3n-3	1.23 ± 0.06 ^a	1.13 ± 0.03 ^{ab}	1.06 ± 0.09 ^b	0.97 ± 0.04 ^a	0.78 ± 0.04 ^b	0.91 ± 0.03 ^a
C20:3n-3	0.2 ± 0.05 ^a	0.13 ± 0.02 ^b	0.1 ± 0.01 ^b	1.07 ± 0.01 ^b	1.42 ± 0.18 ^a	1.11 ± 0.08 ^b
C20:5n-3 (EPA)	0.16 ± 0.03 ^a	0.07 ± 0.03 ^b	0.1 ± 0.02 ^{ab}	1.02 ± 0.12 ^c	2.08 ± 0.09 ^a	1.72 ± 0.03 ^b
C22:6n-3 (DHA)	0.26 ± 0.06 ^a	0.14 ± 0 ^b	0.12 ± 0.02 ^b	0.11 ± 0.01	0.12 ± 0.02	0.12 ± 0.01
Σ n-3 PUFA	1.85 ± 0.09 ^a	1.48 ± 0.01 ^b	1.38 ± 0.12 ^b	3.17 ± 0.09 ^c	4.4 ± 0.28 ^a	3.86 ± 0.02 ^b
Σ PUFA	8.48 ± 0.81 ^a	6.76 ± 0.88 ^{ab}	5.69 ± 0.34 ^b	23.27 ± 0.11 ^a	23.29 ± 1.11 ^a	20.56 ± 0.04 ^b
Σ HUFA	0.82 ± 0.1 ^a	0.51 ± 0.02 ^b	0.43 ± 0.01 ^b	2.54 ± 0.05 ^c	4.11 ± 0.27 ^a	3.54 ± 0.1 ^b
n-3/n-6 PUFA	0.28 ± 0.02	0.29 ± 0.05	0.33 ± 0.05	0.16 ± 0 ^b	0.23 ± 0 ^a	0.23 ± 0 ^a

Control, BSFO1, and BSFO2 represent diets with BSFO inclusion levels of 0%, 1%, and 2%, respectively. All data are expressed as means ± SEMs (n = 3). Means in the same row with different superscripts are significantly different ($p < 0.05$, one-way ANOVA followed by Duncan's multiple-range test).

In the muscle, EPA, HUFA, and Σ n-3 PUFA contents were significantly higher in the BSFO groups compared with the control group ($p < 0.05$), and the contents of DHA and Σ n-6 PUFAs were significantly lower in the control group ($p < 0.05$). The n-3/n-6 PUFA ratios were significantly higher in the BSFO groups than in the control group ($p < 0.05$).

3.4. The Effect of Black Soldier Fly Oil and Glyceryl Monolaurate Supplementation on the Antioxidant Capacity of Muscle

There was no significant difference in T-SOD activity among the groups ($p > 0.05$), as shown in Table 6. The T-AOC in the muscle was not affected by BSFO supplementation ($p > 0.05$) and was significantly lower in the GML0.35 and GML0.7 groups compared with the control group ($p < 0.05$).

Table 6. Effects of black soldier fly oil and glyceryl monolaurate supplementation on the antioxidant capacity of muscle in largemouth bass.

Items	Control	BSFO1	BSFO2	GML0.35	GML0.7
T-SOD (U/mgprot)	33.97 $\pm$ 0.89	33.67 $\pm$ 3.05	31.43 $\pm$ 1.94	33.31 $\pm$ 1.85	31.34 $\pm$ 1.12
T-AOC (U/mgprot)	2.63 $\pm$ 0.12 ^a	2.7 $\pm$ 0.27 ^a	2.95 $\pm$ 0.24 ^a	1.76 $\pm$ 0.18 ^b	1.94 $\pm$ 0.11 ^b

Control, BSFO1, and BSFO2 represent diets with BSFO inclusion levels of 0%, 1%, and 2%, respectively. GML0.35 and GML0.7 represent diets with GML inclusion levels of 0.35% and 0.7%, respectively. All data are expressed as means $\pm$ SEMs ($n = 3$). Means in the same row with different superscripts are significantly different ($p < 0.05$, one-way ANOVA followed by Duncan's multiple-range test).

3.5. The Effect of Black Soldier Fly Oil Supplementation on the Texture of Muscle

As shown in Table 7, the hardness, elasticity, chewiness, and gumminess of muscle significantly increased compared with the control ($p < 0.05$), and cohesiveness and resilience were not affected by BSFO supplementation ($p > 0.05$).

Table 7. Texture of largemouth bass muscle as a result of partially substituting soybean oil with black soldier fly oil.

Items	Control	BSFO1	BSFO2
Hardness, g	218.88 $\pm$ 24.03 ^b	448.18 $\pm$ 46.62 ^a	351.87 $\pm$ 33.88 ^a
Elasticity, mm	0.58 $\pm$ 0.01 ^b	0.64 $\pm$ 0.01 ^a	0.62 $\pm$ 0.01 ^a
Chewiness, mJ	79.55 $\pm$ 11.86 ^b	178.73 $\pm$ 18.88 ^a	142.16 $\pm$ 17.82 ^a
Gumminess, g	135.83 $\pm$ 18.11 ^b	276.26 $\pm$ 27.27 ^a	226.89 $\pm$ 24.68 ^a
Cohesiveness	0.62 $\pm$ 0.02	0.67 $\pm$ 0.01	0.64 $\pm$ 0.03
Resilience	0.47 $\pm$ 0.02	0.5 $\pm$ 0.01	0.48 $\pm$ 0.02

Control, BSFO1, and BSFO2 represent diets with BSFO inclusion levels of 0%, 1%, and 2%, respectively. All data are expressed as means $\pm$ SEMs ($n = 3$). Means in the same row with different superscripts are significantly different ($p < 0.05$, one-way ANOVA followed by Duncan's multiple-range test).

3.6. The Effect of Black Soldier Fly Oil Supplementation on the Histology of Muscle

As shown in Figure 1, compared with the control, the mean diameter of myofibers and the distribution of them in the muscle were not affected by BSFO supplementation ($p > 0.05$). The histological morphological structure of largemouth bass myofibers was similar in both the soybean oil feed group and the BSFO groups.

3.7. The Effect of Fat Source Supplementation on the Relative Gene Expression Levels in the Muscle

The relative expression levels of myogenesis genes, TOR pathway-related genes, and muscle apoptosis-related genes of largemouth bass are shown in Figure 2. Compared with the control group, the relative expression levels of *myf5* related to *myog* were significantly lower in the BSFO1 group ($p < 0.05$), and the relative expression levels of *fbxo32* related to muscle atrophy were significantly higher in the BSFO2 group ($p < 0.05$). Adding BSFO to the diet reduced the relative expression levels of *4ebp1* significantly ($p < 0.05$). The relative

expression levels of *akt1* and *s6k1* were significantly higher in the BSFO2 group ($p < 0.05$). The results showed that the addition of BSFO in the diet had no significant effect on the relative expression levels of apoptosis-related genes ($p > 0.05$).




Figure 1. The effect of black soldier fly oil replacing soybean oil on the histology of muscle in largemouth bass. Control, BSFO1, and BSFO2 represent diets with BSFO inclusion levels of 0%, 1%, and 2%, respectively. (A–C) Transversal section of the white muscle of fish fed different experimental diets. Hematoxylin–eosin staining. Photomicrographs: magnification 100×; scale bar: 100 μm. (D) Average diameter of myofibers. (E) the distribution of myofiber diameters.

Compared with the control group, the relative expression levels of all genes related to myogenesis, the TOR pathway, and muscle apoptosis were significantly higher in the GML0.7 group ($p < 0.05$), and the relative expression levels of *fbxo32*, *s6k1*, *caspase8*, and *caspase9* were significantly lower in the GML0.35 group ($p < 0.05$).




Figure 2. Effects of black soldier oil replacing soybean oil on the regulation of myogenesis, TOR pathway-related genes, and apoptosis-related genes of largemouth bass. Control, BSFO1, and BSFO2 represent diets with BSFO inclusion levels of 0%, 1%, and 2%, respectively. GML0.35 and GML0.7 represent diets with GML inclusion levels of 0.35% and 0.7%, respectively. **(A)** Myogenesis regulation-related genes. **(B)** TOR pathway-related genes. **(C)** Apoptosis-related genes.

4. Discussion

4.1. Effect of BSFO Replacing Soybean Oil on Flesh Quality of Largemouth Bass

Flesh textural characteristics, pivotal in quality assessment, are typically evaluated using the texture profile analysis method [19]. This method quantifies key textural parameters such as hardness, elasticity, chewiness, cohesiveness, gumminess, and resilience, mimicking the human chewing process. These parameters are influenced by several factors, including intramuscular and intermuscular fat content, collagen content in connective tissues, and myofiber diameter and density [20]. Notably, hardness is a critical parameter for consumers, often reflecting the perceived value of meat [21]. Previous studies on fish have shown that the addition of black soldier fly oil alters the texture of flesh in terms of changes in hardness, elasticity, chewiness, and gumminess [22–24]. In this study, we found that hardness, elasticity, chewiness, and gumminess reached higher values in the BSFO1 group, while they decreased in the BSFO2 group, whose diet was characterized by a higher degree of substitution compared with the BSFO1 group, though the values were still higher than in the control group. This result is similar to a previous discovery in largemouth bass, where the addition of BSFO to the diet significantly increased flesh hardness [22]. Among the texture characteristics of flesh, hardness is especially closely related to muscle lipid content [25]. In the present study, the muscle lipid content of largemouth bass did not change significantly when the fish were fed with BSFO instead of soybean oil. Previous studies in *O. mykiss*, *Cyprinus carpio* var. *Jian*, and largemouth bass also found that the usage of BSFO in diets could not vary the lipid content of muscle [6,8,11].

In addition, the characteristics of fish myofibers influence the quality of the flesh, particularly the hardness, which is directly related to myofiber diameter and density [26]. Previous studies have shown that the hardness and chewiness of flesh decrease as myofiber diameter increases and myofiber density decreases [27,28]. However, in this study, the increase in muscle hardness was not attributed to alterations in myofiber diameter, as substituting soybean oil with BSFO did not yield significant changes in either the diameter or density of largemouth bass myofibers. Our results contrast with the findings of Xia et al. [11], who observed a significant increase in the diameter of *Micropterus salmoides* myofibers when BSFO replaced more than 50% of fish oil, whereas replacement levels equal to or below 50% elicited no change. It is possible that the difference in results is due to the difference between the fish oil and soybean oil used in the control group. Stratos et al. [29] demonstrated that caspase-mediated apoptosis was negatively correlated with myofiber diameter. In this study, fish fed BSFO showed no significant impact on the relative expression levels of the apoptosis initiators *caspase8* and *caspase9* and the apoptosis effector *caspase3*. However, the relative expression levels of *caspase8*, *caspase9*, and *caspase3* were significantly upregulated in the GML0.35 and GML0.7 groups. This suggested that lauric acid can induce apoptosis. Interestingly, despite the high lauric acid content in BSFO, its inclusion in fish diets did not exhibit these adverse effects, highlighting the potential of BSFO as a valuable feed component, even with its high concentration of lauric acid.

Furthermore, collagen content also affects the hardness of fish muscles. As a key component of the extracellular matrix, collagen is crucial for sustaining muscle growth and texture in fish [30]. Hydroxyproline, constituting 13.4% of the total amino and imine content in collagen and present in trace amounts in elastin, is absent in other proteins, making it a reliable indicator of collagen levels in muscle tissues [31]. In this study, we found that hydroxyproline content increased in the muscle of largemouth bass fed BSFO instead of soybean oil, which indicated that the increase in collagen amount is a reason for the increase in flesh hardness.

In order to explore the mechanisms behind the observed changes in the muscle texture of largemouth bass, we examined the expression levels of genes associated with muscle

growth. Østbye et al. [32] found a positive correlation between muscle hardness and myogenesis in Atlantic salmon (*Salmo salar*). The process of fish muscle growth is regulated by a variety of factors, including myogenic regulatory factors (MRFs) and atrophy factors. MRFs are a family of basic helix–loop–helix transcription factors, including *myod*, *myf5*, *myf4*, and *myog*, and can convert a large number of different cell types into muscle [33]. When the inhibition of myogenesis and protein degradation by atrophy factors exceed protein synthesis, muscle protein loss occurs, leading to a corresponding decrease in muscle hardness [34]. The genes *mstn*, *fbxo32*, and *MuRF1* are three atrophy signs that reliably indicate muscle protein loss [35–37]. After an 8-week feeding trial with dietary BSFO, we assessed its impact on the relative expression levels of *myod*, *myf5*, *myog*, *mstn*, *fbxo32*, and *MuRF1* in the muscle. Previous studies have shown that some trophic factors affected the expression levels of one or more genes in MRFs, such as dietary methionine levels for rainbow trout juveniles [38], dietary lysine and histidine levels for *Oreochromis niloticus* [39], and dietary plant protein levels for *Solea senegalensis* [40]. Sudha et al. [7] found that replacing fish oil with BSFO increased the expression levels of genes related to muscle myogenesis in *Pangasianodon hypophthalmus* while reducing myostatin expression levels. However, another study on replacing fish oil in largemouth bass diets with BSFO found that there were no significant differences in mRNA expression levels of *myod1*, *myf1*, *myos*, *myog*, *myf5*, and *paxbp-1* [11]. In this study, we replaced soybean oil in largemouth bass diets with BSFO and found that the relative expression levels of genes related to muscle myogenesis and atrophy were not affected in the BSFO2 group compared with the control group. In the BSFO1 group, the expression level of *myf5* decreased significantly, while the expression level of *fbxo32* increased significantly, revealing the primary reason why muscle hardness in the BSFO1 group was higher than that in the control group and the BSFO2 group.

In this study, the relative expression levels of genes related to muscle myogenesis and atrophy were upregulated in the GML0.7 group compared with the control. Despite the substantial lauric acid content in BSFO, the relative expression levels of these genes in the BSFO group were significantly lower than those in the GML group. This suggests that certain components within BSFO may mitigate the adverse effects typically associated with lauric acid, highlighting the potential for BSFO to modulate muscle gene expression in a beneficial manner.

The target of rapamycin (TOR) pathway integrates signals from extracellular and intracellular agents, which can regulate protein synthesis and promote cell survival, proliferation, and growth [41]. In the TOR pathway, kinase akt activates TOR target proteins, phosphorylates kinase S6K1 and eukaryotic translation initiation factor 4E-BP1, and regulates protein synthesis to promote muscle growth [42]. As demonstrated by Goodman et al., sustained activation of Akt1 and its downstream effector S6K1 is sufficient to stimulate muscle protein synthesis and, consequently, promote skeletal muscle growth [43]. In the present study, the relative expression levels of *akt1* and *s6k1* were significantly upregulated, suggesting that BSFO promotes cell growth and may lead to increased hardness in muscle.

4.2. Effect of Dietary BSFO on Fatty Acid Profile of Largemouth Bass

The fatty acid profile in fish is directly affected by the fatty acid composition of the feed [44]. BSFO is predominantly composed of SFAs, the majority of which are lauric acid [45]. The results showed that as the level of BSFO supplementation in the feed increased, the content of lauric acid increased in the whole body and muscle of largemouth bass. Similar results have been reported in gilthead seabream [9]. In addition, feeding BSFO instead of soybean oil can result in an increase in C22:6n-3 (DHA) in muscle in Jian carp [8]. However, the fatty acid profile determined in rainbow trout was not consistent with these

results, the usage of BSFO as a replacement for soybean oil causing no significant change in the DHA content of the muscle [6]. In the present study, replacing soybean oil with BSFO did not cause the DHA content to change in largemouth bass muscle, but significantly elevated C20:5n-3 (EPA) content in muscle, which may be attributed to the fact that lauric acid can be preferentially oxidized, allowing EPA to be better retained in muscle. The result that whole body contains more lauric acid than the muscle seems to support this idea.

In addition, the n-3/n-6 ratio is an important parameter for assessing the nutrition of flesh. Studies have shown that a high n-3/n-6 ratio is favorable for human health because n-3 PUFAs can reduce inflammatory responses, whereas n-6 PUFAs cause inflammation [46–48]. In the present study, n-3/n-6 levels were also significantly higher in the BSFO1 and BSFO2 groups than in the control group. This is in agreement with the results of previous studies [6,10]. Thus, the addition of BSFO to the diet increased EPA levels and n-3/n-6 ratios in largemouth bass muscle, thereby improved fatty acid quality.

4.3. Effect of Dietary BSFO on Antioxidant Capacity of Largemouth Bass

T-AOC is the total level of various antioxidant macromolecules, antioxidant small molecules, and enzymes within a system [49,50]. T-SOD is an antioxidant enzyme that protects cells from peroxidative damage [51]. In the present study, replacing soybean oil with BSFO had no significant effect on T-AOC or T-SOD activities in muscle. The results showed that inclusion of BSFO up to 2% in the diet had no significant effect on the antioxidant capacity of largemouth bass. Similar results have been reported, for instance, in juvenile yellow catfish; BSFO as a substitute for soybean oil has no significant effect on antioxidant capacity [52].

However, another study found that the addition of BSFO to largemouth bass diets resulted in an increase in T-SOD activities in the liver, suggesting that BSFO improves the antioxidant capacity of largemouth bass [11]. Because different studies have produced conflicting results [11,52], to investigate how BSFO affects the antioxidant capacity of largemouth bass, we fed largemouth bass using feed in which soybean oil was partially substituted with GML. The results showed that the total antioxidant capacity of largemouth bass was significantly reduced. However, this may not be a universal phenomenon. For example, a study in hybrid grouper fish indicated that GML increased the activity of SOD [53]. In *Trachinotus ovatus*, 0.15% GML significantly increased total antioxidant capacity and superoxide dismutase activities [54]. Similarly, the dietary addition of GML significantly increased SOD activity in juvenile grouper fish [55]. The different results may be due to the different effects of GML on different species of fish. In the present study, total antioxidant capacity was significantly lower in the GML groups than in the control, whereas in the BSFO groups it remained comparable to that in the control, suggesting that certain components in BSFO might mitigate the GML-induced impairment of antioxidant capacity in largemouth bass.

5. Conclusions

Our results showed that the addition of BSFO to a largemouth bass diet increased muscle hardness, enhanced flesh quality, and increased EPA levels and n-3/n-6 ratios in largemouth bass muscle and improved fatty acid quality. BSFO had no effect on antioxidant capacity. In summary, dietary inclusion of black soldier fly oil at 2% significantly improved the flesh quality of largemouth bass without measurable adverse effects, providing a solid basis for BSFO as a novel, sustainable lipid source for aquafeeds.

Author Contributions: Conceptualization, P.W.; methodology, P.W.; software, Z.W. (Zichuan Wang); formal analysis, Z.W. (Zichuan Wang), Y.C., W.Y., Z.W. (Zeting Wang) and Y.K.; investigation, W.Y., Z.W. (Zeting Wang) and Y.K.; resources, P.W.; data curation, Z.W. (Zichuan Wang), Y.C., W.Y. and P.W.;

writing—original draft preparation, Z.W. (Zichuan Wang) and Y.C.; writing—review and editing, Z.W. (Zichuan Wang), Y.C. and P.W.; supervision, P.W., C.C. and Y.Y.; project administration, P.W., C.C. and Y.Y.; funding acquisition, P.W. and Y.Y. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Natural Science Foundation of the Higher Education Institutions of Jiangsu Province (20KJA240001); the Foundation of ReProtein Biotechnology (Suzhou) Co., Ltd. (H211372); and the Priority Academic Program Development of the Jiangsu Higher Education Institutions (PAPD).

Institutional Review Board Statement: This study was conducted according to the SOPs of the Provincial Aquatic Animal Nutrition Key Laboratory of Soochow University and approved by the Animal Welfare Ethics Committee of Soochow University (Approval No. SUDA20250320A12; 20 March 2025).

Informed Consent Statement: Not applicable.

Data Availability Statement: Data are contained within the article.

Acknowledgments: The authors thank all the students and staff who contributed to and supported the entire study.

Conflicts of Interest: The authors declare that this study received funding from the Foundation of ReProtein Biotechnology (Suzhou) Co., Ltd. (H211372). The funder was not involved in the study design; the collection, analysis, or interpretation of data; the writing of the article; or the decision to submit it for publication.

References

1. Peng, S.M.; Chen, L.Q.; Qin, J.G.; Hou, J.L.; Yu, N.; Long, Z.Q.; Ye, J.Y.; Sun, X.J. Effects of replacement of dietary fish oil by soybean oil on growth performance and liver biochemical composition in juvenile black seabream, *Acanthopagrus schlegeli*. *Aquaculture* **2008**, *276*, 154–161. [\[CrossRef\]](#)
2. Moon, S.J.; Lee, J.W. Current views on insect feed and its future. *Entomol. Res.* **2015**, *45*, 283–285. [\[CrossRef\]](#)
3. Diener, S.; Zurbrugg, C.; Tockner, K. Conversion of organic material by black soldier fly larvae: Establishing optimal feeding rates. *Waste Manag. Res.* **2009**, *27*, 603–610. [\[CrossRef\]](#)
4. Ewald, N.; Vidakovic, A.; Langeland, M.; Kiessling, A.; Sampels, S.; Lalander, C. Fatty acid composition of black soldier fly larvae (*Hermetia illucens*)—Possibilities and limitations for modification through diet. *Waste Manag.* **2020**, *102*, 40–47. [\[CrossRef\]](#)
5. Kumar, V.; Fawole, F.J.; Romano, N.; Hossain, M.S.; Labh, S.N.; Overturf, K.; Small, B.C. Insect (black soldier fly, *Hermetia illucens*) meal supplementation prevents the soybean meal-induced intestinal enteritis in rainbow trout and health benefits of using insect oil. *Fish Shellfish. Immunol.* **2021**, *109*, 116–124. [\[CrossRef\]](#)
6. Fawole, F.J.; Labh, S.N.; Hossain, M.S.; Overturf, K.; Small, B.C.; Welker, T.L.; Hardy, R.W.; Kumar, V. Insect (black soldier fly larvae) oil as a potential substitute for fish or soy oil in the fish meal-based diet of juvenile rainbow trout (*Oncorhynchus mykiss*). *Anim. Nutr.* **2021**, *7*, 1360–1370. [\[CrossRef\]](#) [\[PubMed\]](#)
7. Sudha, C.; Ahilan, B.; Felix, N.; Uma, A.; Chidambaram, P.; Prabu, E. Replacement of fish oil with black soldier fly larvae oil and vegetable oils: Effects of growth, whole-body fatty acid profile, digestive enzyme activity, haemato-biochemical responses and muscle growth-related gene expression of juvenile striped catfish, *Pangasianodon hypophthalmus*. *Aquac. Res.* **2022**, *53*, 3097–3111. [\[CrossRef\]](#)
8. Li, S.L.; Ji, H.; Zhang, B.X.; Tian, J.J.; Zhou, J.S.; Yu, H.B. Influence of black soldier fly (*Hermetia illucens*) larvae oil on growth performance, body composition, tissue fatty acid composition and lipid deposition in juvenile Jian carp (*Cyprinus carpio* var. *Jian*). *Aquaculture* **2016**, *465*, 43–52. [\[CrossRef\]](#)
9. Moutinho, S.; Oliva-Teles, A.; Pulido-Rodríguez, L.; Parisi, G.; Magalhaes, R.; Monroig, O.; Peres, H. Effects of black soldier fly (*Hermetia illucens*) larvae oil on fillet quality and nutritional traits of gilthead seabream. *Aquaculture* **2024**, *579*, 740219. [\[CrossRef\]](#)
10. Goda, A.M.A.; El-Haroun, E.; Nazmi, H.; Van Doan, H.; Aboseif, A.M.; Taha, M.K.S.; Abou Shabana, N.M. Black soldier fly oil-based diets enriched in lauric acid enhance growth, hematological indices, and fatty acid profiles of Nile tilapia, *Oreochromis niloticus* fry. *Aquac. Rep.* **2024**, *37*, 102269. [\[CrossRef\]](#)
11. Xia, M.; Hu, Z.C.; Wang, G.; Yuan, X.T.; Ji, H.; Sun, J.; Yu, H.B. Evaluation of black soldier fly (*Hermetia illucens*) larvae oil as a potential lipid source for largemouth bass (*Micropterus salmoides*): Growth performance, myofiber development and glucolipid metabolism. *Anim. Feed. Sci. Technol.* **2024**, *318*, 116102. [\[CrossRef\]](#)

12. Wang, D.; Yao, H.; Li, Y.H.; Xu, Y.J.; Ma, X.F.; Wang, H.P. Global diversity and genetic landscape of natural populations and hatchery stocks of largemouth bass *Micropterus salmoides* across American and Asian regions. *Sci. Rep.* **2019**, *9*, 16697. [CrossRef]
13. FAO. *The State of World Fisheries and Aquaculture 2024*; FAO: Rome, Italy, 2024; Available online: <https://openknowledge.fao.org/handle/20.500.14283/cd0683en> (accessed on 25 September 2025).
14. *National Food Safety Standard GB5009.3–2016*; Determination of Moisture in Foods. China Standard Press: Beijing, China, 2016.
15. *Feed Industry Standard GB/T6432-2018*; Determination of Crude Protein in Foods-Kjeldahl Method. China Standard Press: Beijing, China, 2018.
16. *Feed Industry Standard GB/T6433-2006*; Determination of Crude Fat in Foods. China Standard Press: Beijing, China, 2006.
17. *National Food Safety Standard GB5009.4-2016*; Determination of Ash in Foods. China Standard Press: Beijing, China, 2016.
18. Pfaffl, M.W. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **2001**, *29*, e45. [CrossRef]
19. Tang, Y.H.; Zhang, Z.; Weng, M.; Shen, Y.A.; Lai, W.C.; Hao, T.T.; Yao, C.W.; Bu, X.Y.; Du, J.L.; Li, Y.R.; et al. Glycerol monolaurate improved intestinal barrier, antioxidant capacity, inflammatory response and microbiota dysbiosis in large yellow croaker (*Larimichthys crocea*) fed with high soybean oil diets. *Fish Shellfish. Immunol.* **2023**, *141*, 109031. [CrossRef]
20. Ye, Y.T. Technology integration and innovation to improve the edible quality of freshwater cultured fish. *Feed. Ind.* **2022**, *43*, 1–11. [CrossRef]
21. de Huidobro, F.R.; Miguel, E.; Blázquez, B.; Onega, E. A comparison between two methods (Warner-Bratzler and texture profile analysis) for testing either raw meat or cooked meat. *Meat Sci.* **2005**, *69*, 527–536. [CrossRef] [PubMed]
22. Yuan, H.L.; Hu, J.R.; Li, X.C.; Sun, Q.X.; Tan, X.H.; You, C.H.; Dong, Y.W.; Huang, Y.H.; Zhou, M. Dietary black soldier fly oil enhances growth performance, flesh quality, and health status of largemouth bass (*Micropterus salmoides*). *Anim. Nutr.* **2024**, *18*, 234–245. [CrossRef] [PubMed]
23. Miron, L.; Montevicchi, G.; Macavei, L.I.; Maistrello, L.; Antonelli, A.; Thomas, M. Effect of black soldier fly larvae protein on the texture of meat analogues. *Lwt-Food Sci. Technol.* **2023**, *181*, 114745. [CrossRef]
24. Chen, Y.K.; Zhong, J.; Chen, X.Q.; Li, X.M.; Pu, H.Q.; Chen, B.Y.; Guo, Y.C.; Chen, A.Q.; Li, W.J.; Hu, P.; et al. Dietary astaxanthin alleviates black soldier fly oil-induced negative changes of fatty acids content and muscle quality on (*Oncorhynchus mykiss*) via mammalian target of rapamycin and AMP-activated protein kinase pathway. *Anim. Nutr.* **2024**, *19*, 313–324. [CrossRef]
25. Boleman, S.J.; Boleman, S.L.; Miller, R.K.; Taylor, J.F.; Cross, H.R.; Wheeler, T.L.; Koohmaraie, M.; Shackelford, S.D.; Miller, M.F.; West, R.L.; et al. Consumer evaluation of beef of known categories of tenderness. *J. Anim. Sci.* **1997**, *75*, 1521–1524. [CrossRef]
26. Sampath, W.; Zhang, Y.; Liu, J.H.; Yang, M.X.; Zhang, W.B.; Mai, K.S. Dietary taurine improves muscle growth and texture characteristics in juvenile turbot (*Scophthalmus maximus*). *Aquac. Rep.* **2020**, *17*, 100305. [CrossRef]
27. Hu, Z.C.; Li, H.D.; Liu, S.; Xue, R.R.; Sun, J.; Ji, H. Assessment of black soldier fly (*Hermetia illucens*) larvae meal as a potential substitute for soybean meal on growth performance and flesh quality of grass carp *Ctenopharyngodon idellus*. *Anim. Nutr.* **2023**, *14*, 425–449. [CrossRef] [PubMed]
28. Valente, L.M.P.; Cornet, J.; Donnay-Moreno, C.; Gouygou, J.P.; Bergé, J.P.; Bacelar, M.; Escórcio, C.; Rocha, E.; Malhao, F.; Cardinal, M. Quality differences of gilthead sea bream from distinct production systems in Southern Europe: Intensive, integrated, semi-intensive or extensive systems. *Food Control* **2011**, *22*, 708–717. [CrossRef]
29. Stratos, I.; Li, Z.D.; Rotter, R.; Herlyn, P.; Mittlmeier, T.; Vollmar, B. Inhibition of caspase mediated apoptosis restores muscle function after crush injury in rat skeletal muscle. *Apoptosis* **2012**, *17*, 269–277. [CrossRef]
30. Lin, F.; Lin, J.L.; Liu, X.; Yuan, Y.Y.; Liu, G.Q.; Ye, X.K. Effects of temperature on muscle growth and collagen deposition in zebrafish (*Danio rerio*). *Aquac. Rep.* **2022**, *22*, 100952. [CrossRef]
31. Burjanadze, T.V. Hydroxyproline Content and Location in Relation to Collagen Thermal-Stability. *Biopolymers* **1979**, *18*, 931–938. [CrossRef]
32. Ostbye, T.K.K.; Ruyter, B.; Standal, I.B.; Stien, L.H.; Bahuaud, D.; Dessen, J.E.; Latif, M.S.; Fyhn-Terjesen, B.; Rorvik, K.A.; Morkore, T. Functional amino acids stimulate muscle development and improve fillet texture of Atlantic salmon. *Aquac. Nutr.* **2018**, *24*, 14–26. [CrossRef]
33. Weintraub, H. The Myod Family and Myogenesis-Redundancy, Networks, and Thresholds. *Cell* **1993**, *75*, 1241–1244. [CrossRef]
34. Kitajima, Y.; Yoshioka, K.; Suzuki, N. The ubiquitin-proteasome system in regulation of the skeletal muscle homeostasis and atrophy: From basic science to disorders. *J. Physiol. Sci.* **2020**, *70*, 40. [CrossRef]
35. Rodriguez, J.; Vernus, B.; Chelch, I.; Cassar-Malek, I.; Gabillard, J.C.; Sassi, A.H.; Seiliez, I.; Picard, B.; Bonniieu, A. Myostatin and the skeletal muscle atrophy and hypertrophy signaling pathways. *Cell. Mol. Life Sci.* **2014**, *71*, 4361–4371. [CrossRef]
36. Coelho, P.D.; Guarnier, F.A.; Figueiredo, L.B.; Zaramela, L.S.; Pacini, E.S.A.; Godinho, R.O.; Gomes, M.D. Identification of potential target genes associated with the reversion of androgen-dependent skeletal muscle atrophy. *Arch. Biochem. Biophys.* **2019**, *663*, 173–182. [CrossRef]
37. Kakareka, K.; Waddell, D. Characterizing the Role of Muscle-Specific RING Finger 1 (MuRF1) as a Regulator of Atrophy-Induced Gene Expression. *Faseb J.* **2017**, *31*, 1021.4. [CrossRef]

38. Alami-Durante, H.; Cluzeaud, M.; Bazin, D.; Schrama, J.W.; Saravanan, S.; Geurden, I. Muscle growth mechanisms in response to isoenergetic changes in dietary non-protein energy source at low and high protein levels in juvenile rainbow trout. *Comp. Biochem. Physiol. A-Mol. Integr. Physiol.* **2019**, *230*, 91–99. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Richter, B.L.; Silva, T.S.D.; Michelato, M.; Marinho, M.T.; Gonçalves, G.S.; Furuya, W.M. Combination of lysine and histidine improves growth performance, expression of muscle growth-related genes and fillet quality of grow-out Nile tilapia. *Aquac. Nutr.* **2021**, *27*, 568–580. [\[CrossRef\]](#)
40. Valente, L.M.P.; Cabral, E.M.; Sousa, V.; Cunha, L.; Fernandes, J.M.O. Plant protein blends in diets for Senegalese sole affect skeletal muscle growth, flesh texture and the expression of related genes. *Aquaculture* **2016**, *453*, 77–85. [\[CrossRef\]](#)
41. Thomas, G.; Hall, M.N. TOR signalling and control of cell growth. *Curr. Opin. Cell Biol.* **1997**, *9*, 782–787. [\[CrossRef\]](#)
42. Lushchak, O.; Strilbytska, O.; Piskovatska, V.; Storey, K.B.; Koliada, A.; Vaiserman, A. The role of the TOR pathway in mediating the link between nutrition and longevity. *Mech. Ageing Dev.* **2017**, *164*, 127–138. [\[CrossRef\]](#)
43. Goodman, C.A.; Frey, J.W.; Mabrey, D.M.; Jacobs, B.L.; Lincoln, H.C.; You, J.S.; Hornberger, T.A. The role of skeletal muscle mTOR in the regulation of mechanical load-induced growth. *J. Physiol.-Lond.* **2011**, *589*, 5485–5501. [\[CrossRef\]](#)
44. Renna, M.; Schiavone, A.; Gai, F.; Dabbou, S.; Lussiana, C.; Malfatto, V.; Prearo, M.; Capucchio, M.T.; Biasato, I.; Biasibetti, E.; et al. Evaluation of the suitability of a partially defatted black soldier fly (*Hermetia illucens* L.) larvae meal as ingredient for rainbow trout (*Oncorhynchus mykiss* Walbaum) diets. *J. Anim. Sci. Biotechnol.* **2017**, *8*, 57. [\[CrossRef\]](#)
45. Henry, M.; Gasco, L.; Piccolo, G.; Fountoulaki, E. Review on the use of insects in the diet of farmed fish: Past and future. *Anim. Feed. Sci. Technol.* **2015**, *203*, 1–22. [\[CrossRef\]](#)
46. Fabrikov, D.; Barroso, F.G.; Sánchez-Muros, M.J.; Hidalgo, M.C.; Cardenete, G.; Tomás-Almenar, C.; Melenchón, F.; Guil-Guerrero, J.L. Effect of feeding with insect meal diet on the fatty acid compositions of sea bream (*Sparus aurata*), tench (*Tinca tinca*) and rainbow trout (*Oncorhynchus mykiss*) fillets. *Aquaculture* **2021**, *545*, 737170. [\[CrossRef\]](#)
47. Dong, Y.Z.; Wei, Y.; Wang, L.; Song, K.; Zhang, C.X.; Lu, K.L.; Rahimnejad, S. Dietary n-3/n-6 polyunsaturated fatty acid ratio modulates growth performance in spotted seabass (*Lateolabrax maculatus*) through regulating lipid metabolism, hepatic antioxidant capacity and intestinal health. *Anim. Nutr.* **2023**, *14*, 20–31. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Grootendorst-van Mil, N.H.; Tiemeier, H.; Steenweg-de Graaff, J.; Koletzko, B.; Demmelmair, H.; Jaddoe, V.W.V.; Steegers, E.A.P.; Steegers-Theunissen, R.P.M. Maternal plasma n-3 and n-6 polyunsaturated fatty acids during pregnancy and features of fetal health: Fetal growth velocity, birth weight and duration of pregnancy. *Clin. Nutr.* **2018**, *37*, 1367–1374. [\[CrossRef\]](#) [\[PubMed\]](#)
49. Gyan, W.R.; Yang, Q.H.; Tan, B.P.; Dong, X.H.; Chi, S.Y.; Liu, H.Y.; Zhang, S. Effects of replacing fishmeal with dietary dried distillers grains with solubles on growth, serum biochemical indices, antioxidative functions, and disease resistance for *Litopenaeus vannamei* juveniles. *Aquac. Rep.* **2021**, *21*, 100821. [\[CrossRef\]](#)
50. Lin, H.X.; Yang, Q.H.; Wang, A.J.; Wang, J.X.; Tan, B.P.; Ray, G.W.; Dong, X.H.; Chi, S.Y.; Liu, H.Y.; Zhang, S. Effects of fish meal under different storage conditions on growth, serum biochemical indices and antioxidant capacity for juvenile grouper *Epinephelus coioides*. *Aquac. Nutr.* **2021**, *27*, 723–733. [\[CrossRef\]](#)
51. Anuar, N.S.; Shafie, S.A.; Maznan, M.A.F.; Zin, N.; Azmi, N.A.S.; Raoof, R.A.; Myrzakozha, D.; Samsulrizal, N. Lauric acid improves hormonal profiles, antioxidant properties, sperm quality and histomorphometric changes in testis and epididymis of streptozotocin-induced diabetic infertility rats. *Toxicol. Appl. Pharmacol.* **2023**, *470*, 116558. [\[CrossRef\]](#)
52. Hu, J.R.; Yi, J.C.; Wang, G.X.; Mo, W.Y.; Huang, Y.H. Effects of dietary soybean oil replace with black soldier fly larvae oil on growth performance, plasma biochemical indexes and liver lipid droplets of juvenile yellow catfish. *Acta Hydrobiol. Sin.* **2020**, *44*, 717–727.
53. Li, X.H.; Yi, Y.M.; Wu, J.H.; Yang, Q.H.; Tan, B.P.; Chi, S.Y. Effects of Plant-Derived Glycerol Monolaurate (GML) Additive on the Antioxidant Capacity, Anti-Inflammatory Ability, Muscle Nutritional Value, and Intestinal Flora of Hybrid Grouper (*Epinephelus fuscoguttatus*♀ × *Epinephelus lanceolatus*♂). *Metabolites* **2022**, *12*, 1089. [\[CrossRef\]](#)
54. Lin, H.X.; Tan, B.P.; Yang, Q.H.; Chi, S.Y.; Wei, H.J.; Wu, Y.C.; Ray, G.W.; Yohana, M.A. Effects of Dietary Glycerol Monolaurate on Growth, Antioxidant Capacity and Lipid Metabolism in Cage-Farmed Pompano (*Trachinotus ovatus*) Juveniles. *Front. Mar. Sci.* **2022**, *9*, 914134. [\[CrossRef\]](#)
55. Li, X.H.; Zhu, D.W.J.; Mao, M.L.; Wu, J.W.; Yang, Q.H.; Tan, B.P.; Chi, S.Y. Glycerol Monolaurate Alleviates Oxidative Stress and Intestinal Flora Imbalance Caused by Salinity Changes for Juvenile Grouper. *Metabolites* **2022**, *12*, 1268. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.