

Editorial

Utilizing Extracorporeal Shockwave Therapy for In-Season Athletes

Hye Chang Rhim ¹, Joanne Borg-Stein ¹, Steven Sampson ^{2,†} and Adam S. Tenforde ^{1,*,†}

¹ Department of Physical Medicine and Rehabilitation, Harvard Medical School, Boston, MA 02115, USA; hrhim@mgh.harvard.edu (H.C.R.)

² David Geffen School of Medicine, University of California, Los Angeles, CA 90095, USA

* Correspondence: atenforde@mgh.harvard.edu

† These authors contributed equally to this work.

Introduction

An athlete's health and availability to train and compete at an optimal performance level is a growing focus for professional sports organizations. Back-to-back competitions with limited recovery time, along with travelling across multiple time zones, are inherent challenges [1]. Studies suggest that a higher rate of injuries are sustained during the pre-season and during competitions [2]. These injuries include, but are not limited to, muscle strains [3], tendinopathies [4], and injuries to bone [1,5]. Athletic injuries can be challenging to treat in-season, with unpredictable healing times following treatment interventions. Surgical management is typically reserved for off-season athletes, as recovery commonly requires a substantial rehabilitation period of 6 to 9 months post-operatively [4]. Identifying methods to treat injuries that exert positive clinical effects within 3 months would be desirable for in-season athletes. Emerging research suggests that extracorporeal shockwave therapy (ESWT) may represent an effective treatment to address sports-related injuries for in-season athletes and accelerate return to play.

What Is ESWT?

Shockwaves are a type of energy that has biological effects at cellular, tissue, and organ levels. Some of the proposed mechanisms of action for ESWT include increased collagen synthesis [6], cellular proliferation and wound healing [7,8], pain reduction [9], and neovascularization [10]. ESWT currently has two primary modes of delivery: radial shockwave therapy (R-SWT) and focused shockwave therapy (F-SWT). R-SWT generates pressure waves that reach lower speeds and have lower peak pressure and therefore work on more superficial structures. On the other hand, F-SWT has the capacity to achieve deeper penetration from the site of application [11]. While both forms of ESWT have been used to treat a variety of conditions, the differences in their mechanistic effects may lead to different outcomes for a given condition.

ESWT has been shown to be effective in common athletic injuries, including plantar fasciitis [12], Achilles tendinopathy [13], medial tibial stress syndrome [14], and proximal hamstring tendinopathy [15]. Some of the potential side effects include post-procedural pain, skin erythema, skin bruising, hematoma formation, nerve irritation, and superficial edema [11]. ESWT has been known to be effective for the longitudinal management of musculoskeletal injuries and has a favorable safety profile, with recent work suggesting feasibility to support utilization of treatment in the care of in-season athletes.

What High-Level Evidence Is Available to Support the Use of ESWT for In-Season Athletes?

Muscle injury: ESWT was shown to increase muscle elasticity, muscular tone, and muscular recruitment in selected muscles of healthy athletes within 30 days after 3 sessions



Citation: Rhim, H.C.; Borg-Stein, J.; Sampson, S.; Tenforde, A.S. Utilizing Extracorporeal Shockwave Therapy for In-Season Athletes. *Healthcare* **2023**, *11*, 1006. <https://doi.org/10.3390/healthcare11071006>

Received: 7 March 2023

Accepted: 30 March 2023

Published: 1 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

of ESWT [16], suggesting its potential role in muscle recovery. A single session of F-SWT was found to provide relief of pain, increase in force, and improve pain-associated impairments to daily living following eccentric exercise-induced, delayed-onset muscle soreness [17].

Chronic injuries: Return to sport may be accelerated by the use of ESWT with medial tibial stress syndrome and bone trauma.

A prospective observational study evaluated athletes treated with a graded running program and F-SWT over 9 weeks for medial tibial stress syndrome. The authors demonstrated significantly faster time to full recovery, defined by running 18 min consecutively without pain, compared to a graded running program alone [18]. A randomized controlled trial in military cadets with medial tibial stress syndrome observed one session of F-SWT and exercise program resulted in longer running capacity and improvement in pain at four weeks [19].

A randomized controlled trial investigated European football players with groin pain with bone edema of the pubic bone (osteitis pubis). Visual analog scale for pain and Hip Disability and Osteoarthritis Outcome Score were significantly improved in the group receiving ESWT at 1 and 3 months. The athletes who received ESWT were also able to return to football significantly earlier (73.2 days vs. 102.6 days) [20].

One randomized controlled trial in jumping athletes identified limited relief at one week and no sustained benefit of 3 sessions of F-SWT for management of patellar tendinopathy during the competitive season [21]. The authors attributed these findings possibly to not adding other exercises or treatments such as eccentric training and not modifying training or competitions.

What Are the Advantages of ESWT over Alternative Injection Treatments Such as Corticosteroids and Platelet-Rich Plasma?

Corticosteroid injections may offer rapid relief, but may increase the risk of atrophy, pain, and tendon or soft tissue rupture [22]. Furthermore, needle injection therapies in an athlete have potential complications, including infection, bleeding, increased pain, and prolonged recovery time [22,23]. Unlike these interventions, athletes may continue activities as tolerated with ESWT [24]. R-SWT for acute muscle injuries was tolerated well by elite European football players, even when applied on a daily basis [25]. Furthermore, while F-SWT alone was not shown to be beneficial in jumping athletes, the athletes continued to participate in both training and matches without noticeable adverse events during treatment [21].

Platelet-rich plasma is commonly used for sports medicine injuries given the mechanism of action to stimulate tissue healing, reduce pain, without concerns of cellular toxicity [26]. A meta-analysis, including five randomized controlled trials, suggests that the use of platelet-rich plasma may result in an earlier return to sport in the treatment of acute grade I or II muscle strains, excluding those isolated to the hamstring muscle [27]. Despite this finding, based on the systematic review of post-procedure protocols following platelet-rich plasma injections, weight bearing limitation or activity limitation has been imposed immediately after the injection of platelet-rich plasma and up to 7 days and return to play restricted up to 4-6 weeks [28]. A similar lack of clinical results is observed in the management of acute hamstring injuries in competitive and recreational athletes with intramuscular platelet-rich plasma [29]. However, this involved substandard platelet concentration and dosage, limiting researchers to the conclusion that plasma concentrates rather than platelet-rich plasma is ineffective for acute hamstring injuries.

Clinical Applications and Future Work

R-SWT is classified as a low-risk device with a good safety profile that can be used for in-season athletes to facilitate their return to sport in selected injuries. The use of ESWT may be favorable for in-season athletes following load management strategies to allow

for healing, or as a bridge for more advanced procedures such as platelet-rich plasma or surgery for off-season use.

There are some case series and reports that support the use of ESWT on physically active or athletic populations with early outcomes for sports injuries including acute muscle injuries [25], myositis ossificans [30,31], plantar fasciitis [32], Achilles tendinosis [33], proximal hamstring tendinopathy [33], and non-union stress fractures [24,34]. Future high-quality studies should aim to capture larger cohorts of athletes as well as early outcomes such as 3 and 6 weeks, not limited to outcomes 12 weeks or beyond that have been commonly reported. ESWT should also be considered an adjunctive tool for managing commonly encountered athletic injuries. Future research will explore its combination with orthobiologics to help with perioperative and post-operative recovery, including anterior cruciate ligament reconstruction [35].

Conclusions

ESWT offers the potential to accelerate recovery from athletic injury and save athletic clubs valuable resources. Further human athlete studies using ESWT on larger cohorts are needed to explore optimal indications, dosing, and frequency parameters.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest. S.S. is a medical consultant for Globus Medical and Bioventus. A.T. serves as Senior editor for PM&R Journal. He gives professional talks, such as grand rounds and medical conference plenary lectures, and receives honoraria from conference organizers. He has participated in research funded by the Arnold P. Gold Foundation (physician and patient care disparities), the Football Player Health Study at Harvard (health in American-Style Football players), the American Medical Society for Sports Medicine (bone density research), the Uniform Health Service and Enovis (Achilles tendinopathy). He is a paid consultant for State Farm Insurance and Strava.

References

1. Torres-Ronda, L.; Gámez, I.; Robertson, S.; Fernández, J. Epidemiology and injury trends in the National Basketball Association: Pre- and per-COVID-19 (2017–2021). *PLoS ONE* **2022**, *17*, e0263354. [[CrossRef](#)] [[PubMed](#)]
2. Hootman, J.M.; Dick, R.; Agel, J. Epidemiology of collegiate injuries for 15 sports: Summary and recommendations for injury prevention initiatives. *J. Athl. Train.* **2007**, *42*, 311–319. [[PubMed](#)]
3. Ekstrand, J.; Häggglund, M.; Waldén, M. Epidemiology of muscle injuries in professional football (soccer). *Am. J. Sport. Med.* **2011**, *39*, 1226–1232. [[CrossRef](#)]
4. Cook, J.L.; Purdam, C.R. The challenge of managing tendinopathy in competing athletes. *Br. J. Sport. Med.* **2014**, *48*, 506–509. [[CrossRef](#)] [[PubMed](#)]
5. Arendt, E.; Agel, J.; Heikes, C.; Griffiths, H. Stress injuries to bone in college athletes: A retrospective review of experience at a single institution. *Am. J. Sport. Med.* **2003**, *31*, 959–968. [[CrossRef](#)] [[PubMed](#)]
6. Bosch, G.; Lin, Y.L.; van Schie, H.T.; van De Lest, C.H.; Barneveld, A.; van Weeren, P.R. Effect of extracorporeal shock wave therapy on the biochemical composition and metabolic activity of tenocytes in normal tendinous structures in ponies. *Equine Vet. J.* **2007**, *39*, 226–231. [[CrossRef](#)]
7. Wang, F.S.; Yang, K.D.; Chen, R.F.; Wang, C.J.; Sheen-Chen, S.M. Extracorporeal shock wave promotes growth and differentiation of bone-marrow stromal cells towards osteoprogenitors associated with induction of TGF-beta1. *J. Bone Jt. Surg. Br.* **2002**, *84*, 457–461. [[CrossRef](#)]
8. Chen, Y.J.; Wang, C.J.; Yang, K.D.; Kuo, Y.R.; Huang, H.C.; Huang, Y.T.; Sun, Y.C.; Wang, F.S. Extracorporeal shock waves promote healing of collagenase-induced Achilles tendinitis and increase TGF-beta1 and IGF-I expression. *J. Orthop. Res.* **2004**, *22*, 854–861. [[CrossRef](#)]
9. Wess, O.J. A neural model for chronic pain and pain relief by extracorporeal shock wave treatment. *Urol. Res.* **2008**, *36*, 327–334. [[CrossRef](#)]

10. Wang, C.J.; Huang, H.Y.; Pai, C.H. Shock wave-enhanced neovascularization at the tendon-bone junction: An experiment in dogs. *J. Foot Ankle Surg.* **2002**, *41*, 16–22. [[CrossRef](#)]
11. Tenforde, A.S.; Borgstrom, H.E.; DeLuca, S.; McCormack, M.; Singh, M.; Hoo, J.S.; Yun, P.H. Best practices for extracorporeal shockwave therapy in musculoskeletal medicine: Clinical application and training consideration. *PM R* **2022**, *14*, 611–619. [[CrossRef](#)]
12. Rhim, H.C.; Kwon, J.; Park, J.; Borg-Stein, J.; Tenforde, A.S. A Systematic Review of Systematic Reviews on the Epidemiology, Evaluation, and Treatment of Plantar Fasciitis. *Life* **2021**, *11*, 1287. [[CrossRef](#)] [[PubMed](#)]
13. Rhim, H.C.; Kim, M.S.; Choi, S.; Tenforde, A.S. Comparative Efficacy and Tolerability of Nonsurgical Therapies for the Treatment of Midportion Achilles Tendinopathy: A Systematic Review with Network Meta-analysis. *Orthop. J. Sport. Med.* **2020**, *8*, 2325967120930567. [[CrossRef](#)] [[PubMed](#)]
14. Forogh, B.; Karimzad, Y.; Babaei-Ghazani, A.; Janbazi, L.; Cham, M.B.; Abdolghaderi, S. Effect of extracorporeal shockwave therapy on medial tibial stress syndrome: A systematic review. *Curr. Orthop. Pract.* **2022**, *33*, 384–392. [[CrossRef](#)]
15. Korakakis, V.; Whiteley, R.; Tzavara, A.; Malliaropoulos, N. The effectiveness of extracorporeal shockwave therapy in common lower limb conditions: A systematic review including quantification of patient-rated pain reduction. *Br. J. Sport. Med.* **2018**, *52*, 387–407. [[CrossRef](#)]
16. Notarnicola, A.; Covelli, I.; Maccagnano, G.; Marvulli, R.; Mastromauro, L.; Ianieri, G.; Boodhoo, S.; Turitto, A.; Petruzzella, L.; Fari, G. Extracorporeal shockwave therapy on muscle tissue: The effects on healthy athletes. *J. Biol. Regul. Homeost.* **2018**, *32*, 185–193.
17. Fleckenstein, J.; Fritton, M.; Himmelreich, H.; Banzer, W. Effect of a Single Administration of Focused Extracorporeal Shock Wave in the Relief of Delayed-Onset Muscle Soreness: Results of a Partially Blinded Randomized Controlled Trial. *Arch. Phys. Med. Rehabil.* **2017**, *98*, 923–930. [[CrossRef](#)] [[PubMed](#)]
18. Moen, M.H.; Rayer, S.; Schipper, M.; Schmikli, S.; Weir, A.; Tol, J.L.; Backx, F.J. Shockwave treatment for medial tibial stress syndrome in athletes; a prospective controlled study. *Br. J. Sport. Med.* **2012**, *46*, 253–257. [[CrossRef](#)]
19. Gomez Garcia, S.; Ramon Rona, S.; Gomez Tinoco, M.C.; Benet Rodriguez, M.; Chaustre Ruiz, D.M.; Cardenas Letrado, F.P.; Lopez-Illescas Ruiz, A.; Alarcon Garcia, J.M. Shockwave treatment for medial tibial stress syndrome in military cadets: A single-blind randomized controlled trial. *Int. J. Surg.* **2017**, *46*, 102–109. [[CrossRef](#)]
20. Schöberl, M.; Prantl, L.; Loose, O.; Zellner, J.; Angele, P.; Zeman, F.; Spreitzer, M.; Nerlich, M.; Krutsch, W. Non-surgical treatment of pubic overload and groin pain in amateur football players: A prospective double-blinded randomised controlled study. *Knee Surg. Sport. Traumatol. Arthrosc.* **2017**, *25*, 1958–1966. [[CrossRef](#)] [[PubMed](#)]
21. Zwerver, J.; Hartgens, F.; Verhagen, E.; van der Worp, H.; van den Akker-Scheek, I.; Diercks, R.L. No effect of extracorporeal shockwave therapy on patellar tendinopathy in jumping athletes during the competitive season: A randomized clinical trial. *Am. J. Sport. Med.* **2011**, *39*, 1191–1199. [[CrossRef](#)] [[PubMed](#)]
22. Nichols, A.W. Complications associated with the use of corticosteroids in the treatment of athletic injuries. *Clin. J. Sport Med.* **2005**, *15*, 370–375. [[CrossRef](#)] [[PubMed](#)]
23. Cheng, J.; Abdi, S. Complications of joint, tendon, and muscle injections. *Tech. Reg. Anesth. Pain Manag.* **2007**, *11*, 141–147. [[CrossRef](#)]
24. Saxena, A.; Fullem, B.; Gerdesmeyer, L. Treatment of Medial Tibial Stress Syndrome with Radial Soundwave Therapy in Elite Athletes: Current Evidence, Report on Two Cases, and Proposed Treatment Regimen. *J. Foot Ankle Surg.* **2017**, *56*, 985–989. [[CrossRef](#)] [[PubMed](#)]
25. Morgan, J.P.M.; Hamm, M.; Schmitz, C.; Brem, M.H. Return to play after treating acute muscle injuries in elite football players with radial extracorporeal shock wave therapy. *J. Orthop. Surg. Res.* **2021**, *16*, 708. [[CrossRef](#)] [[PubMed](#)]
26. Taylor, D.W.; Petrera, M.; Hendry, M.; Theodoropoulos, J.S. A systematic review of the use of platelet-rich plasma in sports medicine as a new treatment for tendon and ligament injuries. *Clin. J. Sport Med.* **2011**, *21*, 344–352. [[CrossRef](#)] [[PubMed](#)]
27. Sheth, U.; Dwyer, T.; Smith, I.; Wasserstein, D.; Theodoropoulos, J.; Takhar, S.; Chahal, J. Does Platelet-Rich Plasma Lead to Earlier Return to Sport When Compared with Conservative Treatment in Acute Muscle Injuries? A Systematic Review and Meta-analysis. *Arthroscopy* **2018**, *34*, 281–288.e1. [[CrossRef](#)]
28. Townsend, C.; Von Rickenbach, K.J.; Bailowitz, Z.; Gellhorn, A.C. Post-Procedure Protocols Following Platelet-Rich Plasma Injections for Tendinopathy: A Systematic Review. *PM R* **2020**, *12*, 904–915. [[CrossRef](#)]
29. Reurink, G.; Goudswaard, G.J.; Moen, M.H.; Weir, A.; Verhaar, J.A.; Bierma-Zeinstra, S.M.; Maas, M.; Tol, J.L. Platelet-rich plasma injections in acute muscle injury. *N. Engl. J. Med.* **2014**, *370*, 2546–2547. [[CrossRef](#)]
30. Buselli, P.; Coco, V.; Notarnicola, A.; Messina, S.; Saggini, R.; Tafuri, S.; Moretti, L.; Moretti, B. Shock waves in the treatment of post-traumatic myositis ossificans. *Ultrasound Med. Biol.* **2010**, *36*, 397–409. [[CrossRef](#)]
31. Torrance, D.A.; Degraauw, C. Treatment of post-traumatic myositis ossificans of the anterior thigh with extracorporeal shock wave therapy. *J. Can. Chiropr. Assoc.* **2011**, *55*, 240–246.
32. Moretti, B.; Garofalo, R.; Patella, V.; Sisti, G.L.; Corrado, M.; Mouhsine, E. Extracorporeal shock wave therapy in runners with a symptomatic heel spur. *Knee Surg. Sport. Traumatol. Arthrosc.* **2006**, *14*, 1029–1032. [[CrossRef](#)] [[PubMed](#)]
33. Reilly, J.M.; Tenforde, A.S. The Role of Extracorporeal Shockwave Therapy in Return to Competition for Endurance Runners: Two Case Reports. *PM R* **2020**, *12*, 516–517. [[CrossRef](#)] [[PubMed](#)]

34. Taki, M.; Iwata, O.; Shiono, M.; Kimura, M.; Takagishi, K. Extracorporeal shock wave therapy for resistant stress fracture in athletes: A report of 5 cases. *Am. J. Sport. Med.* **2007**, *35*, 1188–1192. [[CrossRef](#)] [[PubMed](#)]
35. Wang, C.J.; Ko, J.Y.; Chou, W.Y.; Hsu, S.L.; Ko, S.F.; Huang, C.C.; Chang, H.W. Shockwave therapy improves anterior cruciate ligament reconstruction. *J. Surg. Res.* **2014**, *188*, 110–118. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.