



Article

# Indoor Exposure and Regional Inhaled Deposited Dose Rate during Smoking and Incense Stick Burning—The Jordanian Case as an Example for Eastern Mediterranean Conditions

Tareq Hussein <sup>1,2</sup>

<sup>1</sup> Environmental and Atmospheric Research Laboratory (EARL), Department of Physics, School of Science, The University of Jordan, Amman 11942, Jordan; tareq.hussein@helsinki.fi

<sup>2</sup> Institute for Atmospheric and Earth System Research (INAR/Physics), University of Helsinki, FI-00014 Helsinki, Finland

**Abstract:** Tobacco smoking and incense burning are commonly used in Jordanian microenvironments. While smoking in Jordan is prohibited inside closed spaces, incense burning remains uncontrolled. In this study, particle size distributions (diameter 0.01–25  $\mu\text{m}$ ) were measured and inhaled deposited dose rates were calculated during typical smoking and incense stick-burning scenarios inside a closed room, and the exposure was summarized in terms of number and mass concentrations of submicron ( $PN_{Sub}$ ) and fine particles ( $PM_{2.5}$ ). During cigarette smoking and incense stick-burning scenarios, the particle number concentrations exceeded  $3 \times 10^5 \text{ cm}^{-3}$ . They exceeded  $5 \times 10^5 \text{ cm}^{-3}$  during shisha smoking. The emission rates were  $1.9 \times 10^{10}$ ,  $6.8 \times 10^{10}$ , and  $1.7 \times 10^{10}$  particles/s, respectively, for incense, cigarettes, and shisha. That corresponded to about 7, 80, and 120  $\mu\text{g/s}$ , respectively. Males received higher dose rates than females, with about 75% and 55% in the pulmonary/alveolar during walking and standing, respectively. The total dose rates were in the order of  $10^{12}$ – $10^{13}$  #/h ( $10^3$ – $10^4$   $\mu\text{g/h}$ ), respectively, for  $PN_{Sub}$  and  $PM_{2.5}$ . The above reported concentrations, emissions rates, and dose rates are considered seriously high, recalling the fact that aerosols emitted during such scenarios consist of a vast range of toxicant compounds.

**Keywords:** fine particles; particle losses; particle number size distribution



**Citation:** Hussein, T. Indoor Exposure and Regional Inhaled Deposited Dose Rate during Smoking and Incense Stick Burning—The Jordanian Case as an Example for Eastern Mediterranean Conditions. *Int. J. Environ. Res. Public Health* **2023**, *20*, 587. <https://doi.org/10.3390/ijerph20010587>

Academic Editors: Jolanda Palmisani, Alessia Di Gilio and Gianluigi de Gennaro

Received: 22 November 2022  
Revised: 20 December 2022  
Accepted: 22 December 2022  
Published: 29 December 2022



**Copyright:** © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Incense-burning and aroma products are not widely prohibited worldwide, as some countries still use them extensively [1–8]. In principle, smoking is prohibited in some countries, but this is still not fully obeyed in a large portion of the globe, especially in the Middle East and North Africa (MENA) region [9–23], where tobacco smoking (mainly cigarettes and shisha, which is also known as waterpipe, hookah, narghile, or narghila) is still experienced in homes, restaurants, vehicles, malls, government buildings, and offices; shisha smoking is still widely allowed in coffee shops and offered to teenagers [22,24–41]. Alternatively, other forms of smoking (e.g., IQOS, e-cigarettes, vaping, etc.) have been introduced as safe or less harmful than tobacco or shisha smoking, but these forms of smoking have not been extensively researched for their possible harmful health effects [42–46]. Nevertheless, some research results have confirmed that these types of alternative smoking types do have harmful health effects [43,47–59].

Smoking is not only considered harmful for the smoker themselves, but also harmful for the persons nearby, as passive smokers are exposed to the smoke itself being released and transported in the atmosphere, which is known as being a second-hand smoker [47–50,60–67]. Researchers have also introduced the term “third-hand smokers” [68–79], which is not a new issue in public health but is a less well-understood type of smoke exposure. This happens as residual smoke sorbed onto surfaces (furniture, building materials, clothes, etc.) is re-emitted into the atmosphere even after active smoking is ceased. Third-hand

smoking exposure pathways can be thought of as the inhalation, ingestion, or dermal uptake of pollutants.

Although the Jordanian Ministry of Health has stated that smoking is forbidden in microenvironments (e.g., houses, schools, offices, etc.), people are still violating this law. In fact, Jordan acquires the highest rate of smoking; 8 out of 10 men consume nicotine products such as cigarettes, shisha (waterpipe, hookah, narghile), and e-cigarettes. According to previous statistics [10,11], about 20% of 13-year-old boys (7.5% girls) in Irbid, Jordan consume nicotine products such as shisha and cigarettes compared to about 11.3% in girls. Contrary to smoking, incense burning does not have any regulations in Jordan. These two types of indoor activities impose serious health issues indoors. It is, therefore, very important to quantify the exposure to air pollution produced during smoking and incense burning for Jordanian conditions, which can be a good representative of several nations in the Eastern Mediterranean region.

Tobacco smoke consists of up of 4000 chemicals, including (varying greatly between the type of smoking) volatile organic compounds (VOCs: Acetonitrile, Acrylnitrile, Benzene, 2,5-Dimethylfuran, 3-Ethenylpyridine, Nicotine, Aldehydes (e.g., Formaldehyde, Acetaldehyde, Acrolein, Propionaldehyde, Methacrolein, etc.), Acetone, Pyrrole, Pyrroline, Pyrrolidine, Phenolics, etc.), polycyclic aromatic hydrocarbons (PAHs: Naphthalene, Acenaphthylene, Acenaphthene, Fluorene, Phenanthrene, Anthracene, Fluoranthene, Pyrene, Benz(a)anthracene, Chrysene, benzo(b)-fluoranthene, benzo(k)-fluoranthene, Benzo(a)pyrene, Indeno(1,2,3-cd)pyrene, Dibenz(a,h)anthracene, Benzo(g,h,i)perylene), CO, and heavy metals; these are those ones that have been identified as risk factors for health [32,62,80–101]. Incense burning produces a mixture of particulate matter and a vast range of gases, which includes—and is not limited to—CO, CO<sub>2</sub>, NO<sub>x</sub>, SO<sub>2</sub>, PAHs, VOCs (including benzene, toluene, terpinols, and xylenes), formaldehyde (HCHO), BTEX, and heavy metals [102–114]. As a matter of fact, air pollution produced during tobacco smoking and incense burning is one of the risk factors for cardiovascular diseases, cancer, respiratory illness, and a wide range of other harmful health outcomes [86,87,93,101,115–132]. The World Health Organization (WHO) estimates the mortality rate associated with tobacco smoking as six million people per year, with a projected increase to eight million by the year 2030. Moreover, smoking accounts for six out of eight leading causes of death worldwide and is associated with a large percentage of healthcare costs.

Here, we aim to present the exposure levels and regional inhaled deposited rate in respiratory tracts during the most common combustion activities in Jordanian microenvironments. The investigation included measurements of particle number size distributions (diameter 0.01–25 µm) during typical smoking and incense stick-burning scenarios inside a closed room. The exposure and dose rates were presented in terms of two metrics (number and mass) of submicron and fine particles. In order to put an insight into the exposure scenarios, the regional inhaled deposited dose rates were calculated for an adult male or female during two of the main human activities (walking and standing).

## 2. Materials and Methods

### 2.1. Measurement Setup

The measurement setup was according to the description provided by Hussein et al. [133]. The setup consisted of the following portable instruments: (1) condensation particle counters (CPC, 3007-2, TSI, Shoreview, MN, USA), (2) handheld optical counter (P-Trak 8525, TSI, Shoreview, MN, USA), and (3) and a handheld optical particle counter (AeroTrak 9306-V2, TSI, Shoreview, MN, USA).

The CPC and P-Trak measure submicron particle number concentrations with cutoff sizes of 10 nm and 25 nm, respectively. The sampling flow rate in these two instruments was 0.1 lpm (inlet flow rate 0.7 lpm). The maximum measurable concentrations were  $\sim 10^5 \text{ cm}^{-3}$  and  $5 \times 10^5 \text{ cm}^{-3}$ , respectively. These instruments were operated at 1 s time resolution. Using the AeroTrak extended measured particle diameters to reach 25 µm and

allowed for 6 particle size channels: 0.3, 0.5, 1, 2.5, 10, and 25  $\mu\text{m}$  (optical diameter). The AeroTrak sampling time resolution was 30 s at a flow rate of 2.83 lpm.

## 2.2. Exposure Scenarios

The exposure scenarios included smoking (cigarette and shisha) and incense stick burning. The scenarios were performed inside an office room ( $5 \times 2.7 \times 2.7 \text{ m}^3$ ), which was fully furnished and had its window and door closed. Each scenario was repeated four times. The room was naturally ventilated.

The cigarette smoking was made by burning two cigarettes within a short time (<3 min). Two shishas were smoked at the same time for a period between 30 min and 90 min. The incense stick scenario included burning two sticks (~20 cm long) at the same time for five minutes.

After finishing the smoking and the incense stick burning, the room was kept closed and unoccupied until the particle number concentration reached a level close to what was before the scenario was started. This took a period of longer than three hours.

## 2.3. Data Handling

The measured particle number size distribution (eight channels within 0.01–25  $\mu\text{m}$ ) was generated by processing the data by the three instruments listed in the previous section [22,133]. Basically, the use of three instruments with different cutoffs has the advantage of generating the particle number size distribution by subtracting the concentrations measured with the different instruments. The most important part here is to harmonize the aerosol database measured with different instruments.

The harmonized aerosol database was processed for 1 min average. The particle number size distribution (10 nm–25  $\mu\text{m}$ , 8 channels) was generated by calculating the concentration differences between different instruments: 10–25 nm (CPC and P-Trak difference), 25–300 nm (P-Trak and Aerotrak difference), and 6 channels in the AeroTrak (0.3–0.5  $\mu\text{m}$ , 0.5–1  $\mu\text{m}$ , 1–2.5  $\mu\text{m}$ , 2.5–5  $\mu\text{m}$ , 5–10  $\mu\text{m}$ , and 10–25  $\mu\text{m}$ ).

The size-fractionated number and mass concentrations were calculated over the specified particle size range:

$$PN_{D_{p1}-D_{p2}} = \int_{D_{p1}}^{D_{p2}} n_N^0 \cdot d\log_{10}(D_p) \quad (1)$$

$$PM_{D_{p1}-D_{p2}} = \int_{D_{p1}}^{D_{p2}} n_M^0 \cdot d\log_{10}(D_p) \quad (2)$$

where  $n_N^0 = dN/d\log_{10}(D_p)$  is the lognormal particle number size distribution and  $n_M^0 = dM/d\log_{10}(D_p)$  is the lognormal particle mass size distribution. Here, the particles were assumed spherical, with unit density ( $\rho_p = 1000 \text{ kg/m}^3$ ).

## 2.4. Particle Emission Rate and Loss Rate Calculation—A Simple Indoor Aerosol Model

The size-specific emission rate and losses of aerosol particles were calculated by utilizing a simple indoor aerosol model [23,134–136]. The basic principle is based on the change rate of particle number concentrations calculated for each particle size. At first, the particle loss rate was calculated. Then, the emission rate was calculated and corrected for the particle losses.

While in general, indoor aerosol origin is a combination of indoor or outdoor sources, the indoor aerosol sources in this study are dominated by those sources that are due to smoking and incense burning. The indoor aerosols are eventually lost from the atmosphere by either dry deposition or removal via air exchange. Complex dynamic processes (coagulation, condensation, chemical reactions, etc.) were ignored. Accordingly, indoor aerosols concentrations can be described by the mass balance equation:

$$\frac{dI_i}{dt} = P_i\lambda O_i - (\lambda + \lambda_{d,i})I_i + S_{in,i} \tag{3}$$

where  $t$  is the time;  $I$  and  $O$  are the indoor and outdoor concentrations of the aerosol particles, respectively;  $P$  is the penetration factor of aerosol particles while being transported from the outdoor air into the indoor air;  $\lambda$  is the ventilation rate;  $\lambda_d$  is the dry deposition rate of aerosol particles onto available indoor surfaces; and  $S_{in}$  represents the emission rates from an indoor source. The subscript  $i$  represents a certain particle size range. A well-mixed indoor air domain was ensured because the room size was small enough and a fan was used to produce air convection.

In Equation (3), the second and third terms on the right-hand side are the particle losses (dry deposition and removal by ventilation) and emission rate from an indoor source, respectively. Right after an indoor source is terminated, Equation (3) can be rewritten to calculate the particle losses as

$$\frac{dI_i}{dt} \rightarrow -(\lambda + \lambda_{d,i})I_i \tag{4}$$

and right after a source is started, it can be rewritten to calculate the emission rate as

$$\frac{dI_i}{dt} + \{(\lambda + \lambda_{d,i})I_i\}_{empirical} \rightarrow S_{in,i} \tag{5}$$

Here, Equation (4) is built on the fact that indoor aerosols of indoor origin are dominant (i.e.,  $P\lambda O \ll (\lambda + \lambda_d)I$ ). In Equation (5), the source term is assumed to be dominant (i.e.,  $S_{in} \ll P\lambda O - (\lambda + \lambda_d)I$ ) but it needs to be corrected for the particle losses.

### 2.5. Regional Inhaled Deposited Dose in the Respiratory Tracts

The measured particle number size distribution can be used to calculate the deposited fraction of aerosols in the respiratory tracts for a specific particle diameter range ( $D_{p1}-D_{p2}$ ). Here, we followed our previous approach described elsewhere [137–140]. Here, the respiratory tracts were divided into three main regions: head/throat (H), tracheobronchial (TB), and pulmonary/alveolar (P/Alv), according to the ICRP and MPPD models [141–143], and the calculations require the following:


- The predefined particle deposition fraction ( $DF(D_p)$ ) in the respiratory tracts during exercise or rest (Figure 1) [141–143];
- The minute ventilation ( $V_E, m^3/h$ ), which is the volume of breathed air per time (Table 1);
- The measured particle number size distribution ( $n_N^0(D_p)$ , particles/cm<sup>3</sup>):

$$\text{Dose Rate} = \int_{D_{p1}}^{D_{p2}} V_E DF n_N^0 f \cdot d\log(D_p) \tag{6}$$

Here,  $f$  is a metric conversion for the particle concentration: it is 1 for particle number, and for particle mass it is  $\rho_p D_p^3 \pi / 6$ .

**Table 1.** Minute ventilation ( $V_E, m^3/h$ ) for adult subjects.

Activity	Female	Male	DF Curve Type
Walking (4.0 km/h)	1.20	1.38	Exercise
Standing	0.48	0.66	at rest




**Figure 1.** Deposition fraction (*DF*) curves in an adult: (a) male exercising, (b) male at rest, (c) female exercising, and (d) female at rest.

### 3. Results and Discussion

#### 3.1. Temporal Variation, Size Distributions, and Particle Losses

As examples, the total particle number concentrations during one case of each exposure scenario (cigarette, shisha, and incense) are presented in Figure 2 (left panel). When the scenario started, the fine particle concentrations increased by several factors. After each scenario was over (i.e., smoking or burning), the concentrations decayed gradually as a result of the indoor–outdoor air exchange and dry deposition. During the cigarette and incense scenarios, the concentrations increased continuously until the end of smoking or burning, but during shisha smoking, the concentrations varied as a result of the puffing sequence or the addition of charcoal. The pollution emitted during smoking depends on the puffing and smoking behaviour [144–146]. The emission factor is also dependent on the shisha setup (bowl size, head size, hose length, etc.) [147]. In fact, the temporal variation of the particle concentrations agreed well with previous studies that reported smoking (cigarette and shisha) and incense burning [106,148–150].


Selected particle number size distributions during one case of each exposure scenario are presented in Figure 2 (right panel). The mean particle number and mass size distributions during the almost-steady-state concentration level are shown in Figures 3 and 4. The total particle losses—which are dependent on many factors, including: the particle size distribution, indoor–outdoor air exchange, and dry deposition for each case of the scenarios—are presented in Figure 5. The average total particles losses were  $0.45\text{--}2\text{ h}^{-1}$  for particles smaller than  $0.1\text{ }\mu\text{m}$ . Those for micron particles were  $0.45\text{--}5.5\text{ h}^{-1}$ . According to the losses of particles within the range  $0.1\text{--}1\text{ }\mu\text{m}$ , the ventilation rate ( $\lambda$ ) can be around  $0.5\text{ h}^{-1}$ . This value agrees with typical values of natural ventilation.




**Figure 2.** Particle number concentrations and size distributions during (a,b) incense stick burning, (c,d) cigarette smoking, and (e,f) shisha smoking. The left panel is the time series of different particle size fraction concentrations, where the vertical dashed line at zero hour indicates the start of the scenario, and the right panel is selected particle number size distributions during different time periods.

The three-layer deposition model calculation, as described by Lai and Nazaroff [151], shows underestimation for the total particle losses; room volume  $5 \times 2.7 \times 2.7 \text{ m}^3$  and friction velocity  $u^* = 10 \text{ cm/s}$ . This underestimation is expected because the Lai and Nazaroff calculation [151] is for smooth surfaces. The roughness of indoor surfaces can be taken into account by the three-layer deposition model described by Hussein et al. [152]; the surface roughness parameter ( $F_{rough}$ ) can be smaller than 0.8.


The incense smoke (inside a small stainless-steel test chamber,  $1.2 \times 1.2 \times 1.2 \text{ m}^3$ ) was characterized by a skewed unimodal particle number size distribution to small diameters and varying peak values of 75–170 nm (mean 130 nm) and total particle losses of  $1.0\% \text{ } 2.32 \text{ h}^{-1}$  (mean  $2.4 \text{ h}^{-1}$ ) [150]. In the same study, the cigarette smoke was characterized by a unimodal particle number size distribution, a peak value at around 110 nm, and mean total particle losses of  $3.9 \text{ h}^{-1}$  [150]. In another study inside a small chamber ( $1.016 \times 0.660 \times 1.600 \text{ m}^3$ ), the geometric mean diameter of the incense smoke particles was nearly unimodal, with a geometric mean of 80–15 nm [153]. The  $\text{PM}_{2.5}$  of incense smoke inside a church showed a loss rate in the range of  $1.1 \times 10^{-2}$ – $5.5 \times 10^{-2} \text{ min}^{-1}$  (mean  $3.2 \times 10^{-2} \text{ min}^{-1}$ ) [8].



**Figure 3.** Mean particle number size distributions during each case of the different scenarios: (a) incense stick burning, (b) cigarette smoking, and (c) shisha smoking. (d–f) The corresponding mean particle mass size distributions.



**Figure 4.** Overall mean (a) particle number size distributions and (b) corresponding mass size distributions by assuming spherical particles and unit density.



**Figure 5.** Total particle losses of aerosol particles calculated for each scenario compared to the three-layer model calculations by Lai and Nazaroff [151] for smooth surfaces and Hussein et al. [152] for rough surfaces. The model calculation was made for a room ( $5 \times 2.7 \times 2.7 \text{ m}^3$ ), friction velocity  $u^* = 10 \text{ cm/s}$ , and ventilation rate  $\lambda = 0.5 \text{ h}^{-1}$ . The surface roughness parameter ( $F_{\text{rough}}$ ) is zero for a smooth surface.

### 3.2. Exposure

#### 3.2.1. Incense Sticks

During the incense scenario and right after the two sticks started burning, the fine particle concentrations increased quickly, especially those in the ultrafine particle size range (i.e., UFP, diameter  $< 0.1 \mu\text{m}$ ). The total particle number concentrations were higher than  $3 \times 10^5 \text{ cm}^{-3}$  (corresponding to  $>110 \mu\text{g}/\text{m}^3$ ) (Figure 2a). The duration of the high-concentration period was as long as the two sticks were burning. It is noticed from Figure 2a that the micron particle concentrations increased before starting the incense stick burning; this was basically due to the fact that the room was opened to enter and start the scenario. The evolution of the particle number size distributions is presented in Figure 2b, which clearly shows how the burning affected the fine particles.

In a previous investigation conducted inside a room ( $2.53 \times 2.50 \times 5.15 \text{ m}^3$ ) [106], the total particle number concentration was reported as  $9.1 \times 10^4 \text{ cm}^{-3}$  during incense burning. In another study conducted inside a small chamber ( $1.016 \times 0.660 \times 1.600 \text{ m}^3$ ), the incense smoke (four incense sticks were burned at each of the four corners of the chamber) showed a highest concentration of  $1.1 \times 10^6$ – $2.4 \times 10^6 \text{ cm}^{-3}$  [153]. The total particle number concentrations observed inside an empty room with dimensions of  $2.7 \times 3.8 \times 3.1 \text{ m}^3$  for two types of incense varied within  $1.6 \times 10^5$ – $2.9 \times 10^5 \text{ cm}^{-3}$  (mean  $2.2 \times 10^5 \text{ cm}^{-3}$ ) [108]. The UFP concentrations (inside a stainless-steel test chamber  $1.2 \times 1.2 \times 1.2 \text{ m}^3$ ) [150] had a maximum concentration of the incense smoke within the range of  $5.7 \times 10^4$ – $6.1 \times 10^5 \text{ cm}^{-3}$  (mean  $2.9 \times 10^5 \text{ cm}^{-3}$ ). In the European Accredited (EA) Laboratory of Industrial Measurements (LAMI) at the University of Cassino and Southern Lazio, a test of three incense-burning scenarios yielded  $4.1 \times 10^6$ – $9.8 \times 10^6 \text{ cm}^{-3}$  ( $\text{PM}_{2.5}$  6.3–14.6  $\mu\text{g}/\text{m}^3$ ) [105].

As a cultural and religious custom, incense burning is widely used in some countries [3–8]. The  $\text{PM}_{2.5}$  concentrations inside a church reached values of 36–123  $\mu\text{g}/\text{m}^3$  (mean 56  $\mu\text{g}/\text{m}^3$ ), equivalent to  $5 \times 10^3$ – $1.4 \times 10^4 \text{ cm}^{-3}$  (mean  $6.3 \times 10^3 \text{ cm}^{-3}$ ) [8]. The average  $\text{PM}_{2.5}$  concentration at ten temples was reported to be 660  $\mu\text{g}/\text{m}^3$ , and at two crematoriums, it

was reported to be  $1050 \mu\text{g}/\text{m}^3$ , as a result of incense burning [7]; these values were ten and five times higher than the outdoor concentration reported at the nearby ambient air-monitoring station.

### 3.2.2. Cigarette

During cigarette-smoking scenarios (Figure 2c,d), the total particle number concentrations were higher than  $3 \times 10^5 \text{ cm}^{-3}$ . The duration of smoking two cigarettes was less than three minutes, and consequently, the aerosol emission span time was very short. Initially, fine particles of all sizes were affected as a result of opening the office to start the cigarette-smoking scenario. However, it was evident that particles smaller than  $0.3 \mu\text{m}$  were the most affected, because their concentrations immediately increased significantly right after starting the scenario. Interestingly, after 10 min of starting the smoking, when the UFP concentrations started to decrease, the concentrations of the accumulation mode (diameter  $0.1\text{--}1 \mu\text{m}$ ) started to increase as a result of coagulation of the emitted smoke particles. This was also evident in the evolution of the particle number size distributions shown in Figure 2d. The lifetime of the emitted smoke particles were longer than those emitted from the incense burning.

The UFP concentrations of cigarette smouldering evaluated inside in a stainless-steel test chamber ( $1.2 \times 1.2 \times 1.2 \text{ m}^3$ ) were as high as  $1.2 \times 10^6 \text{ cm}^{-3}$  [150]. The  $\text{PM}_{2.5}$  exposure levels as second-hand smoke (five types of Vogue cigarettes) ranged between  $1.3$  and  $2.2 \text{ mg}/\text{m}^3$  [154]. In another test inside a small chamber ( $2.88 \text{ m}^3$ ), the  $\text{PM}_{2.5}$  exposure levels as second-hand smoke were  $0.7\text{--}1.5 \text{ mg}/\text{m}^3$  during the testing of five types of Marlboro cigarette [155].

Particle concentrations in the mainstream are extremely higher than those inhaled by a passive smoker in typical indoor, outdoor, and occupational environments, as well as vehicles: typically in the range of  $10^3\text{--}10^6 \text{ cm}^{-3}$  [156–179]. It was reported that in the mainstream, the particle number concentration may reach values in the order of  $10^8 \text{ cm}^{-3}$ , as tested for five different brands (B&H, Camel, Marlboro, Merit, Wenston) and characterized by a single lognormal mode with a geometric mean diameter of  $180 \text{ nm}$  and  $220 \text{ nm}$  [180], which agreed well with this study, as well as others reported in the literature [164,181]. In another study, it was shown to be in the range of  $10^5\text{--}10^9 \text{ cm}^{-3}$  in the mainstream [182].

### 3.2.3. Shisha

The scenario of shisha smoking took about an hour, during which the particle number concentrations of all particle sizes smaller than  $10 \mu\text{m}$  were significantly high (total  $> 5 \times 10^5 \text{ cm}^{-3}$ ) when compared to the background concentrations (Figure 2e). Interestingly, the UFP concentrations were steadily high throughout the scenario, but the concentrations of other particle size fractions within the range of  $0.1\text{--}10 \mu\text{m}$  were continuously variable. During shisha smoking, the smokers were frequently moving, and consequently, dust resuspension occurred, as clearly seen in the particle number size distributions shown in Figure 2f. The lifetime of the emitted aerosols from shisha smoking (less than 2 h) was shorter than those emitted from cigarette smoking or incense burning.


The literature about shisha smoke characterization is fairly limited when compared to cigarette smoke or incense burning. However, and as pointed out before, shisha smoke temporal variation in this study was similar to what was previously reported in other studies, and can be explained by puffing intensity and frequency, as well as the addition/changing of charcoal [148,149]. In our previous study reported for indoor air quality inside Jordanian dwellings, houses with smoking activities had aerosol concentrations (both number and mass) higher than houses without smoking activities [22]. Compared to this study, which was inside an office ( $\sim 36.5 \text{ m}^3$ ), these houses reflected real-life conditions inside relatively large indoor environments (apartments and houses), revealing that the submicron particle number concentrations were as high as  $1.5 \times 10^5 \text{ cm}^{-3}$  (equivalent to  $\text{PM}_{2.5} \sim 100 \mu\text{g}/\text{m}^3$ ) inside the dwellings with cigarette smoking. Those with shisha smoking experienced even higher concentrations  $\sim 4 \times 10^5 \text{ cm}^{-3}$  (equivalent to  $\text{PM}_{2.5} \sim 130 \mu\text{g}/\text{m}^3$ ).

As expected and reported previously [148], cigarette smoke particles were generally larger in size than shisha smoke; the count mean diameter (CMD) of the cigarette smoke particles was 130 nm ( $\sigma_g = 2.0$ ) whereas the size distribution of the particles emitted from shisha was bimodal (one peak below 10 nm and another peak at 65 nm). The mean particle number concentration in the aerosol chamber over the entire measurement period was  $1.5 \times 10^5$  and  $1.9 \times 10^5 \text{ cm}^{-3}$  for the cigarette and waterpipe smoke, respectively.

In a similar study, a shisha-smoking session was carried out in a room (floor area  $20 \text{ m}^2$ , 4 h smoking), resulting a median particle number concentration of  $2.9 \times 10^5 \text{ cm}^{-3}$  (95% values  $5.5 \times 10^5 \text{ cm}^{-3}$ ), which was equivalent to a  $\text{PM}_{2.5}$  of about  $390 \mu\text{g}/\text{m}^3$  (95% value  $740 \mu\text{g}/\text{m}^3$ ) [16].

### 3.3. Particle Emission Rates

The size-specific particle emission rate is shown in Figure 6. On average, the total number of particles emitted while burning two incense sticks at the same time was about  $1.9 \times 10^{10}$  particles/s; this is equivalent to about  $7 \mu\text{g}/\text{s}$ . As for smoking, the total particles emitted were about  $6.8 \times 10^{10}$  and  $1.7 \times 10^{10}$  particles/s, respectively, for cigarettes and shisha. Although the total number emitted during cigarette smoking was more than that during shisha smoking, the total particle mass emitted during shisha smoking (about  $120 \mu\text{g}/\text{s}$ ) was more than that during cigarette smoking (about  $80 \mu\text{g}/\text{s}$ ).



**Figure 6.** Mean emission rates in the form of particle number size distributions during each case of the different scenarios: (a) incense stick burning, (b) cigarette smoking, and (c) shisha smoking. (d–f) The corresponding mean particle mass size distributions.

The literature has limited and few studies reporting the emission factors during incense-burning and cigarette-smoking scenarios. To our knowledge, emission factors were not reported before for shisha. Inside a small chamber ( $1.016 \times 0.660 \times 1.600 \text{ m}^3$ ) used to characterize the physical properties of particles in incense smoke (four incense sticks were burned at each of the four corners of the chamber) the emission rate was  $5.1 \times 10^{12}$ – $1.42 \times 10^{13}$  particles/h [153]. In the European Accredited (EA) Laboratory of Industrial Measurements (LAMI, University of Cassino) the emission rate was  $4.5 \times 10^{14}$ – $1.1 \times 10^{15}$  particles/h ( $\text{PM}_{2.5}$  19.4–50.3 mg/h)

during three incense-burning scenarios [105]. It was previously reported for 23 different types of incense (sticks, cones, powder, smudge bundle, coil, rope, and rocks) that the emissions of particulate matter were with a  $PM_{2.5}$  emission rate of 7–202 mg/h [183]. The UFP emission rate of incense burning carried out in a  $1.2 \times 1.2 \times 1.2 \text{ m}^3$  stainless-steel test chamber was  $0.4 \times 10^{11}$  particles/min [150]. The  $PM_{2.5}$  emission rates were  $1.1 \times 10^2$ – $3.9 \times 10^2$  particles/cm<sup>3</sup>min (mean  $2.5 \times 10^2$  particles/cm<sup>3</sup>min), equivalent to 1.4–3.9 µg/m<sup>3</sup>min (mean 56 µg/m<sup>3</sup>min) [8].

The UFP emission rate of cigarette smouldering carried out in a  $1.2 \times 1.2 \times 1.2 \text{ m}^3$  stainless-steel test chamber was  $3.4 \times 10^{11}$  particles/min [150], with a unimodal particle number size distribution and peak value at around 110 nm. The temporal variation was similar to the one observed in this study, with a maximum concentration of  $1.2 \times 10^6 \text{ cm}^{-3}$  and mean total particle losses of  $3.9 \text{ h}^{-1}$ . The  $PM_{2.5}$  emission rate of 1.2–2 mg/m<sup>3</sup>s was derived from the exposure levels as second-hand smoke (five types of Vogue cigarette) [154]. In another test made inside a small chamber (2.88 m<sup>3</sup>), the  $PM_{2.5}$  emission rate was 580–1250 mg/m<sup>3</sup>s while testing five types of Marlboro cigarette [155].

### 3.4. Regional Inhaled Deposited Dose

In order to put an insight into the exposure scenarios, the regional inhaled deposited dose rates were calculated for an adult male or female (Tables 2 and 3) during two of the main human activities (walking 4 km/h and standing) and being exposed to three scenarios: incense burning, cigarette smoking, and shisha smoking. The metrics were the number concentration of submicron particles ( $PN_{Sub}$ , diameter < 1 µm) and particulate matter ( $PM_{2.5}$ ).

**Table 2.** Regional inhaled deposited dose rate ( $\times 10^9 \text{ \#}/\text{h}$ ) calculated based on submicron particle number ( $PN_{Sub}$ , diameter < 1 µm) concentration for adult males and females during two main activities (walking 4 km/h and standing). Abbreviations H, TB, and Alv are for head and throat airways, tracheobronchial, and pulmonary/alveolar regions, respectively.

		Male				Female			
		H	TB	Alv	Total	H	TB	Alv	Total
Walking	Incense	1000	2500	10,200	13,700	900	2300	8700	12,000
	Cigarette	200	400	1700	2300	200	400	1500	2000
	Shisha	800	1800	7700	10,300	700	1700	6600	9000
Standing	Incense	900	1500	3800	6300	700	1300	2500	4600
	Cigarette	200	300	600	1100	100	200	400	800
	Shisha	700	1100	2900	4700	600	1000	1900	3400

**Table 3.** Regional inhaled deposited dose rate (µg/h) calculated based on submicron particle mass concentration ( $PM_{2.5}$ ) for adult males and females during two main activities (walking 4 km/h and standing). Abbreviations H, TB, and Alv are for head and throat airways, tracheobronchial, and pulmonary/alveolar regions, respectively.

		Male				Female			
		H	TB	Alv	Total	H	TB	Alv	Total
Walking	Incense	190	410	1910	2500	160	370	1600	2130
	Cigarette	40	90	400	520	30	80	340	450
	Shisha	410	930	3920	5270	370	820	3390	4580
Standing	Incense	340	220	820	1380	240	200	520	970
	Cigarette	100	40	170	310	70	40	110	220
	Shisha	1670	350	1750	3760	1150	300	1090	2540

In general, and regardless of the type of combustion scenario and concentration metric, the total inhaled dose rate was higher in males than females. This is basically due to the

fact that male minute ventilation ( $V_E$ ) is higher, and the deposition fraction ( $DF$ ) curves are slightly different for males than females. On average, the regional inhaled dose rates during the walking scenario were about 7%, 18%, and 75%, respectively, for the head and throat airways (H), tracheobronchial (TB), and pulmonary/alveolar (Alv) regions. During standing, the corresponding dose rates were 25%, 20%, and 55%, respectively. The differences in the regional dose rates reflect the changes in the  $DF$  curves between resting and exercising.

During standing, and with respect to the combustion scenario, the total dose rate of submicron particles ( $PN_{Sub}$ , Table 2) was the highest during incense burning ( $6.3 \times 10^{12}$  #/h for males versus  $4.6 \times 10^{12}$  #/h for females) and the lowest during cigarette smoking ( $1.1 \times 10^{12}$  #/h for males versus  $0.8 \times 10^{12}$  #/h for females). Considering the total dose rate of fine particles ( $PM_{2.5}$ , Table 3) during standing, the highest was during shisha smoking, with 3.8 mg/h for males (versus 2.5 mg/h for females), and the lowest was 0.3 mg/h for males (versus 0.2 mg/h for females).

Similarly, the total dose rate during walking (4 km/h) was the highest during incense burning ( $13.7 \times 10^{12}$  #/h for males versus  $12 \times 10^{12}$  #/h for females) and the lowest was during cigarette smoking ( $2.3 \times 10^{12}$  #/h for males versus  $2 \times 10^{12}$  #/h for females). With respect to  $PM_{2.5}$ , the total dose rate was the highest during shisha smoking, with 5.3 mg/h for males (versus 4.6 mg/h for females), and the lowest was 0.5 mg/h for both males and females.

It is interesting to mention here that the higher dose rates calculated during walking when compared to standings are attributed to the fact that the minute ventilation ( $V_E$ ) is higher during exercise (i.e., walking) than during resting (i.e., standing). In addition, the  $DF$  curves are slightly different between the two cases. However, the main effect comes from the minute ventilation ( $V_E$ ).

The calculated dose rates can be converted to actual cumulative dose in the respiratory tracts by multiplying the dose rate by the exposure time (hours). Without performing the calculations, the cumulative dose is considerably high that it can impose serious health risk, especially knowing that the chemical combustion of the smoke consists of high contents of toxic compounds and heavy metals.

#### 4. Conclusions

Tobacco smoking and incense burning are commonly practiced in Jordanian microenvironments. While smoking in Jordan is prohibited inside closed spaces, incense burning remains uncontrolled.

In this study, the particle size distributions (diameter 0.01–25  $\mu\text{m}$ ) were measured inside a room replicating real-life conditions in a Jordanian dwelling. The measurement was made during typical smoking and incense stick-burning scenarios as examples of common indoor aerosol sources and exposure. The results were summarized in terms of number and mass concentrations of submicron and fine particles. The particle losses and emission rates were calculated by utilizing a simple indoor aerosols model.

During cigarette smoking and incense stick-burning scenarios, the particle number concentrations exceeded  $3 \times 10^5 \text{ cm}^{-3}$ . During shisha smoking, they exceeded  $5 \times 10^5 \text{ cm}^{-3}$ . The average emission rates were estimated at  $1.9 \times 10^{10}$ ,  $6.8 \times 10^{10}$  and  $1.7 \times 10^{10}$  particles/s, respectively, for incense, cigarettes, and shisha. That corresponded to about 7, 80, and 120  $\mu\text{g/s}$ , respectively. Interestingly, the exposure levels were almost similar for incense and cigarette-smoking aerosols, but the emission rate from cigarette smoking was about three times that during incense burning. This was expected, because the smoking occurred for a shorter time than the burning of the incense sticks.

In general, and regardless to the type of combustion scenario and concentration metric, males received higher dose rates than females, with average percentiles in the different regions of the respiratory tracts of 7%, 18%, and 75%, respectively, for the head and throat airways (H), tracheobronchial (TB), and pulmonary/alveolar (Alv) regions during walking. During standing, the corresponding dose rates were 25%, 20%, and 55%, respectively. The

total dose rates during standing were in the order of  $10^{12}$  #/h and  $10^3$   $\mu\text{g}/\text{h}$ , respectively, for submicron particle number concentration ( $PN_{Sub}$ ) and  $PM_{2.5}$ . During walking (4 km/h), they were in the order of  $10^{13}$  #/h and  $10^4$   $\mu\text{g}/\text{h}$ , respectively.

The exposure levels, emissions rates, and inhaled deposited dose rates during the studied scenarios of smoking and incense burning are considered seriously high, recalling the fact that aerosols emitted during such scenarios consist of a vast range of polycyclic aromatic hydrocarbons (PAHs), volatile organic compounds (VOCs), and heavy metals, in addition to harmful gases such CO, CO<sub>2</sub>, NO<sub>x</sub>, and SO<sub>2</sub>.

By all means, the Jordanian authorities have to take immediate actions to enforce the prohibition of cigarette smoking indoors and develop regulations for incense stick burning.

**Funding:** This research was financially supported by generous funding from the Deanship of Scientific Research at the University of Jordan (grant# 2379) and the Scientific Research Support Fund at the Jordanian Ministry of Higher Education (project# WE-2-2-2017).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Data are available upon request.

**Acknowledgments:** The author acknowledges the support of the Deanship of Scientific Research (DSR) at the University of Jordan. The author also acknowledges the support of the Scientific Research Support Fund at the Jordanian Ministry of Higher Education; the Atmosphere and Climate Competence Center (ACCC) Flagship funded by the Academy of Finland (grant# 337549); and the Eastern Mediterranean and Middle East Climate and Atmosphere Research (EMME-CARE) project, which received funding from the European Union's Horizon 2020 Research and Innovation Programme (Grant Agreement Number 856612) and the Government of Cyprus. The sole responsibility of this publication lies with the authors. Open-access funding was provided by the University of Helsinki.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Cobb, C.O.; Shihadeh, A.; Weaver, M.F.; Eissenberg, T. Waterpipe Tobacco Smoking and Cigarette Smoking: A Direct Comparison of Toxicant Exposure and Subjective Effects. *Nicotine Tob. Res.* **2011**, *13*, 78–87. [[CrossRef](#)] [[PubMed](#)]
2. Prignot, J. Risks and Subjective Effects of Waterpipe versus Cigarette Smoking. *Breathe* **2011**, *7*, 371. [[CrossRef](#)]
3. Goel, A.; Wathore, R.; Chakraborty, T.; Agrawal, M. Characteristics of Exposure to Particles Due to Incense Burning inside Temples in Kanpur, India. *Aerosol Air Qual. Res.* **2017**, *17*, 608–615. [[CrossRef](#)]
4. Fang, G.-C.; Chu, C.-C.; Wu, Y.-S.; Fu, P.P.-C. Emission Characters of Particulate Concentrations and Dry Deposition Studies for Incense Burning at a Taiwanese Temple. *Toxicol. Ind. Health* **2002**, *18*, 183–190. [[CrossRef](#)]
5. Chiang, K.-C.; Chio, C.-P.; Chiang, Y.-H.; Liao, C.-M. Assessing Hazardous Risks of Human Exposure to Temple Airborne Polycyclic Aromatic Hydrocarbons. *J. Hazard. Mater.* **2009**, *166*, 676–685. [[CrossRef](#)]
6. Hu, M.-T.; Chen, S.-J.; Huang, K.-L.; Lin, Y.-C.; Lee, W.-J.; Chang-Chien, G.-P.; Tsai, J.-H.; Lee, J.-T.; Chiu, C.-H. Characterization of, and Health Risks from, Polychlorinated Dibenzo-p-Dioxins/Dibenzofurans from Incense Burned in a Temple. *Sci. Total. Environ.* **2009**, *407*, 4870–4875. [[CrossRef](#)]
7. Bhadauria, V.; Parmar, D.; Ganguly, R.; Rathi, A.K.; Kumar, P. Exposure Assessment of PM<sub>2.5</sub> in Temple Premises and Crematoriums in Kanpur, India. *Environ. Sci. Pollut. Res.* **2022**, *29*, 38374–38384. [[CrossRef](#)]
8. Polednik, B. Particle Exposure in a Baroque Church during Sunday Masses. *Environ. Res.* **2013**, *126*, 215–220. [[CrossRef](#)]
9. Maziak, W. The Global Epidemic of Waterpipe Smoking. *Addict. Behav.* **2011**, *36*, 1–5. [[CrossRef](#)]
10. Maziak, W. The Waterpipe: An Emerging Global Risk for Cancer. *Cancer Epidemiol.* **2013**, *37*, 1–4. [[CrossRef](#)]
11. Azab, M.; Khabour, O.F.; Alkaraki, A.K.; Eissenberg, T.; Alzoubi, K.H.; Primack, B.A. Water Pipe Tobacco Smoking among University Students in Jordan. *Nicotine Tob. Res.* **2010**, *12*, 606–612. [[CrossRef](#)] [[PubMed](#)]
12. Warren, C.W.; Lea, V.; Lee, J.; Jones, N.R.; Asma, S.; McKenna, M. Change in Tobacco Use among 13-15 Year Olds between 1999 and 2008: Findings from the Global Youth Tobacco Survey. *Glob. Health Promot.* **2009**, *16* (Suppl. S2), 38–90. [[CrossRef](#)] [[PubMed](#)]
13. El-Awa, F.; Warren, C.W.; Jones, N.R. Changes in Tobacco Use among 13-15-Year-Olds between 1999 and 2007: Findings from the Eastern Mediterranean Region. *East. Mediterr. Health J.* **2010**, *16*, 266–273. [[CrossRef](#)] [[PubMed](#)]
14. Erguder, T.; Cakir, B.; Babalioglu, N.; Dogusan, H.; Turkoral, E.; Warren, C.W. Tobacco Use among Institutionalized Adolescents in Turkey: Does Social Environment Affect the Risk? *Int. J. Public Health* **2009**, *54*, 379–389. [[CrossRef](#)]
15. Warren, C.W.; Erguder, T.; Lee, J.; Lea, V.; Sauer, A.G.; Jones, N.R.; Bilir, N. Effect of Policy Changes on Cigarette Sales: The Case of Turkey. *Eur. J. Public Health* **2012**, *22*, 712–716. [[CrossRef](#)]

16. Koh, H.K.; Alpert, H.R.; Judge, C.M.; Caughey, R.W.; Elqura, L.J.; Connolly, G.N.; Warren, C.W. Understanding Worldwide Youth Attitudes towards Smoke-Free Policies: An Analysis of the Global Youth Tobacco Survey. *Tob. Control.* **2011**, *20*, 219–225. [[CrossRef](#)]
17. Barbouni, A.; Hadjichristodoulou, C.; Merakou, K.; Antoniadou, E.; Kourea, K.; Miloni, E.; Warren, C.W.; Rahiotis, G.; Kremastinou, J. Tobacco Use, Exposure to Secondhand Smoke, and Cessation Counseling Among Health Professions Students: Greek Data from the Global Health Professions Student Survey (GHPSS). *Int. J. Environ. Res. Public Health* **2012**, *9*, 331–342. [[CrossRef](#)]
18. Warren, C.W.; Jones, N.R.; Peruga, A.; Chauvin, J.; Baptiste, J.P.; Costa de Silva, V.; el Awa, F.; Tsouros, A.; Rahman, K.; Fishburn, B.; et al. Global Youth Tobacco Surveillance, 2000–2007. *MMWR. Surveill. Summ. Morb. Mortal. Wkly. Rep. Surveill. Summ./CDC* **2008**, *57*, 1–28.
19. Basir, F.; Khan, M.S.; Ahmed, B.; Farooq, W.; Virji, R.N. The Frequency of Shisha (Waterpipe) Smoking in Students of Different Age Groups. *J. Coll. Physicians Surg. Pak.* **2014**, *24*, 265–268.
20. Jawaid, A.; Zafar, A.M.; Rehman, T.-U.; Nazir, M.R.; Ghafoor, Z.A.; Afzal, O.; Khan, J.A. Knowledge, Attitudes and Practice of University Students Regarding Waterpipe Smoking in Pakistan. *Int. J. Tuberc. Lung Dis.* **2008**, *12*, 1077–1084.
21. Anjum, Q.; Ahmed, F.; Ashfaq, T. Knowledge, Attitude and Perception of Water Pipe Smoking (Shisha) among Adolescents Aged 14–19 Years. *J. Pak. Med. Assoc.* **2008**, *58*, 312–317. [[PubMed](#)]
22. Hussein, T.; Alameer, A.; Jaghbeir, O.; Albeitshaweesh, K.; Malkawi, M.; Boor, B.E.; Koivisto, A.J.; Löndahl, J.; Alrifai, O.; Al-Hunaiti, A. Indoor Particle Concentrations, Size Distributions, and Exposures in Middle Eastern Microenvironments. *Atmosphere* **2020**, *11*, 41. [[CrossRef](#)]
23. Hussein, T. Indoor-to-Outdoor Relationship of Aerosol Particles inside a Naturally Ventilated Apartment—A Comparison between Single-Parameter Analysis and Indoor Aerosol Model Simulation. *Sci. Total Environ.* **2017**, *596–597*, 321–330. [[CrossRef](#)] [[PubMed](#)]
24. Yeatts, K.B.; El-Sadig, M.; Leith, D.; Kalsbeek, W.; Al-Maskari, F.; Couper, D.; Funk, W.E.; Zoubeidi, T.; Chan, R.L.; Trent, C.; et al. Indoor Air Pollutants and Health in the United Arab Emirates. *Environ. Health Perspect.* **2012**, *120*, 687–694. [[CrossRef](#)]
25. Blank, M.D.; Brown, K.W.; Goodman, R.J.; Eissenberg, T. An Observational Study of Group Waterpipe Use in a Natural Environment. *Nicotine Tob. Res.* **2014**, *16*, 93–99. [[CrossRef](#)]
26. Salloum, R.G.; Thrasher, J.F.; Kates, F.R.; Maziak, W. Water Pipe Tobacco Smoking in the United States: Findings from the National Adult Tobacco Survey. *Prev. Med.* **2015**, *71*, 88–93. [[CrossRef](#)]
27. Salloum, R.G.; Nakkash, R.; Abu-Rmeileh, N.M.E.; Hamadeh, R.R.; Darawad, M.W.; Kheirallah, K.A.; Al-Farsi, Y.; Yusufali, A.; Thomas, J.; Mostafa, A.; et al. Individual-Level Determinants of Waterpipe Smoking Demand in Four Eastern-Mediterranean Countries. *Health Promot. Int.* **2019**, *34*, 1157–1166. [[CrossRef](#)]
28. Amoatey, P.; Omidvarborna, H.; Baawain, M.S.; Al-Mamun, A. Indoor Air Pollution and Exposure Assessment of the Gulf Cooperation Council Countries: A Critical Review. *Environ. Int.* **2018**, *121*, 491–506. [[CrossRef](#)]
29. el Amin, S.E.T. School Smoking Policies and Health Science Students' Use of Cigarettes, Shisha, and Dipping Tombak in Sudan. *Front. Public Health* **2019**, *7*, 290. [[CrossRef](#)]
30. Hensel, E.C.; Sarles, S.E.; Al-Olayan, A.; Difrancesco, A.G.; Jayasekera, S.; Eddingsaas, N.C.; Robinson, R.J. A Proposed Waterpipe Emissions Topography Protocol Reflecting Natural Environment User Behaviour. *Int. J. Environ. Res. Public Health* **2020**, *17*, 92. [[CrossRef](#)]
31. Alali, W.Q.; Longenecker, J.C.; Alwotyan, R.; AlKandari, H.; Al-Mulla, F.; al Duwairi, Q. Prevalence of Smoking in the Kuwaiti Adult Population in 2014: A Cross-Sectional Study. *Environ. Sci. Pollut. Res.* **2021**, *28*, 10053–10067. [[CrossRef](#)] [[PubMed](#)]
32. Erythropel, H.C.; Garcia Torres, D.S.; Woodrow, J.G.; de Winter, T.M.; Falinski, M.M.; Anastas, P.T.; O'Malley, S.S.; Krishnan-Sarin, S.; Zimmerman, J.B. Quantification of Flavorants and Nicotine in Waterpipe Tobacco and Mainstream Smoke and Comparison to E-Cigarette Aerosol. *Nicotine Tob. Res.* **2021**, *23*, 600–604. [[CrossRef](#)] [[PubMed](#)]
33. Hirpa, S.; Fogarty, A.; Addissie, A.; Bauld, L.; Frese, T.; Unverzagt, S.; Kantelhardt, E.J.; Getachew, S.; Deressa, W. An Emerging Problem of Shisha Smoking among High School Students in Ethiopia. *Int. J. Environ. Res. Public Health* **2021**, *18*, 7023. [[CrossRef](#)] [[PubMed](#)]
34. Semple, S.; Dobson, R.; O'Donnell, R.; Zainal Abidin, E.; Tigova, O.; Okello, G.; Fernández, E. Smoke-Free Spaces: A Decade of Progress, a Need for More? *Tob. Control.* **2022**, *31*, 250–256. [[CrossRef](#)]
35. Azagba, S.; Latham, K.; Shan, L. Waterpipe Tobacco Smoking Trends among Middle and High School Students in the United States from 2011 to 2017. *Drug Alcohol Depend.* **2019**, *200*, 19–25. [[CrossRef](#)]
36. Shihadeh, A.; Azar, S.; Antonios, C.; Haddad, A. Towards a Topographical Model of Narghile Water-Pipe Café Smoking: A Pilot Study in a High Socioeconomic Status Neighborhood of Beirut, Lebanon. *Pharm. Biochem. Behav.* **2004**, *79*, 75–82. [[CrossRef](#)]
37. Dalibalta, S.; Elsayed, Y.; Alqtaishat, F.; Gomes, L.; Fernandes, N. A Health Risk Assessment of Arabian Incense (Bakhour) Smoke in the United Arab Emirates. *Sci. Total Environ.* **2015**, *511*, 684–691. [[CrossRef](#)]
38. King, B.A.; Dube, S.R.; Tynan, M.A. Current Tobacco Use among Adults in the United States: Findings from the National Adult Tobacco Survey. *Am. J. Public Health* **2012**, *102*, e93–e100. [[CrossRef](#)]
39. Majeed, B.A.; Sterling, K.L.; Weaver, S.R.; Pechacek, T.F.; Eriksen, M.P. Prevalence and Harm Perceptions of Hookah Smoking among U.S. Adults, 2014–2015. *Addict. Behav.* **2017**, *69*, 78–86. [[CrossRef](#)]

40. Robinson, J.N.; Wang, B.; Jackson, K.J.; Donaldson, E.A.; Ryant, C.A. Characteristics of Hookah Tobacco Smoking Sessions and Correlates of Use Frequency among US Adults: Findings from Wave 1 of the Population Assessment of Tobacco and Health (PATH) Study. *Nicotine Tob. Res.* **2018**, *20*, 731–740. [[CrossRef](#)]
41. Rezk-Hanna, M.; Macabasco-O’Connell, A.; Woo, M. Hookah Smoking among Young Adults in Southern California. *Nurs. Res.* **2014**, *63*, 300–306. [[CrossRef](#)] [[PubMed](#)]
42. Hensel, E.C.; Eddingsaas, N.C.; Saleh, Q.M.; Jayasekera, S.; Sarles, S.E.; Difrancesco, A.G.; Robinson, R.J. Proposed Standard Test Protocols and Outcome Measures for Quantitative Comparison of Emissions from Electronic Nicotine Delivery Systems. *Int. J. Environ. Res. Public Health* **2022**, *19*, 2144. [[CrossRef](#)] [[PubMed](#)]
43. Gordon, T.; Karey, E.; Rebuli, M.E.; Escobar, Y.-N.H.; Jaspers, I.; Chen, L.C. E-Cigarette Toxicology. *Annu. Rev. Pharm. Toxicol.* **2022**, *62*, 301–322. [[CrossRef](#)]
44. Stone, M.D.; DeAtley, T.; Pianin, S.; Strasser, A.A.; Audrain-McGovern, J. Switching from Cigarettes to IQOS: A Pilot Examination of IQOS-Associated Reward, Reinforcement, and Abstinence Relief. *Drug Alcohol Depend.* **2022**, *238*, 109569. [[CrossRef](#)] [[PubMed](#)]
45. Lempert, L.K.; Bialous, S.; Glantz, S. FDA’s Reduced Exposure Marketing Order for IQOS: Why It Is Not a Reliable Global Model. *Tob. Control.* **2022**, *31*, E83–E87. [[CrossRef](#)]
46. Uguna, C.N.; Snape, C.E. Should IQOS Emissions Be Considered as Smoke and Harmful to Health? A Review of the Chemical Evidence. *ACS. Omega* **2022**, *7*, 22111–22124. [[CrossRef](#)]
47. Orzabal, M.R.; Naik, V.D.; Lee, J.; Hillhouse, A.E.; Brashear, W.A.; Threadgill, D.W.; Ramadoss, J. Impact of E-Cig Aerosol Vaping on Fetal and Neonatal Respiratory Development and Function. *Transl. Res.* **2022**, *246*, 102–114. [[CrossRef](#)]
48. Onoue, A.; Inaba, Y.; Machida, K.; Samukawa, T.; Inoue, H.; Kurosawa, H.; Ogata, H.; Kunugita, N.; Omori, H. Association between Fathers’ Use of Heated Tobacco Products and Urinary Cotinine Concentrations in Their Spouses and Children. *Int. J. Environ. Res. Public Health* **2022**, *19*, 6275. [[CrossRef](#)]
49. Farrell, K.R.; Weitzman, M.; Karey, E.; Lai, T.K.Y.; Gordon, T.; Xu, S. Passive Exposure to E-Cigarette Emissions Is Associated with Worsened Mental Health. *BMC Public Health* **2022**, *22*, 1138. [[CrossRef](#)]
50. Çetintaş, E.; Luo, Y.; Nguyen, C.; Guo, Y.; Li, L.; Zhu, Y.; Ozcan, A. Characterization of Exhaled E-Cigarette Aerosols in a Vape Shop Using a Field-Portable Holographic on-Chip Microscope. *Sci. Rep.* **2022**, *12*, 3175. [[CrossRef](#)]
51. Vivarelli, F.; Granata, S.; Rullo, L.; Mussoni, M.; Candeletti, S.; Romualdi, P.; Fimognari, C.; Cruz-Chamorro, I.; Carrillo-Vico, A.; Paolini, M.; et al. On the Toxicity of E-Cigarettes Consumption: Focus on Pathological Cellular Mechanisms. *Pharm. Res.* **2022**, *182*, 106315. [[CrossRef](#)] [[PubMed](#)]
52. Talih, S.; Balhas, Z.; Eissenberg, T.; Salman, R.; Karaoghlanian, N.; Hellani, A.E.; Baalbaki, R.; Saliba, N.; Shihadeh, A. Effects of User Puff Topography, Device Voltage, and Liquid Nicotine Concentration on Electronic Cigarette Nicotine Yield: Measurements and Model Predictions. *Nicotine Tob. Res.* **2015**, *17*, 150–157. [[CrossRef](#)]
53. Amorós-Pérez, A.; Cano-Casanova, L.; Román-Martínez, M.D.C.; Lillo-Ródenas, M.Á. Comparison of Particulate Matter Emission and Soluble Matter Collected from Combustion Cigarettes and Heated Tobacco Products Using a Setup Designed to Simulate Puffing Regimes. *Chem. Eng. J. Adv.* **2021**, *8*, 100144. [[CrossRef](#)]
54. Nitta, N.A.; Sato, T.; Komura, M.; Yoshikawa, H.; Suzuki, Y.; Mitsui, A.; Kuwasaki, E.; Takahashi, F.; Kodama, Y.; Seyama, K.; et al. Exposure to the Heated Tobacco Product IQOS Generates Apoptosis-Mediated Pulmonary Emphysema in Murine Lungs. *Am. J. Physiol. Lung Cell Mol. Physiol.* **2022**, *322*, L699–L711. [[CrossRef](#)] [[PubMed](#)]
55. Wiens, T.; Taylor, J.; Cole, C.; Saravia, S.; Peterson, J.; Lunda, M.; Margetta, J.; D’Heilly, P.; Holzbauer, S.; Lynfield, R. Lessons Learned from the E-Cigarette, or Vaping, Product Use–Associated Lung Injury (EVALI) Outbreak Response, Minnesota, 2019–2020. *Public Health Rep.* **2022**, *137*, 1053–1060. [[CrossRef](#)]
56. Raja, A.; Zelikoff, J.T.; Jaimes, E.A. A Contemporary Review of Nephrotoxicity and E-Cigarette Use. *Curr. Opin. Toxicol.* **2022**, *31*, 100361. [[CrossRef](#)]
57. McKenzie, C.R.L.; Davis, J.; Dunlop, A.J. E-Cigarette or Vaping Product Use–Associated Lung Injury in an Adolescent. *Med. J. Aust.* **2022**, *216*, 374. [[CrossRef](#)]
58. Williams, M.A.; Reddy, G.; Quinn, M.J.; Millikan Bell, A. Toxicological Assessment of Electronic Cigarette Vaping: An Emerging Threat to Force Health, Readiness and Resilience in the U.S. Army. *Drug Chem. Toxicol.* **2022**, *45*, 2049–2085. [[CrossRef](#)]
59. Chan, B.S.; Kiss, A.; McIntosh, N.; Sheppard, V.; Dawson, A.H. E-Cigarette or Vaping Product Use–Associated Lung Injury in an Adolescent. *Med. J. Aust.* **2021**, *215*, 313–314.e1. [[CrossRef](#)]
60. Nafees, A.A.; Taj, T.; Kadir, M.M.; Fatmi, Z.; Lee, K.; Sathiakumar, N. Indoor Air Pollution (PM<sub>2.5</sub>) Due to Secondhand Smoke in Selected Hospitality and Entertainment Venues of Karachi, Pakistan. *Tob. Control.* **2012**, *21*, 460–464. [[CrossRef](#)]
61. Russo, E.T.; Hulse, T.E.; Adamkiewicz, G.; Levy, D.E.; Bethune, L.; Kane, J.; Reid, M.; Shah, S.N. Comparison of Indoor Air Quality in Smoke-Permitted and Smoke-Free Multiunit Housing: Findings from the Boston Housing Authority. *Nicotine Tob. Res.* **2015**, *17*, 316–322. [[CrossRef](#)] [[PubMed](#)]
62. Muñoz, C.; Droppelmann, A.; Erazo, M.; Aceituno, P.; Orellana, C.; Parro, J.; Mesias, S.; Marchetti, N.; Navas-Acien, A.; Iglesias, V. Occupational Exposure to Polycyclic Aromatic Hydrocarbons: A Cross-Sectional Study in Bars and Restaurants in Santiago, Chile. *Am. J. Ind. Med.* **2016**, *59*, 887–896. [[CrossRef](#)]
63. Dobson, R.; Demou, E.; Semple, S. Occupational Exposure to Second-Hand Tobacco Smoke: Development of a Job Exposure Matrix. *Ann. Work Expo. Health* **2021**, *65*, 1133–1138. [[CrossRef](#)] [[PubMed](#)]

64. Puvanesarajah, S.; Tsai, J.; Alexander, D.S.; Tynan, M.A.; Gentzke, A.S. Youth Indoor and Outdoor Exposure to Secondhand Smoke and Secondhand Aerosol. *Am. J. Prev. Med.* **2022**, *62*, 903–913. [[CrossRef](#)]
65. Veerabhadra, S.K.; Yadav, S.; Ramachandra, S.S.; Dicksit, D.D.; Muttalib, K.B.A.; Zamzuri, A.T.B. Secondhand Smoke Exposure from the Indoor and Outdoor Shisha Centers Located at the Perimeter of Educational Institutions in Malaysia: A Cross-Sectional Study. *J. Public Health Policy* **2022**, *43*, 77–88. [[CrossRef](#)] [[PubMed](#)]
66. Sureda, X.; Bilal, U.; Fernández, E.; Valiente, R.; Escobar, F.J.; Navas-Acien, A.; Franco, M. Second-Hand Smoke Exposure in Outdoor Hospitality Venues: Smoking Visibility and Assessment of Airborne Markers. *Environ. Res.* **2018**, *165*, 220–227. [[CrossRef](#)]
67. Nandasena, S.; Wickremasinghe, A.R.; Lee, K.; Sathiakumar, N. Indoor Fine Particle (PM<sub>2.5</sub>) Pollution Exposure Due to Secondhand Smoke in Selected Public Places of Sri Lanka. *Am. J. Ind. Med.* **2012**, *55*, 1129–1136. [[CrossRef](#)]
68. Kuo, H.-W.; Rees, V.W. Third-Hand Smoke (THS): What Is It and What Should We Do about It? *J. Formos. Med. Assoc.* **2019**, *118*, 1478–1479. [[CrossRef](#)]
69. Lidón-Moyano, C.; Fu, M.; Perez-Ortuño, R.; Ballbè, M.; Garcia, E.; Martín-Sánchez, J.C.; Pascual, J.A.; Fernández, E.; Martínez-Sánchez, J.M. Toward a Correct Measure of Third-Hand Exposure. *Environ. Res.* **2021**, *194*, 110686. [[CrossRef](#)]
70. Wu, J.-X.; Lau, A.T.Y.; Xu, Y.-M. Indoor Secondary Pollutants Cannot Be Ignored: Third-Hand Smoke. *Toxics* **2022**, *10*, 363. [[CrossRef](#)]
71. Lidón-Moyano, C.; Fu, M.; Pérez-Ortuño, R.; Ballbè, M.; Garcia, E.; Martín-Sánchez, J.C.; Pascual, J.A.; Fernández, E.; Martínez-Sánchez, J.M. Third-Hand Exposure at Homes: Assessment Using Salivary Cotinine. *Environ. Res.* **2021**, *196*, 110393. [[CrossRef](#)]
72. Özpınar, S.; Demir, Y.; Yazıcıoğlu, B.; Bayçelebi, S. Pregnant Women’s Beliefs about Third-Hand Smoke and Exposure to Tobacco Smoke. *Cent. Eur. J. Public Health* **2022**, *30*, 154–159. [[CrossRef](#)] [[PubMed](#)]
73. Borujeni, E.T.; Yaghmaian, K.; Naddafi, K.; Hassanvand, M.S.; Naderi, M. Identification and Determination of the Volatile Organics of Third-Hand Smoke from Different Cigarettes and Clothing Fabrics. *J. Environ. Health Sci. Eng.* **2022**, *20*, 53–63. [[CrossRef](#)] [[PubMed](#)]
74. Yang, J.; Hashemi, S.; Han, W.; Song, Y.; Lim, Y. Exposure and Risk Assessment of Second- and Third-Hand Tobacco Smoke Using Urinary Cotinine Levels in South Korea. *Int. J. Environ. Res. Public Health* **2022**, *19*, 3746. [[CrossRef](#)]
75. Mahabee-Gittens, E.M.; Merianos, A.L.; Matt, G.E. Comment Regarding Categorization of Third-Hand Smoke Exposure in “Third-Hand Exposure at Homes: Assessment Using Salivary Cotinine”. *Environ. Res.* **2021**, *195*, 110595. [[CrossRef](#)] [[PubMed](#)]
76. Moon, S.Y.; Kim, T.; Kim, Y.-J.; Kim, Y.; Kim, S.Y.; Kang, D. Public Facility Utility and Third-Hand Smoking Exposure without First and Second-Hand Smoking According to Urinary Cotinine Level. *Int. J. Environ. Res. Public Health* **2019**, *16*, 855. [[CrossRef](#)]
77. Jiang, W.; Wu, H.; Yu, X.; Wang, Y.; Gu, W.; Wei, W.; Li, B.; Jiang, X.; Wang, Y.; Hou, W.; et al. Third-Hand Smoke Exposure Is Associated with Abnormal Serum Melatonin Level via Hypomethylation of CYP1A2 Promoter: Evidence from Human and Animal Studies. *Environ. Pollut.* **2021**, *277*, 116669. [[CrossRef](#)]
78. Lai, F.Y.; Lypousi, K.; Been, F.; Benaglia, L.; Udrișard, R.; Delémont, O.; Esseiva, P.; Thomaidis, N.S.; Covaci, A.; van Nuijs, A.L.N. Levels of 4-(Methylnitrosamino)-1-(3-Pyridyl)-1-Butanone (NNK) in Raw Wastewater as an Innovative Perspective for Investigating Population-Wide Exposure to Third-Hand Smoke. *Sci. Rep.* **2018**, *8*, 13254. [[CrossRef](#)]
79. de Granda-Orive, J.I.; Solano-Reina, S.; Jiménez-Ruiz, C.A. Is Smoking Outside an Enclosed Space Enough to Prevent Second and Third-Hand Exposure? ¿Salir a Fumar Fuera de Un Ambiente Cerrado Es Suficiente Para Evitar El Tabaquismo de Segunda y Tercera Mano? *Arch. Bronconeumol.* **2021**, *57*, 83–84. [[CrossRef](#)]
80. Shihadeh, A. Investigation of Mainstream Smoke Aerosol of the Argileh Water Pipe. *Food Chem. Toxicol.* **2003**, *41*, 143–152. [[CrossRef](#)]
81. Shihadeh, A.; Saleh, R. Polycyclic Aromatic Hydrocarbons, Carbon Monoxide, “Tar”, and Nicotine in the Mainstream Smoke Aerosol of the Narghile Water Pipe. *Food Chem. Toxicol.* **2005**, *43*, 655–661. [[CrossRef](#)] [[PubMed](#)]
82. al Rashidi, M.; Shihadeh, A.; Saliba, N.A. Volatile Aldehydes in the Mainstream Smoke of the Narghile Waterpipe. *Food Chem. Toxicol.* **2008**, *46*, 3546–3549. [[CrossRef](#)] [[PubMed](#)]
83. Sepetdjian, E.; Shihadeh, A.; Saliba, N.A. Measurement of 16 Polycyclic Aromatic Hydrocarbons in Narghile Waterpipe Tobacco Smoke. *Food Chem. Toxicol.* **2008**, *46*, 1582–1590. [[CrossRef](#)] [[PubMed](#)]
84. Eissenberg, T.; Shihadeh, A. Waterpipe Tobacco and Cigarette Smoking. Direct Comparison of Toxicant Exposure. *Am. J. Prev. Med.* **2009**, *37*, 518–523. [[CrossRef](#)]
85. Daher, N.; Saleh, R.; Jaroudi, E.; Sheheitli, H.; Badr, T.; Sepetdjian, E.; al Rashidi, M.; Saliba, N.; Shihadeh, A. Comparison of Carcinogen, Carbon Monoxide, and Ultrafine Particle Emissions from Narghile Waterpipe and Cigarette Smoking: Sidestream Smoke Measurements and Assessment of Second-Hand Smoke Emission Factors. *Atmos. Environ.* **2010**, *44*, 8–14. [[CrossRef](#)]
86. Kumar, S.R.; Davies, S.; Weitzman, M.; Sherman, S. A Review of Air Quality, Biological Indicators and Health Effects of Second-Hand Waterpipe Smoke Exposure. *Tob. Control.* **2015**, *24*, i54–i59. [[CrossRef](#)]
87. Kim, K.-H.; Kabir, E.; Jahan, S.A. Waterpipe Tobacco Smoking and Its Human Health Impacts. *J. Hazard Mater.* **2016**, *317*, 229–236. [[CrossRef](#)]
88. Wang, H.; Li, X.; Guo, J.; Peng, B.; Cui, H.; Liu, K.; Wang, S.; Qin, Y.; Sun, P.; Zhao, L.; et al. Distribution of Toxic Chemicals in Particles of Various Sizes from Mainstream Cigarette Smoke. *Inhal. Toxicol.* **2016**, *28*, 89–94. [[CrossRef](#)]
89. Phares, D.J.; Collier, S.; Zheng, Z.; Jung, H.S. In-Situ Analysis of the Gas- and Particle-Phase in Cigarette Smoke by Chemical Ionization TOF-MS. *J. Aerosol Sci.* **2017**, *106*, 132–141. [[CrossRef](#)]

90. Feliu, A.; Fu, M.; Russo, M.; Martinez, C.; Sureda, X.; López, M.J.; Cortés, N.; Fernández, E. Exposure to Second-Hand Tobacco Smoke in Waterpipe Cafés in Barcelona, Spain: An Assessment of Airborne Nicotine and PM<sub>2.5</sub>. *Environ. Res.* **2020**, *184*, 109347. [[CrossRef](#)]
91. Haiduc, A.; Zanetti, F.; Zhao, X.; Schlage, W.K.; Scherer, M.; Pluym, N.; Schlenger, P.; Ivanov, N.V.; Majeed, S.; Hoeng, J.; et al. Analysis of Chemical Deposits on Tooth Enamel Exposed to Total Particulate Matter from Cigarette Smoke and Tobacco Heating System 2.2 Aerosol by Novel GC–MS Deconvolution Procedures. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* **2020**, *1152*, 122228. [[CrossRef](#)] [[PubMed](#)]
92. Jaccard, G.; Tabin Djoko, D.; Korneliou, A.; Belushkin, M. Analysis of Waterpipe Aerosol Constituents in Accordance with the ISO Standard 22486. *Toxicol. Rep.* **2020**, *7*, 1344–1349. [[CrossRef](#)] [[PubMed](#)]
93. Martinasek, M.P.; Calvanese, A.V.; Lipski, B.K. A Naturalistic Study of Carbon Monoxide, Heart Rate, Oxygen Saturation, and Perfusion Index in Hookah Lounge Patrons. *Respir. Care* **2021**, *66*, 269–274. [[CrossRef](#)] [[PubMed](#)]
94. Niu, X.; Jones, T.; Bérubé, K.; Chuang, H.-C.; Sun, J.; Ho, K.F. The Oxidative Capacity of Indoor Source Combustion Derived Particulate Matter and Resulting Respiratory Toxicity. *Sci. Total. Environ.* **2021**, *767*, 144391. [[CrossRef](#)] [[PubMed](#)]
95. Zhang, S.; Wang, Z.; Zhang, J.; Guo, D.; Chen, Y. Inhalable Cigarette-Burning Particles: Size-Resolved Chemical Composition and Mixing State. *Environ. Res.* **2021**, *202*, 111790. [[CrossRef](#)] [[PubMed](#)]
96. Amorós-Pérez, A.; Cano-Casanova, L.; Román-Martínez, M.D.C.; Lillo-Ródenas, M.Á. Solid Matter and Soluble Compounds Collected from Cigarette Smoke and Heated Tobacco Product Aerosol Using a Laboratory Designed Puffing Setup. *Environ. Res.* **2022**, *206*, 112619. [[CrossRef](#)] [[PubMed](#)]
97. Holt, J.C.M.; Mayer-Helm, B.; Gafner, J.; Zierlinger, M.; Hirn, C.; Paschke, T.; Eilenberger, G.; Kuba, M.; Pummer, S.; Charriere, M. Investigating the Transfer Rate of Waterpipe Additives to Smoke as an Integral Part of Toxicological Risk Assessments. *Toxicol. Rep.* **2022**, *9*, 945–950. [[CrossRef](#)]
98. Jung, C.-C.; Syu, Z.-H.; Su, H.-J.; Lian, P.-Y.; Chen, N.-T. Stable C and N Isotopes of PM<sub>2.5</sub> and Size-Segregated Particles Emitted from Incense Stick and Cigarette Burning. *Environ. Res.* **2022**, *212*, 113346. [[CrossRef](#)]
99. Fromme, H.; Dietrich, S.; Heitmann, D.; Dressel, H.; Diemer, J.; Schulz, T.; Jörres, R.A.; Berlin, K.; Völkel, W. Indoor Air Contamination during a Waterpipe (Narghile) Smoking Session. *Food Chem. Toxicol.* **2009**, *47*, 1636–1641. [[CrossRef](#)]
100. Schubert, J.; Müller, F.D.; Schmidt, R.; Luch, A.; Schulz, T.G. Waterpipe Smoke: Source of Toxic and Carcinogenic VOCs, Phenols and Heavy Metals? *Arch. Toxicol.* **2015**, *89*, 2129–2139. [[CrossRef](#)]
101. Bernd, K.; DeGroot, D.; Stadler, H.; Coats, S.; Carmack, D.; Mailig, R.; Lidsky, S.; Hauser, C. Contributions of Charcoal, Tobacco, and Syrup to the Toxicity and Particle Distribution of Waterpipe Tobacco Smoke. *Toxicol. Lett.* **2019**, *313*, 60–65. [[CrossRef](#)] [[PubMed](#)]
102. Ho, S.S.H.; Yu, J.Z. Concentrations of Formaldehyde and Other Carbonyls in Environments Affected by Incense Burning. *J. Environ. Monit.* **2002**, *4*, 728–733. [[CrossRef](#)] [[PubMed](#)]
103. Chuang, H.-C.; Jones, T.; Chen, Y.; Bell, J.; Wenger, J.; Bérubé, K. Characterisation of Airborne Particles and Associated Organic Components Produced from Incense Burning. *Anal. Bioanal. Chem.* **2011**, *401*, 3095–3102. [[CrossRef](#)] [[PubMed](#)]
104. See, S.W.; Balasubramanian, R. Characterization of Fine Particle Emissions from Incense Burning. *Build. Environ.* **2011**, *46*, 1074–1080. [[CrossRef](#)]
105. Stabile, L.; Fuoco, F.C.; Buonanno, G. Characteristics of Particles and Black Carbon Emitted by Combustion of Incenses, Candles and Anti-Mosquito Products. *Build. Environ.* **2012**, *56*, 184–191. [[CrossRef](#)]
106. Manoukian, A.; Quivet, E.; Temime-Roussel, B.; Nicolas, M.; Maupetit, F.; Wortham, H. Emission Characteristics of Air Pollutants from Incense and Candle Burning in Indoor Atmospheres. *Environ. Sci. Pollut. Res.* **2013**, *20*, 4659–4670. [[CrossRef](#)] [[PubMed](#)]
107. Hwang, Y.-H.; Lin, Y.-S.; Lin, C.-Y.; Wang, I.-J. Incense Burning at Home and the Blood Lead Level of Preschoolers in Taiwan. *Environ. Sci. Pollut. Res.* **2014**, *21*, 13480–13487. [[CrossRef](#)]
108. Višić, B.; Kranjc, E.; Pirker, L.; Bačnik, U.; Tavčar, G.; Škapin, S.; Remškar, M. Incense Powder and Particle Emission Characteristics during and after Burning Incense in an Unventilated Room Setting. *Air Qual. Atmos. Health* **2018**, *11*, 649–663. [[CrossRef](#)]
109. Ndong Ba, A.; Verdin, A.; Cazier, F.; Garçon, G.; Thomas, J.; Cabral, M.; Dewaele, D.; Genevray, P.; Garat, A.; Allorge, D.; et al. Individual Exposure Level Following Indoor and Outdoor Air Pollution Exposure in Dakar (Senegal). *Environ. Pollut.* **2019**, *248*, 397–407. [[CrossRef](#)]
110. Chinh, N.N.B.; Fujii, Y.; Hien, T.T.; Takenaka, N. Characteristics of Gas Phase Carbonyl Emission and Excess Risk from Incense Stick Burning. *Water Air Soil Pollut.* **2020**, *231*, 297. [[CrossRef](#)]
111. Vora, A.; Chalbot, M.-C.G.; Shin, J.Y.; Kavouras, I.G. Size Distribution and Lung Deposition of Particle Mass Generated by Indoor Activities. *Indoor Built Environ.* **2021**, *30*, 1344–1352. [[CrossRef](#)]
112. Thuy, N.T.; May, D.T.; Thao, D.N.P.; Thuy, V.T.T.; Thanh, D.V.; Thanh, N.T.; Huy, N.N. Field Study of Visitors' Behavior in Incense Burning and Its Induced Air Pollution Assessment and Treatment. *Environ. Sci. Pollut. Res.* **2022**, *29*, 45933–45946. [[CrossRef](#)] [[PubMed](#)]
113. Yang, T.-T.; Ho, S.-C.; Chuang, L.-T.; Chuang, H.-C.; Li, Y.-T.; Wu, J.-J. Characterization of Particulate-Phase Polycyclic Aromatic Hydrocarbons Emitted from Incense Burning and Their Bioreactivity in RAW264.7 Macrophage. *Environ. Pollut.* **2017**, *220*, 1190–1198. [[CrossRef](#)] [[PubMed](#)]
114. Bu-Olayan, A.H.; Thomas, B.V. Exposition of Respiratory Ailments from Trace Metals Concentrations in Incenses. *Sci. Rep.* **2021**, *11*, 10210. [[CrossRef](#)] [[PubMed](#)]

115. Raad, D.; Gaddam, S.; Schunemann, H.J.; Irani, J.; Abou Jaoude, P.; Honeine, R.; Akl, E.A. Effects of Water-Pipe Smoking on Lung Function: A Systematic Review and Meta-Analysis. *Chest* **2011**, *139*, 764–774. [[CrossRef](#)] [[PubMed](#)]
116. Tse, L.A.; Yu, I.T.-S.; Qiu, H.; Au, J.S.K.; Wang, X.-R. A Case-Referent Study of Lung Cancer and Incense Smoke, Smoking, and Residential Radon in Chinese Men. *Environ. Health Perspect* **2011**, *119*, 1641–1646. [[CrossRef](#)] [[PubMed](#)]
117. Almedawar, M.M.; Walsh, J.L.; Isma'eel, H.A. Waterpipe Smoking and Risk of Coronary Artery Disease. *Curr. Opin. Cardiol.* **2016**, *31*, 545–550. [[CrossRef](#)]
118. El-Zaatari, Z.M.; Chami, H.A.; Zaatari, G.S. Health Effects Associated with Waterpipe Smoking. *Tob. Control.* **2015**, *24*, i31–i43. [[CrossRef](#)]
119. Veen, M. Carbon Monoxide Poisoning Caused by Water Pipe Smoking: A Case Series. *J. Emerg. Med.* **2016**, *51*, e41–e44. [[CrossRef](#)]
120. Chen, Y.C.; Ho, W.C.; Yu, Y.H. Adolescent Lung Function Associated with Incense Burning and Other Environmental Exposures at Home. *Indoor. Air.* **2017**, *27*, 746–752. [[CrossRef](#)]
121. Kuo, S.-C.; Tsai, Y.I. Emission Characteristics of Allergenic Terpenols in PM<sub>2.5</sub> Released from Incense Burning and the Effect of Light on the Emissions. *Sci. Total Environ.* **2017**, *584–585*, 495–504. [[CrossRef](#)] [[PubMed](#)]
122. Eichhorn, L.; Michaelis, D.; Kemmerer, M.; Jüttner, B.; Tetzlaff, K. Carbon Monoxide Poisoning from Waterpipe Smoking: A Retrospective Cohort Study. *Clin. Toxicol.* **2018**, *56*, 264–272. [[CrossRef](#)]
123. Kammoolkon, R.; Taneepanichskul, N.; Pitaknoppakul, N.; Lertmaharit, S.; Lohsoonthorn, V. Incense Smoke and Increasing Carotid Intima Media Thickness: A Cross-Sectional Study of the Thai-Vietnamese Community. *Asia Pac. J. Public Health* **2018**, *30*, 178–187. [[CrossRef](#)] [[PubMed](#)]
124. Wei, C.-F.; Chen, M.-H.; Lin, C.-C.; Guo, Y.L.; Lin, S.-J.; Hsieh, W.-S.; Chen, P.-C. Household Incense Burning and Infant Gross Motor Development: Results from the Taiwan Birth Cohort Study. *Environ. Int.* **2018**, *115*, 110–116. [[CrossRef](#)] [[PubMed](#)]
125. Ghosh, B.; Reyes-Caballero, H.; Akgün-Ölmez, S.G.; Nishida, K.; Chandrala, L.; Smirnova, L.; Biswal, S.; Sidhaye, V.K. Effect of Sub-Chronic Exposure to Cigarette Smoke, Electronic Cigarette and Waterpipe on Human Lung Epithelial Barrier Function. *BMC. Pulm. Med.* **2020**, *20*, 216. [[CrossRef](#)]
126. Guo, S.-E.; Chi, M.-C.; Lin, C.-M.; Yang, T.-M. Contributions of Burning Incense on Indoor Air Pollution Levels and on the Health Status of Patients with Chronic Obstructive Pulmonary Disease. *PeerJ* **2020**, *8*, e9768. [[CrossRef](#)]
127. Wei, C.-F.; Lin, C.-C.; Tsai, M.-S.; Guo, Y.L.; Lin, S.-J.; Liao, H.-F.; Hsieh, W.-S.; Chen, M.-H.; Chen, P.-C. Associations between Household Incense Burning and Delayed Motor Development among Preterm Infants Modified by Gestational Age and Maternal Educational Status. *Indoor Air* **2021**, *31*, 660–672. [[CrossRef](#)]
128. Hung, D.-Z.; Yang, K.-W.; Wu, C.-C.; Hung, Y.-H. Lead Poisoning Due to Incense Burning: An Outbreak in a Family. *Clin. Toxicol.* **2021**, *59*, 756–759. [[CrossRef](#)]
129. Hsieh, S.-W.; Chen, S.-C.; Chen, C.-H.; Wu, M.-T.; Hung, C.-H. Risk of Cognitive Impairment from Exposure to Incense Smoke. *Int. J. Environ. Health Res.* **2021**. [[CrossRef](#)]
130. Lee, C.-W.; Vo, T.T.T.; Wee, Y.; Chiang, Y.-C.; Chi, M.-C.; Chen, M.-L.; Hsu, L.-F.; Fang, M.-L.; Lee, K.-H.; Guo, S.-E.; et al. The Adverse Impact of Incense Smoke on Human Health: From Mechanisms to Implications. *J. Inflamm. Res.* **2021**, *14*, 5451–5472. [[CrossRef](#)]
131. Qasim, H.; Alarabi, A.B.; Alzoubi, K.H.; Karim, Z.A.; Alshbool, F.Z.; Khasawneh, F.T. The Effects of Hookah/Waterpipe Smoking on General Health and the Cardiovascular System. *Environ. Health Prev. Med.* **2019**, *24*, 58. [[CrossRef](#)] [[PubMed](#)]
132. Yadav, V.K.; Malik, P.; Tirth, V.; Khan, S.H.; Yadav, K.K.; Islam, S.; Choudhary, N.; Inwati, G.K.; Arabi, A.; Kim, D.-H.; et al. Health and Environmental Risks of Incense Smoke: Mechanistic Insights and Cumulative Evidence. *J. Inflamm. Res.* **2022**, *15*, 2665–2693. [[CrossRef](#)] [[PubMed](#)]
133. Hussein, T.; Al-Jagheer, O.; Bqour, N.; Zidan, B.; Lahlouh, B. Exposure to Aerosols Emitted from Common Heating Combustion Sources Indoors—The Jordanian Case as an Example for Eastern Mediterranean Conditions. *Atmosphere* **2022**, *13*, 870. [[CrossRef](#)]
134. Hussein, T.; Kulmala, M. Indoor Aerosol Modeling: Basic Principles and Practical Applications. *Water Air Soil Pollut. Focus* **2008**, *8*, 23–34. [[CrossRef](#)]
135. Hussein, T.; Korhonen, H.; Herrmann, E.; Hämeri, K.; Lehtinen, K.E.J.; Kulmala, M. Emission Rates Due to Indoor Activities: Indoor Aerosol Model Development, Evaluation, and Applications. *Aerosol Sci. Technol.* **2005**, *39*, 1111–1127. [[CrossRef](#)]
136. Salthammer, T.; Zhao, J.; Schieweck, A.; Uhde, E.; Hussein, T.; Antretter, F.; Künzel, H.; Pazold, M.; Radon, J.; Birmili, W. A Holistic Modeling Framework for Estimating the Influence of Climate Change on Indoor Air Quality. *Indoor Air* **2022**, *32*, e13039. [[CrossRef](#)]
137. Hussein, T.; Löndahl, J.; Paasonen, P.; Koivisto, A.J.; Petäjä, T.; Hämeri, K.; Kulmala, M. Modeling Regional Deposited Dose of Submicron Aerosol Particles. *Sci. Total Environ.* **2013**, *458–460*, 140–149. [[CrossRef](#)]
138. Hussein, T.; Wierzbicka, A.; Löndahl, J.; Lazaridis, M.; Hänninen, O. Indoor Aerosol Modeling for Assessment of Exposure and Respiratory Tract Deposited Dose. *Atmos. Environ.* **2015**, *106*, 402–411. [[CrossRef](#)]
139. Hussein, T.; Saleh, S.S.A.; dos Santos, V.N.; Boor, B.E.; Koivisto, A.J.; Löndahl, J. Regional Inhaled Deposited Dose of Urban Aerosols in an Eastern Mediterranean City. *Atmosphere* **2019**, *10*, 530. [[CrossRef](#)]
140. Hussein, T.; Boor, B.E.; Löndahl, J. Regional Inhaled Deposited Dose of Indoor Combustion-Generated Aerosols in Jordanian Urban Homes. *Atmosphere* **2020**, *11*, 1150. [[CrossRef](#)]
141. Human Respiratory Tract Model for Radiological Protection. A Report of a Task Group of the International Commission on Radiological Protection. *Ann. ICRP* **1994**, *24*, 1–482.

142. Anjilvel, S.; Asgharian, B. A Multiple-Path Model of Particle Deposition in the Rat Lung. *Toxicol. Sci.* **1995**, *28*, 41–50. [[CrossRef](#)]
143. Löndahl, J.; Massling, A.; Pagels, J.; Swietlicki, E.; Vaclavik, E.; Loft, S. Size-Resolved Respiratory-Tract Deposition of Fine and Ultrafine Hydrophobic and Hygroscopic Aerosol Particles during Rest and Exercise. *Inhal. Toxicol.* **2007**, *19*, 109–116. [[CrossRef](#)] [[PubMed](#)]
144. Maziak, W.; Rastam, S.; Ibrahim, I.; Ward, K.D.; Shihadeh, A.; Eissenberg, T. CO Exposure, Puff Topography, and Subjective Effects in Waterpipe Tobacco Smokers. *Nicotine Tob. Res.* **2009**, *11*, 806–811. [[CrossRef](#)]
145. Maziak, W.; Rastam, S.; Shihadeh, A.L.; Bazzi, A.; Ibrahim, I.; Zaatari, G.S.; Ward, K.D.; Eissenberg, T. Nicotine Exposure in Daily Waterpipe Smokers and Its Relation to Puff Topography. *Addict. Behav.* **2011**, *36*, 397–399. [[CrossRef](#)]
146. Soule, E.K.; Ramôa, C.; Eissenberg, T.; Cobb, C.O. Differences in Puff Topography, Toxicant Exposure, and Subjective Response between Waterpipe Tobacco Smoking Men and Women. *Exp. Clin. Psychopharmacol.* **2018**, *26*, 440–447. [[CrossRef](#)]
147. Hauser, C.D.; Mailig, R.; Stadler, H.; Reed, J.; Chen, S.; Uffman, E.; Bernd, K. Waterpipe Tobacco Smoke Toxicity: The Impact of Waterpipe Size. *Tob. Control.* **2019**, *29*, s90–s94. [[CrossRef](#)]
148. Markowicz, P.; Löndahl, J.; Wierzbicka, A.; Suleiman, R.; Shihadeh, A.; Larsson, L. A Study on Particles and Some Microbial Markers in Waterpipe Tobacco Smoke. *Sci. Total. Environ.* **2014**, *499*, 107–113. [[CrossRef](#)]
149. Travers, M.J.; Kulak, J.A.; Vogl, L. Waterpipe Cafés Are Hazardous to Your Health: Determination of a Waterpipe Specific Calibration Factor. *Int. J. Hyg. Environ. Health* **2018**, *221*, 48–53. [[CrossRef](#)]
150. Wu, C.L.; Chao, C.Y.H.; Sze-To, G.N.; Wan, M.P.; Chan, T.C. Ultrafine Particle Emissions from Cigarette Smouldering, Incense Burning, Vacuum Cleaner Motor Operation and Cooking. *Indoor. Built. Environ.* **2012**, *21*, 782–796. [[CrossRef](#)]
151. Lai, A.C.K.; Nazaroff, W.W. Modeling Indoor Particle Deposition from Turbulent Flow onto Smooth Surfaces. *J. Aerosol Sci.* **2000**, *31*, 463–476. [[CrossRef](#)]
152. Hussein, T.; Smolik, J.; Kerminen, V.-M.; Kulmala, M. Modeling Dry Deposition of Aerosol Particles onto Rough Surfaces. *Aerosol Sci. Technol.* **2012**, *46*, 44–59. [[CrossRef](#)]
153. See, S.W.; Balasubramanian, R.; Man Joshi, U. Physical Characteristics of Nanoparticles Emitted from Incense Smoke. *Sci. Technol. Adv. Mater.* **2007**, *8*, 25–32. [[CrossRef](#)]
154. Kant, N.; Müller, R.; Braun, M.; Gerber, A.; Groneberg, D. Particulate Matter in Second-Hand Smoke Emitted from Different Cigarette Sizes and Types of the Brand Vogue Mainly Smoked by Women. *Int. J. Environ. Res. Public Health* **2016**, *13*, 799. [[CrossRef](#)] [[PubMed](#)]
155. Braun, M.; Koger, F.; Klingelhöfer, D.; Müller, R.; Groneberg, D.A. Particulate Matter Emissions of Four Different Cigarette Types of One Popular Brand: Influence of Tobacco Strength and Additives. *Int. J. Environ. Res. Public Health* **2019**, *16*, 263. [[CrossRef](#)] [[PubMed](#)]
156. Adams, H.S.; Nieuwenhuijsen, M.J.; Colvile, R.N. Determinants of Fine Particle (PM<sub>2.5</sub>) Personal Exposure Levels in Transport Microenvironments, London, UK. *Atmos. Environ.* **2001**, *35*, 4557–4566. [[CrossRef](#)]
157. Chan, L.Y.; Lau, W.L.; Zou, S.C.; Cao, Z.X.; Lai, S.C. Exposure Level of Carbon Monoxide and Respirable Suspended Particulate in Public Transportation Modes While Commuting in Urban Area of Guangzhou, China. *Atmos. Environ.* **2002**, *36*, 5831–5840. [[CrossRef](#)]
158. Kaur, S.; Nieuwenhuijsen, M.J.; Colvile, R.N. Pedestrian Exposure to Air Pollution along a Major Road in Central London, UK. *Atmos. Environ.* **2005**, *39*, 7307–7320. [[CrossRef](#)]
159. Fondelli, M.C.; Chellini, E.; Yli-Tuomi, T.; Cenni, I.; Gasparrini, A.; Nava, S.; Garcia-Orellana, I.; Lupi, A.; Grechi, D.; Mallone, S.; et al. Fine Particle Concentrations in Buses and Taxis in Florence, Italy. *Atmos. Environ.* **2008**, *42*, 8185–8193. [[CrossRef](#)]
160. Sohn, H.; Lee, K. Impact of Smoking on In-Vehicle Fine Particle Exposure during Driving. *Atmos. Environ.* **2010**, *44*, 3465–3468. [[CrossRef](#)]
161. Buonanno, G.; Morawska, L.; Stabile, L.; Viola, A. Exposure to Particle Number, Surface Area and PM Concentrations in Pizzerias. *Atmos. Environ.* **2010**, *44*, 3963–3969. [[CrossRef](#)]
162. Buonanno, G.; Morawska, L.; Stabile, L.; Wang, L.; Giovinco, G. A Comparison of Submicrometer Particle Dose between Australian and Italian People. *Environ. Pollut.* **2012**, *169*, 183–189. [[CrossRef](#)] [[PubMed](#)]
163. Fuoco, F.C.; Stabile, L.; Buonanno, G.; Trassiera, C.V.; Massimo, A.; Russi, A.; Mazaheri, M.; Morawska, L.; Andrade, A. Indoor Air Quality in Naturally Ventilated Italian Classrooms. *Atmosphere* **2015**, *6*, 1652–1675. [[CrossRef](#)]
164. Fuoco, F.C.; Buonanno, G.; Stabile, L.; Vigo, P. Influential Parameters on Particle Concentration and Size Distribution in the Mainstream of E-Cigarettes. *Environ. Pollut.* **2014**, *184*, 523–529. [[CrossRef](#)] [[PubMed](#)]
165. Stabile, L.; Fuoco, F.C.; Marini, S.; Buonanno, G. Effects of the Exposure to Indoor Cooking-Generated Particles on Nitric Oxide Exhaled by Women. *Atmos. Environ.* **2015**, *103*, 238–246. [[CrossRef](#)]
166. Stabile, L.; Dell’Isola, M.; Frattolillo, A.; Massimo, A.; Russi, A. Effect of Natural Ventilation and Manual Airing on Indoor Air Quality in Naturally Ventilated Italian Classrooms. *Built. Environ.* **2016**, *98*, 180–189. [[CrossRef](#)]
167. Mazaheri, M.; Lin, W.; Clifford, S.; Yue, D.; Zhai, Y.; Xu, M.; Rizza, V.; Morawska, L. Characteristics of School Children’s Personal Exposure to Ultrafine Particles in Heshan, Pearl River Delta, China—A Pilot Study. *Environ. Int.* **2019**, *132*, 105134. [[CrossRef](#)]
168. Hussein, T.; Hämeri, K.; Aalto, P.; Asmi, A.; Kakko, L.; Kulmala, M. Particle Size Characterization and the Indoor-to-Outdoor Relationship of Atmospheric Aerosols in Helsinki. *Scand. J. Work Environ. Health* **2004**, *30* (Suppl. S2), 54–62.
169. Hussein, T.; Hämeri, K.; Heikkinen, M.S.A.; Kulmala, M. Indoor and Outdoor Particle Size Characterization at a Family House in Espoo-Finland. *Atmos. Environ.* **2005**, *39*, 3697–3709. [[CrossRef](#)]

170. Hussein, T.; Glytsos, T.; Ondráček, J.; Dohányosová, P.; Ždímal, V.; Hämeri, K.; Lazaridis, M.; Smolík, J.; Kulmala, M. Particle Size Characterization and Emission Rates during Indoor Activities in a House. *Atmos. Environ.* **2006**, *40*, 4285–4307. [[CrossRef](#)]
171. Koivisto, A.J.; Hussein, T.; Niemelä, R.; Tuomi, T.; Hämeri, K. Impact of Particle Emissions of New Laser Printers on Modeled Office Room. *Atmos. Environ.* **2010**, *44*, 2140–2146. [[CrossRef](#)]
172. Hussein, T. Particle Size Distributions inside a University Office in Amman, Jordan. *Jordan. J. Phys.* **2014**, *7*, 73–83.
173. Mølgaard, B.; Viitanen, A.-K.; Kangas, A.; Huhtiniemi, M.; Larsen, S.T.; Vanhala, E.; Hussein, T.; Boor, B.E.; Hämeri, K.; Koivisto, A.J. Exposure to Airborne Particles and Volatile Organic Compounds from Polyurethane Molding, Spray Painting, Lacquering, and Gluing in a Workshop. *Int. J. Environ. Res. Public Health* **2015**, *12*, 3756. [[CrossRef](#)] [[PubMed](#)]
174. Hussein, T.; Dada, L.; Juwhari, H.; Faouri, D. Characterization, Fate, and Re-Suspension of Aerosol Particles (0.3–10 Mm): The Effects of Occupancy and Carpet Use. *Aerosol Air Qual. Res.* **2015**, *15*, 2367–2377. [[CrossRef](#)]
175. Lazaridis, M.; Eleftheriadis, K.; Ždímal, V.; Schwarz, J.; Wagner, Z.; Ondráček, J.; Drossinos, Y.; Glytsos, T.; Vratolis, S.; Torseth, K.; et al. Number Concentrations and Modal Structure of Indoor/Outdoor Fine Particles in Four European Cities. *Aerosol Air Qual. Res.* **2017**, *17*, 131–146. [[CrossRef](#)]
176. Maragkidou, A.; Jaghbeir, O.; Hämeri, K.; Hussein, T. Aerosol Particles (0.3–10 µm) inside an Educational Workshop—Emission Rate and Inhaled Deposited Dose. *Build. Environ.* **2018**, *140*, 80–89. [[CrossRef](#)]
177. Zhao, J.; Birmili, W.; Hussein, T.; Wehner, B.; Wiedensohler, A. Particle Number Emission Rates of Aerosol Sources in 40 German Households and Their Contributions to Ultrafine and Fine Particle Exposure. *Indoor Air* **2021**, *31*, 818–831. [[CrossRef](#)]
178. Viitanen, A.-K.; Kallonen, K.; Kukko, K.; Kanerva, T.; Saukko, E.; Hussein, T.; Hämeri, K.; Säämänen, A. Technical Control of Nanoparticle Emissions from Desktop 3D Printing. *Indoor Air* **2021**, *31*, 1061–1071. [[CrossRef](#)]
179. Wierzbicka, A.; Bohgard, M.; Pagels, J.H.; Dahl, A.; Löndahl, J.; Hussein, T.; Swietlicki, E.; Gudmundsson, A. Quantification of Differences between Occupancy and Total Monitoring Periods for Better Assessment of Exposure to Particles in Indoor Environments. *Atmos. Environ.* **2015**, *106*, 419–428. [[CrossRef](#)]
180. Stabile, L.; Buonanno, G.; Ficco, G.; Scungio, M. Smokers’ Lung Cancer Risk Related to the Cigarette-Generated Mainstream Particles. *J. Aerosol Sci.* **2017**, *107*, 41–54. [[CrossRef](#)]
181. Sahu, S.K.; Tiwari, M.; Bhangare, R.C.; Pandit, G.G. Particle Size Distribution of Mainstream and Exhaled Cigarette Smoke and Predictive Deposition in Human Respiratory Tract. *Aerosol Air Qual. Res.* **2013**, *13*, 324–332. [[CrossRef](#)]
182. Li, X.; Kong, H.; Zhang, X.; Peng, B.; Nie, C.; Shen, G.; Liu, H. Characterization of Particle Size Distribution of Mainstream Cigarette Smoke Generated by Smoking Machine with an Electrical Low Pressure Impactor. *J. Environ. Sci.* **2014**, *26*, 827–833. [[CrossRef](#)] [[PubMed](#)]
183. Jetter, J.J.; Guo, Z.; McBrien, J.A.; Flynn, M.R. Characterization of Emissions from Burning Incense. *Sci. Total. Environ.* **2002**, *295*, 51–67. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.