



Editorial

Questioning Established Theories and Treatment Methods Related to Iron and Other Metal Metabolic Changes, Affecting All Major Diseases and Billions of Patients

George J. Kontoghiorghes

Postgraduate Research Institute of Science, Technology, Environment and Medicine, 3 Ammochostou Street, Limassol 3021, Cyprus; kontoghiorghes.g.j@pri.ac.cy; Tel./Fax: +357-2627-2076

Keywords: metal metabolism; iron chelation; metal related diseases; treatment; metal toxicity; drugs in iron diseases

The medical and scientific literature is dominated by highly cited historical theories and findings. It is important to question these theories and report findings that are deemed to be less significant, as this could introduce new perspectives in theoretical models, which will aim to better understand disease processes and their treatments. The gaps and limitations in theoretical models are usually identified as molecular basic science models are translated into biological models and then further into clinical ones, especially when a wide range of treatment options exist.

There are many examples in various areas of medicine, including diseases related to iron metabolism, where the translation of basic science information, which is potentially insufficient, could lead to errors when operating disease models and suboptimal treatment for patients. Billions of people are affected by iron-related diseases worldwide, and most of these involve iron deficiency and iron overload, which are treated using iron chelator complexes and chelators that eliminate iron, respectively [1–4]. Most of the other common diseases affecting humans, such as cancer, neurodegenerative and haematological diseases, infections including COVID-19 and also many others, appear to cause changes in iron metabolic pathways. In most cases, these changes are characterized by abnormal levels in iron diagnostic indices such as the level of haemoglobin, serum ferritin, serum iron, abnormal focal iron deposition in specific cells or tissues, etc. Despite the fact that iron may not be directly implicated in many of these diseases, interventional therapies that restore or improve iron related imbalances may facilitate the treatment of or even improve the prognosis for patients with such diseases [5].

In general, the control of iron, which is involved in many iron containing enzymes and metabolic pathways in all organisms including humans, may play a central role in therapeutic protocols involving both common and rare diseases [5]. Similar roles and treatment approaches are considered in the case of other essential metals such as copper and zinc [6–11].

Abnormal metal metabolic changes and associated metal interactions can also affect therapeutic interventions and the overall treatment of patients. For example, changes in iron metabolism such as iron overload can also affect the metabolism of many other metals including essential, xenobiotic and diagnostic metals. Most of these metals share and compete for the metal transport properties of transferrin and subsequently interfere with the metabolic, therapeutic or diagnostic effects of other metals [12]. Similarly, the metabolism and therapeutic effects of many drugs, as well as the metabolism and nutritional effects of many nutrients, most of which have metal binding properties, can also be affected [13–16]. All these interactions and their implications on health and disease have not yet been fully investigated.

Citation: Kontoghiorghes, G.J. Questioning Established Theories and Treatment Methods Related to Iron and Other Metal Metabolic Changes, Affecting All Major Diseases and Billions of Patients. *Int. J. Mol. Sci.* **2022**, *23*, 1364. <https://doi.org/10.3390/ijms23031364>

Received: 28 October 2021

Accepted: 16 December 2022

Published: 25 January 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

The lack of in-depth investigations and inaccurate reporting, as well as distorted, misleading or poorly promoted findings, may also lead to insufficient information and suboptimal patient treatments. This influence is related in most cases to the level of academic promotion and the number of citations of particular theories and is also generally observed to take place in relation to the marketing activities of private companies involved mainly in the sale of new, very expensive medicines and medical devices [17].

There are many examples in the literature questioning established theories in the risk/benefit assessment for the use of different drugs, dietary components and other medicines. In particular, the benefits, including the efficacy and low toxicity of traditional/folk medicines, are questioned in the medical literature despite the fact that millions of people are preferably using them in comparison to synthetic drugs. Similarly, millions of people are using routinely old/generic drugs such as the chelating drug ethylenediaminetetraacetic acid (EDTA) because of the perceived multiple health benefits [18,19].

The existence of several drugs for each disease can be beneficial in cases of adverse reactions from one particular drug. In contrast, the selection of a specific drug that is randomly selected from a group of drugs for a particular disease can also lead, in many cases, to suboptimal treatment for patients. For example, the conflict of opinion in the use of different chelating drugs and metal complexes in different clinical conditions and categories of patient is highlighted in the medical literature [17,20–27].

In contrast to conflicts that arise from the selection of drugs, differences of opinion in the biology and metabolism of iron and other metal ions is on a much smaller scale compared to the use of different drugs for diseases related to abnormal iron or other metal metabolism. However, such differences and related discussions are necessary for exploring the role of metal ions and their metabolic pathways in the maintenance of metal homeostasis under normal conditions and also for the treatment of many diseases.

An example of a molecular interaction of physiological and toxicological significance, which has been neglected for more than a half a century, is the ferroxidase and potent antioxidant activity of transferrin [12,28]. Despite the fact that caeruloplasmin is promoted and cited in hundreds of relevant textbooks and papers as the protein required for the oxidation of ferrous to ferric iron before its uptake by transferrin, transferrin has also previously been shown to be able to oxidize ferrous to ferric iron before its chelation and the formation of the transferrin ferric iron complex [29–34]. The extent of the participation of either caeruloplasmin or transferrin in the ferroxidase activity under physiological conditions has not yet been fully investigated. In the meantime, similar properties have been shown by the iron chelating drugs deferiprone, deferasirox and deferoxamine, which can rapidly oxidize ferrous to ferric iron at physiological conditions before the formation of the corresponding ferric iron complexes [15,35–37]. The ferroxidase and iron complex formation activity of the iron chelating drugs is the cornerstone of their antioxidant activity in the Fenton reaction and has been utilized and shown to be related to the powerful antioxidant effects exerted in many *in vitro*, *in vivo* and clinical models of oxidative stress [38]. The therapeutic application of chelating drugs in many diseases associated with free radical pathology, as well as the role of transferrin, is currently under investigation [28,39,40]. One major characteristic of chelators including transferrin and chelating drugs is their pro-oxidant effects under certain conditions [41–43]. In particular, the pro-oxidant effects of some selective chelator iron and copper complexes have been used in different experimental models for the development of new anticancer drugs [44–46].

The transport of iron and other metal ions in and out of different cells is of physiological, pharmacological and toxicological importance. The factors affecting metal ion transport in and out of cells is the subject of continuous investigations and discussions in health and disease. In particular, changes in the regulatory molecules, mechanisms and homeostatic controls involved in the transport of iron and other metal ions could have serious implications for the health statuses of many categories of patients [4,5,47]. Despite the fact that emphasis in the medical literature is primarily given to molecular biol-

ogy effect changes, chelation and other molecular effects involving metal ions could also have major implications on the transportation, distribution and deposition of metal ions, as well as on the causes or progression of different diseases [4,5,47–52]. In general, natural or synthetic lipophilic metal chelator complexes can increase cellular metal ion intake and increase metal ion absorption, whereas hydrophilic chelators can remove intracellular metal ion and increase metal ion excretion [4,5,44,49]. In particular, the increase of iron absorption by lipophilic iron chelator complexes is important for all those affected by iron deficiency and mostly vegetarian populations [53].

In addition to the effects of the chelation of metals, there are many other unexplored areas in relation to metal ion metabolism and associated diseases. One such example is the impact of the regulatory excretion of natural and xenobiotic metal ions. Despite the fact that the homeostatic controls, mechanisms and route of the regulatory excretion of copper and zinc have been partly determined, no such information is available for iron and many of the diagnostic metal ions, all of which are important for the diagnosis and treatment of different diseases including iron overload [54–59]. In particular, the elimination of excess iron in the absence of chelation therapy in post-transplanted thalassaemia patients and some other categories of patients requires further investigation [60,61].

A major unexplored area related to many diseases is the identification of the mechanisms and factors involved in the abnormal distribution and deposition of metal ions in different cells and tissues. In particular, the abnormal distribution and deposition of iron has been observed in many diseases with high morbidity and mortality rates, including many cancer, haematological, neurodegenerative and infectious diseases including COVID-19 [4,5]. Although iron may not directly be implicated in the fatal or serious pathophysiological effects in these diseases, iron abnormalities can constitute a major factor in the progress and outcome of the pathological effects of the underlying disease and also their treatment. Within this context and whenever appropriate, iron chelating drugs and other chelators and chelator iron complexes could be introduced to restore iron balance and improve the overall therapeutic effects, including combination therapies with other drugs [4,5].

The targeting of specific mechanisms and pathways involving iron and other metal ion abnormalities is important for many diseases. An example is the major role of macrophages in the abnormal distribution and deposition of iron in many malignant, inflammatory and infectious diseases including COVID-19, which in most cases is also characterized by an elevation in serum ferritin and a reduction in serum iron levels [62–67]. The mechanisms involved in these changes are not yet fully clarified. Similarly, therapeutic interventions, which may include chelating drugs for the targeting of specific pathways involving macrophage iron and related changes, as well as other relevant targets need further investigation [28,68–70].

The diagnosis and treatment of different conditions related to iron and other metal metabolic abnormalities are among the major areas of conflict and controversy involving marketing tactics for the promotion of different medical diagnostics and drugs by pharmaceutical and other companies. In general, the marketing activities of private companies affect academic research and the reporting of clinical findings and also, most importantly, the morbidity and mortality rate of billions of patients [4,5,25,53,71–73]. For example, in the case of iron deficiency anaemia, hundreds of different chelator and other iron complexes are promoted, especially of new and expensive formulations claiming effective treatments and superiority over other inexpensive, generic formulations [53,74,75].

Similar competition takes place in the case of iron chelation therapy for the removal of excess toxic iron in iron overload conditions, where there is conflicting opinion in the selection of the appropriate drug between deferoxamine, deferiprone and deferasirox. However, only deferiprone and the deferiprone/deferoxamine combination have been shown to remove iron from the heart and all other major organs effectively, as well as returning body iron levels to those similar to normal individuals [76–78]. Furthermore, in a long term follow up study, it has been shown that the deferiprone/deferoxamine combination can increase the long term survival of thalassaemia patients [79]. It should be not-

ed that the combination of drugs with similar pharmacological activity such as the deferiprone/deferioxamine combination is more effective than either of the monotherapies, but such therapeutic strategies are not promoted by pharmaceutical companies or even by drug regulatory authorities [72].

Conflicting theoretical models and research findings involving academic researchers sponsored by pharmaceutical companies are also widely discussed in the medical and other academic literature. In many cases, elite journals are also involved in such promotions by pharmaceutical companies, especially when publishing questionable theoretical models and findings, causing confusion to patients [72]. There are many examples such as the categorization of thalassaemia intermedia in the so called “non-transfusion-dependent-thalassaemia” group, where deferiasirox was promoted as a new therapy despite the fact that deferioxamine and deferiprone have been used in this category of patients for over 50 years [72,80,81].

Another major issue affecting the safety and survival of patients in diseases related to iron and other metal metabolism, as well as chelation therapies, is the interaction of chelating drugs and chelator metal complexes with other drugs and therapies [13,14,82–86]. In this context, such interactions may be critical and established related treatment protocols involving such interactions have to be modified. In particular, the safety and efficacy of newly approved drugs in many cases is questionable because of the lack of sufficient long term clinical outcome data. For example, the introduction of erythropoietic biologics such as luspatercept (Reblozyl) has been considered for reducing red blood cell transfusions in haematological diseases including thalassaemia, where it has been suggested that a 20% reduction is possible [87–89]. However, the use of luspatercept in thalassaemia major is questionable since, even if haematopoiesis can be increased in patients, this can only produce abnormal, non-functional haemoglobin. Similarly, serious concerns remain in other aspects of therapy, such as the safety, efficacy and the cost of luspatercept and also of other biologics, including those used in the treatment of COVID-19, since, in general, the immunogenicity and other toxic side effects of these compounds have not yet been fully investigated [90,91].

In conclusion, established theoretical models and principles in science and medicine have to be scrutinized and re-evaluated at all times, with an emphasis on the improvement of knowledge and, where appropriate, the introduction of better treatments for the increased survival and safety of patients. In cases of pharmaceutical products, the influence of private companies including the promotion of their new products through the medical literature may affect the morbidity and mortality rate of many categories of patients. Herein, several examples examining the validity of a number of established theoretical models and drugs associated with diseases related to iron and other metal metabolic diseases have been questioned. It is hoped that further discussions and re-evaluation could lead in the design of better theoretical models in science and medicine related to iron and other metal metabolism and, most importantly, for safer and more effective treatments for patients.

Funding: This research received no external funding.

Acknowledgments: I thank Christina N. Kontoghiorghe for the useful comments without implying endorsement regarding the manuscript.

Conflicts of Interest: The author declares no conflict of interest. The author has no commercial associations including consultancies, stock ownership, equity interest, patent/licensing arrangement etc. that might pose a conflict of interest in connection with the submitted article.

References

1. Andrews, N.C. Disorders of iron metabolism. *N. Engl. J. Med.* **1999**, *341*, 1986–1995. <https://doi.org/10.1056/NEJM199912233412607>. Erratum in *N. Engl. J. Med.* **2000**, *342*, 364.
2. Gozzelino, R.; Arosio, P. Iron Homeostasis in Health and Disease. *Int. J. Mol. Sci.* **2016**, *17*, 130. [10.3390/ijms17010130](https://doi.org/10.3390/ijms17010130).
3. Cairo, G.; Bernuzzi, F.; Recalcati, S.A.A. Precious metal: Iron, an essential nutrient for all cells. *Genes Nutr.* **2006**, *1*, 25–39.

4. Kontoghiorghes, G.J. Advances on Chelation and Chelator Metal Complexes in Medicine. *Int. J. Mol. Sci.* **2020**, *21*, 2499. <https://doi.org/10.3390/ijms21072499>.
5. Kontoghiorghes, G.J.; Kontoghiorghes, C.N. Iron and Chelation in Biochemistry and Medicine: New Approaches to Controlling Iron Metabolism and Treating Related Diseases. *Cells* **2020**, *9*, 1456.
6. Prasad, A.S. Zinc: An overview. *Nutrition* **1995**, *11*, 93–99.
7. Coleman, J.E. Zinc proteins: Enzymes, storage proteins, transcription factors, and replication proteins. *Annu. Rev. Biochem.* **1992**, *61*, 897–946.
8. Prasad, A.S. Zinc deficiency. *BMJ* **2003**, *326*, 409–410.
9. Daniel, K.G.; Harbach, R.H.; Guida, W.C.; Dou, Q.P. Copper storage diseases: Menkes, Wilsons, and cancer. *Front. Biosci.* **2004**, *9*, 2652–2662.
10. Baldari, S.; Di Rocco, G.; Toietta, G. Current Biomedical Use of Copper Chelation Therapy. *Int. J. Mol. Sci.* **2020**, *21*, 1069.
11. Balsano, C.; Porcu, C.; Sideri, S. Is copper a new target to counteract the progression of chronic diseases? *Metallomics* **2018**, *10*, 1712–1722.
12. Kontoghiorghes, C.N.; Kolnagou, A.; Kontoghiorghes, G.J. Potential clinical applications of chelating drugs in diseases targeting transferrin-bound iron and other metals. *Expert Opin. Investig. Drugs* **2013**, *22*, 591–618.
13. Djaldetti, M.; Fishman, P.; Notti, I.; Bessler, H. The effect of tetracycline administration on iron absorption in mice. *Biomedicine* **1981**, *35*, 150–152.
14. Konstantinou, E.; Pashalidis, I.; Kolnagou, A.; Kontoghiorghes, G.J. Interactions of hydroxycarbamide (hydroxyurea) with iron and copper: Implications on toxicity and therapeutic strategies. *Hemoglobin* **2011**, *35*, 237–246.
15. Timoshnikov, V.A.; Kobzeva, T.V.; Polyakov, N.E.; Kontoghiorghes, G.J. Redox Interactions of Vitamin C and Iron: Inhibition of the Pro-Oxidant Activity by Deferiprone. *Int. J. Mol. Sci.* **2020**, *21*, 3967. <https://doi.org/10.3390/ijms21113967>.
16. Kontoghiorghes, C.N.; Kolnagou, A.; Kontoghiorghes, G.J. Phytochelators Intended for Clinical Use in Iron Overload, Other Diseases of Iron Imbalance and Free Radical Pathology. *Molecules* **2015**, *20*, 20841–20872. <https://doi.org/10.3390/molecules201119725>.
17. Kontoghiorghes, G.J. Ethics in Medicines: Exposing Unethical Practices and Corruption in All Sectors of Medicines Is Essential for Improving Global Public Health and Saving Patients' Lives. *Medicines* **2021**, *8*, 54. <https://doi.org/10.3390/medicines8090054>.
18. Lamas, G.A.; Navas-Acien, A.; Mark, D.B.; Lee, K.L. Heavy Metals, Cardiovascular Disease, and the Unexpected Benefits of Chelation Therapy. *J. Am. Coll. Cardiol.* **2016**, *67*, 2411–2418. <https://doi.org/10.1016/j.jacc.2016.02.066>.
19. Born, T.; Kontoghiorghes, C.N.; Spyrou, A.; Kolnagou, A.; Kontoghiorghes, G.J. EDTA chelation reappraisal following new clinical trials and regular use in millions of patients: Review of preliminary findings and risk/benefit assessment. *Toxicol. Mech. Methods* **2012**, *23*, 11–17.
20. Steensma, D.P. Myelodysplasia paranoia: Iron as the new radon. *Leuk. Res.* **2009**, *33*, 1158–1163.
21. Lucania, G.; Vitranio, A.; Filosa, A.; Maggio, A. Chelation treatment in sickle-cell-anaemia: Much ado about nothing? *Br. J. Haematol.* **2011**, *154*, 545–555.
22. Maximova, N.; Gregori, M.; Simeone, R.; Sonzogni, A.; Boz, G.; Fucile, C.; Marini, V.; Martelli, A.; Mattioli, F. Safety and tolerability of deferasirox in pediatric hematopoietic stem cell transplant recipients: One facility's five years' experience of chelation treatment. *Oncotarget* **2017**, *8*, 63177–63186.
23. Braga, C.C.B.; Benites, B.D.; de Albuquerque, D.M.; Alvarez, M.C.; Seva-Pereira, T.; Duarte, B.K.L.; Costa, F.F.; Gilli, S.C.O.; Saad, S.T.O. Deferasirox associated with liver failure and death in a sickle cell anemia patient homozygous for the -1774delG polymorphism in the Abcc2 gene. *Clin. Case Rep.* **2017**, *5*, 1218–1221.
24. Vill K, Müller-Felber W, Teusch V; Blaschek, A., Gerstl, L., Huetker, S., Albert, M. H. Proximal muscular atrophy and weakness: An unusual adverse effect of deferasirox iron chelation therapy. *Neuromuscul. Disord.* **2016**, *26*, 322–325.
25. Kontoghiorghes, C.N.; Kontoghiorghes, G.J. New developments and controversies in iron metabolism and iron chelation therapy. *World J. Methodol.* **2016**, *6*, 1.
26. Kontoghiorghes, G.J.; Spyrou, A.; Kolnagou, A. Iron chelation therapy in hereditary hemochromatosis and thalassemia intermedia: Regulatory and non regulatory mechanisms of increased iron absorption. *Hemoglobin* **2010**, *34*, 251–264. <https://doi.org/10.3109/03630269.2010.486335>.
27. Nick, H.P. Iron chelation therapy in hereditary hemochromatosis and thalassemia intermedia: regulatory and non regulatory mechanisms of increased iron absorption [Kontoghiorghes GJ, Spyrou A, Kolnagou A. Hemoglobin. 2010;34(3); 251-264]. *Hemoglobin*. **2011**, *35*, 175–179. doi: 10.3109/03630269.2011.557173.
28. Kontoghiorghes, G.J. Iron mobilization from transferrin and non-transferrin-bound-iron by deferiprone. Implications in the treatment of thalassemia, anemia of chronic disease, cancer and other conditions. *Hemoglobin* **2006**, *30*, 183–200. <https://doi.org/10.1080/03630260600642450>.
29. White, K.N.; Conesa, C.; Sánchez, L.; Amini, M.; Farnaud, S.; Lörvorlak, C.; Evans, R.W. The transfer of iron between ceruloplasmin and transferrins. *Biochim. Biophys. Acta (BBA)-Gen. Subj.* **2012**, *1820*, 411–416. <https://doi.org/10.1016/j.bbagen.2011.10.006>.
30. Bates, G.W.; Workman, E.F.; Schlabach, M.R. Does transferrin exhibit ferroxidase activity? *Biochem. Biophys. Res. Commun.* **1973**, *50*, 84–90. [https://doi.org/10.1016/0006-291X\(73\)91067-X](https://doi.org/10.1016/0006-291X(73)91067-X).

31. Huebers, H.A.; Josephson, B.; Huebers, E.; Csiba, E.; Finch, C.A. Occupancy of the iron binding sites of human transferrin. *Proc. Natl. Acad. Sci. USA* **1984**, *81*, 4326–4330. <https://doi.org/10.1073/pnas.81.14.4326>.
32. Kontoghiorghes, G.J. The study of iron mobilisation from transferrin using α -ketohydroxy heteroaromatic chelators. *Biochim. Biophys. Acta (BBA)/Protein Struct. Mol.* **1986**, *869*, 141–146. [https://doi.org/10.1016/0167-4838\(86\)90288-8](https://doi.org/10.1016/0167-4838(86)90288-8).
33. Baldwin, D.A.; Jenny, E.R.; Aisen, P. The effect of human serum transferrin and milk lactoferrin on hydroxyl radical formation from superoxide and hydrogen peroxide. *J. Biol. Chem.* **1984**, *259*, 13391–13394.
34. Huebers, H.A.; Huebers, E.; Csiba, E.; Finch, C.A. Iron uptake from rat plasma transferrin by rat reticulocytes. *J. Clin. Investig.* **1978**, *62*, 944–951. <https://doi.org/10.1172/JCI109223>.
35. Kontoghiorghes, G.J.; Jackson, M.J.; Lunec, J. In Vitro Screening of Iron Chelators Using Models of Free Radical Damage. *Free. Radic. Res. Commun.* **1986**, *2*, 115–124.
36. Timoshnikov, V.A.; Selyutina, O.Y.; Polyakov, N.E.; Didichenko, V.; Kontoghiorghes, G.J. Mechanistic Insights of Chelator Complexes with Essential Transition Metals: Antioxidant/Pro-Oxidant Activity and Applications in Medicine. *Int. J. Mol. Sci.* **2022**, *23*, 1247. <https://doi.org/10.3390/>
37. Timoshnikov, V.A.; Kichigina, L.A.; Selyutina, O.Y.; Polyakov, N.E.; Kontoghiorghes, G.J. Antioxidant Activity of Deferasirox and Its Metal Complexes in Model Systems of Oxidative Damage: Comparison with Deferiprone. *Molecules* **2021**, *26*, 5064. <https://doi.org/10.3390/molecules26165064>.
38. Kontoghiorghes, G.J. Prospects for introducing deferiprone as potent pharmaceutical antioxidant. *Front. Biosci.* **2009**, *1*, 161–178.
39. Korkina, L.G.; Afanas'ev, I.B. Antioxidant and chelating properties of flavonoids. *Adv. Pharmacol.* **1997**, *38*, 151–163.
40. Kontoghiorghes, C.N.; Kolnagou, A.; Kontoghiorghes, G.J. Antioxidant targeting by deferiprone in diseases related to oxidative damage. *Front. Biosci.* **2014**, *19*, 862–885.
41. Klebanoff, S.J.; Waltersdorff, A.M. Prooxidant activity of transferrin and lactoferrin. *J. Exp. Med.* **1990**, *172*, 1293–1303. <https://doi.org/10.1084/jem.172.5.1293>.
42. Reeder, B.J.; Wilson, M.T. Hemoglobin and myoglobin associated oxidative stress: From molecular mechanisms to disease States. *Curr. Med. Chem.* **2005**, *12*, 2741–2751. <https://doi.org/10.2174/092986705774463021>.
43. Chen, C.; Wang, S.; Liu, P. Deferoxamine Enhanced Mitochondrial Iron Accumulation and Promoted Cell Migration in Triple-Negative MDA-MB-231 Breast Cancer Cells Via a ROS-Dependent Mechanism. *Int. J. Mol. Sci.* **2019**, *20*, 4952. <https://doi.org/10.3390/ijms20194952>.
44. Kontoghiorghes, G.J.; Piga, A.; Hoffbrand, A.V. Cytotoxic and DNA-inhibitory effects of iron chelators on human leukaemic cell lines. *Hematol. Oncol.* **1986**, *4*, 195–204.
45. Park, K.C.; Fouani, L.; Jansson, P.J.; Wooi, D.; Sahni, S.; Lane, D.J.; Palanimuthu, D.; Lok, H.C.; Kovačević, Z.; Huang, M.L.; Kalinowski, D.S.; Richardson, D.R. Copper and conquer: Copper complexes of di-2-pyridylketone thiosemicarbazones as novel anti-cancer therapeutics. *Metallomics* **2016**, *8*, 874–886.
46. Jabłońska-Trypuć, A.; Świdorski, G.; Krętowski, R.; Lewandowski, W. Newly synthesized doxorubicin complexes with selected metals-Synthesis, structure and anti-breast cancer activity. *Molecules* **2017**, *22*, 1106. <https://doi.org/10.3390/molecules22071106>.
47. Katsarou, A.; Pantopoulos, K. Basics and principles of cellular and systemic iron homeostasis. *Mol. Aspects Med.* **2020**, *75*, 100866.
48. Pantopoulos, K. TFR2 links iron metabolism and erythropoiesis. *Blood* **2015**, *125*, 1055–1006.
49. Kontoghiorghes, G.J. Structure/red blood cell permeability. Activity of iron(III) chelator complexes. *Inorg. Chim. Acta* **1988**, *151*, 101–106. [https://doi.org/10.1016/S0020-1693\(00\)91889-9](https://doi.org/10.1016/S0020-1693(00)91889-9).
50. Forsbeck, K.; Nilsson, K.; Kontoghiorghes, G.J. Variation in iron accumulation, transferrin membrane binding and DNA synthesis in the K-562 and U-937 cell lines induced by chelators and their iron complexes. *Eur. J. Haematol.* **1987**, *39*, 318–325. <https://doi.org/10.1111/j.1600-0609.1987.tb00776.x>.
51. Thumser, A.E.; Rashed, A.A.; Sharp, P.A.; Lodge, J.K. Ascorbate enhances iron uptake into intestinal cells through formation of a FeCl₃-ascorbate complex. *Food Chem.* **2010**, *123*, 281–285. <https://doi.org/10.1016/j.foodchem.2010.04.031>.
52. Lane, D.J.R.; Robinson, S.R.; Czerwinska, H.; Bishop, G.M.; Lawen, A. Two routes of iron accumulation in astrocytes: Ascorbate-dependent ferrous iron uptake via the divalent metal transporter (DMT1) plus an independent route for ferric iron. *Biochem. J.* **2010**, *432*, 123–132. <https://doi.org/10.1042/BJ20101317>.
53. Kontoghiorghes, G.J.; Kolnagou, A.; Demetriou, T.; Neocleous, M.; Kontoghiorghes, C.N. New Era in the Treatment of Iron Deficiency Anaemia Using Trimaltol Iron and Other Lipophilic Iron Chelator Complexes: Historical Perspectives of Discovery and Future Applications. *Int. J. Mol. Sci.* **2021**, *22*, 5546. <https://doi.org/10.3390/ijms22115546>.
54. Turnlund, J.R. Human whole-body copper metabolism. *Am. J. Clin. Nutr.* **1998**, *67*, 960S–964S. <https://doi.org/10.1093/ajcn/67.5.960S>.
55. Gray, L.W.; Peng, F.; Molloy, S.A.; Pendyala, V.S.; Muchenditsi, A.; Muzik, O.; Lee, J.; Kaplan, J.H.; Lutsenko, S. Urinary copper elevation in a mouse model of Wilson's disease is a regulated process to specifically decrease the hepatic copper load. *PLoS ONE* **2012**, *7*, e38327. <https://doi.org/10.1371/journal.pone.0038327>.
56. Tubek, S. Zinc supplementation or regulation of its homeostasis: Advantages and threats. *Biol. Trace Elem. Res.* **2007**, *119*, 1–9. <https://doi.org/10.1007/s12011-007-0043-7>.
57. Stefanidou, M.; Maravelias, C.; Dona, A.; Spiliopoulou, C. Zinc: A multipurpose trace element. *Arch. Toxicol.* **2006**, *80*, 1. <https://doi.org/10.1007/s00204-005-0009-5>.

58. Kontoghiorghes, G.J.; Kolnagou, A. Molecular factors and mechanisms affecting iron and other metal excretion or absorption in health and disease. The role of natural and synthetic chelators. *Curr. Med. Chem.* **2005**, *12*, 2695–2709.
59. Kontoghiorghes, G.J. Chelators affecting iron absorption in mice. *Arzneimittelforschung* **1990**, *40*, 1332–1335.
60. Lucarelli, G.; Angelucci, E.; Giardini, C.; Baronciani, D.; Galimberti, M.; Polchi, P.; Bartolucci, M.; Muretto, P.; Albertini, F. Fate of iron stores in thalassaemia after bone-marrow transplantation. *Lancet* **1993**, *342*, 1388–1391. [https://doi.org/10.1016/0140-6736\(93\)92753-g](https://doi.org/10.1016/0140-6736(93)92753-g).
61. Kontoghiorghes, G.J. Compliance with iron-chelation treatment after bone marrow transplantation. *Lancet* **1994**, *343*, 604–605.
62. Mertens, C.; Marques, O.; Horvat, N.K.; Simonetti, M.; Muckenthaler, M.U.; Jung, M. The Macrophage Iron Signature in Health and Disease. *Int. J. Mol. Sci.* **2021**, *22*, 8457. <https://doi.org/10.3390/ijms22168457>.
63. Recalcati, S.; Locati, M.; Cairo, G. Systemic and cellular consequences of macrophage control of iron metabolism. *Semin. Immunol.* **2012**, *24*, 393–398. <https://doi.org/10.1016/j.smim.2013.01.001>.
64. Jung, M.; Mertens, C.; Brüne, B. Macrophage iron homeostasis and polarization in the context of cancer. *Immunobiology* **2015**, *220*, 295–304. <https://doi.org/10.1016/j.imbio.2014.09.011>.
65. Nairz, M.; Schroll, A.; Demetz, E.; Tancevski, I.; Theurl, I.; Weiss, G. ‘Ride on the ferrous wheel’--the cycle of iron in macrophages in health and disease. *Immunobiology* **2015**, *220*, 280–294. <https://doi.org/10.1016/j.imbio.2014.09.010>.
66. Corna, G.; Campana, L.; Pignatti, E.; Castiglioni, A.; Tagliafico, E.; Bosurgi, L.; Campanella, A.; Brunelli, S.; Manfredi, A.A.; Apostoli, P.; et al. Polarization dictates iron handling by inflammatory and alternatively activated macrophages. *Haematologica* **2010**, *95*, 1814–1822. <https://doi.org/10.3324/haematol.2010.023879>.
67. Huang, I.; Pranata, R.; Lim, M.A.; Oehadian, A.; Alisjahbana, B. C-reactive protein, procalcitonin, D-dimer, and ferritin in severe coronavirus disease-2019: A meta-analysis. *Ther. Adv. Respir. Dis.* **2020**, *14*, 1753466620937175. <https://doi.org/10.1177/1753466620937175>.
68. Vreugdenhil, G.; Kontoghiorghes, G.J.; Van Eijk, H.G.; Swaak, A.J.G. Impaired erythropoietin responsiveness to the anemia in rheumatoid arthritis. A possible inverse relationship with iron stores and effects of the oral iron chelator 1,2-dimethyl-3-hydroxypyrid-4-one. *Clin. Exp. Rheumatol.* **1991**, *9*, 35–40.
69. Vreughtenhil, G.; Kontoghiorghes, G.J.; Van Eijk, H.G.; Swaak, A.J.G. Efficacy and safety of the oral chelator L1 in anaemic rheumatoid arthritis patients. *Lancet* **1989**, *2*, 1398–1399.
70. Brock, J.H.; Liceaga, J.; Arthur, H.M.; Kontoghiorghes, G.J. Effect of novel 1-alkyl-3-hydroxy-2-methylpyrid-4-one chelators on uptake and release of iron from macrophages. *Am. J. Hematol.* **1990**, *34*, 21–25.
71. Kontoghiorghes, C.N.; Andreou, N.; Constantinou, K.; Kontoghiorghes, G.J. World health dilemmas: Orphan and rare diseases, orphan drugs and orphan patients. *World J. Methodol.* **2014**, *4*, 163–188. <https://doi.org/10.5662/wjm.v4.i3.163>.
72. Kontoghiorghes, G.J. Differences between the European Union and United States of America in Drug Regulatory Affairs Affect Global Patient Safety Standards and Public Health Awareness: The Case of Deferasirox and Other Iron Chelating Drugs. *Medicines* **2021**, *8*, 36. <https://doi.org/10.3390/medicines8070036>.
73. Kontoghiorghes, G.J. Ethical issues and risk/benefit assessment of iron chelation therapy: advances with deferiprone/deferioxamine combinations and concerns about the safety, efficacy and costs of deferasirox. *Hemoglobin* **2008**, *32*, 1–15, doi: 10.1080/03630260701726533.
74. Lopez, A.; Cacoub, P.; Macdougall, I.C.; Peyrin-Biroulet, L. Iron deficiency anaemia. *Lancet* **2016**, *387*, 907–916. [https://doi.org/10.1016/S0140-6736\(15\)60865-0](https://doi.org/10.1016/S0140-6736(15)60865-0).
75. Pasricha, S.R.; Tye-Din, J.; Muckenthaler, M.U.; Swinkels, D.W. Iron deficiency. *Lancet* **2021**, *397*, 233–248. [https://doi.org/10.1016/S0140-6736\(20\)32594-0](https://doi.org/10.1016/S0140-6736(20)32594-0).
76. Farmaki, K.; Tzoumari, I.; Pappa, C.; Chouliaras, G.; Berdoukas, V. Normalisation of total body iron load with very intensive combined chelation reverses cardiac and endocrine complications of thalassaemia major. *Br. J. Haematol.* **2010**, *148*, 466–475. <https://doi.org/10.1111/j.1365-2141.2009.07970.x>.
77. Kolnagou, A.; Kleanthous, M.; Kontoghiorghes, G.J. Reduction of body iron stores to normal range levels in thalassaemia by using a deferiprone/deferioxamine combination and their maintenance thereafter by deferiprone monotherapy. *Eur. J. Haematol.* **2010**, *85*, 430–438.
78. Pennell, D.J. T2* magnetic resonance and myocardial iron in thalassemia. *Ann. N. Y. Acad. Sci.* **2005**, *1054*, 373–378.
79. Telfer, P.; Coen, P.G.; Christou, S.; Hadjigavriel, M.; Kolnakou, A.; Pangalou, E.; Pavlides, N.; Psiloinis, M.; Simamonian, K.; Skordos, G.; Sitarou, M.; Angastiniotis, M. Survival of medically treated thalassemia patients in Cyprus. Trends and risk factors over the period 1980–2004. *Haematologica* **2006**, *91*, 1187–1192.
80. Kontoghiorghes, C.N.; Kontoghiorghes, G.J. Efficacy and safety of iron-chelation therapy with deferioxamine, deferiprone, and deferasirox for the treatment of iron-loaded patients with non-transfusion-dependent thalassemia syndromes. *Drug Des. Devel Ther.* **2016**, *10*, 465–481.
81. Taher, A.T.; Porter, J.B.; Kattamis, A.; Viprakasit, V.; Cappellini, M.D. Comment on and author’s reply. In: Kontoghiorghes, C.N.; Kontoghiorghes, G.J. Efficacy and safety of iron-chelation therapy with deferioxamine, deferiprone, and deferasirox for the treatment of iron-loaded patients with non-transfusion-dependent thalassemia syndromes. *Drug Des. Devel Ther.* **2016**, *10*, 4073–4078.
82. Campbell, N.; Hasinoff, B. Iron supplements: A common cause of drug interactions. *Br. J. Clin. Pharmacol.* **1991**, *31*, 251–255.
83. Tarnacka, B.; Jopowicz, A.; Maślińska, M. Copper, Iron, and Manganese Toxicity in Neuropsychiatric Conditions. *Int. J. Mol. Sci.* **2021**, *22*, 7820. <https://doi.org/10.3390/ijms22157820>.

84. Sheppard, L.N.; Kontoghiorghes, G.J. Competition between deferiprone, desferrioxamine and other chelators for iron and the effect of other metals. *Arzneimittelforschung* **1993**, *43*, 659–663.
85. Iyengar, V.; Pullakhandam, R.; Nair, K.M. Dietary ligands as determinants of iron-zinc interactions at the absorptive enterocyte. *J. Food Sci.* **2010**, *75*, H260–H264.
86. Nkhili, E.; Loonis, M.; Mihai, S.; El Hajji, H.; Dangles, O. Reactivity of food phenols with iron and copper ions: binding, dioxygen activation and oxidation mechanisms. *Food Funct.* **2014**, *5*, 1186–1120.
87. Fenaux, P.; Kiladjian, J.J.; Platzbecker, U. Luspatercept for the treatment of anemia in myelodysplastic syndromes and primary myelofibrosis. *Blood* **2019**, *133*, 790–794. <https://doi.org/10.1182/blood-2018-11-876888>.
88. Schmid, H.; Jelkmann, W. Investigational therapies for renal disease-induced anemia. *Expert Opin. Investig. Drugs* **2016**, *25*, 901–916.
89. Piga, A.; Perrotta, S.; Gamberini, M.R.; Voskaridou, E.; Melpignano, A.; Filosa, A.; Caruso, V.; Pietrangelo, A.; Longo, F.; Tartaglione, I.; et al. Luspatercept improves hemoglobin levels and blood transfusion requirements in a study of patients with β -thalassemia. *Blood* **2019**, *133*, 1279–1289. <https://doi.org/10.1182/blood-2018-10-879247>.
90. Kontoghiorghes, G.J. How to manage iron toxicity in post-allogeneic hematopoietic stem cell transplantation? *Expert Rev. Hematol.* **2020**, *13*, 299–302. <https://doi.org/10.1080/17474086.2020.1719359>.
91. Kontoghiorghes, G.J.; Fetta, S.; Kontoghiorghes C.N. The need for a multi-level drug targeting strategy to curb the COVID-19 pandemic. *Front Biosci.* **2021**, *26*, 1723–1736. doi: 10.52586/5064.