



Article

Integrated Transcriptomic and Metabolomic Analyses Provide Insights into the Response of Red Clover (*Trifolium pratense* L.) to High-Temperature Stress

Wanbin Sun ¹, Weiqiang Yang ², Wei He ¹, Yuandong Xu ¹, Jie Zhou ¹, Qifan Ran ¹, Huizhen Ma ¹, Yan Peng ^{2,*} and Yan Fan ^{1,*}

¹ Chongqing Crop Germplasm Rongchang Forage Resource Bank, Chongqing Academy of Animal Sciences, Chongqing 402460, China; sunwbchina@163.com (W.S.); cqhewei1978@163.com (W.H.); 55813961@163.com (Y.X.); 18408270699@163.com (J.Z.); ranqifan@outlook.com (Q.R.); 18990019772@163.com (H.M.)

² College of Grassland Science and Technology, Sichuan Agricultural University, Chengdu 611130, China; yweiqiang2023@163.com

* Correspondence: 10657@sicau.edu.cn (Y.P.); cqxky@cqaa.cn (Y.F.)

Abstract

High-temperature stress adversely affects the yield and quality of red clover (*Trifolium pratense*). However, the molecular mechanisms underlying red clover's heat tolerance remain incompletely understood, which limits targeted genetic improvement. This study employed a multi-omics approach to compare a heat-tolerant cultivar ("HL") and a heat-sensitive cultivar ("Tp615"). Under high-temperature stress, "HL" exhibited superior physiological adaptation, showing 1.8 times higher chlorophyll retention compared to "Tp615." Transcriptome analysis identified 3104 core differentially expressed genes, including 175 transcription factors, with qRT-PCR validation confirming expression patterns consistent with the transcriptomic data. Broad-targeted metabolomics revealed 1242 differentially accumulated metabolites. Multi-omics integration highlighted the crucial role of the phenylpropanoid biosynthesis pathway in enhancing red clover's heat tolerance. This study deciphers the mechanisms underlying red clover's heat resistance and offers valuable genetic resources and a theoretical foundation for breeding stress-tolerant forage crops.

Keywords: red clover; high-temperature stress; transcriptome; metabolomic

1. Introduction

Plants are unable to escape the effects of environmental stresses due to their sessile nature. Among these stresses, high-temperature (HT) stress is one of the most prevalent abiotic factors impacting agriculture. It is well established that global warming has significantly affected crop yields, thereby posing a serious threat to global food security. Average global temperatures have been steadily rising and are projected to increase by 1.8 to 4.0 °C by the end of the 21st century [1]. This temperature increase has been shown to reduce agricultural productivity; for instance, daily temperatures exceeding 30 °C can cause corn (*Zea mays*) and soybean (*Glycine max*) yields to decline by up to 6% under rain-fed conditions [2]. Furthermore, a 1 °C rise in global temperature is associated with a 6% reduction in the harvest index of crops [3]. Therefore, it is crucial to understand the mechanisms underlying heat tolerance in plants and to develop germplasm capable of withstanding elevated temperatures to address the challenges posed by global climate change.



Academic Editor: Tibor Janda

Received: 5 December 2025

Revised: 29 January 2026

Accepted: 30 January 2026

Published: 10 February 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and

conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

When plants are exposed to HT stress, their cells accumulate significant amounts of reactive oxygen species (ROS), such as hydrogen peroxide (H_2O_2) and superoxide anion (O_2^-), which induce oxidative stress [4]. This stress can inhibit photosynthesis, damage cell membranes, accelerate aging, and ultimately lead to cell death [5]. Furthermore, high-temperature stress disrupts the energy conversion process between the oxygen-evolving complex of photosystem II (PSII) and the photosystem receptor site, resulting in irreversible damage to PSII and a substantial reduction in its activity [6]. Plants have evolved complex and diverse strategies to adapt to heat stress. At the molecular level, various heat shock proteins and heat stress transcription factors respond positively when plants are exposed to high temperatures, thereby enhancing their resistance to heat stress [7]. The accumulation of heat shock proteins (Hsps) is a crucial component of the heat stress response, and these proteins are believed to play a central role in acquired heat tolerance in plants and other organisms [8]. Previous studies have demonstrated that heat shock factors HsfA1a, HsfA1b, and HsfA2 play a crucial role in regulating the heat stress response in Arabidopsis [9–11]. Additionally, the overexpression of WRKY39 in Arabidopsis enhances the plant's heat tolerance [12]. Studies by Li et al. [13] demonstrated that the transcription factor NAC019 binds to the promoters of HSFA1b, HSFA6b, HSFA7a, and HSFC1, thereby positively regulating thermotolerance in Arabidopsis.

Gaining a deeper understanding of the mechanisms by which plants acquire heat tolerance is crucial, as it facilitates the development of heat-tolerant crops through marker-assisted breeding [14]. Although significant progress has been made in recent years in elucidating how plants sense and respond to hormones [15,16], it remains evident that many regulatory pathways remain unexplored. The plant response to high-temperature stress involves a multi-layered, dynamic network of changes encompassing gene expression, metabolic reprogramming, and physiological phenotypes [17,18]. To systematically elucidate this complex mechanism, research approaches capable of simultaneously probing multiple molecular levels are urgently required. Against this backdrop, integrated transcriptomic and metabolomic analyses have emerged as powerful tools for elucidating the core mechanisms of plant stress responses, owing to their capacity to correlate upstream transcriptional regulation with downstream metabolic phenotypes [19]. For instance, Zhou et al. [20] revealed that maintaining normal hormone synthesis and metabolism may represent a crucial pathway for enhancing alfalfa's tolerance to high temperatures. Khan et al. [21] demonstrated that the interconnected pathways of starch–sucrose metabolism and ABC transporters are pivotal for enhancing rhododendron growth under high-temperature stress. Furthermore, the accumulation of sucrose and triacylglycerols, alongside reduced membrane lipid unsaturation, was identified as a primary factor altered in wild-type tomato fruit following exposure to transient heat stress [22].

Red clover (*Trifolium pratense*) is a perennial legume widely cultivated for its resilience to cold and moisture, nutritional benefits, and status as a preferred forage crop. However, its susceptibility to elevated temperatures negatively impacts both the quality and utilization of this plant. The mechanisms that confer heat tolerance in this species remain poorly understood. To address this knowledge gap, we analyzed potted red clover plants under varying temperature stress conditions by examining their physiological, transcriptional, and metabolic responses. This study provides crucial insights into plant heat stress and is particularly vital for breeding new heat-tolerant varieties, especially within the context of climate change.

2. Results

2.1. Differences in Physiological Responses of Two Red Clover Accessions to High-Temperature Stress

Phenotypic changes indicated that HT stress caused wilting and yellowing in red clover leaves of both accessions; however, Tp615 suffered more severe damage than HL (Figure 1A). The electrolyte leakage (EL) and malonaldehyde acid (MDA) levels of both accessions increased significantly, but HL maintained lower EL and MDA values. Similarly, the O_2^- and H_2O_2 contents showed comparable trends (Figure 1B–E), indicating that oxidative damage occurred under HT stress. Under normal conditions, no significant differences were observed in chlorophyll content, as well as in the photochemical efficiency (Fv/Fm) and the photosynthetic performance index (PI_{ABS}) between the two materials. HT stress led to a significant decrease in chlorophyll content in both accessions, along with reductions in PI_{ABS} and Fv/Fm values. This suggests that HT stress caused chlorophyll degradation in red clover leaves and adversely affected their health status. Furthermore, the chlorophyll content of HL was 1.8 times higher than that of Tp615 under HT stress, and the PI_{ABS} value of HL was superior to that of Tp615 (Figure 1F–H). In conclusion, HL exhibited less oxidative damage under HT stress, maintained higher chlorophyll content, and was able to sustain higher photosynthetic activity.




Figure 1. Comparison of physiological indices between two different accessions, HL and Tp615, under normal (CK) and high-temperature (HT) stress conditions. (A) Leaf phenotype; (B) electrolyte osmolality (EL); (C) malondialdehyde (MDA) content; (D) hydrogen peroxide content; (E) superoxide anion content; (F) chlorophyll content; (G) chlorophyll fluorescence parameters; and (H) health index value (Pi). Bars represent the means of four biological replicates $\pm$ standard error (SE), and different letters (a–c) indicate significant differences ($p < 0.05$) as determined by Duncan's multiple range test.

2.2. Transcriptome Quality Control and Differentially Expressed Genes (DEGs) Analysis

To gain a comprehensive understanding of the gene regulatory networks in two red clover germplasms, RNA sequencing (RNA-Seq) was performed on red clover leaves exposed to high temperatures for 0, 6, and 12 h, respectively. The base quality Q30 (the proportion of bases with a sequencing error rate of $\leq 0.1\%$) for all RNA-Seq libraries exceeded 92% (Supplementary Table S1). Correlation coefficient heatmaps and principal component analysis (PCA) confirmed that the three biological replicates for each treatment clustered effectively, demonstrating clear separation between the two red clover accessions (Figure 2A,B). Using the 0-h sample as a control, a total of 8778 DEGs were identified in HL after 6 h (4170 upregulated and 4608 downregulated) and 9481 DEGs after 12 h (6471 upregulated and 3010 downregulated) of high-temperature treatment. In Tp615, 9265 DEGs were detected after 6 h (5903 upregulated and 3362 downregulated) and 8896 DEGs after 12 h (5590 upregulated and 3306 downregulated) (Figure 2C).




Figure 2. Transcriptome analysis of HL and Tp615 red clover under high-temperature (HT) treatment at various time points. **(A)** Principal Component Analysis (PCA). **(B)** Sample correlation heatmap. **(C)** Number of DEGs identified between HL and Tp615 following HT treatment (CK: 0 h).

2.3. Differential Gene Enrichment Analysis

The Gene Ontology (GO) functional annotation analysis revealed that the DEGs were predominantly associated with detoxification and nitrogen utilization responses under varying durations of stress, with significant enrichment within the biological process (BP) category. In the molecular function (MF) category, notable enrichments were observed in catalytic activity, transporter activity, antioxidant activity, and ATP-dependent activity. Furthermore, the cellular component (CC) category showed significant enrichment in cellular anatomical entities and protein-containing complexes. The metabolic categories of DEGs in the two red clover germplasms exhibited distinct differences when subjected to high-temperature stress (Figure 3A–D; Supplemental Table S2), indicating variability in both the number of DEGs and their associated biological functions.

Red clover-specific and time-independent DEGs were further analyzed, revealing 5764 common DEGs identified in HL and 6411 common DEGs identified in Tp615 during the HT treatment. Cross-comparison of these time-independent DEGs between the red clover species indicated that 2660 and 3307 DEGs were specific to HL and Tp615, respectively (Figure 3E; Supplemental Figure S1). KEGG enrichment analyses demonstrated that these species-specific DEGs were predominantly associated with the biosynthesis of secondary metabolites, as well as starch and sucrose metabolism (Figure 3E; Supplemental Figure S2A,B; Supplemental Table S3). The remaining time-independent DEGs (3104) were primarily identified as genes related to metabolic pathways, linoleic acid metabolism, and

protein processing in the endoplasmic reticulum (Figure 3E; Supplemental Figure S2C; Supplemental Table S3).




Figure 3. DEGs were analyzed for functional enrichment. (A–D) GO enrichment circular plots for HL and Tp615 under high-temperature treatments at 6 and 12 h. The GO enrichment circular plots consist of the following: (A): GO terms in the top 20 for enrichment, with the number of differential genes represented outside the circle as a scaling factor. Different colors indicate different ontologies; (B): the number of GO terms in the differential gene background and the Q value. A longer bar indicates a greater number of differential genes in the background, while a smaller Q value corresponds to a redder color; (C): the ratio of up- and down-regulated differential genes, where dark purple represents the up-regulated gene ratio and light purple represents the down-regulated gene ratio; specific values are displayed at the bottom; (D): the Rich Factor value for each GO term (calculated as the number of differential genes in the GO term divided by the total number of genes in that GO term). The background grid lines indicate the number of differential genes in the background, with each cell representing 0.1. (E) Venn diagrams illustrating the DEGs across treatments, highlighting unique and common pathways of DEG enrichment between red clover species and treatment durations.

We further compared the species-independent effects of HT treatment duration on gene expression (Figure 3E; Supplemental Figure S3; Supplemental Table S4) and identified 4655 common DEGs after 6 h and 5487 common DEGs after 12 h. Cross-comparison between the 6-h and 12-h data revealed 1551 time-dependent DEGs unique to the 6-h treatment. KEGG enrichment analysis indicated that these DEGs were primarily associated with the biosynthesis of secondary metabolites and isoflavonoid biosynthesis (Figure 3E; Supplemental Figure S4A; Supplemental Table S4). The 12-h-specific time-dependent DEGs (2383) were predominantly linked to starch and sucrose metabolism (Figure 3E; Supplemental Figure S4B; Supplemental Table S4). In both red clover accessions, HT treatments induced genes related to the biosynthesis of secondary metabolites and starch and sucrose metabolism in a time-dependent manner, whereas the induction of genes related to protein processing in the endoplasmic reticulum occurred in a time-independent manner.

2.4. Transcription Factor Gene Expression Analysis

To further analyze the DEGs between HL and Tp615 red clover germplasm, a total of 3014 DEGs were identified at both stages of HT stress (Figure 3B). These genes may represent core components involved in the red clover's response to HT stress. Additionally, 175 transcripts encoding transcription factors were identified (Supplemental Table S4). Among the differentially expressed TFs, the bHLH, ERF, MYB, Dof, NAC, WRKY, and HSF families accounted for more than 50% of the total. Notably, eight bHLH genes were upregulated after heat stress (Figure 4A). Furthermore, three HSFs (ncbi_123894233, ncbi_123881398, ncbi_123907261) were significantly upregulated in both accessions following heat stress (Figure 4B). Most ERFs exhibited a tendency to be downregulated after heat stress; however, ERF3 (ncbi_123918970) was upregulated, and ncbi_123923905 was upregulated in the Tp615 material but downregulated in HL. Additionally, eight MYB, two NAC, and three WRKY genes were upregulated in both accessions after heat stress. A comprehensive list of these DEGs can be found in Supplemental Table S5. To validate the RNA-Seq data, six DEGs were selected for real-time PCR analysis (Figure 4C). The results from RT-qPCR and RNA-Seq demonstrated consistent expression patterns. Therefore, these transcription factors may play a crucial role in the heat stress response of red clover.




Figure 4. Analysis of Differentially Expressed TFs. (A) Heatmap illustrating the expression of bHLH family genes. (B) Heatmap depicting the expression of HSF family genes. (C) Heatmap representing

the expression of ERF family genes. **(D)** The expression profile of the selected DEGs was determined using RT-qPCR analysis, and the lines indicate the relative gene transcription levels obtained from RNA-Seq. The left y-axis represents the RT-qPCR values, while the right y-axis represents the RNA-Seq values. Data are presented as the SE $\pm$ mean of three biological replicates.

2.5. Metabolomics Measurement and Analysis

2.5.1. QC of Metabolomics Data

Heatmaps generated from normalized data (Figure 5A) reveal significant metabolic differences between the two germplasm lines under high-temperature stress, primarily characterized by marked alterations in metabolite levels. Metabolites from each sample were analyzed using PCA and OPLS-DA. Results showed that samples from different treatments formed distinct clusters, demonstrating good reproducibility and intra-group aggregation. The R^2 and Q^2 values for all control groups across different stages ranged between 0.7 and 1 (Figure 5B,C), indicating that the model possesses good predictive capability and high data reliability (Supplemental Table S6).




Figure 5. Metabolomic analysis. **(A)** clustering heatmap analysis of the DEMs. **(B)** Principal component analysis (PCA) of DEMs, with QC indicating quality control samples prepared by mixing equal amounts of experimental samples. **(C)** OPLS-DA score map of metabolome data.

2.5.2. Statistical Analysis of DEMs

Analysis of metabolome sequencing data identified a total of 1242 metabolites (Figure 6A, Table S7). During the HT stress treatment, 128 differentially expressed metabolites (DEMs) were identified in the heat-tolerant accessions HL, with 72 significantly upregulated and 56 downregulated. In contrast, the heat-sensitive accessions TP615 exhibited

150 DEMs (86 upregulated, 64 downregulated). Notably, 95 DEMs were common to both accessions (Figure 6B). These shared metabolites—including amino acids, flavonoids, carbohydrates, organic acids, and their respective derivatives—may play crucial roles in the response of red clover to HT stress.




Figure 6. Statistical analysis of DEMs. (A) Wayne diagram of differential metabolites. (B) Analysis of common differential metabolites in two accessions.

2.5.3. KEGG Pathway Enrichment Analysis of DEMs

To better understand the metabolic pathways crucial for red clover's response to HT stress, we performed KEGG pathway enrichment analysis of DEMs in two red clover accessions with differing heat tolerance. Results indicated significant enrichment in Glyoxylate and dicarboxylate metabolism and Biosynthesis of alkaloids derived from ornithine. Furthermore, the Glyoxylate and dicarboxylate metabolism pathway contained 19 metabolites, while the Biosynthesis of plant secondary metabolites pathway contained 67 metabolites significantly enriched in the HL germplasm (Figure 7A). Conversely, the Aminoacyl-tRNA biosynthesis pathway contained 21 metabolites, and the Biosynthesis of plant hormones pathway contained 32 metabolites significantly enriched in the TP615 germplasm (Figure 7B). Additionally, enrichment analysis of the 95 metabolites common to both accessions (Figure 7C) showed significant enrichment of 52 metabolites in the Biosynthesis of phenylpropanoids pathway and 19 metabolites in the Phenylpropanoid biosynthesis pathway. These findings indicate that these metabolic pathways play a crucial role in red clover's resistance to HT stress.

2.6. Integrated Analysis of Transcriptomics and Metabolomics

2.6.1. Correlation Analysis Between DEGs and DEMs

Spearman's correlation analysis was employed to investigate the relationship between DEGs and DEMs under high-temperature stress. The results revealed that numerous DEGs and DEMs exhibited strong positive correlations (Figure S5), indicating potential regulatory relationships.

2.6.2. Enrichment Analysis of KEGG Pathways for DEGs and DEMs

Based on transcriptomic and metabolomic analyses, a KEGG pathway co-enrichment analysis was performed for differentially expressed genes (DEGs) and differentially accumulated metabolites (DEMs) (Figure 8A,B). Notably, biosynthesis of secondary metabolites, metabolic pathways, phenylpropanoid biosynthesis, and galactose metabolism were significantly co-enriched in both germplasm sets. Within the phenylpropanoid biosynthesis pathway, significant correlations were observed for ^L-phenylalanine, hydroxycinnamate, ferulic acid, *p*-coumaric acid, ferulate, *p*-hydroxycinnamic acid, and *p*-coumaryl alcohol. Further analysis revealed that the 37 metabolites shared between HL and TP615 were predominantly enriched in pathways including phenylalanine metabolism, phenylpropanoid biosynthesis, and plant hormone biosynthesis. These results indicate that these metabolites play a crucial role in red clover responses to high-temperature (HT) stress (Figure 8C,D).




Figure 7. KEGG pathway enrichment analysis of DEMs. (A) KEGG enrichment analysis of differential metabolites in HL accessions. (B) KEGG enrichment analysis of differential metabolites in TP615 accessions. (C) KEGG enrichment analysis of common differential metabolites in the two accessions.




Figure 8. Enrichment analysis of KEGG pathways for DEGs and DEMs. (A) Co-enrichment analysis of DEGs and DEMs KEGG pathways in HL accessions. (B) Co-enrichment analysis of DEGs and DEMs KEGG pathways in TP615 accessions. (C) Metabolite Venn diagram analysis. (D) KEGG enrichment analysis of 37 core differential metabolites.

<https://doi.org/10.3390/ijms27041704>

A structural gene–metabolite association network for phenylpropanoid biosynthesis was constructed (Figure 9B). Twenty-seven key genes exhibited 29 significant association nodes ($p < 0.05$) with nine metabolites. Among the phenylpropanoid compounds, NNK (Com_271_pos), ferulic acid (Com_664_neg), and ferulate (Com_922_neg) were significantly associated with 10, 7, and 9 structural genes, respectively. Core structural genes included ncbi_123921203, ncbi_123919763, and ncbi_123907091. Together, these genes and their interactions form the molecular regulatory network for phenylpropanoid biosynthesis in red clover, providing key candidate genes for elucidating the mechanism of its response to HT stress.

3. Discussion

Under high-temperature stress, both red clover leaf types exhibited significantly elevated levels of electrolyte leakage (EL), H_2O_2 , O_2^- , and MDA (Figure 1B–E). Notably, Tp615 showed markedly higher H_2O_2 and MDA levels compared to the other genotype, indicating more severe oxidative damage to its cell membranes. Consistent with previous studies in rice, maize, and wheat, high-temperature stress typically elevates MDA and H_2O_2 levels, thereby exacerbating membrane lipid peroxidation and cellular damage [23–26]. The high-temperature treatment also significantly reduced chlorophyll content and PI_{ABS} , with Tp615 exhibiting a greater decline. This decline in photosynthetic pigments is closely linked to excessive ROS accumulation exceeding antioxidant enzyme scavenging capacity, ultimately disrupting photosynthetic structures and photosystem function [27]. Collectively, these findings indicate that Tp615 exhibits greater oxidative stress and photosynthetic damage under high-temperature stress, while the other variety demonstrates relatively higher tolerance. This provides a foundation for subsequent analysis of heat tolerance mechanisms in different germplasm.

Transcription factors play a pivotal role in plant growth, development, and stress responses. In plant heat responses, transcription factors typically function synergistically within regulatory networks rather than regulating individual target genes in isolation [28]. In this study, distinct transcription factor families exhibited markedly differential expression patterns under heat stress, suggesting that they may perform distinct roles in the regulation of heat tolerance in red clover. Basic helix–loop–helix (bHLH) family transcription factors constitute one of the largest transcription factor gene families in plants [29]. bHLH transcription factors not only participate in environmental signal perception but also play a crucial integrative role in coordinating growth and stress responses, as reported in species such as apple, rice, and cucumber [30–32]. In this study, eight members of the bHLH family were significantly induced following high-temperature treatment. These findings suggest that red clover may rapidly modulate transcriptional networks to counteract physiological disturbances caused by elevated temperatures through the induction of specific bHLH factors. In contrast to the bHLH family, most ERF transcription factors exhibited a downregulation trend after heat stress, with only ERF3 being significantly up-regulated. ERFs typically serve as key nodes in ethylene and ROS signaling pathways; thus, their suppression may help reduce unnecessary defense responses under high-temperature conditions, thereby conserving energy [33]. The specific upregulation of ERF3 suggests that it may play a crucial role in maintaining basal stress signaling or cellular homeostasis. Previous studies have shown that ERF74/77 and ERF108/125 enhance rice heat tolerance by synergistically regulating the expression of common target genes and specific heat-responsive genes [34]. This differential regulation indicates that ERF3 in red clover may precisely modulate downstream target genes in a stress-specific manner under heat stress. Furthermore, MYB family transcription factors play crucial roles in plant responses to high temperatures [35,36]. Zhao et al. [37] demonstrated that overexpression of *TaMYB80* in

transgenic *Arabidopsis* enhances plant tolerance to combined heat and drought stress. In this study, eight MYB genes were significantly upregulated under HT stress, suggesting that these MYB factors may promote sustained heat tolerance in red clover by regulating genes involved in antioxidant defense, secondary metabolism, or osmotic adjustment. Heat shock factors (HSFs) serve as central regulators in plant responses to thermal stress. Upon exposure to high temperatures, HSFs bind to cis-regulatory elements known as Heat Shock Elements (HSEs) in the promoters of downstream genes, thereby activating the expression of heat-stress-related genes such as heat shock proteins (HSPs) [38]. In *Arabidopsis*, HSFA1 activates a broad range of target genes, including multiple HSPs, thereby positively regulating heat tolerance. Additionally, HSFA2 can be induced by HSFA1 and plays a specific role in heat stress memory [39]. In this study, three HSF genes were significantly induced following HT treatment, and qPCR results further validated their expression changes, supporting their crucial roles in activating heat shock proteins and maintaining protein homeostasis (Figure 4).

The phenylpropanoid biosynthetic pathway serves as a common precursor for multiple key secondary metabolites, including flavonoids and lignin. Its role in responses to abiotic stress exhibits remarkable functional diversity and regulatory complexity [40]. Previous studies indicate that flavonoids, as potent natural antioxidants, can directly scavenge excess reactive oxygen species (ROS) accumulated under HT stress, thereby mitigating membrane lipid peroxidation and cellular oxidative damage [41,42]. However, activation of the phenylpropanoid pathway extends beyond antioxidant defense and may involve metabolic flux redistribution among different branches. As a crucial downstream branch of this pathway, lignin biosynthesis plays a unique structural protective role under HT stress [43]. Elevated temperatures often lead to cell wall relaxation and reduced structural stability. Enhanced phenylpropanoid pathway activity may promote lignin deposition, thereby increasing cell wall mechanical strength, maintaining cell shape stability, and limiting electrolyte leakage [44]. Phenylalanine ammonia-lyase (PAL), a key and rate-limiting enzyme in phenylalanine metabolism, responds to high-temperature stress by promoting the synthesis of total phenolic compounds and flavonoids [45,46]. Furthermore, studies have shown that PAL and other genes associated with lignin biosynthesis are significantly upregulated under HT stress in wheat, suggesting that activation of these pathways may contribute to improved heat tolerance [47]. In this study, the significant upregulation of genes related to lignin monomer synthesis (e.g., CAD and CCR) in heat-tolerant materials suggests that phenylpropanoid metabolism may preferentially shift toward the lignin branch rather than solely increasing soluble phenolic compounds. This result supports a defense strategy in heat-tolerant materials that counteracts heat-induced cell wall damage, thereby enhancing heat tolerance at the tissue level. Furthermore, mounting evidence indicates that certain intermediate metabolites in the phenylpropanoid pathway (e.g., cinnamic acid derivatives) function not only as metabolic precursors but also as signaling molecules involved in systemic acquired resistance (SAR) and the amplification and integration of stress response signals [48,49]. Therefore, activation of the phenylpropanoid pathway in heat-tolerant materials may exert dual roles in metabolic defense and signal regulation. By enhancing antioxidant capacity, reinforcing cellular structure, and coordinating stress response networks, this pathway collectively promotes overall plant adaptability to HT stress.

4. Materials and Methods

4.1. Plant Material and Treatments

This study was conducted at Sichuan Agricultural University during 2024. The plant materials included one commercial variety, ‘Red Dragon’ (abbreviated as HL), and one

wild accession, Tp615, which was collected from Sichuan Province, China. Seeds were germinated in Petri dishes containing two layers of filter paper and 25 mL of distilled water for 8 days. After germination, seedlings were transplanted into pots (13.3 cm in diameter and 5.6 cm in height) filled with a mixed-nutrient soil (nutrient soil/vermiculite/perlite = 8:1:1). Each pot contained 15 seedlings, and all pots were placed in a plant growth chamber (Jiangnan Instrument Factory, Ningbo, China; GHP-500) for 40 days. At this time, the plants had reached the late vegetative stage, characterized by the presence of 20–30 fully expanded trifoliate leaves and the absence of visible flower buds. To simulate high-temperature stress, the day and night temperatures in the plant growth chamber were adjusted to 38/35 °C for 18 days for the treatment group. Under normal conditions, the day/night temperature was maintained at 25/18 °C, with a 12-h photoperiod, 750 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation (PAR), and 65% relative humidity. During this period, leaf tissue samples were harvested from both the control group (normal conditions) and the treatment group (following 18 days of high-temperature stress). For each treatment, the second and third fully expanded trifoliate leaves were randomly selected from the apex of the main stems of all 15 individual seedlings per pot. Each treatment comprised four biological replicates, with samples within each replicate pooled to ensure representativeness. The collected tissues were immediately flash-frozen in liquid nitrogen and subsequently stored at $-80\text{ }^{\circ}\text{C}$ for further physiological parameter measurements.

4.2. Measurement of Physiological Parameters

To determine the EL value, we immersed 0.1 g of fresh leaves (collected at the late vegetative growth stage) in 20 mL of distilled water and shook the mixture at 25 °C for 24 h. The initial conductivity (C_i) of the extracts was measured using a conductivity meter (DDS-307; Lei-Ci, Shanghai, China). Subsequently, the extracts were sterilized at 121 °C for 20 min, and the maximum conductivity (C_{max}) was measured after the extracts were cooled to room temperature. The values of EL were calculated using the formula $\text{EL} = (C_i/C_{\text{max}}) \times 100\%$. To determine the MDA content, fresh leaf tissue (0.1 g) was ground in 2 mL of 50 mM cold phosphate-buffered saline (pH 7.8). The mixture was then centrifuged at 12,000 rpm for 30 min at 4 °C. Next, 1 mL of the reaction solution (composed of 0.5% thiobarbituric acid and 20% trichloroacetic acid) was mixed with 0.5 mL of the supernatant. This mixture was heated at 95 °C for 15 min, then rapidly cooled in ice water and centrifuged at 8000 rpm for 10 min. The absorbance of the supernatant was measured at 532 nm and 600 nm [50]. O_2^- and H_2O_2 were detected using the methods described by Elstner et al. [51] and Velikova et al. [52], respectively. For the determination of chlorophyll content, leaves were immersed in 15 mL of dimethyl sulfoxide and allowed to stand in the dark for 48 h until they became transparent. The absorbance of the extracts was then measured at 645 and 663 nm using a spectrophotometer (Spectronic Instruments, Rochester, NY, USA) [53]. To assess the F_v/F_m and the PI_{ABS} of PS II, a chlorophyll fluorometer (Pocket PEA, Hansatech, Norfolk, UK) was utilized. PI_{ABS} serves as a crucial indicator of the total number of PS II-activated photochemical reaction centers, reflecting transient photosynthetic performance and plant health under environmental stress. F_v/F_m and PI_{ABS} values were recorded after placing monolayer leaves in the instrumented leaf chamber for 20 min of dark adaptation.

4.3. Transcriptome Analysis

Leaf tissues were collected at 0 h, 6 h, and 12 h of high-temperature treatment, wrapped in aluminum foil, and immediately frozen in liquid nitrogen. Each treatment included three biological replicates. Total RNA was extracted, followed by library construction and Illumina sequencing at Genedenovo Biotechnology Co., Ltd. (Guangzhou, China). Raw

sequencing reads were processed using fastp for quality control, aligned to the reference genome with HISAT2 [54], assembled into transcripts by String Tie [55], and quantified for gene expression using RSEM [56]. Utilizing the expression data, we performed principal component analysis in R (<http://www.r-project.org/> (accessed on 31 May 2024)) and calculated Pearson correlation coefficients to generate heat maps. Differentially expressed genes were analyzed using DESeq2 software (version 1.40.2) with the standard criteria of $|\log_2(\text{fold change})| > 1$ and adjusted for significance with a p -value < 0.05 [57,58]. Gene Ontology (GO) enrichment analysis was conducted by mapping the differentially expressed genes to the GO database (<http://www.geneontology.org/> (accessed on 31 May 2024)). This involved associating the differential genes with each term in the GO database and calculating the number of differentially expressed genes associated with each term, thereby obtaining statistics on the number of differential genes linked to specific GO functions. Pathway significance enrichment analysis was performed using a hypergeometric test to determine the number of differentially expressed genes within the entire background compared to the pathways that were significantly enriched. The pathways were identified as significantly enriched in differentially expressed genes by applying a hypergeometric test to the KEGG Pathway, with a corrected p -value < 0.05 set as the threshold for significance.

4.4. Extensive Targeted Metabolomics Analysis and Data Analysis

Leaf samples were collected under normal conditions and after 18 days of high-temperature stress, promptly frozen in liquid nitrogen and stored for subsequent metabolite extraction. Metabolomic profiling was conducted in collaboration with Gene Denovo Biotechnology Co., Ltd. (Guangzhou, China) using mass spectrometry platforms. The acquired metabolomic data underwent quality control using QC samples [59], with PCA analysis performed using the R package gmodels (v2.18.1) [60]. All metabolites were subjected to z-score standardization using the R package pheatmap (v2.18.1) [61] prior to cluster analysis. OPLS-DA analysis was conducted using the R package ropls [62]. Differential metabolites between comparison groups were identified by combining the VIP values from multivariate OPLS-DA analysis with the p -values from univariate t -tests [63]. The differential threshold was set as: $\text{VIP} \geq 1$ in the OPLS-DA model and t -test $p < 0.05$.

4.5. RT-qPCR Analysis

First-strand cDNA was synthesized using the RevertAid First Strand cDNA Synthesis Kit (All-in-One First-Strand Synthesis MasterMix with dsDNase, iScience, Shanghai, China). Quantitative reverse transcription PCR (RT-qPCR) was performed following the manufacturer's protocol with Taq SYBR[®] Green qPCR Premix (Universal; iScience, Shanghai, China) to validate transcript abundance of the candidate genes. Primer sequences are provided in Supplemental Table S8.

4.6. Statistical Analysis

One-way ANOVA followed by Tukey's Honestly Significant Difference (HSD) test for multiple comparisons was used to analyze the data using SPSS v.25 (IBM, Armonk, NY, USA). Data were examined for normality and homogeneity of variance prior to analysis. Correlation graphs were generated using GraphPad Prism 9. Unless otherwise noted, all experiments were carried out with four biological replicates, and the data are reported as mean $\pm$ standard deviation (SD). Statistical significance was defined as $p < 0.05$.

5. Conclusions

This study elucidates the distinct physiological and multi-omic mechanisms underlying heat stress adaptation in red clover. Our results demonstrate that the heat-tolerant cultivar 'HL' maintains significantly higher membrane stability and approximately 1.8-fold

greater chlorophyll retention compared with the sensitive cultivar ‘Tp615’. Integrated transcriptomic and metabolomic analyses reveal robust systemic reprogramming, identifying 3104 core differentially expressed genes (DEGs) and 1242 differentially accumulated metabolites (DAMs). Notably, the phenylpropanoid biosynthetic pathway emerged as a central regulatory hub, where the coordinated enrichment of key transcripts and metabolites plays a role in mitigating thermal proteotoxicity. These findings highlight specific metabolic checkpoints and candidate genetic loci, providing a solid theoretical foundation and valuable genetic resources for the genomic-assisted breeding of thermotolerant red clover cultivars.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijms27041704/s1>.

Author Contributions: W.S. conducted the experiments; W.Y. authored the original manuscript; W.H. and Y.X. reviewed and edited the manuscript; J.Z., H.M. and Q.R. created the diagrams; Y.F. and Y.P. conceived and designed the experiments and reviewed and edited the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Research Institute Performance Incentive and Guidance Project of the Chongqing Science and Technology Bureau (No. CSTC2022JXJL80013), the Development and Breeding Utilization of High Quality Protein Forage in Southern China (No. CSTB2023JXJL-YFX0034), and the Sichuan Forage Innovation Team Program (No. SCCXTD-2025-16).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The raw sequencing data generated in this study have been deposited in the BIG Submission (BIG Sub) database of the National Genomics Data Center (NGDC) under accession number CRA030495 (<https://ngdc.cncb.ac.cn/gsa/s/o4ZV5jsF>, accessed on 31 June 2025).

Acknowledgments: We extend our gratitude to the Chongqing Academy of Animal Sciences and Sichuan Agricultural University for their support in conducting this research, as well as to all contributors of this article.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Bita, C.E.; Gerats, T.J. Plant tolerance to high temperature in a changing environment: Scientific fundamentals and production of heat stress-tolerant crops. *Front. Plant Sci.* **2013**, *4*, 273. [\[CrossRef\]](#)
2. Schauburger, B.; Archontoulis, S.; Arneth, A.; Balkovic, J.; Ciais, P.; Deryng, D.; Elliott, J.; Folberth, C.; Khabarov, N.; Müller, C.; et al. Consistent negative response of US crops to high temperatures in observations and crop models. *Nat. Commun.* **2017**, *8*, 13931. [\[CrossRef\]](#)
3. Mishra, D.; Shekhar, S.; Agrawal, L.; Chakraborty, S.; Chakraborty, N. Cultivar-specific high temperature stress responses in bread wheat (*Triticum aestivum* L.) associated with physicochemical traits and defense pathways. *Food Chem.* **2017**, *221*, 1077–1087. [\[CrossRef\]](#)
4. Cheng, B.; Zhou, M.; Tang, T.; Hassan, M.J.; Zhou, J.; Tan, M.; Li, Z.; Peng, Y. A *Trifolium repens* flavodoxin-like quinone reductase 1 (*TrFQR1*) improves plant adaptability to high temperature associated with oxidative homeostasis and lipids remodeling. *Plant J.* **2023**, *115*, 369–385. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Priya, M.; Farooq, M.; Siddique, K.H.M. Enhancing tolerance to combined heat and drought stress in cool-season grain legumes: Mechanisms, genetic insights, and future directions. *Plant Cell Environ.* **2025**, eraf237. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Xu, T.; Zhou, H.; Feng, J.; Guo, M.; Huang, H.; Yang, P.; Zhou, J. Involvement of HSP70 in BAG9-mediated thermotolerance in *Solanum lycopersicum*. *Plant Physiol. Biochem.* **2024**, *207*, 108353. [\[CrossRef\]](#)
7. Ohama, N.; Sato, H.; Shinozaki, K.; Yamaguchi-Shinozaki, K. Transcriptional regulatory network of plant heat stress response. *Trends Plant Sci.* **2017**, *22*, 53–65. [\[CrossRef\]](#)

8. Baniwal, S.K.; Bharti, K.; Chan, K.Y.; Fauth, M.; Ganguli, A.; Kotak, S.; Mishra, S.K.; Nover, L.; Port, M.; Scharf, K.-D.; et al. Heat stress response in plants: A complex game with chaperones and more than twenty heat stress transcription factors. *J. Biosci.* **2004**, *29*, 471–487. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Busch, W.; Wunderlich, M.; Schöffl, F. Identification of novel heat shock factor-dependent genes and biochemical pathways in *Arabidopsis thaliana*. *Plant J.* **2005**, *41*, 1–14. [\[CrossRef\]](#)
10. Charng, Y.-Y.; Liu, H.-C.; Liu, N.-Y.; Chi, W.-T.; Wang, C.-N.; Chang, S.-H.; Wang, T.-T. A heat-inducible transcription factor, HsfA2, is required for extension of acquired thermotolerance in *Arabidopsis*. *Plant Physiol.* **2007**, *143*, 251–262. [\[CrossRef\]](#)
11. Lohmann, C.; Eggers-Schumacher, G.; Wunderlich, M.; Schöffl, F. Two different heat shock transcription factors regulate immediate early expression of stress genes in *Arabidopsis*. *Mol. Genet. Genomics* **2004**, *271*, 11–21. [\[CrossRef\]](#)
12. Nishizawa, A.; Yabuta, Y.; Yoshida, E.; Maruta, T.; Yoshimura, K.; Shigeoka, S. *Arabidopsis* heat shock transcription factor A2 as a key regulator in response to several types of environmental stress. *Plant J.* **2006**, *48*, 535–547. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Li, S.; Zhou, X.; Chen, L.; Huang, W.; Yu, D. Functional characterization of *Arabidopsis thaliana* WRKY39 in heat stress. *Mol. Cells* **2010**, *29*, 475–483. [\[CrossRef\]](#)
14. Li, B.; Gao, K.; Ren, H.; Tang, W. Molecular mechanisms governing plant responses to high temperatures. *J. Integr. Plant Biol.* **2018**, *60*, 757–779. [\[CrossRef\]](#)
15. Jin, X.; Lin, Q.; Zhang, X.; Zhang, S.; Ma, W.; Sheng, P.; Zhang, L.; Wu, M.; Zhu, X.; Li, Z.; et al. OsPRMT6b balances plant growth and high temperature stress by feedback inhibition of abscisic acid signaling. *Nat. Commun.* **2025**, *16*, 5173. [\[CrossRef\]](#) [\[PubMed\]](#)
16. Huang, Z.; Lin, R.; Dong, Y.; Tang, M.; Xia, X.; Fang, L.; Yu, J.; Kang, H.; Zhou, Y. MiR164a-targeted NAM3 inhibits thermotolerance in tomato by regulating HSF4b-mediated redox homeostasis. *Plant Physiol.* **2025**, *197*, kiae113. [\[CrossRef\]](#)
17. Wang, Q.; Wu, Y.; Wu, W.; Lyu, L.; Li, W. Research progress on the physiological, biochemical and molecular regulatory mechanisms of fruit tree responses to high-temperature stress. *Hortic. Plant J.* **2025**, *11*, 1–14. [\[CrossRef\]](#)
18. Gao, Z.; Zhou, Y.; He, Y. Molecular epigenetic mechanisms for the memory of temperature stresses in plants. *J. Genet. Genom.* **2022**, *49*, 991–1001. [\[CrossRef\]](#)
19. Li, T.; Wang, Y.H.; Liu, J.X.; Feng, K.; Xu, Z.S.; Xiong, A.S. Advances in genomic, transcriptomic, proteomic, and metabolomic approaches to study biotic stress in fruit crops. *Crit. Rev. Biotechnol.* **2019**, *39*, 680–692. [\[CrossRef\]](#)
20. Zhou, J.; Tang, X.; Li, J.; Dang, S.; Ma, H.; Zhang, Y. Comparative transcriptomic and metabolomic analyses provide insights into the responses to high temperature stress in alfalfa (*Medicago sativa* L). *BMC Plant Biol.* **2024**, *24*, 776. [\[CrossRef\]](#) [\[PubMed\]](#)
21. Khan, Z.; Liu, S.; Peng, J.; Cai, H.; Bai, Y.; Hu, B.; Zhang, L.; Zhuang, P.; Cai, Z.; Xie, L. Growth regulation mechanism of *Rhododendron moulmainense* to high-temperature stress: Integrated physiological, transcriptomic, and metabolomic insights. *Front. Plant Sci.* **2025**, *16*, 1680853. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Almeida, J.; Perez-Fons, L.; Fraser, P.D. A transcriptomic, metabolomic and cellular approach to the physiological adaptation of tomato fruit to high temperature. *Plant Cell Environ.* **2021**, *44*, 2211–2229. [\[CrossRef\]](#)
23. Zhao, Q.; Zhou, L.; Liu, J.; Du, X.; Huang, F.; Pan, G.; Cheng, F. Relationship of ROS accumulation and superoxide dismutase isozymes in developing anther with floret fertility of rice under heat stress. *Plant Physiol. Biochem.* **2018**, *122*, 90–101. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Zhao, F.; Zhang, D.; Zhao, Y.; Wang, W.; Yang, H.; Tai, F.; Li, C.; Hu, X. The difference of physiological and proteomic changes in maize leaves adaptation to drought, heat, and combined both stresses. *Front. Plant Sci.* **2016**, *7*, 1471. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Ru, C.; Hu, X.; Chen, D.; Wang, W.; Zhen, J. Photosynthetic, antioxidant activities, and osmoregulatory responses in winter wheat differ during the stress and recovery periods under heat, drought, and combined stress. *Plant Sci.* **2023**, *327*, 111557. [\[CrossRef\]](#)
26. Rexroth, S.; Mullineaux, C.W.; Ellinger, D.; Sendtko, E.; Rögner, M.; Koenig, F. The plasma membrane of the cyanobacterium *Gloeobacter violaceus* contains segregated bioenergetic domains. *Plant Cell* **2011**, *23*, 2379–2390. [\[CrossRef\]](#)
27. Al-Ashkar, N.M.; Bakry, B.A.; El-Bassiouny, H.M.S.; Abdallah, M.; Sadak, M. Sodium nitroprusside as a signal molecule for up-regulating membrane characteristics, antioxidant defense system to improve flax productivity under water stress. *Oil Crop Sci.* **2024**, *9*, 160–169. [\[CrossRef\]](#)
28. Zhu, J.; Cao, X.; Deng, X. Epigenetic and transcription factors synergistically promote the high temperature response in plants. *Trends Biochem. Sci.* **2023**, *48*, 788–800. [\[CrossRef\]](#)
29. Qian, Y.; Hu, Z.; Tang, Y.; Meng, J.; Tao, J.; Zhao, D. Isolation and function analysis of herbaceous peony bHLH transcription factor PlbHLH87 in response to high-temperature stress. *Sci. Hortic.* **2024**, *329*, 112993. [\[CrossRef\]](#)
30. Fang, Y.; Ma, H.Y.; Wu, Z.M.; Liu, C.F.; Xu, T.Y.; Wang, T.; Li, L.X.; Feng, S.Q. High temperature induces MdGATA15 to suppress anthocyanin accumulation in apple peels. *Plant J.* **2025**, *123*, e70381. [\[CrossRef\]](#)
31. Lu, H.P.; Liu, X.H.; Wang, M.J.; Zhu, Q.Y.; Lyu, Y.S.; Xu, J.H.; Liu, J.X. Author correction: The NAT1-bHLH110-CER1/CER1L module regulates heat stress tolerance in rice. *Nat. Genet.* **2025**, *57*, 763. [\[CrossRef\]](#)
32. Liang, Y.; Yang, C.; Ming, F.; Yu, B.; Cheng, Z.; Wang, Y.; Yan, S. A bHLH transcription factor, CsSPT, regulates high-temperature resistance in cucumber. *Hortic. Plant J.* **2024**, *10*, 503–514. [\[CrossRef\]](#)

33. Zhou, J.; Fang, S.; Liu, X.; Luo, L.; Liu, Y.; Zhang, H. OsERF2 acts as a direct downstream target of OsEIL1 to negatively regulate salt tolerance in rice. *Rice* **2025**, *18*, 29. [\[CrossRef\]](#)
34. Luo, A.; Yang, C.; Yao, Z.W.; Zhang, L.L.; Ye, H.; Xu, J.H.; Lu, H.P.; Liu, J.X. Transcription factors ERF74/77/108/125 enhance thermotolerance in rice by regulating common and distinct heat-responsive gene expression. *Plant Biotechnol. J.* **2025**, *23*, 4539–4551. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Meng, X.; Wang, J.R.; Wang, G.D.; Liang, X.Q.; Li, X.D.; Meng, Q.W. An R2R3-MYB gene, LeAN2, positively regulates thermotolerance in transgenic tomato. *J. Plant Physiol.* **2015**, *175*, 1–8. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Casaretto, J.A.; El-Kereamy, A.; Zeng, B.; Stiegelmeier, S.M.; Chen, X.; Bi, Y.M.; Rothstein, S.J. Expression of OsMYB55 in maize activates stress-responsive genes and enhances heat and drought tolerance. *BMC Genom.* **2016**, *17*, 312. [\[CrossRef\]](#)
37. Zhao, Y.; Tian, X.; Wang, F.; Zhang, L.; Xin, M.; Hu, Z.; Yao, Y.; Ni, Z.; Sun, Q.; Peng, H. Characterization of wheat MYB genes responsive to high temperatures. *BMC Plant Biol.* **2017**, *17*, 208. [\[CrossRef\]](#)
38. Li, Z.; Li, Z.; Ji, Y.; Wang, C.; Wang, S.; Shi, Y.; Le, J.; Zhang, M. The heat shock factor 20-HSF4-cellulose synthase A2 module regulates heat stress tolerance in maize. *Plant Cell* **2024**, *36*, 2652–2667. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Xin, H.; Zhang, H.; Chen, L.; Li, X.; Lian, Q.; Yuan, X.; Hu, X.; Cao, L.; He, X.; Yi, M. Cloning and characterization of HsfA2 from lily (*Lilium longiflorum*). *Plant Cell Rep.* **2010**, *29*, 875–885. [\[CrossRef\]](#)
40. Zhang, L.; Zheng, L.; Wu, J.; Liu, Y.; Liu, W.; He, G.; Wang, N. OsCCRL1 is essential for phenylpropanoid metabolism in rice anthers. *Rice* **2023**, *16*, 10. [\[CrossRef\]](#)
41. Li, J.; Yu, Q.; Liu, C.; Zhang, N.; Xu, W. Flavonoids as key players in cold tolerance: Molecular insights and applications in horticultural crops. *Hortic. Res.* **2025**, *12*, uhae366. [\[CrossRef\]](#) [\[PubMed\]](#)
42. Postiglione, A.E.; Delange, A.M.; Ali, M.F.; Wang, E.Y.; Houben, M.; Hahn, S.L.; Khoury, M.G.; Roark, C.M.; Davis, M.; Reid, R.W.; et al. Flavonols improve tomato pollen thermotolerance during germination and tube elongation by maintaining reactive oxygen species homeostasis. *Plant Cell* **2024**, *36*, 4511–4534. [\[CrossRef\]](#)
43. Wang, Z.; Guo, J.; Luo, W.; Niu, S.; Qu, L.; Li, J.; Chen, Y.; Li, G.; Yang, H.; Lu, D. Salicylic acid cooperates with lignin and sucrose signals to alleviate waxy maize leaf senescence under heat stress. *Plant Cell Environ.* **2025**, *48*, 4341–4355. [\[CrossRef\]](#)
44. Dong, N.Q.; Lin, H.X. Contribution of phenylpropanoid metabolism to plant development and plant–environment interactions. *J. Integr. Plant Biol.* **2021**, *63*, 180–209. [\[CrossRef\]](#) [\[PubMed\]](#)
45. Rawal, H.C.; Singh, N.K.; Sharma, T.R. Conservation, divergence, and genome-wide distribution of gene families in plants. *Int. J. Genom.* **2013**, *2013*, 678969. [\[CrossRef\]](#)
46. Wu, D.G.; Zhan, Q.W.; Yu, H.B.; Huang, B.H.; Cheng, X.X.; Li, W.Y.; Huang, S.C.; Wang, C.J.; Du, J.L. Genome-wide identification and analysis of maize gene family and its expression profile in response to high-temperature stress. *Pak. J. Bot.* **2020**, *52*, 1577–1587. [\[CrossRef\]](#)
47. Feduraev, P.; Riabova, A.; Skrypnik, L.; Pungin, A.; Tokupova, E.; Maslennikov, P.; Chupakhina, G. Assessment of the role of PAL in lignin accumulation in wheat (*Triticum aestivum* L.) at the early stage of ontogenesis. *Int. J. Mol. Sci.* **2021**, *22*, 9848. [\[CrossRef\]](#)
48. Shi, J.; Wang, Y.; Fan, X.; Li, R.; Yu, C.; Peng, Z.; Gao, Y.; Liu, Z.; Duan, L. A novel plant growth regulator B2 mediates drought resistance by regulating reactive oxygen species, phytohormone signaling, phenylpropanoid biosynthesis, and starch metabolism pathways in *Carex breviculmis*. *Plant Physiol. Biochem.* **2024**, *213*, 108860. [\[CrossRef\]](#)
49. Pratyusha, D.S.; Sarada, D.V.L. MYB transcription factors—Master regulators of phenylpropanoid biosynthesis and diverse developmental and stress responses. *Plant Cell Rep.* **2022**, *41*, 2245–2260. [\[CrossRef\]](#) [\[PubMed\]](#)
50. Dhindsa, R.S.; Plumb-Dhindsa, P.; Thorpe, T.A. Leaf senescence: Correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J. Exp. Bot.* **1981**, *32*, 93–101. [\[CrossRef\]](#)
51. Elstner, E.F.; Heupel, A. Inhibition of nitrite formation from hydroxylammonium chloride: A simple assay for superoxide dismutase. *Anal. Biochem.* **1976**, *70*, 616–620. [\[CrossRef\]](#)
52. Velikova, V.; Yordanov, I.; Edreva, A. Oxidative stress and some antioxidant systems in acid rain-treated bean plants: Protective role of exogenous polyamines. *Plant Sci.* **2000**, *151*, 59–66. [\[CrossRef\]](#)
53. Arnon, D.I. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiol.* **1949**, *24*, 1–15. [\[CrossRef\]](#)
54. Chen, S.; Zhou, Y.; Chen, Y.; Gu, J. fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **2018**, *34*, i884–i890. [\[CrossRef\]](#) [\[PubMed\]](#)
55. Pertea, M.; Pertea, G.M.; Antonescu, C.M.; Chang, T.-C.; Mendell, J.T.; Salzberg, S.L. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat. Biotechnol.* **2015**, *33*, 290–295. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Li, B.; Dewey, C.N. RSEM: Accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform.* **2011**, *12*, 323. [\[CrossRef\]](#)
57. Love, M.I.; Huber, W.; Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **2014**, *15*, 550. [\[CrossRef\]](#)
58. Benjamini, Y.; Hochberg, Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Stat. Soc. B* **1995**, *57*, 289–300. [\[CrossRef\]](#)

59. Dunn, W.B.; Broadhurst, D.; Begley, P.; Zelena, E.; Francis-McIntyre, S.; Anderson, N.; Brown, M.; Knowles, J.D.; Halsall, A.; Haselden, J.N.; et al. Procedures for large-scale metabolic profiling of serum and plasma using gas chromatography and liquid chromatography coupled to mass spectrometry. *Nat. Protoc.* **2011**, *6*, 1060–1083. [[CrossRef](#)]
60. Warnes, G.R.; Bolker, B.; Lumley, T.; Johnson, R.C. Gmodels: Various R Programming Tools for Model Fitting. R Package Version 2.18.1. 2025. Available online: <https://CRAN.R-project.org/package=gmodels> (accessed on 12 April 2025).
61. Kolde, R.; Kolde, M.R. Package ‘pheatmap’. *R Package* **2015**, *1*, 790.
62. Thevenot, E.A. ropls: PCA, PLS (-DA) and OPLS (-DA) for multivariate analysis and feature selection of omics data. *R Package Version* **2016**.
63. Saccenti, E.; Hoefsloot, H.C.J.; Smilde, A.K.; Westerhuis, J.A.; Hendriks, M.M.W.B. Reflections on univariate and multivariate analysis of metabolomics data. *Metabolomics* **2014**, *10*, 361–374. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.