



Article

# Genome-Wide Identification and Expression Analysis of the *GRF* Gene Family in *Gossypium hirsutum* L.

Cong-Hua Feng <sup>1,†</sup>, Linlin Liu <sup>1,†</sup>, Di Liu <sup>1</sup> , Junbo Zhen <sup>1</sup> , Mengzhe Li <sup>1</sup>, Mengmeng Jiang <sup>1,2</sup> and Jina Chi <sup>1,\*</sup>

<sup>1</sup> Institute of Cotton, Hebei Academy of Agricultural and Forestry Sciences/Key Laboratory of Biology and Genetic Improvement of Cotton in Huanghuaihai Semiarid Area, Ministry of Agriculture and Rural Affairs, Hebei Key Laboratory of Cotton Bio-Breeding and Cultivation Physiology, Shijiazhuang 050051, China; fch19942024@163.com (C.-H.F.); spp2016@126.com (L.L.); liudijoy@126.com (D.L.); zjb210@126.com (J.Z.); limengzhe0131@126.com (M.L.); jmm18270464849@163.com (M.J.)

<sup>2</sup> Genetics Laboratory, College of Life Science, Hebei University, Baoding 071002, China

\* Correspondence: mhschijina@haafs.org; Tel.: +86-311-87652076

† These authors contributed equally to this work.

## Abstract

Growth Regulating Factors (GRFs) are plant-specific transcription factors that play crucial roles in regulating growth and development throughout the plant life cycle. A total of 34 *Gossypium hirsutum* GRF family genes were identified at the genome-wide level, which were unevenly distributed on 19 chromosomes, and were predicted to be mainly localized in the nucleus and plasma membrane. The number of GRF family genes varied greatly among different species, and they were categorized into four subfamilies (I–IV) according to their phylogenetic relationships. The *G. hirsutum* GRF genes possessed specific highly conserved structural domains, Trp-Arg-Cys motif (WRC) and Gln, Leu, Gln motif (QLQ), and structural analysis of the genes revealed that they contained 1–23 exons, and most of them contained UTRs. Intraspecies covariance analysis revealed that the GRF genes expanded in *G. hirsutum* by segmental duplication. The promoter region of the *G. hirsutum* GRF gene contained a large number of adversity stress response elements, as well as a small number of hormone response elements and growth and development-related response elements. Transcriptome data showed that the expression of *G. hirsutum* GRF genes was significantly higher in leaves than in other tissues, and some GRF genes responded to a variety of abiotic stresses. Additionally, transcriptomic sequencing revealed significantly higher expression levels of *GhGRFs* (e.g., *GhGRF13/14/18*) in embryonic callus (EC) compared to non-embryonic callus (NEC). This differential expression was validated by RT-qPCR, which confirmed that *GhGRF13/14/16/20* were significantly upregulated in EC relative to NEC. These findings provide valuable candidate genes and molecular insights for improving *G. hirsutum* regeneration efficiency and yield-related traits through genetic manipulation, thereby accelerating the molecular breeding of elite *G. hirsutum* varieties.

**Keywords:** growth regulating factors; *Gossypium hirsutum*; growth and development; expression pattern; embryogenic callus



Academic Editors: Setsuko Komatsu and Pedro Martínez-Gómez

Received: 27 January 2026

Revised: 13 February 2026

Accepted: 21 February 2026

Published: 26 February 2026

**Copyright:** © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

## 1. Introduction

As a primary global source of natural fiber, cotton is a vital cash crop and a fundamental raw material for the textile industry [1–3]. However, cotton production faces a number of challenges, including attacks by pests and diseases, increasing environmental pressures, and bottlenecks in genetic improvement [4]. In order to meet these challenges,

the development of cotton breeding technology is particularly important. Among various breeding approaches, the induction and utilization of embryogenic callus (EC) in cotton have garnered significant attention due to its unique biological characteristics and application potential [5,6]. Cotton somatic embryogenesis plays a crucial role in the genetic transformation of cotton [7,8]. Embryogenic callus refers to a proliferative mass of cells derived from induced embryonic or embryogenic tissues, possessing regenerative capacity [9,10]. This type of callus not only retains embryogenic potential but can also proliferate indefinitely under appropriate culture conditions, thereby providing essential experimental material for genetic transformation, variety improvement, and biotechnological research in cotton [11–14]. Through embryogenic callus, efficient genetic transformation can be achieved, facilitating the selection of novel cultivars with desirable traits. In addition, embryonic healing tissues play a key role in the study of developmental biology, cell biology and molecular biology of cotton, which helps to reveal the molecular mechanism of cotton growth and development [6]. During cotton development, a large number of genes are differentially expressed and interact with each other at different stages and in different tissues, and this complex regulatory mechanism ensures the normal development of somatic embryos [7]. Although the induction and application of embryonic healing tissues in cotton are of great significance in cotton breeding and biotechnology research, there are still some technical and theoretical problems that need to be solved.

GRFs typically contain the Trp-Arg-Cys motif (WRC) and Gln, Leu, Gln motif (QLQ) structural domains [15–17]. In addition, three less-conserved structural domains (TQL, FFD, GGPL) have been identified in the C-terminal region of GRF proteins; however, the specific functions of these less-conserved motifs remain unclear [18,19]. GRFs have been extensively studied, and their roles in plant-healing tissue formation have been initially explored [20,21]. Lu et al. previously identified a novel rice gene, growth regulatory factor 1 (*OsGRF1*), which encodes a suspected transcription factor that plays a regulatory role in stem elongation [22,23]. A large number of GRF genes have been identified in *Arabidopsis thaliana*, *Oryza sativa*, *Populus trichocarpa*, and *Triticum aestivum* [24–26]. GRFs function to regulate leaf growth and act as positive regulators of cell proliferation, affecting leaf shape and size [27,28]. *PpnGRF5-1* is overexpressed in poplar and results in larger leaves than the wild type by increasing cell division and cell expansion [29,30]. Overexpression of *BnGRF2a* resulted in 20% leaf elongation in kale-type oilseed rape, and overexpression of *AtGRF1* resulted in larger leaves [19,31]. GRF mutant leaves were smaller, and *AtGRF9* negatively regulated leaf development by inhibiting *A. thaliana* leaf primordial cell proliferation [21,32–35].


This study was designed to characterize the induction mechanisms and genetic stability of cotton embryogenic tissues, with the ultimate goal of facilitating their application in cotton breeding. This was done by optimizing the induction and cultivation conditions of guiding tissues to improve their regeneration efficiency and genetic stability; by using modern molecular biology techniques to analyze the mechanism of guiding tissues in cotton breeding; and by exploring new strategies to combine guiding tissue technology with gene editing and marker-assisted selection, etc., with a view to providing new theoretical and technical support for the genetic improvement and varietal innovation of cotton. The results of this study will not only provide new theories and methods for cotton breeding, but also provide reference for biotechnology research of other crops.

## 2. Results

### 2.1. Genome-Wide Identification and Evolutionary Analysis of GhGRF Genes in Terrestrial *G. hirsutum*

In the whole genome data of *G. hirsutum*, 34 *G. hirsutum* GRF genes were finally screened by BLAST (Table S1). The protein database analysis in the NCBI database determined that all of them contained WRC and QLQ structural domains and belonged to the GRF gene family. The genes were named according to the order of the corresponding chromosome positions (*GhGRF1- GhGRF34*). The *GhGRF* genes were distributed on 19 chromosomes of the *G. hirsutum* genome (Figure S1), with three *GhGRF* genes on five chromosomes, two *GhGRF* genes on another five chromosomes, and one *GhGRF* gene on the remaining nine chromosomes. Next, analysis of the physicochemical properties of the proteins of the *GhGRFs* gene family revealed that the *GhGRF* genes encoded 182–610 amino acids, and the molecular weights of the proteins were relatively large, all of which were greater than 20 kDa (Table S1). The theoretical pIs of the *GhGRFs* gene family ranged from 5.89 to 9.65, with eight genes being acidic proteins and 26 genes being basic proteins. All 34 members of the *GhGRFs* gene family are stable with an instability index greater than 40. All *GhGRF* proteins are hydrophilic. Most of the *GhGRF* proteins are located in the nucleus and three genes are located in the plasma membrane (*GhGRF8/13/30*) as predicted by the Plant-mPLOC database.


Phylogenetic tree analysis classified the *G. hirsutum* GRF genes into four subfamilies, and the genes belonging to the same subfamily were closely related and highly homologous (Figure 1A). These results suggest that the GRFs gene family is highly conserved throughout evolution. We analyzed the number of GRF genes in different species and found that there were significant differences in the number of GRF genes in different species (Figure 1B). There was only one GRF gene in *Marchantia polymorpha*, while *Salix purpurea* possessed 43 GRF genes.



**Figure 1.** Phylogenetic analysis of the GRF gene family in *G. hirsutum* and the number of GRF genes across multiple species. (A) Evolutionary and phylogenetic analysis of the *GhGRFs* gene family in *G. hirsutum*. The GRF gene family in *G. hirsutum* is divided into four subfamilies based on different colors. (B) Number of GRF gene family members across 14 species. *Arabidopsis thaliana*; *Triticum aestivum*; *Brassica napus*; *Glycine soja*; *Gossypium hirsutum*; *Oryza sativa* (japonica); *Populus euphratica*; *Populus trichocarpa*; *Salix purpurea*; *Zea mays*; *Theobroma cacao*; *Glycine max*; *Coffea canephora*; *Marchantia polymorpha*.

## 2.2. Motif, Conserved Structural Domain and Gene Structure Analysis of the GhGRF Genes

All 34 members of the *GhGRFs* gene family possess Motif2, Motif3, and Motif1, and their sequences are very conserved (Table S2). All *GhGRF* genes contain the WRC and QLQ structural domains, and *GhGRF25* also contains the RETICULATA-like super family conserved structural domains. Gene structure analysis of the *GhGRFs* gene family revealed that they contain 1–23 exons, 0–23 introns, and 0–4 UTRs. *GhGRF21* contains the highest number of 23 exons and 23 introns, and *GhGRF10* and *GhGRF32* do not contain UTRs (Figure 2).



**Figure 2.** Analysis of motifs, conserved domains, and gene structures in the *GhGRFs* gene family of *G. hirsutum*. (A) Phylogenetic tree of the *GhGRFs* gene family in *G. hirsutum*; (B) Motif analysis of the *GhGRFs* gene family in *G. hirsutum*, with each motif represented by a differently colored square; (C) Gene structure analysis of the *GhGRFs* gene family in *G. hirsutum*; (D) Gene structure analysis of the *GhGRFs* gene family in *G. hirsutum*, with upstream and downstream non-coding regions indicated by green rectangles, exons by yellow rectangles, and introns by gray lines.


## 2.3. Collinearity Analysis of GhGRF Genes

During the evolutionary process, the *G. hirsutum* *GhGRFs* gene family has undergone several duplication events. As shown in Figure 3, 34 *GhGRF* genes within the *G. hirsutum* genome formed 57 gene pairs by chromosome segmental duplication (SD). Then, we plotted the covariance analysis of the *G. hirsutum* *GhGRFs* gene family with five other plants (*A. thaliana*, *O. sativa*, *T. aestivum*, *P. trichocarpa*, and *Zea mays*) (Figure 4). The *G. hirsutum* *GhGRFs* gene family has 31 pairs of GRF homologues with *A. thaliana*, 14 pairs of homologues with *O. sativa*, 10 pairs of GRF homologues with *T. aestivum*, 34 pairs of GRF homologues with *P. trichocarpa* and nine pairs of GRF homologues with *Z. mays*.

## 2.4. GhGRFs Gene Family Cis-Element Analysis

To determine the expression pattern of the *GhGRFs* gene family, the cis-elements of the *GhGRFs* gene family promoters were analyzed using the PlantCARE database (Figure 5). These elements are involved in cotton growth and development as well as in stress response, including abiotic and biotic stresses, phytohormone responses, and plant growth and development. Stress-related cis-elements (MYB, MYC, ARE) were enriched in some *GhGRF* genes, suggesting that these *GhGRF* genes may play a key role in response to unfavorable conditions. In addition, the promoters of some *GhGRF* genes are enriched in ABRE (involved in ABA response), such as *GhGRF29*, which may be responsive to ABA

hormones. The *GhGRF10/13/29* promoter contains a large number of promoter elements related to growth and development, such as the G-box.




**Figure 3.** Collinearity analysis of the *GhGRFs* gene family. The fan-shaped rings indicate chromosome numbers and sizes. Gray line segments represent genomic collinear blocks, while red lines denote homologous gene pairs of *GhGRF* genes.

### 2.5. Transcriptome Analysis of *GhGRF* Gene Family in *G. hirsutum*


To further investigate the role of the *GhGRFs* gene family in cotton growth, development and abiotic stresses, we downloaded tissue expression and abiotic stress expression transcriptome data of the *GRF* gene family of *G. hirsutum* from the database. Expression patterns of the *GhGRFs* across 10 different tissues and under four abiotic stresses conditions were visualized using heatmaps (Figure 6). The results showed that most *GhGRFs* were downregulated under abiotic stress, including *GhGRF8*, *GhGRF9*, and *GhGRF22*. In contrast, *GhGRF2* and *GhGRF34* were upregulated under three of the stress treatments. Several genes exhibited tissue-specific high expression; for instance, *GhGRF13* and *GhGRF30* were highly expressed in leaves, while *GhGRF18* showed elevated expression in filaments and petals. These findings suggest that members of the *GhGRF* gene family may perform distinct functions at different developmental stages and in response to environmental stresses in cotton.

### 2.6. *GRF* Gene Family Expression Patterns in Embryonic Callus (EC) and Non-Embryonic Callus (NEC)


Next, transcriptomic sequencing was performed on EC and NEC from *G. hirsutum*, which led to the identification of 12 *GhGRFs* genes. The results indicated that all of these genes were upregulated in EC compared to NEC (Figure 7A). Their expression patterns were further validated by RT-qPCR (Figure 7B,C). The results confirmed that *GhGRF1*, *GhGRF13*, *GhGRF14*, *GhGRF16* and *GhGRF20* were significantly highly expressed in EC but at lower levels in NEC, suggesting their potential key roles in EC formation and maintenance.




**Figure 4.** Collinearity analysis of the *GhGRFs* gene family in *G. hirsutum* and 5 plant species. (A): *A. thaliana*; (B): *O. sativa*; (C): *T. aestivum*; (D): *P. trichocarpa*; (E): *Zea mays*. Gray lines indicate the collinearity between the *G. hirsutum* genome and those of *A. thaliana*, *O. sativa*, *T. aestivum*, *P. trichocarpa*, and *Z. mays*, while red lines highlight collinear gene pairs between *G. hirsutum* *GhGRF* genes and genes from other species.



**Figure 5.** Promoter cis-acting elements of *GhGRFs* genes. (A) The heatmap represents the number of cis-acting elements; (B) The histogram shows the number of cis-acting elements of different types per gene, with different colors denoting distinct categories: plant hormone response type, abiotic and biotic stress type, and plant growth and development type.




**Figure 6.** Expression profiles of *GhGRFs* during *G. hirsutum* growth and development and under abiotic stresses. (A) Heatmap of *GhGRFs* gene expression levels in different tissues. (B) Heatmap of *GhGRFs* gene expression levels under cold stress, drought stress, heat stress, and salt stress. The color bars represent high and low relative expression levels, respectively.



**Figure 7.** Transcriptome and RT-qPCR analysis of *GhGRFs* gene expression in EC and NEC. (A) RNA-seq analysis of 12 *GhGRF* genes in EC and NEC. The color bars represent high and low relative expression levels, respectively. (B,C) RT-qPCR results for 12 *GhGRF* genes in *G. hirsutum* EC and NEC. One-way ANOVA with LSD multiple comparison,  $n = 12$ . \*  $p < 0.05$ ; \*\*  $p < 0.01$ .

To further explore the involvement of *GhGRFs* in EC, we focused on seven genes that exhibited marked upregulation in the expression heatmap. A regulatory network analysis revealed that *GhGRF1*, *GhGRF18*, *GhGRF13*, *GhGRF26*, *GhGRF14*, *GhGRF33*, and *GhGRF15* are predicted to be regulated by 8, 8, 12, 10, 10, 3, and 4 transcription factors, respectively (Figure 8A–G). These transcription factors belong to families such as MYB, Dof, ERF, HD-

ZIP, AP2, etc. Additionally, the same gene may be co-regulated by the same transcription factor simultaneously. *GhGRF1* and *GhGRF18* potentially share eight common regulators. Since these transcription factors themselves show tissue-specific expression patterns, our findings imply that *GhGRF* genes may be differentially regulated across tissues. These transcription factors exhibit varying expression levels in different tissues, suggesting that *GhGRF* genes may be regulated by different transcription factors in different tissues.



**Figure 8.** Bioinformatics analysis of upstream transcription factors (TFs) of *GhGRFs*. (A–G) Mutual regulation network analysis of TFs-*GhGRF1/18/13/26/14/33/15*. Green circles indicate upstream transcription factors of *GhGRFs*; yellow circles represent *GhGRF1/18/13/26/14/33/15*. Heatmap showing transcriptional levels of upstream transcription factors in leaf, root, and stem. The color bars represent high and low relative expression levels, respectively.

### 3. Discussion

*GRF* genes constitute a plant-specific transcription factor family widely involved in growth development, and stress response, playing important regulatory roles in plants [22,36]. In this study, 34 *GRF* genes were identified in *G. hirsutum*. Notably, the number of *GRF* genes in *G. hirsutum* exceeds that reported in several model species, such as *A. thaliana* (9), *P. trichocarpa* (26), *O. sativa* (19), and *T. aestivum* (16) [24,26,29]. This variation suggests that the *GRF* family has undergone different expansion through duplication events during evolution. Analysis of the physicochemical properties of the *GhGRFs* indicated that most are basic hydrophilic proteins, with a Grand Average of Hydropathicity (GRAVY)

below zero. Furthermore, an instability index greater than 40 for most members suggests these proteins may be relatively prone to degradation or denaturation (Table S1).

Phylogenetic analyses classified the 34 *GhGRFs* into four distinct clades, with members of the same subfamily showing highly similar homology (Figure 1A). There were significant differences in the number of *GRF* genes in different species, which reflected the diversity and adaptability of the *GRF* genes family during the evolutionary process (Figure 1B) [24]. Whole-genome duplication events have notably expanded the *GRF* family in some species [37]; for example, there are 48 *GRF* genes in tetraploid kale (*Brassica napus*), far more than its diploid ancestors [38], and *Salix purpurea* also exhibits a high *GRF* count likely due to genome duplication [29]. Furthermore, covariance analysis of gene families can explain gene evolution [39]. Tandem duplications and segmental duplications play a key role in species evolution [40]. In cotton, the *GRF* gene family forms two tandem duplication clusters on chromosomes A12 and D12 (Figure S1). Additionally, two segmental duplication pairs (*GhGRF14/15* and *GhGRF31/32*) exhibited upregulation, supporting the role of both duplication types in the expansion of cotton *GRF* genes. Synteny analysis also revealed numerous orthologous gene pairs between cotton and five representative model plants, indicating a degree of evolutionary conservation across herbaceous and woody lineages (Figure 4).

Gene structure diversity often mirrors functional differentiation [41]. Exon–intron architecture analysis of *GhGRFs* revealed considerable variation in exon (1–23) and intron (0–23) numbers and arrangements, implying relatively low structural conservation and potential functional divergence among the 34 genes (Figure 2). This structural diversity is consistent with observations in other plants, highlighting the plasticity of the *GRF* family across species [42]. At the protein level, two conserved structural domains, QLQ and WRC, were predicted at the N-terminus of all *GhGRFs*, aligning with previous reports [43]. The QLQ domain typically mediates protein–protein interactions, including transcription factor complexes formation, whereas the WRC domain is involved in DNA binding [44]. It contributes to the recognition and binding of specific DNA sequences by transcription factors, thereby regulating gene expression. The conservation and functional relevance of these two structural domains underscore the likely importance of *GhGRFs* in regulating cotton growth, development and stress responses.

*GRFs*, as key regulators of plant growth, primarily by controlling cell proliferation and expansion during primary and secondary development [45]. In *A. thaliana*, for example, members of the *GRFs* form complexes with GIF (GRF-Interacting Factor) proteins to regulate cell division and organ formation [43]. Beyond development, *GRFs* also participate in hormone signaling and abiotic stress responses [42,46]. For example, *miR396* regulates root growth and drought tolerance in plants under drought stress by targeting *GRFs* [22]. Gene expression levels are critical for plant growth, development, and response to external stresses [47]. Our expression analysis showed that *GhGRF* genes are broadly expressed across various tissues and developmental stages (Figure 6A), yet individual members exhibited distinct expression patterns in different organs and under abiotic stresses. This specificity may be linked to the varied cis-regulatory elements identified in their promoters (Figures 5 and 6B). Notably, *GhGRF* promoters are enriched in stress-related elements, and their expression is highly responsive to abiotic stress. Furthermore, transcriptomic analysis of embryogenic callus (EC) and non-embryogenic callus (NEC) identified 12 *GhGRFs* that were consistently upregulated in EC (Figure 7). Their expression may be finely tuned by different transcription factors across tissues (Figure 8). Although the *GRF* gene family in upland cotton has been previously identified, we re-identified 34 *GRF* genes using the latest upland cotton genome data [42,48]. These results suggest that upregulation of specific *GhGRFs*, as key developmental regulators, may contribute to establishing and maintaining

the high differentiation potential of embryogenic callus, which is essential for genetic transformation and plant regeneration in cotton.

## 4. Materials and Methods

### 4.1. Cotton Material Cultivation and Processing

The cotton material used in this experiment was sourced from the Cotton Research Institute of Hebei Academy of Agricultural and Forestry Sciences, variety JIN668 [6,49,50]. After removing the seed coats, the cotton seeds were disinfected in a 0.1% mercuric chloride solution for 8–10 min, then rinsed five times with ddH<sub>2</sub>O to thoroughly remove the surface mercuric chloride solution.

The seeds were then cultured in prepared MS medium (MS (Murashige and Skoog) 2.2 g/L, sucrose 20 g/L, pH adjusted to 7.8, agar 7 g/L) for 5–7 days after autoclaving (121 °C, 15 min). After 5–7 days, the cotton hypocotyls were cut into 0.5–1 cm segments and placed in callus induction medium (MS + 0.1 mg/L 2,4-Dichlorophenoxyacetic acid + 0.1 mg/L Kinetin + 3% glucose + 0.3% Phytigel, pH 5.85–5.95) for induction. Finally, culturing was continued until NEC and EC were obtained. All operations were performed in a sterile tissue culture workbench.

### 4.2. Identification of the GhGRF Genes Family in *G. hirsutum* and Construction of an Evolutionary Tree

The GhGRFs gene family was identified and analyzed based on the whole genome of *G. hirsutum*, HAU [51]. The *G. hirsutum* genome sequence was downloaded from the CottonFGD database (<https://cottonfgd.net/>). The GRF gene family files for species such as *A. thaliana*, *T. aestivum*, *B. napus*, *G. soja*, *O. sativa* (japonica), *P. euphratica*, *P. trichocarpa*, *S. purpurea*, and *Z. mays* are sourced from the Plant Transcriptional Regulation Atlas Database (<https://plantregmap.gao-lab.org/>) [52]. We searched for GhGRF genes from the terrestrial cotton genome using the Trp-Arg-Cys motif (WRC) and Gln, Leu, Gln motif (QLQ) structural domains and used the CottonFGD database to examine and obtain the correct GhGRF family genes [53,54]. The physicochemical properties of the GhGRF genes were obtained through the Expasy website ([https://web.expasy.org/compute\\_pi/](https://web.expasy.org/compute_pi/)) (accessed on 4 August 2024)) [55].

For a more comprehensive analysis of the GhGRF genes family, a lineage tree was obtained by comparing all GhGRF protein sequences using ClustalW 2.1 [56,57]. The evolutionary tree was constructed using MEGA5.0 based on the 1000 bootstrap neighbor-joining (NJ) method [58]. Each GhGRF gene was named according to its position on the chromosome, and different subfamilies were indicated by different colors.

### 4.3. Analysis of Gene Structure, Conserved Motifs, and Cis-Acting Elements of the GhGRF Genes Family in *G. hirsutum*

The GhGRF genes family was analyzed bioinformatically as previously described. The GhGRF genes family was examined using The MEME Suite online website (<https://meme-suite.org/meme/index.html>) (accessed on 15 September 2024)) to analyze its structure and identify conserved motifs [59]. Use the One Step Build a ML Tree feature in TBtools software (v2.423) to construct a phylogenetic tree, then obtain the Newick file [60,61]. Simultaneously, submit the amino acid sequences to NCBI-CDD for conserved domain analysis (<https://www.ncbi.nlm.nih.gov/>) [62,63].

In addition, GhGRFs were analyzed for cis-elements using the PlantCARE database, and graphical visualization was performed using Tbttools software [41].

#### 4.4. Analysis of Gene Duplication and Expression Patterns of the GhGRFs Gene Family in *G. hirsutum*

Co-collinearity analyses between the *G. hirsutum* GhGRF gene family and *G. hirsutum* itself as well as other species were assessed using the Multiple collinearity Scan toolkit (MCScanX) [64]. In addition, abiotic stress transcriptome data and tissue expression transcriptome data were obtained from the CottonFGD database. Graphical visualization was performed using Tbttools software.

#### 4.5. RT-qPCR Analysis

RNA extraction was performed using the Nuoto<sup>®</sup> AutoExtractor-32 Nucleic Acid Extractor (KangmaXin (Shanghai) Intelligent Technology Co., Ltd., Shanghai, China), with the 5fz Cytoplasmic RNA Automated Extraction Kit (Product Code: protn\_5fzCeAE96DW3in1) (KangmaXin (Shanghai) Intelligent Technology Co., Ltd., Shanghai, China). Extracted RNA was converted to cDNA using the Reverse Transcription Kit (Tiangen Biotechnology Co., Ltd., Beijing, China). Primers were designed by the Primer 5 software and synthesized by Sangon Ltd. in Shanghai, China. RT-qPCR experiments were performed with *GhUBQ7* (*Ghir\_D12G021700*) as internal reference genes (Table S3) [65]. Three biological replicates and four technical replicates were performed for each sample. RT-qPCR was performed on a CFX96 Touch<sup>™</sup> instrument (Bio-Rad Co. Ltd., Hercules, CA, USA). RT-qPCR procedures and analysis of the results followed previous studies.

#### 4.6. Bioinformatics Analysis and Tissue Expression Analysis of Upstream Transcription Factors of GhGRFs

Potential upstream transcription factors of GhGRFs were identified using the PlantRegMap database (<http://plantregmap.gao-lab.org/network.php> (accessed on 25 October 2024)) [66]. Network diagrams visualizing target genes and their upstream transcription factors were generated using Office software (2013). In the initial phase of the study, gene expression profiles of upstream transcription factors under different treatments were obtained from the CottonFGD database and visualized using TBtools software.

#### 4.7. Statistical Analysis

We analyzed the experimental data using SPSS v.25.0 (SPSS Inc., Chicago, IL, USA). For gene relative expression, one-way analysis of variance (ANOVA) was performed (LSD multiple comparison test, \*  $p < 0.05$ ; \*\*  $p < 0.01$ ). Prior to conducting the ANOVA, data were tested for normality and homogeneity of variance [47].

## 5. Conclusions

In this study, we performed a genome-wide identification of the GRF transcription factor family in *G. hirsutum*, characterizing 34 GhGRF genes. Comprehensive analyses revealed their gene structures, phylogenetic relationships, expression profiles across tissues, responses to abiotic stresses, and specific expression in embryogenic callus (EC). All identified GhGRF proteins contain the characteristic and highly conserved QLQ and WRC domains. Phylogenetic analysis classified the GhGRFs into four distinct clades (I–IV). These genes are distributed asymmetrically across 19 chromosomes, with expansion primarily driven by segmental duplication events, highlighting a key mechanism in the evolution of this family in cotton. Collinearity analysis indicated that the GhGRF family shares closer homology and a stronger evolutionary association with woody plants compared to herbaceous species. Cis-acting element analysis of the GhGRF promoters revealed a strong enrichment of stress-responsive elements (e.g., MYB, MYC, ARE, ABRE), supporting their potential involvement in abiotic stress signaling. Expression profiling demonstrated that GhGRF members exhibit distinct, often tissue-specific expression patterns and differ-

ential responses to various abiotic stresses, suggesting functional diversification and the regulation of separate signaling pathways. Transcriptomic sequencing of embryogenic (EC) and non-embryogenic callus (NEC) showed that 12 *GhGRF* genes were consistently upregulated in EC. RT-qPCR validation confirmed the significantly higher expression of *GhGRF1*, *GhGRF13*, *GhGRF14*, *GhGRF16*, and *GhGRF20* in EC, pointing to their potential key roles in the establishment and maintenance of embryogenic competence. Furthermore, regulatory network analysis suggests that the expression of *GhGRF* genes is likely modulated by different sets of transcription factors in various tissues. In summary, our findings demonstrate that members of the *GhGRF* family are upregulated during somatic embryogenesis in cotton. The sustained and specific expression of different *GhGRF* transcription factors in embryogenic callus, potentially through distinct regulatory pathways, confers diverse functions. This underscores their indispensable role within the complex regulatory network governing callus formation and somatic embryogenesis in *G. hirsutum*.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijms27052191/s1>.

**Author Contributions:** C.-H.F., J.C., D.L., J.Z., M.L., M.J., and L.L. designed the experiments. C.-H.F., J.C., and L.L. drafted the manuscript. C.-H.F. and L.L. conducted RT-qPCR and data analysis. J.C., D.L., L.L., M.L., J.Z. and C.-H.F. critically reviewed the manuscript, L.L. and J.Z. contributed to project administration and funding acquisition. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was supported by Biological Breeding of Stress tolerant and High Yield Cotton Varieties (Project NO.2023ZD04040-2), and HAAFS Agriculture Science and Technology Innovation Project (2022KJCXZX-MHS-7, 2026KJCXZX-MHS-7).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The original contributions presented in this study are included in the article/Supplementary Materials. Further inquiries can be directed to the corresponding author.

**Conflicts of Interest:** The authors declare that they have no competing interests.

## References

1. Du, X.; Huang, G.; He, S.; Yang, Z.; Sun, G.; Ma, X.; Li, N.; Zhang, X.; Sun, J.; Liu, M.; et al. Resequencing of 243 diploid cotton accessions based on an updated A genome identifies the genetic basis of key agronomic traits. *Nat. Genet.* **2018**, *50*, 796–802. [CrossRef] [PubMed]
2. He, S.; Sun, G.; Geng, X.; Gong, W.; Dai, P.; Jia, Y.; Shi, W.; Pan, Z.; Wang, J.; Wang, L.; et al. The genomic basis of geographic differentiation and fiber improvement in cultivated cotton. *Nat. Genet.* **2021**, *53*, 916–924. [CrossRef]
3. Li, L.; Zhang, C.; Huang, J.; Liu, Q.; Wei, H.; Wang, H.; Liu, G.; Gu, L.; Yu, S. Genomic analyses reveal the genetic basis of early maturity and identification of loci and candidate genes in upland cotton (*Gossypium hirsutum* L.). *Plant Biotechnol. J.* **2020**, *19*, 109–123. [CrossRef]
4. Zhang, X.; Liu, S.; Wu, P.; Xu, W.; Yang, D.; Ming, Y.; Xiao, S.; Wang, W.; Ma, J.; Nie, X.; et al. A panoramic view of cotton resistance to *Verticillium dahliae*: From genetic architectures to precision genomic selection. *iMeta* **2025**, *4*, e70029. [CrossRef]
5. Ge, X.; Yu, X.; Liu, Z.; Yuan, J.; Qin, A.; Wang, Y.; Chen, Y.; Qin, W.; Liu, Y.; Liu, X.; et al. Spatiotemporal transcriptome and metabolome landscapes of cotton somatic embryos. *Nat. Commun.* **2025**, *16*, 859, Correction in *Nat. Commun.* **2025**, *16*, 2213. [CrossRef]
6. Zhu, X.; Xu, Z.; Wang, G.; Cong, Y.; Yu, L.; Jia, R.; Qin, Y.; Zhang, G.; Li, B.; Yuan, D.; et al. Single-cell resolution analysis reveals the preparation for reprogramming the fate of stem cell niche in cotton lateral meristem. *Genome Biol.* **2023**, *24*, 194, Correction in *Genome Biol.* **2023**, *24*, 194. [CrossRef]
7. Yang, Z.; He, J.; Yao, S.; Li, X.; Du, C.; Gong, J.; Yan, H.; Zhao, Y.; Li, Y.; Yuan, Y.; et al. Identification and overexpression of RNA-decapping protein GhLSM1BS: Enhancing cotton somatic embryogenesis through up-regulating brassinosteroid biosynthesis. *Plant Biotechnol. J.* **2025**, *23*, 2670–2672. [CrossRef]

8. Yuan, J.; Liu, X.; Zhao, H.; Wang, Y.; Wei, X.; Wang, P.; Zhan, J.; Liu, L.; Li, F.; Ge, X. GhRCD1 regulates cotton somatic embryogenesis by modulating the GhMYC3–GhMYB44–GhLBD18 transcriptional cascade. *New Phytol.* **2023**, *240*, 207–223. [\[CrossRef\]](#)
9. Gao, F.; Peng, C.; Wang, H.; Shen, H.; Yang, L. Selection of culture conditions for callus induction and proliferation by somatic embryogenesis of *Pinus koraiensis*. *J. For. Res.* **2020**, *32*, 483–491. [\[CrossRef\]](#)
10. Koo, D.; Gil Lee, H.; Bae, S.H.; Lee, K.; Seo, P.J. Callus proliferation-induced hypoxic microenvironment decreases shoot regeneration competence in *Arabidopsis*. *Mol. Plant* **2024**, *17*, 395–408. [\[CrossRef\]](#) [\[PubMed\]](#)
11. Wang, Y.; He, S.; Long, Y.; Zhang, X.; Zhang, X.; Hu, H.; Li, Z.; Hou, F.; Ge, F.; Gao, S.; et al. Genetic variations in *ZmSAUR15* contribute to the formation of immature embryo-derived embryonic calluses in maize. *Plant J.* **2021**, *109*, 980–991. [\[CrossRef\]](#)
12. Wittmer, J.; Pijnenburg, M.; Wijsman, T.; Pelzer, S.; Adema, K.; Kerstens, M.; Kutevska, A.-N.; Fierens, J.; Hofhuis, H.; Sevenier, R.; et al. Rational design of induced regeneration via somatic embryogenesis in the absence of exogenous phytohormones. *Plant Cell* **2025**, *37*, koaf252. [\[CrossRef\]](#)
13. Yin, R.; Chen, R.; Xia, K.; Xu, X. A single-cell transcriptome atlas reveals the trajectory of early cell fate transition during callus induction in *Arabidopsis*. *Plant Commun.* **2024**, *5*, 100941. [\[CrossRef\]](#)
14. Zhang, S.; Zhu, C.; Zhang, X.; Liu, M.; Xue, X.; Lai, C.; Xuhan, X.; Chen, Y.; Zhang, Z.; Lai, Z.; et al. Single-cell RNA sequencing analysis of the embryogenic callus clarifies the spatiotemporal developmental trajectories of the early somatic embryo in *Dimocarpus longan*. *Plant J.* **2023**, *115*, 1277–1297. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Cui, D.; Song, Y.; Jiang, W.; Ye, H.; Wang, S.; Yuan, L.; Liu, B. Genome-wide characterization of the GRF transcription factors in potato (*Solanum tuberosum* L.) and expression analysis of StGRF genes during potato tuber dormancy and sprouting. *Front. Plant Sci.* **2024**, *15*, 1417204. [\[CrossRef\]](#)
16. Zhu, R.; Cao, B.; Sun, M.; Wu, J.; Li, J. Genome-Wide Identification and Evolution of the GRF Gene Family and Functional Characterization of *PbGRF18* in Pear. *Int. J. Mol. Sci.* **2023**, *24*, 14690. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Zheng, F.; Xu, Z.; Deng, X.; Wang, X.; Sun, Y.; Shen, X.; Yu, Z. Genome-Wide Transcriptome Analysis Reveals GRF Transcription Factors Involved in Methyl Jasmonate-Induced Flavonoid Biosynthesis in *Hedera helix*. *Plants* **2025**, *14*, 2094. [\[CrossRef\]](#)
18. Zhang, Z.; Quan, S.; Niu, J.; Guo, C.; Kang, C.; Liu, J.; Yuan, X. Comprehensive Identification and Analyses of the GRF Gene Family in the Whole-Genome of Four Juglandaceae Species. *Int. J. Mol. Sci.* **2022**, *23*, 12663. [\[CrossRef\]](#) [\[PubMed\]](#)
19. Chen, G.-Z.; Huang, J.; Zhou, X.-Q.; Hao, Y.; Chen, J.-L.; Zhou, Y.-Z.; Ahmad, S.; Lan, S.; Liu, Z.-J.; Peng, D.-H. Comprehensive Analysis for GRF Transcription Factors in Sacred Lotus (*Nelumbo nucifera*). *Int. J. Mol. Sci.* **2022**, *23*, 6673. [\[CrossRef\]](#)
20. Angulo, J.; Astin, C.P.; Bauer, O.; Blash, K.J.; Bowen, N.M.; Chukwudinma, N.J.; DiNofrio, A.S.; Faletti, D.O.; Ghulam, A.M.; Gusinde-Duffy, C.M.; et al. CRISPR/Cas9 mutagenesis of the *Arabidopsis* GROWTH-REGULATING FACTOR (GRF) gene family. *Front. Genome Ed.* **2023**, *5*, 1251557. [\[CrossRef\]](#)
21. Kim, J.H.; Tsukaya, H. Regulation of plant growth and development by the GROWTH-REGULATING FACTOR and GRF-INTERACTING FACTOR duo. *J. Exp. Bot.* **2015**, *66*, 6093–6107. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Chandran, V.; Wang, H.; Gao, F.; Cao, X.-L.; Chen, Y.-P.; Li, G.-B.; Zhu, Y.; Yang, X.-M.; Zhang, L.-L.; Zhao, Z.-X.; et al. miR396-OsGRFs Module Balances Growth and Rice Blast Disease-Resistance. *Front. Plant Sci.* **2019**, *9*, 1999. [\[CrossRef\]](#)
23. Lu, Y.; Meng, Y.; Zeng, J.; Luo, Y.; Feng, Z.; Bian, L.; Gao, S. Coordination between GROWTH-REGULATING FACTOR1 and GRF-INTERACTING FACTOR1 plays a key role in regulating leaf growth in rice. *BMC Plant Biol.* **2020**, *20*, 200. [\[CrossRef\]](#)
24. Cao, Y.; Han, Y.; Jin, Q.; Lin, Y.; Cai, Y. Comparative Genomic Analysis of the GRF Genes in Chinese Pear (*Pyrus bretschneideri* Rehd), Poplar (*Populus*), Grape (*Vitis vinifera*), *Arabidopsis* and Rice (*Oryza sativa*). *Front. Plant Sci.* **2016**, *7*, 1750. [\[CrossRef\]](#)
25. Cheng, Z.; Wen, S.; Wu, Y.; Shang, L.; Wu, L.; Lyu, D.; Yu, H.; Wang, J.; Jian, H. Comparatively Evolution and Expression Analysis of GRF Transcription Factor Genes in Seven Plant Species. *Plants* **2023**, *12*, 2790. [\[CrossRef\]](#)
26. Huang, W.; He, Y.; Yang, L.; Lu, C.; Zhu, Y.; Sun, C.; Ma, D.; Yin, J. Genome-wide analysis of growth-regulating factors (GRFs) in *Triticum aestivum*. *PeerJ* **2021**, *9*, e10701. [\[CrossRef\]](#)
27. Lazzara, F.E.; Rodriguez, R.E.; Palatnik, J.F. Molecular mechanisms regulating GROWTH-REGULATING FACTORS activity in plant growth, development, and environmental responses. *J. Exp. Bot.* **2024**, *75*, 4360–4372. [\[CrossRef\]](#)
28. Liu, Z.; Li, N.; Zhang, Y.; Li, Y. Transcriptional repression of GIF1 by the KIX-PPD-MYC repressor complex controls seed size in *Arabidopsis*. *Nat. Commun.* **2020**, *11*, 1846. [\[CrossRef\]](#) [\[PubMed\]](#)
29. Wu, W.; Li, J.; Wang, Q.; Lv, K.; Du, K.; Zhang, W.; Li, Q.; Kang, X.; Wei, H. Growth-regulating factor 5 (GRF5)-mediated gene regulatory network promotes leaf growth and expansion in poplar. *New Phytol.* **2021**, *230*, 612–628. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Wu, W.; Liao, T.; Du, K.; Wei, H.; Kang, X. Transcriptome comparison of different ploidy reveals the mechanism of photosynthetic efficiency superiority of triploid poplar. *Genomics* **2021**, *113*, 2211–2220. [\[CrossRef\]](#) [\[PubMed\]](#)
31. Omidbakhshfard, M.A.; Proost, S.; Fujikura, U.; Mueller-Roeber, B. Growth-Regulating Factors (GRFs): A Small Transcription Factor Family with Important Functions in Plant Biology. *Mol. Plant* **2015**, *8*, 998–1010. [\[CrossRef\]](#)
32. Kim, J.H.; Choi, D.; Kende, H. The AtGRF family of putative transcription factors is involved in leaf and cotyledon growth in *Arabidopsis*. *Plant J.* **2003**, *36*, 94–104. [\[CrossRef\]](#)

33. Liu, Y.; Guo, P.; Wang, J.; Xu, Z. Growth-regulating factors: Conserved and divergent roles in plant growth and development and potential value for crop improvement. *Plant J.* **2023**, *113*, 1122–1145. [\[CrossRef\]](#)
34. Wang, L.; Gu, X.; Xu, D.; Wang, W.; Wang, H.; Zeng, M.; Chang, Z.; Huang, H.; Cui, X. miR396-targeted AtGRF transcription factors are required for coordination of cell division and differentiation during leaf development in *Arabidopsis*. *J. Exp. Bot.* **2010**, *62*, 761–773. [\[CrossRef\]](#)
35. Zhang, L.; Li, G.; Li, Y.; Min, J.; Kronzucker, H.J.; Shi, W. Tomato plants ectopically expressing *Arabidopsis* GRF9 show enhanced resistance to phosphate deficiency and improved fruit production in the field. *J. Plant Physiol.* **2018**, *226*, 31–39. [\[CrossRef\]](#)
36. Kim, J.H. Biological roles and an evolutionary sketch of the GRF-GIF transcriptional complex in plants. *BMB Rep.* **2019**, *52*, 227–238. [\[CrossRef\]](#) [\[PubMed\]](#)
37. Zan, T.; Zhang, L.; Xie, T.; Li, L. Genome-Wide Identification and Analysis of the Growth-Regulating Factor (GRF) Gene Family and GRF-Interacting Factor Family in *Triticum aestivum* L. *Biochem. Genet.* **2020**, *58*, 705–724. [\[CrossRef\]](#) [\[PubMed\]](#)
38. Ma, J.-Q.; Jian, H.-J.; Yang, B.; Lu, K.; Zhang, A.-X.; Liu, P.; Li, J.-N. Genome-wide analysis and expression profiling of the GRF gene family in oilseed rape (*Brassica napus* L.). *Gene* **2017**, *620*, 36–45. [\[CrossRef\]](#)
39. Liu, M.; Zhao, S.; Wang, J.; Tu, Z.; Chen, J. Genome-wide identification and analysis of miR396 family members and their target GRF genes in rubber tree (*Hevea brasiliensis*). *BMC Genom.* **2025**, *26*, 985. [\[CrossRef\]](#) [\[PubMed\]](#)
40. He, F.; Shi, Y.-J.; Zhao, Q.; Zhao, K.-J.; Cui, X.-L.; Chen, L.-H.; Yang, H.-B.; Zhang, F.; Mi, J.-X.; Huang, J.-L.; et al. Genome-wide investigation and expression profiling of polyphenol oxidase (PPO) family genes uncover likely functions in organ development and stress responses in *Populus trichocarpa*. *BMC Genom.* **2021**, *22*, 731. [\[CrossRef\]](#)
41. He, F.; Shi, Y.-J.; Li, J.-L.; Lin, T.-T.; Zhao, K.-J.; Chen, L.-H.; Mi, J.-X.; Zhang, F.; Zhong, Y.; Lu, M.-M.; et al. Genome-wide analysis and expression profiling of Cation/H<sup>+</sup> exchanger (CAX) family genes reveal likely functions in cadmium stress responses in poplar. *Int. J. Biol. Macromol.* **2022**, *204*, 76–88. [\[CrossRef\]](#)
42. Cao, J.-F.; Huang, J.-Q.; Liu, X.; Huang, C.-C.; Zheng, Z.-S.; Zhang, X.-F.; Shangguan, X.-X.; Wang, L.-J.; Zhang, Y.-G.; Wendel, J.F.; et al. Genome-wide characterization of the GRF family and their roles in response to salt stress in *Gossypium*. *BMC Genom.* **2020**, *21*, 575. [\[CrossRef\]](#)
43. Bull, T.; Debernardi, J.; Reeves, M.; Hill, T.; Bertier, L.; Van Deynze, A.; Micheltore, R. GRF–GIF chimeric proteins enhance in vitro regeneration and Agrobacterium-mediated transformation efficiencies of lettuce (*Lactuca* spp.). *Plant Cell Rep.* **2023**, *42*, 629–643. [\[CrossRef\]](#)
44. Ferela, A.; Debernardi, J.M.; Rosatti, S.; Liebsch, D.; Schommer, C.; Palatnik, J.F. Interplay among ZF-HD and GRF transcription factors during *Arabidopsis* leaf development. *Plant Physiol.* **2023**, *191*, 1789–1802. [\[CrossRef\]](#)
45. Baucher, M.; Moussawi, J.; Vandeputte, O.M.; Monteyne, D.; Mol, A.; Pérez-Morga, D.; El Jaziri, M. A role for the miR396/GRF network in specification of organ type during flower development, as supported by ectopic expression of *Populus trichocarpa* miR396c in transgenic tobacco. *Plant Biol.* **2012**, *15*, 892–898. [\[CrossRef\]](#)
46. Li, G.; Chen, Y.; Zhao, X.; Yang, J.; Wang, X.; Li, X.; Hu, S.; Hou, H. Genome-Wide Analysis of the Growth-Regulating Factor (GRF) Family in Aquatic Plants and Their Roles in the ABA-Induced Turion Formation of *Spirodela polyrrhiza*. *Int. J. Mol. Sci.* **2022**, *23*, 10485. [\[CrossRef\]](#)
47. He, F.; Shi, Y.-J.; Chen, Q.; Li, J.-L.; Niu, M.-X.; Feng, C.-H.; Lu, M.-M.; Tian, F.-F.; Zhang, F.; Lin, T.-T.; et al. Genome-Wide Investigation of the PtrCHLP Family Reveals That PtrCHLP3 Actively Mediates Poplar Growth and Development by Regulating Photosynthesis. *Front. Plant Sci.* **2022**, *13*, 870970. [\[CrossRef\]](#)
48. Liu, K.; Kabir, N.; Wei, Z.; Sun, Z.; Wang, J.; Qi, J.; Liu, M.; Liu, J.; Zhou, K. Genome-wide identification and expression profile of GhGRF gene family in *Gossypium hirsutum* L. *PeerJ* **2022**, *10*, e13372. [\[CrossRef\]](#)
49. Li, J.; Wang, M.; Li, Y.; Zhang, Q.; Lindsey, K.; Daniell, H.; Jin, S.; Zhang, X. Multi-omics analyses reveal epigenomics basis for cotton somatic embryogenesis through successive regeneration acclimation process. *Plant Biotechnol. J.* **2018**, *17*, 435–450. [\[CrossRef\]](#)
50. Wen, L.; Li, W.; Parris, S.; West, M.; Lawson, J.; Smathers, M.; Li, Z.; Jones, D.; Jin, S.; Saski, C.A. Transcriptomic profiles of non-embryogenic and embryogenic callus cells in a highly regenerative upland cotton line (*Gossypium hirsutum* L.). *BMC Dev. Biol.* **2020**, *20*, 25. [\[CrossRef\]](#)
51. You, J.; Liu, Z.; Qi, Z.; Ma, Y.; Sun, M.; Su, L.; Niu, H.; Peng, Y.; Luo, X.; Zhu, M.; et al. Regulatory controls of duplicated gene expression during fiber development in allotetraploid cotton. *Nat. Genet.* **2023**, *55*, 1987–1997. [\[CrossRef\]](#)
52. Kong, L.; Ma, L.; Jiang, S.; Zhang, X.; Ma, J.; Pan, M.; Wu, W.; Liu, W.; Ren, W.; Ma, W. Genome-Wide Identification, Exogenous Hormone Response, Gene Structure, and Conserved Motif Analysis of the GRF Gene Family in *Cerasus humilis*. *Biology* **2025**, *14*, 763. [\[CrossRef\]](#)
53. Chen, X.; Zhang, J.; Wang, S.; Cai, H.; Yang, M.; Dong, Y. Genome-wide molecular evolution analysis of the GRF and GIF gene families in Plantae (Archaeplastida). *BMC Genom.* **2024**, *25*, 74. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Lee, S.-J.; Lee, B.H.; Jung, J.-H.; Park, S.K.; Song, J.T.; Kim, J.H. GROWTH-REGULATING FACTOR and GRF-INTERACTING FACTOR Specify Meristematic Cells of Gynoecia and Anthers. *Plant Physiol.* **2017**, *176*, 717–729. [\[CrossRef\]](#)

55. Feng, C.-H.; Niu, M.-X.; Liu, X.; Bao, Y.; Liu, S.; Liu, M.; He, F.; Han, S.; Liu, C.; Wang, H.-L.; et al. Genome-Wide Analysis of the FBA Subfamily of the Poplar F-Box Gene Family and Its Role under Drought Stress. *Int. J. Mol. Sci.* **2023**, *24*, 4823. [[CrossRef](#)] [[PubMed](#)]
56. Feng, C.-H.; Niu, M.-X.; Zhao, S.; Guo, S.; Yin, W.; Xia, X.; Su, Y. Aspartyl tRNA-synthetase (AspRS) gene family enhances drought tolerance in poplar through BABA-PtrIBIs-PtrVOZ signaling module. *BMC Genom.* **2023**, *24*, 473. [[CrossRef](#)]
57. Sievers, F.; Wilm, A.; Dineen, D.; Gibson, T.J.; Karplus, K.; Li, W.; Lopez, R.; McWilliam, H.; Remmert, M.; Söding, J.; et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* **2011**, *7*, 539. [[CrossRef](#)]
58. Hall, B.G. Building Phylogenetic Trees from Molecular Data with MEGA. *Mol. Biol. Evol.* **2013**, *30*, 1229–1235. [[CrossRef](#)]
59. Bailey, T.L.; Boden, M.; Buske, F.A.; Frith, M.; Grant, C.E.; Clementi, L.; Ren, J.; Li, W.W.; Noble, W.S. MEME SUITE: Tools for motif discovery and searching. *Nucleic Acids Res.* **2009**, *37*, w202–w208. [[CrossRef](#)]
60. Chen, C.J.; Chen, H.; Zhang, Y.; Thomas, H.R.; Frank, M.H.; He, Y.H.; Xia, R. TBtools: An Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. *Mol. Plant* **2020**, *13*, 1194–1202. [[CrossRef](#)]
61. Chen, C.; Wu, Y.; Li, J.; Wang, X.; Zeng, Z.; Xu, J.; Liu, Y.; Feng, J.; Chen, H.; He, Y.; et al. TBtools-II: A “one for all, all for one” bioinformatics platform for biological big-data mining. *Mol. Plant* **2023**, *16*, 1733–1742. [[CrossRef](#)]
62. Geer, L.Y.; Marchler-Bauer, A.; Geer, R.C.; Han, L.; He, J.; He, S.; Liu, C.; Shi, W.; Bryant, S.H. The NCBI BioSystems database. *Nucleic Acids Res.* **2009**, *38*, D492–D496. [[CrossRef](#)]
63. Schoch, C.L.; Ciufo, S.; Domrachev, M.; Hotton, C.L.; Kannan, S.; Khovanskaya, R.; Leipe, D.; Mcveigh, R.; O’Neill, K.; Robbertse, B.; et al. NCBI Taxonomy: A comprehensive update on curation, resources and tools. *Database* **2020**, *2020*, baaa062. [[CrossRef](#)] [[PubMed](#)]
64. Wang, Y.; Tang, H.; DeBarry, J.D.; Tan, X.; Li, J.; Wang, X.; Lee, T.-H.; Jin, H.; Marler, B.; Guo, H.; et al. MCScanX: A toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Res.* **2012**, *40*, e49. [[CrossRef](#)]
65. Jiang, M.; Feng, C.; Zhen, J.; Liu, L.; Liu, D.; Zhang, S.; Chi, J. Genome-Wide Identification and Characterization of the VOZ Gene Family in *Gossypium hirsutum* L. and Functional Characterization in Abiotic Stress and Somatic Embryogenesis. *Int. J. Mol. Sci.* **2025**, *26*, 10965. [[CrossRef](#)]
66. Tian, F.; Yang, D.-C.; Meng, Y.-Q.; Jin, J.; Gao, G. PlantRegMap: Charting functional regulatory maps in plants. *Nucleic Acids Res.* **2019**, *48*, D1104–D1113. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.