



Article

LncSMIM14 Hijacks Rab3a-Mediated Endocytosis to Promote Bovine Viral Diarrhea Virus Replication

Zhiran Shao ^{1,†}, Siqi Ma ^{1,†}, FengSiyue Gao ¹, Yang Lou ¹, Xinyi Liu ¹, Li Yang ¹, Zhanhai Mai ¹, Lixia Wang ¹, Areayi Haiyilati ¹, Huijun Shi ^{1,2,*} and Qiang Fu ^{1,2,*}

¹ College of Veterinary Medicine, Xinjiang Agricultural University, Urumqi 830052, China

² Xinjiang Key Laboratory of New Drug Research and Development for Herbivores Animals, Urumqi 830052, China

* Correspondence: shj@xjau.edu.cn (H.S.); fuqiang_xjau@xjau.edu.com (Q.F.)

† First Authors.

Abstract

Bovine Viral Diarrhea Virus (BVDV) poses a significant threat to the global cattle industry, causing substantial economic losses. Long non-coding RNAs (lncRNAs) play crucial regulatory roles in various biological processes, including viral infections. However, the specific lncRNAs influencing BVDV replication remain poorly characterized. This study identified lncSMIM14 as a key host factor upregulated during BVDV infection in MDBK cells. Functional analyses demonstrated that lncSMIM14 overexpression significantly enhanced BVDV replication, evidenced by increased viral mRNA levels, progeny virus titers, cytopathic effects, and dsRNA abundance, while its knockdown exerted the opposite effect. Mechanistically, we revealed that lncSMIM14 specifically targets and positively regulates the expression of the endocytosis-related GTPase Rab3a. Importantly, Rab3a itself was shown to be essential for efficient BVDV replication, as its overexpression promoted viral replication, and its knockdown inhibited it. Furthermore, Rab3a co-localized with key endocytic regulators Rab5a and Rab7a, and both lncSMIM14 overexpression and Rab3a overexpression promoted the formation of endocytic vesicles, particularly post-BVDV infection. Our findings unveil a novel mechanism wherein BVDV exploits the host lncRNA lncSMIM14 to hijack Rab3a-mediated endocytosis, facilitating its own replication. This study identifies the lncSMIM14-Rab3a axis as a critical host pathway subverted by BVDV, providing new potential targets for antiviral intervention.

Keywords: bovine viral diarrhea virus; long non-coding RNA; lncSMIM14; Rab3a; viral replication

Academic Editor: Alfredo Ciccodicola

Received: 8 January 2026
Revised: 10 February 2026
Accepted: 11 February 2026
Published: 27 February 2026

Copyright: © 2026 by the authors. Submitted for possible open access publication under the terms and conditions of the [Creative Commons Attribution \(CC BY\)](#) license.

1. Introduction

Bovine viral diarrhea virus (BVDV) is the primary etiological agent responsible for viral diarrhea and mucosal disease in cattle, conditions collectively referred to as bovine viral diarrhea–mucosal disease (BVD/MD) [1]. As a member of the genus *Pestivirus* within the family *Flaviviridae*, BVDV infection manifests clinically through inflammatory diarrhea, enteritis, and mucosal necrosis [2]. BVDV represents a significant global threat to livestock, imposing a substantial economic burden on the cattle industry worldwide [3]. Although BVDV has evolved sophisticated strategies to establish and maintain host

infection, the underlying molecular pathogenic mechanisms remain only partially understood.

Long non-coding RNAs (lncRNAs), once dismissed as mere “transcriptional noise,” are now defined as transcripts exceeding 200 nucleotides in length that lack protein-coding potential [4]. In recent years, lncRNAs have emerged as pivotal regulators of diverse biological processes, including cell proliferation, differentiation, and development [5,6]. Accumulating evidence underscores the essential roles of lncRNAs in the initiation, modulation, and progression of various viral infections, such as influenza [7], hepatitis [8], coronaviruses [9], and human immunodeficiency virus [10]. Consequently, identifying key host lncRNAs is crucial for elucidating their functional roles, infection-related targets, and the complex regulatory networks they govern [11].

Endocytosis is a fundamental cellular process for internalizing membrane lipids, proteins, and extracellular material, thereby playing a vital role in modulating intracellular signaling cascades [11,12]. Pathogens frequently hijack endocytic pathways to facilitate their entry and internalization within host cells [13]. Central to these processes are Rab proteins, a family of small GTP-binding proteins that serve as master regulators of intracellular membrane trafficking [14]. The mammalian genome encodes over 60 Rab genes, among which Rab3a traditionally acts as a molecular switch—binding GTP and GDP—to regulate synaptic vesicle exocytosis and hormone secretion [15]. However, emerging evidence suggests that Rab GTPases possess remarkable functional versatility, coordinating multiple stages of vesicular trafficking through “Rab cascades” or cross-talk between endocytic and exocytic pathways [16]. Other Rab family members, such as Rab5a, contribute to signal transduction, receptor downregulation, and the phagocytosis of pathogens [17], with studies indicating that Rab5a upregulation can suppress antiviral immune responses in epithelial cells [18]. Similarly, Rab7a plays a critical role in the intracellular transport of viruses [19]. Primarily localized to late endosomes, Rab7a regulates viral spread [20] and exerts bidirectional regulatory functions throughout the viral replication cycle, facilitating host–pathogen interactions that promote infection [21,22].

Despite the scarcity of research regarding the interplay between lncRNAs and BVDV replication, our study provides novel insights into this relationship. Through lncRNA sequencing (accession number: [to be provided]), we discovered that lncSMIM14 is significantly upregulated during BVDV infection. We identified lncSMIM14 as a key proviral lncRNA that is strongly induced by BVDV and essential for efficient viral replication. Mechanistically, lncSMIM14 enhances the expression of Rab3a, which in turn co-localizes with key endocytic regulators Rab5a and Rab7a to facilitate the formation of endocytic vesicles. Our findings reveal a novel mechanism wherein BVDV exploits the host lncRNA lncSMIM14 to hijack Rab3a-mediated endocytosis, thereby optimizing its own replication. This study identifies the lncSMIM14-Rab3a axis as a critical host pathway subverted by BVDV, offering a new molecular perspective on BVDV intracellular replication and providing potential targets for future antiviral interventions.

2. Results

2.1. The Expression of lncSMIM14 Is Significantly Elevated in MDBK Cells After BVDV Infection

To explore the involvement of long non-coding RNAs (lncRNAs) in bovine viral diarrhoea virus (BVDV) infection, we performed RNA sequencing (RNA-seq) to identify differentially expressed lncRNAs in MDBK cells at 6, 12 and 24 h post-infection (hpi) (Figure 1A). The top 13 upregulated lncRNAs identified by RNA-seq were further validated using quantitative real-time PCR (qRT-PCR). The results confirmed that the expression patterns of these 13 candidates were highly consistent with the sequencing data (Figure 1B). Notably, lncSMIM14, lncWIPF3, and lncGPC6 exhibited significant upregulation across all three

time points (Figure 1C). Furthermore, Gene Ontology (GO) and KEGG pathway enrichment analyses revealed that these differentially expressed lncRNAs were predominantly associated with the endocytosis pathway (Figure 1D). We used lncLocator to predict the subcellular localizations of lncSMIM14, lncWIPF3, and lncGPC6, which suggested that lncSMIM14 and lncGPC6 are primarily localized in the cytoplasm, while lncWIPF3 is situated within the cytosol (Table 1).

Given that lncSMIM14 expression demonstrated a stable, time-dependent increase throughout the course of BVDV infection (Figure 1C), it was selected as the primary candidate for further functional investigation.




Figure 1. Upregulation of lncSMIM14 in MDBK cells during BVDV infection. (A) Heatmap analysis of differentially expressed lncRNAs in MDBK cells infected with BVDV for 6, 12, and 24 h. (B) qRT-PCR-based validation of representative differentially expressed lncRNAs identified by RNA-seq. (C) Time-course expression profiles of lncSMIM14, lncWIPF3, and lncGPC6 at 6, 12, and 24 h post-infection (hpi), as determined by qRT-PCR. (D) Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analyses of the consistently upregulated gene across all three time points. Data are presented as the mean $\pm$ standard deviation (SD). Statistical significance is indicated as $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$ (**).

Table 1. Predictive analysis of lncRNA positions by lncLocator.

Subcellular Locations Score	Endocytosis Related lncRNAs		
	lncSMIM14	lncWIPF3	lncGPC6
Cytoplasm	0.369671600163	0.181414846852	0.747400390359
Nucleus	0.287521215258	0.0335444274812	0.167841284931
Ribosome	0.157724942096	0.0785417971393	0.012327340771
Cytosol	0.118447661906	0.678341753356	0.0339009144726
Exosome	0.0666345805775	0.028157175172	0.0385300694608
Predicted location	Cytoplasm	Cytosol	Cytoplasm

2.2. lncSMIM14 Influences Intracellular Replication of BVDV

To elucidate the role of lncSMIM14 in intracellular BVDV replication, we first performed qRT-PCR to assess its expression in MDBK cells following BVDV infection. The results revealed that lncSMIM14 expression increased in a time-dependent manner during the course of infection (Figure 2A). Subsequently, we established lncSMIM14-overexpressing (lncSMIM14 + MDBK) cells (Figure 2B,C) and lncSMIM14-knockdown (lncSMIM14 – MDBK) cells to evaluate its functional impact. lncSMIM14-overexpressing cells exhibited significantly elevated BVDV mRNA levels, higher progeny virus titers, more pronounced cytopathic effects (CPE), and enhanced fluorescence intensity and broader distribution of BVDV dsRNA (Figure 2D–G). Conversely, lncSMIM14 knockdown led to a marked reduction in BVDV mRNA levels, lower progeny virus titers, attenuated CPE, and diminished dsRNA fluorescence (Figure 3B–E). Collectively, our findings demonstrate that lncSMIM14 acts as a proviral factor that promotes intracellular BVDV replication.




Figure 2. lncSMIM14 overexpression enhances intracellular BVDV replication. (A) Time-course expression profile of lncSMIM14 in MDBK cells following BVDV infection, as determined by qRT-PCR. (B) Representative fluorescence microscopy images of HEK-293T cells transfected with the pLVX-AcGFP1-N1 plasmid to evaluate transfection efficiency. (C) qRT-PCR validation of lncSMIM14 overexpression levels in transfected MDBK cells. (D) Quantification of BVDV 5'UTR RNA levels in lncSMIM14-overexpressing MDBK cells at the indicated time points. (E) Progeny virus titers in the culture supernatants determined by TCID₅₀ assay. (F) Representative light microscopy images showing BVDV-induced cytopathic effects (CPE) in MDBK cells. (G) Confocal immunofluorescence analysis of BVDV dsRNA in MDBK cells; nuclei were counterstained with DAPI (blue). Data are presented as the mean $\pm$ standard deviation (SD). Statistical significance is indicated as $p < 0.01$ (**) and $p < 0.001$ (***) and $p > 0.05$ (ns).




Figure 3. Knockdown of lncSMIM14 inhibits intracellular BVDV replication. **(A)** Validation of lncSMIM14 knockdown efficiency in MDBK cells transfected with the pLKO.1-copGFP-PURO-SMIM14 plasmid, as assessed by fluorescence microscopy and qRT-PCR. **(B)** qRT-PCR quantification of BVDV 5'UTR RNA levels in lncSMIM14-silenced MDBK cells at the indicated time points. **(C)** Determination of progeny virus titers in culture supernatants using TCID₅₀ assays. **(D)** Representative light microscopy images showing the attenuation of BVDV-induced cytopathic effects (CPE) in lncSMIM14-knockdown cells. **(E)** Confocal immunofluorescence analysis of BVDV dsRNA (red) in MDBK cells; nuclei were counterstained with DAPI (blue). Statistical significance is indicated as $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$ (***) and $p > 0.05$ (ns).

2.3. lncSMIM14 Targets and Regulates the Endocytosis-Related Protein Rab3a

To explore the molecular mechanism underlying lncSMIM14-mediated BVDV replication, we first predicted its potential target genes through bioinformatics analysis (Figure 4A). Subsequently, the binding probabilities between lncSMIM14 and candidate proteins were assessed using the RNA–protein interaction prediction (RPISeq) tool based on Random Forest (RF) and Support Vector Machine (SVM) scores. This analysis identified Rab3a, an endocytosis-related GTPase, as a high-confidence binding target of lncSMIM14 (Figure 4B). To determine whether lncSMIM14 modulates Rab3a expression, MDBK cells were transfected with Myc, Myc-lncSMIM14, shNC, or sh-lncSMIM14. Overexpression of lncSMIM14 significantly upregulated both Rab3a mRNA and protein levels, whereas its knockdown resulted in a marked reduction in Rab3a expression (Figure 4C–F). Following BVDV infection of these modified cells, Rab3a protein levels showed a gradual increase in lncSMIM14-overexpressing cells, peaking at 24–36 h post-infection (hpi) before slightly declining at 48 hpi. Conversely, Rab3a levels progressively decreased in lncSMIM14-knockdown cells post-infection (Figure 4G–H). Finally, RNA immunoprecipitation (RIP) assays demonstrated that Strep–Rab3a specifically enriched lncSMIM14 compared to the control group (Figure 4I). Taken together, these results confirm a direct regulatory interaction between lncSMIM14 and Rab3a.




Figure 4. LncSMIM14 directly targets and modulates the expression of the endocytic GTPase Rab3a. (A) Bioinformatics-based prediction of the interaction network between LncSMIM14 and differentially expressed mRNAs. (B) RNA-protein interaction prediction (RPISeq) analysis showing RF and SVM scores for LncSMIM14 and candidate mRNAs. (C,D) qRT-PCR quantification of Rab3a mRNA levels in MDBK cells following (C) Myc-tagged LncSMIM14 overexpression or (D) shRNA-mediated LncSMIM14 knockdown. (E) Representative Western blot images showing Rab3a protein levels under conditions of LncSMIM14 overexpression or knockdown. (F) Densitometric analysis of Rab3a protein levels normalized to GAPDH. (G,H) Time-course Western blot analysis of Rab3a protein expression in (G) LncSMIM14-overexpressing or (H) LncSMIM14-knockdown MDBK cells at the indicated hours post-infection (hpi) with BVDV. (I) RNA immunoprecipitation (RIP) assay demonstrating the physical interaction between Strep-tagged Rab3a and LncSMIM14. Data are presented as the mean $\pm$ standard deviation (SD). Statistical significance is indicated as $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$ (***).

2.4. The Endocytosis-Related Protein Rab3a Promotes Intracellular Replication of BVDV

To evaluate the involvement of Rab3a in BVDV replication, we first monitored its expression levels in MDBK cells following BVDV infection. qRT-PCR analysis revealed that Rab3a mRNA levels were significantly upregulated at 48 h post-infection (hpi) ($p < 0.05$) (Figure 5A). These findings were corroborated by Western blot analysis, which showed a consistent trend in Rab3a protein expression (Figure 5B). Subsequently, we established Rab3a-overexpressing MDBK cells (mCherry-Rab3a) (Figure 5C–G). Upon BVDV infection, these cells exhibited significantly elevated BVDV mRNA levels, increased progeny virus titers, more pronounced cytopathic effects (CPE), and enhanced fluorescence intensity and broader distribution of BVDV dsRNA (Figure 5H). Conversely, Rab3a-knockdown cells (Rab3a-sg1) (Figure 6A–D) displayed markedly reduced BVDV

mRNA levels, lower progeny virus titers, attenuated CPE, and diminished dsRNA fluorescence (Figure 6E–G). Collectively, these results demonstrate that Rab3a acts as a critical host factor that facilitates the intracellular replication of BVDV.




Figure 5. Rab3a overexpression promotes intracellular BVDV replication. (A) Time-course qRT-PCR analysis of Rab3a mRNA levels in MDBK cells following BVDV infection. (B) Representative Western blot images and densitometric quantification of Rab3a protein levels normalized to GAPDH during BVDV infection. (C) Representative fluorescence images of HEK-293T cells transfected with mCherry or mCherry-Rab3a plasmids (Brightfield, mCherry channel, and merged images), with validation of Rab3a mRNA expression by qRT-PCR. (D) Western blot analysis and densitometric quantification of Rab3a protein levels in MDBK cells following transfection with mCherry or mCherry-Rab3a. (E) qRT-PCR quantification of BVDV 5'UTR RNA levels in Rab3a-overexpressing cells at the indicated time points. (F) Determination of progeny virus titers in culture supernatants via TCID₅₀ assays. (G) Representative light microscopy images showing BVDV-induced cytopathic effects (CPE) in cells transfected with mCherry or mCherry-Rab3a at the indicated hours post-infection (hpi). (H) Confocal immunofluorescence analysis of BVDV dsRNA (red) in transfected cells; nuclei were counterstained with DAPI (blue). Data are presented as the mean ± standard deviation (SD). Statistical significance is indicated as $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$ (***).




Figure 6. Knockdown of Rab3a inhibits intracellular BVDV replication. (A,B) Representative Western blot (A) and densitometric quantification (B) of Rab3a protein levels in MDBK cells transfected with Rab3a-sg1 to evaluate knockdown efficiency. GAPDH was used as a loading control. (C) qRT-PCR validation of Rab3a mRNA silencing in Rab3a-sg1-transfected cells. (D) qRT-PCR quantification of BVDV 5'UTR RNA levels in Rab3a-knockdown cells at the indicated time points. (E) Progeny virus titers in culture supernatants determined by TCID₅₀ assays. (F) Representative light microscopy images showing the attenuation of BVDV-induced cytopathic effects (CPE) in Rab3a-sg1-transfected cells at the indicated hours post-infection (hpi). (G) Confocal immunofluorescence analysis of BVDV dsRNA (red) in Rab3a-sg1-transfected cells; nuclei were counterstained with DAPI (blue). Data are presented as the mean $\pm$ standard deviation (SD). Statistical significance is indicated as $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$ (***).

2.5. Rab3a Co-Localizes with Key Endocytic Regulators Rab5a and Rab7a and Promotes Endocytic Vesicle Formation

To further delineate the role of Rab3a in BVDV replication, we examined its spatial relationship with the key endocytic GTPases, Rab5a and Rab7a. Following the co-transfection of mCherry-tagged Rab3a with either mCherry-tagged Rab5a or Rab7a plasmids, confocal laser scanning microscopy revealed distinct spatial co-localization among these proteins in HEK-293T cells (Figure 7A), confirming the close association of Rab3a with the endocytic pathway.

We next investigated whether Rab3a modulates endocytic vesicle formation during BVDV infection. Cells transfected with mCherry-Rab3a or Myc-IncSMIM14 were subjected to mock or BVDV infection, and endocytic vesicles were visualized using transmission electron microscopy (TEM). While BVDV infection alone significantly increased the

number of endocytic vesicles compared to the mock control, the overexpression of either Rab3a or lncSMIM14 further stimulated intracellular endocytic vesicle formation in the context of infection (Figure 7B).

Collectively, these findings indicate that lncSMIM14 acts directly on Rab3a, which in turn coordinates with Rab5a and Rab7a to promote endosome maturation and formation, thereby facilitating efficient intracellular BVDV replication.




Figure 7. Rab3a co-localizes with endocytic regulators Rab5a and Rab7a and enhances endocytic vesicle formation. (A) Confocal laser scanning microscopy images showing the spatial co-localization of mCherry–Rab3a (green) with mCherry–tagged Rab5a or Rab7a (red) in HEK-293T cells. Nuclei were counterstained with DAPI (blue). (B) Transmission electron microscopy (TEM) micrographs and quantitative analysis of endocytic vesicle formation in MDBK cells overexpressing Rab3a or lncSMIM14, under mock or BVDV infection conditions. Arrows indicate representative endocytic vesicles.

3. Discussion

Endocytosis is a fundamental process for numerous intracellular signaling pathways [23]. Pathogens frequently exploit these pathways to facilitate host cell entry, often leading to severe or persistent infections [24]. In this study, we identified lncSMIM14—a host long non-coding RNA significantly induced by BVDV in MDBK cells—as a novel proviral factor that augments intracellular BVDV replication by modulating host endocytic dynamics. Mechanistically, lncSMIM14 specifically interacts with Rab3a, which in turn orchestrates with key endocytic regulators Rab5a and Rab7a to promote endosome formation.

Our findings are concordant with the expanding paradigm implicating lncRNAs in the complex orchestration of viral life cycles. For instance, lncRNA-CMPK2 and lncRNA NRAV have been shown to promote the replication of HCV and IAV, respectively [25,26], while lncRNA GAS5 acts as an antiviral effector against HCV [27]. In the context of BVDV, previous research suggested that host lncRNAs modulate replication primarily through apoptotic pathways [28]. Our study expands this landscape by demonstrating that lncSMIM14 facilitates the assembly or coordination of endocytosis-related proteins (Rab3a, Rab5a, and Rab7a), thereby catalyzing endocytic vesicle formation. While our current data do not fully delineate the boundaries between initial viral entry and post-entry trafficking, the overall enhancement of the endocytic machinery clearly bolsters the broader BVDV life cycle.

A pivotal discovery of this study is the positive regulatory role of Rab3a in BVDV replication. Traditionally associated with exocytosis and synaptic vesicle trafficking, Rab3a is shown here to be “hijacked” by BVDV via lncSMIM14 induction. By directly

binding to Rab3a, lncSMIM14 leverages Rab3a's spatial association with Rab5a and Rab7a to accelerate endosome maturation, thereby optimizing early viral transport. This non-canonical function is supported by the observed co-localization of Rab3a with endocytic markers (Figure 7A) and is consistent with the "fluid" functional boundaries of Rab proteins in non-neuronal cells. Indeed, the involvement of the Rab3 family in endocytosis is corroborated by findings that Rab3D associates with early endosomes in specific epithelial models [29,30].

The roles of Rab5a and Rab7a in viral infections are multifaceted, ranging from inhibiting rabies virus nucleoprotein expression to facilitating HIV-1 transmission and HBV endocytosis [31,32]. Our observation that Rab3a co-localizes with these proteins suggests a sophisticated reciprocal regulatory relationship among BVDV, lncSMIM14, and the Rab3a-mediated pathway. However, several questions remain. Further studies are required to elucidate the precise interaction sites between lncSMIM14 and Rab3a, as well as the specific molecular triggers for endosome production. Additionally, while we focus on the overall promotion of replication, future research employing direct viral entry assays (e.g., virus attachment and internalization tracking) will be essential to dissect whether the lncSMIM14–Rab3a axis primarily triggers initial entry or governs subsequent intracellular trafficking.

In summary, we describe a novel molecular mechanism wherein BVDV induces host lncSMIM14 to subvert the Rab3a-mediated endocytic pathway, ultimately facilitating efficient viral replication. This lncSMIM14–Rab3a axis represents a potential target for future antiviral interventions against *Pestivirus* infections.

4. Materials and Methods

4.1. Cell Culture and Virus

Madin–Darby bovine kidney (MDBK) cells and human embryonic kidney (HEK-293T) cells were obtained from the Cell Bank of the Chinese Academy of Sciences (Shanghai, China). Cells were maintained in Dulbecco's modified Eagle medium (DMEM; Thermo Fisher Scientific, Waltham, USA) supplemented with 10% fetal bovine serum (FBS, Gibco, Grand Island, USA), 100 U/mL penicillin, and 100 µg/mL streptomycin at 37 °C in a humidified incubator with 5% CO₂. Transfections were performed using a DNA transfection reagent (Roche Diagnostics, Basel, Switzerland) according to the manufacturer's protocol. The bovine viral diarrhea virus (BVDV) TC strain was preserved in our laboratory (Xinjiang University, Urumqi, China).

4.2. Plasmids

The plasmids pLVX-Myc-IRES-Puro, pLVX-Myc-IRES-Puro-SMIM14, pLKO.1-copGFP-PURO-SMIM14-1/-2/-3/-4, pLKO.1-copGFP-PURO-NC, pCDH-CMV-MCS-EF1-Blast-mCherry-puro, pCDH-CMV-MCS-EF1-Blast-mCherry-Rab3a-puro, pCDH-CMV-MCS-EF1-Blast-mCherry-Rab5a-puro, and pCDH-CMV-MCS-EF1-Blast-mCherry-Rab7a-puro were purchased from QIAGEN. Plasmids pLentiCRISPR v2-Rab3a-1-sgRNA, pLentiCRISPR v2-Rab3a-2-sgRNA, and pLentiCRISPR v2-Scramble-sgRNA were synthesized by GenScript (Nanjing, China). Plasmids psPAX2 and pMD2.G were obtained from Hunan Fenghui Biotechnology Co., Ltd. Competent *Escherichia coli* Stbl3 cells were prepared and stored in our laboratory.

4.3. LncRNA Sequencing Analysis

Total RNA was extracted using TRIzol reagent (TIANGEN, Beijing, China) according to the manufacturer's protocol. RNA quality and integrity were assessed by agarose gel electrophoresis to ensure suitability for library construction. Sequencing libraries were

prepared with the TruSeq PE Cluster Kit v3-cBot-HS (Illumina) and sequenced on the Illumina HiSeq 4000 platform by Novogene Company, Ltd. (Beijing, China). Raw reads were quality-filtered to remove low-quality sequences. Differentially expressed lncRNAs and mRNAs (p -value < 0.05) were subjected to GO term enrichment (GOseq) and KEGG pathway analyses. Selected transcripts were validated by qRT-PCR to confirm concordance with RNA-Seq results.

4.4. Bioinformatics Analysis and Target Prediction

Bioinformatics analysis and target prediction: The potential target genes of lncSMIM14 were predicted using the LncTar and RNAplex databases to identify potential RNA-RNA interactions. To further evaluate the binding probability between lncSMIM14 and candidate proteins, we employed the RPISeq (RNA-Protein Interaction Prediction) web server. Predictions were based on two machine learning algorithms: Random Forest (RF) and Support Vector Machine (SVM). A threshold of >0.5 for both RF and SVM scores was used to identify high-confidence interactions. Among the top-ranked candidates, several proteins involved in vesicular trafficking and immune regulation were identified, including Rab3a, Rab11, and Annexin A2.

4.5. Virus Infection

MDBK cells were seeded at 5×10^4 cells/dish in 10 cm dishes. When cells reached ~70% confluence, the medium was replaced with serum-free DMEM, and BVDV was inoculated at 1000 TCID₅₀/mL. After 2 h adsorption, the inoculum was replaced with DMEM containing 2% horse serum. Cells were washed with PBS and collected at 6, 12, and 24 h post-infection (hpi).

4.6. Western Blotting

Cells were lysed in RIPA buffer (Roche Diagnostics, Basel, Switzerland) containing protease inhibitors. Protein concentration was determined using a Bio-Rad protein assay (Bio-Rad Laboratories, Hercules, USA). Equal amounts of protein (30 µg) were separated by SDS-PAGE, transferred to PVDF membranes (GE Healthcare, Chicago, USA), and blocked with 5% skim milk (BD Biosciences, Franklin Lakes, USA). Membranes were incubated with primary antibodies against dsRNA (J2, Scicons, Szirák, Hungary), GAPDH (Wuhan Sanying Biotechnology, Wuhan, China), or Rab3a (Wuhan Sanying Biotechnology, Wuhan, China), followed by HRP-conjugated secondary antibodies (Wuhan Sanying Biotechnology, Wuhan, China). Blots were visualized using BeyoECL Plus (Biosharp Biotechnology, Hefei, China).

4.7. Quantitative Real-Time PCR (qRT-PCR)

Total RNA was extracted using TRNzol reagent (TIANGEN Biotech, Beijing, China) and reverse transcribed with the ReverTra Ace qPCR RT Kit (Toyobo, Osaka, Japan). qRT-PCR was performed using SYBR Green qPCR Master Mix (GenStar, Beijing, China) on a QuantStudio system (Applied Biosystems, Foster City, USA).

4.8. Lentiviral Packaging

HEK-293T cells were seeded in 10 cm dishes one day prior to transfection to reach 80–90% confluence. Packaging was performed by co-transfecting the transfer plasmid (e.g., pLV-EF1α-GFP, 10 µg), psPAX2 (7.5 µg), and pMD2.G (2.5 µg) using polyethyleneimine (PEI). Medium was replaced after 6 h. Viral supernatants were harvested at 48 h and 72 h post-transfection, clarified by centrifugation (3000× g , 10 min, 4 °C), concentrated by ultracentrifugation (70,000× g , 2 h, 4 °C), and resuspended in PBS. Viral titers were determined by fluorescence reporter assay (TU/mL).

4.9. Immunofluorescence Microscopy

MDBK cells on coverslips were fixed in 4% paraformaldehyde (15 min), permeabilized with 0.25% Triton X-100 (5 min), and blocked with 5% BSA. Cells were incubated with primary antibodies, followed by fluorophore-conjugated secondary antibodies. Nuclei were counterstained with DAPI. Images were acquired using a Zeiss LSM 780 confocal microscope.

4.10. RNA Immunoprecipitation (RIP)

Cells were crosslinked with 2% formaldehyde and lysed in RIPA buffer (50 mM Tris, pH 7.4; 150 mM NaCl; 1 mM EDTA; 0.1% SDS; 1% NP-40; 0.5% sodium deoxycholate; 0.5 mM DTT; 1 mM PMSF). Lysates were incubated with antibodies and Protein A/G magnetic beads (Thermo Fisher Scientific) at 4 °C. Beads were washed sequentially with RIPA buffer and 1 M RIPA buffer, then treated with proteinase K. RNA was extracted using TRIzol and analyzed by qRT-PCR.

4.11. Statistical Analysis

All experiments were independently repeated at least three times. Data are presented as mean $\pm$ SEM. Statistical significance was determined using the Mann–Whitney U test in SPSS 19.0 software. Differences were considered significant at $p < 0.05$ ($p < 0.05$; * $p < 0.01$; ns, not significant).

5. Conclusions

In conclusion, this study provides the first evidence that the long non-coding RNA lncSMIM14 acts as a pivotal modulator of Bovine Viral Diarrhea Virus (BVDV) replication by orchestrating the endocytic pathway. Our findings demonstrate that the BVDV-induced upregulation of lncSMIM14 augments viral replication, a process coupled with the elevated expression of the GTPase Rab3a. Conversely, lncSMIM14 knockdown attenuates BVDV replication and concomitantly reduces Rab3a levels, suggesting that Rab3a expression is a critical determinant of BVDV replication dynamics. Furthermore, the spatial colocalization of Rab3a with key endocytic regulators Rab5a and Rab7a supports a mechanistic model in which the lncSMIM14–Rab3a axis facilitates viral entry or intracellular trafficking via the endocytic pathway. Overall, these insights expand our understanding of host lncRNA functions during *Pestivirus* infection and identify the lncSMIM14–Rab3a axis as a promising antiviral target for the development of future therapeutic strategies.

Author Contributions: Z.S. contributed to conceptualization, methodology, investigation, and writing (original draft). S.M. performed investigation, validation, and formal analysis. F.G., Y.L. and X.L. were involved in software development, formal analysis, and visualization. L.Y., Z.M., L.W., and A.H. provided resources, writing (review and editing), and supervision. H.S. and Q.F. were responsible for conceptualization, writing (review and editing), supervision, project administration, and funding acquisition. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by the Autonomous Region's 'Tianshan Talents' Young Top Scientific and Technological Talent Program (Project No.: 2022TSYCCX0049); the Central Guidance Local Science and Technology Development Special Fund Project—'Integration and Demonstration of Rapid Screening Technologies for Common Pathogens in Calves and Lambs'; and the National Natural Science Foundation Regional Fund Project (Project No.: 32160829).

Data Availability Statement: The sequence data supporting this study have been deposited in the NCBI database, with the main accession number being PRJNA1333864. (<https://data-view.ncbi.nlm.nih.gov/object/PRJNA1333864?reviewer=rinuh739tos9nmje5tivcjc3v7>; accessed on 24 August 2025).

Conflicts of Interest: The authors declare that they have no competing interests.

References

1. Yeşilbağ, K.; Alpay, G.; Becher, P. Variability and Global Distribution of Subgenotypes of Bovine Viral Diarrhea Virus. *Viruses* **2017**, *9*, 128.
2. Chase, C.C.L.; Thakur, N.; Darweesh, M.F.; Morarie-Kane, S.E.; Rajput, M.K. Immune response to bovine viral diarrhea virus—Looking at newly defined targets. *Anim. Health Res. Rev.* **2015**, *16*, 4–14.
3. Zhou, Y.; Zhang, J.; Wu, H.; Zhao, S.; Ren, Y.; Chen, Q.; Zhang, Z.; Liao, X.; Mo, Y.; Zhong, Y.; et al. First linear B-cell epitope identified on the nucleocapsid protein of bovine coronavirus. *Virology* **2025**, *610*, 110581.
4. Bertone, P.; Stolc, V.; Royce, T.E.; Rozowsky, J.S.; Urban, A.E.; Zhu, X.; Rinn, J.L.; Tongprasit, W.; Samanta, M.; Weissman, S.; et al. Global identification of human transcribed sequences with genome tiling arrays. *Science* **2004**, *5705*, 2242–2246.
5. Fatica, A.; Bozzoni, I. Long non-coding RNAs: New players in cell differentiation and development. *Nat. Rev. Genet.* **2014**, *15*, 7–21.
6. Yang, X.; Yuan, Z.; Gou, L.; Cheng, L.; Wang, Z.; Wu, P.; Wang, X.; Ma, X.; Ma, T.; Yu, Y.; et al. FEZF1-AS1 drives autophagy-mediated progression of colon cancer and reduces chemosensitivity through inhabiting the PI3K/AKT/mTOR signaling pathway. *Front. Genet.* **2025**, *16*, 1514205.
7. Sarma, A.; Suri, P.; Justice, M.; Angamuthu, R.; Pushparaj, S. An Emphasis on the Role of Long Non-Coding RNAs in Viral Gene Expression, Pathogenesis, and Innate Immunity in Viral Chicken Diseases. *Non-Coding RNA* **2025**, *11*, 42.
8. dos Santos, D.B.; Fernandez, G.J.; Silva, L.T.; Silva, G.F.; Lima, E.O.; Galvani, A.F.; Pereira, G.L.; Ferrasi, A.C. lncRNAs as Biomarkers of Hepatocellular Carcinoma Risk and Liver Damage in Advanced Chronic Hepatitis C. *Curr. Issues Mol. Biol.* **2025**, *47*, 348.
9. Yan, W.; Fu, X.; Li, H.; Wang, K.; Song, C.; Hou, C.; Lei, C.; Wang, H.; Yang, X. The long non-coding RNA lncRNA-DRNR enhances infectious bronchitis virus replication by targeting chicken JMJD6 and modulating interferon-stimulated genes expression via the JAK-STAT signalling pathway. *Vet. Res.* **2024**, *55*, 141.
10. Mahajan, A.; Chandran, S.; Kumar, A.; Chen, L.; Dhillon, N.K. lncRNA-536 and RNA-Binding Protein RBM25 Interactions in Pulmonary Artery Smooth Muscle Cells: Implications in Pulmonary Hypertension. *Arterioscler. Thromb. Vasc. Biol.* **2025**, *45*, e374–e391.
11. Gao, C.; Li, H.; Zhuang, J.; Zhang, H.; Wang, K.; Yang, J.; Liu, C.; Liu, L.; Zhou, C.; Sun, C. The construction and analysis of ceRNA networks in invasive breast cancer: A study based on The Cancer Genome Atlas. *Cancer Manag. Res.* **2019**, *11*, 1–11.
12. Kaplan, N.; Ventrella, R.; Peng, H.; Pal-Ghosh, S.; Arvanitis, C.; Rappoport, J.Z.; Mitchell, B.J.; Stepp, M.A.; Lavker, R.M.; Getsios, S. EphA2/Ephrin-A1 Mediate Corneal Epithelial Cell Compartmentalization via ADAM10 Regulation of EGFR Signaling. *Investig. Ophthalmol. Vis. Sci.* **2018**, *59*, 393–406.
13. Mark, M.; Ari, H. Virus entry: Open sesame. *Cell* **2006**, *124*, 729–740.
14. Sharma, R.; Maity, S.K.; Chakrabarti, P.; Katika, M.R.; Kapettu, S.; Parsa, K.V.L.; Misra, P. PIMT Controls Insulin Synthesis and Secretion through PDX1. *Int. J. Mol. Sci.* **2023**, *24*, 8084.
15. Coppola, T.; Perret-Menoud, V.; Gattesco, S.; Magnin, S.; Pombo, I.; Blank, U.; Regazzi, R. The death domain of Rab3 guanine nucleotide exchange protein in GDP/GTP exchange activity in living cells. *Biochem. J.* **2002**, *362*, 273–279.
16. Stenmark, H. Rab GTPases as coordinators of vesicle traffic. *Nat. Rev. Mol. Cell Biol.* **2009**, *10*, 513–525.
17. Lemahieu, G.; Moreno-Layseca, P.; Hub, T.; Bevilacqua, C.; Gómez-González, M.; Pennarola, F.; Colombo, F.; Massey, A.E.; Barzaghi, L.; Palamidessi, A.; et al. RAB5A Promotes Active Fluid Wetting by Reprogramming Breast Cancer Spheroid Mechanics. *Adv. Sci.* **2025**, *12*, E03569.
18. Mo, S.; Tang, W.; Xie, J.; Chen, S.; Ren, L.; Zang, N.; Xie, X.; Deng, Y.; Gao, L.; Liu, E. Respiratory Syncytial Virus Activates Rab5a to Suppress IRF1-Dependent Lambda Interferon Production, Subverting the Antiviral Defense of Airway Epithelial Cells. *J. Virol.* **2021**, *95*, e02333–20.
19. Wong, W.H.; Liu, S.Z.; Li, A.S.R.; Liu, X.; Manolson, M.F.; Zirngibl, R.A. Evidence for Rab7b and Its Splice Isoforms Having Distinct Biological Functions from Rab7a. *Int. J. Mol. Sci.* **2025**, *26*, 2610.
20. Caillet, M.; Janvier, K.; Pelchen-Matthews, A.; Delcroix-Genête, D.; Camus, G.; Marsh, M.; Berlioz-Torrent, C. Rab7A Is Required for Efficient Production of Infectious HIV-1. *PLoS Pathog.* **2011**, *7*, e1002347.
21. Yang, C.; Cheng, Y.; Zhang, T.; Gebeyew, K.; Fischer-Tlustos, A.; Guan, L.; Steele, M.; Tan, Z.; He, Z. Multi-omics analysis provides new insights into the molecular mechanisms underlying colostral immunoglobulin G absorption in the gut of neonatal goat kids. *Anim. Nutr. (Zhongguo Xu Mu Shou Yi Xue Hui)* **2025**, *21*, 166–178.

22. Serres, M.P.; Shaughnessy, R.; Escot, S.; Hammich, H.; Cuvelier, F.; Salles, A.; Rocancourt, M.; Verdon, Q.; Gaffuri, A.-L.; Sourigues, Y.; et al. MiniBAR/GARRE1 is a dual Rac and Rab effector required for ciliogenesis. *Dev. Cell* **2023**, *58*, 2477–2494.e8.
23. Hancock, L.P.; Allwood, E.G.; Palmer, J.S.; Ayscough, K.R.; Williamson, M.P. Agent-based modelling of the early stages of actin polymerisation required to drive endocytosis in *Saccharomyces cerevisiae*. *Sci. Rep.* **2025**, *15*, 28951.
24. Santamaria, A.; Pereira, D.; Pawar, N.; Kelly, B.T.; Carrascosa-Tejedor, J.; Sardo, M.; Mafra, L.; Fragneto, G.; Owen, D.J.; Marín-Montesinos, I.; et al. Structural molecular details of the endocytic adaptor protein CALM upon binding with phosphatidylinositol 4,5-bisphosphate-containing model membranes. *Commun. Chem.* **2025**, *8*, 219.
25. Kambara, H.; Niazi, F.; Kostadinova, L.; Moonka, D.K.; Siegel, C.T.; Post, A.B.; Carnero, E.; Barriocanal, M.; Fortes, P.; Anthony, D.D.; et al. Negative regulation of the interferon response by an interferon-induced long non-coding RNA. *Nucleic Acids Res.* **2014**, *42*, 10668–10680.
26. Baysal, A.Ç.; Kıymaz, Y.Ç.; Şahin, N.Ö.; Bakır, M. Investigation of Long Noncoding RNA-NRAV and Long Noncoding RNA-Lethe Expression in Crimean–Congo Hemorrhagic Fever. *J. Med. Virol.* **2024**, *96*, e70142; Correction in *J. Med. Virol.* **2025**, *97*, e70245.
27. Guo, Y.; Li, C.; Zhang, R.; Zhan, Y.; Yu, J.; Tu, J.; Zheng, J. Epigenetically-regulated serum GAS5 as a potential biomarker for patients with chronic hepatitis B virus infection. *Cancer Biomark. Sect. A Dis. Markers* **2021**, *32*, 137–146.
28. Ma, Q.; Li, L.; Tang, Y.; Fu, Q.; Liu, S.; Hu, S.; Qiao, J.; Chen, C.; Ni, W. Analyses of long non-coding RNAs and mRNA profiling through RNA sequencing of MDBK cells at different stages of bovine viral diarrhea virus infection. *Res. Vet. Sci.* **2017**, *115*, 508–516.
29. He, H.; Ai, R.; Fang, E.F.; Palikaras, K. The Rab3 family proteins in age-related neurodegeneration: Unraveling molecular pathways and potential therapeutic targets. *npj Aging* **2025**, *11*, 64.
30. Sun, Y.; Zhang, M.; Ge, L. A RAB transition orchestrates membrane trafficking in unconventional protein secretion. *J. Cell Biol.* **2024**, *223*, e202312096.
31. Zeyen, L.; Prange, R. Host Cell Rab GTPases in Hepatitis B Virus Infection. *Front. Cell Dev. Biol.* **2018**, *6*, 154.
32. Bartusch, C.; Döring, T.; Prange, R. Rab33B Controls Hepatitis B Virus Assembly by Regulating Core Membrane Association and Nucleocapsid Processing. *Viruses* **2017**, *9*, 157.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.