

## Article

# Automated Compactness Quantitative Metrics for Wrist Bone on Conventional Radiography in Rheumatoid Arthritis: A Clinical Evaluation Study

Jiajing Zhou <sup>1</sup>, Junmu Peng <sup>1</sup>, Haolin Wang <sup>1</sup>, Hiroshi Kataoka <sup>2</sup>, Masaya Mukai <sup>3</sup>, Tunlada Wiriyanukhroh <sup>1,4</sup> and Tamotsu Kamishima <sup>1,\*</sup>

<sup>1</sup> Faculty/Graduate School of Health Sciences, Hokkaido University, Sapporo 060-0812, Hokkaido, Japan; zzzjjj99@163.com (J.Z.); junmupeng@kamiken.net (J.P.); haolin.wang.k3@elms.hokudai.ac.jp (H.W.); tunlada.wi@gmail.com (T.W.)

<sup>2</sup> Sapporo City General Hospital, Sapporo 060-8604, Hokkaido, Japan; hiroshi.kataoka@doc.city.sapporo.jp

<sup>3</sup> Soen Chuo Hospital, Sapporo 621-1023, Hokkaido, Japan; mmukai@cocoa.ocn.ne.jp

<sup>4</sup> Phramongkutklao Hospital and College of Medicine, Bangkok 10400, Thailand

\* Correspondence: ktamotamo2@hs.hokudai.ac.jp

## Abstract

Rheumatoid arthritis (RA) frequently affects the joints of the hands, with joint space narrowing (JSN) representing an important early marker of structural damage. The semi-quantitative Sharp/van der Heijde (SvdH) scoring system is widely used in clinical practice but is inherently subjective and susceptible to observer variability. Moreover, the complex anatomy of the wrist and substantial overlap of carpal bones pose challenges for automated quantitative assessment of wrist JSN on routine radiographs. This study aimed to introduce a novel quantitative assessment perspective and to clinically validate an automated, compactness-related quantification framework for evaluating wrist JSN in RA. This study initially enrolled 51 patients with RA. After excluding one case with severe carpal fusion that precluded anatomical differentiation, 50 patients (44 females and 6 males) were included in the final analysis. The cohort had a mean age of 61 years (range: 21–82), a median symptom duration of 9 years (IQR: 1–32), and a median follow-up interval for bilateral hand radiographs of 1.06 years (IQR: 0.82–1.30). To quantify global wrist JSN, 10 compactness-related metrics were computed based on the spatial distribution of bone centroids extracted from carpal segmentation masks. These metrics were validated against the wrist JSN subscore of the SvdH score (SvdH-JSN\_wrist) and the total Sharp score (TSS) as gold standards. Several distance-based metrics among the compactness-related metrics showed significant negative correlations with the wrist joint space narrowing subscore of the Sharp/van der Heijde score (SvdH-JSN\_wrist). Specifically, mean-pairwise-distance (MPD), root-mean-square-radius (RMSR), and median-radius (R50) showed moderate to strong correlations ( $r = -0.52$  to  $-0.63$ , all  $p \leq 0.0001$ ) that were consistent at BL and FU. Correlations with TSS were weaker overall, with only R50 and its normalized form showing stable negative correlations ( $r = -0.40$  to  $-0.43$ ,  $p < 0.01$ ). Longitudinal analyses showed limited correlations between metric changes and clinical score changes. The proposed automated compactness quantification framework enables objective and reliable assessment of wrist JSN on standard radiographs and complements conventional scoring systems by supporting automated and standardized evaluation of RA-related wrist structural changes.



Academic Editor: Leonardo Rundo

Received: 10 January 2026

Revised: 11 February 2026

Accepted: 13 February 2026

Published: 18 February 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and

conditions of the [Creative Commons](#)

[Attribution \(CC BY\)](#) license.

**Keywords:** rheumatoid arthritis; computer-aided diagnosis; deep learning; conventional radiography

## 1. Introduction

Rheumatoid arthritis (RA) is a chronic autoimmune disease characterized by synovial inflammation and clinically manifested as symmetrical polyarthritis [1]. It predominantly affects small joints, particularly those of the hands and wrists [2]. As the disease progresses, structural damage such as bone erosion (BE) and joint space narrowing (JSN) develops, eventually leading to joint deformity and functional impairment in severe cases [3,4]. RA substantially compromises quality of life and imposes a significant socioeconomic burden, with a reported global prevalence of approximately 0.5–1% [2,5,6]. Early detection, accurate diagnosis, and longitudinal monitoring are therefore essential to prevent irreversible joint damage and disability.

The wrist joint is among the earliest and most frequently affected anatomical sites in RA [7]. Radiographic findings such as JSN and BE reflect early structural progression and remain central to disease assessment [8]. Conventional radiography continues to be the most widely used imaging modality in RA because of its low cost, broad availability, and standardized role in diagnosis, staging, and long-term follow-up [9,10]. However, quantitative assessment of wrist JSN on radiographs is particularly challenging. The wrist consists of multiple small carpal bones with complex morphology and close spatial relationships, and radiographs provide only two-dimensional projections. As a result, manual evaluation is highly susceptible to projection variability, bone overlap, and reader subjectivity [11,12]. The Sharp/van der Heijde (SvdH) scoring system is currently the most widely accepted radiographic scoring method for RA [10,13]. Its total score (TSS) integrates JSN and BE in the hands and feet, reflecting cumulative structural damage and long-term prognosis [14–16]. Despite its clinical utility, SvdH scoring is semi-quantitative, labor-intensive, and subject to inter- and intra-observer variability [10,17]. Although direct manual measurements can provide quantitative information, they are often impractical in routine workflows and insufficiently sensitive to subtle, continuous morphological changes, particularly within the anatomically complex wrist joint. Consequently, there is a clear need for automated quantitative approaches that can objectively characterize wrist structural changes while maintaining clinical relevance.

In this context, recent studies have increasingly emphasized the importance of explainable artificial intelligence (XAI) and imaging biomarkers in medical image analysis. Imaging biomarkers are defined as quantitative image-derived features with explicit morphological or geometric meaning, enabling transparent and clinically interpretable characterization of disease-related structural changes. Compared with purely data-driven deep learning representations, handcrafted features grounded in predefined anatomical or geometric formulations have been highlighted as inherently explainable and more closely aligned with clinical reasoning [18]. Furthermore, evidence-based XAI frameworks stress that interpretability and empirical clinical relevance should be integral components of quantitative imaging methodologies, particularly in radiological decision support applications [19]. These considerations are especially pertinent in musculoskeletal radiography of peripheral joints, where anatomically interpretable measurements are essential for clinical acceptance and longitudinal assessment.

Recent advances in artificial intelligence have enabled automated joint segmentation and quantitative analysis using radiographs. However, most existing studies have focused on finger joints, where anatomical structures are relatively well separated [20,21]. Automated quantitative assessment of the wrist remains limited due to severe bone overlap, complex joint geometry, and projection-related variability inherent to two-dimensional imaging [22]. Under these conditions, achieving robust and reliable quantitative measurements across different imaging positions remains challenging [23]. To address these limitations, we propose a compactness-based morphological analysis framework for auto-

mated assessment of wrist JSN in RA. Rather than relying on local joint space measurements, the proposed approach characterizes the global spatial distribution of carpal bones using centroid-based compactness metrics derived from automated segmentation. The main contributions of this study are as follows:

- **A compactness-based quantitative perspective:** We introduce a global geometric characterization of wrist JSN based on compactness, shifting from conventional local distance measurements to an integrated assessment of carpal bone spatial configuration.
- **Centroid-based compactness metrics:** Using automated carpal bone segmentation with the AP-DPM framework, centroid coordinates of individual carpal bones are extracted to construct a series of compactness-related morphological metrics that are robust to bone overlap and projection variability.
- **Clinical validation against established scoring systems:** The proposed metrics are validated by analyzing their associations with the wrist JSN subscore and total Sharp/van der Heijde score, to evaluate their ability to capture both local wrist structural changes and overall disease severity.

## 2. Materials and Methods

### 2.1. Patients

This study initially enrolled 51 patients with RA (45 females and 6 males). All patients underwent bilateral posteroanterior (PA) conventional radiography examinations at baseline (BL) and follow-up (FU). Patient demographics are detailed in Table 1. During data preprocessing, one patient was excluded based on pre-defined image usability criteria. Specifically, this case exhibited severe structural joint destruction, including extensive bony fusion and advanced bone erosion of the carpal joints, which precluded reliable anatomical differentiation of individual carpal bones on the PA radiographic view. Under such conditions, centroid localization and subsequent compactness-related metric computation were deemed unreliable. Accordingly, cases with severe carpal fusion or late-stage structural deformity were considered an intrinsic limitation of the proposed centroid-based compactness analysis framework and were excluded from quantitative analysis. Ultimately, 50 patients were included in the subsequent compactness-related quantitative metric calculations. For JSNS, all 50 included cases had complete BL and FU data. TSS data were derived from the same cohort of subjects. However, due to missing raw scoring data, one patient lacked BL scores, and three lacked FU scores. The patient inclusion process and data availability are summarized in Figure 1.


**Table 1.** Baseline (BL) clinical and laboratory characteristics of patients with rheumatoid arthritis (RA).

| Characteristic                                                           | Value            |
|--------------------------------------------------------------------------|------------------|
| Number of patients                                                       | 50               |
| Age, years, mean (range)                                                 | 61 (21–82)       |
| Sex, n (female/male)                                                     | 44/6             |
| Symptom duration, years, median (IQR)                                    | 9 (1–32)         |
| Interval between baseline and follow-up radiographs, years, median (IQR) | 1.06 (0.82–1.30) |
| Structural remission status, n (remission/non-remission)                 | 30/20            |
| ESR, mm/h, median (IQR)                                                  | 15 (4–76)        |
| CRP, mg/dL, median (IQR)                                                 | 0.14 (0–6.6)     |
| Swollen joint count, median (IQR)                                        | 0 (0–5)          |
| Tender joint count, median (IQR)                                         | 0 (0–17)         |
| DAS28-CRP, median (IQR)                                                  | 1.84 (0.99–3.85) |
| DAS28-ESR, median (IQR)                                                  | 2.64 (1.13–4.95) |
| Total SHS at baseline, median (IQR)                                      | 67 (0–494)       |
| Total SHS at follow-up, median (IQR)                                     | 67.5 (0–494)     |

Table 1. Cont.


| Characteristic          | Value |
|-------------------------|-------|
| Biologic agents used, n |       |
| None                    | 0     |
| Infliximab              | 17    |
| Adalimumab              | 7     |
| Tocilizumab             | 3     |
| Etanercept              | 19    |
| Abatacept               | 4     |

CRP, C-reactive protein; DAS28, disease activity score in 28 joints; ESR, erythrocyte sedimentation rate; SHS, Sharp–van der Heijde score; IQR, interquartile range.



**Figure 1.** Flowchart of patient inclusion and data availability. 51 patients with rheumatoid arthritis (RA) underwent bilateral posteroanterior wrist radiography at baseline (BL) and follow-up (FU). 1 subject was excluded due to severe carpal bony fusion, leaving 50 patients for quantitative analysis. The wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) was analyzed at the wrist (200 wrists) and subject levels (100 observations), and paired BL–FU TSS data were available for 46 subjects.

Following the description of patient characteristics, the severity of structural joint damage in the study cohort was further characterized. The distribution of joint-level wrist JSN grades across all evaluated joints is shown in Figure 2. While a substantial proportion of joints exhibited no narrowing, moderate to severe JSN grades were also frequently observed, indicating marked heterogeneity in wrist structural involvement.



**Figure 2.** Distribution of joint-level joint space narrowing (JSN) grades (0–4) across all evaluated wrist joints. The histogram summarizes the frequency of individual JSN grades underlying the summed JSN scores used in the correlation analyses.

In addition, TSS demonstrated substantial inter-patient variability, with a mean  $\pm$  standard deviation of  $98.6 \pm 97.8$ .

## 2.2. Radiograph Acquisition

All BL and FU radiographs were obtained using the same conventional radiography system (Toshiba KXO-50G, Tokyo, Japan) in standardized acquisition settings. The imaging was performed without an aluminum filter, with a film–focus distance of 1 m, a tube voltage of 45 kV, a tube current of 250 mA and an exposure time of 0.014 s. Subsequently, all images were digitized and stored in Digital Imaging and Communications in Medicine (DICOM) format, with a pixel resolution of  $0.15 \times 0.15$  mm and a 10-bit grayscale depth.

## 2.3. Wrist Sharp/Van Der Heijde JSN Scores


JSN progression was independently evaluated using SvdH scoring method by two experienced radiologists. Their consensus scores were used as the reference standard for subsequent analyzes in this study. JSN was graded for the wrist joints on a five-point scale as follows: grade 0, normal joint space; grade 1, focal narrowing or questionable change; grade 2, generalized narrowing with more than 50% of the original joint space remaining; grade 3, generalized narrowing with less than 50% of the original joint space remaining or presence of subluxation; and grade 4, bony ankylosis or complete joint luxation [10].

## 2.4. Total Sharp/Van Der Heijde Scores

TSS were assessed using the SvdH scoring method by two experienced rheumatologists from the affiliated hospitals. Their consensus scores were used as the reference standard for subsequent analyses in this study. The TSS was calculated as the sum of JSN and BE scores across all evaluated joints, including the hands and wrists, and the feet when available, thereby reflecting the overall extent of structural joint damage.

## 2.5. Segmentation Network and Quantitative Metrics

As shown in Figure 3, our proposed quantification framework consists of a wrist-specific segmentation network and a set of wrist bone compactness metrics.



**Figure 3.** Overview of AP-DPM segmentation network and segmentation masks based wrist JSN quantitative metric. (A) illustrates metrics based on local centroid relationships, (B) illustrates metrics based on local-global centroids relationships, (C) illustrates metrics based on bone overlap size relationships, (D,E) presents the normalized-BBox versions of (A,B).

### 2.5.1. AP-DPM Segmentation Network

For segmentation, we performed AP-DPM [24], a model tailored to the anatomical characteristics of the wrist bones. AP-DPM employs a dual-path architecture combined with adversarial anatomical priors to improve delineation in severely overlapped regions. The network separately predicts the global bone structure and the overlapping areas, which are subsequently refined through a lightweight merging module, enabling enhanced sensitivity to blurred boundaries and joint spaces. In parallel, a discriminator imposes anatomical plausibility by penalizing implausible bone shapes, thereby enforcing structurally

realistic predictions. A two-stage training strategy further stabilizes learning and balances segmentation accuracy with anatomical consistency. This design yields contour-precise and occlusion-robust segmentation masks, which serve as reliable inputs for downstream compactness metric computation.

The Quantitative results of AP-DPM consistently demonstrate the superiority of AP-DPM over existing state-of-the-art segmentation approaches across multiple evaluation metrics [24]. In terms of overall region accuracy, AP-DPM achieves the highest Dice score of  $0.9784 \pm 0.0052$ , outperforming strong baselines such as nnUNet ( $0.9765 \pm 0.0058$ ) and UNet ( $0.9231 \pm 0.001$ ). More notably, its advantage becomes more pronounced on the boundary-sensitive normalized surface Dice (NSD), where AP-DPM reaches  $0.7830 \pm 0.0788$ , compared with  $0.7612 \pm 0.0572$  for nnUNet and  $0.7087 \pm 0.0784$  for UNet. This improvement underscores its enhanced capability to preserve fine anatomical interfaces under severe projection overlap. Consistently lower volumetric overlap error (VOE:  $0.0142 \pm 0.0096$ ) and relative absolute volume difference (RAVD:  $0.0090 \pm 0.0021$ ) further confirm the geometric stability and robustness of the proposed framework.

### 2.5.2. Carpal Compactness Quantitative Metrics

To quantify the spatial compactness and overall geometric arrangement of the seven carpal bones, we purposed a set of metrics based on pairwise mask overlap and centroid distributions, as shown in Table 2. All carpal masks were first resampled to a uniform resolution of  $600 \times 600$ , and the centroid of each bone was extracted to form a compact set of quantitative metrics. These metrics can be grouped into three categories: overlap size-based metrics derived from pairwise overlap size, local centroid-based metrics capturing inter-centroid relationships, and local-global centroid-based metrics characterizing the spatial distribution of centroids with respect to the global centroid, as detailed below.

**Table 2.** Compactness quantitative metrics for carpal bone.

| Category                                 | Full Name                         | Abbreviation |
|------------------------------------------|-----------------------------------|--------------|
| Overlap size based                       | Overlap size sum                  | OSS          |
|                                          | Overlap size standard deviation   | OSSD         |
| Local centroid based                     | Root mean square radius           | RMSR         |
|                                          | Mean pairwise distance            | MPD          |
| local-global centroid based              | Median radius                     | R50          |
|                                          | 90th-percentile radius            | R90          |
| Local centroid based (normalized)        | Normalized RMS radius             | $RMSR_n$     |
|                                          | Normalized mean pairwise distance | $MPD_n$      |
| local-global centroid based (normalized) | Normalized median radius          | $R50_n$      |
|                                          | Normalized 90th-percentile radius | $R90_n$      |

#### Overlap Size Based Metrics

The degree of anatomical crowding between any two carpals was quantified using the binary overlap of their segmentation masks. For carpal masks  $M_i$  and  $M_j$ , the overlap size was defined as Equation (1)

$$S_{ij} = |\{x \mid M_i(x) = 1 \wedge M_j(x) = 1\}|, \quad (1)$$

where  $|\cdot|$  denotes the number of pixels. Based on all unordered carpal pairs  $(i, j)$  with  $i \neq j$ , the overlap size sum (OSS) and the overlap size standard deviation (OSSD) were computed as Equation (2)

$$OSS = \sum_{i \neq j} S_{ij}, \quad OSSD = \sqrt{\text{Var}(S_{ij})}. \quad (2)$$

#### Local Centroid Based Metrics

To quantify how tightly the carpal centroids cluster around their mean location, we computed the root-mean-square (RMS) radius. Let  $c_k \in \mathbb{R}^2$  denote the centroid of the  $k$ -th carpal bone and  $\mu$  the global centroid defined as Equation (3). Thus, the RMS radius (RMSR) was then defined as Equation (4)

$$\mu = \frac{1}{n} \sum_{k=1}^n c_k. \quad (3)$$

$$RMSR = \sqrt{\frac{1}{n} \sum_{k=1}^n \|c_k - \mu\|^2}. \quad (4)$$

Inter-bone spacing was further characterized using the mean pairwise Euclidean distance (MPD) between all carpal centroids, defined as Equation (5)

$$MPD = \frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \|c_i - c_j\|. \quad (5)$$

#### Local–Global Centroid Based Metrics

To reduce sensitivity to outliers and asymmetric bone arrangements, robust radius measures were derived from the distribution of radial distances to the global centroid. The robust radius was defined as Equation (6).

$$R50 = \text{quantile}(\|c_k - \mu\|, 0.5), \quad R90 = \text{quantile}(\|c_k - \mu\|, 0.9) \quad (6)$$

**Scale Normalization.** To achieve scale invariance across subjects, all centroid-based compactness metrics were normalized by the diagonal length of the bounding box enclosing all centroids. The bounding box diagonal was computed as Equation (7).

$$L_{\text{bbox}} = \sqrt{(x_{\max} - x_{\min})^2 + (y_{\max} - y_{\min})^2}, \quad (7)$$

where  $(x_{\min}, y_{\min})$  and  $(x_{\max}, y_{\max})$  denote the minimum and maximum centroid coordinates. The normalized metrics were then defined as Equation (8).

$$RMSR_n = \frac{RMSR}{L_{\text{bbox}}}, \quad MPD_n = \frac{MPD}{L_{\text{bbox}}}, \quad R50_n = \frac{R50}{L_{\text{bbox}}}, \quad R90_n = \frac{R90}{L_{\text{bbox}}}. \quad (8)$$

### 3. Results

#### 3.1. Comparison Results with JSN

##### 3.1.1. Entire Cohort

Based on the entire cohort of 50 patients, a total of 200 wrist radiographs (50 subjects  $\times$  bilateral wrists  $\times$  BL and FU) with available JSN data were included in the correlation analyses. Overall, several centroid-based compactness metrics demonstrated moderate to relatively strong negative correlations with JSN (Table 3).

Among all evaluated compactness-related metrics, the centroid-based distance metrics derived from centroid spatial distribution showed the strongest correlations with JSN. Specifically, *RMSR* ( $r = -0.45$ ,  $p < 0.0001$ ), *MPD* ( $r = -0.49$ ,  $p < 0.0001$ ) and *R50* ( $r = -0.49$ ,  $p < 0.0001$ ) showing moderate to strong negative correlations with JSN. In contrast, *R90* showed a weaker but still statistically significant negative correlation ( $r = -0.33$ ,  $p < 0.0001$ ).

Among the normalized centroid-based compactness metrics, significant negative correlations with JSN were consistently observed. Both *MPD<sub>n</sub>* and *RMSR<sub>n</sub>* exhibited moderate correlations ( $r = -0.45$  and  $r = -0.31$ , respectively; both  $p < 0.0001$ ), while *R50<sub>n</sub>* showed a statistically significant but slightly weaker correlation ( $r = -0.30$ ,  $p < 0.0001$ ). Notably, *R90<sub>n</sub>* demonstrated a weak positive correlation with JSN ( $r = 0.16$ ,  $p = 0.027$ ).

In contrast, the overlap size-based metrics (*OSS* and *OSSD*) showed no statistically significant correlations with JSN in the overall dataset ( $p > 0.05$ ).

The complete correlation coefficients, confidence intervals, and significance levels are presented in Table 3. The complete results for all evaluated metrics are provided in Supplementary Figure S1.

**Table 3.** Correlations between compactness-related metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) ( $n = 200$ ).

| Metric            | r      | 95% CI           | p       |
|-------------------|--------|------------------|---------|
| OSS               | −0.050 | −0.187 to 0.090  | 0.485   |
| OSSD              | −0.058 | −0.196 to 0.081  | 0.412   |
| RMSR              | −0.453 | −0.556 to −0.335 | <0.0001 |
| MPD               | −0.485 | −0.585 to −0.371 | <0.0001 |
| R50               | −0.486 | −0.586 to −0.373 | <0.0001 |
| R90               | −0.327 | −0.446 to −0.197 | <0.0001 |
| RMSR <sub>n</sub> | −0.306 | −0.427 to −0.175 | <0.0001 |
| MPD <sub>n</sub>  | −0.446 | −0.551 to −0.327 | <0.0001 |
| R50 <sub>n</sub>  | −0.298 | −0.420 to −0.166 | <0.0001 |
| R90 <sub>n</sub>  | 0.157  | 0.018 to 0.289   | 0.0266  |

### 3.1.2. Comparison of Baseline and Follow-Up Results for the Left and Right Hands

In BL radiographs ( $n = 100$ ) (50 subjects  $\times$  BL and FU), among the centroid-based compactness metrics, strong and statistically significant negative correlations with JSN were observed. Specifically, *RMSR* ( $r = -0.47$ ,  $p < 0.0001$ ), *MPD* ( $r = -0.51$ ,  $p < 0.0001$ ), and *R50* ( $r = -0.48$ ,  $p < 0.0001$ ) exhibited high and consistent correlations, all reaching statistical significance. *R90* also demonstrated a statistically significant but comparatively weaker negative correlation ( $r = -0.34$ ,  $p = 0.0006$ ). Among the normalized centroid-based compactness metrics, statistically significant negative correlations with JSN were observed for *RMSR<sub>n</sub>* ( $r = -0.28$ ,  $p = 0.0046$ ), *MPD<sub>n</sub>* ( $r = -0.41$ ,  $p < 0.0001$ ), and *R50<sub>n</sub>* ( $r = -0.28$ ,  $p = 0.0042$ ). In contrast, *R90<sub>n</sub>* did not show a statistically significant correlation with JSN at BL ( $p > 0.05$ ). In addition, the overlap size-based metrics (*OSS* and *OSSD*) did not demonstrate statistically significant correlations with JSN at BL ( $p > 0.05$ ).

In FU radiographs ( $n = 100$ ), similar but slightly strengthened correlation patterns were observed. Among the centroid-based compactness metrics, *RMSR* ( $r = -0.44$ ,  $p < 0.0001$ ), *MPD* ( $r = -0.47$ ,  $p < 0.0001$ ), and *R50* ( $r = -0.49$ ,  $p < 0.0001$ ) exhibited moderate to strong negative correlations with JSN. *R90* demonstrated a weaker but still statistically significant correlation ( $r = -0.32$ ,  $p = 0.0014$ ). Among the normalized centroid-based compactness metrics, *MPD<sub>n</sub>* showed the strongest correlation with JSN ( $r = -0.48$ ,  $p < 0.0001$ ), followed by *RMSR<sub>n</sub>* ( $r = -0.33$ ,  $p = 0.0008$ ) and *R50<sub>n</sub>* ( $r = -0.31$ ,  $p = 0.0014$ ), all reaching statistical significance. In contrast, *R90<sub>n</sub>* did not demonstrate a significant

correlation with JSN at FU ( $p > 0.05$ ). Similarly, the overlap size-based metrics (OSS and OSSD) did not show statistically significant correlations with JSN in FU radiographs ( $p > 0.05$ ).

In the longitudinal change analysis ( $\Delta$ ,  $n = 50$ ), only the overlap size-based metric  $\Delta$ OSS exhibited a statistically significant negative correlation with  $\Delta$ JSN ( $r = -0.30$ ,  $p = 0.0364$ ).  $\Delta$ OSSD showed a non-significant trend toward a negative correlation ( $p > 0.05$ ).


The complete correlation coefficients, confidence intervals, and significance levels are presented in Tables 4 and 5. Representative scatter plots of the strongest correlated metrics demonstrated a clear linear negative correlation (Figure 4). Detailed scatter plots for all evaluated metrics at BL and FU are shown in Supplementary Figures S2 and S3.

**Table 4.** Correlations between the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) and compactness-related metrics at baseline (BL) and follow-up (FU) ( $n = 100$  per time point).

| Metric            | Baseline (BL) |                  |         | Follow-Up (FU) |                  |         |
|-------------------|---------------|------------------|---------|----------------|------------------|---------|
|                   | r             | 95% CI           | p       | r              | 95% CI           | p       |
| OSS               | −0.017        | −0.212 to 0.181  | 0.871   | −0.085         | −0.276 to 0.114  | 0.403   |
| OSSD              | −0.004        | −0.201 to 0.192  | 0.966   | −0.119         | −0.309 to 0.079  | 0.237   |
| RMSR              | −0.468        | −0.609 to −0.300 | <0.0001 | −0.439         | −0.585 to −0.266 | <0.0001 |
| MPD               | −0.505        | −0.638 to −0.343 | <0.0001 | −0.469         | −0.609 to −0.300 | <0.0001 |
| R50               | −0.482        | −0.620 to −0.316 | <0.0001 | −0.491         | −0.627 to −0.326 | <0.0001 |
| R90               | −0.339        | −0.502 to −0.153 | 0.0006  | −0.316         | −0.483 to −0.128 | 0.0014  |
| RMSR <sub>n</sub> | −0.281        | −0.453 to −0.090 | 0.0046  | −0.331         | −0.496 to −0.144 | 0.0008  |
| MPD <sub>n</sub>  | −0.411        | −0.562 to −0.233 | <0.0001 | −0.484         | −0.622 to −0.318 | <0.0001 |
| R50 <sub>n</sub>  | −0.284        | −0.455 to −0.092 | 0.0042  | −0.315         | −0.482 to −0.126 | 0.0014  |
| R90 <sub>n</sub>  | 0.147         | −0.051 to 0.334  | 0.144   | 0.166          | −0.031 to 0.351  | 0.098   |

**Table 5.** Correlations between changes in the wrist joint space narrowing subscore of the Sharp/van der Heijde score ( $\Delta$ JSNS) and changes in compactness-related metrics ( $\Delta$ metrics) ( $n = 50$ ).

| $\Delta$ Metric            | r      | 95% CI           | p      |
|----------------------------|--------|------------------|--------|
| $\Delta$ OSS               | −0.297 | −0.531 to −0.020 | 0.0364 |
| $\Delta$ OSSD              | −0.221 | −0.470 to 0.062  | 0.1239 |
| $\Delta$ RMSR              | 0.059  | −0.223 to 0.332  | 0.6844 |
| $\Delta$ MPD               | 0.077  | −0.206 to 0.348  | 0.5956 |
| $\Delta$ R50               | 0.079  | −0.204 to 0.350  | 0.5864 |
| $\Delta$ R90               | −0.012 | −0.289 to 0.268  | 0.9363 |
| $\Delta$ RMSR <sub>n</sub> | 0.006  | −0.273 to 0.284  | 0.9651 |
| $\Delta$ MPD <sub>n</sub>  | 0.014  | −0.265 to 0.291  | 0.9226 |
| $\Delta$ R50 <sub>n</sub>  | 0.010  | −0.270 to 0.287  | 0.9473 |
| $\Delta$ R90 <sub>n</sub>  | −0.127 | −0.392 to 0.157  | 0.3793 |



**Figure 4.** Correlations between the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) and compactness-related metrics at baseline (BL) and follow-up (FU). Scatter plots show representative metrics: *RMSR* (A,D), *MPD* (B,E), and *R50* (C,F). The upper row (A–C) shows BL results, and the lower row (D–F) shows FU results. \* indicates statistical significance ( $p < 0.05$ ).

### 3.1.3. Comparison of Bilateral Total Scores at Baseline and Follow-Up

Bilateral total JSNS was calculated by summing left and right wrist subscores for each subject at each time point. Thus, 50 subjects were included in the BL and FU analyses, respectively.


In BL radiographs ( $n = 50$ ), among the centroid-based compactness metrics, strong and statistically significant negative correlations with JSN were observed. Specifically, *RMSR* ( $r = -0.55$ ,  $p < 0.0001$ ), *MPD* ( $r = -0.59$ ,  $p < 0.0001$ ), and *R50* ( $r = -0.53$ ,  $p < 0.0001$ ) exhibited high and consistent correlations, all reaching statistical significance. *R90* also demonstrated a statistically significant but comparatively weaker negative correlation ( $r = -0.40$ ,  $p = 0.0043$ ). Among the normalized centroid-based compactness metrics, statistically significant negative correlations with JSN were observed for *RMSR<sub>n</sub>* ( $r = -0.28$ ,  $p = 0.0486$ ) and *MPD<sub>n</sub>* ( $r = -0.43$ ,  $p = 0.0018$ ). In contrast, *R50<sub>n</sub>* ( $p = 0.0577$ ) and *R90<sub>n</sub>* ( $p > 0.05$ ) did not show statistically significant correlations with JSN at BL. In addition, the overlap size-based metrics (*OSS* and *OSSD*) did not demonstrate statistically significant correlations with JSN at BL ( $p > 0.05$ ).

In FU radiographs ( $n = 50$ ), a similar but slightly strengthened correlation pattern was observed. Among the centroid-based compactness metrics, *RMSR* ( $r = -0.52$ ,  $p = 0.0001$ ), *MPD* ( $r = -0.55$ ,  $p < 0.0001$ ), and *R50* ( $r = -0.58$ ,  $p < 0.0001$ ) exhibited moderate to strong negative correlations with JSN. *R90* demonstrated a weaker but still statistically significant correlation ( $r = -0.39$ ,  $p = 0.0047$ ). Among the normalized centroid-based compactness metrics, *MPD<sub>n</sub>* showed the strongest correlation with JSN ( $r = -0.56$ ,  $p < 0.0001$ ), followed by *RMSR<sub>n</sub>* ( $r = -0.41$ ,  $p = 0.0028$ ) and *R50<sub>n</sub>* ( $r = -0.38$ ,  $p = 0.0072$ ), all reaching statistical significance. In contrast, *R90<sub>n</sub>* did not demonstrate a significant correlation with JSN at FU ( $p > 0.05$ ). Similarly, the overlap size-based metrics (*OSS* and *OSSD*) did not show statistically significant correlations with JSN in FU radiographs ( $p > 0.05$ ).

The complete correlation coefficients, confidence intervals, and significance levels are presented in Table 6. Representative scatter plots of the strongest correlated metrics demonstrated a clear linear negative correlation (Figure 5). Detailed scatter plots for all evaluated metrics at BL and FU are shown in Supplementary Figures S4 and S5.

**Table 6.** Correlations between the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) and compactness-related metrics at baseline (BL) and follow-up (FU), based on summed bilateral (left + right) wrist values ( $n = 50$  per time point).

| Metric   | Baseline (BL) |                  |         | Follow-Up (FU) |                  |         |
|----------|---------------|------------------|---------|----------------|------------------|---------|
|          | $r$           | 95% CI           | $p$     | $r$            | 95% CI           | $p$     |
| OSS      | −0.012        | −0.289 to 0.267  | 0.935   | −0.082         | −0.352 to 0.201  | 0.572   |
| OSSD     | −0.006        | −0.284 to 0.273  | 0.969   | −0.118         | −0.383 to 0.166  | 0.417   |
| RMSR     | −0.552        | −0.720 to −0.323 | <0.0001 | −0.519         | −0.696 to −0.281 | 0.0001  |
| MPD      | −0.593        | −0.748 to −0.377 | <0.0001 | −0.546         | −0.716 to −0.316 | <0.0001 |
| R50      | −0.530        | −0.705 to −0.296 | <0.0001 | −0.580         | −0.739 to −0.360 | <0.0001 |
| R90      | −0.397        | −0.608 to −0.134 | 0.0043  | −0.394         | −0.606 to −0.130 | 0.0047  |
| $RMSR_n$ | −0.280        | −0.518 to −0.002 | 0.0486  | −0.415         | −0.621 to −0.154 | 0.0028  |
| $MPD_n$  | −0.431        | −0.633 to −0.173 | 0.0018  | −0.560         | −0.725 to −0.333 | <0.0001 |
| $R50_n$  | −0.270        | −0.510 to 0.009  | 0.0577  | −0.376         | −0.592 to −0.108 | 0.0072  |
| $R90_n$  | 0.207         | −0.076 to 0.459  | 0.149   | 0.170          | −0.114 to 0.428  | 0.237   |



**Figure 5.** Correlations between centroid-based compactness metrics and the bilateral summed wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) at baseline (BL) and follow-up (FU). Scatter plots show representative metrics:  $RMSR$  (A,D),  $MPD$  (B,E), and  $R50$  (C,F). The upper row (A–C) shows BL results, and the lower row (D–F) shows FU results. \* indicates statistical significance ( $p < 0.05$ ).

### 3.1.4. Comparison of Timepoint-Specific Correlations in the Left and Right Hands

In each subgroup analysis,  $n = 50$  represents the number of wrist radiographs included. Specifically, left and right wrists were analyzed separately, and baseline and follow-up timepoints were evaluated independently, without summation across hands or timepoints.

In the left hand radiographs at BL ( $n = 50$ ), centroid-based compactness metrics demonstrated strong and statistically significant negative correlations with JSN. Specifically,  $RMSR$  ( $r = -0.56$ ,  $p < 0.0001$ ),  $MPD$  ( $r = -0.61$ ,  $p < 0.0001$ ), and  $R50$  ( $r = -0.55$ ,  $p < 0.0001$ ) exhibited strong and consistent correlations with JSN.  $R90$  also showed a statistically significant but comparatively weaker negative correlation ( $r = -0.32$ ,  $p = 0.0235$ ). Among the normalized centroid-based compactness metrics,  $MPD_n$  ( $r = -0.53$ ,  $p < 0.0001$ ) exhibited a strong negative correlation with JSN, while  $RMSR_n$  ( $r = -0.37$ ,  $p = 0.0082$ ) and

$R50_n$  ( $r = -0.33$ ,  $p = 0.0187$ ) also showed statistically significant correlations. In contrast,  $R90_n$  ( $r = 0.34$ ,  $p = 0.0151$ ) demonstrated a weak positive correlation with JSN.


At FU in the left hand ( $n = 50$ ), a largely consistent correlation pattern was observed. Strong and statistically significant negative correlations with JSN were again found for  $R50$  ( $r = -0.57$ ,  $p < 0.0001$ ),  $MPD$  ( $r = -0.55$ ,  $p < 0.0001$ ), and  $RMSR$  ( $r = -0.52$ ,  $p < 0.0001$ ).  $R90$  remained significantly correlated ( $r = -0.40$ ,  $p = 0.0043$ ). Among the normalized metrics,  $MPD_n$  continued to demonstrate a strong negative correlation ( $r = -0.54$ ,  $p < 0.0001$ ), followed by  $RMSR_n$  ( $r = -0.41$ ,  $p = 0.0031$ ) and  $R50_n$  ( $r = -0.31$ ,  $p = 0.0299$ ), whereas  $R90_n$  did not reach statistical significance ( $p > 0.05$ ).

In the right hand radiographs at BL ( $n = 50$ ), a similar correlation pattern was observed, although the overall strength of the correlations was slightly weaker than that of the left hand. Among the centroid-based compactness metrics,  $RMSR$  ( $r = -0.39$ ,  $p = 0.0052$ ),  $MPD$  ( $r = -0.41$ ,  $p = 0.0030$ ), and  $R50$  ( $r = -0.40$ ,  $p = 0.0036$ ) showed moderate and statistically significant negative correlations with JSN.  $R90$  also demonstrated a statistically significant negative correlation ( $r = -0.37$ ,  $p = 0.0080$ ). However, among the normalized metrics, none reached statistical significance at BL in the right hand ( $p > 0.05$ ).

At FU in the right hand ( $n = 50$ ), the correlation strength slightly increased compared with BL. Significant negative correlations with JSN were observed for  $R50$  ( $r = -0.44$ ,  $p = 0.0015$ ),  $MPD$  ( $r = -0.40$ ,  $p = 0.0038$ ), and  $RMSR$  ( $r = -0.37$ ,  $p = 0.0080$ ). Among the normalized metrics,  $MPD_n$  ( $r = -0.41$ ,  $p = 0.0028$ ) and  $R50_n$  ( $r = -0.32$ ,  $p = 0.0257$ ) exhibited statistically significant negative correlations with JSN, whereas  $RMSR_n$  and  $R90_n$  did not reach statistical significance ( $p > 0.05$ ).

Across all timepoints and both hands, OSS and OSSD consistently showed no statistically significant correlations with JSN ( $p > 0.05$ ).


The complete correlation coefficients, confidence intervals, and significance levels are presented in Table 7. Representative scatter plots of the strongest correlated metrics demonstrated a clear linear negative correlation (Figures 6 and 7). Corresponding results for all evaluated metrics in the left and right hands are presented in Supplementary Figures S6–S9.



**Figure 6.** Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) for the left hand at baseline (BL) and follow-up (FU). Scatter plots show representative metrics:  $RMSR$  (A,D),  $MPD$  (B,E), and  $R50$  (C,F). The upper row (A–C) shows results for the left hand, and the lower row (D–F) shows for the right hand. \* indicates statistical significance ( $p < 0.05$ ).

**Table 7.** Correlations between the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) and compactness-related metrics at baseline (BL) and follow-up (FU) for the left and right hands (n = 50 per subgroup).

| Metric            | Left Hand BL |                  |         | Left Hand FU |                  |         | Right Hand BL |                  |        | Right Hand FU |                  |        |
|-------------------|--------------|------------------|---------|--------------|------------------|---------|---------------|------------------|--------|---------------|------------------|--------|
|                   | r            | 95% CI           | p       | r            | 95% CI           | p       | r             | 95% CI           | p      | r             | 95% CI           | p      |
| OSS               | 0.029        | −0.252 to 0.305  | 0.8425  | −0.023       | −0.300 to 0.257  | 0.8735  | −0.058        | −0.331 to 0.224  | 0.6903 | −0.141        | −0.404 to 0.143  | 0.3279 |
| OSSD              | 0.047        | −0.234 to 0.321  | 0.7460  | −0.089       | −0.359 to 0.194  | 0.5382  | −0.058        | −0.331 to 0.224  | 0.6879 | −0.156        | −0.416 to 0.128  | 0.2809 |
| RMSR              | −0.557       | −0.723 to −0.330 | <0.0001 | −0.524       | −0.700 to −0.288 | <0.0001 | −0.390        | −0.603 to −0.125 | 0.0052 | −0.371        | −0.589 to −0.104 | 0.0080 |
| MPD               | −0.608       | −0.758 to −0.396 | <0.0001 | −0.554       | −0.721 to −0.326 | <0.0001 | −0.411        | −0.619 to −0.150 | 0.0030 | −0.402        | −0.612 to −0.139 | 0.0038 |
| R50               | −0.554       | −0.721 to −0.326 | <0.0001 | −0.565       | −0.729 to −0.340 | <0.0001 | −0.404        | −0.613 to −0.141 | 0.0036 | −0.437        | −0.638 to −0.181 | 0.0015 |
| R90               | −0.320       | −0.549 to −0.046 | 0.0235  | −0.397       | −0.609 to −0.134 | 0.0043  | −0.371        | −0.589 to −0.103 | 0.0080 | −0.239        | −0.485 to 0.043  | 0.0954 |
| RMSR <sub>n</sub> | −0.370       | −0.588 to −0.102 | 0.0082  | −0.411       | −0.618 to −0.149 | 0.0031  | −0.177        | −0.434 to 0.106  | 0.2180 | −0.238        | −0.484 to 0.044  | 0.0966 |
| MPD <sub>n</sub>  | −0.526       | −0.702 to −0.290 | <0.0001 | −0.544       | −0.714 to −0.313 | <0.0001 | −0.274        | −0.513 to 0.005  | 0.0541 | −0.414        | −0.621 to −0.154 | 0.0028 |
| R50 <sub>n</sub>  | −0.331       | −0.558 to −0.058 | 0.0187  | −0.307       | −0.540 to −0.032 | 0.0299  | −0.212        | −0.463 to 0.070  | 0.1386 | −0.315        | −0.546 to −0.041 | 0.0257 |
| R90 <sub>n</sub>  | 0.342        | 0.070 to 0.566   | 0.0151  | 0.087        | −0.197 to 0.356  | 0.5497  | 0.003         | −0.276 to 0.281  | 0.9857 | 0.245         | −0.036 to 0.490  | 0.0867 |



**Figure 7.** Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) for the Right hand at baseline (BL) and follow-up (FU). Scatter plots show representative metrics: RMSR (A,D), MPD (B,E), and R50 (C,F). The upper row (A–C) shows results for the left hand, and the lower row (D–F) shows for the right hand. \* indicates statistical significance ( $p < 0.05$ ).

### 3.1.5. Comparison of Pooled Correlations in the Left and Right Hands

For hand-specific analyses, the left and right hands were analyzed separately, and BL and FU radiographs were pooled. Accordingly, 50 subjects contributed two time-point observations, yielding 100 left-hand radiographs and 100 right-hand radiographs, respectively.


In the left hand radiographs (n = 100), centroid-based compactness metrics demonstrated moderate to strong and statistically significant negative correlations with JSN. Specifically, RMSR ( $r = -0.54$ ,  $p < 0.0001$ ), MPD ( $r = -0.58$ ,  $p < 0.0001$ ), and R50 ( $r = -0.56$ ,  $p < 0.0001$ ) exhibited strong and consistent correlations with JSN. R90 also showed a statistically significant but comparatively weaker negative correlation ( $r = -0.36$ ,  $p = 0.0002$ ). Among the normalized centroid-based compactness metrics, significant negative correlations with JSN were observed for RMSR<sub>n</sub> ( $r = -0.39$ ,  $p < 0.0001$ ), MPD<sub>n</sub> ( $r = -0.53$ ,  $p < 0.0001$ ), and R50<sub>n</sub> ( $r = -0.32$ ,  $p = 0.0013$ ). In contrast, R90<sub>n</sub> demonstrated a weak positive correlation with JSN ( $r = 0.21$ ,  $p = 0.036$ ).

In the right hand radiographs (n = 100), a similar correlation pattern was observed, although the overall strength of the correlations was consistently weaker than that of the

left hand. Among the centroid-based compactness metrics,  $RMSR$  ( $r = -0.38$ ,  $p = 0.0001$ ),  $MPD$  ( $r = -0.40$ ,  $p < 0.0001$ ), and  $R50$  ( $r = -0.42$ ,  $p < 0.0001$ ) showed moderate and statistically significant negative correlations with JSN.  $R90$  demonstrated a weaker but still statistically significant correlation ( $r = -0.30$ ,  $p = 0.0021$ ). Among the normalized centroid-based compactness metrics,  $RMSR_n$  ( $r = -0.21$ ,  $p = 0.0396$ ),  $MPD_n$  ( $r = -0.34$ ,  $p = 0.0005$ ), and  $R50_n$  ( $r = -0.27$ ,  $p = 0.0074$ ) exhibited statistically significant negative correlations with JSN. In contrast,  $R90_n$  did not demonstrate a statistically significant correlation with JSN ( $p > 0.05$ ).

For the overlap size-based metrics ( $OSS$  and  $OSSD$ ), no statistically significant correlations with JSN were observed in either the left or right hand ( $p > 0.05$ ).

The complete correlation coefficients, confidence intervals, and significance levels are presented in Table 8. Representative scatter plots of the strongest correlated metrics demonstrated a clear linear negative correlation (Figure 8). Corresponding scatter plots for all evaluated metrics in the left and right hands are shown in Supplementary Figures S10 and S11.



**Figure 8.** Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) in the left and right hands at baseline (BL) and follow-up (FU). Scatter plots show representative metrics:  $RMSR$  (A,D),  $MPD$  (B,E), and  $R50$  (C,F). The upper row (A–C) shows results for the left hand, and the lower row (D–F) shows for the right hand. \* indicates statistical significance ( $p < 0.05$ ).

**Table 8.** Correlations between the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) and compactness-related metrics using pooled baseline and follow-up measurements for the left and right hands (BL and FU pooled;  $n = 100$  per hand).

| Metric      | Left Hand (BL and FU Pooled) |                  |         | Right Hand (BL and FU Pooled) |                  |         |
|-------------|------------------------------|------------------|---------|-------------------------------|------------------|---------|
|             | r                            | 95% CI           | p       | r                             | 95% CI           | p       |
| <i>OSS</i>  | 0.004                        | −0.193 to 0.200  | 0.9712  | −0.099                        | −0.290 to 0.100  | 0.3288  |
| <i>OSSD</i> | −0.013                       | −0.209 to 0.184  | 0.8979  | −0.106                        | −0.296 to 0.092  | 0.2935  |
| <i>RMSR</i> | −0.540                       | −0.666 to −0.384 | <0.0001 | −0.379                        | −0.535 to −0.197 | 0.0001  |
| <i>MPD</i>  | −0.580                       | −0.697 to −0.433 | <0.0001 | −0.404                        | −0.557 to −0.226 | <0.0001 |
| <i>R50</i>  | −0.556                       | −0.679 to −0.404 | <0.0001 | −0.420                        | −0.570 to −0.244 | <0.0001 |
| <i>R90</i>  | −0.360                       | −0.520 to −0.176 | 0.0002  | −0.304                        | −0.472 to −0.114 | 0.0021  |

Table 8. Cont.

| Metric   | Left Hand (BL and FU Pooled) |                  |         | Right Hand (BL and FU Pooled) |                   |        |
|----------|------------------------------|------------------|---------|-------------------------------|-------------------|--------|
|          | r                            | 95% CI           | p       | r                             | 95% CI            | p      |
| $RMSR_n$ | −0.389                       | −0.544 to −0.209 | <0.0001 | −0.206                        | −0.387 to −0.010  | 0.0396 |
| $MPD_n$  | −0.533                       | −0.661 to −0.377 | <0.0001 | −0.341                        | −0.504 to −0.155  | 0.0005 |
| $R50_n$  | −0.318                       | −0.484 to −0.129 | 0.0013  | −0.267                        | −0.440 to −0.074  | 0.0074 |
| $R90_n$  | 0.210                        | 0.0137 to 0.3899 | 0.0364  | 0.128                         | −0.0701 to 0.3165 | 0.2042 |

### 3.2. Comparison Results with TSS

#### 3.2.1. Entire Cohort

Based on the final cohort of 50 patients, only patients with paired BL and FU TSS assessments were included in the correlation analyses. As a result, 4 patients (1 missing BL data and 3 missing FU data) were excluded, yielding a total of 46 patients with complete paired TSS data. Overall, several centroid-based compactness metrics demonstrated weak to moderate negative correlations with TSS (Table 9).

Among all evaluated compactness-related metrics, the centroid-based distance metrics derived from centroid spatial distribution exhibited relatively higher correlations with TSS in the overall analysis.  $R50$  consistently showed moderate negative correlations with TSS and reached strong statistical significance ( $r = -0.43$ ,  $p < 0.0001$ ), whereas  $MPD$  demonstrated weaker but statistically significant correlations with TSS ( $r = -0.30$ ,  $p = 0.0042$ ).  $RMSR$  also showed a statistically significant but comparatively weaker correlation with TSS ( $r = -0.27$ ,  $p = 0.010$ ), while  $R90$  did not reach statistical significance ( $p > 0.05$ ).

Among the normalized centroid-based compactness metrics, stronger correlations with TSS were observed. Both  $MPD_n$  and  $R50_n$  demonstrated moderate negative correlations, with correlation coefficients of  $r = -0.48$  and  $r = -0.47$ , respectively (both  $p < 0.0001$ ).  $RMSR_n$  also exhibited a statistically significant correlation with TSS ( $r = -0.40$ ,  $p < 0.0001$ ), whereas  $R90_n$  showed no significant correlation ( $p > 0.05$ ).

In contrast, the overlap size-based metrics ( $OSS$  and  $OSSD$ ) showed no statistically significant correlations with TSS in the overall dataset ( $p > 0.05$ ).

The complete correlation coefficients, confidence intervals, and significance levels are presented in Table 9. The complete results for all evaluated metrics are provided in Supplementary Figure S12.

**Table 9.** Correlations between compactness-related metrics and the total Sharp/van der Heijde score (TSS) ( $n = 92$ ).

| Metric   | r      | 95% CI           | p       |
|----------|--------|------------------|---------|
| OSS      | −0.052 | −0.255 to 0.154  | 0.620   |
| OSSD     | −0.055 | −0.257 to 0.152  | 0.605   |
| RMSR     | −0.267 | −0.448 to −0.066 | 0.010   |
| MPD      | −0.296 | −0.472 to −0.097 | 0.0042  |
| R50      | −0.425 | −0.579 to −0.241 | <0.0001 |
| R90      | −0.148 | −0.342 to 0.059  | 0.159   |
| $RMSR_n$ | −0.403 | −0.561 to −0.216 | <0.0001 |
| $MPD_n$  | −0.483 | −0.626 to −0.309 | <0.0001 |
| $R50_n$  | −0.465 | −0.611 to −0.287 | <0.0001 |
| $R90_n$  | 0.030  | −0.176 to 0.233  | 0.779   |

### 3.2.2. Comparison of Baseline and Follow-Up

In BL radiographs ( $n = 46$ ), among the centroid-based compactness metrics,  $R50$  showed a statistically significant negative correlation with TSS ( $r = -0.41$ ,  $p = 0.0044$ ), while  $MPD$  also demonstrated a statistically significant negative correlation ( $r = -0.31$ ,  $p = 0.0386$ ). In contrast, both  $RMSR$  and  $R90$  did not show statistically significant correlations with TSS at BL ( $p > 0.05$ ). Among the normalized centroid-based compactness metrics,  $MPD_n$  ( $r = -0.38$ ,  $p = 0.0084$ ) and  $R50_n$  ( $r = -0.41$ ,  $p = 0.0051$ ) demonstrated statistically significant negative correlations with TSS, whereas the remaining normalized metrics ( $RMSR_n$  and  $R90_n$ ), did not reach statistical significance ( $p > 0.05$ ). In addition, the overlap size-based metrics ( $OSS$  and  $OSSD$ ) did not show statistically significant correlations with TSS at BL ( $p > 0.05$ ).

In FU radiographs (46), stronger and more consistent correlations with TSS were observed. Among the centroid-based compactness metrics,  $R50$  showed a statistically significant negative correlation with TSS ( $r = -0.44$ ,  $p = 0.0023$ ), whereas  $MPD$  demonstrated a borderline but non-significant association ( $r = -0.29$ ,  $p = 0.051$ ). Both  $RMSR$  and  $R90$  did not reach statistical significance ( $p > 0.05$ ). Among the normalized centroid-based compactness metrics, strong negative correlations with TSS were observed.  $RMSR_n$  ( $r = -0.54$ ,  $p = 0.0001$ ),  $MPD_n$  ( $r = -0.59$ ,  $p < 0.0001$ ), and  $R50_n$  ( $r = -0.53$ ,  $p = 0.0002$ ) all demonstrated statistically significant negative correlations with TSS. In contrast,  $R90_n$  did not show a significant correlation ( $p > 0.05$ ). Similarly, the overlap size-based metrics ( $OSS$  and  $OSSD$ ) did not demonstrate statistically significant correlations with TSS at FU ( $p > 0.05$ ).

In the longitudinal change analysis ( $\Delta$ ), only a limited number of centroid-based compactness metrics, including  $\Delta RMSR$ ,  $\Delta MPD$ , and  $\Delta R90$ , exhibited statistically significant negative correlations with  $\Delta TSS$  ( $p < 0.05$ ), whereas the normalized centroid-based compactness metrics and overlap size-based metrics did not demonstrate significant correlations ( $p > 0.05$ ).


The complete correlation coefficients, confidence intervals, and significance levels are presented in Tables 10 and 11. Representative scatter plots of the strongest correlated metrics demonstrated a clear linear negative correlation (Figure 9). Complete correlation results for all metrics with TSS at BL and FU are provided in Supplementary Figures S13 and S14.

**Table 10.** Correlations between compactness-related metrics and the total Sharp/van der Heijde score (TSS) at baseline (BL,  $n = 46$ ) and follow-up (FU,  $n = 46$ ).

| Metric   | Baseline (BL, $n = 46$ ) |                  |       | Follow-Up (FU, $n = 46$ ) |                  |         |
|----------|--------------------------|------------------|-------|---------------------------|------------------|---------|
|          | $r$                      | 95% CI           | $p$   | $r$                       | 95% CI           | $p$     |
| $OSS$    | −0.029                   | −0.317 to 0.263  | 0.848 | −0.076                    | −0.359 to 0.219  | 0.614   |
| $OSSD$   | −0.010                   | −0.300 to 0.281  | 0.947 | −0.103                    | −0.382 to 0.193  | 0.494   |
| $RMSR$   | −0.265                   | −0.516 to 0.027  | 0.075 | −0.271                    | −0.520 to 0.021  | 0.069   |
| $MPD$    | −0.306                   | −0.548 to −0.017 | 0.039 | −0.290                    | −0.535 to 0.001  | 0.051   |
| $R50$    | −0.413                   | −0.628 to −0.139 | 0.004 | −0.438                    | −0.646 to −0.169 | 0.002   |
| $R90$    | −0.137                   | −0.411 to 0.159  | 0.363 | −0.158                    | −0.429 to 0.139  | 0.294   |
| $RMSR_n$ | −0.275                   | −0.523 to 0.017  | 0.065 | −0.540                    | −0.718 to −0.296 | 0.0001  |
| $MPD_n$  | −0.384                   | −0.607 to −0.106 | 0.008 | −0.588                    | −0.750 to −0.359 | <0.0001 |
| $R50_n$  | −0.406                   | −0.623 to −0.131 | 0.005 | −0.528                    | −0.710 to −0.281 | 0.0002  |
| $R90_n$  | 0.088                    | −0.207 to 0.369  | 0.559 | −0.026                    | −0.314 to 0.266  | 0.863   |

**Table 11.** Correlations between changes in compactness-related metrics ( $\Delta$ Metric) and changes in the total Sharp/van der Heijde score ( $\Delta$ TSS) ( $n = 46$ ).

| $\Delta$ Metric | $r$    | 95% CI           | $p$   |
|-----------------|--------|------------------|-------|
| $\Delta OSS$    | 0.056  | −0.238 to 0.340  | 0.713 |
| $\Delta OSSD$   | −0.170 | −0.439 to 0.127  | 0.259 |
| $\Delta RMSR$   | −0.367 | −0.594 to −0.086 | 0.012 |
| $\Delta MPD$    | −0.352 | −0.583 to −0.069 | 0.016 |
| $\Delta R50$    | −0.168 | −0.437 to 0.129  | 0.265 |
| $\Delta R90$    | −0.335 | −0.570 to −0.050 | 0.023 |
| $\Delta RMSR_n$ | −0.048 | −0.334 to 0.246  | 0.751 |
| $\Delta MPD_n$  | −0.068 | −0.351 to 0.227  | 0.653 |
| $\Delta R50_n$  | 0.061  | −0.224 to 0.345  | 0.689 |
| $\Delta R90_n$  | 0.147  | −0.150 to 0.419  | 0.331 |

**Figure 9.** Scatter plots showing correlations between representative compactness-related metrics and the total Sharp/van der Heijde score (TSS) at baseline (BL) and follow-up (FU). Scatter plots show representative metrics: *RMSR* (A,D), *MPD* (B,E), and *R50* (C,F). The upper row (A–C) shows BL results, and the lower row (D–F) shows FU results. \* indicates statistical significance ( $p < 0.05$ ).

## 4. Discussion

### 4.1. Overall Performance of Centroid-Based Compactness Metrics for Wrist JSN Assessment

This study proposes a novel quantitative framework for objectively characterizing wrist joint space narrowing (JSN) in patients with rheumatoid arthritis (RA). Chronic synovitis in RA leads to progressive cartilage loss and gradual joint space narrowing, which underlie the radiographic assessment of structural damage [25,26]. However, most previous studies have relied on traditional semi-quantitative scoring systems, such as the Sharp/van der Heijde (SvdH) method, which depend on manual interpretation, are inherently subjective, and are relatively insensitive to subtle structural changes [27].

Existing automated approaches have mainly focused on local joint boundary detection or overall disease classification, which limits their ability to capture the global spatial contraction of the wrist during disease progression [21,28]. This limitation is particularly relevant for the wrist, where anatomical complexity and overlapping carpal bones reduce the robustness of local joint space measurements on two-dimensional radiographs [29]. To address these challenges, the present study introduces centroid-based compactness

metrics derived from the spatial distribution of carpal bone centroids, aiming to characterize the overall contraction trend of the wrist as JSN progresses.

Our results demonstrate that several distance-based centroid metrics showed statistically significant negative correlations with manual JSN scores across the overall cohort and at baseline (BL) and follow-up (FU). Among all evaluated metrics, mean pairwise distance (MPD), root mean square radius (RMSR), and median radius (R50) consistently exhibited the strongest correlations with JSN. These findings indicate that increasing JSN is associated with greater global spatial contraction of carpal bone centroids, supporting the biological plausibility and clinical relevance of the proposed compactness framework.

#### *4.2. Sensitivity to Early and Progressive Structural Changes at Baseline and Follow-Up*

At baseline, several centroid-based compactness metrics demonstrated significant negative correlations with JSN, suggesting that even subtle early joint space narrowing can influence the global spatial configuration of the carpal bones. This finding indicates that compactness metrics may be sensitive to early structural alterations that are difficult to quantify using conventional radiographic measurements.

At follow-up, the same negative correlation trends were maintained, although longitudinal changes in compactness metrics showed weaker associations with changes in JSN. This observation is likely attributable to the relatively short follow-up interval and the long-standing disease duration in the studied cohort, where substantial structural progression within one year is not always expected. Importantly, the consistency of correlation strength between BL and FU suggests that centroid-based compactness reflects a stable, reproducible imaging feature related to structural narrowing rather than measurement noise.

#### *4.3. Bilateral Robustness and Symmetry Considerations*

Bilateral analyses showed that centroid-based compactness metrics were consistently negatively correlated with JSN in both wrists. Although slight differences in correlation strength were observed between the left and right hands, these variations may reflect differences in functional loading, positioning during radiographic acquisition, or image quality [23,30]. Overall, the consistent bilateral correlation patterns indicate that the proposed metrics robustly capture JSN-related structural changes across both wrists, supporting their applicability in routine bilateral assessment.

#### *4.4. Comparison with TSS and Implications for Assessment Scope*

When compared with the total Sharp/van der Heijde score (TSS), centroid-based compactness metrics generally showed weaker correlations than those observed with wrist-specific JSN scores. This difference is expected, as TSS reflects cumulative, multi-joint structural damage represented as JSN and BE in the bilateral hand and foot joints, whereas the proposed compactness metrics focus on local wrist geometry [31]. Structural changes at the wrist do not necessarily progress synchronously with systemic joint damage [32]. Therefore, the relatively weaker association with TSS highlights the complementary nature of the proposed metrics: they are better suited for capturing local wrist-specific structural alterations related to JSN rather than serving as a surrogate for whole-body disease burden.

#### *4.5. Advantages over Traditional Scoring Systems and Existing Automated Methods*

Traditional radiographic scoring systems remain the clinical reference standard but are time-consuming and limited in their ability to detect subtle or continuous structural changes [33]. Recent deep learning-based methods have improved objectivity by automating joint space width estimation or texture-based assessment [12]; however, these approaches primarily rely on localized joint measurements and remain vulnerable to projection variability and bone overlap in the wrist [34].

In contrast, the proposed compactness framework quantifies the global spatial distribution of multiple carpal bones using centroid-based metrics. By focusing on overall geometric contraction rather than individual joint boundaries, this approach is inherently less sensitive to local segmentation errors and projection-related variability. As demonstrated in this study, centroid-based distance metrics showed stable performance across BL, FU, and bilateral analyses, with correlations that were generally comparable to or superior to those of previously reported computer-aided diagnosis metrics based on local joint characteristics [35].

#### 4.6. Biological Interpretation, Limitations, and Future Perspectives

RA is characterized by heterogeneous, sometimes asymmetric, structural progression, particularly in the early stages. Localized and diffuse JSN may coexist, and early changes can be difficult to detect using conventional joint-level scoring alone. By integrating centroid information from multiple carpal bones, the proposed compactness metrics provide a global representation of wrist structural remodeling, capturing an aspect of disease progression that has received limited attention in prior radiographic studies.

Several limitations should be acknowledged. First, the sample size and relatively short follow-up period limit the assessment of sensitivity to short-term progression. Larger cohorts with longer longitudinal follow-up are required to fully evaluate the responsiveness of compactness metrics to disease progression. Second, in advanced RA with severe carpal fusion or marked deformity, reliable segmentation and centroid extraction become infeasible. In such cases, quantitative geometric assessment is subject to ceiling effects and has limited clinical utility. Accordingly, the proposed framework is most suitable for early to moderate RA, where carpal anatomy remains sufficiently distinguishable and quantitative monitoring of structural change is clinically meaningful.

Despite these limitations, this study demonstrates that automated centroid-based compactness metrics can reliably capture global structural changes in the wrist associated with JSN. When used in conjunction with established semi-quantitative scoring systems, this approach provides a complementary, objective tool for standardized radiographic assessment and may facilitate future applications in clinical follow-up and clinical trials.

## 5. Conclusions

This study proposes an automated compactness-related quantitative assessment framework based on AP-DPM for the objective assessment of wrist JSN in patients with RA. The results demonstrate that compactness-related quantitative metrics derived from the spatial distribution of carpal bone centroids can stably capture changes in wrist compactness correlated with JSN and exhibit good measurement stability. In clinical validation, this method shows strong consistency with the JSNS, and comparative analysis with the TSS further clarifies its assessment focus and scope of application. Overall, this automated compactness-related quantitative assessment framework provides a feasible and objective approach for quantifying joint space narrowing-related structural changes in the wrist of patients with RA and lays the foundation for future standardized automated imaging assessments.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/jimaging12020087/s1>, Figure S1: Correlations between centroid-based compactness metrics and joint space narrowing (JSN) in the entire cohort. Figure S2: Correlations between the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) and compactness-related metrics at baseline (BL) for the combined left and right hands. Figure S3: Correlations between the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) and compactness-related metrics at follow-up (FU) for the combined left and right hands. Figure S4: Corre-

lations between bilateral summed wrist joint space narrowing subscores of the Sharp/van der Heijde score (JSNS) and all evaluated compactness-related metrics at baseline (BL). Figure S5: Correlations between bilateral summed wrist joint space narrowing subscores of the Sharp/van der Heijde score (JSNS) and all evaluated compactness-related metrics at follow-up (FU). Figure S6: Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) in the left hand at baseline (BL). Figure S7: Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) in the left hand at follow-up (FU). Figure S8: Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) in the right hand at baseline (BL). Figure S9: Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) in the right hand at follow-up (FU). Figure S10: Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) in the left hand at baseline (BL) and follow-up (FU). Figure S11: Correlations between centroid-based compactness metrics and the wrist joint space narrowing subscore of the Sharp/van der Heijde score (JSNS) in the right hand at baseline (BL) and follow-up (FU). Figure S12: Correlations between centroid-based compactness metrics and the total Sharp/van der Heijde score (TSS) in the entire cohort. Figure S13: Correlations between centroid-based compactness metrics and the total Sharp/van der Heijde score (TSS) at baseline (BL). Figure S14: Correlations between centroid-based compactness metrics and the total Sharp/van der Heijde score (TSS) at follow-up (FU).

**Author Contributions:** Conceptualization, T.K. and H.W.; methodology, H.W., J.P. and J.Z.; software, H.W.; validation, J.Z. and J.P.; formal analysis, J.Z.; investigation, J.Z. and H.W.; resources, H.K., M.M., T.W. and T.K.; data curation, J.Z.; writing—original draft preparation, J.Z.; writing—review and editing, J.P., H.W. and T.K.; visualization, J.Z.; supervision, T.K. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki, and approved by the institutional ethics committee of the Faculty of Health Sciences, Hokkaido University (protocol code 24-104 and approval date 4 March 2025).

**Informed Consent Statement:** Patient consent was waived because the data were collected in an earlier period, and it was difficult to contact individual patients; instead, the study information was disclosed publicly on the institutional website in accordance with the ethics committee's requirements.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author, as the legal procedures regarding permission for public data sharing at the institution where the data were generated have not yet been completed; the data will be made publicly available once these procedures are finalized.

**Acknowledgments:** The authors would like to thank ChatGPT-5.2 (OpenAI) for assistance with language polishing and improvement of academic writing. The authors also gratefully acknowledge the participating institutions and the clinical staff involved in data acquisition and clinical evaluation for their support and cooperation in this study. All individuals who made substantial intellectual contributions are appropriately included in the author list.

**Conflicts of Interest:** The authors declare no conflicts of interests.

## References

1. Sweeney, S.E.; Firestein, G.S. Rheumatoid arthritis: Regulation of synovial inflammation. *Int. J. Biochem. Cell Biol.* **2004**, *36*, 372–378.
2. Sharif, K.; Sharif, A.; Jumah, F.; Oskouian, R.; Tubbs, R.S. Rheumatoid arthritis in review: Clinical, anatomical, cellular and molecular points of view. *Clin. Anat.* **2018**, *31*, 216–223.
3. Trieb, K. Treatment of the wrist in rheumatoid arthritis. *J. Hand Surg.* **2008**, *33*, 113–123.
4. Smolen, J.S.; Van der Heijde, D.; Aletaha, D.; Xu, S.; Han, J.; Baker, D.; St Clair, E. Progression of radiographic joint damage in rheumatoid arthritis: independence of erosions and joint space narrowing. *Ann. Rheum. Dis.* **2009**, *68*, 1535–1540.

5. Finckh, A.; Gilbert, B.; Hodgkinson, B.; Bae, S.C.; Thomas, R.; Deane, K.D.; Alpizar-Rodriguez, D.; Lauper, K. Global epidemiology of rheumatoid arthritis. *Nat. Rev. Rheumatol.* **2022**, *18*, 591–602.
6. Almutairi, K.B.; Nossent, J.C.; Preen, D.B.; Keen, H.I.; Inderjeeth, C.A. The prevalence of rheumatoid arthritis: A systematic review of population-based studies. *J. Rheumatol.* **2021**, *48*, 669–676.
7. Leak, R.S.; Rayan, G.M.; Arthur, R.E. Longitudinal radiographic analysis of rheumatoid arthritis in the hand and wrist. *J. Hand Surg.* **2003**, *28*, 427–434.
8. Van der Heijde, D. Joint erosions and patients with early rheumatoid arthritis. *Rheumatology* **1995**, *34*, 74–78.
9. Colebatch, A.N.; Edwards, C.J.; Østergaard, M.; van der Heijde, D.; Balint, P.V.; d’Agostino, M.A.; Forslind, K.; Grassi, W.; Haavardsholm, E.A.; Haugeberg, G.; et al. EULAR recommendations for the use of imaging of the joints in the clinical management of rheumatoid arthritis. *Ann. Rheum. Dis.* **2013**, *72*, 804–814.
10. Van der Heijde, D.M. Plain X-rays in rheumatoid arthritis: Overview of scoring methods, their reliability and applicability. *Bailliere’s Clin. Rheumatol.* **1996**, *10*, 435–453.
11. James, M.F.; Heald, G.; Shorter, J.H.; Turner, R.A. Joint space measurement in hand radiographs using computerized image analysis. *Arthritis Rheum. Off. J. Am. Coll. Rheumatol.* **1995**, *38*, 891–901.
12. Huo, Y.; Vincken, K.L.; Van Der Heijde, D.; De Hair, M.J.; Lafeber, F.P.; Viergever, M.A. Automatic quantification of radiographic wrist joint space width of patients with rheumatoid arthritis. *IEEE Trans. Biomed. Eng.* **2017**, *64*, 2695–2703.
13. Bruynesteyn, K.; van der Heijde, D.; Boers, M.; van der Linden, S.; Lassere, M.; van der Vleuten, C. The Sharp/van der Heijde method out-performed the Larsen/Scott method on the individual patient level in assessing radiographs in early rheumatoid arthritis. *J. Clin. Epidemiol.* **2004**, *57*, 502–512.
14. Sharp, J.T.; Lidsky, M.D.; Collins, L.C.; Moreland, J. Methods of scoring the progression of radiologic changes in rheumatoid arthritis. Correlation of radiologic, clinical and laboratory abnormalities. *Arthritis Rheum. Off. J. Am. Coll. Rheumatol.* **1971**, *14*, 706–720.
15. Van der Heijde, D. How to read radiographs according to the Sharp/van der Heijde method. *J. Rheumatol.* **2000**, *27*, 261–263.
16. Wen, J.; Liu, J.; Xin, L.; Wan, L.; Jiang, H.; Sun, Y.; Sun, Y.; Wang, X.; Wang, J. Effective factors on Sharp Score in patients with rheumatoid arthritis: A retrospective study. *BMC Musculoskelet. Disord.* **2021**, *22*, 865.
17. Genant, H.K. Methods of assessing radiographic change in rheumatoid arthritis. *Am. J. Med.* **1983**, *75*, 35–47.
18. Rundo, Leonardo and Militello, Carmelo. Image biomarkers and explainable AI: Handcrafted features versus deep learned features. *Eur. Radiol. Exp.* **2024**, *8*, 130.
19. Famigliani, Lorenzo and Campagner, Andrea and Barandas, Marilia and La Maida, G and Gallazzi, Enrico and Cabitza, Federico and others. Evidence-based XAI: An empirical approach to design more effective and explainable decision support systems. *Comput. Biol. Med.* **2024**, *170*, 108042.
20. Zieliński, B. Hand radiograph analysis and joint space location improvement for image interpretation. *Schedae Informaticae* **2009**, *1718*, 45–61.
21. Huo, Y.; Vincken, K.L.; van der Heijde, D.; De Hair, M.J.; Lafeber, F.P.; Viergever, M.A. Automatic quantification of radiographic finger joint space width of patients with early rheumatoid arthritis. *IEEE Trans. Biomed. Eng.* **2015**, *63*, 2177–2186.
22. Kang, B.K.; Han, Y.; Oh, J.; Lim, J.; Ryu, J.; Yoon, M.S.; Lee, J.; Ryu, S. Automatic segmentation for favourable delineation of ten wrist bones on wrist radiographs using Convolutional Neural Network. *J. Pers. Med.* **2022**, *12*, 776.
23. Ichikawa, S.; Kamishima, T.; Sutherland, K.; Fukae, J.; Katayama, K.; Aoki, Y.; Okubo, T.; Okino, T.; Kaneda, T.; Takagi, S.; et al. Computer-based radiographic quantification of joint space narrowing progression using sequential hand radiographs: Validation study in rheumatoid arthritis patients from multiple institutions. *J. Digit. Imaging* **2017**, *30*, 648–656.
24. Yang, S.; Wang, H.; Ikebe, M.; Kamishima, T.; Ou, Y.; Okutomi, M. AP-DPM: A Dual-Path Merging Network via Adversarial Anatomical Prior Guidance for Wrist Bone Segmentation. In Proceedings of the 2025 IEEE International Conference on Bioinformatics and Biomedicine (BIBM), Wuhan, China, 15–18 December 2025.
25. Ostrowska, M.; Maśliński, W.; Prochorec-Sobieszek, M.; Nieciecki, M.; Sudoł-Szopińska, I. Cartilage and bone damage in rheumatoid arthritis. *Rheumatologia* **2018**, *56*, 111–120.
26. Kirwan, J.; Byron, M.; Watt, I. The relationship between soft tissue swelling, joint space narrowing and erosive damage in hand X-rays of patients with rheumatoid arthritis. *Rheumatology* **2001**, *40*, 297–301.
27. Swinkels, H.L.; Laan, R.F.; van’t Hof, M.A.; van der Heijde, D.M.; de Vries, N.; van Riel, P.L. Modified Sharp method: Factors influencing reproducibility and variability. In *Seminars in Arthritis and Rheumatism*; Elsevier: Amsterdam, The Netherlands, 2001; Volume 31, pp. 176–190.
28. Maziarz, K.; Krason, A.; Wojna, Z. Deep learning for rheumatoid arthritis: Joint detection and damage scoring in x-rays. *arXiv* **2021**, arXiv:2104.13915.
29. Bhat, A.K.; Kumar, B.; Acharya, A. Radiographic imaging of the wrist. *Indian J. Plast. Surg.* **2011**, *44*, 186–196.
30. Schenk, O.; Huo, Y.; Vincken, K.L.; van de Laar, M.A.; Kuper, I.H.; Slump, K.C.; Lafeber, F.P.; Bernelot Moens, H.J. Validation of automatic joint space width measurements in hand radiographs in rheumatoid arthritis. *J. Med Imaging* **2016**, *3*, 044502.

31. Van der Heijde, D.; Van Leeuwen, M.; Van Riel, P.; Van de Putte, L. Radiographic progression on radiographs of hands and feet during the first 3 years of rheumatoid arthritis measured according to Sharp's method (van der Heijde modification). *J. Rheumatol.* **1995**, *22*, 1792–1796.
32. Janta, I.; Stanciu, D.; Hinojosa, M.; Nieto-González, J.C.; Valor, L.; Bello, N.; Serrano, B.; Mata-Martínez, C.; Martínez-Barrio, J.; Ovalles-Bonilla, J.G.; et al. Structural damage in rheumatoid arthritis: Comparison between tendon damage evaluated by ultrasound and radiographic damage. *Rheumatology* **2016**, *55*, 1042–1046.
33. Yazici, Y.; Sokka, T.; Pincus, T. Radiographic measures to assess patients with rheumatoid arthritis: advantages and limitations. *Rheum. Dis. Clin.* **2009**, *35*, 723–729.
34. Pietka, E.; Kaabi, L.; Kuo, M.; Huang, H. Feature extraction in carpal-bone analysis. *IEEE Trans. Med Imaging* **1993**, *12*, 44–49.
35. Gundry, M.; Knapp, K.; Meertens, R.; Meakin, J. Computer-aided detection in musculoskeletal projection radiography: A systematic review. *Radiography* **2018**, *24*, 165–174.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.