



Supplementary Material

Comparison of rumen and manure microbiomes and implications for the inoculation of anaerobic digesters

Emine Gozde Ozbayram^{1,3}, Orhan Ince¹, Bahar Ince², Hauke Harms³, Sabine Kleinsteuber^{3,*}

¹ Department of Environmental Engineering, Faculty of Civil Engineering, Istanbul Technical University, Maslak, 34469 Istanbul, Turkey; gozbayram@itu.edu.tr, inceor@itu.edu.tr

² Institute of Environmental Sciences, Boğaziçi University, Bebek, 34342 Istanbul, Turkey; bahar.ince@boun.edu.tr

³ Department of Environmental Microbiology, Helmholtz Centre for Environmental Research - UFZ, 04318 Leipzig, Germany; hauke.harms@ufz.de, sabine.kleinsteuber@ufz.de

* Correspondence: sabine.kleinsteuber@ufz.de; Tel.: +49-341-235-1325

Supplementary Figures:

Figure S1. Krona charts illustrating the bacterial community composition of (a) the rumen samples and (b) the manure samples from three individuals on phylum, class, order and family levels.

Figure S2. Krona charts illustrating the methanogenic community composition of (a) the rumen samples and (b) the manure samples from three individuals on phylum, class, order and family levels.




Figure S1. Krona charts illustrating the bacterial community composition of (a) the rumen samples and (b) the manure samples from three individuals on phylum, class, order and family levels.




Figure S2. Krona charts illustrating the methanogenic community composition of (a) the rumen samples and (b) the manure samples from three individuals on phylum, class, order and family levels.