



Article

Genotypic and Phenotypic Resistance Profiles of Invasive *Acinetobacter baumannii* Isolates at Bulgarian University Hospital

Teodora Marinova-Bulgaranova ^{1,*} , Preslava Hristova ¹, Polyana Marinovska ¹, Nikolay Balgaranov ², Vladislav Nankov ³ , Slavi Peykov ^{4,5,6} and Hristina Hitkova ^{1,*}

¹ Department of Microbiology and Virology, Medical University of Pleven, 5800 Pleven, Bulgaria; preslava.hristova@mu-pleven.bg (P.H.); polyana.marinovska@mu-pleven.bg (P.M.)

² Department of Pediatric Diseases, Medical University of Pleven, 5800 Pleven, Bulgaria; nikolai.balgaranov@mu-pleven.bg

³ Department of Anatomy, Histology, Cytology and Biology, Medical University Pleven, 5800 Pleven, Bulgaria; vladislav.nankov@mu-pleven.bg

⁴ Department of Medical Microbiology, Faculty of Medicine, Medical University of Sofia, 2 Zdrave Str., 1431 Sofia, Bulgaria; spejgov@biofac.uni-sofia.bg

⁵ Department of Genetics, Faculty of Biology, University of Sofia ‘St. Kliment Ohridski’, 8 Dragan Tzankov Blvd., 1164 Sofia, Bulgaria

⁶ BioInfoTech Laboratory, Sofia Tech Park, 111 Tsarigradsko Shose Blvd., 1784 Sofia, Bulgaria

* Correspondence: teodora.marinova@mu-pleven.bg (T.M.-B.); hristinka.hitkova@mu-pleven.bg (H.H.); Tel.: +359-899-11-60-21 (H.H.)

Abstract

Acinetobacter baumannii is a formidable nosocomial pathogen responsible for severe hospital-acquired infections. This study aimed to investigate the relationship between genotypic and phenotypic resistance profiles in 36 non-duplicate invasive *A. baumannii* isolates collected at a tertiary care hospital in Bulgaria over a six-year period (2019–2024). Identification was performed by MALDI-TOF MS and PCR-based detection of *bla*_{OXA-51-like} and *gyrB*. Antimicrobial susceptibility testing to ten agents was performed by gradient diffusion and broth microdilution, and selected resistance genes were detected by conventional PCR. Whole-genome sequencing of 17 representative isolates was conducted to confirm the resistance genotype–phenotype associations and perform MLST. Carbapenem resistance was detected in 34/36 isolates (94.4%), mediated by *bla*_{OXA-23-like} (29/34) or *bla*_{OXA-72} (5/34). The *armA* gene associated with high-level resistance to all three aminoglycosides was identified exclusively in *bla*_{OXA-23-like}-positive isolates (18/29). In *armA*-negative isolates, gentamicin and tobramycin MICs decreased markedly, whereas amikacin MICs remained elevated. In addition, minocycline MICs ≥ 8 mg/L were observed in the presence of *armA*. Seven distinct genotypic–phenotypic profiles were defined based on these gene–MIC associations. Resistome and phylogenetic analyses confirmed these relationships, with MLST assigning the isolates to established clonal lineages ST2 and ST636 (International Clone II). These findings support the use of targeted genes, such as *armA*, combined with phenotypic testing to guide empirical therapy and strengthen local surveillance of high-risk *A. baumannii* clones in hospital settings.

Keywords: *A. baumannii*; minimal inhibitory concentration; antimicrobial resistance genes; resistance genotype; resistance phenotype



Academic Editor: Haihua Liang

Received: 6 June 2026

Revised: 18 July 2026

Accepted: 20 July 2026

Published: 24 July 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and

conditions of the [Creative Commons](#)

[Attribution \(CC BY\)](#) license.

1. Introduction

Acinetobacter baumannii has emerged over the past decade as a major nosocomial pathogen owing to its capacity to withstand last-line antimicrobials and disseminate clonally [1,2]. As a member of the ESKAPE group (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *A. baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* spp.), it ranks among the most critical threats to global public health [3]. Its resistance to desiccation enables prolonged survival in the hospital environment, thereby facilitating endemic persistence [2].

The European Antimicrobial Resistance Surveillance Network (EARS-Net) data for 2024 indicate that carbapenem resistance in *A. baumannii* exceeds 90% in several Southeast European EU Member States, particularly in Balkan countries [4]. Moreover, the Balkan region showed the highest rates of combined resistance: Romania (89.5%), Croatia (89.0%), and Greece (87.9%) [4]. In Bulgaria, the proportion of invasive carbapenem-resistant *A. baumannii* (CRAB) isolates increased from 80% in 2022 to 90.5% in 2024, following the COVID-19 pandemic period [4]. During the same period, combined resistance rose from 75.0% to 86.7%, markedly exceeding the EU/EEA population-weighted mean of 30.2% in 2024 [4].

Antibiotic resistance in Gram-negative bacteria is mediated by three principal strategies: enzymatic inactivation or complete hydrolysis of the antimicrobial molecule, modification of the target, and reduced membrane permeability resulting from the loss or alteration of porins and/or the activity of efflux pumps [5,6]. These mechanisms operate in clinical *A. baumannii* isolates and are driven by well-defined genetic determinants or mutations. In this regard, it is important to highlight the remarkable genomic plasticity of this species, which facilitates chromosomal rearrangements and the integration of foreign genetic elements [7]. This intrinsic adaptability, coupled with antibiotic selective pressure, has contributed to the establishment of endemic multidrug-resistant *A. baumannii* strains in healthcare settings [8].

Since the introduction of sequencing technologies in microbiology, considerable efforts have been devoted to elucidating the mechanisms underlying multidrug resistance in *A. baumannii*. The genomes of both collection and endemic strains have been subjected to detailed comparative analysis [9–11]. Resistance islands, plasmids, and mobile genetic elements have been identified and characterized [9–12], while the genetic environment of resistance determinants has been studied in detail [11,13]. Based on these investigations, the authors have drawn conclusions regarding the dissemination of genetic determinants among strains and species.

Fournier et al. [9] reported the presence of genes within a resistance island acquired from *Pseudomonas*, *Salmonella*, and *Escherichia* species.

Zhu et al. [11] concluded that horizontal gene transfer of mobile genetic elements between chromosomes and plasmids may be a significant contributor to the rapid spread of resistance among *A. baumannii* strains.

Holt et al. [14] provided experimental evidence that natural transformation is a major driver of resistance gene acquisition in Global Clone 1.

Frenk et al. [8] observed that sequence type and clade distribution remained stable over time, suggesting that horizontal gene transmission and the de novo emergence of multidrug-resistant *A. baumannii* are rare events, yet crucial to the evolutionary success of specific clades and sequence types.

To date, large-scale systematic investigations of invasive *A. baumannii* isolates integrating phenotypic and genotypic data remain scarce in Bulgaria. The present study aimed to characterize the prevalent phenotypes and genotypes among 36 invasive *A. baumannii* isolates recovered from inpatients at a university hospital over a six-year period (2019–2024)

and elucidate the relationships between phenotypic resistance profiles and the underlying genetic determinants.

2. Materials and Methods

2.1. Bacterial Isolates

During the study period (2019–2024), a total of 67 non-duplicate *Acinetobacter calcoaceticus–baumannii* complex (Acb) isolates were recovered from blood cultures at the University Hospital. The isolates were routinely identified to the Acb-complex level using the VITEK 2 Compact system (bioMérieux, Marcy-l'Étoile, France) and tested for antimicrobial susceptibility by disk diffusion and VITEK 2. They were then stored in skim-milk broth at -80°C pending further study. From these 67 non-duplicate isolates, 36 (53.7%) were selected for further investigation. The selection was based on two criteria: (i) representation of every calendar year of the study period and (ii) coverage of defined susceptibility phenotypes, based on the routine susceptibility results for eight agents (MEM, IMI, GEN, AMK, TOB, CIP, LEV, and COL) (Supplementary Table S1). All rare phenotypes were retained, whereas the predominant phenotypes were proportionally subsampled to reduce their overrepresentation relative to the rarer phenotypes, resulting in the exclusion of 31 isolates from further analysis. One of the two isolates susceptible only to tobramycin and colistin was additionally excluded owing to non-viability, yielding a final collection of 36 isolates. For the cohort of 36 selected isolates, species-level identification was confirmed by MALDI-TOF MS (VITEK MS, bioMérieux, Marcy-l'Étoile, France) and two PCR-based methods targeting the *bla*_{OXA-51-like} gene [15] and *gyrB* [16]. *A. baumannii* ATCC 19606 was used as the positive control strain for PCR-based species identification.

The study was approved by the Ethics Committee of the Medical University of Pleven (Approval No. 794, approved in June 2024) and conducted in accordance with the Declaration of Helsinki and its later amendments. As the research focused solely on bacterial isolates and no personal patient information was used, individual informed consent was not required.

2.2. Antimicrobial Susceptibility Testing

Minimum inhibitory concentrations (MICs) were determined by the gradient diffusion method (MIC Test Strip, Liofilchem, Roseto degli Abruzzi, Italy) for imipenem (IPM), meropenem (MEM), gentamicin (GEN), amikacin (AMK), tobramycin (TOB), levofloxacin (LEV), trimethoprim/sulfamethoxazole (SXT), and minocycline (MIN). MICs for cefiderocol (FDC) and colistin (COL) were determined by the broth microdilution method using ComASP™ Cefiderocol (0.008–128 mg/L) and ComASP™ Colistin (0.25–16 mg/L) panels (Liofilchem, Roseto degli Abruzzi, Italy), respectively. *Escherichia coli* ATCC 25922, *P. aeruginosa* ATCC 27853, and *A. baumannii* ATCC 19606 served as quality control strains. MICs were interpreted according to EUCAST clinical breakpoints (v16.0, 2026). For minocycline, CLSI criteria (M100, 35th edition, 2025) [17] were applied.

2.3. Definition for MDR-AB, XDR-AB, and PDR-AB

Multidrug-resistant (MDR) *A. baumannii* isolates were defined as non-susceptible to at least one agent in three or more antimicrobial categories, whereas extensively drug-resistant (XDR) isolates were defined as non-susceptible to at least one agent in all but two or fewer antimicrobial categories. Pandrug-resistant (PDR) isolates were defined as non-susceptible to all agents in all antimicrobial categories, according to previously described criteria [18].

2.4. DNA Isolation

Total DNA from all investigated strains was isolated using the DNeasy Blood and Tissue Kit (QIAGEN, Hilden, Germany), according to the manufacturer's instructions, from 3 mL of overnight culture inoculated with a single colony.

2.5. PCR Screening for Antimicrobial Resistance Genes (ARGs)

Amplifications were performed on a qTOWER³ G touch thermal cycler (Analytik Jena, Jena, Germany). Genes encoding carbapenemase production (*bla*_{OXA-23-like}, *bla*_{OXA-24/40-like}, *bla*_{OXA-58-like}, *bla*_{NDM-like}, *bla*_{VIM-like}, *bla*_{IMP-like}, and *bla*_{KPC-like}) were investigated. The specific allelic variant within the *bla*_{OXA-24/40-like} group was subsequently determined by whole-genome sequencing (WGS) (see Section 3.7, Resistome Analysis). The association of IS*Aba1* with *bla*_{OXA-23-like} was assessed by PCR using an IS*Aba1*-specific forward primer and a *bla*_{OXA-23-like}-specific reverse primer. In addition, PCR assays were performed for the *armA* gene, encoding a 16S rRNA methylase associated with high-level aminoglycoside resistance, and the *sul1* gene, associated with sulfonamide resistance and located within class 1 integrons.

PCR assays were performed using previously described protocols with minor modifications, mainly to the annealing temperatures [18–23]. All reactions were carried out in a final volume of 20 µL containing OneTaq[®] 2× Master Mix (New England Biolabs, Ipswich, MA, USA) and 20 ng of template DNA. The class D carbapenemase genes were detected in a multiplex hot-start reaction (94 °C for 4 min; 36 cycles of 94 °C for 30 s, 52 °C for 35 s, 70 °C for 75 s; final extension 71 °C for 3 min). The *bla*_{VIM-like}, *bla*_{IMP-like}, *bla*_{NDM-like}, and *bla*_{KPC-like} genes were amplified by touchdown PCR (95 °C for 3 min; 5 cycles with the annealing temperature decreasing by 1.6 °C per cycle from 63 °C, followed by 20–25 cycles at 54 °C; extension 70 °C for 1 min). The *armA* gene was amplified at an annealing temperature of 50 °C (30 cycles), and *sul1* by touchdown PCR (62 °C decreasing by 1 °C per cycle for 7 cycles, followed by 25 cycles at 54 °C). The IS*Aba1*–*bla*_{OXA-23-like} assay was performed at 52 °C for 30 cycles. The expected amplicon size of the IS*Aba1*–*bla*_{OXA-23-like} PCR assay was verified in silico using Primer-BLAST (National Center for Biotechnology Information (NCBI), Bethesda, MD, USA; <https://www.ncbi.nlm.nih.gov/tools/primer-blast/>, accessed on 4 July 2025).

Primer sequences and expected amplicon sizes are provided in Table 1. PCR products were separated by agarose gel electrophoresis and visualized by ethidium bromide staining. The following strains were used as controls:

- *P. aeruginosa* 4782 (*bla*_{IMP-100}/*bla*_{VIM-4})
- *P. aeruginosa* 1793 (*bla*_{NDM}/*bla*_{GES})
- *K. pneumoniae* ATCC 2814 (*bla*_{KPC})
- *A. baumannii* strains (*bla*_{OXA23-like}/*bla*_{OXA24/40-like})
- *A. baumannii* (*bla*_{OXA58})

Table 1. PCR primer sequences and expected amplicon sizes used for antimicrobial resistance gene detection in this study.

Target	Primer Sequence (5'–3')	Product Size (bp)	Reference
<i>bla</i> _{OXA-23-like}	F: GATCGGATTGGAGAACCAGA R: ATTTCTGACCGCATTTCAT	501	[19]
<i>bla</i> _{OXA-24/40-like}	F: GGTTAGTTGGCCCCCTTAAA R: AGTTGAGCGAAAAGGGGATT	246	[19]

Table 1. Cont.

Target	Primer Sequence (5'–3')	Product Size (bp)	Reference
<i>bla</i> _{OXA-58-like}	F: AAGTATTGGGGCTTGTGCTG R: CCCCTCTGCGCTCTACATAC	599	[19]
<i>bla</i> _{NDM-like}	F: AACACAGCCTGACTTTTCG R: TGATATTGTCACTGGTGTGG	111	[20]
<i>bla</i> _{VIM-like}	F: GATGGTGTGTTGGTCGCATA R: CGAATGCGCAGCACCAG	390	[21]
<i>bla</i> _{IMP-like}	F: TTGACACTCCATTTACTGCTA R: TCATTTGTAAATTCAGATGCATA	172	[19]
<i>bla</i> _{KPC-like}	F: TCTGGACCGCTGGGAGCTGG R: TGCCCGTTGACGCCCAATCC	399	[22]
<i>armA</i>	F: ATTCTGCCTATCCTAATTGG R: ACCTATACTTTATCGTCGTC	315	[23]
<i>sul1</i>	F: ACGGTGTTTCGGCATTCT R: TTTGAAGGTTTCGACAGC	580	[24]
IS _{Aba1} - <i>bla</i> _{OXA-23-like}	F: CACGAATGCAGAAGTTG R: ATTTCTGACCGCATTTCAT	1404	This study

F, forward primer; R, reverse primer; bp, base pairs; IS, insertion sequence. Gene abbreviations are as defined in the main text.

The *A. baumannii* strains are in-house reference strains provided by the National Center of Infectious and Parasitic Diseases (NCIPD), Sofia, Bulgaria, previously characterized by PCR and sequencing.

2.6. Whole-Genome Sequencing (WGS)

Based on the combined phenotypic susceptibility and PCR screening results, seven distinct genotypic–phenotypic resistance profiles were defined. A subset of 17 isolates, representative of these profiles, was selected for WGS to enable comprehensive resistome analysis. Genomic DNA was randomly fragmented, size-selected, ligated to adapters, and PCR-amplified to generate sequencing libraries. The libraries were sequenced on an Illumina NovaSeq X Plus platform (Novogene, Cambridge, UK), producing 2 × 150 bp paired-end reads.

2.7. Draft Genome Assembly

The genome assembly procedure involved several steps: quality control (FastQC v0.12.1, <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>, accessed on 15 April 2026), raw read preprocessing (Trimmomatic v0.39) [25], and genome assembly (SPAdes v4.2.0; --isolate and --cov-cutoff 5.0 flags) [26]. The resulting scaffold sequences were uploaded to the Galaxy online platform [27], where draft genome metrics were evaluated (Galaxy v5.3.0). All software tools were run with default parameters unless otherwise specified.

2.8. Resistome and Multilocus Sequence Typing (MLST) Analyses

All uploaded draft genome sequences were screened for AMR genes using the ABRIcate tool (Galaxy v1.4.0) with the following settings: the Comprehensive Antibiotic Resistance Database (CARD), minimum DNA identity of 70%, and minimum DNA coverage of 60%. In addition, the *gyrA* and *parC* coding sequences were manually analyzed for mutations in their quinolone resistance-determining regions (QRDRs).

Finally, MLST analysis was performed on the assembled scaffolds using the MLST tool (Galaxy v2.22.0, <https://usegalaxy.eu/>) and the two available *A. baumannii* schemes.

2.9. Phylogenomic Analysis

Clinical *A. baumannii* genomes were retrieved from the Isolates Browser integrated into the NCBI Pathogen Detection system for Albania, Bosnia and Herzegovina, Bulgaria, Croatia, Greece, Kosovo, Montenegro, North Macedonia, Romania, Serbia, and Turkey. To minimize sampling bias, a single representative genome was selected from each available single-nucleotide polymorphism (SNP) cluster for every country, preferentially choosing bloodstream isolates when available. All selected draft genome sequences were annotated using Prokka v1.14.6 [28].

Pangenome analysis was subsequently performed using Roary v3.13.0 [29] with a minimum BLASTP identity threshold of 95% and a core genome definition of 99%, generating a core gene alignment for downstream analyses. All SNPs were extracted from the resulting multi-FASTA alignment using the Find SNP Sites tool (Galaxy v2.5.1), and pairwise SNP distances were calculated using the SNP Distance Matrix tool (Galaxy v0.8.2). The resulting distance matrix was subjected to hierarchical clustering using the Python data visualization library seaborn (v0.13.2).

A phylogenetic tree was reconstructed from the core genome SNP alignment using RAxML v8.2.12 [30], with branch support estimated from 1000 bootstrap replicates. The resulting tree was visualized using the Interactive Tree of Life (iTOL) web server [31].

2.10. Statistical Analysis

Statistical analyses were conducted using jamovi (v2.7; The jamovi Project, 2025) [32], based on R (v4.5; R Core Team, 2025) [33]. Associations between the presence of resistance genes and antimicrobial susceptibility were assessed using Fisher's exact test for categorical variables. Differences in MIC values between gene-positive and gene-negative groups were assessed using the Mann–Whitney U test. Differences among aminoglycosides within *armA*-negative, *bla*_{OXA-23-like}-positive isolates were assessed using the Friedman test, with pairwise comparisons by Wilcoxon signed-rank test. A *p*-value < 0.05 was considered statistically significant. Genotype subgroups with very small sample sizes ($n \leq 4$) were summarized descriptively, without formal statistical testing, owing to insufficient statistical power.

3. Results

3.1. Patient Demographic and Clinical Characteristics

Supplementary Table S2 presents the demographic and clinical characteristics of the 36 patients (16 men, 20 women). The age distribution by category was as follows: 0–18 years (1/36), 19–40 years (6/36), 41–65 years (17/36), and >65 years (12/36). Most patients (28/36) were treated in intensive care units, followed by surgical units (6/36), with one case each in the Gastroenterology Clinic and the Acute Cardiac Care Unit. The mean length of hospital stay was 22.6 days. Patients were diagnosed with polytrauma, including one case of second-degree burns and one of toxic gas exposure (8/36); acute abdominal disease, including peritonitis, pancreatitis, ileus, colon malignancies with complications, and liver failure (11/36); vascular and cardiac disease (5/36); pulmonary infections (6/36), five of them associated with COVID-19; soft-tissue infections (3/36); systemic lupus erythematosus (1/36); thrombocytopenia (1/36); and viral meningitis (1/36). Prior to the isolation of *A. baumannii*, broad-spectrum antibiotic therapy was documented in 32/36 patients. Mortality was high (26/36), with 10 patients discharged following clinical improvement.

3.2. Identification

A total of 36 invasive isolates were confirmed as *A. baumannii* by MALDI-TOF MS (VITEK MS) and PCR-based methods targeting the intrinsic *bla*_{OXA-51}-like gene and the *gyrB* gene.

3.3. Antimicrobial Susceptibility

The results of antimicrobial susceptibility testing are presented in Table 2. Carbapenem resistance was observed in 34 of 36 isolates (94.4%), with MICs of ≥ 32 mg/L for both meropenem and imipenem. All isolates were susceptible to colistin. With respect to aminoglycosides, resistance rates varied by agent. Gentamicin showed the lowest resistance, with 16/36 isolates (44.4%) categorized as susceptible and 20/36 (55.6%) as resistant. In contrast, tobramycin and amikacin resistance was considerably higher, with only 6/36 (16.7%) and 2/36 (5.6%) isolates susceptible and 30/36 (83.3%) and 34/36 (94.4%) resistant, respectively. Resistance to trimethoprim/sulfamethoxazole was observed in 35/36 isolates (97.2%). Non-susceptibility (resistant plus intermediate) to levofloxacin was observed in 35/36 isolates (97.2%). Minocycline resistance (MIC range 8–24 mg/L) was observed in 18/36 isolates (50.0%), 10/36 (27.8%) showed intermediate susceptibility (MIC range 1.5–2 mg/L), and 8/36 (22.2%) were susceptible (MIC ≤ 1 mg/L).

Table 2. Antimicrobial susceptibility of the 36 *A. baumannii* isolates, n (%).

Category	MEM	IPM	GEN	AMK	TOB	LEV	SXT	MIN	COL
Susceptible	2 (5.6)	2 (5.6)	16 (44.4)	2 (5.6)	6 (16.7)	1 (2.8)	1 (2.8)	8 (22.2)	36 (100)
Intermediate	–	–	–	–	–	13 (36.1)	–	10 (27.8)	–
Resistant	34 (94.4)	34 (94.4)	20 (55.6)	34 (94.4)	30 (83.3)	22 (61.1)	35 (97.2)	18 (50.0)	0 (0)

n, number of isolates; MEM, meropenem; IPM, imipenem; GEN, gentamicin; AMK, amikacin; TOB, tobramycin; LEV, levofloxacin; SXT, trimethoprim/sulfamethoxazole; MIN, minocycline; COL, colistin. Dashes (–) indicate that no intermediate category is defined for that agent by the applied breakpoint standard. Susceptibility was interpreted according to EUCAST v16.0 clinical breakpoints, except for minocycline (CLSI M100, 35th ed., 2025) [17].

As an exploratory sub-analysis, cefiderocol susceptibility testing was performed on a subset of nine isolates, selected to represent different genotypic profiles identified in this study. MIC values were interpreted using the non-species-specific PK/PD cutoff ($S \leq 2$ mg/L, EUCAST v16.0). Seven of the nine isolates fell within this range, whereas two (MIC 4 mg/L) exceeded the PK/PD susceptibility cutoff.

Overall, 34/36 isolates (94.4%) met the criteria for XDR classification, remaining susceptible only to colistin. The findings regarding cefiderocol should be considered preliminary, given the limited number of isolates tested.

3.4. PCR Screening for Antimicrobial Resistance Genes

The intrinsic *bla*_{OXA51-like} determinant was detected in all 36 isolates (100%), confirming their identification as *A. baumannii*. The prevalence of *bla*_{OXA23-like} and *bla*_{OXA24/40-like} was 80.6% (29/36) and 13.9% (5/36), respectively. Using a primer combination consisting of the forward primer targeting *IsaAba1* and reverse primer specific for *bla*_{OXA23-like}, all *bla*_{OXA23-like} positive isolates yielded a 1404 bp amplicon, indicating the presence of *IsaAba1* upstream of the gene and suggesting promoter-mediated activation. No other *bla* genes were detected. The *armA* gene, encoding a 16S rRNA methylase, was identified in 50.0% (18/36) of the isolates. Notably, it was detected exclusively in combination with *bla*_{OXA23-like}. No isolate simultaneously harbored *bla*_{OXA24/40-like} and *armA*, and no isolate harbored *armA* as a sole resistance determinant. The *sul1* gene, associated with resistance to sulfamethoxazole, was detected in 80.6% (29/36) of the isolates.

3.5. Association Between Resistance Phenotypes and Genotypes

Table 3 summarizes the relationship between antimicrobial resistance phenotypes and the detected resistance genes. The complete MIC values and detected resistance genes for all 36 isolates are provided in Table S2, and the detailed statistical output is in Table S3 (Supplementary Materials). The presence of either *bla*_{OXA-23}-like or *bla*_{OXA-24/40}-like was associated with high MIC values for meropenem and imipenem (≥ 32 mg/L), whereas their absence corresponded to MIC values within the susceptible range (Fisher's exact test, $p = 0.002$).

Table 3. Relationship between genotypic and phenotypic resistance profiles in the 36 *A. baumannii* isolates (MIC in mg/L).

Genotype	Genotypic Resistance Profiles	Phenotypic Resistance Profiles (MIC in mg/L)								No.	Percent
		MEM	IPM	GEN	AMK	TOB	LEV	SXT	MIN		
GT1	<i>bla</i> _{OXA-23} + <i>bla</i> _{OXA-24} − <i>armA</i> + <i>sul1</i> +	≥ 32	≥ 32	≥ 1024	≥ 256	≥ 1024	1–32	≥ 32	12–24	13	36.1%
GT2	<i>bla</i> _{OXA-23} + <i>bla</i> _{OXA-24} − <i>armA</i> − <i>sul1</i> +	≥ 32	≥ 32	2–6	96–256	12–24	1–8	≥ 32	2	11	30.6%
GT3	<i>bla</i> _{OXA-23} + <i>bla</i> _{OXA-24} − <i>armA</i> + <i>sul1</i> −	≥ 32	≥ 32	≥ 1024	≥ 256	≥ 1024	12–32	1.5–4	8–12	5	13.9%
GT4	<i>bla</i> _{OXA-23} − <i>bla</i> _{OXA-24} + <i>armA</i> − <i>sul1</i> +	≥ 32	≥ 32	1.5–2	64–256	0.5–1	0.75–1	6–8	0.5	4	11.1%
GT5	<i>bla</i> _{OXA-23} − <i>bla</i> _{OXA-24} + <i>armA</i> − <i>sul1</i> −	≥ 32	≥ 32	3	64	1.5	1	0.5	0.5	1	2.8%
GT6	<i>bla</i> _{OXA-23} − <i>bla</i> _{OXA-24} − <i>armA</i> − <i>sul1</i> +	2	1	4	8	16	1	≥ 32	2	1	2.8%
GT7	<i>bla</i> _{OXA-23} − <i>bla</i> _{OXA-24} − <i>armA</i> − <i>sul1</i> −	0.125	0.5	0.75	4	1	0.47	0.25	0.38	1	2.8%
										36	100%

No., number of isolates; MEM, meropenem; IPM, imipenem; GEN, gentamicin; AMK, amikacin; TOB, tobramycin; LEV, levofloxacin; SXT, trimethoprim/sulfamethoxazole; MIN, minocycline. +, gene detected; −, gene not detected. *bla*_{OXA-23} denotes *bla*_{OXA-23}-like and *bla*_{OXA-24} denotes *bla*_{OXA-24/40}-like. MIC values are given as single values or observed ranges.

The *armA* gene was associated with significantly higher MICs for all three aminoglycosides tested (Mann–Whitney U test, $p < 0.001$), with *armA*-positive isolates reaching 768–1024 mg/L for gentamicin, 256 mg/L for amikacin, and 512–1024 mg/L for tobramycin. In *armA*-negative, *bla*_{OXA-23}-like-positive isolates, gentamicin MICs decreased to the susceptible or low-level resistance range (2–6 mg/L), and tobramycin MICs dropped markedly from ≥ 1024 to 12–24 mg/L (Mann–Whitney U test, $p < 0.0001$ for both). Amikacin MIC values, however, remained elevated compared with gentamicin and tobramycin (Friedman test, $p < 0.0001$; pairwise Wilcoxon signed-rank tests, $p = 0.001$ for all comparisons).

Notably, elevated minocycline MICs were also significantly associated with *armA* carriage (Mann–Whitney U test, $p < 0.001$; 8–24 mg/L), and a strong concordance was observed between *armA* carriage and minocycline resistance (Fisher's exact test, $p < 0.001$), with susceptibility to minocycline observed only among *armA*-negative isolates. These findings suggest a potential co-linkage between *armA* and minocycline resistance determinants. Similar associations extended to fluoroquinolones: among the 29 *bla*_{OXA-23}-like-positive isolates, those additionally carrying *armA* (18/29) showed significantly higher levofloxacin MICs (median 16, range 1–32 mg/L) than *armA*-negative *bla*_{OXA-23}-like carriers (11/29; median 1, range 1–8 mg/L; Mann–Whitney U test, $p < 0.001$), with a corresponding significant association between *armA* carriage and levofloxacin resistance (Fisher's exact test, $p = 0.001$).

A significant difference was also found between *armA*-negative isolates harboring *bla*_{OXA-23}-like (11/29) and *bla*_{OXA-24/40}-like-positive ones (5/36, all *armA*-negative). The former group showed significantly higher gentamicin MICs (median 4 vs. 2 mg/L; $p = 0.003$) and significantly higher tobramycin MICs (median 16 vs. 0.75 mg/L; $p = 0.001$). Amikacin MICs did not differ significantly between the two genotypes ($p = 1.000$), remaining elevated in both. Minocycline MICs were also significantly higher among *bla*_{OXA-23}-like carriers than among *bla*_{OXA-24/40}-like-positive isolates (median 1.5 vs. 0.5 mg/L; $p = 0.002$).

The isolate carrying only *sul1* remained susceptible to carbapenems and amikacin and showed intermediate minocycline susceptibility and low-level resistance to gentamicin, tobramycin, and levofloxacin, but was highly resistant to trimethoprim/sulfamethoxazole (MIC $\geq$ 32 mg/L). The isolate lacking all screened genes was fully susceptible to all agents. Among the nine isolates tested for cefiderocol, the *armA*-positive genotypes GT1 and GT3 showed MICs of 0.25–0.50 mg/L, whereas the *armA*-negative genotypes GT2, GT4, and GT6 showed higher values (1–4 mg/L). The gene-negative isolate (GT7) had the lowest MIC of the subset (0.125 mg/L). Given the limited size of the subset, this observation should be regarded as preliminary rather than a firm conclusion. In addition, these results were excluded from the general XDR susceptibility statement. Detailed cefiderocol MIC values by isolate and genotype are presented separately in Table S4 (Supplementary Materials).

3.6. Draft Genome Assembly and Evaluation

In total, 17 of the 36 isolates were subjected to whole-genome sequencing. The resulting data were used to perform MLST and resistome analysis. The draft genome assembly metrics are summarized in Table 4.

Table 4. Draft genome assembly metrics of the studied *A. baumannii* isolates.

Isolate	Genome Size (Mbp)	GC%	N50 (bp)	No. of Contigs (≥kbp)	ST	Alleles						
						<i>cpn60</i>	<i>fusA</i>	<i>gltA</i>	<i>pyrG</i>	<i>recA</i>	<i>rplB</i>	<i>rpoB</i>
1	4.24	40.08	288,607	97	2	2	2	2	2	2	2	2
5	4.09	39.69	288,607	72	2	2	2	2	2	2	2	2
9	4.24	40.10	341,347	77	2	2	2	2	2	2	2	2
10	4.19	39.80	480,198	58	2	2	2	2	2	2	2	2
11	4.28	40.39	340,884	73	2	2	2	2	2	2	2	2
15	4.04	39.40	461,212	51	2	2	2	2	2	2	2	2
25	4.13	39.35	461,311	60	2	2	2	2	2	2	2	2
32	4.24	40.18	292,960	82	2	2	2	2	2	2	2	2
41	4.11	39.29	630,075	54	2	2	2	2	2	2	2	2
71	4.06	39.53	323,167	53	636	2	1	2	2	2	1	1
75	4.11	39.25	461,102	55	2	2	2	2	2	2	2	2
86	4.10	39.80	323,167	76	636	2	1	2	2	2	1	1
97	4.42	41.25	461,411	66	2	2	2	2	2	2	2	2
111	4.00	39.50	434,910	82	2	2	2	2	2	2	2	2
112	3.93	39.01	461,078	79	2	2	2	2	2	2	2	2
120	3.93	39.67	2,183,114	15	-	3	~3	2	1	6	1	~60
121	4.10	39.89	387,851	56	636	2	1	2	2	2	1	1

ST, sequence type.

The genome sizes ranged from 3.93 to 4.42 Mbp, with a GC content of 39.01–41.25%. The N50 values ranged from 288,607 to 2,183,114 bp, and the number of contigs (≥1 kbp) varied from 15 to 97. Notably, isolate 120—the only susceptible isolate in the collection—exhibited markedly superior assembly metrics (only 15 contigs; N50 = 2,183,114 bp). This isolate could not be assigned to a sequence type by the Pasteur MLST scheme and was classified only by the Oxford MLST scheme as ST 1061.

3.7. Resistome Analysis

Table 5 presents the resistance determinants identified in the seven genotypes. *bla*_{OXA-72} was the only *bla*_{OXA-24/40-like} variant detected. All *armA*-positive isolates also harbored *tet(B)*, the main determinant for minocycline resistance. The *bla*_{OXA-23-like}-positive isolates and the isolate positive only for *sul1* and lacking *armA* carried *aac(6′)-Ib7*, *aph(3′)-*

Vla. One of these additionally harbored *aph(3')-Ia*. The *bla*_{OXA-24/40-like}-positive isolates carried only *aph(3')-Vla*.

Table 5. Whole-genome sequencing data and genotype classification of the studied isolates.

Genotype	Isolate No.	Department	Date of Isolation	Resistance Genes	QRDR Mutations	ST
GT1	32	ICU1	November 2019	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-73} , <i>armA</i> , <i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>tet(B)</i> , <i>sul1</i> , <i>catA1</i> , <i>mphE</i> , <i>msrE</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	2
	5	GEC	July 2020	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-73} , <i>armA</i> , <i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>tet(B)</i> , <i>sul1</i> , <i>catA1</i> , <i>mphE</i> , <i>msrE</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	2
	111	AbS	November 2024	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-73} , <i>armA</i> , <i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>aph(3')-Vla</i> , <i>tet(B)</i> , <i>sul1</i> , <i>catA1</i> , <i>mphE</i> , <i>msrE</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	2
	112	ICU1	December 2024	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-73} , <i>armA</i> , <i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>aph(3')-Vla</i> , <i>tet(B)</i> , <i>sul1</i> , <i>catA1</i> , <i>mphE</i> , <i>msrE</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	2
GT2	41	ICU1	February 2020	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-11} , <i>bla</i> _{OXA-20} , <i>aac(6')-Ib7</i> , <i>aph(3')-Vla</i> , <i>aph(3')-Ia</i> , <i>sul1</i>	<i>gyrA</i> : S81L	2
	75	ICU1	April 2022	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-11} , <i>bla</i> _{OXA-20} , <i>aac(6')-Ib7</i> , <i>aph(3')-Vla</i> , <i>sul1</i>	<i>gyrA</i> : S81L	2
	10	ICU1	November 2023	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-11} , <i>bla</i> _{OXA-20} , <i>aac(6')-Ib7</i> , <i>aph(3')-Vla</i> , <i>sul1</i>	<i>gyrA</i> : S81L	2
	25	ICU1	March 2023	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-11} , <i>bla</i> _{OXA-20} , <i>aac(6')-Ib7</i> , <i>aph(3')-Vla</i> , <i>sul1</i>	<i>gyrA</i> : S81L	2
	97	DSS	February 2024	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-11} , <i>bla</i> _{OXA-20} , <i>aac(6')-Ib7</i> , <i>aph(3')-Vla</i> , <i>sul1</i>	<i>gyrA</i> : S81L	2
GT3	1	ICU1	January 2019	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-73} , <i>armA</i> , <i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>aph(3')-Vla</i> , <i>tet(B)</i> , <i>mphE</i> , <i>msrE</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	2
	9	ICU1	April 2021	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-73} , <i>armA</i> , <i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>aph(3')-Vla</i> , <i>tet(B)</i> , <i>catA1</i> , <i>mphE</i> , <i>msrE</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	2
	11	ICU1	April 2021	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-73} , <i>armA</i> , <i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>aph(3')-Vla</i> , <i>tet(B)</i> , <i>catA1</i> , <i>mphE</i> , <i>msrE</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	2
GT4	71	ICU1	July 2021	<i>bla</i> _{OXA-72} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-74} , <i>aph(3')-Vla</i> , <i>catA1</i> , <i>sul1</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	636
	86	ICU1	December 2023	<i>bla</i> _{OXA-72} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-74} , <i>aph(3')-Vla</i> , <i>catA1</i> , <i>sul1</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	636
GT5	121	ICU1	February 2024	<i>bla</i> _{OXA-72} , <i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-74} , <i>aph(3')-Vla</i>	<i>gyrA</i> : S81L; <i>parC</i> : S84L	636
GT6	15	ICU1	August 2021	<i>bla</i> _{OXA-66} , <i>bla</i> _{ADC-11} , <i>bla</i> _{OXA-20} , <i>aac(6')-Ib7</i> , <i>aph(3')-Ia</i> , <i>sul1</i>	<i>gyrA</i> : S81L	2
GT7	120	ICU1	July 2024	<i>bla</i> _{OXA-314} , <i>bla</i> _{ADC-7}	none	1061 Oxf

GT, genotype; ST, sequence type; QRDR, quinolone resistance-determining region; ICU1, Intensive Care Unit 1; GEC, Department of Gastroenterology; AbS, Department of Abdominal Surgery; DSS, Department of Suppurative Surgery.

Comparison of the QRDRs of the isolates with the corresponding region of *A. baumannii* ATCC 19606 revealed amino-acid substitutions previously reported to play a role in quinolone resistance [34]. All but one isolate possessed S81L substitution in *GyrA*. The single isolate lacking this mutation was also the only one negative for all resistance genes screened by PCR. Isolates co-harboring *bla*_{OXA-23}-like and *armA*, as well as those positive for *bla*_{OXA-72}, additionally carried S84L substitution in *ParC*.

3.8. MLST Analysis

MLST analysis (Pasteur scheme) assigned all but one of the sequenced isolates to either ST2 or its triple-locus variant ST636. Thirteen isolates were ST2, the globally dominant high-risk clone corresponding to International Clone II (IC2) [35–37], whereas three isolates, all harboring *bla*_{OXA-72} (a *bla*_{OXA-24/40}-like variant), were ST636. The remaining isolate could not be classified by the Pasteur scheme but was assigned to ST1061 by the Oxford scheme. It was negative for all tested acquired resistance genes, susceptible to all antimicrobials tested, and harbored two distinct intrinsic β -lactamase genes: *bla*_{ADC-7} and *bla*_{OXA-314}.

3.9. Phylogenomic Analysis of Clinical *A. baumannii* Isolates from the Balkans

To investigate the epidemiological context of the studied isolates, a collection of 200 non-duplicate clinical *A. baumannii* genomes representing isolates recovered from 11 Balkan countries was assembled as described above. The geographic distribution of these strains is shown in Figure 1A.



Figure 1. Cont.




Figure 1. Phylogenomic analysis of clinical *A. baumannii* isolates from the Balkans retrieved from the NCBI Pathogen Detection system. (A) Geographic origins of the included isolates. (B) Hierarchical clustering based on the SNP distance matrix of the isolates analyzed in this study. (C) Maximum-likelihood phylogenetic tree reconstructed from single-nucleotide polymorphisms (SNPs) identified in the core genome alignment generated by pangenome analysis. Isolates sequenced in this study are highlighted in red.

Hierarchical clustering of the SNP distance matrix revealed that the isolates analyzed in this study formed three distinct clusters: cluster 1 (isolates 1, 5, 9, 11, 32, 111, and 112), cluster 2 (isolates 10, 15, 25, 41, 75, and 97), and cluster 3 (isolates 71, 86, and 121) (Figure 1B). Pairwise SNP differences within these clusters ranged from 1 to 16 SNPs, indicating close relatedness, with the exception of isolates 111 and 112, which were genetically identical in this assay (0 SNP differences) but differed from the remaining members of cluster 1 by up to 203 SNPs. Isolate 120 remained the only isolate separated by more than 20,000 SNPs from all other isolates, indicating that it is not closely related to any of them.

The phylogenetic tree (Figure 1C) revealed that the closest relative of cluster 1 was genome GCA_021572415, which was recovered from a patient at Henry Dunant Hospital in Athens, Greece. The closest relatives of clusters 2 and 3 were genomes GCA_058149165.1 and GCA_058149125.1, respectively. Both genomes were generated as part of the 2026 ECDC genomic survey of carbapenem-resistant *A. baumannii* in Europe; however, no additional epidemiological information is currently available.

4. Discussion

4.1. Carbapenem Resistance Associated with *bla*_{OXA-23}-like and *bla*_{OXA-72}

Our data revealed the relationship between genotypic and phenotypic resistance profiles of 36 *Acinetobacter baumannii* invasive isolates recovered from blood cultures. The tested isolates showed a high prevalence of carbapenem resistance (94.4%). They were positive for *bla*_{OXA-23}-like (29/36) and *bla*_{OXA-72}, a *bla*_{OXA24/40}-like variant (5/36). No isolate harbored both genes simultaneously. The *bla*_{OXA-58}-like, *bla*_{KPC}-like, and metallo- β -lactamase genes (*bla*_{NDM}, *bla*_{IMP}, *bla*_{VIM}) were not detected.

These findings are consistent with the increasing trend of carbapenem resistance observed in Bulgaria. The first identification of OXA-23-producing isolates from our hospital was documented by Stoeva et al. [38] among isolates recovered between 1999 and 2006. However, *bla*_{OXA-24/40}-like genes were not detected during this period. Over approximately two decades, carbapenem resistance rose from 61.7% [39] to 94.4% within the same hospital setting. A similar increase from 59.4% (2004–2011) to 93–98.7% (2014–2022) was documented by Strateva et al. in a multicenter study [40]. The same authors reported a high prevalence of *bla*_{OXA-24/40}-like as the sole acquired carbapenem-hydrolyzing class D β -lactamase (CHDL) in isolates from Varna and two Sofia university hospitals during 2014–2022, indicating expansion of strain lineages producing this enzyme. In addition, they identified strains co-producing both CHDLs, which were not observed in our study. Across the Balkans, various surveys report *bla*_{OXA-24/40}-like (including *bla*_{OXA-72}) and *bla*_{OXA-23}-like occurring either as mutually exclusive determinants or in simultaneous co-occurrence [41–43]. These reports illustrate the dissemination of *bla*_{OXA-72} across the region.

Current ECDC surveillance data (2023) confirm a pronounced east–west gradient in carbapenem resistance among invasive *A. baumannii* isolates across Europe. Resistance rates exceed 50% throughout southern and eastern Europe (Italy 80.8%, Poland 78.2%), with striking data from the Balkans (Greece 94.6%, Romania 92.7%) compared with below 5% in northern and western Europe (Germany 3.5%, Norway 1.6%) [4].

Globally, *bla*_{OXA-23}-like remains the most prevalent acquired CHDL gene, accounting for approximately 88% of carbapenemase genes in carbapenem-resistant *A. baumannii*, while *bla*_{OXA-24/40}-like and *bla*_{OXA-58}-like are detected less frequently [8,35,44,45]. Unlike Greece, Turkey, and other countries in the Mediterranean region, where a temporal shift from *bla*_{OXA-58} to *bla*_{OXA-23} predominance occurred between the first and second decades of the century [45–48], Bulgaria has consistently demonstrated *bla*_{OXA-23}-like dominance throughout this period, with only sporadic detections of *bla*_{OXA-58}-like [38,40].

4.2. Aminoglycoside Resistance Associated with *armA*

In the present study, gentamicin demonstrated the highest activity among the aminoglycosides. Isolates carrying *armA* exhibited high-level resistance to all three tested aminoglycosides. In *armA*-negative, *bla*_{OXA-23-like}-positive isolates, MIC values for gentamicin and tobramycin decreased substantially, whereas amikacin MICs remained elevated. In *bla*_{OXA-72}-positive isolates, gentamicin and tobramycin MICs fell within the susceptibility range. Notably, *armA* was detected exclusively in combination with the *bla*_{OXA-23-like} gene. Multiple publications support the co-persistence of *bla*_{OXA-23-like} and *armA* in MDR/XDR *A. baumannii* associated with ST2 (CC2/IC2) [49–53]. Co-occurrence of *bla*_{OXA-72} and *armA* was not observed. Although *armA* has been reported in combination with other carbapenemases, including *bla*_{OXA-72}, *bla*_{NDM-1}, and *bla*_{OXA-58}, these associations remain rare compared with the *armA*–*bla*_{OXA-23-like} combination [54].

APH(3′)-VIa is a highly prevalent aminoglycoside-modifying enzyme, conferring resistance to amikacin and kanamycin. In the present study, whole-genome sequencing confirmed the presence of the encoding gene, *aph*(3′)-VIa, in *armA*-negative isolates, which may explain the persistently elevated amikacin MICs despite susceptibility to gentamicin and tobramycin observed in this group. A study from Iran similarly reported a predominance of *aph*(3′)-VIa in sequence groups not carrying *armA* [55].

4.3. *armA* and Minocycline Resistance Association

A substantial revision of the CLSI breakpoints for minocycline was introduced in 2025. In the SENTRY study, the susceptibility rate of XDR *A. baumannii* isolates to minocycline was 72.7% according to the former breakpoints; re-evaluation using the updated CLSI criteria reduces this estimate to approximately 40% [56]. The impact of breakpoint revision is further illustrated in other studies [57,58].

To the best of our knowledge, this is the first study in Bulgaria reporting minocycline susceptibility data for *A. baumannii*. We observed resistance in half of our isolates, using CLSI M100, 35th edition (2025) [17] breakpoints, with a further 27.8% in the intermediate range.

Minocycline resistance in *A. baumannii* is primarily mediated by the narrow-spectrum efflux pump TetB, a member of the major facilitator superfamily. In our collection, all *tet*(B)-positive isolates exhibited high MIC values (≥ 8 mg/L), consistent with previous reports identifying *tet*(B) as a strong predictor of minocycline resistance [58]. Sequence analysis revealed that all *armA*-positive isolates also carried *tet*(B) and exhibited high MICs (8–24 mg/L), whereas *armA*-negative isolates uniformly lacked *tet*(B) and fell within the intermediate or susceptible range.

Applying the revised 2025 CLSI minocycline breakpoints to our isolates, the susceptibility rate fell from 50.0% to 22.2%: 10 isolates (27.8%) previously classified as susceptible were reclassified as intermediate, and one previously intermediate isolate became resistant, yielding 18 resistant (50.0%), 10 intermediate (27.8%), and 8 susceptible (22.2%). Because inappropriate initial therapy is an independent predictor of mortality in XDR *A. baumannii*, interpretation under the former breakpoints could have led to falsely reassuring susceptibility reports and inappropriate use of minocycline, with an attendant risk of treatment failure and poor outcome. Based on the aforementioned findings, *armA* may serve as a reliable surrogate marker for minocycline resistance in this hospital setting, given the clear delineation of the resistance category. Furthermore, the identification of genotypes 4 and 5, which carry the *bla*_{OXA-72} gene, may predict true susceptibility to the agent. However, this association reflects the clonal structure of the local *A. baumannii* population and may not be generalizable to other settings.

4.4. Determinants of Trimethoprim/Sulfamethoxazole and Fluoroquinolone Resistance

The *sul1* gene, part of a class 1 integron, confers resistance to the sulfonamide component of trimethoprim/sulfamethoxazole [59]. The MIC variability among our trimethoprim/sulfamethoxazole-resistant isolates may be explained by additional mechanisms, including active efflux by RND- and MFS-type pumps, as well as multiple copies of *sul1* [60–62]. The absence of all investigated resistance genes in one isolate corresponded to full susceptibility to trimethoprim/sulfamethoxazole.

Both *bla*_{OXA-23-like}/*armA*-positive and *bla*_{OXA-24/40-like}-positive isolates harbored dual QRDR mutations in *gyrA* and *parC*. However, the former exhibited markedly higher levofloxacin MICs. This discordance suggests the involvement of additional resistance mechanisms that may differ between the two clonal lineages, ST2 and ST636 [63].

4.5. Cefiderocol Susceptibility in a Subset of Nine Isolates

Five of the nine isolates tested exceeded the ECOFF (0.5 mg/L) and were therefore classified as non-wild-type, indicating reduced cefiderocol activity relative to the wild-type population. All five were *armA*-negative. Notably, only two of them (MIC 4 mg/L) exceeded the clinical resistance breakpoint ($R > 2$ mg/L), meaning the remaining three would be reported as susceptible despite already showing a non-wild-type phenotype, an early signal of emerging resistance that routine susceptibility reporting would not capture. Given the limited size of the subset, these observations remain preliminary and require confirmation in a larger series. Detailed values are presented in Table S4 (Supplementary Materials).

4.6. Genotypic–Phenotypic Profiles

By correlating detected resistance genes with the MIC values for each isolate, distinct genotypic–phenotypic resistance profiles were established (Tables 3 and 5). The most prevalent genotypic profiles were as follows: GT1 (36.1%) > GT2 (30.6%) > GT3 (13.9%) > GT4 (11.1%), reflecting the composition of the diversity-selected cohort rather than natural clinical prevalence. Overall, *bla*_{OXA-23-like} was present in 80.6% of isolates, the combination of *bla*_{OXA-23-like} with *armA* in 50.0%, and *bla*_{OXA-23-like} with *sul1* in 66.7%.

4.7. Current Treatment Options for Multidrug-Resistant *A. baumannii* Infections

The present study comprised 34/36 XDR *A. baumannii* isolates. XDR *A. baumannii* is recognized as a WHO critical-priority pathogen, associated with high treatment failure rates and mortality ranging from 18% to 89%, depending on the infection source [64]. For this reason, it is worth noting that treatment options for infections caused by XDR strains remain limited. A few agents, mostly used in combination, have been proposed. Sulbactam–durlobactam combined with a carbapenem is now recommended as first-line therapy, whereas high-dose ampicillin–sulbactam represents an alternative combined with a second active agent such as polymyxin B, minocycline, tigecycline, or cefiderocol [65,66]. Colistin, although retaining full in vitro activity in our cohort, should likewise be used in combination rather than as a monotherapy, given the risk of resistance emergence and treatment failure [48]. The revised, more conservative CLSI (2025) [17] minocycline breakpoints support its use with optimized, PK/PD-guided dosing in combination regimens, thereby reducing the risk of treatment failure [67,68]. Given the severity of XDR *A. baumannii* infections and the need for individualized treatment regimens, the genotype–phenotype associations proposed in this study, combining *bla*_{OXA-23-like}, *bla*_{OXA-24/40-like}, and *armA* as markers for resistance to aminoglycosides and minocycline, may contribute to a better understanding of local epidemiology and support treatment decision-making within a hospital setting.

4.8. Practical Recommendations

Based on our findings, the following practical recommendations can be proposed. In settings with a high prevalence of *bla*_{OXA-23-like}/*armA* co-carriage, empirical avoidance of minocycline may be warranted. Given colistin susceptibility in this cohort, colistin-based combination regimens remain a viable empirical option for XDR *A. baumannii* in this hospital setting. Combined PCR screening for *bla*_{OXA-23-like}, *bla*_{OXA-24/40-like}, and *armA* could serve as a practical tool to guide antibiotic selection in laboratories lacking full genomic sequence capacity.

4.9. Phylogenetic Analysis

Phylogenetic analysis identified three clusters, corresponding entirely to the proposed genotypes. Clusters 1 and 2 comprised genotypes 1–3. All belonged to ST2 and were *bla*_{OXA-23-like}-positive, distinguished by *armA* presence, highlighting the value of *armA* as a sensitive marker for specific resistance phenotypes. The single genotype 6 isolate also clustered here as an ST2 member but was *bla*_{OXA-23-like}-negative and carbapenem-susceptible (discussed below). Cluster 3, positive for *bla*_{OXA-72} and *armA*-negative, belonged to ST636, a triple-locus variant of ST2.

ST2 (CC2/IC2) is the dominant *A. baumannii* lineage worldwide [69,70], strongly linked to carbapenem resistance via *bla*_{OXA-23-like} [69], and typically shows high multidrug/extensive drug resistance beyond carbapenems [49]. Notably, an Italian study of ST2 isolates causing bloodstream infections in COVID-19 patients found uniform resistance not only to carbapenems but also to colistin across all 43 strains [71]. In addition, all but one of the tested ST2 isolates from Ukrainian combat wounds were reported to be cefiderocol-resistant [72]. These two investigations highlight the potential of the clone to acquire a pan-resistant phenotype.

ST636 is mainly reported in Southeast Europe, ranking third among European STs after ST2 and ST1, and has not been reported outside Europe [73]. Furthermore, the clone is uniformly associated with *bla*_{OXA-72} production [37], consistent with our finding. In addition, plasmid localization of *bla*_{OXA-72} within ST636 has been reported [35,49]. In our collection, *bla*_{OXA-72} was confined to ST636 isolates lacking *bla*_{OXA-23-like}, whereas *bla*_{OXA-23-like} was restricted to ST2, with no co-occurrence of the two determinants. This contrasts with reports of *bla*_{OXA-23-like}/*bla*_{OXA-72} co-occurrence within a single clone, described both in ST2 in neighboring Romania [41] and in an ST636 isolate from Varna, Bulgaria [49]. Our ST636/*bla*_{OXA-72} profile more closely resembles the OXA-72-producing ST636 lineage reported in Serbia, both in a neonatal ICU outbreak [74] and among nationally circulating isolates [75]. However, the isolates from the Serbian studies were not subjected to WGS and therefore could not be included in our phylogenomic comparison. This suggests that the ST636/*bla*_{OXA-72} combination has arisen independently in more than one genetic background across the region, rather than spreading from a single common ancestor. ST2 and ST636 co-circulate across the Balkans, sometimes within the same hospital [41,49,75], confirmed by the results of our study.

The close phylogenetic relation of our cluster 1 isolates to a reference genome from Athens, Greece, suggests regional spread of the resistant ST2 clone across the Balkans. Isolates 111 and 112, genetically identical, were collected from the abdominal surgery department (November 2024) and ICU1 (December 2024), respectively, suggesting strain transfer between these territorially close, patient-exchanging departments.

One of the two carbapenem-susceptible isolates, 15 (cluster 2, genotype 6), may be a variant or defective strain within the cluster, while the other, 120 (genotype 7), belongs to no cluster. The significance of such strains remains unclear within this study and requires investigation of a greater number of carbapenem-susceptible isolates.

4.10. Limitations of the Study

This study has several limitations. First, the small number of isolates, 36 with only 17 sequenced, limits the statistical power of the gene–phenotype associations. Second, isolate selection prioritized phenotypic diversity over random sampling, so the reported genotype frequencies (e.g., GT1–GT4) reflect this diversity-enriched subset rather than true prevalence among bloodstream *A. baumannii* isolates. Third, this was a single-center, retrospective study. Consequently, certain information regarding the clinical course of the disease and the precise duration of antibiotic exposure prior to isolate collection could not be reliably reconstructed from medical records. In addition, only bloodstream isolates were included, which may limit generalizability, although bloodstream infections represent the most life-threatening manifestation of *A. baumannii* resistance. Finally, the limited number of isolates tested for cefiderocol could not provide conclusive information regarding the susceptibility of isolates in our hospital setting to the agent. Larger, multicenter studies are needed to confirm and extend these findings.

5. Conclusions

At present, whole-genome sequencing remains predominantly a research tool, with limited implementation in routine clinical microbiology. Routine laboratory practice requires practical and accessible approaches that can inform individualized treatment decisions and facilitate the identification of clonal lineages during outbreak investigations and surveillance of endemic strains. The seven genotypic–phenotypic profiles proposed in this study, together with the role of *armA* as a potential marker for resistance phenotypes, including not only gentamicin and tobramycin but also minocycline, represent a new contribution to the existing data on *A. baumannii* clinical isolates in Bulgaria. Combined with susceptibility testing, they may serve as a practical tool for both clinical decision-making and local epidemiological practice. It is worth mentioning that cefiderocol, as one of the few treatment options for XDR *A. baumannii* infection, requires thorough investigations with clinical isolates. Future investigations addressing pathogenic potential, such as virulence factors, biofilm formation, and closer correlation with clinical data, may further expand the knowledge and practical application of these findings.

Supplementary Materials: The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/microorganisms14081618/s1>: Table S1: Distribution of selected vs. non-selected Acb bloodstream isolates; Table S2: Demographic and clinical characteristics of the patients, MIC values, and detected resistance genes for the 36 *A. baumannii* isolates; Table S3: Statistical Analyses of Associations Between Resistance Genotypes and Antimicrobial Susceptibility Phenotypes; Table S4: Cefiderocol susceptibility results and *armA* status among the nine tested isolates.

Author Contributions: Conceptualization, T.M.-B. and H.H.; methodology, T.M.-B.; software, S.P.; validation, P.M., V.N., and S.P.; formal analysis, P.M. and V.N.; investigation, T.M.-B., H.H., P.M., P.H., V.N., S.P., and N.B.; resources, T.M.-B.; data curation, T.M.-B. and H.H.; writing—original draft preparation, T.M.-B.; writing—review and editing, H.H., S.P., and N.B.; visualization, T.M.-B. and H.H.; supervision, H.H. and S.P.; project administration, T.M.-B.; funding acquisition, T.M.-B. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by the Medical University of Pleven under Intramural Research Project No. 7/2024.

Institutional Review Board Statement: This study was approved by the Ethics Committee of the Medical University of Pleven (approval no. 794/14 June 2024).

Informed Consent Statement: Patient consent was waived because the study used only bacterial isolates and no identifiable personal data. All patients had signed the standard hospital admission consent form; no study-specific informed consent was required.

Data Availability Statement: The whole-genome shotgun sequencing project of 17 isolates of *A. baumannii* is deposited in GenBank under BioProject accession number PRJNA1479364.

Acknowledgments: The authors would like to thank the Medical University of Pleven for providing administrative, financial, and technical support. The authors also gratefully acknowledge the National Centre of Infectious and Parasitic Diseases, Sofia, for providing methodological guidance and reference strains. The authors declare that an artificial intelligence tool (OpenEvidence, Inc., Miami, FL, USA; <https://www.openevidence.com>, accessed on 30 May 2026) was used during manuscript preparation for text formatting and literature search assistance. The authors take full responsibility for the accuracy and integrity of the final manuscript content.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Peleg, A.Y.; Seifert, H.; Paterson, D.L. *Acinetobacter baumannii*: Emergence of a Successful Pathogen. *Clin. Microbiol. Rev.* **2008**, *21*, 538–582. [CrossRef] [PubMed]
2. Müller, C.; Reuter, S.; Wille, J.; Xanthopoulou, K.; Stefanik, D.; Grundmann, H.; Higgins, P.G.; Seifert, H. A Global View on Carbapenem-Resistant *Acinetobacter baumannii*. *mBio* **2023**, *14*, e02260. [CrossRef] [PubMed]
3. Miller, W.R.; Arias, C.A. ESKAPE Pathogens: Antimicrobial Resistance, Epidemiology, Clinical Impact and Therapeutics. *Nat. Rev. Microbiol.* **2024**, *22*, 598–616. [CrossRef] [PubMed]
4. Surveillance Atlas of Infectious Diseases. Available online: <https://atlas.ecdc.europa.eu/public/index.aspx> (accessed on 14 July 2026).
5. McGowan, J.E. Resistance in Nonfermenting Gram-Negative Bacteria: Multidrug Resistance to the Maximum. *Am. J. Infect. Control* **2006**, *34*, S29–S37. [CrossRef] [PubMed]
6. Breijyeh, Z.; Jubeh, B.; Karaman, R. Resistance of Gram-Negative Bacteria to Current Antibacterial Agents and Approaches to Resolve It. *Molecules* **2020**, *25*, 1340. [CrossRef] [PubMed]
7. Kyriakidis, I.; Vasileiou, E.; Pana, Z.D.; Tragiannidis, A. *Acinetobacter baumannii* Antibiotic Resistance Mechanisms. *Pathogens* **2021**, *10*, 373. [CrossRef] [PubMed]
8. Frenk, S.; Temkin, E.; Lurie-Weinberger, M.N.; Keren-Paz, A.; Rov, R.; Rakovitsky, N.; Wullfhart, L.; Nutman, A.; Daikos, G.L.; Skiada, A.; et al. Large-Scale WGS of Carbapenem-Resistant *Acinetobacter baumannii* Isolates Reveals Patterns of Dissemination of ST Clades Associated with Antibiotic Resistance. *J. Antimicrob. Chemother.* **2022**, *77*, 934–943. [CrossRef] [PubMed]
9. Fournier, P.-E.; Vallenet, D.; Barbe, V.; Audic, S.; Ogata, H.; Poirol, L.; Richet, H.; Robert, C.; Mangenot, S.; Abergel, C.; et al. Comparative Genomics of Multidrug Resistance in *Acinetobacter baumannii*. *PLoS Genet.* **2006**, *2*, e7. [CrossRef] [PubMed]
10. Zhou, H.; Zhang, T.; Yu, D.; Pi, B.; Yang, Q.; Zhou, J.; Hu, S.; Yu, Y. Genomic Analysis of the Multidrug-Resistant *Acinetobacter baumannii* Strain MDR-ZJ06 Widely Spread in China. *Antimicrob. Agents Chemother.* **2011**, *55*, 4506–4512. [CrossRef] [PubMed]
11. Zhu, L.; Yan, Z.; Zhang, Z.; Zhou, Q.; Zhou, J.; Wakeland, E.K.; Fang, X.; Xuan, Z.; Shen, D.; Li, Q.-Z. Complete Genome Analysis of Three *Acinetobacter baumannii* Clinical Isolates in China for Insight into the Diversification of Drug Resistance Elements. *PLoS ONE* **2013**, *8*, e66584. [CrossRef] [PubMed]
12. Ou, H.-Y.; Kuang, S.N.; He, X.; Molgora, B.M.; Ewing, P.J.; Deng, Z.; Osby, M.; Chen, W.; Xu, H.H. Complete Genome Sequence of Hypervirulent and Outbreak-Associated *Acinetobacter baumannii* Strain LAC-4: Epidemiology, Resistance Genetic Determinants and Potential Virulence Factors. *Sci. Rep.* **2015**, *5*, 8643. [CrossRef] [PubMed]
13. Karah, N.; Dwibedi, C.K.; Sjöström, K.; Edquist, P.; Johansson, A.; Wai, S.N.; Uhlin, B.E. Novel Aminoglycoside Resistance Transposons and Transposon-Derived Circular Forms Detected in Carbapenem-Resistant *Acinetobacter baumannii* Clinical Isolates. *Antimicrob. Agents Chemother.* **2016**, *60*, 1801–1818. [CrossRef] [PubMed]
14. Holt, K.; Kenyon, J.J.; Hamidian, M.; Schultz, M.B.; Pickard, D.J.; Dougan, G.; Hall, R. Five Decades of Genome Evolution in the Globally Distributed, Extensively Antibiotic-Resistant *Acinetobacter baumannii* Global Clone 1. *Microb. Genom.* **2016**, *2*, e000052, Erratum in *Microb. Genom.* **2019**, *5*, e000280. [CrossRef] [PubMed]
15. Turton, J.F.; Woodford, N.; Glover, J.; Yarde, S.; Kaufmann, M.E.; Pitt, T.L. Identification of *Acinetobacter baumannii* by Detection of the blaOXA-51-like Carbapenemase Gene Intrinsic to This Species. *J. Clin. Microbiol.* **2006**, *44*, 2974–2976. [CrossRef] [PubMed]
16. Higgins, P.G.; Lehmann, M.; Wisplinghoff, H.; Seifert, H. *gyrB* Multiplex PCR to Differentiate between *Acinetobacter calcoaceticus* and *Acinetobacter* Genomic Species 3. *J. Clin. Microbiol.* **2010**, *48*, 4592–4594. [CrossRef] [PubMed]

17. M100; Performance Standards for Antimicrobial Susceptibility Testing. Clinical and Laboratory Standards Institute: Wayne, PA, USA, 2025.
18. Magiorakos, A.-P.; Srinivasan, A.; Carey, R.B.; Carmeli, Y.; Falagas, M.E.; Giske, C.G.; Harbarth, S.; Hindler, J.F.; Kahlmeter, G.; Olsson-Liljequist, B.; et al. Multidrug-Resistant, Extensively Drug-Resistant and Pandrug-Resistant Bacteria: An International Expert Proposal for Interim Standard Definitions for Acquired Resistance. *Clin. Microbiol. Infect.* **2012**, *18*, 268–281. [\[CrossRef\]](#) [\[PubMed\]](#)
19. Woodford, N.; Ellington, M.J.; Coelho, J.M.; Turton, J.F.; Ward, M.E.; Brown, S.; Amyes, S.G.B.; Livermore, D.M. Multiplex PCR for Genes Encoding Prevalent OXA Carbapenemases in *Acinetobacter Spp.* *Int. J. Antimicrob. Agents* **2006**, *27*, 351–353. [\[CrossRef\]](#)
20. Goudarzi, H.; Mirsamadi, E.S.; Ghalavand, Z.; Hakemi Vala, M.; Mirjalali, H.; Hashemi, A. Rapid Detection and Molecular Survey of blaVIM, blaIMP and blaNDM Genes among Clinical Isolates of *Acinetobacter baumannii* Using New Multiplex Real-Time PCR and Melting Curve Analysis. *BMC Microbiol.* **2019**, *19*, 122. [\[CrossRef\]](#) [\[PubMed\]](#)
21. Poirel, L.; Walsh, T.R.; Cuvillier, V.; Nordmann, P. Multiplex PCR for Detection of Acquired Carbapenemase Genes. *Diagn. Microbiol. Infect. Dis.* **2011**, *70*, 119–123. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Cole, J.M.; Schuetz, A.N.; Hill, C.E.; Nolte, F.S. Development and Evaluation of a Real-Time PCR Assay for Detection of Klebsiella Pneumoniae Carbapenemase Genes. *J. Clin. Microbiol.* **2009**, *47*, 322–326. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Yamane, K.; Wachino, J.; Suzuki, S.; Shibata, N.; Kato, H.; Shibayama, K.; Kimura, K.; Kai, K.; Ishikawa, S.; Ozawa, Y.; et al. 16S rRNA Methylase-Producing, Gram-Negative Pathogens, Japan. *Emerg. Infect. Dis.* **2007**, *13*, 642–646. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Barbolla, R.; Catalano, M.; Orman, B.E.; Famiglietti, A.; Vay, C.; Smayevsky, J.; Centrón, D.; Piñeiro, S.A. Class 1 Integrins Increase Trimethoprim-Sulfamethoxazole MICs against Epidemiologically Unrelated *Stenotrophomonas maltophilia* Isolates. *Antimicrob. Agents Chemother.* **2004**, *48*, 666–669. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Bolger, A.M.; Lohse, M.; Usadel, B. Trimmomatic: A Flexible Trimmer for Illumina Sequence Data. *Bioinformatics* **2014**, *30*, 2114–2120. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Bankevich, A.; Nurk, S.; Antipov, D.; Gurevich, A.A.; Dvorkin, M.; Kulikov, A.S.; Lesin, V.M.; Nikolenko, S.I.; Pham, S.; Pribelski, A.D.; et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J. Comput. Biol.* **2012**, *19*, 455–477. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Gurevich, A.; Saveliev, V.; Vyahhi, N.; Tesler, G. QUAST: Quality Assessment Tool for Genome Assemblies. *Bioinformatics* **2013**, *29*, 1072–1075. [\[CrossRef\]](#) [\[PubMed\]](#)
28. Seemann, T. Prokka: Rapid Prokaryotic Genome Annotation. *Bioinformatics* **2014**, *30*, 2068–2069. [\[CrossRef\]](#) [\[PubMed\]](#)
29. Page, A.J.; Cummins, C.A.; Hunt, M.; Wong, V.K.; Reuter, S.; Holden, M.T.G.; Fookes, M.; Falush, D.; Keane, J.A.; Parkhill, J. Roary: Rapid Large-Scale Prokaryote Pan Genome Analysis. *Bioinformatics* **2015**, *31*, 3691–3693. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Kozlov, A.M.; Darriba, D.; Flouri, T.; Morel, B.; Stamatakis, A. RAXML-NG: A Fast, Scalable and User-Friendly Tool for Maximum Likelihood Phylogenetic Inference. *Bioinformatics* **2019**, *35*, 4453–4455. [\[CrossRef\]](#) [\[PubMed\]](#)
31. Letunic, I.; Bork, P. Interactive Tree of Life (iTOL) v6: Recent Updates to the Phylogenetic Tree Display and Annotation Tool. *Nucleic Acids Res.* **2024**, *52*, W78–W82. [\[CrossRef\]](#) [\[PubMed\]](#)
32. Jamovi—Open Statistical Software. Available online: <https://www.jamovi.org/index.html> (accessed on 19 July 2026).
33. The Comprehensive R Archive Network. Available online: <https://cran.r-project.org/> (accessed on 19 July 2026).
34. Park, S.; Lee, K.M.; Yoo, Y.S.; Yoo, J.S.; Yoo, J.I.; Kim, H.S.; Lee, Y.S.; Chung, G.T. Alterations of gyrA, gyrB, and parC and Activity of Efflux Pump in Fluoroquinolone-Resistant *Acinetobacter baumannii*. *Osong Public. Health Res. Perspect.* **2011**, *2*, 164–170. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Valcek, A.; Nesporova, K.; Whiteway, C.; De Pooter, T.; De Coster, W.; Strazisar, M.; Van der Henst, C. Genomic Analysis of a Strain Collection Containing Multidrug-, Extensively Drug-, Pandrug-, and Carbapenem-Resistant Modern Clinical Isolates of *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2022**, *66*, e0089222. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Shelenkov, A.; Akimkin, V.; Mikhaylova, Y. International Clones of High Risk of *Acinetobacter baumannii*-Definitions, History, Properties and Perspectives. *Microorganisms* **2023**, *11*, 2115. [\[CrossRef\]](#) [\[PubMed\]](#)
37. Hummel, D.; Juhasz, J.; Kamotsay, K.; Kristof, K.; Xavier, B.B.; Koster, S.D.; Szabo, D.; Kocsis, B. Genomic Investigation and Comparative Analysis of European High-Risk Clone of *Acinetobacter baumannii* ST2. *Microorganisms* **2024**, *12*, 2474. [\[CrossRef\]](#) [\[PubMed\]](#)
38. Stoeva, T.; Higgins, P.G.; Bojkova, K.; Seifert, H. Clonal Spread of Carbapenem-Resistant OXA-23-Positive *Acinetobacter baumannii* in a Bulgarian University Hospital. *Clin. Microbiol. Infect.* **2008**, *14*, 723–727. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Stoeva, T.J. Microbiological and Molecular-Genetic Investigations on the Epidemiology and Antimicrobial Resistance of Clinical *A. baumannii* Isolates. Doctoral Thesis, Medical University of Varna, Varna, Bulgaria, 2009. (In Bulgarian)
40. Strateva, T.V. Antibiotic Resistance, Virulence and Epidemiology of Therapeutically Problematic Isolates from the Group of “Non-Glucose-Fermenting Gram-Negative Bacteria” (2004–2023). Habilitation Thesis, Medical University of Sofia, Sofia, Bulgaria, 2023. (In Bulgarian)

41. Gheorghe-Barbu, I.; Surleac, M.; Barbu, I.C.; Paraschiv, S.; Bănică, L.M.; Rotaru, L.-I.; Vrâncianu, C.O.; Niță Lazăr, M.; Oțelea, D.; Chifiriuc, M.C. Decoding the Resistome, Virulome and Mobilome of Clinical versus Aquatic *Acinetobacter baumannii* in Southern Romania. *Heliyon* **2024**, *10*, e33372. [[CrossRef](#)] [[PubMed](#)]
42. Goic-Barisic, I.; Kovacic, A.; Medic, D.; Jakovac, S.; Petrovic, T.; Tonkic, M.; Novak, A.; Rubic, Z.; Radic, M.; Milosavljević, B.; et al. Endemicity of OXA-23 and OXA-72 in Clinical Isolates of *Acinetobacter baumannii* from Three Neighbouring Countries in Southeast Europe. *J. Appl. Genet.* **2021**, *62*, 353–359. [[CrossRef](#)] [[PubMed](#)]
43. Papadopoulou, M.; Deliolanis, I.; Polemis, M.; Vatopoulos, A.; Psychogiou, M.; Giakkoupi, P. Characteristics of the Genetic Spread of Carbapenem-Resistant *Acinetobacter baumannii* in a Tertiary Greek Hospital. *Genes* **2024**, *15*, 458. [[CrossRef](#)] [[PubMed](#)]
44. Pournaras, S.; Dafopoulou, K.; Del Franco, M.; Zarkotou, O.; Dimitroulia, E.; Protonotariou, E.; Poulou, A.; Zarrilli, R.; Tsakris, A.; Greek Study Group on Acinetobacter Antimicrobial Resistance. Predominance of International Clone 2 OXA-23-Producing *Acinetobacter baumannii* Clinical Isolates in Greece, 2015: Results of a Nationwide Study. *Int. J. Antimicrob. Agents* **2017**, *49*, 749–753. [[CrossRef](#)] [[PubMed](#)]
45. Mugnier, P.D.; Poirel, L.; Naas, T.; Nordmann, P. Worldwide Dissemination of the bla_{OXA-23} Carbapenemase Gene of *Acinetobacter baumannii*1. *Emerg. Infect. Dis.* **2010**, *16*, 35–40. [[CrossRef](#)] [[PubMed](#)]
46. Metan, G.; Sariguzel, F.; Sumerkan, B.; van der Reijden, T.; Dijkshoorn, L. Clonal Diversity and High Prevalence of OXA-58 among *Acinetobacter baumannii* Isolates from Blood Cultures in a Tertiary Care Centre in Turkey. *Infect. Genet. Evol.* **2013**, *14*, 92–97, Correction in *Infect. Genet. Evol.* **2013**, *16*, 447–448. [[CrossRef](#)] [[PubMed](#)]
47. Ahmed, S.S.; Alp, E.; Ulu-Kilic, A.; Dinc, G.; Aktas, Z.; Ada, B.; Bagirova, F.; Baran, I.; Ersoy, Y.; Esen, S.; et al. Spread of Carbapenem-Resistant International Clones of *Acinetobacter baumannii* in Turkey and Azerbaijan: A Collaborative Study. *Eur. J. Clin. Microbiol. Infect. Dis.* **2016**, *35*, 1463–1468. [[CrossRef](#)] [[PubMed](#)]
48. Karampatakis, T.; Tsergouli, K.; Behzadi, P. Carbapenem-Resistant *Acinetobacter baumannii*: Virulence Factors, Molecular Epidemiology, and Latest Updates in Treatment Options. *Microorganisms* **2025**, *13*, 1983. [[CrossRef](#)] [[PubMed](#)]
49. Strateva, T.V.; Sirakov, I.; Stoeva, T.J.; Stratev, A.; Peykov, S. Phenotypic and Molecular Characteristics of Carbapenem-Resistant *Acinetobacter baumannii* Isolates from Bulgarian Intensive Care Unit Patients. *Microorganisms* **2023**, *11*, 875. [[CrossRef](#)] [[PubMed](#)]
50. Hong, S.B.; Shin, K.S.; Ha, J.; Han, K. Co-Existence of bla_{OXA-23} and armA in Multidrug-Resistant *Acinetobacter baumannii* Isolated from a Hospital in South Korea. *J. Med. Microbiol.* **2013**, *62*, 836–844. [[CrossRef](#)] [[PubMed](#)]
51. Lim, J.; Cho, H.H.; Kim, S.; Kim, J.; Kwon, K.C.; Park, J.W.; Koo, S.H. The Genetic Characteristics of Multidrug-resistant *Acinetobacter baumannii* Coproducing 16S rRNA Methylase armA and Carbapenemase OXA-23. *J. Bacteriol. Virol.* **2013**, *43*, 27–36. [[CrossRef](#)]
52. Palmieri, M.; D'Andrea, M.M.; Pelegrin, A.C.; Perrot, N.; Mirande, C.; Blanc, B.; Legakis, N.; Goossens, H.; Rossolini, G.M.; van Belkum, A. Abundance of Colistin-Resistant, OXA-23- and ArmA-Producing *Acinetobacter baumannii* Belonging to International Clone 2 in Greece. *Front. Microbiol.* **2020**, *11*, 668. [[CrossRef](#)] [[PubMed](#)]
53. Findlay, J.; Nordmann, P.; Bouvier, M.; Kerbol, A.; Poirel, L. Dissemination of ArmA- and OXA-23-Co-Producing *Acinetobacter baumannii* Global Clone 2 in Switzerland, 2020–2021. *Eur. J. Clin. Microbiol. Infect. Dis.* **2023**, 1–5. [[CrossRef](#)] [[PubMed](#)]
54. Tada, T.; Miyoshi-Akiyama, T.; Shimada, K.; Shimojima, M.; Kirikae, T. Dissemination of 16S rRNA Methylase ArmA-Producing *Acinetobacter baumannii* and Emergence of OXA-72 Carbapenemase Coproducers in Japan. *Antimicrob. Agents Chemother.* **2014**, *58*, 2916–2920. [[CrossRef](#)] [[PubMed](#)]
55. Hasani, A.; Sheikhalizadeh, V.; Ahangarzadeh Rezaee, M.; Rahmati-Yamchi, M.; Hasani, A.; Ghotaslou, R.; Goli, H.R. Frequency of Aminoglycoside-Modifying Enzymes and armA Among Different Sequence Groups of *Acinetobacter baumannii* in Iran. *Microb. Drug Resist.* **2016**, *22*, 347–353. [[CrossRef](#)] [[PubMed](#)]
56. Gales, A.C.; Seifert, H.; Gur, D.; Castanheira, M.; Jones, R.N.; Sader, H.S. Antimicrobial Susceptibility of *Acinetobacter calcoaceticus*-*Acinetobacter baumannii* Complex and *Stenotrophomonas maltophilia* Clinical Isolates: Results from the SENTRY Antimicrobial Surveillance Program (1997–2016). *Open Forum Infect. Dis.* **2019**, *6*, S34–S46. [[CrossRef](#)] [[PubMed](#)]
57. Hsueh, S.-C.; Huang, Y.-T.; Ko, W.-C.; Liu, I.-M.; Hsieh, P.-C.; Jean, S.-S. In Vitro Antimicrobial Susceptibility Data of Global Meropenem-Resistant *Acinetobacter baumannii* Isolates Causing Pneumonia: Data from the Antimicrobial Testing Leadership and Surveillance Program, 2014–2021, and Re-Estimations of Susceptibility Breakpoints and Appropriate Dosages of Important Antibiotics for Pneumonia Treatment. *J. Glob. Antimicrob. Resist.* **2024**, *36*, 411–418. [[CrossRef](#)] [[PubMed](#)]
58. Yu, X.; Liu, Y.; Du, J.; Hu, F.; Yin, D. Assessment of the Revision of the 2025 CLSI Breakpoints for the Interpretation of Minocycline Susceptibility for *Acinetobacter baumannii* Complex. *Microbiol. Spectr.* **2026**, *14*, e02700-25. [[CrossRef](#)] [[PubMed](#)]
59. Bala, A.; Uhlin, B.E.; Karah, N. Insights into the Genetic Contexts of Sulfonamide Resistance among Early Clinical Isolates of *Acinetobacter baumannii*. *Infect. Genet. Evol.* **2023**, *112*, 105444. [[CrossRef](#)] [[PubMed](#)]
60. Venkatesan, M.; Fruci, M.; Verellen, L.A.; Skarina, T.; Mesa, N.; Flick, R.; Pham, C.; Mahadevan, R.; Stogios, P.J.; Savchenko, A. Molecular Mechanism of Plasmid-Borne Resistance to Sulfonamide Antibiotics. *Nat. Commun.* **2023**, *14*, 4031. [[CrossRef](#)] [[PubMed](#)]

61. Nowak, J.; Schneiders, T.; Seifert, H.; Higgins, P.G. The Asp20-to-Asn Substitution in the Response Regulator AdeR Leads to Enhanced Efflux Activity of AdeB in *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2016**, *60*, 1085–1090. [[CrossRef](#)] [[PubMed](#)]
62. Gaona, M.; Corral, J.; Campoy, S.; Barbé, J.; Pérez–Varela, M.; Aranda, J. The Novel MFS Efflux Pump SxtP, Regulated by the LysR-Type Transcriptional Activator SxtR, Is Involved in the Susceptibility to Sulfamethoxazole/Trimethoprim (SXT) and the Pathogenesis of *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2024**, *68*, e00712–24. [[CrossRef](#)] [[PubMed](#)]
63. Roy, S.; Chatterjee, S.; Bhattacharjee, A.; Chattopadhyay, P.; Saha, B.; Dutta, S.; Basu, S. Overexpression of Efflux Pumps, Mutations in the Pumps' Regulators, Chromosomal Mutations, and AAC(6')-Ib-Cr Are Associated With Fluoroquinolone Resistance in Diverse Sequence Types of Neonatal Septicaemic *Acinetobacter baumannii*: A 7-Year Single Center Study. *Front. Microbiol.* **2021**, *12*, 602724. [[CrossRef](#)] [[PubMed](#)]
64. Dubey, V.; Reza, N.; Hope, W. Drug-Resistant *Acinetobacter baumannii*: Mortality, Emerging Treatments, and Future Pharmacological Targets for a WHO Priority Pathogen. *Clin. Microbiol. Rev.* **2025**, *38*, e00279–24. [[CrossRef](#)] [[PubMed](#)]
65. Paul, M.; Carrara, E.; Retamar, P.; Tängdén, T.; Bitterman, R.; Bonomo, R.A.; de Waele, J.; Daikos, G.L.; Akova, M.; Harbarth, S.; et al. European Society of Clinical Microbiology and Infectious Diseases (ESCMID) Guidelines for the Treatment of Infections Caused by Multidrug-Resistant Gram-Negative Bacilli (Endorsed by European Society of Intensive Care Medicine). *Clin. Microbiol. Infect.* **2022**, *28*, 521–547. [[CrossRef](#)] [[PubMed](#)]
66. Tamma, P.D.; Aitken, S.L.; Bonomo, R.A.; Mathers, A.J.; van Duin, D.; Clancy, C.J. Infectious Diseases Society of America 2023 Guidance on the Treatment of Antimicrobial Resistant Gram-Negative Infections. *Clin. Infect. Dis.* **2023**, ciad428. [[CrossRef](#)] [[PubMed](#)]
67. Kim, T.; Jeon, E.H.; Hong, Y.-K.; Jung, J.; Kim, M.J.; Sung, H.; Kim, M.-N.; Kim, S.-H.; Choi, S.-H.; Lee, S.-O.; et al. Minocycline Susceptibility of Carbapenem-Resistant *Acinetobacter baumannii* Blood Isolates from a Single Center in Korea: Role of tetB in Resistance. *Infect. Chemother.* **2025**, *57*, 111–118. [[CrossRef](#)] [[PubMed](#)]
68. Shields, R.K.; Paterson, D.L.; Tamma, P.D. Navigating Available Treatment Options for Carbapenem-Resistant *Acinetobacter baumannii-calcoaceticus* Complex Infections. *Clin. Infect. Dis.* **2023**, *76*, S179–S193. [[CrossRef](#)] [[PubMed](#)]
69. Hamidian, M.; Nigro, S.J. Emergence, Molecular Mechanisms and Global Spread of Carbapenem-Resistant *Acinetobacter baumannii*. *Microb. Genom.* **2019**, *5*, e000306. [[CrossRef](#)] [[PubMed](#)]
70. Boutzoukas, A.; Doi, Y. The Global Epidemiology of Carbapenem-Resistant *Acinetobacter baumannii*. *JAC Antimicrob. Resist.* **2025**, *7*, dlaf134. [[CrossRef](#)] [[PubMed](#)]
71. Cherubini, S.; Perilli, M.; Segatore, B.; Fazii, P.; Parruti, G.; Frattari, A.; Amicosante, G.; Piccirilli, A. Whole-Genome Sequencing of ST2 *A. baumannii* Causing Bloodstream Infections in COVID-19 Patients. *Antibiotics* **2022**, *11*, 955. [[CrossRef](#)] [[PubMed](#)]
72. Kondratiuk, V.M.; Jones, B.T.; Luo, T.L.; Fomina, N.S.; Lebreton, F.; Bennett, J.W.; Mc Gann, P.; Kovalchuk, V.P. Phenotypic and Genotypic Analysis of *Acinetobacter baumannii* Isolated from Combat Wounds in Ukraine during 2022 and 2023. *JAC Antimicrob Resist.* **2025**, *7*, dlaf140. [[CrossRef](#)] [[PubMed](#)]
73. Pearl, S.; Anbarasu, A. Genomic Landscape of Nosocomial *Acinetobacter baumannii*: A Comprehensive Analysis of the Resistome, Virulome, and Mobilome. *Sci. Rep.* **2025**, *15*, 18203. [[CrossRef](#)] [[PubMed](#)]
74. Gajic, I.; Jovicevic, M.; Milic, M.; Kekic, D.; Opavski, N.; Zrnica, Z.; Dacic, S.; Pavlovic, L.; Mijac, V. Clinical and Molecular Characteristics of OXA-72-Producing *Acinetobacter baumannii* ST636 Outbreak at a Neonatal Intensive Care Unit in Serbia. *J. Hosp. Infect.* **2021**, *112*, 54–60. [[CrossRef](#)] [[PubMed](#)]
75. Lukovic, B.; Gajic, I.; Dimkic, I.; Kekic, D.; Zornic, S.; Pozder, T.; Radisavljevic, S.; Opavski, N.; Kojic, M.; Ranin, L. The First Nationwide Multicenter Study of *Acinetobacter baumannii* Recovered in Serbia: Emergence of OXA-72, OXA-23 and NDM-1-Producing Isolates. *Antimicrob. Resist. Infect. Control* **2020**, *9*, 101. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.