


Short Note

# 5,9,11-Trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one from *Calophyllum pseudomole*

Mulyadi Tanjung \*, Ratih Dewi Saputri and Tjitjik Srie Tjahjandarie 

Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60115, Indonesia; duffputri@gmail.com (R.D.S.); tjitjiktjahjandarie@fst.unair.ac.id (T.S.T.)

\* Correspondence: mulyadi-t@fst.unair.ac.id; Tel.: +62-31-5936501; Fax: +62-31-5936502


Received: 1 September 2017; Accepted: 27 September 2017; Published: 30 September 2017

**Abstract:** A new pyranocoumarin, namely 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one **1**, was isolated from the stem bark of *Calophyllum pseudomole*. The structure of compound **1** was elucidated based on its ultraviolet (UV); infrared (IR); high resolution electro spray ionization mass spectrometry (HRESIMS); 1D and 2D nuclear magnetic resonance (NMR) spectral data.

**Keywords:** 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one; *Calophyllum pseudomole*; pyranoxanthone

## 1. Introduction

The genus *Calophyllum* belongs to the Clusiaceae family which comprises about 180 species found mainly in Southeast Asia. This plant is endemic to Kalimantan Island, Indonesia. This genus has been shown to possess a number of secondary metabolites such as xanthenes [1–3], coumarins [4,5], and chromanone acids [6,7]. Many of these compounds have shown a wide range of biological and pharmacological properties such as anti-HIV [8,9], anticancer [10,11], and antimalarial properties [12].



**Figure 1.** Structures of 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one.

In continuation of our phytochemical investigation on bioactive xanthone, we wish to report the isolation and structural elucidation of a new pyranoxanthone, 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-

one (Figure 1) from the stem bark of *Calophyllum pseudomole*. The cytotoxicity against murine leukemia P-388 and antiplasmodial activity against *Plasmodium falciparum* of **1** are also briefly described.


## 2. Result and Discussion

5,9,11-Trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one was isolated as a yellow solid, and its UV absorption maxima at  $\lambda_{\max}$  245, 268, and 320 nm which is characteristic of xanthone chromophore [2]. Based on HRESIMS (see data supplementary Figure S1), compound **1** showed deprotonated molecule ion  $[M - H]^-$  at  $m/z$  477.1917 corresponding to the molecular formula  $C_{28}H_{30}O_7$ . The IR spectrum (data supplementary Figure S2) showed absorption for hydroxyl ( $3423\text{ cm}^{-1}$ ), conjugated carbonyl ( $1645\text{ cm}^{-1}$ ), and aromatic ( $1577$  and  $1460\text{ cm}^{-1}$ ) groups [13]. The  $^1\text{H-NMR}$  spectrum (Table 1, data supplementary Figure S3) of **1** showed the presence of a singlet of chelated hydroxyl at  $\delta_{\text{H}}$  13.82 (11-OH), a singlet of hydroxyl at  $\delta_{\text{H}}$  6.14 (5-OH), and two singlets of the isolated aromatic proton at  $\delta_{\text{H}}$  6.81 (H-6) and  $\delta_{\text{H}}$  6.39 (H-8) which is typical for a xanthone with six substituents [1]. The  $^1\text{H-NMR}$  spectrum also revealed the presence of a substituent of a monosubstituted 2,2-dimethylpyrano, 2-methylbut-3-en-2-yl, and 2-hydroxy-3-methylbut-3-en-1-yl group. The presence of a monosubstituted 2,2-dimethylpyrano was determined from resonances of vinyl at  $\delta_{\text{H}}$  8.13 (s, 1H), and two signals of methyl at  $\delta_{\text{H}}$  1.54 (s, 3H) and 1.52 (s, 3H). The  $^1\text{H-NMR}$  spectrum of 2-methylbut-3-en-2-yl showed the presence of a methyl at  $\delta_{\text{H}}$  1.43 (s, 6H), a vinyl at  $\delta_{\text{H}}$  5.98 (dd, 1H), and a methylene at  $\delta_{\text{H}}$  5.13 (d, 1H) and  $\delta_{\text{H}}$  5.08 (d, 1H). The  $^1\text{H-NMR}$  spectrum of 2-methylbut-3-en-2-yl also showed the presence of a methyl at  $\delta_{\text{H}}$  1.87 (s, 3H), two proton signals of methylene at  $\delta_{\text{H}}$  3.16 (dd, 1H) and 2.93 (dd, 1H), a methyne at  $\delta_{\text{H}}$  4.43 (d, 1H), and two proton signals of vinylic methylene at  $\delta_{\text{H}}$  4.97 (s, 1H) and 4.86 (s, 1H).

The  $^{13}\text{C-NMR}$  spectrum (Table 1, Figure S4 data supplementary) of **1** showed 28 carbon signals and their assignments were determined by heteronuclear multiple quantum coherence (HMQC) and heteronuclear multiple bond correlation (HMBC) spectra. The placement of hydroxyl, a monosubstituted 2,2-dimethylpyrano, 2-methylbut-3-en-2-yl, and 2-hydroxy-3-methylbut-3-en-1-yl group was confirmed by HMQC and HMBC spectra (Table 1). Based on the HMBC spectrum, a chelated hydroxyl signal at  $\delta_{\text{H}}$  13.82 (11-OH) was correlated with two quaternary carbons [ $\delta_{\text{C}}$  107.4 (C-10); 103.4 (C-11a)], and an oxyaryl carbon  $\delta_{\text{C}}$  161.1 (C-11) indicated that C-10 has a substituent. Two proton signals of methylene (H-1'') of the 2-hydroxy-3-methylbut-3-en-1-yl group at  $\delta_{\text{H}}$  3.18 and  $\delta_{\text{H}}$  2.95 showed long-range correlations with two oxyaryl carbons [ $\delta_{\text{C}}$  161.1; 163.2 (C-9)], two quaternary carbons [ $\delta_{\text{C}}$  107.4; 146.7 (C-3'')], and a methine carbon of alcohol [ $\delta_{\text{C}}$  77.5 (C-2'')]. The correlations showed that the 2-hydroxy-3-methylbut-3-en-1-yl group attached at C-10. The position of 2-hydroxy-3-methylbut-3-en-1-yl group at C-10 is reinforced by the correlation of the isolated aromatic proton signals at  $\delta_{\text{H}}$  6.39 (H-8) with two oxyaryl carbons [ $\delta_{\text{C}}$  155.8 (C-7a); 163.2 (C-9)], and two quaternary carbons [ $\delta_{\text{C}}$  107.4 (C-10); 103.4 (C-11a)]. Furthermore, the proton signal of isolated aromatic at  $\delta_{\text{H}}$  6.81 (H-6) correlates with two oxyaryl carbons [ $\delta_{\text{C}}$  153.4 (C-5), 150.6 (C-6a)] and a quaternary carbon signal ( $\delta_{\text{C}}$  108.5, C-4a), which showed that 2,2-dimethylpyrano was fused at C-4a and C-12b and the hydroxyl group attached at C-5 from the xanthone skeleton. The position of the hydroxyl group at C-5 is reinforced by the correlation of the hydroxyl proton signal at  $\delta_{\text{H}}$  6.15 (5-OH) with an oxyaryl carbon [ $\delta_{\text{C}}$  153.4 (C-5)], and a methine carbon [ $\delta_{\text{C}}$  101.8 (C-6)]. The methine signal of the vinyl group at  $\delta_{\text{H}}$  8.13 (H-4) of a monosubstituted 2,2-dimethylpyrano group showed long-range correlations with four quaternary carbons [ $\delta_{\text{C}}$  80.6 (C-2), 135.7 (C-3), 108.5 (C-4a), 42.1 (C-2')] which showed that a 2-methylbut-3-en-2-yl group attached at C-3. The HMBC correlation of compound **1** can be seen on Figure 2. Therefore, compound **1** was identified as 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one and a novel compound.

**Table 1.** NMR spectroscopic data of 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one in CDCl<sub>3</sub>.

No.C	$\delta_H$ (Mult, J in Hz)	$\delta_C$	HMBC
2	-	80.6	-
2a	1.54 (s, 3H)	27.4	C-2; C-2b; C-12b
2b	1.53 (s, 3H)	27.4	C-2; C-2a; C-12b
3	-	135.7	-
4	8.13 (s, 1H)	118.0	C-2; C-3; C-4a; C-2'
4a	-	108.5	-
5	-	150.6	-
6	6.81 (s, 1H)	101.8	C-3; C-4a; C-5; C-6a
6a	-	153.4	-
7a	-	155.8	-
8	6.39 (s, 1H)	94.5	C-7a; C-9; C-10; C-11a
9	-	163.2	-
10	-	107.4	-
11	-	161.1	-
11a	-	103.4	-
12	-	182.5	-
12a	-	121.7	-
12b	-	148.8	-
1'	1.43 (s, 3H)	28.3	C-2'; C-3'; 2'-CH <sub>3</sub>
2'	-	42.7	-
2'-CH <sub>3</sub>	1.43 (s, 3H)	28.2	C-1'; C-2'; C-3'
3'	5.98 (dd, 10.6; 17.6, 1H)	146.8	C-1'; C-2'; 2'-CH <sub>3</sub>
4'	5.08 (d, 10.6, 1H) 5.13 (d, 17.6, 1H)	112.0	C-2'; C-3'
1''	3.16 (d, 15.1, 1H) 2.93 (dd, 7.6; 15.1, 1H)	28.1	C-9; C-10; C-11; C-2''; C-3''
2''	4.42 (d, 7.3, 1H)	77.5	C-10; C-1''; C-3''; C-4''; C-5''
3''	-	146.7	-
4''	4.97 (s, 1H) 4.86 (s, 1H)	110.2	C-2''; C-5''
5''	1.87 (s, 3H)	18.7	C-2''; C-3''; C-4''
5-OH	6.15 (s, 1H)	-	C-4a; C-5; C-6
9-OH	9.15 (s, 1H)	-	-
11-OH	13.48 (s, 1H)	-	C-10; C-11; C-11a

**Figure 2.** Selected HMBC correlation for 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one.

The cytotoxic activity of compound **1** against murine leukemia P-388 cells, as determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, showed IC<sub>50</sub> values of 4.10 µg/mL and was categorized as moderate activity. The antiplasmodial activity of **1** against *P. falciparum* showed IC<sub>50</sub> values of 2.88 µg/mL and was categorized as moderate activity.

### 3. Experiment Section

#### 3.1. General

NMR spectra were recorded on a JEOL 400 ECA spectrophotometer (Tokyo, Japan) in CDCl<sub>3</sub> at 400 (<sup>1</sup>H) and 100 (<sup>13</sup>C) MHz using tetramethyl silane (TMS) as the internal standard. The mass spectra were recorded using a Waters LCT Premier XE (Santa Clara, CA, USA). The UV spectra were measured with a Shimadzu series 1800 spectrophotometer (Kyoto, Japan). The IR spectra were recorded with Perkin-Elmer spectrum-100 FT-IR (Waltham, MA, USA). Column chromatography and planar radial chromatography were carried out using silica gel 60 G 1.07734.1000 and Si gel 60 PF<sub>254</sub> 1.07749.1000 (Merck, Darmstadt, Germany).

#### 3.2. Plant Material

The stem bark of *C. Pseudomole* was collected in Sungai Mendawak, anak Sungai Kapuas, District Kubu Raya, Kalimantan, Indonesia on April 2015. The specimen was identified at the Herbarium Bogoriense, Center of Biological Research and Development, National Institute of Science, Bogor, Indonesia.

#### 3.3. Extraction and Isolation

The dried stem bark of *C. pseudomole* (3.0 kg) was macerated in MeOH twice for 2 days each time. After evaporation of the solvent in a rotary evaporator, 260 g of pale brown semi-solid was obtained. Further, the MeOH extract was partitioned first with *n*-hexane (1:1 *v/v*); water was added (10% *v/v*) in the second step to increase the polarity; then the MeOH extract was partitioned with EtOAc (1:1 *v/v*). The EtOAc extract (35 g) was subjected to column chromatography over silica gel and eluted with *n*-hexane-ethyl acetate (from 9:1 to 1:1) to give fractions A–C. Fraction C was then subjected further to column chromatography and eluted with *n*-hexane-EtOAc (from 9:1 to 1:1) to produce three subfractions C<sub>1</sub>–C<sub>3</sub>. Subfraction C<sub>2</sub> was purified by planar radial chromatography using *n*-hexane-CHCl<sub>3</sub> (from 3:7 to 7:3), CHCl<sub>3</sub> and CHCl<sub>3</sub>-EtOAc 9:1 to yield compound 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one (18 mg).

#### 3.4. Cytotoxic Assay

The cytotoxic properties of **1** against murine leukemia P-388 cells were evaluated according to the MTT method as previously described [14–16]. Artonin E was used as the positive control.

#### 3.5. Antiplasmodial Assay

The antiplasmodial properties of **1** against *P. falciparum* were evaluated according to the Trager–Jensen method as previously described [17,18]. Chloroquine was used as the positive control.

**Supplementary Materials:** The following are available online at <http://www.mdpi.com/1422-8599/2017/4/M961/s1>, IR, HRESIMS, 1D and 2D NMR spectra are reported in the supplementary materials as Figures S1–S6.

**Acknowledgments:** This research was supported by Universitas Airlangga, Ministry of Research, Technology and Higher Education, Republic of Indonesia (Penelitian Hibah Kompetensi, Universitas Airlangga, 2017).

**Author Contributions:** Mulyadi Tanjung designed the whole experiment of bioactivity and wrote the manuscript. Tjitjik Srie Tjahjandarie researched data, analyzed the NMR and HRESIMS spectra and contributed to the manuscript, Ratih Dewi Saputri designed the whole experiment. All authors read and approved the final manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Tanjung, M.; Saputri, R.D.; Tjahjandarie, T.S. 5,9,11-Trihydroxy-2,2-dimethyl-10-(3'-methyl-2'-butenyl)-3-(2''-methyl-3''-butenyl)pyrano[2,3-a]xanthen-12(2H)-one from the stem bark of *Calophyllum pseudomole*. *Molbank* **2016**, M906. [[CrossRef](#)]
2. Tjahjandarie, T.S.; Saputri, R.D.; Tanjung, M. 5,9,11-Trihydroxy-2,2-dimethyl-3-(2-methylbut-3-en-2-yl)pyrano[2,3-a]xanthen-12(2H)-one from the stem bark of *Calophyllum tetrapterum* Miq. *Molbank* **2017**, M936. [[CrossRef](#)]
3. Wei, D.J.; Mei, W.L.; Zhong, H.M.; Zeng, Y.B.; Wu, X.D.; Dai, H.F. A new prenylated xanthone from the branches of *Calophyllum inophyllum*. *J. Asian Nat. Prod. Res.* **2011**, *13*, 265–269. [[CrossRef](#)] [[PubMed](#)]
4. Daud, S.B.; Ee, G.C.L.; Malek, E.A.; Teh, S.S.; See, I. A new coumarin from *Calophyllum hosei*. *Nat. Prod. Res.* **2014**, *28*, 1534–1538. [[CrossRef](#)] [[PubMed](#)]
5. Joshi, S.P.; Kulkarni, S.R.; Phalgune, U.D.; Puranik, V.G. New dipyrancoumarin from the leaves of *Calophyllum apetalum* Willd. *Nat. Prod. Res.* **2013**, *27*, 1896–1901. [[CrossRef](#)] [[PubMed](#)]
6. Ha, L.D.; Hansen, P.E.; Duus, F.; Pham, H.D.; Nguyen, L.D. A new chromanone acid from the bark of *Calophyllum dryobalanoides*. *Phytochem. Lett.* **2012**, *5*, 287–291. [[CrossRef](#)]
7. Lim, C.K.; Subramaniam, H.; Say, Y.H.; Jong, V.Y.M.; Khaledi, H.; Chee, C.F. A new chromanone acid from the stem bark of *Calophyllum teysmannii*. *Nat. Prod. Res.* **2015**, *29*, 1970–1977. [[CrossRef](#)] [[PubMed](#)]
8. McKee, T.C.; Covington, C.D.; Fuller, R.W.; Bokesch, H.R.; Young, S.; Cardellina, J.H.; Kadushin, M.R.; Soejarto, D.D.; Stevens, P.F.; Cragg, G.M. Pyranocoumarins from tropical species of the genus *Calophyllum*: Chemotaxonomic study of extracts in the national cancer institute collection. *J. Nat. Prod.* **1998**, *61*, 1252–1256. [[CrossRef](#)] [[PubMed](#)]
9. Reyes, M.H.; Basualdo, M.C.; Abe, F.; Estrada, M.J.; Soler, C.; Chilpa, R.R. HIV-1 inhibitory compounds from *Calophyllum brasiliense* leaves. *Biol. Pharm. Bull.* **2004**, *27*, 1471–1475. [[CrossRef](#)]
10. Xiao, Q.; Zeng, Y-B.; Mei, W.L.; Zhao, Y-X.; Deng, Y-Y.; Dai, H-F. Cytotoxic prenylated xanthenes from *Calophyllum inophyllum*. *J. Asian Nat. Prod. Res.* **2008**, *10*, 993–997. [[CrossRef](#)] [[PubMed](#)]
11. Mah, S.H.; Ee, G.C.L.; Teh, S.S.; Sukari, M.A. *Calophyllum inophyllum* and *Calophyllum soulattri* source of anti-proliferative xanthenes and their structure-activity relationships. *Nat. Prod. Res.* **2013**, *27*, 98–101.
12. Hay, A.; Helesbeux, J.; Duval, O.; Labaied, M.; Grellier, P.; Richomme, P. Antimalarial xanthenes from *Calophyllum caledonicum* and *Garcinia vieillardii*. *Life Sci.* **2004**, *75*, 3077–3085. [[CrossRef](#)] [[PubMed](#)]
13. Tanjung, M.; Saputri, R.D.; Tjahjandarie, T.S. Antioxidant activity of two isomeric benzoxepinderivatives from the stem bark of *Bauhinia acuelata* L. *J. Chem. Pharm. Res.* **2014**, *6*, 705–708.
14. Tanjung, M.; Hakim, E.H.; Syah, Y.M. Prenylated dihydrostilbenes from *Macaranga rubiginosa*. *Chem. Nat. Compd.* **2017**, *53*, 215–218. [[CrossRef](#)]
15. Tanjung, M.; Tjahjandarie, T.S.; Sentosa, M.H. Antioxidant and cytotoxic agent from the rhizomes of *Kaempferia pandurata*. *Asian Pac. J. Trop. Dis.* **2013**, *3*, 401–404. [[CrossRef](#)]
16. Tanjung, M.; Hakim, E.H.; Elfahmi; Latip, J.; Syah, Y.M. Dihydroflavonol and flavonol derivatives from *Macaranga recurvata*. *Nat. Prod. Commun.* **2012**, *10*, 1309–1310.
17. Tanjung, M.; Saputri, R.D.; Wahjoedi, R.A.; Tjahjandarie, T.S. 4-Methoxy-3-(3-methylbut-2-en-1-yl)-7-(3-methylbut-2-en-1-yl)oxyquinolin-2(1H)-one from *Melicope moluccana* T.G. Hartley. *Molbank* **2017**, M939. [[CrossRef](#)]
18. Tjahjandarie, T.S.; Pudjiastuti, P.; Saputri, R.D.; Tanjung, M. Antimalaria and antioxidant activity of phenolic compounds isolated from *Erythrina crista-galli* L. *J. Chem. Pharm. Res.* **2014**, *6*, 786–790.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).