

Comparative Chemical Profiles of Essential Oils and Hydrolate Extracts from Fresh Flowers of Eight *Paeonia suffruticosa* Andr. Cultivars from Central China

Gaoming Lei ^{1,*}, Jie Li ¹, Tao Zheng ², Junqiao Yao ³, Jingjing Chen ¹ and Lengxin Duan ¹

¹ Department of Pharmaceutical Sciences, Medical College, Henan University of Science and Technology, Luoyang 471023, China; lijie@haust.edu.cn (J.L.); 170317200708@stu.haust.edu.cn (J.C.); lengxinduan@haust.edu.cn (L.D.)

² National Flower Garden Administration of Luoyang, Luoyang 471022, China; tony7811@163.com

³ Department of Scientific Research, Peony Institute of Luoyang, Luoyang 471022, China; mdsyj@163.com

* Correspondence: leigm@haust.edu.cn; Tel.: +86-187-3630-3548




Figure S1. Mass spectra (GC-MS, 70 eV) of representative oxygenated compounds from hydrolate extracts of eight *Paeonia suffruticosa* Andr. cultivars: (a) 2-phenylethanol; (b) geraniol; (c) citronellol; (d) 1,3,5-trimethoxybenzene; (e) (Z)-3-hexen-1-ol.