

Article

Isolation of a Melanoblast Stimulator from *Dimocarpus longan*, Its Structural Modification, and Structure–Activity Relationships for Vitiligo

Jae-Won Song¹, Sunju Choi², Gayeong Kim², Hyang Bok Lee², P. Sankara Rao¹, Jeonghyun Shin³ , Eun Ki Kim² and Dong-Gyu Cho^{1,*} 

¹ Department of Chemistry and Chemical Engineering, Inha University, Incheon 22212, Korea; qkwldpdy@naver.com (J.-W.S.); p.sankarmsc89@gmail.com (P.S.R.)

² Department of Biological Engineering, Inha University, Incheon 22212, Korea; tjsw1174@naver.com (S.C.); teclra94@naver.com (G.K.); hbjbb@hanmail.net (H.B.L.); ekkim@inha.ac.kr (E.K.K.)

³ Department of Dermatology, School of Medicine, Inha University, Incheon 22212, Korea; jshin@inha.ac.kr

* Correspondence: dgcho@inha.ac.kr; Tel.: +82-32-860-7686

Abstract: A novel melanoblast stimulator (**1**) was isolated from *Dimocarpus longan*. Its analogs were also synthesized to support a new furan-based melanoblast stimulator scaffold for treating vitiligo. Isolated 5-(hydroxymethyl)furfural (HMF, **1**) is a well-known compound in the food industry. Surprisingly, the melanogenic activity of HMF (**1**) was discovered here for the first time. Both HMF and its synthetic analog (**16**) promote the differentiation and migration of melanoblasts in vitro. Typically, stimulator (**1**) upregulated MMP2 expression, which promoted the migration of melanoblasts in vitro.

Keywords: melanoblast; stimulator; melanogenesis; vitiligo structure–activity relationships



Citation: Song, J.-W.; Choi, S.; Kim, G.; Lee, H.B.; Rao, P.S.; Shin, J.; Kim, E.K.; Cho, D.-G. Isolation of a Melanoblast Stimulator from *Dimocarpus longan*, Its Structural Modification, and Structure–Activity Relationships for Vitiligo. *Molecules* **2022**, *27*, 2135. <https://doi.org/10.3390/molecules27072135>

Academic Editor: Jamshed Iqbal

Received: 3 March 2022

Accepted: 24 March 2022

Published: 25 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Vitiligo is an acquired depigmentation disease of the skin [1,2]. Histologically, loss of epidermal melanocytes results in white patches in the skin. The cause is still not clear, although multiple pathogeneses such as autoimmune, neurological, self-destruction, stress, and viral hypotheses are being considered [3–6]. Patients with vitiligo account for approximately 0.05–1% of the world's population. Patients often experience difficulties in interpersonal or social relationships due to their appearance, and desperately need an effective therapy to cure vitiligo.

Treatment for vitiligo requires halting of melanocyte destruction and repigmentation of the affected skin. Various methods have been developed to treat vitiligo, including topical or oral steroids [7], immune suppressants [8], phototherapy [9,10], and laser therapy [11]. However, the remaining problems with repigmentation therapies are that repigmentation takes a long time and is usually incomplete. Thus, a more effective medicine is necessary to repigment the lesions. In vitiligo, epidermal melanocytes are lost, but melanoblasts of the outer root sheath of the hair follicles are not affected. Melanoblasts are non-pigmented precursors of melanocytes. They lack tyrosinase and do not stain DOPA or produce melanin. However, upon activation, melanoblasts migrate into the epidermis and differentiate into melanocytes. They then produce melanin pigments. During phototherapy, the repigmentation spots begin from the follicle opening and slowly spread out. Thus, an effective therapeutic agent could be proposed that promotes migration and differentiation of inactive melanoblasts from the outer root sheath of hair follicles to near the epidermis [12,13]. Therefore, melanoblast activation is an ideal target for vitiligo [14,15].

There have been many potential molecules (melanocyte stimulators) that stimulate melanogenesis, mostly at the melanocyte level [16]. However, none of the melanoblast stimulators as single molecules are known except 8-methoxypsoralen (8-MOP) of furocoumarins [17]. Thus far, PUVA therapy (8-MOP plus ultraviolet (UV)A radiation) is still

effectively used, but it is being replaced by UVB or laser therapies due to the synergistic side effects of 8-MOP (carcinogenesis) [18–20]. In addition, the central role of furocoumarins is elusive due to their diverse therapeutic effects [21] and complex nature of pathogenesis. Here, we report a new melanoblast stimulator scaffold (isolated and characterized from *Dimocarpus longan* and its synthetic analogs) as a furan derivative and a potential substitute for 8-MOP. This new melanoblast activation scaffold can guide us to view previously reported coumarin derivatives (melanocyte stimulators) [22–24] as furan and its analogs having different coumarins (Figure 1). We also showed in vitro that furan derivatives (1 and 16) can induce the secretion of matrix metalloproteinase (MMP)-2, which helps in melanoblast migration.




Figure 1. Coumarin (in blue) vs. furan scaffolds for melanogenesis stimulators.

2. Results and Discussion

2.1. Isolation of HMF

To isolate an active compound of *Dimocarpus longan*, its dried fruit was extracted with methanol at 22 ± 3 °C (rt) to obtain a crude extract, which was suspended in water and hexane. The separated water layers were successively partitioned using CHCl_3 and EtOAc. Each fraction, including the hexane layer, was evaluated for its ability to induce melanoblast migration, while the extract with CHCl_3 displayed promising activity (2.1-fold increase in cell migration (%) at 50 $\mu\text{g}/\text{mL}$). The CHCl_3 extract was further purified by column chromatography on silica gel using a gradient mobile phase (MeOH/ CHCl_3), thereby obtaining three fractions. Among them, the first fraction was further purified by preparative HPLC, which showed the highest migration activity (Figure S1 in ESI). To identify the active chemical structure of the migration activity in *D. longan*, typical characterization techniques, including NMR spectroscopy and high-resolution mass analysis, were conducted. Surprisingly, a few carbon signals were observed in the ^{13}C -NMR spectrum (Figure S4 in ESI) and an ion peak was found at m/z 126.0319 $[\text{M}]^+$ in the high-resolution mass spectrum (Figure S9 in ESI). These results indicated that the molecule is very small. The coupling constant of the two signals ($J = 3.5$ Hz at 7.45 and 6.56 ppm) in the aromatic region corresponds to a five-membered aromatic ring, while one sharp singlet proton at 9.51 ppm appears to be similar to the proton of the aldehyde group (Figure S3 in ESI). Another set of protons has strong cross-peaks in the COSY NMR spectrum and one of the two peaks at 5.51 ppm is likely attached to a heteroatom because it does not have a cross-peak in HSQC (Figures S5 and S6 in ESI). This structural information enabled the identification of 5-(hydroxymethyl)furfural (HMF). The stacked ^1H and ^{13}C -NMR spectra of the isolated and purchased materials matched exactly (Figures S7 and S8 in ESI). The efficacy profiles of the products isolated from the extract were also similar to those of HMF. Thus, we concluded that the melanogenesis activity of the dried fruit of *D. longan* is responsible for HMF. HMF is a well-known compound in the food industry and is isolated from various plants such as *Schisandra*, *Cornus officinalis*, and *Laurencia undulata* [25]. It is often formed in processed foods such as coffee and dried plum. The toxicity of HMF is not severe, as it is a part of our daily diet, unless it is overconsumed. It has also been reported

that there is no adverse effect in the range of 80–100 mg/Kg body weight per day in various animal experiments [26]. Thus, it is a great surprise to us that the melanogenesis activity of HMF was observed for the first time. This finding could be significant from a food and medicinal perspective.

2.2. SAR Study

To support the therapeutic effect of HMF on vitiligo and its preventive effect on gray hair, systematic derivatives were conducted by replacing the two substituents at the α -position of the furan ring in Table 1. The detailed synthetic schemes are in Schemes S1 and S2 (ESI). As stated earlier, melanoblast activation is an ideal target for therapy for vitiligo. Thus, the melanin content of melanoblasts can indicate their differentiation into melanocytes with treatment. Furthermore, measuring the melanin content and migration ability of melanoblasts can be an indicator of the efficacy of the stimulator. In addition, a lower chemical toxicity is necessary to ensure the safety of the stimulator. All these indices were scrutinized through the SAR study. First, the efficacy profile of HMF (**1**) appeared promising as shown in Table 1. The efficacy of the 2, 5-substituted furans was also assessed at a concentration of 10 μ M. In the first round, while R_1 was fixed as the aldehyde group, R_2 was screened. Replaced ether and amide groups had no effect on melanin content, while cell migration increased slightly, and there was no cytotoxicity (**2–6**) as compared to the negative control (DMSO). The ester group increased both melanin content and migration rate (%) compared to the negative control. Surprisingly, the aldehyde group on R_2 (furan-2,5-dicarbaldehyde (**10**)) decreased both the melanin content and cell migration. Interestingly, the cell viability assay of **10** revealed that it was the most toxic compound in Table 1. The best compound of the first round was the methyl ester (**9**), although both its activities were slightly lower than those of HMF (**1**). A larger ester (**7** and **8**) led to a decrease in both activities compared with compound (**9**). These results indicated that CH_2OH was the best substitute for R_2 (**1**). Therefore, CH_2OH was fixed for R_1 , whereas the other functional groups were examined for R_2 in the second round. When carboxylic acid derivatives were introduced for R_2 , smaller substituents were better than larger substituents (**13** > **12** > **11**) in terms of melanin content and cell migration. This trend was also observed for the three esters (**7–9**). In addition, a one-carbon elongated analog (**15**) of 2,5-furandimethanol (**14**) has an activity profile similar to those of acid derivatives (**11–13**) and slightly improved cytotoxicity. It was apparently better than **14**. After the second round, the activities of HMF (**1**) were still the best, and we thought that further derivatives could be limited because smaller derivatives were always better than larger ones. However, one exception was found in the case of **14** vs. **15**. These observations led to the design of compound **16** (one carbon elongation from HMF (**1**)) in the third round, which showed comparable therapeutic indices to HMF. Furthermore, the acetylated compound exhibited a lower activity than compound **16**. Two-carbon elongated compounds (**17**) from HMF decreased the activity compared to one-carbon elongated compounds (**16**). The activities of the acylated compound (**19**) from **17** remained almost the same.

Table 1. SAR studies of furan derivatives.













					
No.	R ₁	R ₂	Melanin Content ^a	Cell Migration ^a	Cell Viability ^a
1	CHO	CH ₂ OH	147 (4) ^b	318 (17) ^b	108 (4) ^b
2	CHO	CH ₂ OCH ₂ Ph	82 (4)	114 (19)	104 (2)
3	CHO		95 (8)	134 (12)	106 (1)
4	CHO		97 (4)	115 (20)	109 (2)
5	CHO		94 (2)	229 (12)	100 (1)
6	CHO		96 (3)	139 (18)	106 (3)
7	CHO	CO ₂ CH ₂ Ph	134 (2)	202 (62)	100 (4)
8	CHO		122 (5)	178 (20)	109 (2)
9	CHO	CO ₂ CH ₃	136 (1)	296 (19)	110 (3)
10	CHO	CHO	95 (6)	117 (22)	23 (6)
11	CH ₂ OH	CO ₂ CH ₂ CH ₃	104 (5)	111 (14)	71 (4)
12	CH ₂ OH	CO ₂ CH ₃	145 (6)	161 (29)	80 (1)
13	CH ₂ OH	CO ₂ H	158 (1)	264 (8)	93 (1)
14	CH ₂ OH	CH ₂ OH	139 (2)	173 (30)	86 (2)
15	CH ₂ OH		139 (20)	214 (14)	97 (1)
16	CHO		160 (1)	356 (25)	97 (1)
17	CHO		112 (2)	261 (17)	98 (1)

Table 1. Cont.

					
No.	R ₁	R ₂	Melanin Content ^a	Cell Migration ^a	Cell Viability ^a
18	CHO		116 (3)	230 (25)	100 (1)
19	CHO		115 (3)	260 (17)	101 (2)

^a The values (%) were obtained by comparing the observed effects of negative control (DMSO) for each therapeutic index. ^b The numbers in parentheses are measurement errors (\pm). α -MSH (0.1 μ M) was used as a positive control. All assays were conducted in Melb-a (melanoblast) and treated with a 10 μ M concentration of each.

2.3. Biological Studies

Two potential stimulators (**1** and **16**) were selected as representatives from the extract of *D. longan* and the synthetic stimulator, respectively. The cytotoxicity of both compounds at various concentrations was assessed using a typical MTT assay with Melb-a cells (melanoblasts). Stimulator (**1**) showed significant cytotoxicity at 1000 μ M, while stimulator (**16**) did at 100 μ M in Melb-a (Figure 2a). This observed cytotoxicity should affect the dose dependence of the two therapeutic indices. Thus, the dose dependence of stimulators (**1** and **16**) for the indices was observed in the different concentration ranges (Figure 2). As shown in Figure 2b, stimulator (**1**) displayed dose dependence for the melanin content at a concentration range of 1–100 μ M. On the other hand, stimulator (**16**) was dose dependent at a concentration range of 1–10 μ M. These trends were also observed in cell migration assays of the two compounds. Compared to the results of the melanin content assay, the dose dependence of cell migration upon treatment with stimulators **1** and **16** was steeper (Figure 2c). Typically, cell migration (%) induced by stimulator **16** was significantly better than that of stimulator **1** below 10 μ M. For example, the cell migration rate with stimulator **16** was increased by approximately 2.8 times at a concentration of 10 μ M as compared to the negative control, while the migration was increased by about 2.4 times in the case of stimulator **1** under the same conditions. These trends in the cell migration assay were also found with a stimulator (**1** or **16**) at various concentrations. Images were magnified 40 times under a microscope for Figure 2d.




Figure 2. Cont.




Figure 2. A graph showing the results of evaluating (a) cell survival, (b) the intracellular melanin content, (c) the melanoblast migration at various concentrations of 1 and 16, (d) a set of photographs (magnified 40 \times) taken after migration assay at various concentrations of 1 and 16.

The observed steep dose dependence of cell migration rate may be related to metalloproteinases (MMPs) that play an important role in cleaving components of the extracellular matrix. Previous studies have reported that MMP2 is upregulated when 8-MOP or α -MSH were used in melanoblasts (Melb-a) [17]. Specifically, MMP2, MMP9, and MT1-MMP transcripts are expressed, but MMP2 is overexpressed. In the same study, the percentage of cell migration increased with the overexpression of MMP2, whereas cell migration was reduced upon treatment with additional GM6001 (MMP inhibitor). Thus, we speculated that the steep dose dependence of our stimulators for cell migration was highly related to the overexpression of MMP2. This possibility was thoroughly examined with stimulator 1 as a representative furan derivative and compared with α -MSH (positive control). To identify whether MMP2 is responsible for the migration of melanoblasts in vitro, we first quantified their mRNA expression levels in melanoblasts using semiquantitative RT-PCR when treated with the stimulator (1). Both stimulator 1 and α -MSH led to a significant

increase in MMP2 mRNA levels (34% and 29%, respectively) compared to the negative control (Figure 3a). In addition, the expression of MMP2 was determined by Western blotting (active form at 62 kDa) with a stimulator (**1**, 100 μ M) and α -MSH (0.1 μ M) (Figure 3b).




Figure 3. (a) mRNA expression of MMP2, (b) MMP2 expression, (c) gelatinase activity of MMP2 in melanoblasts treated with DMSO, α -MSH, or **1**.

MMP2 expression level increased by 86% compared to that in the control when the stimulator (**1**, 100 μ M) was used in Melb-a melanoblasts. These results confirm that the expression of MMP2 is induced by this stimulator (**1**). Gelatin zymography was also performed to confirm the proteolytic activity of MMP2 (Figure 3c). When the stimulator (**1**, 100 μ M) was used, the protein hydrolysis activity of MMP2 was 60% higher than that of the negative control. Under our experimental conditions, the major MMP secreted by Melb-a melanoblasts was detected using zymography at 62 kDa. This was consistent with the Western blot results (Figure 3b). In contrast, in the presence of both GM6001 (10 nM) and the stimulator (**1**, 100 μ M), the protein activity dropped significantly, but not completely,

compared to the same experiments performed without GM6001 (Figure 4a). Thus, MMP2 is expressed by melanoblasts, and this expression can be stimulated by the stimulator (**1**, 100 μ M), with α -MSH (0.1 μ M) as a positive control. To determine whether inhibition of MMP2 affects melanoblast migration, a cell migration assay was conducted with or without GM6001 (10 nM) in the presence of either the stimulator (**1**, 100 μ M) or α -MSH (0.1 μ M) as a positive control (Figure 4b). Melanoblast migration was enhanced by the stimulator (**1**) and suppressed by GM6001 (1.7 times). These results support that MMP2 is upregulated by the stimulator (**1**) and is crucial for the migration of melanoblasts. It should also be noted that the migratory activity of the stimulator (**1**) is at least 50% higher than that of 8-MOP in Melb-a melanoblasts, compared to the relative value of 8-MOP to α -MSH in the literature [17].




Figure 4. (a) MMP2 expression, (b) cell migration rate (%) of MMP2 in melanoblasts treated with only **1** (or **1** and GM6001).

3. Materials and Methods

3.1. Isolation of Stimulator (**1**)

Isolation procedures of the stimulator (**1**) are summarized in Figure S1 (ESI). It was fully characterized by standard methods, including ^1H NMR, ^{13}C NMR, 2D NMR, and high-resolution mass spectroscopy in the ESI.

3.2. Materials and Synthesis

Reagents were purchased at the highest commercial quality and used without further purification, unless otherwise stated. The yields of the synthesized compounds were measured after chromatographic purification.

The syntheses of all the new compounds are summarized in Schemes S1 and S2 (ESI). All new compounds are characterized by standard methods, including ^1H NMR, ^{13}C NMR,

and high-resolution mass spectroscopy. The ^1H and ^{13}C NMR spectra are included in the ESI.

3.3. Plant Material

The dried fruits of *Dimocarpus longan* (Vietnam) were purchased from Kyungdong herbal market (Seoul, Korea) in June 2019 and identified by one of the authors (E.K.K.). A voucher specimen (IUI-2019-06-01) was deposited at the Herbarium of Inha University (IUI), Inha University, Republic of Korea.

3.4. Biological Studies

Biological experimental details are available in the ESI (cell culture, MTT assay, cell migration assay, cell differentiation assay, RT-PCR, Western blot analysis, and gelatin zymography).

4. Conclusions

We report a new melanoblast stimulator (**1**) scaffold isolated from *Dimocarpus longan*, and SAR studies of its synthetic analogous support that a new furan scaffold possibly induces melanogenesis in vitiligo and gray hair because both medical conditions lack melanocytes rather than melanoblasts. Isolated 5-(hydroxymethyl)furfural (HMF, **1**) is a well-known compound in the food industry. Thus, it is surprising that such a small molecule, HMF (**1**), induces melanogenesis in melanoblasts. This is the first report regarding the melanogenic activity of HMF (**1**). Typically, HMF and its synthetic analog (**16**) promote the differentiation and migration of melanoblasts in vitro. Furthermore, the migration rates of the two compounds were at least 50% higher than that of the positive control (8-MOP). Both compounds displayed a step-dose dependence on cell migration rate. Stimulator **1**, a representative furan derivative, promoted cell migration and upregulated MMP2, which is related to the cleavage of the components of the extracellular matrix. In contrast, the cell migration rate decreased when both the stimulator (**1**) and GM6001 (MMP inhibitor) were used simultaneously. Therefore, the mechanism of stimulator **1** in cell migration is suggested to be related to MMP2 upregulation.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules27072135/s1>, Synthetic details, biological experimental, characterization data of stimulator (**1**), and ^1H and ^{13}C NMR spectra for all new compounds are available online.

Author Contributions: Conceptualization of the compounds, J.-W.S. and D.-G.C.; Synthesis, methodology, P.S.R. and J.-W.S.; biological studies, S.C., G.K. and H.B.L.; supervision over the project, E.K.K. and D.-G.C.; writing and preparation of the original draft, J.-W.S. and D.-G.C.; writing, reviewing, and editing, J.S., E.K.K. and D.-G.C.; acquisition of funding, D.-G.C. All authors have read and agreed to the published version of the manuscript.

Funding: Inha University Research Grant.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available in supplementary material.

Acknowledgments: This research was supported by an Inha University Research Grant.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the compounds are not available from the authors.

References

1. Ezzedine, K.; Eleftheriadou, V.; Whitton, M.; van Geel, N. Vitiligo. *Lancet* **2015**, *386*, 74–84. [[CrossRef](#)]
2. Picardo, M.; Dell'Anna, M.L.; Ezzedine, K.; Hamzavi, I.; Harris, J.E.; Parsad, D.; Taieb, A. Vitiligo. *Nat. Rev. Dis. Primers* **2015**, *1*, 15011. [[CrossRef](#)] [[PubMed](#)]

3. Grimes, P.E.; Halder, R.M.; Jones, C.; Chakrabarti, S.G.; Enterline, J.; Minus, H.R.; Kenney, J.A., Jr. Autoantibodies and Their Clinical Significance in a Black Vitiligo Population. *Arch. Dermatol.* **1983**, *119*, 300–303. [[CrossRef](#)] [[PubMed](#)]
4. Koga, M. Vitiligo: A new classification and therapy. *Br. J. Dermatol.* **1977**, *97*, 255–261. [[CrossRef](#)]
5. Laddha, N.C.; Dwivedi, M.; Mansuri, M.S.; Gani, A.R.; Ansarullah, M.; Ramachandran, A.V.; Dalai, S.; Begum, R. Vitiligo: Interplay between oxidative stress and immune system. *Exp. Dermatol.* **2013**, *22*, 245–250. [[CrossRef](#)]
6. Lerner, A.B. On the etiology of vitiligo and gray hair. *Am. J. Med.* **1971**, *51*, 141–147. [[CrossRef](#)]
7. Imamura, S.; Tagami, H. Treatment of Vitiligo with Oral Corticosteroids. *Dermatology* **1976**, *153*, 179–185. [[CrossRef](#)]
8. Grimes, P.E.; Soriano, T.; Dytoc, M.T. Topical tacrolimus for repigmentation of vitiligo. *J. Am. Acad. Dermatol.* **2002**, *47*, 789–791. [[CrossRef](#)]
9. Ortonne, J.P.; Schmitt, D.; Thivolet, J. PUVA-induced Repigmentation of Vitiligo: Scanning Electron Microscopy of Hair Follicles. *J. Invest. Dermatol.* **1980**, *74*, 40–42. [[CrossRef](#)]
10. Taieb, A.; Alomar, A.; Böhm, M.; Dell’Anna, M.L.; De Pase, A.; Eleftheriadou, V.; Ezzedine, K.; Gauthier, Y.; Gawkrödger, D.J.; Jouary, T.; et al. Guidelines for the management of vitiligo: The European Dermatology Forum consensus. *Br. J. Dermatol.* **2013**, *168*, 5–19. [[CrossRef](#)]
11. Alhowaish, A.K.; Dietrich, N.; Onder, M.; Fritz, K. Effectiveness of a 308-nm excimer laser in treatment of vitiligo: A review. *Lasers Med. Sci.* **2013**, *28*, 1035–1041. [[CrossRef](#)]
12. Nishimura, E.K. Melanocyte stem cells: A melanocyte reservoir in hair follicles for hair and skin pigmentation. *Pigment Cell Melanoma Res.* **2011**, *24*, 401–410. [[CrossRef](#)]
13. Paus, R. Migrating melanocyte stem cells: Masters of disaster? *Nat. Med.* **2013**, *19*, 818–819. [[CrossRef](#)]
14. Babitha, S.; Nguyen, D.H.; Park, S.-J.; Shin, J.-H.; Reyes, G.A.; Caburian, A.; Kim, E.-K. Potential of *Cassia alata* leaf extract in inducing differentiation and migration of mouse melanoblasts. *Biotechnol. Bioprocess Eng.* **2010**, *15*, 1071–1076. [[CrossRef](#)]
15. Babitha, S.; Shin, J.-H.; Nguyen, D.H.; Park, S.-J.; Reyes, G.A.; Caburian, A.; Kim, E.K. A stimulatory effect of *Cassia occidentalis* on melanoblast differentiation and migration. *Arch. Dermatol.* **2011**, *303*, 211–216. [[CrossRef](#)]
16. Niu, C.; Aisa, H.A. Upregulation of Melanogenesis and Tyrosinase Activity: Potential Agents for Vitiligo. *Molecules* **2017**, *22*, 1303. [[CrossRef](#)]
17. Lei, T.C.; Vieira, W.D.; Hearing, V.J. In Vitro Migration of Melanoblasts Requires Matrix Metalloproteinase-2: Implications to Vitiligo Therapy by Photochemotherapy. *Pigment Cell Res.* **2002**, *15*, 426–432. [[CrossRef](#)]
18. Fitzsimons, C.P.; Long, J.; MacKie, R.M. Synergistic carcinogenic potential of methotrexate and PUVA in psoriasis. *Lancet* **1983**, *1*, 235–236. [[CrossRef](#)]
19. Marx, J.L.; Auerbach, R.; Possick, P.; Myrow, R.; Gladstein, A.H.; Kopf, A.W. Malignant melanoma in situ in two patients treated with psoralens and ultraviolet A. *J. Am. Acad. Dermatol.* **1983**, *9*, 904–911. [[CrossRef](#)]
20. Stern, R.S.; Laird, N.; Melski, J.; Parrish, J.A.; Fitzpatrick, T.B.; Bleich, H.L. Cutaneous squamous-cell carcinoma in patients treated with PUVA. *N. Engl. J. Med.* **1984**, *310*, 1156–1161. [[CrossRef](#)]
21. Carbone, A.; Montalbano, A.; Spanò, V.; Musante, I.; Galiotta, L.J.V.; Barraja, P. Furocoumarins as multi-target agents in the treatment of cystic fibrosis. *Eur. J. Med. Chem.* **2019**, *180*, 283–290. [[CrossRef](#)]
22. Matsuda, H.; Hirata, N.; Kawaguchi, Y.; Yamazaki, M.; Naruto, S.; Shibano, M.; Taniguchi, M.; Baba, K.; Kubo, M. Melanogenesis Stimulation in Murine B16 Melanoma Cells by Umberiferiae Plant Extracts and Their Coumarin Constituents. *Biol. Pharm. Bull.* **2005**, *28*, 1229–1233. [[CrossRef](#)]
23. Niu, C.; Pang, G.X.; Li, G.; Dou, J.; Nie, L.F.; Himit, H.; Kabas, M.; Aisa, H.A. Synthesis and biological evaluation of furocoumarin derivatives on melanin synthesis in murine B16 cells for the treatment of vitiligo. *Bioorg. Med. Chem.* **2016**, *24*, 5960–5968. [[CrossRef](#)]
24. Pang, G.X.; Niu, C.; Mamat, N.; Aisa, H.A. Synthesis and in vitro biological evaluation of novel coumarin derivatives containing isoxazole moieties on melanin synthesis in B16 cells and inhibition on bacteria. *Bioorg. Med. Chem. Lett.* **2017**, *27*, 2674–2677. [[CrossRef](#)]
25. van Putten, R.-J.; van der Waal, J.C.; de Jong, E.; Rasrendra, C.B.; Heeres, H.J.; de Vries, J.G. Hydroxymethylfurfural A Versatile Platform Chemical Made from Renewable Resources. *Chem. Rev.* **2013**, *113*, 1499–1597. [[CrossRef](#)]
26. Abraham, K.; Gürtler, R.; Berg, K.; Heinemeyer, G.; Lampen, A.; Appel, K.E. Toxicology and risk assessment of 5-Hydroxymethylfurfural in food. *Mol. Nutr. Food Res.* **2011**, *55*, 667–678. [[CrossRef](#)]