

From Chemistry to Bioactivity: HS-SPME-GC-MS Profiling and Bacterial Growth Inhibition of Three Different Propolis Samples from Romania, Australia, and Uruguay

Radosław Balwierz^{1*}, Katarzyna Kasperkiewicz², Martyna Straszak², Daria Siodłak¹, Katarzyna Pokajewicz¹, Ibtissem Ben Hammouda¹, Piotr Paweł Wieczorek¹, Anna Kurek-Górecka^{3*}, Zenon Czuba³, and Tomasz Baj⁴

¹ Institute of Chemistry, University of Opole, Oleska 48, 45-052, Opole, Poland

² Faculty of Natural Sciences, Institute of Biology, Biotechnology and Environmental Protection, University of Silesia in Katowice, Jagiellonska St. 28, 40-032 Katowice, Poland

³ Department of Microbiology and Immunology, Faculty of Medical Sciences in Zabrze, Medical University of Silesia in Katowice, Jordana 19, 41-808, Zabrze, Poland

⁴ Department of Pharmacognosy with the Medicinal Plants Garden, Medical University of Lublin, 20-001, Lublin, Poland.

Table of Content

Figure S1. Heatmap with hierarchical clustering (HCA) of volatile compounds in propolis samples. Color intensity reflects the z-value, highlighting regional chemical markers.....	2
Figure S2. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on <i>Escherichia coli</i> . a – 200 mg/ml concentration of the propolis extract, b – 100 mg/ml concentration of the propolis extract, e – ethanol 70%.....	3
Figure S3. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on <i>Staphylococcus aureus</i> . a – 200 mg/ml concentration of the propolis extract, b – 100 mg/ml concentration of the propolis extract, e – ethanol 70%	4
Figure S4. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on <i>Streptococcus mutans</i> . a – 200 mg/ml concentration of the propolis extract, b – 100 mg/ml concentration of the propolis extract, e – ethanol 70%	5
Figure S5. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on <i>Yersinia enterocolitica</i> . a – 200 mg/ml concentration of the propolis extract, b – 100 mg/ml concentration of the propolis extract, e – ethanol 70%	6
Figure S6. Antimicrobial activity of tetracycline (30 µg) on (A) <i>Escherichia coli</i> , (B) <i>Staphylococcus aureus</i> , (C) <i>Streptococcus mutans</i> and (D) <i>Yersinia enterocolitica</i>	7




Figure S1. Heatmap with hierarchical clustering (HCA) of volatile compounds in propolis samples. Color intensity reflects the z-value, highlighting regional chemical markers.

Figure S2. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on *Escherichia coli*. **a** – 200 mg/ml concentration of the propolis extract, **b** – 100 mg/ml concentration of the propolis extract, **e** – ethanol 70%




Figure S3. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on *Staphylococcus aureus*. **a** – 200 mg/ml concentration of the propolis extract, **b** – 100 mg/ml concentration of the propolis extract, **e** – ethanol 70%




Figure S4. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on *Streptococcus mutans*. **a** – 200 mg/ml concentration of the propolis extract, **b** – 100 mg/ml concentration of the propolis extract, **e** – ethanol 70%




Figure S5. Antimicrobial activity of ethanolic propolis extract from (01) Poland, (02) Turkey, (03) Romania, (04) Australia, (05 and 06) Uruguay on *Yersinia enterocolitica*. **a** – 200 mg/ml concentration of the propolis extract, **b** – 100 mg/ml concentration of the propolis extract, **e** – ethanol 70%




Figure S6. Antimicrobial activity of tetracycline (30 μ g) on (A) *Escherichia coli*, (B) *Staphylococcus aureus*, (C) *Streptococcus mutans* and (D) *Yersinia enterocolitica*

