

## Section A – Taxa List

**Table S1.** Full list of taxa recorded through BRUV sampling around UK coastal waters and their presence in different habitats, arrival times and times of MaxN (min). SG = Seagrass, K = Kelp, S = Sand, MCS = Mixed Coarse Sediments, MW = Midwater, MB = Mussel Beds, AR = Artificial Reef, RR = Rocky Reef. Note- Arrival times and time of MaxN have been recorded where access to raw footage / data is available.

Family	Species	Habitat								First Arrival Time (mins) ( $\pm$ 1SE)	Time of MaxN (mins) ( $\pm$ 1 SE)
		SG	K	S	MCS	MW	MB	AR	RR		
Ammodytidae	-	x	x	x	x					46.15 $\pm$ 11.71	52.06 $\pm$ 10.90
Ammodytidae	<i>Ammodytes tobianus</i>		x							-	-
Ammodytidae	<i>Hyperoplus lanceolatus</i>	x		x						-	-
Anguillidae	<i>Anguilla anguilla</i>	x								27.03 $\pm$ 2.99	27.03 $\pm$ 2.99
Atherinidae	-	x								158.59 $\pm$ 56.45	168.78 $\pm$ 61.42
Atherinidae	<i>Atherina presbyter</i>	x	x	x	x					52.74 $\pm$ 12.56	75.23 $\pm$ 15.36
Balistidae	<i>Balistes capriscus</i>	x		x						31.82 $\pm$ 0.00	31.82 $\pm$ 0.00
Belonidae	<i>Belone belone</i>	x				x				126.59 $\pm$ 0.00	126.59 $\pm$ 0.00
Blenniidae	<i>Lipophrys pholis</i>		x							-	-
Blenniidae	<i>Parablennius gattorugine</i>		x							-	-
Callionymidae	<i>Callionymus lyra</i>	x				x				151.09 $\pm$ 51.21	151.09 $\pm$ 51.21
Cancridae	-			x						-	-
Cancridae	<i>Cancer pagurus</i>		x	x	x		x			45.73 $\pm$ 13.79	60.20 $\pm$ 11.82
Clupeidae	<i>Sprattus sprattus</i>		x	x	x					27.13 $\pm$ 13.27	41.67 $\pm$ 4.74

**Table S1.** Full list of taxa recorded through BRUIV sampling around UK coastal waters and their presence in different habitats, arrival times and times of MaxN (min). SG = Seagrass, K = Kelp, S = Sand, MCS = Mixed Coarse Sediments, MW = Midwater, MB = Mussel Beds, AR = Artificial Reef, RR = Rocky Reef. Note- Arrival times and time of MaxN have been recorded where access to raw footage / data is available.

Family	Species	Habitat								First Arrival Time (mins) ( $\pm$ 1SE)	Time of MaxN (mins) ( $\pm$ 1 SE)
		SG	K	S	MCS	MW	MB	AR	RR		
Cottidae	-	x								19.88 $\pm$ 0.00	190.88 $\pm$ 0.00
Cottidae	<i>Taurulus bubalis</i>	x	x							-	-
Congridae	<i>Conger conger</i>	x								-	-
Dasyatidae	<i>Dasyatis pastinaca</i>	x		x						-	-
Gadidae	-	x	x	x	x		x			39.54 $\pm$ 7.98	51.48 $\pm$ 8.90
Gadidae	<i>Gadus morhua</i>	x	x							23.15 $\pm$ 14.95	23.15 $\pm$ 14.95
Gadidae	<i>Merlangius merlangus</i>	x		x	x	x	x			35.55 $\pm$ 6.00	53.96 $\pm$ 9.06
Gadidae	<i>Pollachius pollachius</i>	x	x	x					x	29.93 $\pm$ 5.78	43.77 $\pm$ 7.88
Gadidae	<i>Pollachius virens</i>	x			x	x				30.85 $\pm$ 8.54	37.13 $\pm$ 6.85
Gadidae	<i>Trisopterus luscus</i>	x	x	x	x	x		x	x	23.77 $\pm$ 4.94	31.97 $\pm$ 5.18
Gadidae	<i>Trisopterus minutus</i>	x		x	x	x	x			31.66 $\pm$ 8.34	67.46 $\pm$ 12.27
Gasterosteidae	<i>Spinachia spinachia</i>	x	x		x					87.76 $\pm$ 20.11	112.68 $\pm$ 20.01
Gastropoda*	-	x		x	x					24.79 $\pm$ 7.94	35.47 $\pm$ 8.15
Gobiidae	-	x	x	x	x					17.60 $\pm$ 4.18	29.49 $\pm$ 7.64
Gobiidae	<i>Gobius niger</i>	x	x		x					9.18 $\pm$ 0.00	31.8 $\pm$ 0.00
Gobiidae	<i>Gobius paganellus</i>	x	x							-	-
Gobiidae	<i>Gobiusculus flavescens</i>	x	x		x	x		x	x	19.34 $\pm$ 2.85	26.06 $\pm$ 2.69

**Table S1.** Full list of taxa recorded through BRUIV sampling around UK coastal waters and their presence in different habitats, arrival times and times of MaxN (min). SG = Seagrass, K = Kelp, S = Sand, MCS = Mixed Coarse Sediments, MW = Midwater, MB = Mussel Beds, AR = Artificial Reef, RR = Rocky Reef. Note- Arrival times and time of MaxN have been recorded where access to raw footage / data is available.

Family	Species	Habitat								First Arrival Time (mins) ( $\pm$ 1SE)	Time of MaxN (mins) ( $\pm$ 1 SE)
		SG	K	S	MCS	MW	MB	AR	RR		
Gobiidae	<i>Pomatoschistus microps</i>	x				x				-	-
Gobiidae	<i>Pomatoschistus minutus</i>	x	x		x			x	x	12.19 $\pm$ 4.75	36.37 $\pm$ 14.82
Gobiidae	<i>Pomatoschistus pictus</i>		x							-	-
Goneplacidae	<i>Goneplax rhomboides</i>			x						-	-
Inachidae	<i>Macropodia</i>			x						46.06 $\pm$ 0.00	46.06 $\pm$ 0.00
Labridae	-	x	x	x	x					31.56 $\pm$ 4.09	37.56 $\pm$ 4.51
Labridae	<i>Centrolabrus exoletus</i>	x	x							14.38 $\pm$ 7.21	14.39 $\pm$ 7.21
Labridae	<i>Crenilabrus melops</i>	x	x							24.12 $\pm$ 14.85	27.05 $\pm$ 14.64
Labridae	<i>Ctenolabrus rupestris</i>	x	x	x	x				x	70.46 $\pm$ 48.67	70.46 $\pm$ 48.67
Labridae	<i>Labrus bergylta</i>	x	x	x	x	x		x	x	47.48 $\pm$ 11.81	53.48 $\pm$ 12.17
Labridae	<i>Labrus mixtus</i>	x	x						x	-	-
Labridae	<i>Symphodus melops</i>		x					x	x	-	-
Lotidae	-	x								238.44 $\pm$ 0.00	238.44 $\pm$ 0.00
Lotidae	<i>Ciliata mustela</i>	x							x	215.53 $\pm$ 33.36	215.53 $\pm$ 33.36

**Table S1.** Full list of taxa recorded through BRUIV sampling around UK coastal waters and their presence in different habitats, arrival times and times of MaxN (min). SG = Seagrass, K = Kelp, S = Sand, MCS = Mixed Coarse Sediments, MW = Midwater, MB = Mussel Beds, AR = Artificial Reef, RR = Rocky Reef. Note- Arrival times and time of MaxN have been recorded where access to raw footage / data is available.

Family	Species	Habitat								First Arrival Time (mins) ( $\pm$ 1SE)	Time of MaxN (mins) ( $\pm$ 1 SE)
		SG	K	S	MCS	MW	MB	AR	RR		
Majidae	<i>Maja squinado</i>	x	x	x	x		x	x		47.28 $\pm$ 8.34	53.91 $\pm$ 8.41
Moronidae	<i>Dicentrarchus labrax</i>		x	x				x		52.71 $\pm$ 0.00	52.71 $\pm$ 0.00
Mugilidae	<i>Chelon labrosus</i>			x				x		32.65 $\pm$ 0.00	32.65 $\pm$ 0.00
Mugilidae	-	x								131.39 $\pm$ 119.83	176.39 $\pm$ 74.83
Mullidae	<i>Mullus surmuletus</i>	x		x	x				x	18.10 $\pm$ 5.78	22.82 $\pm$ 10.50
Nephropidae	-			x	x					17.13 $\pm$ 8.41	21.35 $\pm$ 9.28
Nephropidae	<i>Homarus gammarus</i>	x	x	x	x			x	x	66.30 $\pm$ 15.78	66.76 $\pm$ 15.70
Octopodidae	-						x			41.28 $\pm$ 37.49	54.78 $\pm$ 23.99
Paguridae	-	x		x	x		x	x		26.86 $\pm$ 4.94	38.59 $\pm$ 6.33
Palaemonidae	<i>Palaemon serratus</i>	x			x			x		25.41 $\pm$ 6.28	28.48 $\pm$ 6.13
Pectinidae	-				x					58.03 $\pm$ 0.00	58.03 $\pm$ 0.00
Phocidae	<i>Halichoerus grypus</i>	x	x							30.29 $\pm$ 18.52	34.76 $\pm$ 18.19
Pleuronectidae	-	x		x						39.13 $\pm$ 9.41	48.25 $\pm$ 9.40
Pleuronectidae	<i>Limanda limanda</i>	x		x						15.28 $\pm$ 0.00	15.28 $\pm$ 0.00
Pleuronectidae	<i>Pleuronectes platessa</i>	x	x	x		x				82.99 $\pm$ 38.65	82.99 $\pm$ 38.65
Portunidae	-	x	x	x	x					22.00 $\pm$ 7.58	37.37 $\pm$ 12.16
Portunidae	<i>Carcinus maenas</i>	x			x			x		20.36 $\pm$ 3.80	30.24 $\pm$ 4.12
Portunidae	<i>Necora puber</i>	x	x		x				x	23.16 $\pm$ 9.71	23.69 $\pm$ 9.45

**Table S1.** Full list of taxa recorded through BRUIV sampling around UK coastal waters and their presence in different habitats, arrival times and times of MaxN (min). SG = Seagrass, K = Kelp, S = Sand, MCS = Mixed Coarse Sediments, MW = Midwater, MB = Mussel Beds, AR = Artificial Reef, RR = Rocky Reef. Note- Arrival times and time of MaxN have been recorded where access to raw footage / data is available.

Family	Species	Habitat								First Arrival Time (mins) ( $\pm$ 1SE)	Time of MaxN (mins) ( $\pm$ 1 SE)
		SG	K	S	MCS	MW	MB	AR	RR		
Rajidae	<i>Raja clavata</i>	x		x		x				40.91 $\pm$ 0.00	40.91 $\pm$ 0.00
Scombridae	<i>Scomber scombrus</i>	x								-	-
Scophthalmidae	<i>Scophthalmus rhombus</i>	x								-	-
Scyliorhinidae	-	x	x	x	x		x			58.24 $\pm$ 12.08	66.98 $\pm$ 13.58
Scyliorhinidae	<i>Scyliorhinus</i>	x								18.00 $\pm$ 4.90	17.60 $\pm$ 4.50
Scyliorhinidae	<i>Scyliorhinus canicula</i>	x	x	x	x	x				23.29 $\pm$ 2.11	27.74 $\pm$ 2.11
Scyliorhinidae	<i>Scyliorhinus stellaris</i>	x								120.1 $\pm$ 0.00	120.1 $\pm$ 0.00
Scyphozoa*	-		x							8.79 $\pm$ 0.00	8.79 $\pm$ 0.00
Sepiidae	<i>Sepiola atlantica</i>	x	x							152.26 $\pm$ 52.75	177.45 $\pm$ 44.02
Sepiidae	<i>Sepia officinalis</i>	x		x				x		26.89 $\pm$ 13.29	26.89 $\pm$ 13.29
Sepiolidae	-	x								155.14 $\pm$ 126.97	174.12 $\pm$ 145.95
Soleidae	-			x						27.57 $\pm$ 8.22	27.57 $\pm$ 8.22
Soleidae	<i>Solea solea</i>			x						47.27 $\pm$ 20.09	47.27 $\pm$ 20.09
Sparidae	-			x						23.93 $\pm$ 1.84	23.93 $\pm$ 1.84
Sparidae	<i>SpondylIOSoma cantharus</i>			x				x	x	18.36 $\pm$ 13.66	23.16 $\pm$ 18.46
Syngnathidae	-	x								125.08 $\pm$ 87.75	125.08 $\pm$ 87.75
Syngnathidae	<i>Entelurus aequoreus</i>	x								3.69 $\pm$ 0.50	17.19 $\pm$ 13.00

**Table S1.** Full list of taxa recorded through BRUIV sampling around UK coastal waters and their presence in different habitats, arrival times and times of MaxN (min). SG = Seagrass, K = Kelp, S = Sand, MCS = Mixed Coarse Sediments, MW = Midwater, MB = Mussel Beds, AR = Artificial Reef, RR = Rocky Reef. Note- Arrival times and time of MaxN have been recorded where access to raw footage / data is available.

Family	Species	Habitat								First Arrival Time (mins) ( $\pm$ 1SE)	Time of MaxN (mins) ( $\pm$ 1 SE)
		SG	K	S	MCS	MW	MB	AR	RR		
Syngnathidae	<i>Syngnathus acus</i>	x			x					158.19 $\pm$ 65.52	158.19 $\pm$ 65.52
Syngnathidae	<i>Syngnathus typhle</i>	x	x							-	-
Trachinidae	<i>Echiichthys vipera</i>	x								57.74 $\pm$ 24.78	192.74 $\pm$ 110.22
Triglidae	-			x						16.36 $\pm$ 3.92	16.36 $\pm$ 3.92
Triglidae	<i>Chelidonichthys cuculus</i>	x								-	-
Triglidae	<i>Chelidonichthys lucerna</i>	x		x					x	32.53 $\pm$ 7.48	32.75 $\pm$ 7.30
Triakidae	<i>Mustelus mustelus</i>			x						20.81 $\pm$ 2.71	23.50 $\pm$ 3.61
Unknown	-	x	x	x	x		x			46.83 $\pm$ 10.08	51.21 $\pm$ 9.87

\* Identified to Class.

## Section B – Poisson Regression Coefficients

**Table S2.** Coefficients from the Poisson model for species richness. Reference levels for the categorical predictors were as follows: Broad scale habitat = sand, time of day = day, bait type = none, image quality = excellent, spring/neap tide = mid. **Bold values indicate  $P = \leq 0.05$ .**

Predictor	Coef	SE Coef	Z-Value	P-Values
<b>Constant</b>	-0.348	0.281	-1.24	0.214
<b>Bait Type</b>				
Crab	0.358	0.402	0.89	0.373
Mackerel	0.830	0.299	2.77	<b>0.006</b>
Oily fish meal and fish oils	1.160	0.324	3.58	<b>&lt;0.001</b>
Sardines	0.672	0.566	1.19	0.235
Squid	0.655	0.343	1.91	0.056

**Table S3.** Coefficients from the Poisson model for relative abundance (MaxN). Reference levels for the categorical predictors were as follows: Broad scale habitat = sand, time of day = day, bait type = none, image quality = excellent, spring/neap tide = mid. **Bold values indicate  $P = \leq 0.05$ .**

Predictor	Coef	SE Coef	Z-Value	P-Values
<b>Constant</b>				
<b>Duration of Deployment</b>	0.00310	0.00104	2.99	<b>0.003</b>
<b>Bait Type</b>				
Crab	0.512	0.351	1.46	0.145
Mackerel	0.970	0.267	3.63	<b>&lt;0.001</b>
Oily fish meal and fish oils	0.992	0.335	2.96	<b>0.003</b>
Sardines	0.811	0.517	1.57	0.117
Squid	0.680	0.307	2.21	<b>0.027</b>
<b>Image Quality</b>				
Good	-0.486	0.405	-1.20	0.230
Poor	-0.894	0.418	-2.14	<b>0.032</b>

## Section C – Multivariate Analysis Pairwise Comparisons

**Table S4.** Pairwise comparisons for faunal assemblage composition. \*Note\* N/A has been excluded from this statistical analysis Bold values  $P \leq 0.01$ .

Source	<i>t</i>	<i>P</i> (perm)	Unique Perms
<b>Habitat Observed * Bait Type</b>			
<i>Mixed Coarse Sediment</i>			
Mackerel vs No bait	1.7949	<b>0.001</b>	9034
Mackerel vs Squid	1.5046	0.0137	9060
Mackerel vs Sardines	1.2402	0.1187	1762
Mackerel Oily fish meal	1.2225	0.1244	1762
No bait vs Squid	1.036	0.439	126
No bait vs Sardines	1.4385	0.1756	56
No bait vs Oily fish meal	1.6249	0.0344	56
Squid vs Sardines	0.73517	0.7730	41
Squid vs Oily fish meal	1.5668	0.0737	56
Sardines vs Oily fish meal	1.6173	0.2013	10
<i>Sand</i>			
Mackerel vs No bait	2.9881	<b>&lt;0.001</b>	9931
Mackerel vs Squid	2.0463	<b>&lt;0.001</b>	9943
Mackerel vs Crab	1.9546	<b>0.001</b>	9944
No bait vs Squid	2.9169	<b>0.001</b>	9624
No bait vs Crab	2.0300	<b>0.009</b>	3728
Squid vs Crab	1.3740	0.1222	9870
<i>Seagrass</i>			
Mackerel vs No bait	2.3400	<b>&lt;0.001</b>	9802
Mackerel vs Squid	1.5085	0.0175	8304
Mackerel vs Oily fish meal	3.8963	<b>&lt;0.001</b>	9924
Mackerel vs Prawn	2.3680	<b>&lt;0.001</b>	8259
Squid vs Oily fish meal	1.9675	<b>0.0010</b>	3112
Squid vs Prawn	1.5043	0.1059	10
Squid vs No bait	2.0670	0.0273	35
Oily fish meal vs Prawn	2.1780	<b>&lt;0.001</b>	3138
Oily fish meal vs No bait	2.4533	<b>&lt;0.001</b>	8082
Prawn vs No bait	2.7367	0.0298	35
<i>Kelp</i>			
Mackerel vs Oily fish meal	2.3184	<b>&lt;0.001</b>	9780



*Midwater*

Mackerel vs No bait	2.3687	<b>0.010</b>	191
Mackerel vs Squid	2.0674	0.0237	992
Mackerel vs Prawn	2.2428	0.0115	280
Squid vs Prawn	1.2241	0.3958	6
Squid vs No bait	1.271	0.4742	3
Prawn vs Nothing	1.0000	1.0000	1

**Habitat Observed \* Image Quality***Mixed Coarse Sediment*

Poor vs Good	1.3509	0.0594	9923
Poor vs Excellent	1.4802	0.049	22
Good vs Excellent	1.3046	0.1274	16

*Sand*

Poor vs Good	1.8104	<b>0.0025</b>	9927
--------------	--------	---------------	------

*Kelp*

Poor vs Good	1.6619	0.0182	5114
--------------	--------	--------	------

*Seagrass*

Poor vs Good	2.6896	<b>&lt;0.001</b>	9943
Poor vs Excellent	1.223	0.1066	28
Good vs Excellent	0.88415	0.6112	47

**Habitat Observed \* Time of Day***Mixed Coarse Sediment*

Day vs Evening	0.71279	0.8853	9012
Day vs Night	1.0467	0.450	33
Evening vs Night	0.9647	0.7963	5

*Sand*

Day vs Evening	2.0880	<b>&lt;0.001</b>	9918
Day vs Dawn	0.78797	0.7617	83
Evening vs Dawn	0.88683	0.6508	9

*Seagrass*

Day vs Evening	2.0439	<b>&lt;0.001</b>	9937
Day vs Dawn	2.4133	<b>&lt;0.001</b>	9936
Day vs Night	2.2477	<b>&lt;0.001</b>	9922
Evening vs Dawn	2.8219	<b>&lt;0.001</b>	9930
Evening vs Night	1.3174	0.0701	9927
Dawn vs Night	3.3442	<b>&lt;0.001</b>	9865

*Kelp*

Day vs Evening	1.0777	0.3623	11
----------------	--------	--------	----

Day vs Night	2.0131	<b>0.0028</b>	1001
Evening vs Night	2.3285	0.2026	5
<b>Habitat Observed * Tide (Spring / Neap)</b>			
<i>Mixed Coarse Sediment</i>			
Spring vs Neap	1.8047	<b>0.0021</b>	9944
Spring vs Mid	2.1161	<b>&lt;0.001</b>	9625
Neap vs Mid	1.7190	<b>&lt;0.001</b>	9644
<i>Sand</i>			
Spring vs Neap	1.6237	0.020	9959
Spring vs Mid	2.3531	<b>&lt;0.001</b>	9952
Neap vs Mid	2.2792	<b>&lt;0.001</b>	9943
<i>Seagrass</i>			
Spring vs Neap	2.2807	<b>&lt;0.001</b>	9926
Spring vs Mid	2.3253	<b>&lt;0.001</b>	9940
Neap vs Mid	2.1497	<b>&lt;0.001</b>	9938
<i>Kelp</i>			
Spring vs Neap	1.7477	0.034	119
Spring vs Mid	2.0015	<b>0.0066</b>	792
Neap vs Mid	2.0605	0.0168	56
<i>Midwater</i>			
Spring vs Neap	2.0002	0.0272	550
Spring vs Mid	1.25	0.3137	57
Neap vs Mid	1.8477	0.0282	86

## Section D – Multivariate Analysis SIMPER Results

*Table S5. SIMPER analysis in groups outlined by PERMANOVA showing the top five organisms which most contributed to the observed differences among broad habitats observed. MSC = Mixed coarse sediment, OFM = Oily fish meal and fish oils.*

Species	Av. Abun.	Av. Abun.	Av. Diss.	Diss./SD	Contrib. %	Cum %
<b>Habitat Observed x Bait Type</b>						
<b>Av. Diss.: 87.43</b>	<b>MCS</b>	<b>Sand</b>				
Gobiidae	0.33	0.40	9.51	0.71	10.88	10.88
<i>Scyliorhinus canicula</i>	0.62	0.34	7.93	0.84	9.07	19.95
<i>Pollachius pollachius</i>	0.61	0.05	7.04	0.47	8.06	28.00

Paguridae	0.11	0.53	6.37	0.58	7.29	35.29
<i>Merlangius merlangius</i>	0.15	0.30	5.57	0.67	6.37	41.66
<b>Av. Diss.: 88.87</b>	<b>MCS</b>	<b>Seagrass</b>				
<i>Gobiusculus flavescens</i>	0.04	0.61	7.76	0.87	8.73	8.73
<i>Pollachius pollachius</i>	0.61	0.44	7.30	0.84	8.21	16.94
<i>Scyliorhinus canicula</i>	0.62	0.56	6.44	0.87	7.25	24.19
<i>Atherina presbyter</i>	0.08	0.64	5.86	0.56	6.59	30.79
Labridae	0.26	0.22	5.37	0.59	6.04	36.83
<b>Av. Diss.: 94.10</b>	<b>MCS</b>	<b>Kelp</b>				
<i>Gobiusculus flavescens</i>	0.04	1.04	10.01	0.89	10.63	10.63
<i>Pollachius pollachius</i>	0.61	0.77	8.84	0.83	9.39	20.02
<i>Scyliorhinus canicula</i>	0.62	0.16	6.08	0.93	6.46	26.48
Gobiidae	0.33	0.03	5.46	0.55	5.80	32.29
<i>Labrus bergylta</i>	0.03	0.47	4.68	0.86	4.98	37.27
<b>Av. Diss.:93.55</b>	<b>MCS</b>	<b>Midwater</b>				
<i>Pollachius pollachius</i>	0.61	0.10	15.03	0.65	16.06	16.06
<i>Scyliorhinus canicula</i>	0.62	0.22	12.49	0.76	13.36	29.42
<i>Trisopterus luscus</i>	0.25	0.11	8.68	0.56	9.28	38.70
<i>Merlangius merlangius</i>	0.15	0.34	8.15	0.73	8.7	47.41
Gobiidae	0.33	0.00	5.09	0.42	5.44	52.85
<b>Av. Diss.: 95.50</b>	<b>MCS</b>	<b>Art. Reef</b>				
<i>SpondylIOSoma cantharus</i>	0.00	2.87	22.36	1.32	23.41	23.41
<i>Maja squinado</i>	0.19	1.16	10.70	1.04	11.21	34.62
<i>Gobiusculus flavescens</i>	0.04	0.90	6.82	0.72	7.14	41.77
<i>Scyliorhinus canicula</i>	0.62	0.00	6.08	0.93	6.37	48.13
Gobiidae	0.33	0.00	5.38	0.52	5.64	53.77
<b>Av. Diss.:98.06</b>	<b>MCS</b>	<b>Rocky reef</b>				
<i>Ctenolabrus rupestris</i>	0.06	2.89	18.59	3.25	18.96	18.96
<i>Trisopterus luscus</i>	0.25	2.33	15.56	2.45	15.87	34.83
<i>Parablennius gattorugine</i>	0.00	1.24	8.35	2.18	8.52	43.35
<i>SpondylIOSoma cantharus</i>	0.00	1.46	8.15	0.88	8.31	51.66
<i>Symphodus melops</i>	0.00	0.71	4.23	1.14	4.31	55.97
<b>Av. Diss.: 92.93</b>	<b>Sand</b>	<b>Seagrass</b>				
<i>Gobiusculus flavescens</i>	0.00	0.61	8.73	0.87	9.39	9.39
<i>Scyliorhinus canicula</i>	0.34	0.56	7.60	0.85	8.18	17.57
Gobiidae	0.40	0.06	6.56	0.53	7.06	24.63
<i>Merlangius merlangius</i>	0.30	0.45	5.67	0.69	6.10	30.73

<i>Pollachius pollachius</i>	0.05	0.44	5.59	0.88	6.01	36.74
<b>Av. Diss.: 97.00</b>	<b>Sand</b>	<b>Kelp</b>				
<i>Gobiusculus flavescens</i>	0.00	1.04	11.06	0.89	11.41	11.41
<i>Pollachius pollachius</i>	0.05	0.77	7.97	0.79	8.22	19.63
Gobiidae	0.40	0.03	6.68	0.56	6.89	26.52
<i>Scyliorhinus canicula</i>	0.34	0.16	5.60	0.75	5.77	32.29
<i>Labrus bergylta</i>	0.00	0.47	5.11	0.86	5.27	37.55
<b>Av. Diss.: 94.78</b>	<b>Sand</b>	<b>Midwater</b>				
Paguridae	0.53	0.00	11.38	0.52	12.00	12.00
<i>Merlangius merlangius</i>	0.30	0.34	10.55	0.82	11.13	23.14
<i>Scyliorhinus canicula</i>	0.34	0.22	9.94	0.74	10.49	33.63
Gobiidae	0.40	0.00	8.00	0.47	8.44	42.07
<i>Mustelus mustelus</i>	0.18	0.00	6.26	0.38	6.60	48.68
<b>Av. Diss.: 96.0</b>	<b>Sand</b>	<b>Art. Reef</b>				
<i>SpondylIOSoma cantharus</i>	0.00	2.87	23.98	1.32	24.95	24.95
<i>Maja squinado</i>	0.12	1.16	12.24	1.00	12.73	37.68
<i>Gobiusculus flavescens</i>	0.00	0.90	7.32	0.72	7.61	45.30
Gobiidae	0.40	0.00	6.65	0.51	6.92	52.22
Paguridae	0.53	0.25	5.93	0.60	6.17	58.39
<b>Av. Diss.: 98.71</b>	<b>Sand</b>	<b>Rocky reef</b>				
<i>Ctenolabrus rupestris</i>	0.00	2.89	20.56	3.54	20.80	20.80
<i>Trisopterus luscus</i>	0.10	2.33	16.50	2.39	16.69	37.49
<i>Parablennius gattorugine</i>	0.00	1.24	8.89	2.15	8.99	46.48
<i>SpondylIOSoma cantharus</i>	0.03	1.46	8.60	0.89	8.70	55.18
<i>Symphodus melops</i>	0.00	0.57	3.94	0.80	3.98	67.89
<b>Av. Diss.: 84.33</b>	<b>Seagrass</b>	<b>Kelp</b>				
<i>Gobiusculus flavescens</i>	0.61	1.04	9.70	0.92	11.51	11.51
<i>Pollachius pollachius</i>	0.44	0.77	7.24	0.89	8.59	20.10
<i>Scyliorhinus canicula</i>	0.56	0.16	5.67	0.83	6.73	26.82
<i>Atherina presbyter</i>	0.64	0.23	4.95	0.49	5.87	32.70
<i>Labrus bergylta</i>	0.10	0.47	4.32	0.82	5.13	37.82
<b>Av. Diss.: 91.20</b>	<b>Seagrass</b>	<b>Midwater</b>				
<i>Merlangius merlangius</i>	0.45	0.34	10.86	0.91	11.91	11.91
<i>Scyliorhinus canicula</i>	0.56	0.22	10.83	0.88	11.87	23.78
<i>Gobiusculus flavescens</i>	0.61	0.04	8.74	0.76	9.58	33.36
<i>Pomatoschistus microps</i>	0.15	0.11	6.24	0.42	6.84	40.21
<i>Pollachius pollachius</i>	0.44	0.10	6.22	0.66	6.82	47.03

<b>Av. Diss.: 94.64</b>	<b>Seagrass</b>	<b>Art. Reef</b>				
<i>Spondyllosoma cantharus</i>	0.00	2.87	21.02	1.27	22.21	22.21
<i>Maja squinado</i>	0.09	1.16	10.77	1.02	11.38	33.59
<i>Gobiusculus flavescens</i>	0.61	0.90	9.45	0.94	9.98	43.58
<i>Scyliorhinus canicula</i>	0.56	0.00	6.00	0.78	6.34	49.91
<i>Pollachius pollachius</i>	0.44	0.00	4.65	0.86	4.91	54.2
<b>Av. Diss.: 93.94</b>	<b>Seagrass</b>	<b>Rocky reef</b>				
<i>Ctenolabrus rupestris</i>	0.00	2.89	18.28	3.09	19.46	19.46
<i>Trisopterus luscus</i>	0.24	2.33	12.55	1.69	13.36	32.82
<i>Parablennius gattorugine</i>	0.00	1.24	7.89	2.06	8.40	41.22
<i>Spondyllosoma cantharus</i>	0.00	1.46	7.75	0.87	8.25	49.47
<i>Gobiusculus flavescens</i>	0.61	0.04	5.27	0.92	5.61	55.07
<b>Av. Diss.: 95.72</b>	<b>Kelp</b>	<b>Midwater</b>				
<i>Gobiusculus flavescens</i>	1.04	0.04	11.98	0.82	12.51	12.51
<i>Merlangius merlangius</i>	0.00	0.34	9.39	0.81	9.81	22.32
<i>Pollachius pollachius</i>	0.77	0.10	9.05	0.73	9.46	31.78
<i>Scyliorhinus canicula</i>	0.16	0.22	6.96	0.84	7.27	39.05
<i>Labrus bergylta</i>	0.47	0.04	5.35	0.82	5.59	44.64
<b>Av. Diss.: 92.17</b>	<b>Kelp</b>	<b>Art. Reef</b>				
<i>Spondyllosoma cantharus</i>	0.00	2.87	20.61	1.25	22.36	22.36
<i>Gobiusculus flavescens</i>	1.04	0.90	10.83	0.95	11.75	34.11
<i>Maja squinado</i>	0.14	1.16	10.11	0.99	10.97	45.08
<i>Pollachius pollachius</i>	0.77	0.00	6.51	0.73	7.06	52.14
<i>Pomatoschistus minutus</i>	0.29	0.40	4.65	0.59	5.04	57.18
<b>Av. Diss.: 87.09</b>	<b>Kelp</b>	<b>Rocky reef</b>				
<i>Ctenolabrus rupestris</i>	0.39	2.89	15.40	2.07	17.10	17.10
<i>Trisopterus luscus</i>	0.10	2.33	14.21	2.13	15.78	32.88
<i>Spondyllosoma cantharus</i>	0.00	1.46	7.63	0.86	8.47	41.36
<i>Parablennius gattorugine</i>	0.04	1.24	7.58	1.92	8.42	49.78
<i>Gobiusculus flavescens</i>	1.04	0.04	6.68	0.92	7.42	57.19
<b>Av. Diss.: 98.99</b>	<b>Midwater</b>	<b>Art. Reef</b>				
<i>Spondyllosoma cantharus</i>	0.00	2.87	25.10	1.24	25.36	25.36
<i>Maja squinado</i>	0.00	1.16	14.66	0.87	14.81	40.17
<i>Merlangius merlangius</i>	0.34	0.00	9.39	0.81	9.48	49.65
<i>Gobiusculus flavescens</i>	0.04	0.90	8.05	0.74	8.13	57.78
<i>Scyliorhinus canicula</i>	0.22	0.00	5.83	0.83	5.89	63.68

<b>Av. Diss.:95.77</b>	<b>Midwater</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.00	2.89	21.15	2.85	22.09	22.09
<i>Trisopterus luscus</i>	0.11	2.33	15.06	1.72	15.72	37.81
<i>Parablennius gattorugine</i>	0.00	1.24	9.14	1.89	9.55	47.36
<i>SpondylIOSoma cantharus</i>	0.00	1.46	8.70	0.86	9.08	56.44
<i>Merlangius merlangius</i>	0.34	0.00	7.01	0.85	7.32	63.76
<b>Av. Diss.: 88.00</b>	<b>Art. Reef</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.00	2.39	17.88	2.95	20.32	20.32
<i>SpondylIOSoma cantharus</i>	2.87	1.46	14.47	1.32	16.44	36.76
<i>Trisopterus luscus</i>	0.04	2.33	14.17	2.13	16.10	52.87
<i>Parablennius gattorugine</i>	0.00	1.24	7.72	2.01	8.77	61.64
<i>Maja squinado</i>	1.16	0.00	7.18	1.28	8.16	69.80
<b>Av. Diss.: 96.31</b>	<b>Mackerel</b>	<b>No Bait</b>				
Gobiidae	0.21	0.00	11.99	0.47	12.45	12.45
<i>Scyliorhinus canicula</i>	0.41	0.09	10.33	0.71	10.73	23.18
Paguridae	0.16	0.12	8.75	0.52	9.08	32.36
<i>Merlangius merlangius</i>	0.28	0.00	7.41	0.58	7.69	39.95
Unknown	0.06	0.12	5.76	0.44	5.99	45.93
<b>Av. Diss.: 85.75</b>	<b>Mackerel</b>	<b>Squid</b>				
Paguridae	0.16	0.44	10.69	0.69	12.47	12.47
Gobiidae	0.21	0.13	9.30	0.52	10.85	23.32
<i>Scyliorhinus canicula</i>	0.41	0.45	9.12	0.80	10.64	33.95
<i>Mustelus mustelus</i>	0.03	0.33	8.26	0.61	9.64	43.59
<i>Merlangius merlangius</i>	0.28	0.09	6.95	0.61	8.11	51.70
<b>Av. Diss.:80.95</b>	<b>Mackerel</b>	<b>Sardines</b>				
<i>Pollachius pollachius</i>	0.32	1.05	15.93	1.21	19.68	19.68
<i>Scyliorhinus canicula</i>	0.41	1.00	9.03	0.83	11.15	30.83
Gobiidae	0.21	0.00	7.81	0.56	9.65	40.47
<i>Homarus gammarus</i>	0.08	0.00	5.91	0.54	7.30	47.77
Labridae	0.11	0.00	5.85	0.43	7.23	55.00
<b>Av. Diss.:91.75</b>	<b>Mackerel</b>	<b>OFM</b>				
Unknown	0.06	0.93	7.28	1.11	7.93	7.93
<i>Atherina presbyter</i>	0.22	0.70	7.05	0.74	7.69	15.62
<i>Gobiusculus flavescens</i>	0.49	0.16	6.65	0.81	7.74	22.86
<i>Scyliorhinus canicula</i>	0.41	0.13	5.20	0.78	5.67	28.54

<i>Pollachius pollachius</i>	0.32	0.31	4.77	0.87	5.20	33.74
<b>Av. Diss.: 94.34</b>		<b>Mackerel</b>	<b>Prawn</b>			
<i>Merlangius merlangius</i>	0.28	0.75	15.62	1.36	16.56	16.56
<i>Callionymus lyra</i>	0.03	0.47	9.39	1.54	9.95	26.51
<i>Pomatoschistus microps</i>	0.02	0.44	8.43	1.49	8.93	35.44
<i>Belone belone</i>	0.01	0.00	6.44	0.27	6.83	42.26
<i>Scomber scombrus</i>	0.00	0.42	6.29	0.55	6.67	48.93
<b>Av. Diss.: 90.92</b>		<b>Mackerel</b>	<b>Crab</b>			
Paguridae	0.16	1.01	16.67	0.97	18.33	18.33
Gobiidae	0.21	0.23	13.27	0.58	14.59	32.93
<i>Scyliorhinus canicula</i>	0.41	0.08	8.97	0.68	9.87	42.79
<i>Merlangius merlangius</i>	0.28	0.00	6.87	0.59	7.56	50.35
<i>Mustelus mustelus</i>	0.03	0.08	4.29	0.41	4.72	60.59
<b>Av. Diss.: 92.54</b>		<b>Squid</b>	<b>No Bait</b>			
Paguridae	0.44	0.12	22.73	0.82	24.56	24.56
<i>Mustelus mustelus</i>	0.33	0.00	18.44	0.75	19.93	44.49
<i>Scyliorhinus canicula</i>	0.45	0.09	11.49	0.54	12.42	56.91
Unknown	0.06	0.12	8.01	0.46	8.65	65.56
<i>Pollachius pollachius</i>	0.23	0.12	6.75	0.29	7.30	72.86
<b>Av. Diss.: 49.28</b>		<b>Squid</b>	<b>Sardines</b>			
<i>Pollachius pollachius</i>	0.23	1.05	17.00	1.21	34.50	34.50
<i>Trisopterus luscus</i>	0.18	0.33	12.34	1.13	25.05	59.55
<i>Trisopterus minutus</i>	0.09	0.00	5.71	0.48	11.58	71.13
<b>Av. Diss.: 91.97</b>		<b>Squid</b>	<b>OFM</b>			
<i>Scyliorhinus canicula</i>	0.45	0.19	11.47	1.43	12.47	12.47
<i>Pomatoschistus microps</i>	0.13	0.00	7.07	0.92	7.69	20.16
<i>Merlangius merlangius</i>	0.09	0.45	6.79	0.74	7.39	27.55
Unknown	0.06	0.93	6.73	0.98	7.32	34.86
<i>Spinachia spinachia</i>	0.00	0.54	5.13	0.69	5.58	40.44
<b>Av. Diss.: 91.60</b>		<b>Squid</b>	<b>Prawn</b>			
<i>Pollachius pollachius</i>	0.23	0.00	39.02	0.81	42.60	42.60
<i>Trisopterus luscus</i>	0.18	0.00	19.51	0.50	21.30	63.90
<i>Labrus bergylta</i>	0.00	0.13	17.78	0.49	19.41	83.32
<b>Av. Diss.: 83.88</b>		<b>Squid</b>	<b>Crab</b>			
Paguridae	0.44	1.01	26.92	1.05	32.09	32.09
<i>Mustelus mustelus</i>	0.33	0.08	16.91	0.81	20.16	52.26
<i>Scyliorhinus canicula</i>	0.45	0.08	9.37	0.53	11.17	63.42
Gobiidae	0.13	0.23	5.66	0.66	6.75	70.17

<b>Av. Diss.: 98.28</b>	<b>OFM</b>	<b>No Bait</b>				
<i>Pomatoschistus microps</i>	0.00	0.14	11.68	0.86	11.88	11.88
Unknown	0.93	0.12	8.71	1.09	8.86	20.75
<i>Belone belone</i>	0.00	0.10	8.33	0.96	8.47	29.22
<i>Spinachia spinachia</i>	0.54	0.00	6.68	0.58	6.80	36.02
<i>Scyliorhinus canicula</i>	0.19	0.09	5.96	0.78	6.06	42.08
<b>Av. Diss.: 87.01</b>	<b>OFM</b>	<b>Sardines</b>				
<i>Pollachius pollachius</i>	0.31	1.05	15.87	1.02	18.23	18.23
<i>Scyliorhinus canicula</i>	0.19	1.00	15.40	0.95	17.70	35.94
<i>Spinachia spinachia</i>	0.54	0.00	15.40	0.95	17.70	53.64
Labridae	0.20	0.00	8.39	1.30	9.65	63.29
<i>Gobiusculus flavescens</i>	0.16	0.00	6.16	0.66	7.08	70.37
<b>Av. Diss.: 92.44</b>	<b>OFM</b>	<b>Prawn</b>				
<i>Merlangius merlangius</i>	0.45	0.75	12.24	1.91	13.24	13.24
<i>Pomatoschistus microps</i>	0.00	0.44	8.94	2.79	9.67	22.91
<i>Callionymus lyra</i>	0.10	0.47	8.90	2.11	9.63	32.54
<i>Scomber scombrus</i>	0.00	0.42	7.93	0.68	8.58	41.12
<i>Pleuronectes platessa</i>	0.00	0.33	6.66	1.27	7.20	48.32
<b>Av. Diss.: 68.12</b>	<b>Sardines</b>	<b>No Bait</b>				
<i>Scyliorhinus canicula</i>	1.00	0.09	25.65	1.03	37.66	37.66
<i>Pollachius pollachius</i>	1.05	0.12	21.30	1.17	31.27	68.93
<i>Trisopterus luscus</i>	0.33	0.10	14.20	0.93	20.84	89.77
<b>Av. Diss.: 87.62</b>	<b>Prawn</b>	<b>No Bait</b>				
<i>Labrus bergylta</i>	0.13	0.00	40.00	0.80	45.65	45.65
<i>Merlangius merlangius</i>	0.75	0.00	12.00	1.12	13.69	59.35
<i>Callionymus lyra</i>	0.47	0.00	7.90	1.07	9.02	68.37
<i>Scomber scombrus</i>	0.42	0.00	6.34	0.49	7.24	75.61
<b>Av. Diss.: 93.17</b>	<b>Crab</b>	<b>No Bait</b>				
Paguridae	1.01	0.12	39.84	1.18	42.76	42.76
Gadidae	0.08	0.07	13.60	0.46	14.60	57.36
Unknown	0.08	0.12	10.56	0.50	11.34	63.42
<i>Sepia officianalis</i>	0.08	0.00	4.38	0.33	4.71	70.17
<b>Habitat Observed x Image Quality</b>						
<b>Av. Diss.: 91.66</b>	<b>MCS</b>	<b>Sand</b>				
<i>Scyliorhinus canicula</i>	0.60	0.34	10.59	0.75	11.55	11.55
<i>Pollachius pollachius</i>	0.62	0.05	9.86	0.66	10.76	22.31
Paguridae	0.10	0.53	7.47	0.60	8.15	30.46



Gobiidae	0.32	0.39	6.38	0.56	6.97	37.42
<i>Homarus gammarus</i>	0.20	0.06	5.38	0.40	5.87	43.30
<b>Av. Diss.: 88.70</b>	<b>MCS</b>	<b>Seagrass</b>				
<i>Gobiusculus flavescens</i>	0.15	0.81	7.00	0.77	7.89	7.89
<i>Pollachius pollachius</i>	0.62	0.42	6.82	0.76	7.69	15.58
<i>Atherina presbyter</i>	0.08	0.70	5.89	0.52	6.64	22.22
<i>Scyliorhinus canicula</i>	0.60	0.36	5.82	0.77	6.56	28.78
Labridae	0.28	0.33	4.56	0.58	5.14	33.92
<b>Av. Diss.: 85.97</b>	<b>MCS</b>	<b>Kelp</b>				
Ammodytidae	0.09	1.44	8.79	0.58	10.22	10.22
<i>Pollachius pollachius</i>	0.62	0.28	7.86	0.82	9.14	19.36
<i>Scyliorhinus canicula</i>	0.60	0.46	7.47	0.90	8.69	28.05
Unknown	0.05	0.64	7.38	0.99	8.58	36.63
<i>Atherina presbyter</i>	0.08	0.53	6.40	0.68	7.45	44.07
<b>Av. Diss.: 92.13</b>	<b>MCS</b>	<b>Mussel Beds</b>				
<i>Trisopterus minutus</i>	0.14	1.41	11.55	4.50	12.54	12.54
Scyliorhinidae	0.08	1.21	9.69	2.81	10.52	23.06
<i>Merlangius merlangius</i>	0.17	1.21	8.85	2.37	9.60	32.66
Octopodidae	0.00	1.00	8.24	5.15	8.95	41.61
Paguridae	0.10	1.00	7.87	3.48	8.55	50.15
<b>Av. Diss.: 90.87</b>	<b>MCS</b>	<b>Art. Reef</b>				
Gadidae	0.21	0.00	29.36	5.25	32.31	32.31
<i>Spondyllosoma cantharus</i>	0.00	287	12.55	1.36	13.81	46.12
<i>Merlangius merlangius</i>	0.17	0.00	12.15	5.25	13.38	59.50
<i>Gobius niger</i>	0.04	0.00	7.02	5.25	7.72	67.22
Paguridae	0.10	0.25	6.48	3.06	7.14	74.35
<b>Av. Diss.: 93.99</b>	<b>MCS</b>	<b>Rocky Reef</b>				
Gadidae	0.21	0.00	24.28	7.05	24.28	24.28
<i>Ctenolabrus rupestris</i>	0.05	2.89	11.62	4.51	11.62	35.90
<i>Merlangius merlangius</i>	0.17	0.00	10.05	7.05	10.05	45.96
<i>Trisopterus luscus</i>	0.24	2.33	9.36	2.71	9.36	5.32
<i>Gobius niger</i>	0.04	0.00	5.80	7.05	5.80	61.12
<b>Av. Diss.: 94.61</b>	<b>Sand</b>	<b>Seagrass</b>				
<i>Gobiusculus flavescens</i>	0.00	0.81	6.69	0.68	7.07	7.07
Paguridae	0.53	0.08	6.27	0.61	6.63	13.70
<i>Atherina presbyter</i>	0.03	0.70	6.14	0.51	6.49	20.19

Unknown	0.18	0.36	5.90	0.67	6.23	26.42
<i>Merlangius merlangius</i>	0.30	0.42	5.60	0.64	5.92	32.34
<b>Av. Diss.: 91.96</b>	<b>Sand</b>	<b>Kelp</b>				
Unknown	0.18	0.64	9.11	1.06	9.90	9.90
Ammodytidae	0.00	1.44	8.46	0.56	9.20	9.20
<i>Scyliorhinus canicula</i>	0.34	0.46	8.33	0.74	9.06	28.16
<i>Atherina presbyter</i>	0.03	0.53	7.66	0.70	8.33	36.50
Scyliorhinidae	0.12	0.36	6.43	0.62	7.00	43.49
<b>Av. Diss.: 85.99</b>	<b>Sand</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.02	1.41	12.64	4.78	14.70	14.70
Scyliorhinidae	0.12	1.21	10.66	2.66	12.40	27.09
<i>Merlangius merlangius</i>	0.30	1.21	9.45	2.15	10.99	38.09
Octopodidae	0.00	1.00	8.97	5.08	10.43	48.51
Unknown	0.18	1.00	8.24	2.71	9.59	58.10
<b>Av. Diss.: 88.14</b>	<b>Seagrass</b>	<b>Kelp</b>				
Ammodytidae	0.25	1.44	10.04	0.57	11.39	11.39
<i>Atherina presbyter</i>	0.70	0.53	6.61	0.60	7.50	18.90
<i>Gobiusculus flavescens</i>	0.81	0.07	6.11	0.77	6.93	25.83
<i>Scyliorhinus canicula</i>	0.36	0.46	4.63	0.77	5.25	31.08
Unknown	0.36	0.64	4.58	0.74	5.19	36.27
<b>Av. Diss.: 86.57</b>	<b>Seagrass</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.10	1.41	9.32	3.12	10.76	10.76
<i>Merlangius merlangius</i>	0.42	1.21	7.12	1.59	8.22	18.98
Scyliorhinidae	0.18	1.21	7.02	2.26	8.11	27.09
Octopodidae	0.00	1.00	6.88	4.07	7.95	35.04
Paguridae	0.08	1.00	6.56	2.90	7.58	42.62
<b>Av. Diss.: 93.99</b>	<b>Seagrass</b>	<b>Art. Reef</b>				
<i>Spondyllosoma cantharus</i>	0.00	2.87	14.78	1.37	15.72	15.72
<i>Centrolabrus exoletus</i>	0.06	0.00	8.50	4.32	9.04	24.76
<i>Scyliorhinus canicula</i>	0.36	0.00	8.50	4.32	9.04	33.80
<i>Maja squinado</i>	0.14	1.16	6.75	1.36	7.18	40.99
<i>Gobiusculus flavescens</i>	0.81	0.90	6.04	1.71	6.42	47.41
<b>Av. Diss.: 85.65</b>	<b>Seagrass</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.00	2.89	13.48	4.59	15.74	15.74
<i>Centrolabrus exoletus</i>	0.06	0.00	6.76	6.05	7.89	23.63

<i>Scyliorhinus canicula</i>	0.36	0.00	6.76	6.05	7.89	31.52
<i>Trisopterus luscus</i>	0.25	2.33	6.08	1.48	7.09	38.61
<i>SpondylIOSoma cantharus</i>	0.00	1.46	6.03	0.87	7.04	45.65
<b>Av. Diss.: 84.04</b>	<b>Kelp</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.00	1.41	10.14	4.84	12.07	12.07
<i>Merlangius merlangius</i>	0.00	1.21	8.55	3.53	10.17	22.24
Ammodytidae	1.44	0.00	7.33	0.51	8.73	30.96
Octopodidae	0.00	1.00	7.17	4.84	8.53	39.49
Paguridae	0.00	1.00	7.17	4.84	8.53	48.02
<b>Av. Diss.: 88.00</b>	<b>Art. Reef</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.00	2.89	17.88	2.95	20.32	20.32
<i>SpondylIOSoma cantharus</i>	2.87	1.46	14.47	1.32	16.44	36.76
<i>Trisopterus luscus</i>	0.04	2.33	14.17	2.13	16.10	52.87
<i>Parablennius gattorugine</i>	0.00	1.24	7.72	2.01	8.77	61.64
<i>Maja squinado</i>	1.16	0.00	7.18	1.28	8.16	69.80
<b>Av. Diss.: 87.76</b>	<b>Poor</b>	<b>Good</b>				
Gobiidae	0.19	0.33	10.81	0.50	12.32	12.32
Paguridae	0.30	0.27	9.91	0.56	11.29	23.61
<i>Scyliorhinus canicula</i>	0.39	0.38	6.22	0.73	7.09	30.70
Unknown	0.36	0.12	5.05	0.54	5.76	36.45
<i>Merlangius merlangius</i>	0.33	0.25	4.94	0.56	5.63	42.09
<b>Av. Diss.: 89.50</b>	<b>Poor</b>	<b>Excellent</b>				
Gadidae	0.13	0.11	16.41	1.01	18.33	18.33
<i>Merlangius merlangius</i>	0.33	0.05	7.47	1.10	8.35	26.68
<i>Scyliorhinus canicula</i>	0.39	0.05	5.30	1.51	5.92	32.60
<i>Centrolabrus exoletus</i>	0.00	0.03	4.46	1.06	4.98	37.59
Paguridae	0.30	0.15	4.10	1.00	4.58	42.16
<b>Av. Diss.: 73.44</b>	<b>Good</b>	<b>Excellent</b>				
Gadidae	0.16	0.11	8.36	0.64	11.38	11.38
<i>Centrolabrus exoletus</i>	0.03	0.03	5.97	1.54	8.13	19.50
<i>Scyliorhinus canicula</i>	0.38	0.05	5.06	1.32	6.88	26.39
<i>Merlangius merlangius</i>	0.25	0.05	4.47	0.83	6.09	32.48
<i>Chelidonichthys lucerna</i>	0.00	0.02	4.43	1.64	6.03	38.50

**Habitat Observed x Time of Day**

**Av. Diss.: 92.59**      **MCS**      **Sand**

<i>Pollachius pollachius</i>	0.62	0.04	10.42	0.64	11.25	11.25
<i>Scyliorhinus canicula</i>	0.60	0.28	10.11	0.76	10.92	22.17
Gobiidae	0.32	0.31	7.99	0.55	8.63	30.79
Paguridae	0.10	0.42	7.13	0.58	7.70	38.50
<i>Homarus gammarus</i>	0.20	0.05	4.55	0.38	4.91	43.41
<b>Av. Diss.: 91.21</b>	<b>MCS</b>	<b>Seagrass</b>				
<i>Gobiusculus flavescens</i>	0.15	0.77	9.33	0.86	10.23	10.23
<i>Atherina presbyter</i>	0.08	0.83	9.06	0.55	9.93	20.17
Ammodytidae	0.09	0.66	7.03	0.43	7.70	27.87
<i>Pollachius pollachius</i>	0.62	0.40	6.88	0.78	7.55	35.42
<i>Scyliorhinus canicula</i>	0.60	0.29	6.08	0.84	6.67	42.08
<b>Av. Diss.: 86.84</b>	<b>MCS</b>	<b>Kelp</b>				
Ammodytidae	0.09	1.44	10.69	0.52	12.31	12.31
<i>Scyliorhinus canicula</i>	0.60	0.46	7.24	0.87	8.33	20.64
<i>Pollachius pollachius</i>	0.62	0.28	7.19	0.73	8.28	28.92
<i>Maja squinado</i>	0.19	0.46	6.53	0.99	7.52	35.45
Scyliorhinidae	0.08	0.36	5.59	0.60	6.44	42.88
<b>Av. Diss.: 92.86</b>	<b>MCS</b>	<b>Mussel Beds</b>				
<i>Trisopterus minutus</i>	0.14	1.41	11.42	4.74	12.29	12.29
<i>Merlangius merlangius</i>	0.17	1.21	9.81	3.21	10.57	22.86
Octopodidae	0.00	1.00	8.07	4.74	8.69	31.56
Paguridae	0.10	1.00	8.07	4.74	8.69	40.25
Scyliorhinidae	0.08	1.21	7.85	1.71	8.46	48.71
<b>Av. Diss.: 96.69</b>	<b>MCS</b>	<b>Art. Reef</b>				
<i>SpondylIOSoma cantharus</i>	0.00	2.87	23.63	1.29	24.44	24.44
<i>Maja squinado</i>	0.19	1.16	12.22	0.96	12.64	37.08
<i>Gobiusculus flavescens</i>	0.15	0.90	7.60	0.74	7.87	44.95
<i>Pollachius pollachius</i>	0.62	0.00	6.49	0.60	6.71	51.66
<i>Scyliorhinus canicula</i>	0.60	0.00	6.21	0.79	6.42	58.08
<b>Av. Diss.: 95.63</b>	<b>MCS</b>	<b>Rocky Reef</b>				
<i>Ctenolabrus rupestris</i>	0.05	2.89	19.76	3.11	20.66	20.66
<i>Trisopterus luscus</i>	0.24	2.33	14.86	1.96	15.54	36.20
<i>Parablennius gattorugine</i>	0.00	1.24	8.74	2.08	9.14	45.34
<i>SpondylIOSoma cantharus</i>	0.00	1.46	8.43	0.88	8.82	54.15
<i>Symphodus melops</i>	0.00	0.71	4.40	1.13	4.60	58.75
<b>Av. Diss.: 96.97</b>	<b>Sand</b>	<b>Seagrass</b>				

<i>Gobiusculus flavescens</i>	0.00	0.77	10.59	0.82	10.92	10.92
<i>Atherina presbyter</i>	0.12	0.83	10.42	0.53	10.74	21.67
Ammodytidae	0.04	0.66	7.87	0.41	8.11	29.78
Paguridae	0.42	0.06	4.89	0.54	5.04	34.82
<i>Scyliorhinus canicula</i>	0.28	0.29	4.59	0.66	4.73	39.55
<b>Av. Diss.: 92.81</b>	<b>Sand</b>	<b>Kelp</b>				
Ammodytidae	0.04	1.44	11.26	0.49	12.14	12.14
<i>Scyliorhinus canicula</i>	0.28	0.46	9.70	0.81	10.46	22.59
Scyliorhinidae	0.10	0.36	7.55	0.60	8.13	30.72
<i>Maja squinado</i>	0.10	0.46	7.40	0.99	7.97	38.69
Unknown	0.14	0.64	6.43	0.68	6.93	45.62
<b>Av. Diss.: 99.40</b>	<b>Sand</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.01	1.41	14.09	5.70	14.18	14.18
<i>Merlangius merlangius</i>	0.24	1.21	12.13	3.51	12.21	26.38
Scyliorhinidae	0.10	1.21	11.92	4.83	11.99	38.38
Octopodidae	0.00	1.00	9.96	5.70	10.02	48.40
Paguridae	0.42	1.00	9.96	5.70	10.02	58.43
<b>Av. Diss.:96.99</b>	<b>Sand</b>	<b>Art. Reef</b>				
<i>Spondyllosoma cantharus</i>	0.03	2.87	27.42	1.30	28.27	28.27
<i>Maja squinado</i>	0.10	1.16	16.15	0.86	16.65	44.92
<i>Gobiusculus flavescens</i>	0.00	0.90	8.33	0.72	8.59	53.50
Paguridae	0.42	0.25	7.10	0.60	7.32	60.82
<i>Symphodus melops</i>	0.00	0.37	4.28	0.54	4.42	65.24
<b>Av. Diss.: 99.11</b>	<b>Sand</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.01	2.89	22.68	3.22	22.88	22.88
<i>Trisopterus luscus</i>	0.08	2.33	18.41	2.35	18.57	41.46
<i>Parablennius gattorugine</i>	0.00	1.24	9.88	1.99	9.97	51.52
<i>Spondyllosoma cantharus</i>	0.03	1.46	9.31	0.89	9.40	60.82
<i>Symphodus melops</i>	0.00	0.71	4.88	1.13	4.93	65.75
<b>Av. Diss.: 89.68</b>	<b>Seagrass</b>	<b>Kelp</b>				
Ammodytidae	0.66	1.44	12.73	0.63	14.19	14.19
<i>Atherina presbyter</i>	0.83	0.53	7.62	0.52	8.49	22.68
<i>Gobiusculus flavescens</i>	0.77	0.07	7.61	0.81	8.49	22.68
<i>Scyliorhinus canicula</i>	0.29	0.46	5.55	0.81	6.19	37.36
<i>Maja squinado</i>	0.12	0.46	4.69	0.95	5.23	42.59

<b>Av. Diss.: 82.42</b>	<b>Seagrass</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.08	1.41	8.95	2.45	10.86	10.86
Scyliorhinidae	0.14	1.21	7.46	1.97	9.05	19.92
<i>Merlangius merlangius</i>	0.33	1.21	7.00	1.61	8.50	28.41
Octopodidae	0.00	1.00	6.93	3.39	8.41	36.83
Paguridae	0.06	1.00	6.13	1.98	7.43	44.26
<b>Av. Diss.: 93.95</b>	<b>Seagrass</b>	<b>Art. Reef</b>				
<i>SpondylIOSoma cantharus</i>	0.00	2.87	19.89	1.30	21.18	21.18
<i>Gobiusculus flavescens</i>	0.77	0.90	9.73	0.98	10.36	31.54
<i>Maja squinado</i>	0.12	1.16	9.64	1.01	10.26	41.80
<i>Atherina presbyter</i>	0.83	0.00	7.46	0.49	7.94	49.74
Ammodytidae	0.66	0.00	5.84	0.40	6.21	55.96
<b>Av. Diss.: 96.25</b>	<b>Seagrass</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.02	2.89	17.23	3.28	17.90	17.90
<i>Trisopterus luscus</i>	0.08	2.33	18.41	2.35	18.57	41.46
<i>Parablennius gattorugine</i>	0.00	1.24	7.53	2.21	7.83	39.57
<i>SpondylIOSoma cantharus</i>	0.00	1.46	7.48	0.87	7.78	47.34
<i>Gobiusculus flavescens</i>	0.77	0.04	5.84	0.86	6.07	53.41
<b>Av. Diss.: 83.94</b>	<b>Kelp</b>	<b>Mussel</b>				
		<b>Beds</b>				
Labridae	0.28	0.00	13.91	17.64	16.57	16.57
<i>Halichoerus grypus</i>	0.09	0.00	11.35	17.64	13.53	30.09
<i>Trisopterus minutus</i>	0.00	1.41	11.35	17.64	13.53	43.62
<i>Merlangius merlangius</i>	0.00	1.21	9.76	3.36	11.63	55.25
Scyliorhinidae	0.36	1.21	9.63	5.34	11.47	66.71
<b>Av. Diss.:90.35</b>	<b>Kelp</b>	<b>Art. Reef</b>				
<i>SpondylIOSoma cantharus</i>	0.00	2.87	21.10	1.30	23.36	23.36
Ammodytidae	1.44	0.00	9.30	0.49	10.30	33.65
<i>Maja squinado</i>	0.46	1.16	8.45	0.93	9.35	43.01
<i>Scyliorhinus canicula</i>	0.46	0.00	6.50	0.78	7.19	50.20
<i>Gobiusculus flavescens</i>	0.07	0.90	6.45	0.71	7.13	57.34
<b>Av. Diss.: 97.87</b>	<b>Kelp</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.07	2.89	17.94	2.92	18.33	18.33
<i>Trisopterus luscus</i>	0.13	2.33	14.79	2.37	15.11	33.43
<i>Parablennius gattorugine</i>	0.00	1.24	7.93	2.15	8.10	41.54

<i>SpondylIOSoma cantharus</i>	0.00	1.46	7.80	0.87	7.97	49.50
Ammodytidae	1.44	0.00	7.60	0.49	7.77	57.27
<b>Av. Diss.: 88.00</b>	<b>Art. Reef</b>	<b>Rocky</b>				
		<b>Reef</b>				
<i>Ctenolabrus rupestris</i>	0.00	2.89	17.88	2.95	20.95	20.32
<i>SpondylIOSoma cantharus</i>	2.87	1.46	14.47	1.32	16.44	36.76
<i>Trisopterus luscus</i>	0.04	2.33	14.17	2.13	16.10	52.87
<i>Parablennius gattorugine</i>	0.00	1.24	7.72	2.01	8.77	61.64
<i>Maja squinado</i>	1.16	0.00	7.18	1.28	8.16	69.80
<b>Av. Diss.: 93.49</b>	<b>Day</b>	<b>Evening</b>				
<i>Atherina presbyter</i>	0.22	0.62	14.40	0.66	15.40	15.40
Ammodytidae	0.23	0.57	6.73	0.37	7.19	22.59
Paguridae	0.25	0.17	6.21	0.43	6.64	29.23
Gobiidae	0.19	0.15	5.91	0.38	6.32	35.56
<i>Scyliorhinus canicula</i>	0.29	0.17	4.90	0.49	5.25	40.80
<b>Av. Diss.: 79.69</b>	<b>Day</b>	<b>Dawn</b>				
<i>Atherina presbyter</i>	0.22	0.17	6.36	0.55	7.99	7.99
<i>Gobiusculus flavescens</i>	0.31	0.92	6.03	0.93	7.57	15.56
<i>Merlangius merlangius</i>	0.16	0.82	5.63	1.10	7.06	22.62
Gobiidae	0.19	0.06	5.61	0.31	7.04	29.66
<i>Scyliorhinus canicula</i>	0.29	0.69	5.16	0.92	6.48	36.14
<b>Av. Diss.: 87.64</b>	<b>Day</b>	<b>Night</b>				
<i>Atherina presbyter</i>	0.22	1.44	10.59	1.05	12.08	12.08
Ammodytidae	0.23	0.75	8.20	0.81	9.35	21.44
Unknown	0.11	1.06	7.05	1.72	8.04	29.48
<i>Merlangius merlangius</i>	0.16	0.63	6.16	0.63	7.03	36.51
<i>Gobiusculus flavescens</i>	0.31	0.23	5.66	0.77	6.46	42.97
<b>Av. Diss.: 86.61</b>	<b>Evening</b>	<b>Dawn</b>				
<i>Gobiusculus flavescens</i>	0.13	0.92	6.08	1.13	7.02	7.02
<i>Atherina presbyter</i>	0.62	0.17	5.93	0.64	6.84	13.86
<i>Trisopterus luscus</i>	0.07	0.74	5.26	1.21	6.08	19.94
<i>Merlangius merlangius</i>	0.19	0.82	5.20	1.17	6.00	25.94
<i>Scyliorhinus canicula</i>	0.17	0.69	5.16	1.23	5.96	31.89
<b>Av. Diss.: 79.51</b>	<b>Evening</b>	<b>Night</b>				
<i>Atherina presbyter</i>	0.62	1.44	8.15	1.12	10.25	10.25
Ammodytidae	0.57	0.75	7.01	0.82	8.82	19.07
<i>Merlangius merlangius</i>	0.19	0.63	7.00	0.70	8.81	27.87
Unknown	0.40	1.06	6.18	1.10	7.77	35.64

Gadidae	0.27	0.53	5.54	0.80	6.97	42.62
<b>Av. Diss.: 85.15</b>	<b>Night</b>	<b>Dawn</b>				
<i>Merlangius merlangius</i>	0.82	0.63	7.80	1.26	9.16	9.16
<i>Atherina presbyter</i>	0.17	1.44	7.16	1.21	8.41	17.57
Unknown	0.00	1.06	7.01	3.03	8.23	25.81
<i>Gobiusculus flavescens</i>	0.92	0.23	5.32	1.18	6.25	32.05
Gadidae	0.15	0.53	4.91	0.89	5.77	37.82

**Habitat Observed x Spring / Neap Tide**

<b>Av. Diss.: 89.82</b>	<b>MCS</b>	<b>Sand</b>				
<i>Scyliorhinus canicula</i>	0.60	0.34	10.31	0.71	11.48	11.48
<i>Pollachius pollachius</i>	0.62	0.05	9.83	0.61	10.94	22.42
Paguridae	0.10	0.53	8.52	0.66	9.49	31.91
Gobiidae	0.32	0.39	8.41	0.61	9.37	41.27
Labridae	0.28	0.00	4.90	0.38	5.45	46.72
<b>Av. Diss.: 89.59</b>	<b>MCS</b>	<b>Seagrass</b>				
<i>Scyliorhinus canicula</i>	0.60	0.40	6.77	0.85	7.56	7.56
<i>Atherina presbyter</i>	0.08	0.62	6.58	0.57	7.34	14.90
<i>Gobiusculus flavescens</i>	0.15	0.72	6.35	0.76	7.09	21.98
<i>Pollachius pollachius</i>	0.62	0.40	6.06	0.70	6.76	28.74
<i>Merlangius merlangius</i>	0.17	0.47	5.03	0.66	5.62	34.36
<b>Av. Diss.: 84.70</b>	<b>MCS</b>	<b>Kelp</b>				
Unknown	0.05	0.64	7.64	1.02	9.02	9.02
Ammodytidae	0.09	1.44	7.55	0.57	8.92	17.94
<i>Atherina presbyter</i>	0.08	0.53	7.39	0.75	8.73	26.67
<i>Scyliorhinus canicula</i>	0.60	0.46	7.26	0.90	8.57	35.24
<i>Pollachius pollachius</i>	0.62	0.28	7.17	0.77	8.47	43.71
<b>Av. Diss.: 95.19</b>	<b>MCS</b>	<b>Mussel Beds</b>				
<i>Trisopterus minutus</i>	0.14	1.41	12.70	5.60	13.34	13.34
<i>Merlangius merlangius</i>	0.17	1.21	10.93	3.53	11.48	24.83
Scyliorhinidae	0.08	1.21	10.27	3.13	10.79	35.62
Octopodidae	0.00	1.00	8.98	5.60	9.44	45.05
Paguridae	0.10	1.00	8.98	5.60	9.44	54.49
<b>Av. Diss.: 97.36</b>	<b>MCS</b>	<b>Midwater</b>				
<i>Pollachius pollachius</i>	0.62	0.10	16.23	0.72	16.67	16.67
<i>Scyliorhinus canicula</i>	0.60	0.22	15.58	0.74	16.00	32.67
Gobiidae	0.32	0.00	8.09	0.47	8.31	40.98



Labridae	0.28	0.00	7.30	0.39	7.50	48.48
<i>Trisopterus luscus</i>	0.24	0.11	6.03	0.46	6.20	54.67
<b>Av. Diss.: 93.06</b>	<b>Sand</b>	<b>Seagrass</b>				
<i>Gobiusculus flavescens</i>	0.00	0.72	6.62	0.70	7.11	7.11
<i>Atherina presbyter</i>	0.03	0.62	6.58	0.48	7.07	14.19
<i>Scyliorhinus canicula</i>	0.34	0.40	5.75	0.74	6.18	20.37
<i>Merlangius merlangius</i>	0.30	0.47	5.73	0.69	6.16	26.53
Paguridae	0.53	0.06	5.05	0.47	5.42	31.95
<b>Av. Diss.: 93.30</b>	<b>Sand</b>	<b>Kelp</b>				
<i>Scyliorhinus canicula</i>	0.34	0.46	9.30	0.83	9.96	9.96
Scyliorhinidae	0.12	0.36	9.13	0.78	9.78	19.75
Ammodytidae	0.00	1.44	8.90	0.52	9.54	29.20
Unknown	0.18	0.64	8.03	0.88	8.61	37.89
Paguridae	0.53	0.00	6.36	0.65	6.82	44.71
<b>Av. Diss.: 87.07</b>	<b>Sand</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.02	1.41	13.83	4.83	15.88	15.88
Scyliorhinidae	0.12	1.21	11.70	4.28	13.44	29.33
<i>Merlangius merlangius</i>	0.30	1.21	11.64	2.87	13.37	42.69
Octopodidae	0.00	1.00	9.78	4.83	11.23	53.92
Unknown	0.18	1.00	8.45	2.08	9.70	63.63
<b>Av. Diss.: 99.07</b>	<b>Sand</b>	<b>Midwater</b>				
Paguridae	0.53	0.00	16.50	0.69	16.65	16.65
Gobiidae	0.39	0.00	10.89	0.48	10.99	27.65
<i>Mustelus mustelus</i>	0.18	0.00	9.95	0.44	10.04	37.69
<i>Scyliorhinus canicula</i>	0.34	0.22	8.77	0.55	8.85	46.54
<i>Merlangius merlangius</i>	0.30	0.34	7.46	0.54	7.53	54.07
<b>Av. Diss.: 93.06</b>	<b>Seagrass</b>	<b>Kelp</b>				
Ammodytidae	0.29	1.44	14.74	0.74	15.83	15.83
<i>Atherina presbyter</i>	0.62	0.53	5.93	0.58	6.37	22.20
<i>Pomatoschistus minutus</i>	0.02	0.43	5.83	1.04	6.26	28.46
<i>Maja squinado</i>	0.13	0.46	5.37	1.08	5.77	34.23
<i>Gobiusculus flavescens</i>	0.72	0.07	5.17	0.74	5.55	39.78
<b>Av. Diss.: 95.30</b>	<b>Seagrass</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.08	1.41	0.41	4.04	10.93	10.93
Scyliorhinidae	0.15	1.21	8.83	3.59	9.26	20.19
<i>Merlangius merlangius</i>	0.47	1.21	8.44	2.30	8.86	29.05

Octopodidae	0.00	1.00	7.36	4.04	7.73	36.78
Paguridae	0.06	1.00	7.36	4.04	7.73	44.51
<b>Av. Diss.: 95.72</b>	<b>Seagrass</b>	<b>Midwater</b>				
<i>Atherina presbyter</i>	0.62	0.00	7.86	0.43	8.21	8.21
<i>Gobiusculus flavescens</i>	0.72	0.04	6.97	0.63	7.28	15.49
<i>Merlangius merlangius</i>	0.47	0.34	6.76	0.59	7.06	22.56
<i>Scyliorhinus canicula</i>	0.40	0.22	5.90	0.57	6.16	28.72
<i>Pomatoschistus microps</i>	0.15	0.11	5.11	0.33	5.34	34.06
<b>Av. Diss.: 81.82</b>	<b>Kelp</b>	<b>Mussel</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	0.00	1.41	9.78	7.05	11.96	11.96
<i>Merlangius merlangius</i>	0.00	1.21	8.40	3.91	10.27	22.22
<i>Atherina presbyter</i>	0.53	0.00	7.32	1.09	8.94	31.16
Octopodidae	0.00	1.00	6.92	7.05	8.45	39.62
Paguridae	0.00	1.00	6.92	7.05	8.45	48.07
<b>Av. Diss.: 97.95</b>	<b>Kelp</b>	<b>Midwater</b>				
<i>Scyliorhinus canicula</i>	0.46	0.22	13.77	0.86	14.06	14.06
Scyliorhinidae	0.36	0.00	13.18	0.77	13.45	27.51
Unknown	0.64	0.00	11.42	1.03	11.66	39.17
<i>Atherina presbyter</i>	0.53	0.00	7.89	0.63	8.06	47.23
Ammodytidae	1.44	0.00	7.32	0.50	7.47	54.71
<b>Av. Diss.: 95.40</b>	<b>Mussel</b>	<b>Midwater</b>				
		<b>Beds</b>				
<i>Trisopterus minutus</i>	1.41	0.03	14.48	3.57	15.18	15.18
<i>Merlangius merlangius</i>	1.21	0.34	13.24	3.52	13.88	29.05
Scyliorhinidae	1.21	0.00	12.24	3.35	12.83	41.89
Octopodidae	1.00	0.00	10.24	3.57	10.73	52.62
Paguridae	1.00	0.00	10.24	3.57	10.73	63.35
<b>Av. Diss.: 88.96</b>	<b>Spring</b>	<b>Neap</b>				
Paguridae	0.29	0.34	13.20	0.70	14.83	4.83
Gobiidae	0.08	0.42	9.28	0.54	10.43	25.26
<i>Scyliorhinus canicula</i>	0.49	0.25	8.90	0.65	10.00	35.26
<i>Mustelus mustelus</i>	0.05	0.15	7.43	0.43	8.35	43.62
Unknown	0.26	0.18	6.58	0.50	7.39	51.01
<b>Av. Diss.: 90.08</b>	<b>Spring</b>	<b>Mid</b>				
<i>Scyliorhinus canicula</i>	0.49	0.42	7.89	0.75	8.76	8.76
<i>Merlangius merlangius</i>	0.31	0.42	6.89	0.75	7.64	16.40
Paguridae	0.29	0.12	6.23	0.41	6.92	23.32

<i>Atherina presbyter</i>	0.14	0.46	4.96	0.46	5.51	28.83
Gobiidae	0.08	0.13	4.64	0.29	5.15	33.98
<b>Av. Diss.: 90.36</b>	<b>Neap</b>	<b>Mid</b>				
Gobiidae	0.42	0.13	9.21	0.51	10.19	10.19
Paguridae	0.34	0.12	8.23	0.53	9.11	19.30
<i>Merlangius merlangius</i>	0.24	0.42	7.07	0.73	7.82	27.13
<i>Scyliorhinus canicula</i>	0.25	0.42	6.91	0.73	7.65	34.78
Scyliorhinidae	0.15	0.16	5.93	0.70	6.56	41.34

---