

Review

Redox-Based Mechanisms of O₂ Sensing in Hypoxic Pulmonary Vasoconstriction: Where Are We Now?

Philip I. Aaronson ^{1,*}, Jeremy P. T. Ward ², Asuncion Rocher ^{3,4,*} and Jesus Prieto-Lloret ^{3,4}

¹ Department of Inflammation Biology, School of Immunology and Microbial Sciences, Faculty of Life Sciences and Medicine, King's College London, London SE1 9RT, UK

² King's Centre for Lung Health, School of Immunology and Microbial Sciences, King's College London, London SE1 9RT, UK

³ Unidad de Excelencia Instituto de Biomedicina y Genética Molecular (IBGM), Universidad de Valladolid-CSIC, 47005 Valladolid, Spain

⁴ Departamento de Bioquímica y Biología Molecular y Fisiología, Universidad de Valladolid, 47004 Valladolid, Spain

* Correspondence: paaronson987@gmail.com (P.I.A.); asun.rocher@uva.es (A.R.)

Abstract

Hypoxic pulmonary vasoconstriction (HPV) is a rapid and reversible constrictor response of the pulmonary vasculature, and especially its small muscular precapillary arteries, which is initiated by episodes of local alveolar hypoxia. Acting as a protective homeostatic vasomotor mechanism, HPV enables maximal gas exchange by diverting blood from poorly ventilated alveoli into those rich in oxygen, thereby optimizing oxygen uptake and the ventilation–perfusion (V/Q) ratio so as to maintain the arterial oxygen partial pressure (P_aO₂) within the physiological range. HPV is an intrinsic mechanism of pulmonary artery smooth muscle cells (PASMCs), and requires an O₂ sensor which acts through mediator(s) to trigger effector mechanisms within these cells to evoke constriction. Whereas HPV effector mechanisms are reasonably well defined, the nature of the O₂ sensor and mediators remains in dispute, and a number of proposals have been developed to account for these. Some (but not all) of these share a focus on the concept that hypoxia activates effector mechanisms by inducing a change in the PASMC cytoplasmic redox state. Of these, the Redox Theory, first proposed by Kenneth Weir and Stephen Archer in 1995, proposes that hypoxia inhibits mitochondrial production of reactive oxygen species (ROS), thereby causing the cytoplasm to become more reduced. This inhibits ongoing vasorelaxation maintained by the opening of voltage-gated K⁺ channels. In contrast, according to the Mitochondrial ROS hypothesis, introduced by Paul Schumacker and Naveen Chandel in 2001, hypoxia increases mitochondrial ROS production, causing an oxidizing shift in the cytoplasmic redox state that activates several vasoconstricting pathways. In a third redox-based scenario, developed by Michael Wolin and Sachin Gupte, hypoxia evokes contraction by causing a fall in H₂O₂ production by NADPH oxidase and by activating the pentose phosphate pathway. These effects inhibit basal vasorelaxation maintained by the guanylate cyclase and protein kinase G and also stimulate vasoconstricting mechanisms. In this comprehensive review, we first provide a detailed summary of the key studies contributing to the development of these proposals and then subject the evidence supporting them to a critical appraisal, based in part on how well they accord with the wider literature and recent developments in our understanding of how cells shape and deploy redox mechanisms in order to regulate cell function.

Academic Editor: John T. Hancock

Received: 20 December 2025

Revised: 3 February 2026

Accepted: 15 February 2026

Published: 22 February 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

Keywords: O₂ sensing; hypoxia; hypoxic pulmonary vasoconstriction; K_v channel; mitochondria; NADPH oxidase; reactive oxygen species; redox; hydrogen peroxide; NADH

1. Introduction: Definition and Description of HPV

1.1. Basic Properties of HPV

Hypoxic pulmonary vasoconstriction (HPV) is a rapid and reversible constrictor response of the pulmonary vasculature, and especially its small muscular precapillary arteries, to episodes of local alveolar hypoxia [1–3]. Acting as a protective vasomotor mechanism which enables maximal gas exchange by diverting blood from poorly ventilated alveoli into those rich in oxygen content on a breath-to-breath basis, HPV optimizes oxygen uptake and the ventilation to perfusion (V/Q) ratio, thereby maintaining the arterial oxygen partial pressure (PO₂) within the physiological range [4]. In parallel with HPV, acute hypoxia provokes systemic arterial vasodilation, thereby promoting oxygen delivery to body tissues by increasing blood flow [5].

Studies in isolated perfused whole lungs or lung lobes from a range of species indicate that HPV begins to develop when PO₂ falls below ~80 Torr, becomes half-maximal at ~50 Torr, and reaches its maximum amplitude at ~20 Torr [3]. Depending on the experimental model and conditions used, HPV manifests as a rapidly developing and sustained increase in pulmonary artery pressure (PAP) or pulmonary vascular constriction, which in many studies exhibits a biphasic profile in which an initial transient development of force (Phase 1) is superimposed upon or followed by a slow and progressive increase in tension (Phase 2). The biphasic time course is typically observed when more severe levels of hypoxia are used to evoke the response [3]. In humans, the HPV evoked when end-tidal PO₂ (a measure of the alveolar PO₂) was halved demonstrated an initial rapid increase in pulmonary vascular resistance which reached a steady state within minutes. This was followed by a second slower increase that began after 30–60 min and plateaued after ~3 h [6,7]. HPV is a function of the PO₂ in both the alveoli and the blood perfusing the pulmonary arteries (PAs), with the former being of greater importance [8]. It does not require, but is modulated by, the autonomic nervous system and diverse vasoactive factors [3,9].

The mechanisms responsible for the two phases of HPV appear to be, at least to some extent, different [10,11]. Phase 1 is mediated by an increase in the cytoplasmic Ca²⁺ concentration ([Ca²⁺]_{cyt}) due to a rapid decrease in K⁺ channel activity, leading to membrane depolarization and opening of L-type voltage-gated Ca²⁺ channels [10,12], as well as release of Ca²⁺ from the sarcoplasmic reticulum and the resulting stimulation of store-operated Ca²⁺ entry (SOCE) [10,13,14] in pulmonary artery smooth muscle cells (PASMCS). Most studies of HPV in isolated perfused lungs or PA have used short periods of hypoxia (≤15 min), suggesting that their conclusions apply mainly to Phase 1. Studies of O₂ sensing in isolated or cultured PASMCS have also typically employed short hypoxic challenges, but the extent to which their results are representative of one or the other phase of HPV in more intact systems is unclear.

Phase 2 HPV is less well characterized but probably involves multiple pathways contributing to an increase in [Ca²⁺]_{cyt} as well as rho kinase-mediated Ca²⁺ sensitization [15]. Unlike Phase 1, sustained HPV has generally (but not always) been found to be endothelium-dependent [16]. The evidence for an involvement of membrane depolarization in Phase 2 HPV is mixed [10,14,17,18]. Both phases are reversible, but if global hypoxia lasts longer than 24 h, persistent HPV of the entire pulmonary circulation causes a sustained increase in PAP. This, together with phenotypical, biochemical, and functional changes in

each pulmonary vascular cell type, gradually leads to pulmonary vascular remodeling, which can be irreversible.

1.2. Proposed Models of O₂ Sensing in HPV

HPV is conceptualized as requiring an O₂ sensor, which is coupled through mediator(s) to effector mechanisms that evoke constriction by raising PASM C [Ca²⁺]_{cyt} and/or causing Ca²⁺ sensitization [19]. It is generally (but not universally [20]) thought that all of these components of the response reside within the PASM C, although extrinsic influences (e.g., endothelial factors) can modulate HPV [3].

This review focuses on describing the three well-established proposals for O₂ sensing in PASM C in which ROS play pivotal roles. These theories are variations on a common theme: the concept that the mitochondria and/or NADPH oxidase (Nox) are O₂ sensors that modify the function of contractile effectors to cause HPV by altering the concentration of cytoplasmic reactive oxygen species (ROS) or the state of cellular redox couples.

1. According to the *Redox Theory*, introduced by Kenneth Weir and Stephen Archer in 1995 [21], hypoxia decreases the production of ROS by the mitochondrial electron transport chain (ETC), leading to a fall in [ROS] and/or the reduction of redox couples in the cytoplasm. This evokes the closure of voltage-gated K⁺ (K_v) channels, which are activated by basal ROS production under normoxic conditions, thus causing membrane depolarization, the opening of voltage-gated Ca²⁺ channels (VGCC), and therefore contraction. Although the Redox Theory stresses the importance of K_v channels, there are also other effectors which could potentially respond to a fall in ROS or a reduction of cytoplasmic redox couples in such a way as to cause HPV. For example, there is evidence that the reduction of Cys42 on protein kinase G α1 (PKG1α), which may exert a tonic vasorelaxing influence on PASM C, diminishes its activity [22].
2. A proposal which we will refer to as the *Mitochondrial ROS hypothesis*, developed by Paul Schumacker, Naveen Chandal, and colleagues and first described in 2001 [23], proposes that hypoxia causes an increase in mitochondrial ROS production, leading to a higher ROS concentration and/or the oxidation of redox couples, in the cytoplasm. This hypoxia-induced rise in cytoplasmic ROS may be supplemented by a PKCε-mediated stimulation of Nox, which is triggered by the mitochondrial ROS [24]. The rise in cytoplasmic [ROS] might evoke contraction through multiple effector pathways, potentially including Ca²⁺ release from the sarcoplasmic reticulum, an increase in store-operated Ca²⁺ influx, and RhoA/Rho kinase-mediated Ca²⁺ sensitization [13–15].
3. The processes responsible for O₂ sensing and HPV have also been the subject of an extensive series of papers by the laboratories of Michael Wolin and Sachin Gupte (see [25] for a review). In agreement with the Redox Theory, these authors propose that hypoxia causes contraction by removing a normoxic vasodilating influence, although this is seen as being maintained largely by oxidation-induced activation of soluble guanylate cyclase and protein kinase G (PKG1α) rather than by the opening of K_v channels. In addition, they have presented evidence that hypoxia activates the pentose phosphate pathway, thereby increasing the production of nicotinamide adenine dinucleotide phosphate (NADPH), and that this contributes to the inhibition of PKG1α and also activates other Ca²⁺-dependent and -independent contractile mechanisms.

Two additional O₂-sensing mechanisms have been suggested to bring about HPV. According to the *AMP kinase* (adenosine monophosphate kinase, AMPK) hypothesis, introduced by Mark Evans and co-workers [26,27], HPV is caused by a small but meaningful

inhibition of mitochondrial respiration, which results in an increase in the cellular [AMP]/[ATP] ratio. This activates AMP kinase, which causes PASMC contraction through several mechanisms, the best defined of which is phosphorylation-induced inhibition of Kv1.5 channels, leading to membrane depolarization and Ca^{2+} influx [26,27]. Furthermore, Kenneth Olson's laboratory [28–31] has suggested that O_2 -dependent metabolism of hydrogen sulfide (H_2S ; aka sulfide) constitutes an O_2 sensor in many tissues, including pulmonary arteries and carotid body chemoreceptor cells (CBCC). According to this proposal, hypoxia suppresses the oxidation of sulfide, causing increased cellular concentrations of sulfide and/or reactive sulfur species. These activate effector mechanisms which cause, for example, HPV. Neither of these hypotheses, as proposed by their originators, incorporates the involvement of ROS. However, a possible role for ROS in both schemes is possible, as set out in Sections 6.2 and 6.3. We do not, however, cover any other aspects of these proposals.

Further redox-independent O_2 -sensing mechanisms have also been proposed to be important in triggering HPV [32,33] and also in evoking acute responses to hypoxia in other types of cells (e.g., in CBCC [34–36]). These are not discussed in this review. We also do not cover either the extensive set of known or putative HPV effectors, which have been described in detail elsewhere [3,37,38] (except to discuss those which are proposed to be regulated by redox mechanisms where appropriate), or the role of ROS/redox mechanisms in pulmonary hypertension resulting from chronic pulmonary hypoxia (see, e.g., [39–47] for reviews of this subject).

We note that the term 'reactive oxygen species' has been criticized in light of evidence that the various types of ROS exert their own specific actions, typically on different sets of biomolecules, with 'oxidants' suggested to be a better alternative when the relevant oxidizing species has not been identified [48]. However, to be consistent with other reviews of O_2 sensing in HPV [37,49–51], and because the term 'oxidants' also encompasses reactive sulfur species, we prefer to refer to changes in superoxide/ H_2O_2 production as increases or decreases in ROS, except where the specific use of either term is more appropriate.

1.3. Analytical Framework and Literature Search Strategy

The analytical framework we employed in designing this review was to first decide on which subjects we would cover in the review (in essence the three hypotheses described above, as well as supportive information about redox biology). This was based on our pre-existing knowledge of the literature pertaining to the role of cell redox mechanisms in O_2 sensing in HPV, which suggested that the vast bulk of this literature was linked to one or more of these proposals, and that this situation had emerged from observations by different laboratories, which were often contradictory. This had resulted in a literature characterized by deep and unresolved controversies. Our further inspection of this literature, supplemented by additional papers identified via the search strategy described below, confirmed the validity of these suppositions. We therefore felt that it would be useful to systematically collate and critically examine these observations and the hypotheses which were formulated from them, in an attempt to assess their strengths and weaknesses, especially in light of important developments in the larger literature on redox biology which have occurred since many of the findings underpinning these proposals were made. We have attempted in this review to carry out this analysis in an unbiased manner. However, because we believe that it is important that readers should be able to make up their own minds about the quality of the data underpinning these hypotheses, and also wanted to provide the context in which they developed, we precede our analyses of these hypotheses with comprehensive, detailed, and chronologically ordered accounts of the investigations which contributed to their evolution. We also include introductory

sections which provide summaries of current thinking about relevant aspects of the larger redox literature, including subjects such as the regulation of mitochondrial ROS production and the mechanisms governing the control of the intracellular H_2O_2 concentration. We direct readers who are looking for a more succinct coverage of this literature to a number of excellent previous reviews [3,19,37,41,43,49–57].

With regard to our literature search strategy, in order to identify relevant papers from the HPV literature on ROS/redox-dependent O_2 sensing, we initially searched PubMed using combinations of the search terms ‘hypoxia’, ‘hypoxic pulmonary vasoconstriction’, ‘pulmonary artery’, ‘reactive oxygen species’, ‘hydrogen peroxide’, ‘superoxide’, ‘oxygen sensing’, ‘peroxynitrite’, ‘NADH’, ‘NADPH’, ‘redox’, ‘NADPH oxidase’ and ‘mitochondria’. Initial PubMed searches were also carried out using relevant author names. Additional references were then drawn from the bibliographies of the papers we had identified. Papers covering relevant but non-core topics for which an exhaustive discussion of the literature was not necessary were identified using appropriate search terms, with further references sourced from bibliographies and PubMed searches using author names when appropriate. On a few occasions where the use of search terms was not productive, we interrogated Google/Gemini using questions (e.g., ‘what is the extracellular H_2O_2 concentration?’). We then consulted the references, which were provided in replies. No information was taken directly from Google/Gemini responses, and no references provided in these responses were cited unless we found the paper using PubMed and examined it.

1.4. Normoxia, Physoxia, and Hypoxia

The O_2 concentrations (18–21% O_2) which are used almost invariably in physiological experiments to simulate ‘normoxia’ are substantially higher than ‘physoxic’ concentrations (i.e., those experienced under baseline conditions by cells *in vivo*) [58]. There is a growing literature [59–61] attesting to how this can distort many aspects of cell phenotype. Additional problems inherent in the routine use of effectively hyperoxic conditions in studies of acute O_2 sensing have been discussed in excellent reviews by Alva et al. [62] and Olson [30], the former of which also presents an authoritative and perceptive discussion of many of the other issues around the role of ROS in O_2 sensing, which we cover in this paper. The intra-alveolar PO_2 is ~13.3%, and that in arterial blood averages 12.3%. Measurements of lung tissue PO_2 in anesthetized animals breathing room air have yielded lower values. For example, O_2 electrodes placed in peri-bronchial lymph nodes in dogs reported PO_2 levels of ~53 Torr [63] and ~38 Torr [64], and an electron paramagnetic resonance sensor inserted into lung tissue of rats also detected a PO_2 of ~38 Torr, which fell to ~26 Torr in animals breathing 10% O_2 [65]. It is not clear why the PO_2 in lung tissue is so much lower than it is in the alveoli, but one can speculate that this could be due to the influence of the PO_2 in the blood within PAs, which enters the lung with a mixed venous PO_2 of ~40 Torr. Importantly, it was shown in anesthetized dogs that the mixed venous PO_2 exerts a marked influence on HPV [8], presumably because the blood within a substantial fraction of the PA responsible for HPV has not yet been fully oxygenated and is affecting the PO_2 within the vascular wall. The effective PO_2 stimulus for HPV (P_{sO_2}) in this study was calculated to be determined by the alveolar and mixed venous O_2 concentrations according to the equation: $\text{P}_{\text{sO}_2} = (\text{P}_{\text{AO}_2}^{0.59} \times \text{P}_{\text{mvO}_2}^{0.41})$. A similar weighting was observed in an isolated perfused rat lung [66]. In this case, the P_{sO_2} *in vivo* when breathing room air would be roughly 70 Torr, suggesting that this might be an appropriate PO_2 to use in experiments to simulate physoxia under normoxic conditions when conducting studies of HPV. The mitochondria and Noxs, which are likely to be the two main sources of superoxide/hydroperoxide in PASMC, produce ROS in a manner which is linearly (in mitochondria, at least when they are isolated [67]) or hyperbolically (Nox [62,68]) dependent on the PO_2 . For example, Nox4, which is thought to be constitutively active and therefore

could be an important source of H_2O_2 in PASMC under basal conditions [69], has a $K_m\text{O}_2$ of ~136 Torr [68]. Thus, PASMC and other cells in preparations used to study O_2 sensing in HPV are likely to have been undergoing a degree of oxidative stress under experimental ‘normoxic’ conditions.

A related issue is that the O_2 concentration experienced by cultured cells in standard O_2 -controlled incubators is usually supra-physoxic when the PO_2 is set to 18%. Even so, under certain conditions (e.g., high cell density), the PO_2 in the medium can be much lower than the physoxic level due to cellular O_2 consumption [70]. Cellular respiration can have an especially marked effect on the pericellular PO_2 in cells in culture [71], an effect which has been shown to be inhibited by blocking of the ETC [72]. This is a concern because experimental interventions used to study hypoxia often decrease the activity of the ETC, in which case they may also increase the pericellular PO_2 and therefore diminish the degree of hypoxia experienced by cells. Unfortunately, the extent to which these factors might have had an impact on the results of the investigations we will discuss is unclear.

Hypoxia can be defined as a fall in the PO_2 , which results in a limitation of tissue function [73]. However, the biological effects of a fall in PO_2 are strongly context-dependent. We suggest that a definition of hypoxia which better applies to HPV is that it constitutes a reduction in PO_2 below its normal physiological value, which is sufficient in a particular context/tissue/cell to elicit a response, whether it be HPV, carotid body signaling, changes in gene transcription, or impaired energy production. The question of the range of levels of hypoxia which are appropriate for studying HPV has probably not been taken as seriously as it should have been in some papers. In many cases, for example, the level of hypoxia is defined as the % O_2 content in the gas mixture used to bubble the solution or in the incubator housing the cell chamber, with no information provided about the PO_2 in the solution. In an influential review of HPV, Moudgil et al. [19] stressed the necessity of using physiological levels of hypoxia in studies of HPV, suggesting that the level of hypoxia at the summit of Mount Everest (8488 m; alveolar and arterial PO_2 of ~43 Torr recorded during the Operation Everest II expedition [74]) might be an appropriate guideline, since the PO_2 in the small PAs, which are the main site of HPV, should be very close to that in the alveoli.

On the other hand, if one takes into account the influence of the mixed venous PO_2 on HPV as described above, and factors in the resting $P_{mv}\text{O}_2$ (22 Torr) observed in another study by the Everest II expedition team [75] at a simulated altitude of 8488 m, the $P_{s}\text{O}_2$ for HPV at this height would be ~33 Torr. Moreover, other studies have indicated that elevation has a larger effect on the arterial and alveolar PO_2 ; for example, $P_{A}\text{O}_2$ and $P_{mv}\text{O}_2$ values obtained at 8488 m during an earlier expedition to Everest were 35 and 21 Torr, respectively [76], and Grocott et al. [77] obtained arterial PO_2 values of 48, 42, 34, and 24 Torr at elevations of 5300, 6400, 7100, and 8400 m, with an alveolar PO_2 of 30 Torr at 8400 m, although the corresponding $P_{mv}\text{O}_2$ levels were not reported. Taking into account that even moderate levels of activity further lower the $P_{mv}\text{O}_2$ at altitude [75], it can be argued that lower levels of hypoxia, perhaps down to those approximating the PO_2 range at which HPV reaches its maximum amplitude (~20 Torr), are also suitable for studying physiologically relevant O_2 -sensing mechanisms in HPV. Nonetheless, since the mechanisms responsible for O_2 sensing in PA may well vary according to the level of hypoxia, as is thought to occur in chemoreceptor cells of the carotid body (CBCC) [78], ideally, the mechanisms of O_2 sensing in HPV should be assessed over a range of low PO_2 levels. This has not been done in any organized way.

2. Reactive Oxygen Species as Signaling Molecules

2.1. ROS Definition and Function

ROS include oxygen radicals such as superoxide (O_2^-), hydroxyl ($\cdot OH$) and peroxy ($RO_2\cdot$), as well as oxidants such as hydrogen peroxide (H_2O_2) and hypochlorous acid ($HClO$). These are formed by cell metabolism and can oxidize biological molecules, although their reactivity varies between species [79,80]. O_2^- , the first species formed by single- or multi-enzyme reactions through the single-electron reduction of O_2 , rapidly dismutates to H_2O_2 , a process which occurs spontaneously and is also catalyzed by superoxide dismutase (SOD). $\cdot OH$, which can be formed by the reaction of H_2O_2 with Fe^{2+} (Fenton reaction) or of water with Fe^{3+} [81], is extremely reactive and is scavenged immediately upon its production. H_2O_2 is relatively stable and is able to cross membranes via aquaporins [82], properties which allow it to act as a signaling molecule. O_2^- can also react with nitric oxide (NO) to form the reactive nitrogen species peroxynitrite ($ONOO^-$), which has a signaling function [83]. The two cellular sources of ROS which have received the most attention with regard to O_2 sensing and HPV are the ETC and Nox.

ROS can cause oxidative modifications of nucleic acids, carbohydrates, lipids and proteins. Their excessive production therefore results in deleterious effects on cells, a situation referred to as *oxidative stress*. However, cells use smaller spatiotemporally controlled increases in H_2O_2 to regulate their normal function and initiate adaptive responses (oxidative *eustress*) [84], with HPV being a possible example.

2.2. H_2O_2 Signaling

H_2O_2 signaling depends on its ability to react with thiolates (S^-), the anionic form of cysteine sulfhydryl ($-SH$ groups), leading to modifications of protein structure and therefore function (Figure 1). Since the pK_a value of the thiol in free cysteine is ~ 8.5 , the fraction of cysteines in the reactive thiolate form should be small at the physiological cytoplasmic pH (e.g., ~ 7.3 in PASMCM, [85]). However, the pK_a of cysteine thiols can be lowered by their proximity to positively charged amino acids [86] and by other factors related to their local environment within the protein structure [87,88]. Thus, a fraction of cysteine residues are able to react with H_2O_2 and other oxidizing species, including reactive nitrogen or sulfur species, which cause S-nitrosation and S-sulfhydration, respectively. Reactive cysteines can also undergo S-thiolation, most importantly by oxidized glutathione (GSSG), causing S-glutathionylation. It is thought that disulfide bond formation and S-glutathionylation are the most important oxidative modifications for H_2O_2 signaling [89]. These modifications can alter three-dimensional protein structure, leading to protein translocation [90,91] or substrate targeting [90,92], ultimately causing rapid functional changes that can induce or fine-tune cell signaling during acute responses.




Figure 1. Oxidative modifications of reactive cysteines. Deprotonated cysteine thiols (thiolates) can react with a variety of oxidizing species, leading to post-translational modifications, which can affect protein function. Reactions of thiolates with glutathione (GSH), reactive sulfur species, and nitric oxide (NO) lead to their (a) glutathionylation, (b) persulfidation, and (c) nitrosylation, which can further lead (d) to thiol exchange reactions such as glutathionylation or the formation of an intra- or inter-protein disulfide bond. Thiolate oxidation by H₂O₂ can also lead to the formation of intra- or inter-protein disulfide bonds (e and f, respectively). The reaction of H₂O₂ with thiolates can also produce sulfenic acid (g). Although not shown in the diagram, cysteine sulfenic acid can react with GSH, reactive sulfur species, and NO, leading to its glutathionylation, persulfidation, and nitrosylation respectively, and can also react with thiols to form disulfide bonds [93]. Notably, cysteine sulfenic acid can be formed (j) by the oxidation of thiols with peroxynitrite (ONOO⁻), or by the reaction of peroxynitrous acid (ONOOH) with thiolates. These oxidative modifications can be reversed by cellular antioxidant mechanisms. Under more highly oxidizing conditions, sulfenic acid is further converted to (h) sulfinic acid and then (i) sulfonic acid; these reactions are largely irreversible. The types of oxidative modifications of protein thiolates depicted in green have been shown to alter the activities of many proteins (e.g., [94]).

In order to minimize its potential damaging effects on cells, the basal cytoplasmic concentration of H₂O₂ is maintained in the low nanomolar range by the thioredoxin (Trx) and glutaredoxin (Grx) antioxidant systems, acting in concert with peroxiredoxins (Prxs), and Grx-dependent glutathione peroxidases (Gpxs) (Figure 2) [95–97]. These systems, which are also largely responsible for terminating redox signaling by reversing oxidative thiol modifications, ultimately require the oxidation of NADPH, which is regenerated from NADP⁺ by the pentose phosphate pathway and other enzymatic mechanisms [98]. The Prx family, comprising six isoforms which are differentially expressed in the various cell compartments, is generally thought to be particularly important for the removal of H₂O₂; it was estimated, for example, that 78% and 21% of H₂O₂ scavenging in cells is dependent on Prx and Gpx, respectively [99]. Intriguingly, however, a recent study using the novel NADPH sensor NAPStar3b has shown that the glutathione system plays the predominant role in antioxidant defense in HEK293 cells [100]. H₂O₂ is also scavenged by

the enzyme catalase, which is located mainly but not exclusively in peroxisomes. It is believed that the antioxidant effect of catalase, which does not require cofactors, is more important at higher H_2O_2 concentrations, at which the activities of Prx and Gpx may be limited by the necessity for their recycling by Trx and Grx [101].

The extent to which the direct reactions between H_2O_2 and protein thiols described above can account for redox signaling has been questioned on kinetic grounds. Scavenging of H_2O_2 by Prx is rapid, with a 2nd order rate constant in the range of 10^5 – 10^8 M^{-1} [99,102], whereas H_2O_2 reacts much more slowly with cysteine thiols ($k \approx 10^1$ – 10^2 M^{-1}).




Figure 2. Antioxidant mechanisms. Superoxide spontaneously dismutates to H_2O_2 at a high rate, but this process is reinforced by cellular (and extracellular) SODs. Once formed H_2O_2 is thought to have a half-life of $\sim 10 \text{ s}$. It is reduced to H_2O , most importantly by peroxiredoxins (Prxs), with the oxidizing equivalent then being passed ultimately to NADPH via thioredoxin (Trx) and thioredoxin reductase (TrxR). H_2O_2 is also reduced by glutathione peroxidase, which also passes the oxidizing equivalent to NADPH, in this case via glutathione peroxidase, glutathione (GSH) and glutathione reductase (Gr). NADP^+ is reduced back to NADPH by several enzymes, including malic enzyme (ME) and isocitrate dehydrogenase (IDH) in the mitochondria and by glucose-6-phosphate-dehydrogenase (G-6-PD) and also 6-phosphogluconate dehydrogenase in the cytoplasm. NADP^+ reduction to NADPH in the mitochondria is also coupled to the oxidation of NADH to NAD^+ by the NAM nucleotide transhydrogenase in the inner mitochondrial membrane, a reaction which is driven by the protein gradient across this membrane (see Section 3.1.1). These processes ensure that the $\text{NADP}^+/\text{NADPH}$ redox couple is very highly reduced in both cytoplasm and mitochondria [98], thereby maintaining the effectiveness of the NADPH-requiring antioxidant systems. At higher concentrations, H_2O_2 is also scavenged by catalase, an enzyme mainly expressed in lysosomes. Moreover, Prxs are present in micromolar concentrations in cells, while H_2O_2 concentrations probably remain submicromolar during signaling [84]. Consequently, antioxidant mechanisms are predicted to scavenge H_2O_2 before it can react with thiols on target proteins, unless these are very near to its site of production.

Despite these constraints, there is evidence that cells can use H_2O_2 as a highly specific signaling molecule if it is formed very close to its target protein (e.g., [103]). Longer range specific targeting can also be achieved by way of the formation of thiol peroxidase-based redox relays (Figure 3) in which Prx (or Gpx) is used to convey oxidizing equivalents from H_2O_2 to a target protein, rather than sending them down an antioxidant pathway to be

consumed by NADPH [104,105]. In a one-step relay, once oxidized by H_2O_2 , Prx oxidizes the target protein rather than Trx. In a two-step relay, Prx oxidizes Trx, which then oxidizes the target protein rather than oxidizing Trx reductase (TrxR). The fact that Prx is able to act in both antioxidant and peroxidase capacities can reconcile the apparent paradox that H_2O_2 , despite being subject to swift and efficient scavenging by cellular antioxidant pathways, can initiate the rapid and reversible oxidation of relatively unreactive thiols. The initial reaction between H_2O_2 and Prxs is very fast, with rate constants in the range of 10^7 – $10^8 \text{ M}^{-1} \text{ s}^{-1}$ for all five isoforms; similar rate constants are also seen for the reaction of Prxs and peroxynitrite [106,107]. The factors which determine the balance between the antioxidant and oxidant/signaling activities of Prxs are still being worked out. One possibility described in an elegant study by Portillo-Ledesma et al. [106] is that the antioxidant capacity of Prx becomes saturated at a certain threshold concentration of oxidant (e.g., H_2O_2), above which it can oxidize target proteins. They showed that this threshold is particularly low (4 nM H_2O_2) for the cytoplasmic isoform Prx2, suggesting that it can oxidize target protein at relatively low H_2O_2 concentrations while the antioxidant activity of Prx isoforms with higher thresholds (e.g., Prx1) continues. The activity of Prx is also regulated by several post-translational modifications [108].

The specificity of Prx-dependent redox relays is fostered by the rapid association of Prxs, once they are oxidized, with discrete sets of target proteins. This occurs through the formation of disulfide bonds at its active site, which happens during target protein oxidation. For example, in experiments carried out in HEK293T cells, Prx1 and Prx2, which are both expressed in the cytoplasm, were shown to associate with 735 and 165 proteins, respectively [109]. Of these, only 53 bound to both isoforms. This highlights the specificity of these interactions, which was shown to depend on the local amino acid sequence around the target cysteine. Of the 1233 proteins found to bind to one or more of the five Prx isoforms expressed in these cells, ~80% had previously been identified [86] as harboring redox-sensitive cysteines, suggesting that Prx-dependent oxidative modifications of protein thiols probably play a widespread role in H_2O_2 -signaling. It has also been proposed that redox relays which utilize Gpx and GSSG to cause S-glutathionylation of target cysteines are important in the redox regulation of many proteins, especially within mitochondria [110].




Figure 3. Redox relays. Rather than oxidizing reactive cysteines directly, H_2O_2 generally does so indirectly via redox relays, which transfer oxidizing equivalents from H_2O_2 (and also peroxynitrite) to signaling proteins (SP). Prxs can oxidize signaling proteins directly or can do so by passing an oxidizing equivalent to Trx, which is generally responsible for restoring several Prx isoforms to their reduced state following their oxidation. Oxidizing equivalents can also be passed from H_2O_2 to signaling proteins via glutathione peroxidase [104].

3. Cellular Mechanisms of ROS Production

Mitochondrial complexes, together with a microsomal ETC, the Noxs, and other oxidases (e.g., glucose oxidase, xanthine oxidase), peroxisomes, and cyclooxygenases, comprise the main biological machinery that directly produces or is causally involved in oxidant formation [111]. Of these various sources of ROS, research on HPV has focused only on the potential roles of mitochondria and Nox.

3.1. Mitochondrial Regulation of Cytoplasmic H_2O_2

Although mitochondria can release H_2O_2 into the cytoplasm, e.g., during ischemia–reperfusion [112] and inflammation [113], they can also act as an H_2O_2 sink [67,114–116]. Thus, their net effect on cytoplasmic H_2O_2 levels depends on the balance between these two processes.

3.1.1. Mitochondrial ROS Production

The mechanisms responsible for mitochondrial ROS production have been described by numerous excellent reviews (e.g., [67,117–121]). Here, we will focus mainly on aspects relevant to the possible effects of hypoxia.

Oxidative Phosphorylation

Within mitochondria, the energy released by the metabolic oxidation of glucose, fatty acids, and amino acids is captured in the form of a highly negative redox potential by the transfer of electrons to nicotinamide adenine dinucleotide (NAD^+) and, via succinate, to flavin adenine dinucleotide (FAD), producing NADH and FADH_2 , respectively. These electrons are transferred from NADH/ FADH_2 into the ETC, which consists of four protein complexes with successively more positive redox potentials (Figure 4), and are then used to reduce O_2 to H_2O . The energy released as the electrons flow down this redox potential gradient is used by complexes I, III, and IV to pump protons outwardly across the inner mitochondrial membrane (IMM). The energy stored in the resulting transmembrane electrochemical gradient (ΔP or protonmotive force) is used by ATP synthase to phosphorylate ADP, generating ATP, which provides energy for cellular processes. ΔP has two components: $\Delta\Psi_m$, the membrane potential across the IMM, and ΔpH , the pH gradient across the IMM. It has recently been proposed that $\Delta\Psi_m$ is generated, not only by the proton gradient across the IMM, but also by a Na^+ gradient which also exists across this membrane due to the operation of Na^+/H^+ exchange, which is mediated by complex I [122]. The process by which the energy lost by electrons as they are transferred from metabolic substrates to O_2 through the ETC is coupled via ΔP to the production of ATP is termed oxidative phosphorylation.




Figure 4. Oxidative phosphorylation. The electron transport chain consists of four protein complexes (I–V). Complex V, the ATP synthase, is not part of the ETC per se. Complexes I and II accept electrons (e⁻) from NADH and FADH₂, respectively, which have been reduced by reactions occurring during glycolysis, fatty acid metabolism, and the Krebs cycle. NADH can also enter the mitochondrial matrix from the cytoplasm via the malate-aspartate shuttle (MAS) and the glycerol-3-phosphate shuttle (G-3-P). The electrons from complexes I and II reduce co-enzyme Q (CoQ; ubiquinone) to ubiquinol (CoQH₂). Electrons from CoQH₂ are then passed to complex III, and then via (cyt c) to complex IV, where they combine with O₂ to form H₂O. This process is driven by the successively more positive redox potentials of the complexes, and the energy lost by the electrons as they fall down this energetic redox gradient is used by complexes 1, 3, and 4 to create an electrochemical proton gradient across the inner mitochondrial wall (ΔP). The energy stored in ΔP is used by complex V to phosphorylate ADP, thereby producing ATP, which is used by cells to fuel energy-requiring reactions.

Electron Flow Through the ETC

Electrons from NADH and FADH₂ enter the ETC at complexes I and II, respectively, which use them to reduce coenzyme Q (CoQ; ubiquinone), a small-molecule constituent of the IMM, yielding CoQH₂ (ubiquinol). CoQ can also be reduced to CoQH₂ by electrons transferred during the oxidation of a number of substrates, including dihydroorotate, proline, fatty acids, glycerol 3-phosphate, and H₂S. Each substrate is oxidized by a specific dehydrogenase (e.g., proline dehydrogenase reduces CoQ by oxidizing proline to pyrroline-5-carboxylate).

CoQH₂ binds to complex III (aka Q-cytochrome *c* oxidoreductase or cytochrome *bc*₁ complex) at the Q_o site, which is close to the outer side of the IMM. One electron from CoQH₂ bound at this site is transferred to the nearby Rieske iron–sulfur cluster (and thence, in succession, to cytochrome *c*₁, cytochrome *c*, and complex IV, which utilizes pairs of electrons to reduce O₂ to water). This releases two protons to the intermembrane space (IMS). The other electron is transferred to the b_L (b₅₆₆) heme group, which is also on the outer side of the IMM. From there, it moves to the b_H (b₅₆₂) heme group, which is closer

to the matrix side of the IMM. This movement is driven by the more positive redox potential of b_H compared to b_L , but is opposed by $\Delta\Psi_m$; thus, a high $\Delta\Psi_m$ will retard this electron transfer. A second ubiquinone binding site, Q_i , is close to b_H , and the electron on b_H is transferred to UQ bound to this site, forming ubisemiquinone ($\text{CoQ}\cdot$). When the next molecule of CoQH_2 is oxidized at Q_o , the electron that it donates to b_L and b_H reduces the ubisemiquinone at Q_i to CoQH_2 , which returns to the CoQH_2 pool in the IMM. This causes the uptake of two protons from the matrix. The net result of this process, referred to as the *Q cycle*, is that for each CoQH_2 oxidized, two electrons flow via cytochrome *c* to complex IV, two protons are taken up from the matrix, and four protons enter the IMS, helping to generate ΔP [123,124].

Factors Governing Mitochondrial ROS Production

There are more than a dozen sites within the mitochondria at which electrons flowing into or through the ETC can react with O_2 to form superoxide and/or H_2O_2 (see Figure 5, and the legend to Figure for a list of the sites and a definition of their abbreviations). Many of these sites are locations at which dehydrogenases transfer electrons from metabolic substrates to NAD^+ or CoQ. Superoxide produced is rapidly reduced to H_2O_2 by MnSOD (SOD2) in the mitochondrial matrix or Cu/ZnSOD (SOD1) in the IMS. Nevertheless, superoxide evidently exists long enough to exert local effects, for example, oxidizing the Fe-S cluster of mitochondrial aconitase in the matrix, leading to its loss of catalytic function [125]. The ratio of mitochondrial superoxide + H_2O_2 production to O_2 consumption is in the range of 0.1 to 0.5% [126].

The rate of superoxide/ H_2O_2 production at each site is a function of the O_2 concentration ($[\text{O}_2]$), the concentration of the site in the reduced state ($X^{\cdot-}$), and a second-order rate constant (k):

$$(\text{dROS}/\text{dt}) = ([\text{O}_2])(X^{\cdot-})(k) \quad (1)$$

where $X^{\cdot-}$ is the product of the concentration of the site and the fraction of the site which is in the reduced state [67,118]. Overall mitochondrial ROS production is the sum of that produced at each site. Whereas some sites are incorporated into proteins as prosthetic groups (e.g., flavin mononucleotide in the I_F site on complex I), I_Q in complex I and III_Q in complex III serve as sites within the complexes at which ubisemiquinone is formed and can donate electrons to O_2 to form superoxide or H_2O_2 . The mitochondrial ROS-producing sites fall into two isopotential groups defined by the redox couple involved in ROS production. The NADH-linked sites (the α -keto ketoacid dehydrogenase complexes and Complex I_F) operate at a redox potential of ~ -280 mV, whereas the coenzyme-Q linked sites (the CoQ-linked dehydrogenases, complex I_{CoQ} , complex II_F , and complex III_{CoQ}) [127] operate at a potential of $\sim +20$ mV [121]. The relative amounts of ROS produced by each site vary between cell types and under different conditions [121,126,128]. However, complex I_{CoQ} , complex II_F , α -KGDH, PDH, OADH, and III_{CoQ} are thought to be important mitochondrial ROS sources [67,129], with III_{CoQ} having the highest capacity for ROS production, at least in skeletal muscle [121].




Figure 5. Mitochondrial ROS production. There are more than a dozen sites (e.g., I_Q, G_{CoQ}) at which electrons can ‘leak out’ of the ETC and react with O₂ to form superoxide and/or H₂O₂. In addition to the classical sites of ROS production at complex I (I_F and I_Q), complex II (II_F), and complex III (II_{CoQ}), ROS are also produced by dehydrogenase complexes, including pyruvate dehydrogenase (PDH), α-ketoglutarate dehydrogenase (α-KGDH), branched chain α-ketoacid dehydrogenase (BCKDH), and 2-oxoadipate dehydrogenase (OADH), which are associated with the inner mitochondrial membrane during the oxidation of metabolites. Several of these sites (I_F and also P_F, K_F, B_F, and O_F, which represent the ROS-producing loci of these dehydrogenase complexes) are in redox equilibrium with the mitochondrial NAD⁺:NADH couple. The other sites, which include I_Q (also termed I_{CoQ}), II_F, II_{CoQ}, G_{CoQ}, E_F, D_{CoQ}, P_{CoQ} and the ROS-producing site on sulfide quinone oxidoreductase (SQOR), are in redox equilibrium with the CoQ:CoQH₂ couple in the inner mitochondrial membrane. The G_{CoQ}, E_F, D_{CoQ}, and P_{CoQ} sites are located on the dehydrogenases *sn*-glycerol-3-phosphate (*sn*-GPD), flavoprotein electron-transferring co-enzyme Q oxidoreductase (ETFQO), dihydroorotate dehydrogenase (DHODH), and proline dehydrogenase (PRODH). The content of the figure and the names of the ROS-producing sites are based on references [67,130].

Whereas all of these sites release ROS into the mitochondrial matrix, a substantial fraction of ROS produced by Complex III_{CoQ} and G_{CoQ} is also released into the IMS [121], from which H₂O₂ is able to diffuse into the cytoplasm through voltage-dependent anion channels (VDAC) in the outer mitochondrial membrane (OMM) [131,132]. H₂O₂ released into the matrix can also access the cytoplasm, as evidenced by observations that ROS released into the matrix by complex I can be detected in the cytoplasm [133] and in the medium around isolated mitochondria [127] and intact cells [134], but is to some extent consumed by intramitochondrial antioxidant mechanisms before it can do so. The ability of Complex III_{CoQ} to release ROS into the IMS, and also its position as the ‘bottleneck’ through which all electrons from the CoQ pool pass on their way to O₂ [67], has led to suggestions that it is the main site of mitochondrial ROS production involved in regulating the redox balance of the cytoplasm [67,135].

The sites proposed to be responsible for hypoxia-induced changes in ROS production in PSMC (and CBCC, the other extensively studied cell type which responds acutely to physiological hypoxia) include I_{CoQ} [136,137]; II_F [138], and III_{CoQ} [23,139–142]. I_Q can

produce ROS during either forward electron transport (when the electron is coming from NADH via I_F) or during reverse electron transport (RET), which occurs when the Q pool is highly reduced, causing a backflux of electrons from CoQH₂ into complex I [143]. The mechanism by which the semiquinone is formed and donates an electron to O₂ at III_{CoQ} is controversial. It is thought that superoxide/H₂O₂ is produced at this site via the reduction of O₂ by ubisemiquinone, which exists transiently at III_{CoQ} after CoQH₂ has lost one electron to the Rieske complex (referred to as the ‘semi-forward reaction’), or due to the backflow of an electron from b_L (the ‘semi-reverse reaction’) [144–146]. Thus, superoxide production by III_{CoQ} is promoted by an increase in $\Delta\Psi_m$, which slows the loss of electrons from b_L [147], and by the drug antimycin, which causes an accumulation of electrons at b_L and b_H by preventing the binding of CoQ to Q_i [145]. Note that III_{CoQ} is located at the site where CoQH₂ binds to complex III, which is generally termed Q_o.

Since ROS production by I_{CoQ} and III_{CoQ} is linked to co-enzyme Q, the extent to which the CoQ pool is reduced is a key determinant of mitochondrial ROS production. Importantly, it is thought that ROS production at III_{CoQ} is maximal when the CoQ pool is partially but not fully reduced [146,148,149].

A number of compounds which affect mitochondrial ROS production by binding to specific sites within the ETC are available. Of these, rotenone, myxothiazol, antimycin, cyanide, and sodium azide have been used most widely to study the role of the mitochondria in HPV. Rotenone binds to I_o, thereby blocking the ETC at this point. Myxothiazol blocks the ETC at complex III by binding to III_{CoQ}. Antimycin binds to Q_i in complex III, stopping electron flow by interrupting the Q cycle. Cyanide and azide, which bind to the Fea3-CuB binuclear center in complex IV [150], block the ETC by preventing the reduction of O₂ to H₂O₂.

Mechanisms by Which Hypoxia May Increase Mitochondrial ROS Production

The ‘metabolic hypothesis’, initially developed to explain O₂ sensing in CBCC [151], proposes that hypoxia, by causing a perturbation of the state of the ETC, generates a signal which engages with effector pathways to initiate appropriate cellular responses. According to the mitochondrial ROS hypothesis, a variant of this proposal, one way in which hypoxia can affect the ETC is that a decreased availability of O₂ tends to slow the rate of its reaction with electrons at complex IV (aka cytochrome *aa3*, cytochrome *c* oxidase or Cox). This reaction, the rate-limiting process in mitochondrial electron transport, comprises a series of steps in which two pairs of electrons transferred from cytochrome *c* are used to reduce an O₂ molecule to two H₂O molecules. This reaction is irreversible, whereas the upstream series of redox reactions which couple the oxidation of NADH and FADH₂ to the reduction of cytochrome *c*, are at near-equilibrium with each other. As a result, a decrease in the activity of Cox will cause all parts of the ETC upstream of complex IV, and also the NAD⁺:NADH and FAD:FADH₂ redox couples, to become more reduced [152]. The consequent reduction of the CoQ pool leads to higher ROS production by Complex III_{CoQ} [144].

In accordance with this concept, levels of physiological hypoxia which initiate HPV in PSMC (and transmitter release by CBCC) cause an immediate increase in cellular NAD(P)H fluorescence which is probably due mainly to a rise in the mitochondrial NADH/NAD ratio. For example, half maximal increases in NAD(P)H fluorescence were observed at PO₂ levels of 21 and 15 Torr in PSMC and CBCC, respectively [73,153]. This implies that hypoxia is impeding the flow of electrons through the ETC (although, as described below, this might not result in a commensurate inhibition of O₂ consumption). Similarly, moderate hypoxia was shown to cause the reduction of cytochrome *c* in CBCC [154] and of CoQ and cytochromes *c* and *aa3* in PSMC [141,155]. Hypoxia also alters the

magnitude of $\Delta\Psi_m$ in both types of cells (apparently, although puzzlingly, in opposite directions [142,156,157]).

Notably, however, the reduction of O_2 at Cox has an apparent P_{50} (i.e., the PO_2 at which the activity of the complex, as reflected by cellular O_2 consumption, is half maximal) of ~0.8 Torr in PASMCM [155]. This is ~60 times lower than the PO_2 level which half-maximally stimulates HPV. This raises the question of how mild-to-moderate levels of hypoxia, which cause HPV, could perturb the ETC, thereby causing a signaling effect (e.g., reduction of the CoQ pool and increased ROS production by Complex III_{CoQ}), even though they are too high to affect its rate-limiting step.

Several explanations have been advanced to explain this apparent discrepancy [3,50,73]. One is that the basal mitochondrial PO_2 may be lower than that in the extracellular space (ECS) [158,159] and can therefore fall during moderate hypoxia to a level small enough to significantly limit the activity of Cox. However, the ECS to mitochondria PO_2 gradient in PASMCM would have to be very substantial for a fall in PO_2 associated with moderate hypoxia to exert a meaningful effect on the activity of Cox [73].

Gnaiger and colleagues proposed another explanation for this conundrum, based on their finding that the apparent P_{50} for Cox is proportional to its activity [160]. According to their model, oxidase activity is controlled by the rate at which electrons are being fed into it by cytochrome *c*. This in turn depends on the relative expression of Cox compared to that of the upstream complexes which provide cytochrome *c* with electrons; in effect, the individual cytochrome *c* enzymes compete with each other for the pool of reduced cytochrome *c*. Thus, if the expression of Cox relative to that of complexes 1–3 is low, the supply of electrons to each cytochrome *c*, and therefore its activity and the apparent P_{50} , will be higher. Thus, cells with a relatively low expression of Cox will be more sensitive to moderate hypoxia.

The P_{50} is also increased by nitric oxide (NO) which competes with O_2 for binding to Cox [161], especially under hypoxic conditions. The presence of endogenous nitric oxide synthase (NOS) was shown to be associated with increases in the reduction of cytochromes B_H , *c*1, and *aa*3 and also ROS production under hypoxic conditions in RAW246.7 cells [162]. In line with a role for NO-mediated block of Cox in O_2 sensing, the drug L-NAME (N(ω)-nitro-L-arginine methyl ester), a blocker of endothelial NOS (eNOS) blocked hypoxia-induced ROS production and O_2 sensing in endothelial cells [163–165].

Determining whether an interaction between NO and Cox in PASMCM is involved in O_2 sensing during HPV is not straightforward, since NO production by the pulmonary vascular endothelium exerts a powerful vasodilating influence which inhibits HPV by activating the guanylate cyclase/cyclic GMP/protein kinase G pathway. This has been demonstrated by studies in which knockout of eNOS and application of non-selective NOS antagonists increased the amplitude of HPV in studies carried out in perfused lung [16,166–169], and in humans in vivo [170]. The fact that HPV persisted [171] or was increased [169] when the synthesis of NO by NOS in PA was prevented seems to militate against the possibility that NO could be playing a role in O_2 sensing in HPV. Even so, there is evidence that at least some types of cells possess NOS-independent pathways for synthesizing NO, and that these are stimulated by hypoxia. For example, hypoxia was shown to induce mitochondria from rat liver to produce NO through a Cox-mediated reduction of nitrite (NO_2^-) [172]. Although the importance of this mechanism has been disputed [173], hypoxia is also thought to activate other heme-dependent pathways within cells which can also generate NO from nitrite [174]. Furthermore, it has been shown that Cox plays an important role in metabolizing NO and that this process is inhibited by hypoxia [173]. This would also help to maintain NO levels during hypoxia, in which case there might be enough present in PASMCM during hypoxia to compete with O_2 for binding

to Cox. Therefore, although there is currently no evidence that NO is involved in O₂ sensing in HPV, it remains an intriguing possibility.

The mitochondrial concentration of H₂S has also been proposed to increase during hypoxia [175], and since it also inhibits Cox, with a submicromolar K_i for the isolated enzyme [176], it might have a similar effect to that proposed for NO. However, much higher concentrations of H₂S appear to be required to block Cox in intact cells and tissues [177,178]. These concentrations are unlikely to exist under physiological conditions, although this cannot be ruled out [179,180].

Another mechanism by which moderate levels of hypoxia could affect oxidative phosphorylation is described in a study carried out in isolated mitochondria from rat liver by Wilson et al. [181], and further supported by work using mitochondria from cardiac myocytes [182]. Wilson and colleagues presented evidence that the actual P₅₀ for Cox is much higher than its apparent value, which is typically derived by plotting the rate of respiration (measured as mitochondrial O₂ consumption) vs. PO₂. According to this scheme, even relatively moderate levels of hypoxia (<40 Torr in this study) decrease the activity of Cox. This would tend to slow the flow of electrons from cytochrome *c* to Cox, causing cytochrome *c* to become more reduced. However, this reduction of cytochrome *c* increases its tendency to ‘push’ electrons onto Cox. These two tendencies, which have opposing effects on electron flux, cancel each other out. Thus, hypoxia does not alter the rate at which electrons flow into Cox, or on the rate at which it reduces O₂ to water (i.e., the rate of O₂ consumption, which is equivalent to respiration). Instead, the inhibitory effect of moderate hypoxia on Cox is reflected by an increased reduction of cytochrome *c* rather than by a decrease in respiration. Wilson and colleagues observed a reduction of cytochrome *c* which was half-maximal at a PO₂ of 12 Torr, a value which they proposed was a reflection of the actual P₅₀ for the activity of Cox. This compensatory mechanism would fail as the PO₂ falls towards zero because cytochrome *c* becomes maximally reduced. Thus, with very severe hypoxia, O₂ consumption falls steeply, and the PO₂—dependency of this decline would give rise to the very low apparent P₅₀ for the reaction between Cox and O₂ which is observed when measuring O₂ consumption as a function of PO₂. Because the reactions of oxidative phosphorylation upstream of Cox are at near-equilibrium [152,183], the increased reduction of cytochrome *c* at PO₂ levels well above the apparent P₅₀ for Cox would ‘back up’ into complexes 1–3 and the co-enzyme Q pool, causing these to become more reduced, and would also raise the NADH/NAD⁺ ratio. This would elevate the degree of reduction for both CoQ- and NADH-linked sites of ROS production, tending to increase H₂O₂ production.

This mechanism, which would be further promoted if hypoxia also decreased the cell energy state [184], is not necessarily restricted to any particular type of cell. However, it has also been suggested that Cox in PASM and CBCC is unusually sensitive to hypoxia (i.e., has a higher P₅₀) because they both express high levels of Cox4i2, an atypical subunit of Cox which has been shown to increase its sensitivity to O₂ [185] and is negligibly expressed in other types of cell [186]. If the P₅₀ for Cox was raised sufficiently, this would enable physiological levels of hypoxia to decrease the activity of COX and therefore slow the flux of electrons through the ETC and increase the degree of reduction at CoQ- and NADH-linked ROS-producing sites. Supporting this possibility, the P₅₀ for the hypoxia-induced rise in NAD(P)H in sympathetic neurons was 0.3 Torr, as compared to 15 Torr in CBCC (*n.b.* the rise in NAD(P)H reflects the inhibition of Cox, since this carries out the rate-limiting step of respiration and therefore also of the oxidation of NADH to NAD) [187].

Another factor which may cause a hypoxia-induced increase in mitochondrial ROS production, in this case independently of any effect on Cox, is a rise in the concentration of the Krebs cycle intermediate succinate. Couchani et al. [112] showed that ischemia led

to the reversal of the reaction carried out by succinate dehydrogenase (complex II), causing an accumulation of succinate in myocardial cells. Cardiac reperfusion then evoked a burst of ROS production due to the oxidation of the accumulated succinate by complex II, and this was blocked by the drug rotenone, which prevents the binding of quinones to complex I. This suggested that the ROS burst was mediated by complex I and was due to RET, which causes a high level of ROS production by complex I, probably at the 1_{CoQ} site [121,188,189].

The role of succinate and RET in O_2 sensing during HPV is currently unknown. However, based in part on the observation that the complex II blocker dimethyl malonate depressed the response of CBCC to hypoxia, López-Barneo's laboratory [190,191] initially suggested that succinate-linked RET contributes to acute O_2 -sensing in these cells by causing ROS production by complex I; a similar finding was reported by Swiderska et al. [192], who also used dimethyl malonate. Nevertheless, it has more recently been reported that O_2 sensing in CBCC was not obviously affected in mice in which RET was strongly depressed due to the introduction of a mutation into the ancillary complex I subunit ND6 [193]. The involvement of RET in O_2 sensing in CBCC can also be questioned because, whereas ROS generated at complex I due to RET are released into the mitochondrial matrix [121], the matrix redox balance in CBCC was reduced, rather than oxidized, by hypoxia [191]. In addition, the use of dimethyl malonate is problematic because it could potentially depress ROS production at complex III by decreasing succinate-mediated reduction of the CoQ pool [144].

Hernansanz-Agustin and colleagues [194,195] have proposed a mechanism by which hypoxia can increase complex III-mediated ROS production, which is based on evidence that complex I has intrinsic Na^+/H^+ antiport activity (pumping Na^+ from the IMS into the matrix in exchange for protons), which is greatly increased when it undergoes a hypoxia-induced transition from its normal active state to a 'de-active' conformation in which it loses its usual NADH-ubiquinone oxidoreductase/proton-pumping activity [196,197]. A deficit of O_2 is proposed to promote this 'A/D transition' by causing a net reduction of the CoQ pool, thus inhibiting electron flow through complex I because there is less oxidized CoQ (ubiquinone) available to accept electrons in the quinone binding site [198]. Interestingly, the active/de-active ratio varies over a broad PO_2 range, and deactivation can occur quickly, rendering it a feasible target for moderate hypoxia [199,200].

According to this scheme, A/D transition induced by hypoxia causes matrix acidification by promoting H^+ influx via the complex I Na^+/H^+ antiport and by inhibiting the normal Complex I-mediated pumping of H^+ from matrix to IMS [194]. This acidification causes the partial solubilization of the calcium phosphate complex which is present at a very high concentration in the matrix [201]. The resulting increase in free $[Ca^{2+}]_{mito}$ promotes Na^+ entry into the matrix via the mitochondrial $Na^+/Ca^{2+}/Li^+$ antiporter (NCLX). This causes a rise in matrix $[Na^+]$ which decreases the fluidity of the IMM, especially in the leaflet of the membrane adjacent to the matrix. The authors presented evidence that, whereas an increase in $[Na^+]$ did not affect the function of the individual respiratory complexes when they were studied in isolation (i.e., in a mitochondrial membrane preparation), it did decrease the flow of electrons from complex II and G3PD to cytochrome in intact mitochondria. They proposed that the decreased fluidity of the IMM due to the rise in matrix Na^+ induced by hypoxia would retard the mobility of $CoQH_2$ in this membrane, thereby slowing electron flow from complex II and G3PD to complex III. This would slow the Q cycle, thereby increasing ROS production by prolonging the lifetime of ubisemiquinone at the Q_o site. This mechanism is suggested to contribute to O_2 sensing in HPV [194] (see Section 5.2.2).

There is also evidence that an increased Ca^{2+} concentration in the mitochondrial matrix, which has been shown to occur in PSMC during hypoxia (see Section 5.2.1),

stimulates pyruvate dehydrogenase, α -ketoglutarate dehydrogenase, and complex II [202]. This increases the flux of electrons into the CoQ pool, thereby stimulating oxidative phosphorylation and increasing ROS production [203,204]. It has alternatively been proposed that an increased $[\text{Ca}^{2+}]_{\text{mito}}$ elevates $\Delta\Psi_m$ by reversing a brake on Cox exerted by ongoing ATP-dependent phosphorylation of this complex [205]. Regardless of the mechanism by which $[\text{Ca}^{2+}]_{\text{mito}}$ or other factors elevate $\Delta\Psi_m$, there is abundant evidence that an increased (i.e., hyperpolarized) mitochondrial membrane potential powerfully stimulates ROS production at complexes 1 and 3 [144,147,206]. Ramzan et al. [207] suggested that an increase in $[\text{Ca}^{2+}]_{\text{cyto}}$, which raises $[\text{Ca}^{2+}]_{\text{mito}}$ via influx through the mitochondrial Ca^{2+} uniporter [203,208], accounts for the rise in mitochondrial ROS production evoked by a range of stressors, including hypoxia, in a variety of cell types. However, in this case, increased mitochondrial ROS production could not be the initial O_2 -sensing signal (which would be needed to generate the rise in $[\text{Ca}^{2+}]_{\text{cyto}}$), although it could act to amplify this signal.

Hajnóczky and colleagues have presented an intriguing proposal that an increase in the Ca^{2+} concentration in the mitochondrial matrix may cause it to swell due to the increased influx of K^+ (and therefore water) into the matrix via mitoK_{Ca} channels in the IMM [103]. This then causes a compression of the mitochondrial cristae, which are continuous with the IMS. This would decrease the volume of the cristae/IMS compartment, increasing its $[\text{H}_2\text{O}_2]$ and thereby promoting the leakage of H_2O_2 into the cytoplasm.

Notably, the concept that an increase in $[\text{Ca}^{2+}]_{\text{mito}}$ raises mitochondrial ROS production has been vigorously contested [209].

Possible Mechanisms by Which Hypoxia Could Decrease Mitochondrial ROS Production

According to Eqn 1, a fall in PO_2 would tend to decrease mitochondrial ROS production by the law of mass action, and most papers which have described hypoxia-induced falls in mitochondrial ROS production in cells or isolated mitochondria seem to have implicitly assumed that they occurred for this reason. This idea is consistent with evidence that ROS production by isolated mitochondria is linearly related to the O_2 concentration over a wide PO_2 range [210,211]. Although others have reported a hyperbolic relationship between the PO_2 and ROS production [127,212], this does not necessarily conflict with the idea that the reactions between O_2 and electrons leaving ROS-producing sites in the ETC are linearly dependent on the $[\text{O}_2]$ [149].

Conversely, Archer and colleagues have suggested that a fall in PO_2 also diminishes ROS production by decreasing the rate of electron flow through the ETC, thereby limiting the availability of electrons at the sites in the proximal ETC at which they can react with O_2 [21,52], see also [213]. This would appear to imply that the location at which hypoxia is slowing electron flow must be either at or upstream of the site where the ROS which regulate HPV are being produced, since a block of the ETC downstream of this site would be predicted to increase its electron occupancy and therefore promote its ROS production. This is consistent with evidence from this laboratory (but not others, see [3]) that the effects of hypoxia on ROS production are mimicked by the I_Q-blocker rotenone [214]. Similarly, these effects have been reported to require the complex I subunit Ndufs2 [137], which forms part of I_{CoQ} and interacts with the N2 Fe-S cluster that is believed to mediate the formation of semiquinone at this site [215–217] (see Section 5.1.1). Nevertheless, the nature of the interactions between hypoxia, Ndufs2, and a slowing of electron flow, which could result in decreased ROS production by complex I remain to be defined.

3.1.2. Mitochondrial ROS Consumption

Mitochondrial matrix consumption of H_2O_2 depends on the GSH and Trx2 redoxin systems (see Section 2.2), and on Prxs and catalase [67,116,218]. ROS released into the IMS are subject to degradation by SOD1, GSH, and cytochrome *c* [219]. The antioxidant

activities of GSH, Trx2, and peroxiredoxins require that they be returned to the reduced state by NADPH after having been oxidized due to the scavenging of H_2O_2 . The maintenance of sufficient matrix NADPH to perform this function depends on the activity of the mitochondrial enzymes malic enzyme (ME3) and isocitrate dehydrogenase (IDH2), which reduce NADP^+ , and also on nicotinamide nucleotide transhydrogenase (NNT), an enzyme in the IMM which couples the oxidation of NADH to the reduction of NADP^+ within the matrix.

These antioxidant intra-mitochondrial mechanisms can affect the redox balance of the cytoplasm in two ways. Firstly, they limit the extent to which H_2O_2 generated within the matrix reaches the cytoplasm [115,218]. Secondly, the mitochondria can act as a sink for cytoplasmic H_2O_2 . A study [116] which examined the effect of knocking out and over-expressing mitochondrial antioxidant enzymes on the redox state of the cytoplasm in H9c2 cells exposed to the organic peroxide tert-butyl hydroperoxide (t-BOOH) (1 mM) reported that knockout of NNT, ME3, or IDH2, and also the mitochondrial antioxidant enzyme thioredoxin2, increased the extent to which t-BOOH oxidized the cytoplasm. In contrast, knockout of cytoplasmic antioxidant enzymes did not ameliorate the effects of t-BOOH on the mitochondria. The authors suggested that the mitochondria may function to degrade H_2O_2 which leaks into cells from the ECS. Interestingly, another paper reported that application of 25 and 500 μM extracellular H_2O_2 to EA.hy926 cells caused a rise in cellular H_2O_2 which was larger and occurred more rapidly in the mitochondria compared to the cytoplasm and nucleus [220]. These reports suggest that the mitochondria play a crucial role in the ability of cells to scavenge H_2O_2 leaking in from the extracellular space, although whether this occurs at the relatively low extracellular H_2O_2 concentrations thought to exist under physiological conditions can be questioned, especially given the antioxidant prowess and high concentrations of cytoplasmic Prxs [221] (see Section 2.2).

Oxidation of the matrix leads to S-glutathionylation of complex I and other proteins associated with ETC ROS production, including pyruvate dehydrogenase and α -ketoglutarate dehydrogenase [89]. This depresses oxidative phosphorylation and ROS production, thus stabilizing the redox state of the matrix. Another factor promoting mitochondrial redox homeostasis is that the activity of NNT is crucially dependent on ΔP , meaning that both the production and NADPH-dependent scavenging of ROS within the matrix are enhanced by increases in [NADH] and ΔP . Nevertheless, Treberg et al. [218] demonstrated that the effect of $\Delta\Psi_{\text{m}}$ on mitochondrial ROS production was greater than that on peroxiredoxin-mediated ROS scavenging, suggesting that increasing $\Delta\Psi_{\text{m}}$ would produce a net rise in matrix [H_2O_2].

3.2. ROS Production by NADPH Oxidases

The seven members of the Nox family of oxidoreductases (Noxs 1–5 and Duox 1 and 2) function as adjustable sources of superoxide and/or H_2O_2 which regulate the activities of nearby target proteins. Four Nox isoforms (Nox 1, 2, 4, and 5) are expressed in vascular smooth muscle cells [69]. Nox2 has been shown not to be required for HPV, at least in mice [222], and the role in HPV of Nox 5 in HPV, which is expressed in primates but not rodents, is unknown. However, Noxs 1 and 4 have been proposed to be involved in HPV (see Sections 5.2.5 and 8). Both of these Nox isoforms are heterodimers comprising a membrane-spanning oxidoreductase subunit, which transfers electrons from NADPH to O_2 via FAD and two heme groups, and p22^{phox}, a smaller transmembrane scaffolding protein. Nevertheless, they differ in important ways.

3.2.1. Nox1

Nox1 is present in the plasmalemma, localized to the caveoli in vascular smooth muscle cells [223]. It releases superoxide into the ECS, from where it can enter cells either via

plasmalemmal anion channels or, after being converted to H_2O_2 by EC-SOD, through aquaporins [224]. Nox1 can also be endocytosed upon its activation by certain stimuli, potentially leading to redox signaling by superoxide and/or H_2O_2 both within the endosome and the cytoplasm [225]. Activation of Nox1 is triggered by the binding of the cytoplasmic proteins p47^{phox}, p67^{phox}, and p40^{phox}. Its recruitment of these proteins is triggered by the phosphorylation of p47^{phox} by protein kinase C (PKC) and/or a Src family tyrosine kinase [69,226]. Alternatively, p47^{phox} and p67^{phox} may be replaced by their homologues NoxO1 and NoxA1, respectively. In this case, NoxO1 is constitutively tethered to p22^{phox}, and superoxide production is stimulated by the binding of NoxA1 to NoxO1 [227], which is promoted by PKC [228], and also requires the phosphorylation of threonine 429 on Nox1 (e.g., by PKC β 1; [229]. The combination of p47^{phox} and NoxA1 has also been shown to stimulate Nox in vascular smooth muscle cells [230].

Activation of PKC and Src kinases can be initiated by diverse stimuli, including lipid mediators produced by phospholipases A₂, C, and D following their activation by the binding of agonists to multiple types of G-protein coupled receptors. PKC and Src family kinases can also be activated by each other, and by ROS [226], thereby enabling positive feedback. Full activation of Nox1 occurs with the subsequent binding of activated monomeric G protein Rac1. This in turn necessitates the stimulation of Rac guanine exchange factors following their tyrosine phosphorylation and binding to phosphatidyl inositol triphosphate, which is generated by transactivation of the epidermal growth factor receptor and the stimulation of phosphatidyl inositol-3 kinase [69].

3.2.2. Nox4

Nox4 is thought to be the most highly expressed isoform in vascular smooth muscle cells, in which it is localized to the plasma membrane in association with focal adhesions [223]. It is also expressed in the sarcoplasmic reticulum [231], mitochondria [232], and nucleus [233]. Nox4 primarily produces H_2O_2 , which presumably gains access to the cytoplasm via aquaporins in the plasmalemma and intracellular membranes. The Nox4/p22^{phox} dimer is constitutively active, suggesting that it can produce appreciable levels of H_2O_2 under basal conditions, and that its ability to do so may be regulated by its level of expression [234], and, acutely, by the availability of NADPH and also O_2 [68]. H_2O_2 production by plasmalemmal Nox4 is increased by its association with polymerase delta interacting protein 2 (Poldip2) [235] and, in vascular smooth muscle cells, can be enhanced acutely by G protein-coupled vasoconstrictors through mechanisms that are not well understood. Little is known about the regulation of intracellular Nox4 [69].

One important property of Nox4 which makes it well suited to act as a sensor for acute changes in the PO_2 is that its $K_m(\text{O}_2)$ (the PO_2 at which its production of H_2O_2 is half maximal) has been reported to be ~136 Torr, such that its activity would be decreased by even mild hypoxia. The $K_m(\text{O}_2)$ for Nox1 seems not to have been measured, although that for Nox2 is ~25 Torr [68].

4. Intra- and Extracellular H_2O_2 Concentrations

Recent reviews state that the basal intracellular $[\text{H}_2\text{O}_2]$ is in the range of 1–5 nM [48,111,236], citing as evidence an elegant study by Oshino et al. [226] which took advantage of the fact that the enzyme catalase reacts with H_2O_2 to form the intermediate Compound 1 (which can be monitored spectrophotometrically) to calculate that the intracellular concentration of H_2O_2 in rat liver under resting conditions was on the order of 10^{-9} M [237,238], see also [239].

A similar estimate was reported by Lyublinskaya & Antunes [240], who, using an approach based on measuring the kinetics of oxidation of the H_2O_2 sensor HyPer, determined that the basal cytoplasmic $[\text{H}_2\text{O}_2]$ in K562 cells is in the low nanomolar range [240].

Their study was designed to determine the involvement of extracellular H_2O_2 in setting the basal cytoplasmic $[\text{H}_2\text{O}_2]$. The concept that extracellular $[\text{H}_2\text{O}_2]$ is an important determinant of the cytoplasmic $[\text{H}_2\text{O}_2]$ had emerged from studies showing that a bolus of H_2O_2 added to the solution bathing cells in culture is taken up (via aquaporins [82,241]) and then rapidly scavenged by intracellular antioxidant systems [242–244]. Because the removal of H_2O_2 by cellular antioxidants is much faster than its uptake, a large H_2O_2 concentration gradient across the cell membrane develops immediately upon its extracellular application [245]. The magnitude of this gradient is maintained at a constant level as the H_2O_2 in the bath is taken up by the cells, such that intra- and extracellular $[\text{H}_2\text{O}_2]$ undergo a parallel exponential decline [246].

The steady-state transmembrane gradient can be assessed by keeping the cell number low enough to ensure that the uptake of H_2O_2 has a negligible effect on its concentration in the bath, or by using an H_2O_2 —regulating system such as glucose/glucose oxidase/catalase to stabilize the $[\text{H}_2\text{O}_2]$ in the bathing solution [247,248]. Using the former approach, Lyublinskaya & Antunes reported that the cytoplasmic $[\text{H}_2\text{O}_2]$, which was 2.2 nM when no H_2O_2 was added to the bath, rose to 3.5 and 8.9 nM, respectively, when 1 and 2.5 μM H_2O_2 were applied to the solution. The authors calculated that an average trans-plasma-lemmal H_2O_2 concentration gradient of ~390-fold was present over this range of extracellular $[\text{H}_2\text{O}_2]$.

The existence of even larger (~1000-fold) $[\text{H}_2\text{O}_2]$ gradients between extracellular and cytoplasmic $[\text{H}_2\text{O}_2]$ had previously been predicted by mathematical models developed by Adimora et al. [246] and Lim et al. [249], which took into account information about the rates of transmembrane H_2O_2 flux, cellular production, and scavenging, e.g., [244,250]. Importantly, whereas cellular H_2O_2 was initially thought to be scavenged by catalase and glutathione peroxidase [243], yielding calculated transmembrane gradients which were small (e.g., ~7 in Jurkat T cells [242]), these models took into account more recent evidence that Prxs, which are present in micromolar concentrations in cells and can have a very high affinity for H_2O_2 (e.g., Prx2), are predominantly responsible for removing it from the cytoplasm [99]. Importantly, Lim et al. [249] went on to show that the scavenging of H_2O_2 following its entry into the cell would also set up a significant peroxide concentration gradient *within* the cytoplasm which would develop within milliseconds and remain stable if the extracellular $[\text{H}_2\text{O}_2]$ was constant. Thus, they calculated that $[\text{H}_2\text{O}_2]$ at the inner face of the plasmalemma in a spherical cell with a radius of 10 microns would be ~0.3% of that in the extracellular medium, falling to ~0.08% and 0.02% at 2 and 4 microns into the cell, respectively. These studies imply that the basal intracellular $[\text{H}_2\text{O}_2]$ in the subplasmalemmal region is strongly influenced by the extracellular $[\text{H}_2\text{O}_2]$.

The physiological implications of these findings are difficult to parse, firstly because the $[\text{H}_2\text{O}_2]$ in solutions bathing biological preparations in vitro is typically not controlled or measured during investigations of its intracellular dynamics or effects, and secondly because little is known about the extracellular $[\text{H}_2\text{O}_2]$ in vivo. Although the basal extracellular $[\text{H}_2\text{O}_2]$ has been suggested to be 1–5 μM [48,111,236], this seems to have been measured only by one laboratory, which reported mean in vivo values of 3 and 5 μM for extracellular $[\text{H}_2\text{O}_2]$ in rat ventrolateral medulla and hippocampus, respectively [251–253]. Additional indirect evidence supporting the possibility that cells experience extracellular H_2O_2 concentrations in the low nM range came from an investigation in which the extracellular $[\text{H}_2\text{O}_2]$ immediately adjacent to the extracellular surface of the plasmalemma in A549 (tumor-derived) cells in culture was measured using a membrane-tethered sensor [254]. This showed that average extracellular $[\text{H}_2\text{O}_2]$ close to the membrane was ~2 μM , and also revealed the presence of discrete H_2O_2 hotspots, presumably associated with the presence of a higher local expression of Nox in that area of the membrane, with concentrations of up to ~12 μM . [255]. The authors suggested that these areas were likely to give

rise to corresponding H_2O_2 hotspots in the subplasmalemmal cytoplasm, although they did not confirm this. The results of the study also imply that the basal extracellular $[\text{H}_2\text{O}_2]$ is set locally, implying that it may vary between different tissues and conditions.

Parenthetically, basal plasma $[\text{H}_2\text{O}_2]$ is also generally thought to be 1–5 μM [256]. However, Sousa et al. [257] contend that the basal plasma $[\text{H}_2\text{O}_2]$ has been overestimated, and that due to its uptake by erythrocytes, it is in the range of 10 nM. They note, for example, that taking into account measurements of the permeability of erythrocytes to H_2O_2 , Prx2, which should be heavily oxidized when the plasma $[\text{H}_2\text{O}_2]$ is ≥ 80 nM, is instead found to be almost completely reduced under basal conditions [258].

Although the evidence is very limited, the agreement between the measured values of intra- and extracellular $[\text{H}_2\text{O}_2]$ and the ~ 1000 -fold $[\text{H}_2\text{O}_2]$ gradient predicted by mathematical models makes a neat story. Nevertheless, it has also been suggested that the basal cytoplasmic $[\text{H}_2\text{O}_2]$ may be in the sub-nanomolar range [259]. Lim et al. [221] concluded on theoretical grounds that, absent H_2O_2 influx from the extracellular space, cellular H_2O_2 production and scavenging should set the basal cytoplasmic $[\text{H}_2\text{O}_2]$ at ~ 80 pM. Since, according to their later model [249], the influence of the extracellular $[\text{H}_2\text{O}_2]$ decreases steeply moving away from the cell membrane, $[\text{H}_2\text{O}_2]$ would be predicted to approach this level in the core of large cells. Moreover, using HEK293-MSR cells expressing the novel high-sensitivity H_2O_2 indicator Apex2 [260], Eid et al. [261] detected a sustained increase in intracellular $[\text{H}_2\text{O}_2]$ upon the application to the bathing solution of H_2O_2 concentrations down to 25 nM in HEK293-MSR cells. Sommer et al. [142] found that applying 124 nM H_2O_2 caused membrane depolarization and inhibited the K_v current in PASMNC, and this laboratory has recently reported that extracellular concentrations of H_2O_2 of as little as 10 nM inhibited the current through $\text{K}_v1.5$ channels co-expressed with $\text{K}_v\beta1.4$ ancillary subunits in oocytes [262]. This implies that the basal $[\text{H}_2\text{O}_2]$ even adjacent to the cell membrane could have been in the subnanomolar range, although whether this applies to cells in vivo would presumably depend on the balance between the rates of H_2O_2 influx from the ECS and its intracellular scavenging.

Another implication of the presence of powerful antioxidant mechanisms present in the cytoplasm is that increases in $[\text{H}_2\text{O}_2]$ emanating from intracellular sites are highly localized. This was demonstrated by Mishina et al. [263], using cells expressing HyPer fused to growth factor receptors in the plasmalemma and protein tyrosine phosphatase 1B (PTP1B) in the endoplasmic reticulum (ER) membrane. They found, for example, that applying epidermal growth factor (EGF) to HeLa cells expressing an EGF receptor—HyPer fusion construct led to the internalization of a fraction of the EGF receptor into endosomes, while the rest remained in the plasmalemma. The endosomal EGF receptors reported an increase in $[\text{H}_2\text{O}_2]$ due to Nox activity associated with receptor activation, while those in the plasmalemma did not. This demonstrated that only the internalized EGF receptor was activated, and also that H_2O_2 was not able to diffuse from far enough from the endosomes to reach the plasmalemma. Likewise, H_2O_2 generated by Nox in response to platelet-derived growth factor (PDGF) was sensed by PTP1B-HyPer in the ER membrane but not by PDGF receptor-HyPer in the endosomal membrane. In a subsequent study in HeLa cells, this group demonstrated the presence of a steep cytoplasmic H_2O_2 concentration gradient when a H_2O_2 -generating system (D-amino acid oxidase/alanine) was localized to the nucleus [264].

However, an investigation utilizing the highly sensitive indicator HyPer7 failed to detect cytoplasmic H_2O_2 gradients, or indeed any rise in the global cytoplasmic $[\text{H}_2\text{O}_2]$, associated with an increase in mitochondrial ROS production induced by D-amino acid oxidase/alanine in K562 cells [265]. On the other hand, by applying antimycin to cells fed with galactose, which enhances mitochondrial ROS production, Hoehne et al. [133] were able to demonstrate using HyPer7 that an increase in mitochondrial ROS production at

complex III did cause a small rise in the cytoplasmic $[H_2O_2]$ in HEK293 cells, although this was not observed in cells fed with glucose. Both papers concluded that H_2O_2 was being released by the mitochondria, but that it (and any HyPer7 molecules it may have oxidized [266]) were being reduced before they could diffuse far enough to generate a measurable signal in the overall cytoplasm. Notably, recent experiments using HyPer7 also detected an increase in cytoplasmic $[H_2O_2]$ elicited by mitochondrial D-amino acid oxidase/alanine in HeLa cells, leading the authors to suggest that these cells may have less effective anti-oxidant mechanisms compared to K562 cells. Interestingly, using the novel H_2O_2 indicator HyPerFLEX, which is more sensitive than HyPer7, they also observed an accompanying increase in $[H_2O_2]$ in the nucleus and endoplasmic reticulum [267].

Notably, studies by Hajnoczky and colleagues have shown that the close apposition of mitochondria to the endo/sarcoplasmic reticulum means that mitochondrial ROS release into the nanodomain between these organelles can oxidize a group of cysteines in the cytoplasmic-facing suppressor domain of the IP₃R, inducing Ca^{2+} release [103,268,269]. Using a compartmentally targeted redox sensor (Grx1-Ro-GFP2), they demonstrated in HEPG2 cells [268] that the oxidative signal due to mitochondrial ROS release was restricted to the mitochondria/ER nanodomain, and did not result in a general cytoplasmic oxidation. Intriguingly, they also presented evidence that the ER-mitochondrial interface nanodomain was more oxidized than the bulk cytoplasm even under basal conditions, suggesting the existence of a standing intracellular redox gradient which may function to enable ER Ca^{2+} release.

5. Models of O₂ Sensing in HPV

Although they have opposed views on the effects of hypoxia on cytoplasmic oxidant signaling in PASM, the Redox and Mitochondrial ROS theories both propose that the mitochondria are the primary O₂ sensors responsible for causing HPV. This concept is supported by evidence that 1. blockers of the proximal ETC abolish HPV (see Section 7.1), 2. cultured PASM depleted of their mitochondria by ethidium bromide treatment [270] do not demonstrate hypoxia-induced rises in $[Ca^{2+}]_{cyt}$ [23], and 3. mitochondria in PASM are distributed more peripherally compared to those in mesenteric artery SMC (MASMC), so are better placed to influence effectors located in the plasmalemma [271].

We will first describe the experiments leading to the development of each theory separately in Sections 5.1 and 5.2. Because these proposals share a focus on mitochondrial ROS production, the studies carried out by their proponents have often used similar approaches (e.g., assessing the effects of ETC blockers or pro- and antioxidants, measurement of ROS), and have therefore yielded information relevant to both. We will therefore consider this evidence as a whole and discuss its implications in Section 7.

5.1. The Redox Theory

The Redox Theory of hypoxia of HPV, proposed by Kenneth Weir and Stephen Archer in the mid-1990s [21,272], is based on the concept that HPV is due to the removal by hypoxia of a tonic PASM vasorelaxation [273], which is maintained under normoxic conditions by a level of cytoplasmic ROS that is higher than that present in systemic artery vascular smooth muscle cells. Recent work by Archer's laboratory ascribes this difference to the higher expression of the complex I subunit Ndufs2 in PASM [137].

According to this hypothesis, which is summarized in Figure 6, the relatively high ongoing PASM production of H_2O_2 by mitochondria under normoxic conditions leads to a basal activation of voltage-gated K⁺ (K_v) channels which maintains the cells in a relaxed state by ensuring that the membrane potential (E_m) is negative enough to suppress the activity of L-type voltage-gated Ca^{2+} channels, thereby minimizing Ca^{2+} influx. By decreasing the flux of electrons through the ETC, hypoxia diminishes the production of

mitochondrial superoxide/H₂O₂ [49]. The resulting fall in the cytoplasmic [H₂O₂] depresses K_v channel activity, causing membrane depolarization and an increased Ca²⁺ influx which evokes HPV. The mechanism by which ROS activate the K_v channels is unknown but could involve oxidation of K_v channel protein thiol-containing residues or an action on K_vβ-subunits. ROS are seen as exerting a similar effect on K_v channels in systemic artery vascular smooth muscle cells, although this influence is smaller under normoxic conditions because mitochondrial ROS production is lower. In contrast to PA, systemic arteries dilate to hypoxia, and this is proposed to be due, at least in part, to increased mitochondrial ROS production and the activation of K_v channels [156]. Thus, the opposite responses of pulmonary and systemic arteries to hypoxia are viewed as being due to its contrary effect on mitochondrial ROS production, rather than to fundamental differences between the hypoxia-sensitive effector pathways controlling vascular smooth muscle cell force development in these arteries and/or their responsiveness to ROS [49].

Although the proponents of the Redox Theory have generally stressed the effect of hypoxia on ROS, they also view hypoxia-induced reduction of cellular redox couples (e.g., NAD⁺/NADH, NADP⁺/NADPH, GSSG/GSH) as potentially contributing to HPV [21,137]. This could occur, for example, through an action of NAD(P)H on K_v channel β-subunits, which can modulate channel gating by virtue of their activity as aldehyde dehydrogenases [274]. However, taking into account that the cytoplasmic redox potential of the NADP⁺/NADPH couple in the cytoplasm is approximately −390 mV [275], this couple is already almost entirely reduced under normoxic conditions. This suggests that a fall in [ROS] in and of itself is unlikely to increase the free [NADPH], making it a poor candidate for mediating hypoxia-induced changes in K_v channel activity associated with decreased ROS production. On the other hand, the NAD⁺/NADH couple, which has a cytoplasmic redox potential of ~−240, is highly oxidized [275]. This results in a low free cytoplasmic [NADH] [275,276] which is in the appropriate range for regulating K_v channels [277]. Given its involvement in controlling the oxidation of cellular thiol switches, the GSSG/GSH redox couple is of particular interest with regard to signaling. Importantly, the GSSG/GSH ratio has been shown to be altered by the application of low concentrations of extracellular [H₂O₂], and also by hypoxia [278,279] (see Section 6.4 for a further discussion of the role of these redox couples in HPV).

Similarly, although they have emphasized the role of K_v channels as the crucial ROS-sensitive effectors, the involvement of other redox-sensitive downstream pathways which could also respond to a fall in ROS by promoting contraction (or inhibiting relaxation) are not ruled out [280]. There is, for example, evidence from Michael Wolin's laboratory that a decrease in H₂O₂ production contributes to HPV by suppressing basal vasodilation due to activity of soluble guanylate cyclase (sGC) and protein kinase G1α (see Section 8.2).




Figure 6. The Redox Theory of HPV. This is based on the concept that under normoxic conditions, mitochondria in PASMCs generate a higher level of ROS than do systemic vascular smooth muscle cells. The resulting higher level of H₂O₂ and/or oxidation of cytoplasmic redox couples in PASMC activates K_v channels, causing a relatively hyperpolarized E_m which limits Ca²⁺ influx through voltage-gated Ca²⁺ channels, thereby creating an ongoing vasodilating influence under normoxic conditions. Other as-yet-unidentified redox-regulated vasodilating pathways might also be similarly recruited by cytoplasmic oxidation. By diminishing mitochondrial ROS production, hypoxia decreases this vasodilating influence, thereby increasing vascular tone. See the text for further details.

5.1.1. Evidence for a Fall in Mitochondrial ROS Production and PASMC Reduction as the O₂ Sensor in HPV

A series of studies in the 1980s, carried out mainly by Weir, Archer, and colleagues, investigated the effects of oxidizing and reducing agents on HPV. Weir et al. [281] examined the effect of IV injection of the oxidizing agent diamide on HPV recorded in anesthetized dogs when the inspired O₂ level was reduced to 12%, and on the pressor response to PGF_{2α}. Diamide markedly diminished the increases in PAP and pulmonary vascular resistance elicited by both hypoxia and PGF_{2α}. Burghuber et al. [282] examined the effect of H₂O₂, generated by adding glucose/glucose oxidase to the perfusate, on HPV in isolated perfused rat lung, finding that it did not alter PAP under normoxic conditions, but abolished HPV and halved the amplitude of the pressor response to angiotensin 2. These responses were prevented if catalase was added to the solution and were mimicked by

perfusing the lungs with physiological saline solution (PSS) containing $\sim 10^{-6}$ M H_2O_2 . Together, these studies showed that oxidants tended to suppress HPV.

Subsequent studies from Weir and colleagues were also performed in isolated perfused rat lungs in which HPV was evoked by reducing the PO_2 in the perfusate to ~ 40 Torr. Weir et al. [283] found that diamide applied during hypoxia reversed HPV in this preparation, and that adding xanthine/xanthine oxidase (X/XO) to the solution to generate H_2O_2 had a similar effect, which was prevented if the X/XO blocker allopurinol was present. They suggested that X/XO might be acting by oxidizing cellular sulfhydryl groups, and soon afterwards proposed that hypoxia was inducing PA constriction by causing a similar effect [284]. Archer et al. [285] confirmed this effect of X/XO on HPV and found that it was attenuated if SOD and catalase were added to the perfusate. Application of SOD and catalase which had been incorporated into liposomes to facilitate their entry into PA cells similarly suppressed the effect of X/XO. These results supported the concept that reduction of redox couples in PASMC tended to promote PA constriction, both under normoxic and hypoxic conditions. Both diamide and X/XO suppressed HPV, although it is difficult to interpret this effect because the response to angiotensin 2, which was used as a pre-tone agent to amplify HPV [286] was similarly inhibited by X/XO, and the effect of diamide on the angiotensin 2 response was not described.

Archer et al. [287] performed simultaneous measurements of PAP and pulmonary ROS production in isolated and ventilated rat lungs while changing the PO_2 in the inspired gas. Luminol or lucigenin were included in the perfusion solution as an ROS indicator, and the PO_2 was varied from the normoxic baseline (20%) to 0, 1, 2.5, 10, or 95%. The key result of this paper was that lowering the PO_2 to $\leq 2.5\%$ evoked a fall in luminescence with both indicators that paralleled the decrease in PO_2 and preceded the onset of HPV. Including catalase in the perfusate did not affect lucigenin luminescence under normoxic conditions, whereas this was strongly diminished by SOD. SOD similarly depressed luminol luminescence, which was, however, somewhat increased by catalase. SOD slightly raised basal PAP and strongly enhanced HPV and the pressor response to angiotensin 2, whereas catalase had no effect on PAP under any condition. Based on their observations, the authors suggested that hypoxia was decreasing pulmonary superoxide levels. The ability of SOD to increase basal and stimulated vascular tone echoed earlier findings and was seen to support the idea that the oxidation of cell redox couples exerts a tonic vasodilating influence on PA. However, from a contemporary perspective, which views H_2O_2 rather than superoxide as mediating most ROS-dependent intracellular signaling [48], the observations that SOD caused contraction and enhanced HPV appear to be inconsistent with the Redox Theory.

A crucial step in the development of the Redox Theory came with the observation by Post et al. that hypoxia inhibited the K^+ current in pulmonary (but not renal) artery smooth muscle cells [288]. The authors then investigated whether hypoxia was inhibiting K^+ channels via the decrease in ROS which they had observed in the study described above. Based on the observation that blockers of the ETC cause PA contraction [289] and would be predicted to mimic hypoxia by suppressing mitochondrial ROS production and reducing cell redox couples such as NADH/NAD^+ and GSH/GSSG , Archer et al. [272] investigated the effects of several ETC blockers on PAP and ROS production in their rat lung model, and on the K^+ current in freshly isolated rat PASMC. Rotenone, antimycin, and cyanide, which act at complexes 1, 3, and 4, respectively, were applied in the perfusate inflow, and all produced an increase in PAP similar to that evoked by hypoxia (ventilation with 2.5% O_2). Pulmonary ROS production, assessed using luminol and lucigenin, was decreased by both rotenone and antimycin. Both blockers also abolished HPV. The fall in lung ROS production caused by hypoxia was strongly attenuated by pre-treatment with antimycin, but was unaffected by rotenone (n.b.: this appears to be inconsistent with later evidence from this

laboratory [137] that HPV is due to decreased ROS production by complex I). Cyanide caused a dose-dependent increase in PAP comparable to that caused by rotenone and antimycin, but unlike these blockers had no effect on HPV and caused a transient increase in lucigenin luminescence. Both of these responses were depressed by SOD, suggesting that they were superoxide-dependent. The K^+ current they recorded in PASMCM was strongly blocked by both TEA and 4-AP and was attenuated by hypoxia. The current was also reversibly suppressed by rotenone and antimycin but was not significantly affected by this concentration of cyanide.

In discussing their findings, the authors noted that whereas Rounds & McMurtry [289] had previously highlighted the resemblance between the effects of hypoxia and ETC blockers on pulmonary artery contraction and had speculated that hypoxia might cause a pressor response in PA by depleting ATP, this seemed unlikely in light of evidence that high-energy phosphate levels in PASMCM are little changed by hypoxia [290–292]. They proposed instead that rotenone and antimycin were causing PA contraction by suppressing mitochondrial ROS production and pointed out that the effects of these drugs on PAP, ROS production, and the PASMCM K^+ current resembled those of hypoxia. The ability of both ETC blockers to abolish HPV also suggested that they were acting via the same mechanism as hypoxia. The observation that cyanide, which did not diminish ROS, had no effect on either HPV or the K^+ current, also aligned with the idea that a fall in ROS was linked to K^+ channel inhibition and contraction. They therefore concluded that hypoxia was causing HPV by decreasing mitochondrial ROS production, which then decreased the opening of voltage-dependent K^+ channels. These papers established the basic outline of the Redox Theory, which was then set forth in detail in a seminal review by Weir and Archer in 1995 [21].

Observations that PA constrict in response to hypoxia, whereas systemic arteries dilate (e.g., [171]) could potentially be explained by differences in either the O_2 sensor or the mediators/effectors to which it is coupled. Michelakis et al. [156] therefore examined whether the hypoxia-induced fall in ROS which had been observed in PA also occurred in small renal arteries, and whether mitochondria from the smooth muscle cells of these two types of artery responded differently to hypoxia.

They observed that both hypoxia ($pO_2 \sim 40$ Torr) and rotenone inhibited the whole cell K^+ current in rat PASMCM, whereas they increased its amplitude in renal artery smooth muscle cells (RASMC). Accordingly, both stimuli increased PAP in isolated lungs but reduced the arterial pressure in perfused kidneys. ROS production, as assessed using lucigenin, Amplex Red, or 2',7'-dichlorodihydrofluorescein (DCFH) as indicators, was markedly higher in isolated pulmonary as compared to renal arteries under normoxic conditions. Both hypoxia and rotenone strongly decreased the lucigenin signal in PA but had the opposite effect in renal arteries.

In further experiments, they characterized mitochondria isolated from homogenates of whole lung (LM) and kidney (KM), using glutamate and succinate to support respiration. In line with their observations in the arteries, ROS production, assessed using lucigenin, was higher in LM vs. KM. Using Western blotting, they found higher expression of complexes I and 3 in renal compared to pulmonary arteries, and pointed out that this was consistent with their additional observations that O_2 consumption, indicative of respiration, was greater in KM compared to LM, and that the mitochondrial membrane potential, assessed using JC-1 and TMRM, was more hyperpolarized in RASMC than in PASMCM. Moreover, hypoxia increased the mitochondrial membrane potential in PASMCM but had the opposite effect in RASMC. Based on their findings that glutathione levels and Mn SOD protein expression were substantially higher in the PA compared to RA, which was consistent with an adaptation of PASMCM to the relatively O_2 -rich pulmonary milieu, the authors also proposed that PASMCM are more oxidized than RASMC, although this

conclusion is questionable because the concentrations of oxidized and reduced glutathione, which reflect the cytoplasmic redox potential, were not reported.

Wu et al. [293] used dihydroethidium (DHE) to compare the effect of hypoxia ($pO_2 \sim 30$ Torr) on ROS production by cultured human smooth muscle cells from PAs and coronary arteries (CASM). In accordance with the Redox Theory, they found that hypoxia decreased superoxide levels in PASMC. However, in contrast to the findings of Michelakis et al. [156], hypoxia also caused a similar fall in DHE oxidation in CASMC. The authors pointed out that coronary arteries dilate to hypoxia, and that the fall in superoxide in both types of cells took several minutes to occur, whereas HPV in lungs develops within seconds. This led them to speculate that ROS may not be involved in HPV, or that the opposite contractile response of pulmonary and coronary arteries to hypoxia might be due to differences in the effector mechanisms coupled to ROS. Similar observations were made by Mehta et al. [294], who found that moderate hypoxia (5% O_2) significantly decreased ROS production in both primary cultured human PASMC and CASMC. This effect was seen with three indicators (DCFH, DHE, and Amplex Red). The fall in ROS showed a (non-significant) trend to be more substantial in the PASMC with each indicator.

Notably, the physiological relevance of the effects of acute hypoxia observed in each of these studies which compared the responses of pulmonary and systemic arteries to a decrease in PO_2 are uncertain, since whereas the physoxic PO_2 for systemic arteries is much lower than it is for pulmonary arteries, there is no indication in any of these papers that different levels of PO_2 were used to simulate either normoxia or hypoxia in the two types of arteries.

In 2019, Dunham-Snary and colleagues [137] published a study of the role of ROS in HPV in rats and mice, which included a further comparison of the properties of pulmonary and renal artery mitochondria. This paper presents the most recent and comprehensive evidence currently available to support the Redox Theory, and its results relating to the acute effects of hypoxia will therefore be considered in detail.

The authors incubated isolated LM and KM at a concentration of ~ 2.5 mg protein/mL in physiological medium for 15 min under normoxic or hypoxic conditions. Using Amplex Red, they measured the H_2O_2 concentration in the conditioned media, finding that under normoxic conditions, this was much higher in medium conditioned with LM (~ 1 μM) compared to KM (< 0.2 μM). These results were consistent with their earlier findings [156].

Micropolarimetry experiments were carried out to compare the O_2 consumption rate in cultured PASMC and RASMC, finding that it was about twice as high in the former. Since mitochondria contribute the vast bulk of cellular O_2 consumption, this appears to contradict their earlier observation that the rate of respiration in isolated LM was about a third of that in KM [156]. The authors did not discuss possible reasons for these divergent results, but it is possible that they may have been due to cellular influences on mitochondrial function, which would be absent in isolated mitochondria. If so, this suggests that the effects of hypoxia on isolated mitochondria cannot simply be extrapolated to more intact preparations [62,295].

Isolated PSS-perfused rat lung was used to examine the effect of oxidants on HPV evoked using perfusate gassed with 2.5% O_2 . Injection of 0.1 mL of LM-conditioned medium into the perfusion inflow caused a rapid and marked reversal of HPV (by 56%), whereas injection of KM-conditioned medium, or medium conditioned with LM in the presence of catalase, had no effect on HPV. Perfusion with PSS containing 100 μM t-BOOH also reversed HPV (by 75%), although 10 μM t-BOOH had a negligible effect.

Cultured rat PASMC and RASMC were transfected with HyPer-dMito or HyPer-dCyto to monitor H_2O_2 levels in the mitochondria and cytoplasm, respectively [296]. In PASMC, hypoxia ($pO_2 \sim 20$ Torr) decreased H_2O_2 levels in both the mitochondria and cytoplasm. In contrast, hypoxia had no effect on mitochondrial ROS and increased

cytoplasmic ROS in RASMC. Hypoxia also increased the $[Ca^{2+}]_{cyt}$ in PASMC. This rise in $[Ca^{2+}]_{cyt}$ was completely reversed by 1 μ M t-BOOH and subsequently restored by application of catalase. The effects of hypoxia on $[Ca^{2+}]_{cyt}$ and mitochondrial [ROS] were mimicked by rotenone, which also diminished both the concentration of H_2O_2 generated by LM, and the reversal of HPV induced LM-conditioned medium. Taken together, these results support the idea that mitochondria in PASMC produce more ROS than those in RASMC under normoxic conditions, that hypoxia causes a fall in ROS production which increases PASMC $[Ca^{2+}]_{cyt}$, and that reducing and oxidizing stimuli cause changes in $[Ca^{2+}]_{cyt}$ in the cells which are consistent with the Redox Theory, as does block of the proximal ETC with rotenone.

Ndufs2 is a subunit of complex I which plays an important role in the binding and reduction of ubiquinone by complex I [297–300]. It had been demonstrated [190] that conditional knockout of Ndufs2 in CBCC prevented activation of the carotid body by hypoxia, leading Dunham-Snary et al. [123] to investigate its involvement in HPV. They found that the mRNA expression of Ndufs2 was higher in PASMC than in RASMC, as was its protein expression (normalized to TOMM20, a protein expressed in the outer mitochondrial membrane). Moreover, complex I in PASMC was enriched in Ndufs2s compared to RASMC.

siRNA knockdown of Ndufs2 in PASMC inhibited respiration and increased cellular levels of NADH, consistent with its role in the enabling of electron flow in the ETC. Although the expression of complex 1 was not affected by the knockdown of Ndufs2, its activity, assessed using protein immunocaptured from cell homogenates, was depressed. Knockdown of Ndufs2 also abolished the hypoxia-induced rise in $[Ca^{2+}]_{cyt}$ and decreased the production of H_2O_2 (assessed using Amplex Red) by PASMC under normoxic conditions. In contrast, siRNA knockdown of Uqcrrf1, the protein containing the Rieske Fe-S cluster in complex III, or of Cox4i2, a Cox subunit proposed to contribute to O_2 sensing [142] did not prevent the hypoxia-induced increase in $[Ca^{2+}]_{cyt}$ in PASMC. Knockdown of Ndufs1, a protein in the N-domain of complex I, was also without effect on the rise in $[Ca^{2+}]_{cyt}$ evoked by hypoxia.

In further experiments, siRNA directed against Ndufs2 or control siRNA was administered once to rats via airway nebulization. After 48 h, compared to the controls, the si-Ndufs2 rats demonstrated diminished levels of Ndufs2 mRNA in lung homogenates and exhibited a profound fall in Ndufs2 levels co-localizing with smooth muscle actin, as revealed by immunofluorescence studies of small PA in lung sections. The animals were anesthetized and PAP was measured in vivo under normoxic and hypoxic (10% O_2) conditions. PAP under normoxic conditions was not different in the two groups, but hypoxia caused a much smaller increase in PAP in the si-Ndufs2 rats compared to the controls. Similarly, treatment of mice with Ndufs2 siRNA over a period of more than a week had no effect on basal PAP, but profoundly depressed HPV, compared to animals treated with control siRNA. Likewise, the increase in PAP evoked by rotenone in the controls was almost absent in the si-Ndufs2 rats, whereas the response to phenylephrine was enhanced in the latter group. This may have been due to the down-regulation of Kv1.5 which also occurred in the si-Ndufs2 animals. They also found that exposing mouse lungs in vitro to hypoxia for 30 min increased the ratio of reduced to total Ndufs2. Hypoxia also inhibited complex I activity, an effect mimicked by treatment with the reducing agent dithiothreitol (DTT). They speculated that this might be due to the reduction of the mitochondrial milieu caused by hypoxia.

Based on the specific inhibitory effect of Ndufs2s knockdown on the rise in $[Ca^{2+}]_{cyt}$ evoked by hypoxia, as well as their evidence that hypoxia caused a fall in cytoplasmic H_2O_2 and that HPV was reversed by oxidants, the authors proposed that HPV was

triggered by a fall in ROS production which they suggested was due to inhibition of the production of uncoupled electrons (i.e., electrons leaking from the ETC) at Ndufs2.

In accordance with the concept that Ndufs2 is involved in producing ROS in a PO₂-dependent manner, a subsequent paper from this laboratory [301] reported that the I_Q site, in which Ndufs2 plays a vital role in the transfer of electrons from complex I to ubiquinone [298], is responsible for the increase in ROS production associated with a rise in PO₂ in the ductus arteriosus.

A letter by Hüttemann et al. [302] criticized the conclusion that COX4i2 was not involved in HPV on the basis that its knockdown in the Dunham-Snary study, which amounted to ~55%, was insufficient to significantly affect its role in O₂ sensing. Their contention was based on their unpublished finding that HPV was largely intact in COX4i2^{+/-} mice, whereas it was abolished in homozygous knockouts [142].

In a response letter, Dunham-Snary & Archer [303] expressed their belief in the validity of their findings, and criticized the study by Hüttemann and colleagues [142] based on its use of a very low pO₂ to elicit HPV, and on the small amplitude of the HPV they obtained. They also noted that their conclusion that Ndufs2 was crucial in O₂ sensing in PASM C was consistent with the earlier demonstration by Fernandez-Aguera et al. [190] that it was the O₂ sensor in CBCC, supporting the concept that common O₂ sensing mechanisms are operative in various oxygen-sensing tissues [280]. However, they failed to mention that the proposed mechanism by which Ndufs2 was contributing to O₂ sensing in CBCC [190] was opposite to theirs, since Fernandez-Aguera et al. had shown that Ndufs2 contributes to O₂ sensing in these cells by virtue of its ability to increase, rather than decrease, mitochondrial ROS production in response to hypoxia. Based on subsequent work [191], this laboratory proposed that hypoxia increases ROS production in CBCC by causing reduction of the ubiquinone pool, thereby lengthening the dwell-time of electrons at the Ndufs2 quinone binding site and increasing the probability that they will react with O₂ at this point. It has also been reported that an increase in ROS production dependent on NDUF2.1, the *C. elegans* ortholog of Ndufs2, is responsible for the locomotor response to hypoxia in these organisms [304]. Similarly, Hernansanz-Agustin et al. [195] presented evidence that, whereas hypoxia caused a marked increase in cytoplasmic [H₂O₂], which they measured using HyPer, in bovine aortic endothelial cells transfected with scrambled siRNA, the opposite effect occurred if Ndufs2 was knocked out. They also found that Ndufs2 knockout increased the basal (normoxic) cytoplasmic [H₂O₂].

The results and implications of the experiments in which protein components of the mitochondrial complexes were knocked down can also be questioned. Ndufs2 knockdown depressed respiration and raised the cellular NADH concentration, indicative of a decrease in the activity of the ETC, and has been shown to cause additional and widespread effects on mitochondrial function, including the depolarization of $\Delta\Psi_m$ [305]. This may occur because the reduction of CoQ, which is thought to initiate proton translocation by complex I, requires electrons donated by an aspartate-histidine pair in the β_1 - β_2 loop of Ndufs2 [306]. A change in the configuration of Ndufs2 may also play a role in the active-to-deactive transition of complex I, which was proposed [194] to mediate the increase in ROS responsible for HPV [217]. Thus, it is not clear that the effects of Ndufs2 knockdown on the responses to hypoxia observed by Dunham-Snary and colleagues can be specifically ascribed to the changes in cellular H₂O₂ dynamics which it caused.

It is also curious that the knockdown of Ndufs1 and Uqcrcf1 apparently had no effect on the response to hypoxia in PASM C. Ndufs 1 is one of the 14 catalytic-core subunits required for complex I to function, and contains three (N1b, N4, N5) of the seven Fe-S clusters which form a 'wire' carrying electrons from the NADH binding pocket to the ubiquinone binding site [307]. Since Ndufs1 is upstream of N2, the Fe-S cluster with which Ndufs2 interacts, the lack of effect of its knockdown on the response to hypoxia would

also appear to rule out a role for *Ndufs2* in regulating ROS production at either I_Q or I_F . More generally, knockout of either *Ndufs1* or *Uqcrrf1*, which is required for electron flow from Q_o to cytochrome *c* and thence to *Cox* and O_2 , should suppress mitochondrial respiration and depolarize $\Delta\Psi_m$, thereby presumably causing the dysfunction of any redox-mediated mitochondrial O_2 sensing mechanism [308,309].

5.1.2. The Involvement of ROS in PASM C K^+ Channel Inhibition During Hypoxia

The Redox Theory posits that membrane depolarization due to the hypoxia-induced closure of K_v channels, particularly those incorporating the $K_v1.5$ subunit, is a primary effector mechanism for HPV, and that this inhibition is mediated by a fall in $[H_2O_2]$ and/or an increase in $[NAD(P)H]$ in the cytoplasm.

The concept that K_v channel closure contributes to HPV is supported by abundant evidence [17,288,310–320], although other processes, including intracellular Ca^{2+} release, store-operated Ca^{2+} entry (SOCE), and Ca^{2+} sensitization, are also important [3,10,12,14], and additional depolarizing mechanisms (e.g., TASK-1; non-selective cation channels) [33,321,322] have been identified.

On the other hand, the proposal that this inhibition of K_v channels during HPV requires a reduction in the cytoplasmic redox balance remains controversial. In keeping with the focus of this review on the O_2 sensing rather than effector mechanisms of HPV, and because the question of how K_v channels in PASM C are regulated by redox mechanisms has been critiqued extensively [3,53,323,324] but seems not to have moved any closer to being resolved over the past decade, we will consider this issue only briefly.

A number of laboratories investigated the redox regulation of K_v channels in PASM C by monitoring whole cell or membrane patch currents while applying oxidants, reductants, or ETC blockers, which were used to decrease mitochondrial ROS production. Both rotenone and antimycin (10 μM) were shown to block the K^+ current in PASM C [156,272]. The effect of rotenone is in accordance with Redox Theory, although that of antimycin is difficult to interpret, since it is generally thought to increase mitochondrial ROS production [121]. A more detailed study [325] showed that although antimycin, myxothiazol, and rotenone depressed the K_v current in PASM C at positive potentials, they caused a hyperpolarizing shift in its activation threshold, implying that these blockers should open rather than shut K_v channels in the physiological range of membrane potentials. The authors provided evidence that the ETC blockers were exerting these effects on the K_v current by increasing the cytoplasmic cellular $[Mg^{2+}]$. This could be due to mitochondrial depolarization, an effect exerted by all ETC inhibitors, which could explain why both rotenone and antimycin inhibited the current at positive potentials. To complicate matters further, both rotenone and antimycin were shown to inhibit the K^+ current with IC_{50} values of $\sim 1 \mu M$ in H146 (small cell lung carcinoma) cells by a mechanism which was independent of the mitochondria [326].

The effects of reductants and oxidants on K^+ currents are summarized in Tables 1 and 2, respectively. The majority of studies produced evidence supporting the viewpoint of the Redox Theory with regard to the redox regulation of PASM C K_v channels by hypoxia. However, other investigations reported observations inconsistent with this view [142,327,328] and there is also evidence that hypoxia-induced K_v channel inhibition is mediated by a different mechanism: a rise in $[Ca^{2+}]_{cyt}$ due to Ca^{2+} release from the sarcoplasmic reticulum [329,330]. Moreover, LY83583, which causes intracellular superoxide production, increased the K_v current in the physiological range of membrane potentials in PASM C, as predicted by the Redox Theory, but caused PA constriction rather than the relaxation that would be predicted to arise from K_v channel activation [331]. This suggests that the contractile response of these cells to alterations in the cytoplasmic redox state might not depend exclusively on changes in K_v channel activity.

Yandjue et al. [262] recently showed that the effect of H₂O₂ on the Kv1.5 current expressed in oocytes is strongly dependent, not only on its concentration, but on the potential at which the current is evoked. Application of low concentrations of extracellular [H₂O₂] (0.01–100 µM), a range which is likely to cause changes in its cytoplasmic concentration which are physiologically relevant (see Section 4), tended not to affect or inhibit the current at −20 mV, whereas it was increased by concentrations above 1 mM. Although the implications of these results for the redox regulation of Kv1.5 during HPV remain unclear, the observations made in this paper highlight the need to study the effect of redox mechanisms on channel activity during conditions which more realistically simulate the effects of hypoxia on specific redox couples and oxidant concentrations. More generally, little is known at the molecular level about the redox regulation of the Kv1.x and Kv2.x isoforms thought to play a role in HPV, and the results of the few studies which have examined this in more depth [276,332,333] suggest that an accurate determination of how redox mechanisms (and other factors such as [Ca²⁺]_{cyt} or channel phosphorylation [27,329,334]) contribute to Kv channel closure during HPV will require a more refined and concerted approach to unraveling the effects of hypoxia and redox agents on these channels than has previously been attempted. It will be particularly important to characterize the effects of hypoxia and perturbations on specific redox couples on Kvβ-subunit function, the oxidation of cytoplasmic cysteine residues on the Kv1.5α subunits, and the membrane trafficking of Kv channels (which can change rapidly [332,335]) in native PASMCM.

Table 1. (a) Effects of oxidants on K⁺ currents in PASMCMs. See next page. (b) Effects of reductants on K⁺ currents in PASMCMs.

(a)					
Source of PASMCM	Oxidant (in µM)	Current	Effect on K ⁺ Current	Comments	Ref.
Rabbit intralobar PA	GSSG (5000)	Single-channel BK _{Ca}	P _{open} ↑ by 450%		[336]
4th- & 5th-order rat PA	diamide (100)	Whole cell K ⁺	~120% ↑	Current activated by diamide was very noisy (i.e., mainly BK _{Ca} ?)	[337]
Rabbit intralobar PA	GSSG (5000)	Single-channel BK _{Ca}	P _{open} ↑ by 455%		[338]
Rabbit intralobar PA	DTNB (1000)	Single-channel BK _{Ca}	P _{open} ↑ by 504%		[338]
Rabbit intralobar PA	DTNB (200)	Whole-cell Kv	↑ ~260% at +20 mV	Effect of DTNB shown to be intracellular.	[339]
Rabbit fetal resistance PA	DTNB (1000)	Whole-cell Kv	↑ ~70% at +70 mV	K ⁺ current not identified but appearance (low noise) suggests that it was entirely or predominantly Kv.	[340]
Rat small PA	GSSG (2000)	Whole-cell K ⁺	↑ ~40% at +50 mV in hypoxic cells	Basal and enhanced K ⁺ currents not identified, but low noise suggests they were Kv.	[21]
Rat small PA (cultured)	H ₂ O ₂ (50)	Whole-cell Kv	↑ by ~55% at +60 mV	BK _{Ca} and K _{ATP} currents blocked to isolate Kv.	[341]
Rat small PA (cultured)	diamide (100)	Whole-cell Kv	↑ by ~45% at +60 mV	BK _{Ca} and K _{ATP} currents blocked to isolate Kv.	[341]
Rat 2nd- & 3rd-order PA	t-BOOH (5 and 100)	Whole-cell Kv	↓ by 30% at +60 mV	Similar inhibition of Kv with 5 and 100 µM t-BOOH. [Ca ²⁺] _{cyt} strongly buffered to block BK _{Ca} , thereby isolating Kv.	[327]

Rat 3rd- & 4th-order PA	LY83583 (10)	Whole cell K _v	Activation shifted ~11 mV in hyperpolarizing direction	BK _{Ca} and K _{ATP} currents blocked to isolate K _v . Shift in activation reflects increased K _v current amplitude at physiological membrane potentials.	[331]
Rat PA 0.2–0.5 mm dia.	t-BOOH (10)	Whole-cell K _v	↓ by ~35% at +60 mV	[Ca ²⁺] _{cyt} strongly buffered to block BK _{Ca} , thereby isolating K _v .	[328]
Non-passaged cultured PASM from mouse precapillary PA	H ₂ O ₂ (0.124)	Whole-cell K _v	↓ by ~60% at +60 mV	BK _{Ca} and K _{ATP} currents blocked to isolate K _v .	[142]
(b)					
Source of PASM	Reductant (Conc. in mM)	Current	Effect on K ⁺ current	Comments	Refs.
Rat main and 1st/2nd-order PA (cultured)	GSH (10)	Whole-cell K _v	~25% decrease in steady-state K ⁺ current, 16 mV depolarization	10 mM EGTA in pipette solution to suppress BK _{Ca} current, thereby isolating K _v current.	[342]
Rabbit small intra-lobar PA	GSH (5)	Single-channel BK _{Ca}	P _{open} variably decreased	GSH applied to cytoplasmic face of the membrane	[336]
Rat small PA	GSH (2)	Whole-cell K ⁺	~40% ↓ at +70 mV	K ⁺ current not identified but low noise suggests it was K _v .	[21]
Rat 4th- & 5th-order PA	Co-enzyme Q (0.1)	Whole-cell K ⁺	~40% ↓ at +40 mV	K ⁺ current not identified but low noise suggests it was K _v .	[337]
Rat 4th- & 5th-order PA	Duroquinone (0.1)	Whole-cell K ⁺	~40% ↓ at +40 mV	K ⁺ current not identified but low noise suggests it was K _v .	[337]
Rabbit small intra-lobar PA	DTT (5)	Single-channel BK _{Ca}	19% ↓ in P _{open}	DTT applied to cytoplasmic face of membrane.	[338]
Rabbit small intra-lobar PA	GSH (5)	Single-channel BK _{Ca}	34% ↓ in P _{open}	GSH applied to cytoplasmic face of membrane.	[338]
Rabbit small intra-lobar PA	DTT (5)	Whole-cell K _v	~30% ↓ at +20 mV	10 mM BAPTA in pipette solution to suppress BK _{Ca} current, thereby isolating K _v current.	[339]
Rat PA	GSH (2)	Whole-cell K ⁺	~50% ↓ at +70 mV	GSH applied in pipette solution. K ⁺ current not identified but low noise suggests it was K _v . (n = 1) no statistical analysis.	[343]
Rat PA	GSH (5)	Single K ⁺ channels, not identified	~70% ↓ in P _{open}	(n = 1) no statistical analysis. GSH applied in pipette solution.	[343]
Rabbit fetal resistance PA	DTT (3)	Whole-cell K ⁺	33% ↓ at +50 mV ~20% ↓ at −10 mV	K ⁺ current not identified but low noise suggests it was K _v .	[340]
Rabbit adult resistance PA	DTT (3)	Whole-cell K ⁺	↓ 19% at +50 mV	K ⁺ current not illustrated or ascribed to any specific type(s) of channel.	[340]
Rat 2nd- & 3rd-order PA	Catalase (100U/mL)	Whole-cell, predominantly or entirely K _v	Hypoxia ↓ the K _v current by 33%. This effect was prevented by including catalase in the pipette solution.	Result provides indirect evidence that hypoxia inhibited K _v current by raising cytoplasmic H ₂ O ₂ concentration. 10 mM EGTA in pipette solution to suppress BK _{Ca} current, thereby isolating K _v current.	[328]

n.b.: Except where noted, PSMCs for these studies were freshly isolated on the day that currents were recorded.

5.2. The Mitochondrial ROS Theory of HPV

The proposal that HPV is caused by a fall in cellular [ROS] was challenged by Marshall et al. [344], who reported that hypoxia increased ROS in cultured PASM from fetal calves, and by Killilea et al. [345], who made a similar observation in cultured PASM from rats. Abdalla & Will [346] had also observed that HPV in guinea pig PA was enhanced by two SOD inhibitors, and suggested that HPV was due to a rise in cellular superoxide levels.

Whereas Marshall et al. ascribed their results to an activation of Nox, Paul Schumacker and colleagues proposed in 2001 that hypoxia promotes ROS production by PASM mitochondria, leading to a rise in cytoplasmic [ROS] that results in the stimulation of redox-sensitive effector pathways which cause HPV [23]. This concept grew out of observations by Schumacker's laboratory that hypoxia increased mitochondrial ROS production in cardiac myocytes [347] and HEP3b cells [348].

This 'Mitochondrial ROS hypothesis' was supported and extended in subsequent papers from this and other laboratories, several of which have presented evidence that the site of ROS production is complex III. It has also been proposed that the increase in mitochondrial ROS leads to additional ROS production by Nox which enhances the pressor response to hypoxia (see Section 5.2.5). A schematic of the mitochondrial ROS hypothesis, which has grown out of the investigations described in Section 5.2, is shown in Figure 7.




Figure 7. The mitochondrial ROS hypothesis of HPV. According to this proposal, hypoxia causes an increased ROS production by the Qo site of complex 3. This raises the concentration of ROS in the mitochondrial intermembrane space, thereby generating a cytoplasmic ROS signal which causes contraction in PSMCs by acting on several HPV effectors to increase Ca²⁺ release from the sarcoplasmic reticulum, Ca²⁺ influx, and Ca²⁺ sensitization (see Section 5.2.6 for a brief discussion of these

effectors and references supporting their ROS-sensitivity). A comprehensive description of the evidence implicating these effectors in HPV can be found in [3].

Waypa et al. [23] studied HPV recorded in isolated ventilated rat lung and microvascular PASM in primary culture. Hypoxia was imposed by reducing the $[O_2]$ in the ventilating gas from 21 to 2%. HPV was abolished by rotenone. Myxothiazol, which blocks electron transport in complex III at the ubiquinol oxidation site (Q_o), had a similar effect. Conversely, 1 ng/mL antimycin, which binds to complex III more distally at the Q_i site, had no effect on HPV, and cyanide augmented the amplitude of HPV. All of the ETC blockers except myxothiazol caused a transient contraction.

Also, the antioxidant pyrrolidinedithiolcarbamate (PDTC) reversibly decreased HPV by 65% and blocked hypoxic contraction of PASM. The glutathione peroxidase mimetic ebselen caused a profound and irreversible block of HPV, and diethyldithiocarbamate (DDC), a cytosolic Cu, Zn SOD blocker, inhibited HPV by 68%. Application of DIDS, an anion channel blocker used here to prevent mitochondrial superoxide from entering the cytoplasm through the mitochondrial inner membrane anion channel (IMAC), inhibited HPV to a similar extent, and also blunted the contraction caused by antimycin. At the concentrations used in these experiments, none of these blockers affected the pressor response to the TXA_2 agonist U46619, which was used as a negative control to detect non-specific effects on contraction. In contrast, 3 μM apocynin (used to block Nox) abolished both HPV and the U46619 contraction, whereas it had no effect on either response at a lower concentration (0.3 μM). They also found that hypoxia caused a largely reversible increase in the production of ROS by PASM (monitored using DCHF-DA), which was attenuated by myxothiazol. In addition, HPV was diminished by 70% in PASM p^o cells, which had been incubated in ethidium bromide for two weeks to deplete cellular mitochondria [270].

Based on these results, the authors proposed that hypoxia induces an increase in mitochondrial ROS production upstream of the Q_i site of complex III, and that this causes an increase in the cytoplasmic $[H_2O_2]$ which acts through several effector mechanisms to cause HPV. The authors pointed out that drugs acting distal to ubiquinone, including antimycin, tend to increase ROS during normoxia by increasing the reduction of the ubiquinone pool [349], whereas a more proximal block would suppress ROS production by oxidizing this pool. The lack of HPV in p^o cells, and the inability of apocynin to exert a selective block of HPV, supported the idea that ROS produced by the mitochondria rather than Nox were regulating HPV (n.b.: however, apocynin is not a dependable blocker of Nox; [350]). Based on the block of HPV by DIDS, they also suggested that hypoxia might alternatively increase the entry of superoxide into the cytoplasm by opening IMAC, although in subsequent papers this idea was not pursued.

Leach et al. [351] employed a pharmacological approach to evaluate the role of the mitochondrial complexes in HPV using isolated rat intrapulmonary arteries. They observed that the ETC blockers rotenone, myxothiazol, cyanide and the complex II substrate succinate had no effect on basal tone under normoxic conditions. Rotenone caused an increase in NAD(P)H (assessed by measuring tissue autofluorescence) similar in amplitude to that induced by anoxia. This was not reversed by application of 5 mM succinate, consistent with a complete block of the ETC at complex I by rotenone. Hypoxia (PO_2 of 15–18 Torr) applied to arteries which were slightly pre-constricted with $PGF_{2\alpha}$ caused a biphasic HPV, as typically observed in this preparation under these conditions. Application of either rotenone or myxothiazol abolished both phases of HPV. Importantly, rotenone did not block either phase of HPV if it was co-applied with succinate, whereas succinate did not reverse the inhibition of HPV by myxothiazol. These results indicated that HPV in these arteries required the provision of electrons to the myxothiazol binding site (i.e., Q_o)

of complex III, which succinate would be able to restore in the presence of rotenone but not myxothiazol. The authors pointed out that, whereas this effect of succinate was consistent with the concept that HPV was caused by an increase in ROS production by the ETC at or distal to Q_o , it was difficult to explain whether hypoxia was causing HPV by mimicking the effect of rotenone and inducing a decrease in mitochondrial ROS production, as proposed by the Redox Theory. The lack of effect of rotenone on basal tension was also inconsistent with the Redox model. Subsequent studies have shown that succinate can also increase ROS production by causing reversed electron transport (RET) through complex I [112,190]. Although this would have been favored by the combination of succinate and myxothiazol, which should highly reduce the ubiquinone pool [188], it seems unlikely that ROS production by RET could explain the restoration of HPV by succinate in this study, since this would have been prevented by rotenone [188,352].

Waypa et al. [353] used fura 2 to investigate the effect of ETC blockers and other agents on the hypoxia-induced rise in $[Ca^{2+}]_{cyt}$ in PASM. Exposure to medium equilibrated with 1.5% O_2 caused a sustained rise in $[Ca^{2+}]_{cyt}$. This was attenuated by DPI (diphenyleneiodonium, here used to block complex 1), and by rotenone and myxothiazol, but was unaffected by antimycin and cyanide. None of these drugs significantly affected the increases in $[Ca^{2+}]_{cyt}$ evoked by angiotensin 2 or 50 μM H_2O_2 . The responses to hypoxia and exogenous H_2O_2 , but not angiotensin 2, were diminished in PASM in which catalase had been overexpressed by recombinant adenovirus transduction. Application of cyanide caused a rise in $[Ca^{2+}]_{cyt}$ which was antagonized by myxothiazol and was also smaller in PASM overexpressing catalase. The contrasting effects on HPV of the proximal (rotenone, DPI, and myxothiazol) as compared to distal (antimycin and cyanide) ETC blockers led them to propose that mitochondrial ROS production at the Q_o/III_{CoQ} site in complex III was crucial for triggering the rise in $[Ca^{2+}]_{cyt}$ during HPV. The observation that HPV was blocked by catalase overexpression suggested that H_2O_2 rather than superoxide was the important ROS for HPV, consistent with the idea that H_2O_2 is generated from superoxide by MnSOD in the intermembrane space of the mitochondria, and then crosses the outer membrane to access the cytoplasm.

Weissmann et al. [354] carried out a comprehensive study of the effects on HPV of blockers of complexes I-III, and also of mitochondrial uncouplers, which should decrease ROS production. The aim of the investigation was to test the predictions of the Mitochondrial ROS model as set out by Waypa et al. [36]. In this and other investigations carried out by this laboratory [155,355–357], HPV was recorded in isolated perfused rabbit lungs, which were ventilated via the trachea, and the PO_2 in the ventilating gas mixture was lowered from 21 to 3% for 10 min to reduce the alveolar PO_2 from ~160 to ~23 Torr and evoke the initial transient component of HPV (Phase 1). The thromboxane A_2 agonist U46619 was used as a negative control. Experiments were repeated in the presence of the eNOS blocker L- N^G -monomethyl-L-arginine to determine whether there were any confounding effects of the mitochondrial blockers which were being exerted due to interference with NO release. Whereas some of the results supported the Mitochondrial ROS hypothesis (e.g., blocking complexes I and II antagonized HPV), others did not (e.g., antimycin did not cause a sustained contraction). The authors were able to conclude only that mitochondria are somehow involved in Phase 1 HPV. Importantly, they also observed that most of the blockers they studied strongly attenuated the response to U46619, and/or were interacting with the NO system, suggesting that these drugs have off-target actions on contraction which could obscure the effects they exert on HPV by altering mitochondrial ROS production.

Liu et al. [358] used perfusion myography to evaluate the role of ROS in HPV (evoked by a pO_2 of 29 Torr) in 7th-generation branches of porcine PA. HPV was abolished when the solution contained 150 U/mL SOD or SOD + 200 U/mL catalase. They used lucigenin

and also EPR spectroscopy with the spin probe 5-5-dimethyl-1-pyrroline-N-oxide (DMPO) in an attempt to determine the effect of hypoxia on cellular ROS in the arteries, but although both approaches yielded results suggestive of an increase, in neither case were the results significant (n.b.: the use of DMPO may also have been problematic [359]). However, hypoxia (PO_2 of 34 Torr) did cause a rapid and largely reversible increase in DCFH fluorescence in primary cultured PASM. This was strongly suppressed by SOD and completely prevented by both catalase and SOD + catalase. The authors confirmed, using FITC-labeled SOD and catalase, that both enzymes were entering the cells. Notably, the observation that SOD and SOD + catalase similarly suppressed HPV suggested it was mainly an increase in cellular [superoxide] rather than $[\text{H}_2\text{O}_2]$ which was causing contraction, a finding which appears to be inconsistent with the current view [48] that H_2O_2 is the crucial ROS for cellular signaling.

Weissmann et al. [357] compared the effects of ETC blockers on the 1st and 2nd phases of HPV in rabbit lungs. The complex I blocker MPP⁺ (1-methyl-4 phenylpyridinium), applied at a concentration which slightly blocked Phase 1, strongly blocked Phase 2. Similar semi-selective effects on Phase 2 were exerted by rotenone and cyanide. In contrast, 3-NPA (3-nitropropionic acid; blocks complex III) and antimycin caused a similar and marked suppression of both phases.

In view of the shortcomings of ROS indicators such as lucigenin, DCFH, and luminol (see Section 7.4), Guzy et al. [360] developed a novel fluorescence resonance energy transfer probe, HSP-FRET, in order to evaluate the role of mitochondrial ROS in hypoxia-induced stabilization of HIF-1 α . Their study, carried out in 143B cells, showed that application of exogenous H_2O_2 and hypoxia, but not NO, increased the FRET signal, and that stigmatellin (which blocks the ETC at the Q_o site in complex III), suppressed the response to hypoxia. Their results suggested that HSP-FRET was able to detect H_2O_2 , and that, at least in these cells, hypoxia increases the production of mitochondrial ROS, which then enters the cytoplasm.

This laboratory [361] then used HSP-FRET to determine whether hypoxia caused an increase in ROS in PASM. They described HSP-FRET as being predominantly expressed in the cytoplasm, although specific evidence for this was not provided. In PASM transfected with HSP-FRET, hypoxia (1.5% O_2) caused a gradual increase in ROS which started after ~3 min. A simultaneous increase in $[\text{Ca}^{2+}]_{\text{cyt}}$, assessed using fura 2, also occurred, and similar results were obtained in experiments using YC2.3, a FRET-based Ca^{2+} sensor. Hypoxia also decreased the GSH/GSSG ratio, considered to reflect the general cell redox state [362], from ~50 to ~30. The effects of hypoxia on oxidant signaling and $[\text{Ca}^{2+}]_{\text{cyt}}$ were suppressed by the antioxidants PDTC and N-acetyl-L-cysteine (NAC), neither of which affected the signal from either probe under basal conditions. Overexpression of glutathione peroxidase in order to deplete cytoplasmic H_2O_2 suppressed the effects of hypoxia on both $[\text{Ca}^{2+}]_{\text{cyt}}$ and oxidant signaling, as did overexpression of catalase in either the mitochondria or the cytoplasm. Conversely, overexpression of cytosolic Cu, Zn SOD (SOD 1) had no significant effect on the responses of HSP-FRET or YC2.3 to hypoxia, while overexpression of mitochondrial Mn SOD (SOD 2) similarly did not affect oxidant signaling and enhanced the $[\text{Ca}^{2+}]_{\text{cyt}}$ rise induced by hypoxia. In a subsequent paper from this group, overexpression of Prx5 in the mitochondrial IMS was similarly found to diminish the rise in $[\text{Ca}^{2+}]_{\text{cyt}}$ evoked by hypoxia in these cells [363].

These results supported the concept that hypoxia causes an increase in cytoplasmic H_2O_2 and an oxidizing shift in the redox balance of PASM resulting from mitochondrial ROS production. The observation that PDTC and NAC blocked the effects of hypoxia but did not themselves cause any response argued against the proposal of the Redox Theory that cytoplasmic oxidation under basal conditions maintains vasorelaxation by causing an ongoing stimulation of Kv channels.

A study by Wang et al. [364] which examined the role of mitochondrial ROS in the hypoxia-induced rise in $[Ca^{2+}]_{cyt}$ in freshly isolated mouse PASMC yielded results similar to those reported by Waypa and colleagues. They found that hypoxia (1% O_2 for 5 min) increased ROS levels (assessed using DCFH and lucigenin) and $[Ca^{2+}]_{cyt}$. Pharmacological blockade of complexes 1, 2, and the Q_o site of complex III did not affect the basal ROS or $[Ca^{2+}]_{cyt}$ signals, but strongly antagonized the effects of hypoxia. In contrast, block of complex III at the Q_i site with antimycin or at complex 4 with NaN_3 had no effect on ROS levels or $[Ca^{2+}]_{cyt}$ under either normoxic or hypoxic conditions. The effects of hypoxia on ROS levels and $[Ca^{2+}]_{cyt}$ in PASMC from mice overexpressing glutathione peroxidase or catalase were smaller than those observed in control mice, whereas they were enhanced in PASMC from glutathione peroxidase knockout mice. They also found that application of 51 μM H_2O_2 caused an increase in the DCFH signal similar to that evoked by hypoxia but resulted in a much smaller increase in $[Ca^{2+}]_{cyt}$. This led them to propose that mitochondrial H_2O_2 production was necessary but not sufficient to evoke the full hypoxia-induced increase in $[Ca^{2+}]_{cyt}$, or alternatively that this response might depend on the mitochondrial compartmentalization of ROS, which would not occur when peroxide was applied exogenously. A later investigation [365] similarly reported that hypoxia-induced increases in $[Ca^{2+}]_{cyt}$ in cultured human PASMC were prevented by blocking complexes 1, 2, and Q_o , but were unaffected by block of Q_i or complex IV, and that HPV in small PA from humans showed the same pattern of responsiveness to these drugs.

In order to determine whether hypoxia-induced redox signaling in PASMC was compartmentalized, Waypa et al. [366] transfected PASMC with Ro-GFP, a genetically encoded and ratiometric redox indicator. Ro-GFP expression was targeted to the mitochondrial matrix (Mito-Ro-GFP) or the mitochondrial intermembrane space (IMS-Ro-GFP) and non-targeted Ro-GFP was used to assess the cytoplasmic redox state. Rather than providing a readout of $[H_2O_2]$, Ro-GFP fluorescence is sensitive to the redox state of the GSH/GSSG couple [367].

Hypoxia (1.5% O_2) caused a progressive increase in cytoplasmic RoGTP oxidation from 19 to 35% over 30 min. IMS-Ro-GFP had a higher level of oxidation under normoxic conditions (48%) but was still significantly oxidized (to 62%) by 30 min of hypoxia. In contrast, Mito-Ro-GFP, which was even more highly oxidized under normoxic conditions (63%), became more reduced during hypoxia (to 44%). Overexpression of catalase in the cytoplasm had no effect on the normoxic oxidation level of Ro-GFP but prevented its hypoxia-induced oxidation. Overexpression of catalase in the mitochondrial matrix had no effect on the oxidation of Mito-Ro-GFP under either normoxic or hypoxic conditions. These results imply that changes in H_2O_2 production during hypoxia influence the redox potential in the cytoplasm but not in the mitochondrial matrix. Hypoxia caused the same pattern of compartmentalized changes in Ro-GFP oxidation in renal artery smooth muscle cells. However, experiments with YC2.3 showed that whereas hypoxia caused an increase in $[Ca^{2+}]_{cyt}$ in PASMC, it did not do so in the renal arteriolar cells. This implies that the interaction between increases in cytoplasmic ROS and the various effector mechanisms which regulate vascular tone must differ in the two types of arteries.

These investigations can be criticized because they were carried out in cultured smooth muscle cells which may have undergone phenotypic changes from their in situ state. In order to characterize oxidant signaling under more physiological conditions, Desireddi et al. [368] transfected mouse lung slices with Cyto-Ro-GFP, a process which required their incubation in culture medium for two days. The vascular endothelium was labeled with Cell Tracker Red so the vascular lumen could be imaged and used to measure contraction. The effect of hypoxia on $[Ca^{2+}]_{cyt}$ was assessed in parallel experiments using fura 2, although this was done after only one day of organ culture, as Ca^{2+} signaling was lost thereafter.

Small PA in lung slices contracted when exposed to angiotensin 2, high K^+ PSS, and hypoxia after the two-day period in organ slice culture. Hypoxia (1.5% O_2) caused a rapid and sustained increase in Ro-GFP oxidation from 11 to 17%, which reversed upon re-oxygenation. Overexpression of catalase in lung slices had no effect on basal Ro-GFP oxidation but almost abolished the effect of hypoxia. Hypoxia caused an immediate increase in $[Ca^{2+}]_{cyt}$ which was diminished by catalase overexpression, and was abolished by EUK-134, a SOD-catalase mimic [369]. The increase in $[Ca^{2+}]_{cyt}$ was absent in slices superfused with Ca^{2+} -free PSS containing 2.5 mM EGTA, suggesting that it was due to the influx of extracellular Ca^{2+} . However, this concentration of EGTA can deplete releasable intracellular Ca^{2+} stores [370], so these might also have been involved.

These results demonstrated that hypoxia caused a reversible increase in Ro-GFP oxidation (i.e., oxidation of the GSH/GSSG redox couple) in small PA which, although exposed to culture conditions for two days, retained the ability to contract. It is noteworthy, however, that the size of the contractions was small compared to those previously described in freshly prepared mouse lung slices [371,372], and the proportion of arteries demonstrating a contraction to hypoxia was not stated. Also, the loss of Ca^{2+} signaling caused by the two-day incubation period suggests that changes in the PASM phenotype had occurred at the time when the ROS measurements were carried out. In addition, a potential danger inherent in the use of cultured tissue slices is that PO_2 in the core of the slice may be much lower than that in the bulk solution due to cellular oxygen consumption, which could progressively diminish the viability of the preparation [373]. Cellular O_2 consumption might also result in anoxia instead of hypoxia in the slice core during an acute exposure to a low PO_2 .

5.2.1. Role of Mitochondrial Ca^{2+} and the Rieske Iron-Sulfur Protein in Hypoxia-Induced ROS Production

The Rieske iron-sulfur protein (RISP; also termed ubiquinol-cytochrome c reductase, Rieske iron-sulfur protein 1, and Uqcrcfs1) is a subunit of complex III. RISP contains a 2Fe-2S cluster which accepts electrons from the Q_o site and passes them on to cytochrome c_1 , and has been proposed to play a pivotal role in mediating a hypoxia-induced increase in ROS production by complex III [374].

Korde et al. [139] examined the effect of RISP knockdown on the response to hypoxia in isolated mouse PA, cultured PASM, mitochondria isolated from PA homogenates, and complex III obtained from these using immunocapture. PASM were transfected with the genetically encoded ratiometric H_2O_2 indicator pHyPer-dMito [296] which the authors described as being expressed in the mitochondrial IMS. Hypoxia ($PO_2 = 13$ Torr) markedly increased the pHyPer-dMito signal and enhanced ROS production (assessed using DCFH) by isolated mitochondria and complex III. Hypoxia increased the activity of isolated complex III but had no effect on O_2 consumption or ATP production by the isolated mitochondria. Mitochondrial H_2O_2 under hypoxic conditions was greatly diminished in PASM in which RISP expression was strongly depressed by siRNA treatment, as was hypoxia-induced ROS production in isolated mitochondria and Complex III obtained from these cells. RISP knockdown also strongly depressed HPV and the associated rise in $[Ca^{2+}]_{cyt}$ in isolated PA. Overexpression of RISP had opposite effects. Based on these results, the authors proposed that complex III in PASM is intrinsically sensitive to hypoxia and speculated that this might involve an effect on the transfer of electrons from ubisemiquinone to cytochrome c_1 by RISP.

Based on their observation [375] that hypoxia induces Ca^{2+} release from the sarcoplasmic reticulum in PASM which is dependent on ROS released from the mitochondria, and evidence that Ca^{2+} causes increased ROS production by mitochondria (Section 3.1.1), this laboratory investigated whether hypoxia induces a positive feedback process

involving Ca^{2+} and ROS release in PASMC [376]. They found that the release of Ca^{2+} from the sarcoplasmic reticulum induced by caffeine (0.2 mM to 2M) or norepinephrine (1 μM –10 mM) caused increases in mitochondrial $[\text{Ca}^{2+}]$ and cytoplasmic $[\text{ROS}]$, which was monitored using several ROS probes, including the cytoplasmic H_2O_2 indicator pHyPer-cyto. In addition, raising the $[\text{Ca}^{2+}]$ in a solution bathing mitochondria isolated from these cells also stimulated their ROS production. Hypoxia also increased mitochondrial $[\text{Ca}^{2+}]$ and ROS production by PASMC and isolated mitochondria. These effects were antagonized by RU360, a blocker of the mitochondrial Ca^{2+} uniporter (MCU). RISP knockdown attenuated basal as well as Ca^{2+} - and hypoxia-induced ROS production by PASMC and mitochondria isolated from these cells. These results suggested that hypoxia stimulates complex III-dependent ROS production by increasing mitochondrial $[\text{Ca}^{2+}]$, and that this involves Ca^{2+} release from the sarcoplasmic reticulum and the opening of the MCU. The authors also reported that smooth muscle cells cultured from mesenteric arteries had a much lower expression of RISP compared to PASMC and did not increase their ROS production in response to hypoxia, supporting the concept that RISP-dependent ROS production is a unique property of PASMC.

Evidence for a role of RISP in O_2 sensing was also provided by Waypa et al. [140], who used compartmentally targeted Ro-GFP to compare the effects of hypoxia (30 min: 1.5% O_2) on oxidant signaling in pulmonary and renal artery smooth muscle cells from conditional knockout mice in which RISP expression was either normal or decreased by ~60%. In control PASMC, hypoxia caused a pattern of compartmental redox alterations resembling that which had been seen in rat PASMC: the cytoplasm and IMS became more oxidized and the mitochondrial matrix became more reduced [366]. In contrast, Cyto-Ro-GFP in RISP-depleted cells was more oxidized under normoxic conditions and hypoxia caused the cytoplasm to become more reduced. Hypoxia did not alter the redox state of IMS-Ro-GFP in RISP-depleted cells, and the redox state of the mitochondrial matrix, which was more reduced under normoxic conditions compared to controls, was also unaffected by hypoxia. Echoing their earlier observation in rats [366], they found that the pattern of compartmental changes in Ro-GFP oxidation in renal artery smooth muscle cells induced by hypoxia was similar to that in PASMC, as were the effects of RISP-deletion. In additional in vivo experiments, injection of smooth muscle cell-MHC-Cre/RISP^{flox/flox} mice with tamoxifen was used to knock down smooth muscle RISP expression by ~60%. Measurements of PA diameter and $[\text{Ca}^{2+}]_{\text{cyt}}$ in lung slices revealed that HPV and the associated rise in PASMC $[\text{Ca}^{2+}]_{\text{cyt}}$ were suppressed in tamoxifen- vs. vehicle-treated animals. Moreover, whereas PA wall thickness, RV mass, Fulton index, cardiac function and baseline normoxic PAP were similar in tamoxifen- vs. vehicle-treated animals, the increase in right ventricular systolic pressure caused by ventilating the animals with a hypoxic gas mixture (5% O_2 /95% N_2) was strongly reduced in the tamoxifen group. The authors proposed that RISP ablation acted at site Q_0 to inhibit the formation of ubisemiquinone, which can donate an electron to O_2 to form superoxide. Based on the effects of RISP deletion on the differential compartmental oxidation of Ro-GFP, they suggested that hypoxia might change the configuration of complex III, such that the ROS it produced were directed into the IMS rather than the mitochondrial matrix.

Importantly, however, the depletion of RISP would be predicted to disrupt the Q cycle and the redox state of CoQ bound at Q_0 , which governs the production of ROS by complex III. Because the flow of electrons through RISP is necessary for the reduction of cytochrome *c*, its removal has also been shown to strongly inhibit respiration [377–379], and can cause mitochondrial depolarization [380,381], the upregulation of glycolysis, an increase in the NADH/NAD ratio [377,379], and the decreased expression of multiple isoforms of NADPH oxidase [378]. Although there is also countervailing evidence that cells can maintain a normal mitochondrial membrane potential and rate of glycolysis

following RISP knockdown [204], its potential for causing disruption of mitochondrial function is such that the results of these studies should be interpreted with caution.

5.2.2. Mitochondrial Hyperpolarization, Reduction of the Quinone Pool, and Cox4i2

Several studies by Sommer and colleagues [141,142,155] presented evidence that hyperpolarization of $\Delta\Psi_m$ and reduction of components of the ETC promote hypoxia-induced mitochondrial ROS production in PASM. C.

Sommer et al. [155] evaluated the effect of hypoxia on the redox state of cytochromes $b_{L/H}$, c , and $aa3$ in isolated perfused lung and PASM. C. from rabbit using remission spectrophotometry, a technique in which the redox state of multiple cytochromes is inferred from the wavelength-dependency of light absorbance by cells. They found that HPV was evoked when the pO_2 was lowered to ≤ 75 Torr and increased progressively as the level of hypoxia was deepened. A significant change in absorbance was observed at $pO_2 \leq 53$ Torr, and analysis of the spectra indicated that reduction of cytochromes c , $aa3$, and $b_{L/H}$ occurred at pO_2 values of 38, 23, and 8 Torr respectively in the lungs. Based on measurements of the P_{50} for oxygen consumption, they calculated (see [160]) that respiration was significantly inhibited at ambient PO_2 levels of <100 Torr (e.g., by ~ 2 and $\sim 10\%$ at pO_2 of 38 and 8 Torr, respectively). Experiments in primary cultured microvascular PASM. C. showed that both $[Ca^{2+}]_{cyt}$ and $\Delta\Psi_m$ were increased at a pO_2 of 23 and 8 Torr. Mitochondrial ROS production (monitored using MitoSox) was also increased at a pO_2 of 8 Torr. Application of either the complex IV blocker cyanide or the complex III/ Q_i site blocker HQNQ to isolated lungs caused a reduction of all of the cytochromes and an inhibition of HPV, with both effects developing over the same concentration ranges of each blocker.

The authors concluded that HPV was due to an increase in mitochondrial ROS production by complex III, which was linked to an increased reduction of components of the ETC associated with a subtle but significant inhibition of respiration evoked by hypoxia. They highlighted their observation that cytochrome $b_{L/H}$ remained more oxidized than the other cytochromes, speculating that this might reflect its involvement in ROS production, as suggested previously in a report showing that superoxide production was associated with a similar pattern of hypoxia-induced cytochrome reduction in a monocyte cell line [162].

Cox4 is one of the 10 regulatory subunits of complex IV encoded by nuclear DNA. When phosphorylated, Cox4 functions to inhibit complex IV at a high ATP/ADP ratio [382]. Cox4 exists as two isoforms, Cox4i1 and Cox4i2. The expression of Cox4i1 strongly predominates in most tissues, but Cox4i2 is also highly expressed in lung [186] and CBCC [383,384]. Cox4i2 has been shown to promote respiration [351] and therefore potentially increase $\Delta\Psi_m$, a key determinant of mitochondrial ROS production.

Sommer et al. [142] therefore explored the relationship between $\Delta\Psi_m$, ROS production, and downstream mechanisms causing HPV in Cox4i2 knockout and wildtype mice. They found that hypoxia (1% O_2 for 5 or 10 min) caused a hyperpolarization of $\Delta\Psi_m$ in primary cultured PASM. C. from wildtype mice. This was associated with increased mitochondrial superoxide production, which was measured using electron spin resonance spectroscopy (ESR), as well as a rise in cellular $[H_2O_2]$, the inhibition of plasmalemmal K_v channels, membrane depolarization, and Ca^{2+} influx. All of these effects of hypoxia were greatly diminished in PASM. C. from Cox4i2^{-/-} mice. Hypoxia-induced membrane depolarization was also inhibited in human PASM. C. treated with anti-Cox4i2 siRNA.

The results presented in this paper are too extensive to describe in detail, but are summarized in Figure 8. In essence, they showed that both phases of HPV, which this laboratory had first observed in isolated perfused lung from wildtype mice [357], were virtually absent in Cox4i2^{-/-} mice. Notably, however, since their studies in PASM. C. utilized

only brief exposures to hypoxia, the specific effector mechanisms shown in Figure 8 are not necessarily involved in Phase 2 HPV, although it too was Cox4i2-dependent.

In accordance with their model, the authors found that hypoxia-induced superoxide production was restored by overexpressing wildtype Cox4i2 in PASM from Cox4i2^{-/-} mice. However, this was not the case when Cox4i2, in which cysteine 109 had been replaced by a serine or alanine, was overexpressed in these cells. Additional experiments suggested that cysteines 41 and 45 on Cox4i2 were also required for superoxide production. These results suggested that these cysteines might act as thiol switches, which would allow Cox4i2 to be redox-regulated during hypoxia in such a way as to increase $\Delta\Psi_m$. However, their original hypothesis that Cox4i2 might be increasing $\Delta\Psi_m$ by enhancing respiration during hypoxia was not borne out by their experiments, and they also found that mitoTEMPO did not prevent hypoxia from increasing $\Delta\Psi_m$, which was not consistent with redox regulation of Cox4i2 being responsible for this effect. Thus, whereas the study presents evidence that the hypoxia-induced increase in $\Delta\Psi_m$ in PASM was dependent on the presence of Cox4i2, it did not show that a specific effect of hypoxia on Cox4i2 was causing the rise in $\Delta\Psi_m$. It is particularly noteworthy that HPV, recorded as an increase in PAP in isolated perfused mouse lung, was inhibited in a concentration-dependent manner by S3QEL2, which blocks ROS production by Complex III without affecting respiration.




Figure 8. A summary of the experimental evidence reported in Sommer et al. [142]. Black text in the flow chart on the left shows results related to the characterization of HPV mechanisms, and text in red shows the sequence of events proposed to cause HPV. The evidence supporting each step in the sequence is shown to the right in black font, and the red font shows the results of negative control experiments.

A subsequent paper from this group [Pak, 2018 #29721] again employed ESR to show that hypoxia (1% O₂) increased superoxide levels in mouse PASM_C, and also recorded a rapid increase in cytoplasmic H₂O₂ using Hyper_{cyto}. In contrast, hypoxia strongly decreased superoxide levels in whole lung homogenates. A similar effect occurred in cultured lung fibroblasts, leading them to speculate that this might explain the hypoxia-induced fall in ROS in whole lung homogenate, which had also been seen previously in perfused lungs [287,385]. Using MitoSox, they confirmed their earlier finding [155] that hypoxia increased ROS in the mitochondrial matrix, a finding which, however, runs counter to measurements made using genetically encoded ROS indicators [137,366]. The authors did not discuss this discrepancy, which could conceivably reflect the drawbacks of MitoSox [386].

To further investigate mitochondrial ROS production by the distal ETC, this laboratory [141] utilized AOX mice, which express an alternative oxidase derived from the marine invertebrate *Ciona intestinalis*. AOX becomes active if the flux of electrons through the distal ETC is slowed enough to cause substantial reduction of the quinone pool [387], creating an alternative pathway for electrons to flow from QH₂ to O₂ which bypasses complexes 3 and 4. This prevents over-reduction of the quinone pool while maintaining oxidative phosphorylation, with no apparent deleterious effects on mice [388]. As initially described in tobacco cells, AOX suppresses basal ROS production [389].

They observed that the pressor response to cyanide and both phases of HPV evoked by hypoxia (1% O₂ in the ventilating gas mixture) in isolated perfused lungs from WT mice were almost abolished in AOX expressing animals, but were restored by applying the AOX inhibitor *n*-propyl gallate. Hypoxia-induced PASM_C depolarization was decreased by >50% in AOX compared to wildtype mice, and *n*-propyl gallate caused further depolarization in the PASM_C from AOX mice, but not from controls. HPV in isolated PA recorded in control PSS was depressed in the AOX mice, but if arteries were incubated in PSS containing 20 mM K⁺ to evoke a degree of resting membrane depolarization, HPV in the WT and AOX mice was of similar amplitude. Perfusion of isolated lungs from AOX with high K⁺ PSS also produced some restoration of HPV, although it remained far smaller than that seen in wildtype mouse lung perfused with this solution. They also used ESR to show that the hypoxia-induced increase in superoxide seen in PASM_C from wildtype mice was absent in those from AOX mice. Similarly, hypoxia hyperpolarized ΔΨ_m in PASM_C from WT but not AOX mice.

In further experiments, they transfected rat PASM_C with either an AOX-containing or empty expression vector and saw that the presence of AOX slightly enhanced respiration at very low PO₂. This observation that AOX increased O₂ consumption under hypoxic conditions supported the notion that hypoxia was causing reduction of the quinone pool, since AOX should not be able to donate electrons to O₂ unless this pool is substantially reduced [387]. In accordance with this possibility, and with their earlier observations [155], they found using Raman spectroscopy that hypoxia reduced the quinone pool and cytochrome *c*, but not cytochrome *b*, in WT PASM_C. The quinone pool was also reduced in the AOX PASM_C, but the redox state of cytochromes *b* and *c* was unchanged, supporting the idea that electron flow through AOX was bypassing complexes 3 and 4.

These investigations by Sommer and colleagues suggest a model in which hypoxia increases ROS production by complex III, possibly via the escape of electrons from

cytochrome $b_{H/L}$, and that increased reduction of the quinone pool due to a slight inhibition of respiration, as well as mitochondrial hyperpolarization, which had previously been observed by other laboratories [26,156] contribute to this response.

It is noteworthy that Cox4i2A has also been implicated in O_2 sensing in the carotid body. Moreno-Dominguez et al [390], reported that the normal hypoxic ventilatory response, which is dependent on O_2 sensing by CBCC, was virtually abolished in Cox4i2 null mice, as was the rise in $[Ca^{2+}]_{cyt}$ in CBCC which underlies the secretory response of these cells to a fall in PO_2 . The rise in NADH evoked by hypoxia in CBCC from wildtype controls was also absent in the knockouts. Similar effects were seen when the expression of Cox4i2 was reduced by knocking out *EPas1*, the gene coding for Hif1a. *EPas1* knockout also diminished the expression of two other mitochondrial proteins, COX8b, another regulatory subunit of complex IV, and NDUFA4L2, which interacts with complex IV [382]. These two proteins, together with Cox4i2, have been proposed to contribute to the unusually high $K_m[O_2]$ of CBCC that increases their sensitivity to hypoxia [383,384], although how they would do so is unclear.

The observation that hypoxia had no effect on [NADH] in CBCC from Cox4i2 knockouts, which suggested that it was not inhibiting respiration, is consistent with evidence from a study by Pajuelo-Riguera and colleagues which compared the PO_2 sensitivity of Cox in HEK293 cells exclusively expressing either Cox4i1 or Cox4i2 [185]. Their key finding was that the $K_m[O_2]$ for mitochondrial respiration was almost ~2-fold higher in the cells expressing Cox4i2, indicating that respiration would be more depressed in these cells at low PO_2 levels. The authors pointed out that this should promote reduction of the proximal ETC and of the $NAD^+/NADH$ couple, both factors known to increase mitochondrial ROS production, and was consistent with the loss of HPV seen in PA from Cox4i2 knockout mice.

Notably, however, the effect of Cox4i2 on the $K_m[O_2]$ described in this paper appears to be insufficient to itself cause a meaningful depression of respiration except during severe hypoxia (see Equation (1) in [160]), implying that its expression in O_2 -sensing cells might be necessary but in and of itself insufficient to promote mitochondrial ROS production at moderate levels of hypoxia. The authors suggested that the expression of COX8b and NDUFA4L2 might also be important for the hypoxia-induced ROS production, possibly through their interaction with Cox4i2. It would therefore be of interest to see if these proteins are expressed in PASM, and if the combined knockout of all three proteins affects HPV at moderate levels of hypoxia.

5.2.3. HPV and Mitochondrial NCLX

As described in Section 3.1.1, hypoxia is proposed by Hernansanz-Agustin and colleagues [194,195] to cause complex I deactivation, leading to acidification of the mitochondrial matrix. Since mitochondria contain a very substantial amount of Ca^{2+} bound to phosphate as a $Ca_3(PO_4)_2$ complex, the dissociation of which is highly pH-sensitive [201], acidification causes an increase in free $[Ca^{2+}]_{mito}$. According to their model, this promotes Na^+ entry into the matrix through the mitochondrial NCLX. The resulting rise in $[Na^+]_{mito}$ decreases the fluidity of the IMM, impeding electron transfer from complex II and G3PD to complex III and thereby increasing ROS production at Complex III_{CoQ} by slowing electron flow within the Q cycle.

Although this scheme was based on data obtained from bovine aortic endothelial cells and mouse embryonic fibroblasts, they presented evidence that this mechanism is involved in O_2 sensing during HPV by showing that hypoxia (1% O_2) strongly increased superoxide production by mouse PASM, and that this response was absent in cells in which the mitochondrial NCLX was knocked down using siRNA [194]. Furthermore, the

NCLX inhibitor CGP-37157 (30 μ M) suppressed sustained HPV in isolated rat PA, although the initial peak of the response was unaffected.

However, CGP-37157 at this concentration also blocks L-type voltage-gated Ca^{2+} channels and K_v channels [391,392], both of which contribute importantly to HPV. Also arguing against a role for NCLX in HPV, Becker and colleagues found that replacement of extracellular Na^+ with Li^+ , which should profoundly reduce intracellular $[\text{Na}^+]$ [393] and thereby prevent Na^+ influx into the mitochondria through NCLX, did not inhibit HPV in rat pulmonary arteries. Similarly, application of the $\text{Na}_v\text{K-ATPase}$ inhibitor ouabain, which should greatly increase the $[\text{Na}^+]$ in the cytoplasm and therefore in the mitochondrial matrix [394] did not inhibit HPV. Thus, HPV persisted in spite of significant perturbations of the intracellular Na^+ concentration which would be predicted to disrupt the Na^+ dependent regulation of the fluidity of the mitochondrial inner membrane proposed to be responsible for the hypoxia—induced rise in mitochondrial ROS production. Nevertheless, the Becker et al. study was not designed to evaluate the role of NCLX in HPV, and the evidence presented by Hernansanz-Agustin et al. that their scheme is involved in O_2 sensing is extensive; thus, a more thorough investigation focusing on the involvement of this pathway in HPV would be desirable.

5.2.4. Hypoxia and Increased ROS Production by Complex II

Paddenberg et al. [138] used DCFH-DA to characterize the effect of hypoxia (1% O_2) on ROS levels in PA from mouse lung slices and cultured rabbit PASM. They observed that three hours of hypoxia increased the number of PASM and pulmonary artery endothelial cells in PA from the lung slices which exhibited an increase in the DCFH signal above the threshold level they used to indicate ROS production (DCFH-positive cells), and that one hour of hypoxia also significantly increased DCFH fluorescence in rabbit PASM. The hypoxia-induced increase in DCFH-positive cells was abolished by DPI, rotenone, and the complex II blockers 3-NPA and TTFA. Rotenone decreased the number of DCFH-positive cells under normoxic conditions, as did antimycin, but 3-NPA did not, leading them to propose that hypoxia was acting at complex II to increase ROS production, and also that complex III was contributing to ROS production during both normoxia and hypoxia. Using a histochemical assay, they showed that hypoxia decreased the activity of succinate dehydrogenase in the lung slices. Furthermore, application of exogenous succinate prevented the hypoxia-induced increase in ROS. Based on these results they proposed that hypoxia causes complex II, which normally accepts electrons from succinate to generate fumarate, to switch to using electrons it receives from NADH via complex I and the ubiquinol pool to reduce fumarate to succinate, as a result generating ROS.

5.2.5. Evidence for NAD(P)H Oxidase as a Source of Increased ROS During HPV

In 1996, Marshall et al. [344] presented evidence that hypoxia increases Nox-mediated superoxide production in PAs. They employed immunohistochemistry and spectrophotometry, respectively, to detect the presence of gp91-*phox* and the unique Nox cytochrome (*b-245*) in cultured PASM from fetal calves, and immunoblotting to demonstrate its expression in both PA and systemic arteries. Using lucigenin as a nominal superoxide indicator, they examined the effect of hypoxia (~40 Torr) on ROS production by primary cultured smooth muscle cells from small and large PA, and also aorta and ear arteries. Hypoxia greatly increased superoxide production in each type of smooth muscle cell, particularly those from small PA. This effect was absent if extracellular SOD was applied, leading the authors to conclude that lucigenin was detecting an increase in extracellular superoxide. Hypoxia-induced superoxide production was virtually abolished by the Nox antagonist DPI. Although DPI has many other effects, including block of complex I [395]), superoxide production was unaffected by myxothiazol, suggesting it was not of

mitochondrial origin. HPV in isolated PA from cats was blocked dose-dependently by DPI, with an ID_{50} of $\sim 0.81 \mu\text{M}$, whereas the pressor response to norepinephrine was ~ 15 -fold less sensitive to DPI. The authors concluded that hypoxia was causing an activation of Nox, rather than the inhibition which had been described previously [396] (see Section 8.1), and suggested that the K_m of the oxidase for O_2 ($10 \mu\text{M}$ or ~ 7 Torr) was low enough to ensure that superoxide production would not be limited by the fall in PO_2 associated with moderate hypoxia.

The role of Nox in HPV was subsequently challenged by the observation that HPV was intact in lungs from mice with chronic granulomatous disease, which lack the phagocytic form of gp91-phox (Nox2) [196]. Lungs from these mice also exhibited a profound suppression of normoxic ROS production (detected using lucigenin), and in contrast to the observations of Marshall et al. [344], ROS production in isolated PA from control mice was strongly depressed by moderate (~ 40 Torr) hypoxia.

However, Weissmann et al. [355] thereafter reported the expression of Mox-1, a different isoform of Nox (now called Nox1), in PASMC from rabbit lung. In order to study its role in HPV while avoiding problems associated with the non-selectivity of the Nox blockers DPI and apocynin [397], they used AEBSF, a serine protease inhibitor which had been shown to block Nox in phagocytes [398] and to inhibit the response to hypoxia in the carotid body [399]. AEBSF was also a flawed blocker, as it caused a substantial increase in PAP in isolated perfused rabbit lung. However, this effect was transient, and when the PAP had returned to baseline level, the pressor response to 10 min of hypoxia (O_2 of ~ 23 Torr in the ventilating gas mixture) was depressed by AEBSF in a concentration-dependent manner. The authors also demonstrated that HPV, but not the pressor response to U46619, was attenuated by the SOD inhibitor TETA, which itself did not affect PAP. These results led them to conclude that ROS production by Nox was activated by hypoxia and contributed to HPV.

Weissmann et al. [356] reported that HPV in isolated perfused rabbit lung is biphasic, observing that PAP rose to reach a peak ~ 5 min after the imposition of hypoxia (Phase 1), then fell by $\sim 50\%$ to reach a nadir at ~ 15 min, and thereafter rose gradually to reach a plateau after 2–3 h (Phase 2). The antioxidant nitro blue tetrazolium (NBT), applied at an initial concentration of $0.9 \mu\text{M}$, strongly depressed both phases, as did TETA, applied at an initial concentration of 25 mM . DPI (initial concentration $1.5 \mu\text{M}$), applied in the presence of L-NMMA, diminished Phase 1 by $\sim 50\%$, whereas Phase 2 was almost abolished. DPI itself evoked a minor increase in baseline PAP, whereas TETA and NBT caused a small fall. However, Weissmann et al. 2006 [357] observed that AEBSF (initial concentration $500 \mu\text{M}$) abolished Phase 1 HPV in the rabbit lung, without significantly affecting Phase 2. Like Archer and colleagues [222], they found that Phase 1 was of similar amplitude in gp91-phox knockout mice and wildtype controls. However, Phase 1 was significantly depressed in p47-phox knockouts, in which ROS production by both Nox1 and Nox2 would be depressed. In contrast, the pressor response to U46619 was similar in all three groups of mice. Baseline normoxic PAP was also not different in the p47-phox knockouts, suggesting that basal ROS production by NOX1 or 2 was not contributing to ongoing normoxic vasodilation. These data suggested that ROS production by Nox1 was partly responsible for Phase 1 HPV.

In 2008, Rathore et al. [24] found that protein for Nox1, Nox4, and the p22-, p47-, and p67-phox components of Nox was similarly expressed in small endothelium-denuded pulmonary and mesenteric arteries from mice, but saw no expression of Nox2 in either artery. Normoxic arteries or those subjected to 5 min of hypoxia ($1\% O_2$) were homogenized and centrifuged to isolate a plasma membrane-containing fraction and the activity of Nox was monitored by adding NADPH and measuring the SOD-inhibitable (i.e., superoxide-dependent) reduction of cytochrome C. Nox activity was increased in pulmonary but not

mesenteric arteries, presumably reflecting an effect caused by hypoxia which had persisted during the isolation process. This was associated with an increased plasmalemmal expression of p47-phox, consistent with Nox activation.

Nox activity during hypoxia was strongly suppressed by the nominal Nox blocker apocynin (1 μM) and was smaller in PA from p47-phox knockouts compared to wildtype controls. Similar effects were also observed in PA from PKC ϵ knockout mice compared to wildtype controls and in PA treated with a peptide inhibitor of PKC ϵ translocation (10 μM) but not Gö6796 (0.1 μM), a blocker of conventional PKC isoforms. Although it is not clear that these interventions diminished the effect of hypoxia per se, since their effects on Nox activity under normoxic conditions were not described, parallel experiments utilizing DCFH as an indicator showed that in each case hypoxia-induced ROS production was attenuated in freshly isolated PASMC with no effect apparent during normoxia. Cell shortening and rises in $[\text{Ca}^{2+}]_{\text{cyt}}$ in response to hypoxia were also decreased in PASMC treated with apocynin and in those from p47-phox $^{-/-}$ mice, whereas $[\text{Ca}^{2+}]_{\text{cyt}}$ and the contractile response to caffeine were not affected under normoxic conditions.

The authors also observed that Nox activity during hypoxia was strongly suppressed by rotenone and myxothiazol. They had previously found that hypoxia increased the activity of PKC ϵ in PASMC, that this effect was also sensitive to these ETC blockers, and that the expression of PKC ϵ in PA was much higher than in MA [400]. They therefore proposed that increased mitochondrial ROS production in PASMC due to hypoxia was activating PKC ϵ , causing a stimulation of Nox1 that generated additional superoxide which acted via effector pathways to cause HPV. They suggested that the Nox2 expression which had previously been reported in mouse lung [222] might occur mainly in non-smooth muscle cells or in pulmonary veins. They also pointed out that the lack of effect of blocking or knocking out p47-phox on basal $[\text{Ca}^{2+}]_{\text{cyt}}$ or the caffeine-induced contraction went against the concept, introduced by Michael Wolin's laboratory (see Section 8) that ongoing Nox-mediated ROS production maintains a tonic normoxic vasodilation which can be reversed by hypoxia.

In 2011, Frazziano et al. [328] similarly detected a role for both mitochondrial and Nox-derived ROS production in HPV evoked in small rat PA by 10 min of hypoxia (22–26 Torr). This group had earlier proposed that HPV is due to the activation of neutral sphingomyelinase 2 (nSMase-2), this causing an increased production of ceramide and a consequent stimulation of PKC ζ which acts to depress the K_v current [401,402]. They observed that the increase in cellular ceramide caused by hypoxia was abolished by rotenone, suggesting its dependency on mitochondrial ROS production. Application of exogenous ceramide caused the production of ROS (monitored using DCFH and DHE), and this was blocked by a PKC ζ —inhibitory peptide. Hypoxia also caused the phosphorylation of p47-phox and its association with caveolin-1, and this was blocked by the PKC ζ —inhibitory peptide [328]. They also found that 300 μM apocynin strongly antagonized HPV. Based on these results, they proposed that nSMase-2 is stimulated by mitochondrial ROS, generating ceramide which activates Nox via a PKC ζ —mediated phosphorylation of p47phox, leading to the production of ROS which inhibit K_v channels to cause HPV. The evidence that hypoxia induced a PKC ζ -dependent phosphorylation of p47^{phox}, which was blocked by rotenone, supports the model developed by Rathore and colleagues, in that mitochondrial ROS would act through PKC to promote further oxidant signaling by Nox, although the two schemes differ with regard to which PKC isoform causes p47phox activation.

It is notable, however, that the effects of apocynin described in these papers are uninterpretable, since Heumüller et al. [350] showed that it blocks Nox only after being converted by myeloperoxidase into a dimer and that this does not occur in vascular smooth

muscle cells, which lack this enzyme. Apocynin was also found to act as an antioxidant which interacts with both biomolecules and with ROS indicators such as lucigenin.

In contrast to the findings of these papers, Connolly et al. (2013) [18], who studied HPV in rat intrapulmonary arteries in the absence of pre-tone (under which condition Phase 1 HPV is small), found that the Nox inhibitor VAS2870 (10 μ M) had no effect on HPV, although the broad-spectrum PKC antagonist Gö6983 (3 μ M), which would be expected to block both PKC ϵ and PKC ζ [403], reduced HPV by ~30%.

Veith and colleagues [404] compared the effect of hypoxia (1% O₂) in perfused lungs from wildtype and Nox4^{-/-} mice, finding that neither phase of HPV was different in these two groups. The basal right ventricular systolic pressure (RSVP), an indicator of the PAP, was also not significantly affected by Nox4 knockout. Murtaza et al. [405] subsequently compared the effect of hypoxia on pre-acinar and smaller intra-acinar arterioles in lung slices from these two groups, and saw that wildtype intra-acinar arteries demonstrated a substantial sustained constriction which was halved in the Nox4 knockouts, whereas the pre-acinar arteries showed a smaller response which was similar in the two groups. In view of the results from Veith et al., they suggested that constriction of intra-acinar arteries does not contribute to the overall hypoxia-induced rise in PAP [406] but might be involved in the redistribution of blood flow responsible for ventilation–perfusion matching.

Nagaraj et al. [407] recorded HPV (evoked by 1% O₂) in perfused lungs from wildtype and p22^{phox}^{-/-} mice, reporting that whereas the basal normoxic PAP and initial transient phase of HPV were similar between the two groups, the sustained phase was decreased by ~40%. They also showed that hypoxia increased the activity of rho kinase in lung slices from wildtype mice, and that this effect was absent in the p22^{phox} knockouts. Since Noxs 1 and 4 are the isoforms most likely to be involved in HPV, and both require p22^{phox} for their activity, these results suggest that ROS production by Nox1 may contribute to sustained HPV by stimulating Rho kinase-mediated Ca²⁺ sensitization.

The Role of NADPH Oxidase in HPV: Summary and Conclusions

Although Nox2 is not involved in HPV [222], at least in mice, the studies in rodents described above generally support the possibility that HPV depends to some extent on a hypoxia-induced activation of ROS production by Nox1 which is dependent on PKC, a well-characterized activator of Nox1/2 [69]. The possibility that activation of Nox1 by 5 min of hypoxia is dependent on mitochondrial ROS production acting via PKC ϵ , as evidenced by Rathore and colleagues [24,400], is also consistent with observations that the amplitude of Phase 1 HPV in mice was decreased by ~1/3 when p47-*phox* was knocked out (Weissmann et al., 2006) [357], but was abolished in Cox4i2^{-/-} mice, along with superoxide production [142]. The involvement of a mitochondrial ROS-PKC ϵ -Nox1 axis in the initial phase of HPV is further supported by evidence that the pressor response to a 5 min hypoxic challenge in isolated perfused rat lung was decreased by ~50% [408] in PKC ϵ knockout mice, and that the rapid increase in [Ca²⁺]_{cyt} evoked by hypoxia was strongly depressed in PASMC from these knockouts and from wildtype mice treated with PKC ϵ inhibitor peptide [400]. Moreover, the finding that Ro-31-8220, which blocks multiple PKC isoforms including PKC ϵ , depressed Phase 1 HPV in rat intrapulmonary arteries but had no effect on Phase 2 [409] supports the lack of involvement of this pathway in sustained HPV. The investigation by Nagaraj et al. [407] also implicates Nox1 in HPV. However, they observed that it was Phase 2 rather than Phase 1 HPV which was Nox-dependent. The reason for this discrepancy is unknown.

Importantly, with the exception of the initial paper by Marshall et al. [344], the investigations described above ignored the question of whether the K_m(O₂) for Nox has an impact on ROS production during hypoxia. The K_m(O₂) for Nox1 has not been reported, but if it is similar to that of Nox2 (~25 Torr) [68], its ability to generate ROS would be

diminished at the levels of hypoxia used by most laboratories to study its involvement in HPV. This would call into question the plausibility of the findings that ROS production by Nox1 was increased at this level of hypoxia, but at the same time would imply that if these observations are valid, these studies may have greatly underestimated the ability of Nox1 to produce ROS and therefore contribute to HPV at more moderate levels of hypoxia.

This issue is even more salient with regard to the $K_m(O_2)$ for Nox4, which has been shown to be ~136 Torr [68,410]. This implies that Nox 4 is very unlikely to increase its production of H_2O_2 in response to acute hypoxia, but is instead well suited to act as an O_2 sensor which responds to acute hypoxia by having the opposite effect [68,410]. Indeed, as described in Section 8, it has been suggested that a fall in H_2O_2 production by Nox4 is instrumental in causing HPV in bovine pulmonary arteries.

5.2.6. HPV Effector Mechanisms Coupled to an Increase in PASMC [ROS]

The Redox Theory from the outset incorporated K_v channel inhibition as the essential HPV effector mechanism. In contrast, Schumacker's laboratory did not include any specific effector mechanisms in their initial presentation of the Mitochondrial ROS hypothesis [23], although they later suggested [353] that increases in cytoplasmic ROS might be working through activation of phospholipase C [411], which would cause an increase in IP_3 production leading to Ca^{2+} release from the sarcoplasmic reticulum mediated by the IP_3R (see also [412]). Subsequently, it has been demonstrated that Ca^{2+} release from the sarcoplasmic reticulum mediated by the ryanodine receptor (RyR) [413] and SOCE mediated by STIM1/TRPC1/Orai1 [414] are both activated in a ROS-dependent manner during HPV. H_2O_2 was also shown to cause an increase in $[Ca^{2+}]_{cyt}$ in PASMC mediated by Ca^{2+} release from the sarcoplasmic reticulum through IP_3R and RyR, as well as Ca^{2+} influx through Ni^{2+} -sensitive cation channels [415].

Other pro-contractile mechanisms mobilized by acute hypoxia in PASMC, including Ca^{2+} influx via TRPC6 channels [11,416,417] and the activation of several protein kinases [418], such as PKC [41,328], Src-family kinases [419–421] and rho kinase [421], have also been shown to be stimulated by ROS. It has additionally been demonstrated that the IP_3R , which may contribute to Ca^{2+} release from the sarcoplasmic reticulum during HPV [3] is activated by ROS released from the mitochondria via oxidation of a group of cysteines located in its N-terminal suppressor domain [103,268,269]. Furthermore, as described in Section 5.2.5, Cogolludo and colleagues have provided evidence that an increase in ROS production by the mitochondria and Nox inhibits the opening of $K_v1.5$ channels in PASMCs through a mechanism involving nSMase-2, ceramide, and PKC ζ [328,401].

6. Other Aspects of Redox Regulation of HPV

6.1. H_2O_2 vs. Superoxide (vs. Peroxynitrite?) as Mediators in HPV

Although most papers have made no distinction between superoxide and H_2O_2 as possible HPV mediators, it is increasingly accepted that H_2O_2 , acting directly or, more likely, via intermediates (e.g., Prxs and Grxs), is primarily responsible for the oxidative post-translational modifications to protein thiols which underpin cellular ROS-induced signaling. In contrast, the involvement of superoxide in ROS-induced signal transduction remains enigmatic [48,111,422]. Nonetheless, hypoxia was found to increase superoxide production by PASMC [142,423], and HPV has been shown to be inhibited by SOD and enhanced by SOD antagonists [346,358]. There is also a great deal of evidence that superoxide production is increased and that SOD is downregulated in animal models of pulmonary hypertension, leading to an increased ratio of superoxide to H_2O_2 concentrations in PASMC, which contributes to the pulmonary vasoconstriction and remodeling which

characterizes this condition [424]. In addition, Knock and colleagues [420] found that LY83583, which generates superoxide intracellularly, caused a contraction of PA, which was abolished by SOD but not catalase. They subsequently showed [421] that LY83583 caused thiol oxidation and autophosphorylation of Src-TK, resulting in its activation. Src-TK then activated the guanine exchange factor ARHGEF1, leading to a stimulation of rhoA and rho kinase, Ca^{2+} sensitization, and consequent PA contraction. Although application of 30 μM H_2O_2 also caused autophosphorylation of Src-TK, the authors argued that the effect of LY83583 on rhoA/rho kinase was superoxide-dependent, since H_2O_2 had previously been shown not to activate rhoA/rho kinase in these cells [420]. Furthermore, LY83583 caused a marked (and SOD-inhibitable) enhancement of the contraction to $\text{PGF}_{2\alpha}$, whereas application of H_2O_2 relaxed $\text{PGF}_{2\alpha}$ -constricted PA [425]. Hypoxia (1% O_2) also stimulated Src-TK autophosphorylation and Ca^{2+} sensitization, and both effects were abolished by the application of either rotenone or myxothiazol (both 100 nM), suggesting that Ca^{2+} sensitization due to superoxide production by complex III could play a role in HPV [421]. Along similar lines, the contraction of rat small IPA to LY83583 but not H_2O_2 was blocked by the phosphatidylcholine-specific phospholipase C (PC-PLC) antagonist D609, which also strongly suppressed HPV, again implying that superoxide can contract PA via H_2O_2 -independent mechanisms and that this occurs during HPV [412].

One mechanism by which superoxide might cause contraction is by promoting the synthesis of peroxynitrite [426]. Various observations support the possibility that an increase in peroxynitrate, which is formed by the very rapid reaction of superoxide and NO, could play a role in O_2 sensing in PASM. This could, for example, arise from a hypoxia-induced increase in mitochondrial or cytoplasmic NO production [427], as described in Section 3.1.1. Reactive nitrogen species such as peroxynitrite can induce both S-nitrosylation and S-glutathionylation of cysteine residues [428,429]. There is evidence that Src-TK is activated by S-nitrosylation [421] and other peroxynitrate-induced mechanisms [430], that IP_3R can be stimulated by S-glutathionylation [431], and that the activity of the RyR_2 can be increased by peroxynitrite [432]. Peroxynitrate has also been shown to deplete the SR of Ca^{2+} in cardiac myocytes, an effect which could potentially lead to the activation of store operated Ca^{2+} entry [433] in PASM if it also occurs in these cells. Superoxide, possibly acting via peroxynitrite, may also contribute to the contraction of PA by 5-HT [426]. In addition, peroxynitrite could potentially act on target proteins via peroxiredoxins; for example, it has been shown to react with Prx2 with a rate constant similar to that of H_2O_2 [146,433].

6.2. Do ROS Cause AMPK Activation During HPV?

The heterotrimeric protein AMP-activated protein kinase (AMPK) is an intracellular energy sensor which, when stimulated by a decrease in the cell energy state, orchestrates compensatory adjustments to metabolic pathways so as to maintain ATP levels. The activity of the catalytic α subunit requires its phosphorylation at Thr-172, which is mediated by the following kinases: liver kinase B1 (LKB1) and Ca^{2+} /calmodulin-dependent protein kinase kinase 2 (CAMKK2). α -subunit activity is also regulated allosterically by the competitive binding of ATP and AMP to two sites on the γ -subunit, such that it is stimulated by increases in the AMP/ATP ratio.

Evans' laboratory [434,435] has proposed that hypoxia elicits a small but significant increase in the AMP/ATP ratio in PASM by causing a subtle inhibition of mitochondrial respiration. This stimulates AMPK, leading to the phosphorylation and engagement of HPV effectors such as K_v channels [26,27]. However, Chandel's laboratory found in mouse embryonic fibroblasts that hypoxia did not alter the AMP/ATP ratio and that hypoxic activation of AMPK was prevented by the superoxide/ H_2O_2 scavenger EUK-134, leading them to suggest that hypoxic activation is mediated by complex 3-derived H_2O_2 through

an AMP/ATP-independent pathway [436]. They later reported [437] that hypoxia-induced activation of AMPK in alveolar epithelial cells depends on a rise in $[Ca^{2+}]_{\text{cyt}}$ which is mediated by mitochondrial ROS acting via the opening of I_{CRAC} channels and a consequent stimulation of CAMKK2. Notably, this conclusion was based largely on experiments which employed Reiske-Fe-S protein knockdown to prevent ROS production by complex 3, an approach which may have been problematic since it could conceivably have caused an inhibition of respiration, thereby affecting the regulation of AMPK by altering the basal AMPK/ATP ratio.

Hawley et al. [438] argued that compartmentalization of AMP and ATP means that their overall cell contents may not accurately reflect the AMP/ATP ratio to which AMPK is exposed. Using HEK293 cells expressing either wildtype γ -subunits or γ -subunit mutants unable to bind ATP/AMP, they showed that the stimulation of AMPK by H_2O_2 required that these binding sites were functional, implying that peroxide was not affecting kinase activity through any other mechanisms. They also found that H_2O_2 depressed mitochondrial respiration and increased the ADP/ATP ratio (indicative of a rise in the AMP/ATP ratio), and concluded that H_2O_2 stimulates AMPK indirectly by increasing the AMP/ATP ratio. Notably, although the $[H_2O_2]$ used in these experiments (1 mM) was very high, application of a more physiologically relevant $[H_2O_2]$ (30 μM) had been shown to raise the AMP concentration in NIH-3T3 cells [439]. Evan's group subsequently demonstrated that HPV in mice was not significantly diminished in CAMKK2 knockout mice [26]. Thus, the available evidence suggests that if ROS are indeed contributing to AMPK activation during HPV, this is likely to be because they are increasing the AMP/ATP ratio.

6.3. H_2S , ROS, and HPV

Olson and colleagues proposed that hypoxia, by inhibiting the O_2 -dependent metabolism of H_2S , causes a rise in its cellular concentration which constitutes the PO_2 -sensitive signal triggering HPV. This was based on observations that 1: applying exogenous sulfide caused constricting or dilating effects mimicking those of hypoxia in multiple blood vessels from diverse species; 2: blockers of sulfide-producing enzymes inhibit HPV; and 3: sulfide production by lung homogenates and pulmonary vascular preparations is increased at low PO_2 [29]. However, the contention that sulfide is an O_2 sensor in HPV has been challenged [440,441]. Subsequently, Olson suggested that hypoxia might also increase cellular levels of polysulfides and other reactive sulfur species, which exist in equilibrium with cellular sulfide, but at much higher concentrations [30]. Like ROS, these are feasible HPV mediators since they can regulate protein function by causing oxidative modification of reactive thiols via sulfuration [442]. However, evidence that hypoxia can elevate sulfide or polysulfide levels in PASM C quickly enough to trigger HPV is lacking. Indeed, production of sulfide and polysulfides in bovine pulmonary artery smooth muscle cells was not potentiated during the first 5 h of a hypoxic (5% O_2) challenge [443].

Nevertheless, the mechanisms of the sustained contractions of rat PA induced by applying exogenous sulfide and hypoxia demonstrate many similarities [18,177]. In both cases, contraction was insensitive to the voltage-gated Ca^{2+} channel antagonist nifedipine, but was inhibited by the antioxidant TEMPOL, and by blockers of the RyR, rho kinase, protein kinase C, and complex III $_{CoQ}$ (myxothiazol). Moreover, sulfide mimicked hypoxia by causing mitochondrial hyperpolarization [142,155]. Sulfide also increased ROS production by PASM C, an effect that was abolished by the knockdown of sulfide quinone oxidoreductase (SQOR), a protein located in the inner mitochondrial membrane. SQOR forms part of the 'sulfide oxidation unit' which metabolizes sulfide and in doing so feeds electrons into the CoQ pool and stimulates respiration [178,444]. The contraction of trout gill to sulfide was also antagonized by the antioxidant diethyldithiocarbamic acid (DETCA) and blockers of mitochondrial complexes 1, 3, and 4 [445]. In light of the

proposal that sulfide stored within the mitochondria as thiosulfide can be mobilized by hypoxia [175], as well as the high affinity of SQOR for sulfide [178], it is conceivable that a small hypoxia-induced rise in intramitochondrial [sulfide] could act through SQOR to increase mitochondrial ROS production at complex III_{CoQ}, thereby causing HPV.

6.4. Effects of Hypoxia on Redox Couples

Cells contain several important redox couples, each of which regulates different aspects of cellular function. Although the three main redox couples (GSSG/GSH, NADP⁺/NADPH, NAD⁺/NADH) interact in important ways, the flow of electrons between them is regulated enzymatically and by membrane transport systems, so that they are not in thermodynamic equilibrium [275]. The redox potential of each couple also differs between cell compartments.

The GSH/GSSG couple functions as a conduit to transfer electrons from NADPH to cell proteins, causing their reduction, but GSSG is also used for protein glutathionylation, thereby contributing to oxidative redox signaling. Although its involvement in maintaining overall cellular antioxidant defense has been unclear and has been overshadowed by that of the thioredoxin/peroxiredoxin pathway, a recent investigation suggests that it may be more important than has been previously thought, at least in some types of cells [100]. As discussed in Section 7.4, Connolly et al. (2013) [18] did not detect a difference between the normoxic redox potentials of the GSH/GSSG couple in rat PA and aorta, which are exposed to a similar normoxic plasma PO₂ in vivo, [58]. They also observed that 45 min of hypoxia caused the oxidation of this couple in PASM. Similarly, the GSH/GSSG ratio in cultured rat PASM was decreased by 2 h of hypoxia [361]. These observations are predicted by the mitochondrial ROS model and do not support the Redox Theory, which posits that the cytoplasm of PASM is relatively oxidized under normoxic conditions and becomes more reduced during hypoxia, whereas in comparison, systemic artery smooth muscle cells are more reduced in normoxia and become oxidized during hypoxia. However, the lengthy exposures to hypoxia used in both studies mean that their results may not reflect the immediate response to hypoxia.

Measurements of pyridine nucleotide (i.e., NAD(P)H) fluorescence in rat and bovine pulmonary artery suggest that the NAD⁺/NADH and/or the NADP⁺/NADPH redox couples may become more reduced during hypoxia [73,351,446,447]. However, these fluorescence measurements did not distinguish between NADH and NADPH, or between their free and bound forms. Since it is the concentrations of the free forms of both pyridine nucleotides that determine their redox potentials, and these constitute only a small fraction of their total contents [448], the effects of hypoxia on the individual redox potentials of both couples remain unclear.

The NADPH/NADP⁺ couple is the ultimate source of electrons which are used by redox relays to reduce oxidized proteins but also provides the electrons which are used by Nox to produce superoxide/H₂O₂. The role of NADPH in determining pulmonary vascular tone during HPV has been investigated in the bovine pulmonary artery by Michael Wolin, Sachin Gupta, and colleagues. This work, which is described in Section 8, has shown, somewhat paradoxically, that a hypoxia-induced increase in NADPH may contribute to HPV both by causing the reduction of PKG1 α via Trx-1/TrxR-1 and by increasing ROS production via Nox, presumably leading to the oxidation of other as yet unidentified effector proteins.

The NADH/NAD⁺ couple is mainly involved in controlling respiration by providing reducing equivalents to the ETC, and also acts to reduce mitochondrial NADP⁺ via the NNT, thereby helping to maintain the mitochondrial matrix in a highly reduced state [67,98]. Because its redox potential is much less negative than that of NADP⁺: NADPH, the NAD⁺:NADH couple is not directly involved in regulating thiol oxidation [448].

However, via the NNT, it exerts an important indirect influence on the $\text{NADP}^+/\text{NADPH}$ ratio, and therefore thiol signaling, in the mitochondrial matrix. An increase in mitochondrial $[\text{NADH}]$ also promotes superoxide production by Complex I, both directly and indirectly, through its effects on KGDH and PDH [98] (Section 3.1.1).

Although an increase in NADH has been suggested to cause HPV through a redox-independent pathway (inhibition of cyclic ADPR hydrolase) [449], the impact of changes in the cytoplasmic or mitochondrial NAD^+/NADH ratio on redox signaling during HPV remains unexplored. However, Lopez-Barneo's laboratory has proposed that a rise in cytoplasmic $[\text{NADH}]$ due to a direct effect of hypoxia makes an important contribution to O_2 sensing in femoral artery myocytes [450] by suppressing voltage-gated Ca^{2+} currents, and in CBCC by inhibiting K_v channel opening [190,191]. A role for NADH in O_2 sensing is supported by evidence that hypoxia can increase the cytoplasmic free $[\text{NADH}]$ in coronary artery myocytes [276], and that NADH binds to K_v channel β subunits with a high affinity, thereby affecting channel activity [451]. However, the hypoxia-induced increase in the cytoplasmic $[\text{NADH}]/[\text{NAD}^+]$ ratio, at least in coronary myocytes, was quite small, and the concentration of NADH used to characterize its effect on the K_v current in CBCC (200 μM) was almost certainly much higher than those present in cells, since the NAD^+/NADH couple in the cytoplasm is almost entirely oxidized (i.e., $\text{NAD}^+ \gg \text{NADH}$). This is potentially problematic for the following reasons: 1. NADPH also binds to K_v channel β -subunits. 2. The cytoplasmic $\text{NADP}^+/\text{NADPH}$ redox couple is almost completely reduced so that $\text{NADPH} \gg \text{NADP}^+$ [275]. 3. The affinity of NADPH for the β subunit is even higher than that of NADH. As a result, it has been questioned whether physiological levels of cytoplasmic free NADH can regulate K_v channel activity, since they may not be sufficient to compete with NADPH for binding to K_v β -subunits [277].

It is noteworthy, however, that because the redox state of the cytoplasmic NAD^+/NADH couple is strongly influenced by the lactate/pyruvate ratio, increases in cellular lactate, which can occur under conditions such as systemic hypoxia, can lead to a significant rise in cellular $[\text{NADH}]$, which may contribute to O_2 sensing in CBCC [452].

7. Critique of the Redox and Mitochondrial ROS Theories

As described in Sections 5.1 and 5.2, the Redox and Mitochondrial ROS theories propose opposite effects of hypoxia on mitochondrial ROS production in PSMCs. The Redox model is based mainly on work by Stephen Archer, Kenneth Weir, Evangelos Michelakis, and colleagues, whereas the most salient findings supporting the Mitochondrial ROS hypothesis have come from the labs of Paul Schumacker, Norbert Weissmann, and Yong-Xiao Wang. Although the papers from the proponents of these two theories present apparently irreconcilable bodies of evidence, other authors have also published results which can be used to assess the extent to which the literature as a whole supports the predictions of either theory.

Predictions of the Redox Theory: The key tenet of the Redox Theory is that hypoxia causes contraction by decreasing mitochondrial ROS production. In setting forth the Redox Theory [19,49,214], Archer and colleagues have invoked observations in the literature which are in line with the following predictions:

1. Blockers of the ETC which induce a fall in mitochondrial ROS production should mimic hypoxia by causing inhibition of PSMC K^+ currents and contraction and should also prevent HPV.
2. Similarly, antioxidants should cause a sustained contraction. Their effect on HPV is more difficult to predict, since a small antioxidant effect might add to that of hypoxia to enhance HPV, while a large antioxidant effect could abolish any further response to hypoxia.
3. Pro-oxidants should prevent and reverse HPV.

Most importantly, hypoxia should diminish mitochondrial production of ROS in PASM, thus causing a fall in cytoplasmic [ROS] and/or a reduction of cytoplasmic redox couples involved in regulating reactive thiols, particularly on Kv channels.

The Redox Theory also predicts that hypoxia, reductants, and blockers of the ETC, which decrease mitochondrial ROS production, should inhibit PASM Kv currents, whereas oxidants should have the opposite effects. Whereas it is well established that these currents are indeed suppressed by hypoxia in PASCs, as described in Section 5.1.2, there is no direct evidence (e.g., changes in reactive thiol oxidation or interactions between Kv α and α subunits in native PASCs) that this is due to redox mechanisms.

Predictions of the mitochondrial ROS model: The Mitochondrial ROS hypothesis is predicated on the concept that hypoxia increases mitochondrial ROS production in PASM, causing a rise in [ROS] in the mitochondrial IMS and the cytoplasm, which induces the oxidation of cytoplasmic protein thiols so as to stimulate contractile effector pathways. This effect of hypoxia is seen as not being unique to PASM, and indeed the Mitochondrial ROS hypothesis grew out of observations that hypoxia increased ROS production in cardiomyocytes [347]. In their reviews, Schumacker and colleagues have focused mainly on evidence that hypoxia increases ROS in PASM and in other types of cells rather than emphasizing the effects of ETC blockers, antioxidants, and pro-oxidants on HPV. However, the responses of PASM to these agents predicted by this hypothesis are reasonably clear:

1. Blockers of the ETC, which diminish mitochondrial ROS production, should not cause contraction under normoxic conditions.
2. ETC blockers like myxothiazol or rotenone, which act at or upstream of the Q_o site of complex III, should inhibit HPV. Antimycin, shown by investigators to increase complex III-mediated ROS production in many types of cells, would be expected to cause PASM to contract during normoxia, but its effects on HPV are difficult to predict.
3. Antioxidants should have no effect on PASM tension in normoxia but should block HPV, and pro-oxidants would be predicted to cause contraction.

In accordance with this theory, there is evidence that multiple contractile effector pathways in PASM are activated by oxidation (Section 5.2.6), although identification of specific reactive cysteines oxidized in the constituent effector proteins of these pathways in these cells is lacking.

In the next several sections, we critically assess these theories by examining the results from the substantial number of papers which have investigated the effects of ETC blockers, and anti- and pro-oxidants on PASM contraction and cytoplasmic Ca²⁺ levels under normoxic and hypoxic conditions. We also review the effects of hypoxia on PASM [H₂O₂], superoxide, and the ambient cytoplasmic thiol/disulfide equilibrium, which have been obtained from studies employing genetically encoded ROS indicators or ESR. Finally, having considered this evidence, as well as the literature relating to the effects of ETC blockers and pro- and antioxidants on PASM Kv currents, we describe what we feel are the strengths and weaknesses of the Redox and Mitochondrial ROS theories.

7.1. Pulmonary Effects of ETC Blockers

Observations that blockers of the proximal ETC decrease PASM [ROS] production [156,272], increase PAP in perfused lung or cause PA constriction [156,272,289], block HPV [272], and mimic the effects of hypoxia on K⁺ currents in smooth muscle cells from PA [156,272], renal arteries [156], and ductus arteriosus [453] have consistently been invoked to support the Redox Theory [19,21,49,137,272]. It is noteworthy, however, that the meaning of the term 'proximal ETC' used in these papers varied; in some cases it referred to the portion of the ETC upstream of ubiquinone [19], whereas in others it included the Q_i site in complex III, since antimycin is described as a blocker of the proximal ETC [49,137].

This ambiguity is worth keeping in mind, since, as described below, it is unlikely that antimycin would produce the same effect on mitochondrial ROS production as drugs such as rotenone which act upstream of ubiquinone.

Results of studies of the effects of ETC blockers on PASM, isolated PA, or lung preparations are summarized in Table 2. Rotenone has been used more than any other ETC blocker to examine the role of mitochondrial ROS production in HPV and was seen to decrease basal ROS levels in most studies where this was examined. Rotenone generally increased basal tension in normoxia and was almost always observed to strongly inhibit HPV or reverse hypoxia-induced rises in $[Ca^{2+}]_{cyt}$. If rotenone was truly decreasing ROS production (which is, however, questionable [121,129,148,191,265,454,455]), these effects are predicted by the Redox Theory. However, as noted previously [3], the generally observed transience of the rotenone-induced contraction does not accord with this model. The ability of rotenone to block HPV is also consistent with the Mitochondrial ROS hypothesis (since it would oxidize the CoQ pool and thereby diminish ROS production by complex III), although the rise in basal tension it caused is not.

Table 2. Effects of ETC blockers on PASM, PA, or perfused lung preparations.

Preparation	Blocker $\mu\text{mol/L}$	Effect on Basal ROS or the Response of Basal ROS to Hypoxia (RTH) [ROS Indicator]	Effect on Basal Tension or $[Ca^{2+}]_{cyt}$ Amplitude Compared to That of HPV	Effect on HPV or the Hypoxia-Induced Increase in $[Ca^{2+}]_{cyt}$ (HIIC)	Ref.
Perfused rat lung	Rot 0.5		transient contraction; ~70% HPV	HPV abolished	[289]
Perfused rat lung	Rot 1.0	↓ 21% basal ROS [luminol]	transient ↑ in PAP; similar to HPV	HPV abolished	[272]
Perfused rat lung	Rot 1.0	↓ 12% basal ROS [lucigenin]	transient ↑ in PAP; similar to HPV	HPV abolished	[272]
Endo-isolated bovine PA	Rot 50			no effect on HPV	[396]
Perfused mouse lung	Rot 50		contraction; similar size to HPV		[222]
Isolated rat PA	Rot 0.1		no effect on basal tone	HPV abolished	[351]
Perfused rat lung	Rot 10.0		transient contraction; ~60% HPV		[456]
Perfused rat lung	Rot 0.013		transient contraction; ~40% HPV	HPV abolished	[23]
Endo-isolated rat PA ¹	Rot 5.0	↓ ~40% basal ROS [DCF]			[156]
Endo-isolated rat PA	Rot 5.0	↓ ~60% basal ROS [lucigenin]	transient contraction; ~60% HPV		[156]
Cultured rat PASM	Rot 1.3			~55% ↓ in HIIC	[353]
Perfused rabbit lung	Rot 0.03–0.35		transient ↑ in PAP abolished by eNOS blocker NG-monomethyl-L-arginine (L-NMMA)	Selective (vs. U46619) dose-dependent block of HPV	[354]
Perfused rabbit lung	Rot 0.04			Phase1 HPV ↓ ~40%, Phase2 HPV abolished	[357]
Perfused rat lung	Rot 50	↓ ~60% basal ROS [LO-12]	↑ in PAP; ~65% HPV		[457]
Mouse PASM	Rot 1.27	↔ basal ROS, RTH ↓ ~60% [DCF]	no effect on basal $[Ca^{2+}]_{cyt}$	~40% ↓ in HIIC	[364]
Perfused rabbit lung	Rot 0.35			↓ ~90%	[155]
Endo-isolated bovine PA	Rot 10	↓ ~50% basal ROS [lucigenin]	no effect on response to 20 mM K^+	No effect on HPV	[458]
Isolated rat PA	Rot 1.0			HPV ↓ ~85%	[328]
Isolated rat PA	Rot 1.0		no effect on basal tone		[18]
Perfused rat lung	Rot 50		↑ basal tension; similar to HPV		[137]

Rat PASMCMC	Rot 0.05		~60% ↑ in basal $[Ca^{2+}]_{Cyt}$	~50% ↓ in <i>HIIC</i>	[137]
Perfused rabbit lung	MPP+ 100		↑ in PAP; abolished by L-NMMA	Specific block of HPV (vs. U46619)	[354]
Mouse PASMCMC	MPP+ 5.0	↔ basal ROS, RTH ↓~85% [DCF]	no effect on basal $[Ca^{2+}]_{Cyt}$	~60% ↓ in <i>HIIC</i>	[364]
Cultured human PASMCMC	MPP+ 5.0	RTH ↓ [Redox Red CC1]	no effect on basal $[Ca^{2+}]_{Cyt}$		[365]
Preparation	Blocker μ mol/L	Effect on basal ROS or the response to hypoxia (RTH) [indicator]	Effect on basal tension or $[Ca^{2+}]_{Cyt}$ amplitude compared to that of HPV	Effect on HPV or Hypoxia-induced increase in $[Ca^{2+}]_{Cyt}$ (HIRiC)	Ref.
Perfused rabbit lung	NPA 2.5–5500		No effect	Blocked HPV and to a lesser extent the constriction to U46619	[354]
Perfused rabbit lung	NPA 600			Phase 1 HPV ↓~60; Phase 2 HPV ↓~75%	[357]
Mouse PASMCMC	NPA 5000	↔ basal ROS, RTH ↓~90% [DCF]	no effect on basal $[Ca^{2+}]_{Cyt}$	~60% ↓ in <i>HIIC</i>	[364]
Cultured human PASMCMC	TTFA 10	RTH ↓ [Redox Red CC1]	no effect on basal $[Ca^{2+}]_{Cyt}$	~60% ↓ in <i>HIIC</i>	[365]
Perfused rabbit lung	TTFA 40–300		no effect on basal tone	Similar block of HPV and constriction to U46619	[354]
Isolated rat PA	Myx 0.1		no effect on basal tone	HPV abolished	[351]
Perfused rat lung	Myx 0.1		no effect on basal tone	HPV ↓ by 85%	[23]
¹ Endo– isolated rat PA	Myx N/A	↔ basal ROS [DCF]			[156]
Cultured rat PASMCMC	Myx 10.3			~85% ↓ in <i>HIIC</i>	[353]
Perfused rabbit lung	Myx		no effect on basal tone	Similar block of HPV and response to U46619	[354]
Rat PASMCMC	Myx 10	↔ basal ROS, RTH ↓~85 [HSP-FRET]	no effect on basal tone	~85% ↓ in <i>HIIC</i>	[361]
Mouse PASMCMC	Myx 10	↔ basal ROS, RTH ↓~85 [DCF]	no effect on basal $[Ca^{2+}]_{Cyt}$	~60% ↓ in <i>HIIC</i>	[364]
Isolated rat PA	Myx 1.0			HPV ↓ by 52%	[328]
Isolated human PA	Myx 10			HPV abolished	[365]
Cultured human PASMCMC	Myx 10	RTH ↓~85 [Redox Red CC1]	no effect on basal $[Ca^{2+}]_{Cyt}$	60% ↓ in <i>HIIC</i>	[365]
Perfused rat lung	Ant 5		transient ↑ in PAP, ~60% of HPV	HPV abolished	[289]
Perfused rat lung	Ant 5	↓ 38% basal ROS [luminol]			[272]
Perfused rat lung	Ant 5	↓ 24% basal ROS [lucigenin]	↑ basal tone, ≥ HPV		[272]
Perfused rabbit lung	Ant 10	↓ ~50% basal ROS [lucigenin]			[459]
Perfused rat lung	Ant 0.0018		no effect on basal tone	No effect on HPV	[23]
Cultured rat PASMCMC	Ant 1.82			No effect on HPV	[353]
Endo– isolated rat PA	Ant 50	↓ 35% basal ROS [Amplex Red]			[156]
Perfused rabbit lung	Ant 0.001–0.004		No effect on basal tone	Specific block of HPV vs. U46619; less specific after L-NMMA	[354]
Perfused rat lung	Cyn 1000		Transient ↑ in PAP; ~75% of HPV		[289]
Perfused pig lung	Cyn 50		Large transient and small sustained ↑ PAP		[292]

Perfused rat lung	Cyn 1.5–15	Transient ↑ of 15–35% [luminol]	↑ PAP, same amplitude as HPV	No effect on HPV	[272]
Perfused rat lung	Cyn 10		Transient ↑ in PAP; ~110% of HPV		[23]
Isolated rat PA	Cyn 100		No effect on basal tone	Phase 2 HPV ↑ by ~50%	[351]
Endo–isolated rat PA	Cyn 1.0	↔ basal ROS [DCF]			[156]
Rat PASMC	Cyn 10		↑ basal $[Ca^{2+}]_{cyt}$	No effect on HPV	[353]
Isolated Endo– and ¹ Endo + rat PA	Cyn 100		Small increase in tension in pre-constricted PA	HPV abolished	[460]
Rat PASMC	Cyn 1.0		No effect on basal tone	Small ↑ in HIIC	[361]
Perfused rabbit lung	Cyn 50			Phase 1 HPV ↓ by ~25%, Phase 2 abolished	[357]
Perfused rabbit lung	Cyn 10			HPV abolished	[155]
Isolated rat PA	Cyn 10		No effect on basal tone		[18]
Perfused rat lung	NaN ₃ 10000		Transient ↑ in PAP, ~33% of HPV		[289]
Cultured rat PASMC	NaN ₃ 1000	↔ basal ROS ↔ RTH [DCF]	No effect on basal $[Ca^{2+}]_{cyt}$	No effect on HIIC	[364]
Cultured human PASMC	NaN ₃ 1000	↔ RTH [Redox Red CC]		$[Ca^{2+}]_{cyt}$ during hypoxia unchanged	[365]
Isolated human PA	NaN ₃ 1000			No effect on HPV	[365]

Complex I blockers: Rotenone (ROT), 1-methyl-4-phenylpyridinium (MPP⁺); Complex II blockers: 3-nitropropionic acid (NPA), 2-tenoyltrifluoroacetone (TTFA); Complex III blockers: myxothiazol (Myx; blocks at Qo), antimycin (Ant; blocks at Qi), 2-n-heptyl-4-hydroxyquinoline-N-oxide (HQNO; blocks at Qi). Complex IV blockers: cyanide (Cyn), sodium azide (NaN₃). ¹ Endo— and Endo + indicate that measurements were made from endothelium-denuded and endothelium intact arteries, respectively. Abbreviations: RTH = response of ROS signal to hypoxia; DCF = dichlorofluorescein; HIIC = hypoxia-induced increase in $[Ca^{2+}]$ signal. ↑, ↓, and ↔ arrows refer to an increase, decrease, or no change in, respectively.

Unlike rotenone, myxothiazol was not observed to alter basal ROS production in any of the three studies in which this was studied. Studies of mitochondria from various tissues show that myxothiazol generally has no effect on, or increases, basal mitochondrial ROS production, depending on the cell type and substrates used to fuel oxidative phosphorylation [126,132,349,461,462]. Myxothiazol strongly antagonized the hypoxia-induced increase in ROS recorded in several studies and was consistently seen to depress HPV without itself altering basal tone or $[Ca^{2+}]_{cyt}$. Since myxothiazol blocks ROS production by complex III (although it can increase ROS release upstream of complex III), this combination of effects is generally consistent with the Mitochondrial ROS hypothesis. In light of the observations that myxothiazol did not change basal [ROS], its lack of effect on basal tone also fits with the Redox Theory. However, since the potential effects of myxothiazol on any hypoxia-induced decrease in ROS production by complex I in PASMC are uncertain, the implications for the Redox Theory of the block of HPV it exerted are unclear.

Antimycin was seen to lower (six studies) or not affect (two studies) ROS levels. These results are perplexing, since there is a widespread consensus, emerging from numerous studies in multiple types of cells, that antimycin increases ROS production by complex III, and can also do so at more proximal sites in the ETC by causing reduction of the CoQ pool [126,133]. If antimycin was indeed causing a paradoxical fall in basal ROS

production, the increase in basal contraction it usually (in 4 of 5 studies) evoked was in accord with the Redox Theory. However, although antimycin blocked HPV in 3 studies, as predicted by the Redox Theory, it more often (six studies) had no effect on or increased the response to hypoxia, which does not conform to this model. On the other hand, the normoxic contraction it often caused is consistent with the Mitochondrial ROS hypothesis [23,347], although the apparent fall in ROS production is not. In any case, since no single investigation of antimycin in PA or PASMC evaluated its effects on all of the key parameters (basal [ROS] and tension development or $[Ca^{2+}]_{cyt}$ in normoxia and hypoxia), it is difficult to draw any meaningful conclusions from its observed effects.

The effect on HPV of the complex IV blockers cyanide and NaN_2 is difficult to predict from either the Redox Theory or the Mitochondrial ROS hypothesis. However, since these drugs would be expected to mimic the inhibition of complex IV caused by hypoxia, they can induce reduction of the CoQ pool, which might increase ROS production by complex III. Thus, evocation of contraction by these drugs would be consistent with the Mitochondrial ROS hypothesis [346], although this would probably depend on whether the degree of the block was similar to that caused by hypoxia. Cyanide was observed to cause contraction in most studies in which it was used, but this was never seen with sodium azide. The reasons for this difference are not obvious.

In summary, the use of conventional ETC blockers to study HPV has yielded inconsistent results which, when taken together, do not provide convincing support for either the Redox or Mitochondrial ROS models. More importantly, although these drugs, if employed intelligently, can provide important information about how hypoxia influences mitochondrial ROS production, their use has a number of drawbacks that need to be taken into account when interpreting their effects. Firstly, the effects of ETC blockers on overall mitochondrial ROS production are difficult to predict. For example, rotenone inhibits ROS production at complex I_{CoQ} and downstream sites of the ETC, but tends to have the opposite effect at complex I_F, O_F, and P_F [121]. Secondly, inhibition of respiration caused by ETC blockers has manifold impacts on mitochondrial function, which go beyond their direct effects on ROS production [463]. One of these, which is likely to have immediate consequences for vascular tone, is that ETC blockers depolarize $\Delta\Psi_m$ in isolated mitochondria and intact cells from many cells/tissues, including PA [464–466]. This inhibits mitochondrial Ca^{2+} uptake, leading to complex effects on Ca^{2+} signaling in smooth muscle cells, which generally tend to inhibit contraction by interfering with the generation of Ca^{2+} waves [466–468] and by causing the opening of BK_{Ca} channels, even though global $[Ca^{2+}]_{cyt}$ may not change [468] or even increase [466]. Also, by depressing O_2 consumption, inhibition of respiration can increase the intracellular O_2 concentration [469–471]. This could potentially decrease the effect of hypoxia on extramitochondrial O_2 sensors, thereby attenuating HPV. In addition, there is evidence from Hernansanz-Agustin et al. [194] that rotenone can both mimic and prevent hypoxia-induced complex I deactivation. Furthermore, rotenone at concentrations of $>0.5 \mu M$, as used in many but not all HPV studies, can block K^+ channels [326]. Finally, it is apparent from the comprehensive investigation by Weissmann and colleagues that the observed effects of ETC blockers on PAP in perfused lungs are greatly distorted by their effects on NO release [354].

7.2. Pulmonary Vascular Effects of Antioxidants

As shown in Table 3, antioxidants have often been used to characterize how ROS affect pulmonary vascular tone under normoxic and hypoxic conditions.

The enzymes SOD and catalase have been used more frequently than any other antioxidant agents to study HPV. In most investigations, these enzymes, used either alone or in combination, had no effect on basal tone or HPV. It is unclear whether this reflected a genuine lack of response or was due to their poor entry into cells. However, application

of high concentrations of SOD and catalase together to isolated perfused rat lung greatly increased HPV with no effect on basal PAP, and treatment with liposomes containing a lower concentration of both enzymes, a procedure designed to increase their entry into cells, increased basal tone and caused a smaller enhancement of HPV [285]. These responses are generally in accord with the Redox Theory. Conversely, HPV in pig PA was inhibited by the combination of SOD and catalase [358], which supports the mitochondrial ROS hypothesis. Interestingly, in both preparations, SOD on its own exerted an effect similar to that of SOD and catalase in combination, which implies that superoxide itself must have been acting as a mediator in HPV.

Most of the studies that employed chemical antioxidants found that they inhibited HPV without affecting basal tone. Nevertheless, an antioxidant-induced increase in basal tension, which would be predicted by the Redox Theory, was seen by some investigators. This was observed most often in bovine pulmonary arteries (BPA), in which ebselen, DTT, and DHLA raised basal tone (although NBT and Tiron did not). HPV in BPA was inhibited by NBT and ebselen, not affected by Tiron, and increased by DTT and DHLA. It is difficult to fit these results into a coherent pattern, although Wolin and colleagues, who carried out most of these investigations, concluded that a fall in [ROS] contributes to HPV (see Section 8). However, they also found that HPV in BPA is not blocked by rotenone [458], implying that O₂ sensing in BPA is independent of mitochondrial ROS production. Likewise, Olson's laboratory, which reported the effects of DTT and DHLA, ascribed these to an increase in cellular [H₂S] rather than to an effect on mitochondrial [ROS] [175].

Apart from the experiments carried out in BPA, Table 3 shows that studies employing antioxidants have mostly reported that they inhibit HPV but do not affect basal PA tone or [Ca²⁺]_{cyt}, observations which tend to support the Mitochondrial ROS hypothesis.

Nevertheless, although if employed correctly antioxidants can provide information about how ROS regulate cellular processes, their use for this purpose, especially if not supported by parallel measurements of cellular superoxide and H₂O₂, has come under increasing criticism [386]. Each antioxidant has a unique mechanism of action, which is often complex and/or not fully understood (e.g., TEMPOL [472]; TIRON [473] N-acetylcysteine [474]; ebselen [475]). Also, some antioxidants (e.g., N-acetylcysteine, manganese(III)tetrakis(1-methyl-4-pyridyl)porphyrin—MnTMPyP) can increase the formation of reactive sulfur species [474,476] which could potentially mimic the effect of H₂O₂ on reactive cysteines [31]. In addition, SOD-mimetics such as MnTMPYP would be predicted to cause an increase in cellular H₂O₂, along with the fall in superoxide [477], although this may not be the case for TEMPOL and Tiron, which, although often described as SOD-mimetics, also apparently suppress the production or effect of H₂O₂ [472,478]. This is an important consideration with regard to the effects of antioxidants on HPV, since most studies, particularly those using perfused lungs, did not include measurements of ROS (and those that did used ROS indicators which have since been largely discredited).

The overexpression or knockdown of cellular antioxidant enzymes allows the manipulation of redox tone in specific cellular compartments and avoids some of the uncertainties inherent in the use of chemical antioxidants or proteins such as SOD and catalase [386]. Investigations using cultured PASM C from mice in which catalase was overexpressed by transfection [353,368] found that this blunted the rise in [Ca²⁺]_{cyt} evoked by hypoxia in PASM C. Freshly isolated PASM C from mice which had been genetically modified to overexpress catalase also demonstrated a decreased hypoxia-induced increase in [Ca²⁺]_{cyt} compared to cells from wildtype controls [364]. A similar effect was seen in PASM C from mice in which glutathione peroxidase had been overexpressed in order to enhance the metabolism of H₂O₂, whereas cells from glutathione peroxidase knockouts demonstrated an augmented response of [Ca²⁺]_{cyt} to hypoxia. It could be argued, in line with the Redox Theory, that catalase or glutathione peroxidase overexpression was

causing a basal increase in $[Ca^{2+}]_{\text{cyt}}$ by reducing cytoplasmic redox couples, thereby preventing a further similar effect of hypoxia. This possibility, however, is not supported by the observation [364] that the basal level of $[Ca^{2+}]_{\text{cyt}}$ was not altered in PASMC from any of the genetically altered mice. Therefore, these results consistently agree with the predictions of the Mitochondrial ROS hypothesis.

More recently, Sommer et al. [141] reported that HPV was absent in isolated perfused lungs from mice expressing AOX, which has an antioxidant effect since it diminishes ROS production by Complex III [388]. Basal superoxide production was similar in the PASMC from AOX mice and wildtype controls, as was the resting membrane potential, indicating that baseline ROS production by Complex III was not influencing the cytosolic redox state or K^+ channel activity under normoxic conditions. These results provide strong evidence favoring the Mitochondrial ROS hypothesis.

Table 3. Effects of antioxidants on HPV.

Preparation	Antioxidant	Effect on [ROS]	ROS Indicator	Effect on Basal Tension/PAP or $[Ca^{2+}]_{\text{cyt}}$	Effect on Response to Hypoxia	Ref.
In situ blood-perfused rat lung	Catalase 5000 U				No effect on HPV	[479]
Pig PASMC	Catalase 200U/mL	Hypoxia-induced $\uparrow$ in ROS abolished	DCFH		No effect on HPV	[358]
In situ-perfused rat lung 0% O_2	Catalase 2×10^{-3} U/kg body wgt				No effect on HPV	[480]
Perfused mouse lung 2% O_2	Catalase 200 U/mL					[481]
¹ Endo-bovine PA	Catalase 1 μ M		Luminol/HRP	² No effect on 25 mM K contraction	No effect on HPV	[482]
Endo-bovine PA	Catalase 1 μ M	³ ROS signal virtually abolished	Luminol/HRP			[458]
Isolated mouse PA	Catalase 1 μ M				No significant effect on HPV	[483]
Pig proximal PA	Catalase + PEG-Catalase each 500 U/mL				HPV $\downarrow$ by ~85%	[484]
Isolated sheep PA	SOD 150 U/mL				No effect on HPV at 6 and 21 min	[485]
Perfused rat lung	SOD 100 U/mL			$\uparrow$ in PAP ~25% of the amplitude of HPV		[272]
Perfused rabbit lung	SOD 100 mg/mL	~50% $\downarrow$ in basal ROS	lucigenin			[459]
In situ blood-perfused rat lung	SOD 2750 U				No effect on HPV	[479]
Perfused rabbit lung	SOD 100–2000 U/mL			No effect on PAP	No effect on acute HPV	[486]
Endo + isolated pig PA	SOD 150 U/mL			No effect on basal tension	HPV $\downarrow$ by ~75%	[358]
Pig PASMC	SOD 150 U/mL	Hypoxia-induced $\uparrow$ in ROS decreased by ~75%	DCFH			[358]
In situ-perfused rat lung 0% O_2	SOD 2mg/kg body weight				No effect on HPV	[480]

Endo-bovine PA	SOD 1 μ M	No effect on 25 mM K contraction		HPV $\downarrow$ by ~40%	[482]
Isolated rat 2nd- & 3rd-order PA	SOD 200 U/mL	No effect on ba-DHE, Mito-sal ROS]	Sox Red		[420]
Isolated pig proximal PA	SOD 500 U/mL + MnTMPyP 100 μ M			non-significant $\downarrow$ in HPV	[484]
Perfused rat lung	SOD 245 U/mL + CAT 50 U/mL	No effect on basal PAP		No effect on HPV	[285]
Perfused rat lung	SOD 490 U/mL + CAT 100 U/mL	No effect on basal PAP		$\uparrow\uparrow$ HPV	[285]
Perfused rat lung	SOD + CAT in liposomes low dose	$\uparrow$ normoxic PAP		$\uparrow$ HPV	[285]
Perfused rat lung	SOD + CAT in liposomes high dose	$\uparrow\uparrow$ normoxic PAP		No effect on HPV	[285]
Endo + pig PA	SOD 150 U/mL + CAT 200 U/mL	No effect on basal tension		HPV abolished	[358]
Pig PASMC	SOD 150 U/mL + CAT 200 U/mL	Hypoxia-induced increase in [ROS] abolished			[358]
Perfused rat lung	Ebselen 50 μ M	No effect on basal PAP		HPV $\downarrow$ by ~85%	[23]
Endo-isolated bovine PA	Ebselen 100 μ M	Basal ROS (attributed to H ₂ O ₂) $\downarrow$ ~80%	Luminol/HRP	20 mM K contraction (used by this laboratory as a measure of basal tension) $\uparrow$ by ~100%	HPV $\downarrow$ by ~65% [482]
Endo-isolated bovine PA	Ebselen 100 μ M	Basal ROS (attributed to O ₂ ⁻) $\downarrow$ ~80%	Lucigenin	25 mM K contraction $\uparrow$ 30–50%	HPV $\downarrow$ by ~65% [458]
Isolated rat 2nd- & 3rd-order PA	Ebselen 100 μ M			No effect on Phase 1 HPV Phase 2 HPV $\downarrow$ ~40%	[18]
Perfused rat lung	⁴ DETCA 1 mM	No effect		HPV $\downarrow$ by 70%	[23]
Perfused rabbit lung	DETCA 0.1–10 mM			Dose-dependent $\downarrow$ HPV but also of U46619 contraction	[355]
Endo-isolated bovine PA	DETCA 10 mM	'O ₂ ⁻ ' $\uparrow$ ~220%	Lucigenin	20 mM K contraction (used by this laboratory as a measure of basal tension) $\uparrow$ ~400%	HPV $\downarrow$ by 70% [482]
Isolated extra-lobar guinea pig PA	DETCA 5 mM	'H ₂ O ₂ ' $\downarrow$ ~80%	Luminol/HRP	$\uparrow$ basal tension; $\leq$ 25% of the amplitude of HPV	HPV $\uparrow\uparrow$ [346]
Endo-isolated bovine PA	NBT 0.3 mM	Basal ROS $\downarrow$ ~55%	Lucigenin	No effect on basal tone	HPV $\downarrow$ by 90% [487]
Isolated rat PA	NBT 0.5 μ M			Slight $\downarrow$ of phase 2 HPV	[488]
Perfused rabbit lung	⁵ NBT + ⁶ L- NMMA 0.2–1.0 mM			No effect on basal tone	Dose-dependent $\downarrow$ in HPV [486]
Perfused rabbit lung	NBT 0.2–1 mM + L-NMMA + acetylsalicylic acid			No effect on basal tone	Dose-dependent $\downarrow$ in HPV [486]
Isolated extra-lobar guinea pig PA	⁷ TETA 5 mM			Sustained contraction; $\leq$ 10% of the amplitude of HPV	HPV $\uparrow\uparrow$ [346]
Perfused rabbit lung	TETA 1–25 mM			No effect on basal tension	Dose-dependent $\downarrow$ in HPV [355]

Perfused rabbit lung	TETA 1–25 mM +L-NMMA			No effect on basal tension	Dose-dependent ↓ in HPV	[355]
Perfused rabbit lung	TETA 25 mM +L-NMMA				↓↓ both phases of HPV	[356]
Endo-isolated bovine PA	Tiron 10 mM	basal ROS ↓ ~55%	Lucigenin	No effect on basal tension	No effect on HPV	[487]
Perfused rabbit lung	Tiron 0.2–0.8 mM	basal ROS (said to be superoxide) ↓ ~90%	Lucigenin			[459]
Perfused rabbit lung	Tiron 0.01–10 mM			No effect on basal tension	No effect on Phase 1 HPV	[486]
Rabbit PA	Dithiothreitol 3 mM			↑ basal tension		[340]
Isolated rabbit PA	Dithiothreitol 3 mM			↑ basal $[Ca^{2+}]_{cyt}$ and membrane depolarization		[340]
Rat PASMCMC	Dithiothreitol 3 mM			↑↑ basal tension		[340]
Rat PASMCMC	Dithiothreitol 3 mM			↑↑ basal $[Ca^{2+}]_{cyt}$		[340]
Isolated rat PA	Dithiothreitol 1 mM				Phase 2 HPV completely reversed	[488]
Isolated rat PA	Dithiothreitol 1 mM				Phase 2 HPV abolished	[441]
Isolated bovine PA	Dithiothreitol 0.001–10 mM				↑ HPV with ≥ 100 mM DTT	[175]
Isolated bovine PA	⁸ DHLA 0.1 mM			↑ basal tension in some arteries	↑ HPV in some arteries	[175]
Perfused rat lung	⁹ PDTC 0.5 μM			No effect on basal PAP	HPV ↓ by ~65%	[23]
Rat PASMCMC	PDTC 10 μM	no basal effect	HSP-FRET	No effect on basal $[Ca^{2+}]_{cyt}$	↓ hypoxic-induced increase in $[Ca^{2+}]_{cyt}$ by ~90%	[361]
Isolated endo+ and endo- rat PA	Co-enzyme Q 500 μM			Small ↑ in basal tension, cell depolarization		[337]
Isolated endo+ and endo- rat PA	Duroquinone 5–500 μM			Large ↑ in basal tension, cell depolarization		[337]
Rat PASMCMC	N-acetylcysteine 500 μM	no basal effect	HSP-FRET		↓ hypoxic-induced increase in $[Ca^{2+}]_{cyt}$ by ~95%	[361]
Perfused rabbit lung	Trolox 20 μM				HPV ↓ by ~47% at 5 min AND abolished at 30 min	[489]
Perfused rat lung	TEMPOL ~2.9 mM				HPV ↓ by ~90%	[490]
	TEMPOL ~2.9 mM + L-NAME				HPV abolished	[490]
Isolated 2nd- and 3rd-order rat PA	TEMPOL 3 mM				No effect on Phase 1 HPV, Phase 2 abolished	[18]
Perfused mouse lung	MitoQ 0.5 mM				HPV ↓ by ~50%	[423]
Perfused mouse lung	MnTMPyP 1 mM			No effect on basal PAP	No effect on HPV	[481]
Mouse lung slices	EUK-34 20 mM				Abolished hypoxia-induced rise in $[Ca^{2+}]_{cyt}$	[368]
Isolated sheep PA	N-t-butyl-a-phenylnitrone 300 mM				No effect on HPV at 6 and 21 min	[485]

In situ-perfused rat lung	Dimethylthiourea 100 mg/kg bw			HPV ↓ by ~40%	[480]
In situ-perfused rat lung	Dimethylsulfoxide 1 mL/kg bw			HPV ↓ by ~40%	[480]
In situ-perfused rat lung	Allopurinol 30 mg/kg bw			HPV ↓ by ~40%	[480]
In situ blood-perfused rat lung	Desferrioxamine 2 mg/kg bw			No effect on HPV	[479]
Primary cultured rat micro-vascular PSMCs	Catalase overexpression			↓ hypoxia-induced rise in [Ca ²⁺] _{cyt} by ~40%	[353]
Freshly isolated mouse PSMCs	Catalase overexpression	Basal ROS ↔, but hypoxia-induced rise attenuated		↓ hypoxia-induced rise in [Ca ²⁺] _{cyt} by ~50%	[364]
Mouse lung slices	Catalase overexpression	Abolished hypoxia-induced rise in Ro-GFP oxidation	Ro-GFP	Attenuated hypoxia-induced rise in [Ca ²⁺] _{cyt}	[368]
Primary cultured rat micro-vascular PSMCs	Cytosolic catalase overexpression	Abolished hypoxia-induced rise in Ro-GFP oxidation	Ro-GFP		[366]
Primary cultured rat micro-vascular PSMCs	Mitochondrial catalase overexpression	No effect on hypoxia-induced fall in mitochondrial Ro-GFP oxidation			[366]
Primary cultured rat micro-vascular PSMCs	Glutathione peroxidase overexpression	↓ hypoxia-induced rise in HSP-FRET oxidation	HSP-FRET	↓ hypoxia-induced rise in [Ca ²⁺] _{cyt}	[361]
Freshly isolated mouse PSMCs	Gpx overexpressing mice	↓ hypoxia-induced rise in ROS		↓ hypoxia-induced rise in [Ca ²⁺] _{cyt}	[364]
Freshly isolated mouse PSMCs	Gpx knockout mice	↑ hypoxia-induced rise in ROS		↑ hypoxia-induced rise in [Ca ²⁺] _{cyt}	[364]
Mouse PA	AOX expression	↓↓ hypoxia-induced rise in ROS		HPV almost abolished	[141]

¹ Endo + and Endo – refer to the use of endothelium-intact and -denuded arteries in the experiment, respectively. ² The contraction induced by 25 mM K in this study was used by the authors as an index of basal reactivity. ³ The ROS signal provided by the lucigenin/HRP assay used in this study was considered by the authors to reflect extracellular ROS. ⁴ diethyldithiocarbamic acid; ⁵ nitro blue tetrazolium; ⁶ (N(G)-monomethyl L-arginine); antagonist of endothelial nitric oxide synthase; ⁷ triethylenetetramine; ⁸ dihydrolipoic acid; ⁹ pyrrolidinedithiocarbamate.

7.3. Effects of Oxidants on Basal Tone and HPV

Table 4a,b list the effects of H₂O₂ and other oxidants on pulmonary vascular tone in normoxia and also on HPV and contractions caused by various vasoconstrictors. The most

widely studied oxidant, H_2O_2 , was used over a wide concentration range (0.5 mM–10 mM), raising the question of which concentrations are likely to produce physiologically relevant increases in intracellular $[\text{H}_2\text{O}_2]$.

As discussed in Section 4, the extracellular $[\text{H}_2\text{O}_2]$ in vivo is thought to be 1–5 μM [84], whereas the average cytoplasmic $[\text{H}_2\text{O}_2]$ is generally said to be in the low nM range [48,111,491] but may be higher in the subplasmalemmal region. According to the mathematical model [249] developed by Sykes and colleagues, and assuming that PASMCM have a width of roughly four microns [492], in the presence of an extracellular $[\text{H}_2\text{O}_2]$ of 2 μM the basal subplasmalemmal and core $[\text{H}_2\text{O}_2]$ would be ~6 and ~2 nM, respectively, and applying 10 μM H_2O_2 would increase these H_2O_2 concentrations by ~30 and ~10 nM. It has also been reported [278,279] that the half-maximal oxidation of the cytoplasmic GSSG/GSH redox couple in HeLa cells and cardiac myocytes was induced by extracellular application of ~10 and ~20 μM H_2O_2 , respectively. In both types of cells, the threshold for oxidation of GSH was near ~5 μM $[\text{H}_2\text{O}_2]$ and oxidation became maximal at ~50 μM $[\text{H}_2\text{O}_2]$. Interestingly, diamide oxidized GSH over a similar range of concentrations [279]. Since the cellular GSSG/GSH ratio is a key factor in setting the cytoplasmic redox tone [362] and is crucial for maintaining cellular thiol redox balance [116], these considerations suggest that application of exogenous H_2O_2 or diamide in the concentration range below ~50 mM may be suitable for mimicking physiologically relevant levels of oxidant signaling.

Table 4. (a) Effects of H_2O_2 on basal tension/ $[\text{Ca}^{2+}]_{\text{cyt}}$, and contractile responses to hypoxia and vasoconstrictors in PASMCM, PA, and perfused lung; (b) effects of other oxidants on basal tension/ $[\text{Ca}^{2+}]_{\text{cyt}}$ and contractile responses to hypoxia and vasoconstrictors in PASMCM, PA, and perfused lung.

(a)					
Preparation	$[\text{H}_2\text{O}_2]$ $\mu\text{mol/L}$	Effect of H_2O_2 on Basal Tension or $[\text{Ca}^{2+}]_{\text{cyt}}$	Effect of H_2O_2 on HPV	Effect of H_2O_2 on Response to Vasoconstrictor	Ref.
Perfused rat lung	0.5–1.0	No effect on basal PAP	HPV abolished	Rise in PAP caused by angiotensin 2 ↓ by 72%	[282]
Bovine PA	0.1–100			Partial, concentration-dependent inhibition of 5-HT, histamine and high K^+ contractions	[493]
¹ Endo + and – rabbit PA	100–1000	Large, slow, concentration-dependent contraction			[494]
Rat large PA	200	Small biphasic contraction			[495]
Rat PA	30	Small contraction	No effect on HPV	No effect on contraction to 5 mM $\text{PGF}_{2\alpha}$	[496]
Rat PA	500	Small contraction	HPV almost abolished	Contraction to $\text{PGF}_{2\alpha}$ abolished; high K^+ contraction ↓ >50%	[496]
Rat PA	10,000	Small slow contraction			[497]
Porcine intra-parenchymal PA	100–3100	Slow, sustained, concentration-dependent contraction			[497]
Rat pulmonary vascular pericytes	10–1000	Concentration-dependent relaxation followed by contraction			[498]

Perfused rat lung	100	Contraction; amplitude = ~55% of HPV		[23]	
Rat cultured PASMCM	50	↑in [Ca ²⁺] _{Cyt} ≥ that caused by hypoxia		[140]	
Rat main PA	50	Large increase in tension		[499]	
Endo—bovine PA	100	No significant effect on contraction to 30 K ⁺		[500]	
Endo—bovine PA	1000	↑ 30 mM K ⁺ -induced contraction by ~40%		[500]	
Rat PASMCM	10–1000	Concentration-dependent ↑in [Ca ²⁺] _{Cyt}		[415]	
Endo—rat PA	1–100	30 mM H ₂ O ₂ caused sustained contraction with EC ₅₀ 12 mM, amplitude ² 17% max. Also ↑ in [Ca ²⁺] _{Cyt}	30 mM H ₂ O ₂ caused transient endothelium-dependent 29% relaxation of constriction induced by 10 mM PGF _{2α}	[425]	
Pig PA	100	Small transient ↑ in [Ca ²⁺] _{Cyt} and tension; then, large slow ↑ tension		[501]	
Endo—bovine PA	1–1000	Conc.-dependent relaxation of contraction to 100 nM 5-HT + hypoxia		[502]	
Rat PA	10–100	Conc.-dependent sustained contraction		[328]	
Rat PA	100	Small, slowly developing contraction		[412]	
Rat PA	110	Contraction, 23% max		[503]	
Rat PA	30	Small transient and sustained contractions (12 & 5% max.)		[421]	
Rat cultured PASMCM	10	2-fold ↑ in store-operated Ca ²⁺ entry		[414]	
(b)					
Preparation	Oxidant & conc. in μmol/L	Effect on basal tension or [Ca ²⁺] _i	Effect on HPV or hypoxia-induced ↑in [Ca ²⁺] _{Cyt}	Effect on response to constrictor	Ref.
perfused rabbit lung	t-BOOH 200	Sustained contraction due to unidentified prostanoid			[282]
Endo—rat PA	t-BOOH 10.0	Transient followed by smaller sustained ↑ in tension			[327]
Rat PA	t-BOOH 10–100	Concentration-dependent sustained ↑ in tension			[328]
Perfused rat lung	t-BOOH 100	75% reversal of ongoing HPV			[137]
Rat PASMCM	t-BOOH 1.0	Reversal of hypoxia-induced increase in [Ca ²⁺] _{Cyt}			[137]
In vivo PVR in dogs	Diamide 5mg/kg	Almost complete reversal of ongoing HPV		Increase in PVR caused by PGF _{2α} was ↓ by >50%	[281]
Perfused rat lung	Diamide ~200	Transient ~70% ↓ in HPV			[283]

Endo–bovine PA	Diamide 1.0	Slow ↑ in basal force; due to activation of ERK		[500]
Endo + & – rat PA	Diamide 100		Relaxation of contraction to coenzyme Q/duroquinone, E_m hyperpolarization	[337]
Endo + & – Rat PA	diamide 1–1000		Relaxed contractions to phenylephrine (IC_{50} = 58 mM) & 40 mM K^+ (IC_{50} = 335 mM)	[341]
Perfused rat lung	glu/glu ox	No effect on control PAP, but ↑ PAP after indomethacin	HPV abolished	[282]
Rat main PA	glu/glu ox	Large increase in tension		[499]
Endo- rat small PA	glu/glu ox	Rapid-onset rise in $[Ca^{2+}]_{cyt}$ (22% of high K^+ response)		[425]
Perfused rat lung	x/xo	No obvious effect	30–50% ↓ in HPV	[283]
Perfused rat lung	x/xo	No effect	~55% ↓ in HPV Angiotensin 2 contraction ↓ by 58%	[285]
Rat PA	x/xo	Sustained dose-dependent contraction (~9% max.)		[420]
Endo + & – rat PA	LY83583 10.0	Contraction of ~15% high K^+ , ↑↑ by 5 mM $PGF_{2\alpha}$ pre-tone		[420]
Rat PA	Ly83583 10.0	Contracted U46619-preconstricted PA in presence of eNOS blocker		[331]
Rat PA	LY83583 10.0	Moderate sustained contraction		[412]
Rat PA	Ly83583 0.1–10	No contraction, but ↑↑ responses to high K^+ and U46619		[421]
Rat PA	Menadione 10.0	Very small sustained contraction		[420]
Rat PA	BCNU 100	No effect, but ↑ contractile response to LY83583		[420]
Rabbit fetal resistance PA	DTNB 1000	No effect on basal tone under normoxic conditions	Relaxation of PA pre-constricted by hypoxia + phenylephrine	[340]

¹ E^- and E^+ refer to endothelium-denuded and -intact arteries. ² 'max' refers to the amplitude of the high K^+ solution. Abbreviations: t-BOOH = tert-butyl hydroperoxide; glu/glu ox = glucose/glucose oxidase; x/xo = xanthine/xanthine oxidase; BCNU = 1,3-bis(2-dichloroethyl)-1-nitrosourea; DTNB = 5,5-dithio-bis-2-nitrobenzoic acid; 5-HT = 5-hydroxytryptamine; ERK = extracellular signal-regulated kinase. ↑ and ↓ arrows refer to an increase or decrease in, respectively.

As shown in the tables, the results from almost all studies, including those that applied H_2O_2 within or near this concentration range (Table 4a), form a consistent pattern, which was also observed when other oxidants were tested (Table 4b): H_2O_2 caused a sustained contraction/rise in $[Ca^{2+}]_{cyt}$ in PA/PASMC if applied under baseline normoxic conditions, but induced relaxation or reversed rises in $[Ca^{2+}]_{cyt}$ in PA/PASMC which had been constricted/stimulated by hypoxia or vasoconstrictors.

These opposing actions of H_2O_2 were not necessarily due to the use of different preparations or experimental set-ups by various investigators, since both were recorded when all conditions except the presence or absence of pre-constriction were identical [425].

Notably, a similar pattern has been observed upon application of H_2O_2 to systemic arteries [504,505].

An investigation of the constricting effects of H_2O_2 in PA [425] found that the EC_{50} for H_2O_2 -induced contraction in rat PA was $12\ \mu\text{M}$. $30\ \mu\text{M}$ H_2O_2 caused a rapidly developing and sustained contraction of small rat PA, which was somewhat larger than that evoked by hypoxia (~ 22 – 25 Torr) under similar conditions [441]. The H_2O_2 -induced contraction was endothelium-independent and associated with a rise in $[\text{Ca}^{2+}]_{\text{cyt}}$, which was reversed by ryanodine but insensitive to the removal of extracellular Ca^{2+} , indicating that it was due to Ca^{2+} release from the sarcoplasmic reticulum (see also [415]). The initial phase of the contraction (5–10 min) was partly blocked by the RhoK antagonist Y-27632 and was only slightly depressed by the membrane-permeable Ca^{2+} chelator BAPTA-AM, suggesting a role for Ca^{2+} sensitization. The sustained contraction was BAPTA-sensitive but not affected by Y-27632. H_2O_2 contracted α -toxin-permeabilized PA; this effect was abolished by blockers of conventional PKC isoforms, which also strongly depressed the contraction in intact arteries, suggesting that it was also partially due to PKC-dependent Ca^{2+} sensitization. It was also reported that application of $10\ \mu\text{M}$ H_2O_2 increased store-operated Ca^{2+} influx in cultured rat PASMCM by stimulating the association of STIM1 with both Orai1 and TRPC1 [414].

Although there have been relatively few investigations of how H_2O_2 and other oxidants cause vasorelaxation in PA, the mechanisms which have been characterized in these arteries (opening of K_v channels and activation of protein kinase G [337,341,502,506]) are in line with the evidence available from studies carried out in systemic arteries [22,507–510]. Schach et al. [341] also reported that $100\ \mu\text{M}$ diamide inhibited store-operated Ca^{2+} entry in rat PA, although a later and more detailed investigation reported that $10\ \mu\text{M}$ H_2O_2 had the opposite effect in rat PASMCM [414].

Conclusions drawn on the basis of the effects of extracellular H_2O_2 application on PASMCM contraction or $[\text{Ca}^{2+}]_{\text{cyt}}$ should be interpreted cautiously, since cellular H_2O_2 signaling is highly localized [103,263,511], whereas the external application of H_2O_2 to PASMCM increases its global intracellular concentration [264]. A further complication is that adding exogenous H_2O_2 (and diamide) causes oxidation of the mitochondrial matrix [220,279], whereas hypoxia apparently has the opposite effect in PASMCM (Section 7.4). Taken together with the evidence that H_2O_2 and other oxidants can activate both constricting and dilating mechanisms in PA, and that which of these dominates depends on the pre-existing level (or type [504]) of contraction, these considerations imply that whereas responses to exogenous oxidants can provide meaningful information about the redox regulation of specific contractile mechanisms if these are studied in isolation, the effect of adding exogenous oxidants on vascular tone or $[\text{Ca}^{2+}]_{\text{cyt}}$ in PASMCM under normoxic or hypoxic conditions may not necessarily reflect the role of endogenous ROS production in HPV.

7.4. Effects of Hypoxia on PASMCM ROS Levels

Numerous investigations, many of which are described in Sections 5.1 and 5.2, examined the effects of hypoxia on ROS in PASMCM, PA, and lung tissue. As set forth in previous reviews [3,323], studies which utilized chemical ROS probes such as lucigenin and DCF presented mixed results: some laboratories found that hypoxia decreased ROS, and others found the opposite effect. Possible explanations for these discordant findings include the use of different preparations, levels of hypoxia, and ROS indicators. Indicator signaling arising from extracellular ROS, which could far outweigh that emanating from cells [48], may have obscured effects on cellular ROS in studies carried out in perfused lungs and isolated arteries [3]. Evidence that hypoxia causes a fall in ROS production by pulmonary artery endothelial cells [512,513] and isolated lung fibroblasts [423] also

suggests that the effects of hypoxia on overall [ROS] production by lung tissue may not accurately reflect what is happening in PASMC.

The chemical indicators used in these studies, most of which were said to detect superoxide levels, have also been criticized on the basis that they are in reality non-specific [514,515], although with the proper controls, they can provide useful qualitative information about cellular oxidant levels [99,386]. However, there are additional difficulties in interpreting the signals arising from these probes, which may arise specifically when they are used to measure the effects of changes in PO_2 [99]. For example, DCFH (and DCFH-DA, its more cell permeable derivative), lucigenin, luminol, and L-012 are converted by cellular oxidants (DCFH, luminol, L-012) or enzyme reductases (lucigenin) to intermediates which generate a signal when they react with superoxide. Unfortunately, these intermediates also react with O_2 to form superoxide, a process termed 'redox cycling' [515–517]. DCFH itself has only a low reactivity to superoxide, H_2O_2 and peroxynitrite, and is instead primarily oxidized by iron-dependent reactions with free radicals such as carbonate and nitrogen dioxide, which can be generated by the reaction of hydroxyl radical with bicarbonate and nitrite, respectively [515]. Similarly, although DHE does react with superoxide to form the diagnostic fluorescent marker 2-hydroxyethidium, it also reacts with one-electron oxidants in an iron-dependent manner, adding a spurious component to the diagnostic signal by generating ethidium, which is also fluorescent [516]. Lucigenin has been reported to accumulate in the mitochondria and primarily detect intra-mitochondrial superoxide [518,519]. An additional confounding factor with lucigenin is that its redox cycling is promoted by NADH [520], the concentration of which is increased by hypoxia [351]. Apart from the lack of selectivity of both DHE and MitoSox for superoxide due to their iron-dependent reaction with non-specific oxidation to ethidium, [516], their fluorescence is also affected by their distribution into the mitochondria, which is sensitive to $\Delta\psi_m$ [386,521].

Several of these investigations also utilized the Amplex Red/horseradish peroxidase (HRP) assay, which measures extracellular $[H_2O_2]$ and can be used to detect changes in cellular $[H_2O_2]$, since these lead to its altered leakage out of cells [522,523]. Amplex Red is considered to provide a high-quality, specific, and very sensitive measure of overall cellular H_2O_2 production [359]. It is therefore noteworthy that three studies using this assay reported that hypoxia decreased ROS production by PASMC [156,294,457]. However, it is not clear whether this was due to an effect on mitochondrial ROS production, since the activity of Nox is probably curtailed at the PO_2 levels used in most studies of HPV [68], and has been shown to make a substantial contribution to the overall release of H_2O_2 from a range of cell types [524].

In view of the shortcomings of the conventional intracellular ROS indicators, several laboratories have examined the effect of hypoxia in PASMC using genetically encoded ROS or redox indicators such as Ro-GFP [525], which responds to the ambient thiol/disulfide equilibrium, and HyPer [296], which reacts with H_2O_2 . In addition to responding to specific ROS or redox changes, these can be targeted to specified cell compartments and do not undergo redox cycling, so are therefore seen as greatly superior to chemical ROS indicators [367,386,526,527]. The results of these investigations are summarized in Table 5.

Table 5. Effects of hypoxia on cell H₂O₂, indices of redox balance, and mitochondrial membrane potential.

Preparation	Level of Hypoxia	Indicator/ Compartment	Indicator Responds to...	Effect of Hypoxia	Additional Comments	Ref.
Cultured rat PASM	1.5% O ₂	HSP-FRET, mainly in cytoplasm	H ₂ O ₂	↑ oxidation	Hypoxia caused gradual ↑ in HSP-FRET and Ca ²⁺ signals with similar timecourses	[361]
Cultured rat PASM	1.5% O ₂	Mito-Ro-GFP mitochondria	GSSG/GSH	↓ oxidation		[366]
		IMS—Ro-GFP IMS		↑ oxidation		
		Cyto—Ro-GFP cytoplasm		↑ oxidation	Hypoxia-induced oxidation of cyto-Ro-GFP abolished by overexpression of cytosolic catalase	
Cultured rat renal artery ¹ SMC	1.5% O ₂	Mito-Ro-GFP mitochondria	GSSG/GSH	↓ oxidation		[366]
		IMS—Ro-GFP IMS		↑ oxidation		
		Cyto—Ro-GFP cytoplasm		↑ oxidation		
Cultured mouse lung slice	1.5% O ₂	Cyto—Ro-GFP cytoplasm	GSSG/GSH	↑ oxidation	Hypoxia-induced oxidation of cyto-Ro-GFP strongly suppressed in lung slices overexpressing cytosolic catalase	[368]
Cultured mouse PASM	PO ₂ = 13 Torr	IMS-targeted Ro-GFP		↑ oxidation		[139]
Cultured mouse PASM	1.5% O ₂	Mito-Ro-GFP mitochondria	GSSG/GSH	↓ oxidation		[140]
		² IMS—Ro-GFP		↑ oxidation	Hypoxia-induced oxidation of Cyto- and IMS-Ro-GFP strongly suppressed in RISP-depleted PASM	
		Cyto—Ro-GFP cytoplasm		↑ oxidation		
Isolated mouse PA	1.5% O ₂	Cyto—Ro-GFP cytoplasm	GSSG/GSH	↑ oxidation		[140]
		Mito-Ro-GFP mitochondria		↓ oxidation		
Rat renal artery SMC	2.5% O ₂	Cyto—Ro-GFP cytoplasm	GSSG/GSH	↑ oxidation	Hypoxia-induced oxidation of Cyto-Ro-GFP strongly suppressed in RISP-depleted renal artery SMC	[140]
Perfused mouse lung	1% O ₂	HyPer Cyto	H ₂ O ₂	↑ oxidation		[423]
Wildtype mouse PASM	1% O ₂	HyPer expressed in cytoplasm	H ₂ O ₂	↑ oxidation		[142]
Cox4i2 knock-out mouse PASM	1% O ₂	HyPer expressed in cytoplasm	H ₂ O ₂	↓ oxidation		[142]
Rat PASM	20 Torr	pHyper-dCyto	H ₂ O ₂	↓ oxidation		[137]
		pHyper-dMito	H ₂ O ₂	↓ oxidation		
Rat renal artery SMC	20 Torr	pHyper-dCyto	H ₂ O ₂	↑ oxidation		[137]
		pHyper-dMito		No effect		
Isolated pig PA 100–150mm diameter	4% O ₂ PO ₂ = 29 TorrO ₂	Electron spin resonance with DMPO	Superoxide, peroxynitrite; see [528]	↑ in three of six hypoxic PA but in zero of six normoxic PA	Effect of hypoxia was non-significant. Use of nitron spin probes such as DMPO has been strongly criticized	[358]

Mouse PASMCMC	1% O ₂	Electron spin resonance with CMH	Superoxide, peroxynitrite; see [529]	↑ ROS	Signal was due to superoxide, defined by its abolition by SOD	[142]
Mouse PASMCMC	1,5,10, 15% O ₂	Electron spin resonance with CMH	Superoxide, peroxynitrite	↑ ROS in 1% O ₂ but not in ≥ 5% O ₂	Signal was due to superoxide, defined by its abolition by SOD	[423]
Cultured wildtype mouse PASMCMC	PO ₂ = 15–22 Torr	Electron spin resonance with CMH	Superoxide, peroxynitrite	↑ ROS	Signal was due to superoxide, defined by its abolition by SOD	[141]
Cultured AOX-expressing mouse PASMCMC	PO ₂ = 15–22 Torr	Electron spin resonance with CMH	Superoxide, peroxynitrite	↔ ROS	AOX expression abolished the ↑ ROS observed in wildtype controls	[141]
Cultured rat microvascular PA	1.5% O ₂	GSSG/GSH-412 kit from Oxis Health Products	Mainly cytoplasmic [GSSG]/[GSH]	↑ GSSG/GSH over 2 hrs		[361]
Isolated rat PA	0% O ₂ PO ₂ = 15–20 Torr	GSH/GSSG See [530]	Mainly cytoplasmic GSSG: GSH redox state	↓ GSH/GSSG redox potential (= oxidation) over 45 min	GSH/GSSG redox potential fell from −176 to −158	[18]
Isolated rat aorta	0% O ₂ PO ₂ = 15–20 Torr	GSH/GSSG See [530]	Mainly cytoplasmic GSSG: GSH redox state	↔ GSH/GSSG redox potential (=oxidation) over 45 min	GSH/GSSG redox potential under normoxic conditions was not significantly different in PA and aorta	[18]
1st passage cultured rat PASMCMC	PO ₂ = 40 Torr	JC-1, ³ TMRM	ΔΨ _m (E _m of mitochondria)	ΔΨ _m increased by hypoxia	ΔΨ _m was more depolarized in PASMCMC compared to renal artery SMC under normoxic conditions	[156]
1st passage cultured rat renal PASMCMC	PO ₂ = 40 Torr	JC-1, TMRM	ΔΨ _m	ΔΨ _m decreased by hypoxia		[156]
Primary cultured human ductus arteriosus cells	PO ₂ = 40 Torr	JC-1, TMRM	ΔΨ _m	ΔΨ _m increased by ↑ PO ₂ from 45 to 100 Torr		[156]
Cultured rabbit precapillary PASMCMC	PO ₂ = 8, 23 or 38 Torr	JC-1	ΔΨ _m	↑ ΔΨ _m with PO ₂ of 8 and 23 but not 38 Torr	Hyperpolarization of ΔΨ _m due to PO ₂ of 8 Torr became significant after ~2 min	[155]
Cultured rabbit aorta SMC	PO ₂ = 8 Torr	JC-1	ΔΨ _m	↑ ΔΨ _m		[155]
Cultured rat renal artery SMCs	PO ₂ = 8 Torr	JC-1	ΔΨ _m	↔ ΔΨ _m		[155]
Isolated mouse precapillary PASMCMCs	1% O ₂	JC-1	ΔΨ _m	↑ ΔΨ _m	Progressive hyperpolarization over 8 min period of hypoxia	[142]
Isolated mouse precapillary PASMCMCs	PO ₂ = 15–22 Torr	JC-1	ΔΨ _m	↑ ΔΨ _m	Hyperpolarization of ΔΨ _m was not observed in PASMCMC from Cox4i2 knockout mice	[141]
Isolated mouse PASMCMC from 2nd- and 3rd-order PA	6% O ₂	Rhodamine 123	ΔΨ _m	↑ ΔΨ _m	ΔΨ _m hyperpolarized rapidly after 1–2 min of hypoxia	[26]

Abbreviations: ¹ SMC—smooth muscle cells; ² IMS = intramembrane space (of mitochondria); ³ TMRM = Tetramethylrhodamine methyl ester.

Dunham-Snary et al. [137], using HyPer in cultured rat PASMCMC, found that hypoxia decreased [H₂O₂] in both the mitochondrial matrix and the cytoplasm. They suggested

that since a redox mediator produced in the mitochondria must enter the cytoplasm to modulate contractile effector proteins, the fact that they detected similar effects of hypoxia on $[H_2O_2]$ in both compartments supported the validity of their findings. They also speculated that the hypoxia-induced reduction of Ndufs2 thiols which they had observed (Section 5.1) was responsible for inhibiting its production of ROS, implying that the reduction of the mitochondrial matrix might be a key factor in causing HPV.

In contrast, all of the other studies using genetically encoded indicators, including one which used Hyper in mouse PASM C [142], reported that hypoxia increased $[H_2O_2]$ or oxidized the thiol/disulfide equilibrium in the cytoplasm of mouse and rat PASM C, although, in agreement with Dunham-Snary and colleagues, the mitochondrial matrix became more reduced. Several papers from Schumacker's laboratory also reported that hypoxia increased redox tone in the mitochondrial IMS, consistent with their hypothesis that the increased ROS production by Complex III triggered by hypoxia is directed into the IMS, from which H_2O_2 can enter the cytoplasm. Korde et al. [139] reported a similar finding.

Likewise, several papers from Sommer and colleagues reported that hypoxia increased superoxide production in cultured mouse PASM C. Although the spin probe they used (1-hydroxy-3-methoxycarbonyl-2,2,5, 5-tetramethylpyrrolidine; CMH) detects both mitochondrial and extra-mitochondrial superoxide [531], since they (and others) consistently found that the redox poise in the mitochondrial matrix tone was reduced during hypoxia, it seems likely that the increase in superoxide detected by ESR was cytoplasmic. Importantly, Sommer et al. [141] also demonstrated using ESR that the hypoxia-induced increase in superoxide production was absent in PASM C from mice expressing AOX, consistent with evidence that mitochondrial ROS production associated with an increased reduction of the ubiquinone pool is depressed in these animals [388,532].

The generation of ROS by both complexes 1 and 3 is strongly influenced by $\Delta\Psi_m$ [67,352,533]. As shown in Table 5, hypoxia has been observed to increase $\Delta\Psi_m$ in PASM C in each of the three studies in which it has been measured. This is consistent with an increase in mitochondrial ROS production.

In summary, experiments utilizing genetically encoded ROS indicators and ESR have, with only one exception [137], supported the notion that hypoxia increases ROS production and cytoplasmic oxidant levels in PASM C. The hypoxia-induced hyperpolarization of $\Delta\Psi_m$ which has been reported consistently also provides indirect support for this idea.

7.5. Concluding Remarks: The Redox and Mitochondrial Models

Overall, compared to the mitochondrial ROS hypothesis, the evidence supporting the Redox Theory is less extensive. It is also generally older, and partly as a consequence of this, is based to a larger extent on results arising from experimental approaches (the use of ETC blockers as hypoxia surrogates, the knockout of mitochondrial complex subunits to characterize their involvement in O_2 sensing) which have been criticized as being unreliable [523,534], or supportive but not definitive (the use of isolated mitochondria, pro- and antioxidants, and conventional small molecule ROS indicators [48]).

These considerations also apply to much of the evidence supporting the mitochondrial ROS hypothesis and may well account for some of the discrepancies between the results reported by different laboratories. However, placing this type of evidence to the side, the observations that remain predominantly favor the Mitochondrial ROS hypothesis. These include, most notably, the inhibition of HPV by S3QEL2 [142] and by the expression of AOX [141], as well of the effects of hypoxia on redox state of the cytoplasm using newer genetically coded indicators and electron spin resonance measurements shown in Table 5.

The Mitochondrial ROS hypothesis is also indirectly supported by evidence that Cox4i2, proposed to be crucial for increased hypoxia-induced ROS production during HPV (Section 5.2.2) [142], plays a similar role in the carotid body. In contrast, although the implications of Cox4i2 for the Redox Theory have not been discussed by its proponents, the evidence that its high expression promotes O₂ sensing in both PASMC and CBCC seems at variance with the idea that hypoxia decreases ROS production, since it is widely accepted that reduction of the CoQ pool is a key stimulus for ROS production. However, whilst the comparatively high expression of Cox4i2 in the two most well-characterized types of cells seen to respond acutely to hypoxia suggests that it could be playing an important role O₂-sensing, possibly by promoting mitochondrial ROS production, it is also the case that hypoxia, at levels similar to those generally used by experimenters to evoke HPV and CBCC activation, induces rapid increases in ROS in many other types of native cells/tissues or cell lines which are not known to express high levels of Cox4i2 [156,190,347,348,360,535–544]. Therefore, it remains to be established that the level of expression of Cox4i2 is a pivotal determinant of hypoxia-induced ROS production.

Despite the evidence for the mitochondrial ROS hypothesis being more compelling at this point, questions remain. For example, the observation that HPV recorded in isolated PA from AOX mice was restored if the arteries were incubated in PSS containing 20 mM K⁺ [141] appears to sit uneasily with evidence that an increase in ROS production by complex III is causing HPV by causing Ca²⁺ release from the sarcoplasmic reticulum and Ca²⁺ sensitization (see Sections 5.2.1 and 5.2.6). Since 20 mM K⁺, which promotes vascular smooth muscle cell contraction by causing depolarization, was able to replace the effect of ROS production by complex III which was absent in the AOX mice, the implication is that ROS produced by complex III contribute to HPV by causing membrane depolarization. If so, the contractile effects over and above those caused by depolarization (i.e., Ca²⁺ sensitization and release from the sarcoplasmic reticulum) which are evoked by hypoxia are not due to ROS production by complex III. If nothing else, these apparently contradictory observations highlight the fact that the HPV effector mechanisms linked to an increase in ROS remain imperfectly understood.

Another unresolved aspect of the mitochondrial ROS hypothesis is that the nature of the oxidizing signal responsible for HPV remains to be convincingly identified. Although H₂O₂ is generally seen as being the most important oxidizing species responsible for cell signaling, the evidence that this is the case for HPV is decidedly mixed. Its specific involvement in HPV is supported by evidence from three investigations showing that over-expression of catalase [361,364,368] prevented the hypoxia-induced rise in [Ca²⁺]_{cyt} in PASMC, and one showing that hypoxia oxidized HyPer [142]. On the other hand, hypoxia was seen to cause a fall in the cytoplasmic concentration or cellular production of H₂O₂ in four other studies using AmplexRed/HRP or HyPer [156,294,303,457], and the mechanisms by which H₂O₂ caused contraction when applied at a concentration which probably caused a quasi-physiological intracellular signal did not closely match those thought to be responsible for HPV [425]. Also arguing against H₂O₂ as the HPV mediator is the carefully conducted study by Liu and colleagues showing that the application of SOD, which they verified led to its penetration into PASMC, strongly suppressed HPV in pig PA [358].

The question of the redox mediator responsible for HPV is particularly relevant in light of recent experiments carried out in HEK 293 cells by Sen et al., which suggest that hypoxia may exert opposite effects on two types of cell oxidant species [523]. This investigation aimed to determine whether the opposing effects of hypoxia on ROS production observed in PASMC and many other types of cells were genuine or might, for example, be due to the use of different indicators with non-identical responses to particular oxidant species.

The effects of hypoxia (1% O₂) in HEK293 cells were recorded using the Amplex UltraRed + SOD assay, both in the presence of HRP to record H₂O₂, and in the absence of HRP, in which case the assay detects other oxidant species instead of H₂O₂. They also used DCF to measure general cell oxidation. They found that hypoxia decreased H₂O₂ release. Importantly, this was associated with a fall in H₂O₂ production by complexes 1 and 3, and also by Nox, plus other sources which were not characterized. This supports the idea that the decrease in H₂O₂ production was caused by a deficiency of O₂ rather than a decrease in the availability of electrons at any specific site. Hypoxia also decreased DCF oxidation. On the other hand, hypoxia increased the release from the cells of (an)other oxidant(s), as sensed by Amplex UltraRed in the absence of HRP. Whereas they did not identify these other oxidant(s), work by Kalyanaranam's laboratory showed that the oxidizing species involved is probably peroxynitrite [516], or, more likely, products of its decomposition [545]. Notably, peroxynitrite (or its products) can also increase the fluorescence of AmplexRed + HRP [545]. Even so, this reaction occurs much more slowly than that with H₂O₂, suggesting that the two assays were genuinely picking up opposing effects of hypoxia on H₂O₂ vs. (an)other oxidant species.

These results, therefore, imply that hypoxia alters the balance between cellular production of H₂O₂ and other oxidizing species, thereby providing a potential explanation for the apparent indicator-dependency of the effect of hypoxia reported in both the HPV literature and by studies carried out in other types of cells and tissues.

If the unidentified oxidant was peroxynitrite, the observations of Sen et al. could potentially be explained by the hypothesis developed by Poyton and colleagues [427] that hypoxia increases the production of both NO and superoxide by the mitochondria, generating a cytoplasmic peroxynitrite signal capable of orchestrating a cellular response. One implication of this model is that since the rate at which NO and SOD react with superoxide is similar [546], they would compete for superoxide released into the IMS by complex III. Thus, an increase in [NO] evoked by hypoxia, whether originating in the mitochondria or elsewhere in the cell [174], could enhance the production of peroxynitrite while at the same time diminishing that of hydrogen peroxide. Furthermore, since Ro-GFP, ESR, and HSP-FRET can sense peroxynitrite directly or indirectly through its effects on the ambient redox state, whereas HyPer responds specifically to H₂O₂, some of the apparently contradictory observations which set the Mitochondrial ROS hypothesis and the Redox Theory apart could in fact represent two facets of the same process.

In any case, we suggest that the role of mitochondrial ROS production in O₂ sensing during HPV remains sufficiently uncertain so as to warrant further studies. For example, the observation by Sommer et al. [142] that S3QEL2 inhibits HPV, which we believe is currently the most convincing evidence that it requires ROS production at complex III, has not been confirmed independently, and would be strengthened by corresponding evidence that S3QEL2 also suppresses hypoxia-induced cytoplasmic oxidation. It would also be useful if the current evidence for cytoplasmic oxidation could be broadened, for example, by using newer and more sensitive indicators such as Ro-GFP2-Tsa2DeltaCRm, which is based on a modified form of peroxiredoxin [511]. The possible involvement of NO and peroxynitrite in HPV appears to be another area worthy of investigation.

In addition, an analysis of the effects of hypoxia on reactive cysteines in putative ROS targets such as IP₃R, RyR, and Kv1.5 would confirm that any alteration in cytoplasmic redox state is not merely an epiphenomenon. This approach could be implemented using isolated PA or perfused lung, circumventing the use of cultured cells and avoiding attendant complications arising from their altered phenotype and isolation from their normal environment.

8. The Role of the Pentose Phosphate Pathway and the Withdrawal of Normoxic Vasodilation Maintained by Nox4, H₂O₂, and Protein Kinase G in HPV

Michael Wolin and colleagues developed a complex scheme for O₂ sensing in HPV which resembles the Redox Theory in that HPV is proposed to be due, at least in part, to hypoxia-induced withdrawal of ongoing ROS-dependent normoxic vasodilation. However, whereas according to the Redox Theory this vasodilation is caused by mitochondrial ROS production and a resulting activation of K_v channels, the model developed by Wolin's laboratory envisions that normoxic vasodilation is maintained mainly by H₂O₂-mediated activation of the soluble guanylate cyclase (sGC)/protein kinase G (PKG) pathway, and that Nox4 rather than the mitochondrial ETC is responsible for H₂O₂ production.

Additional studies by Sachin Gupte, mostly carried out in collaboration with Wolin, presented evidence that hypoxia also regulates arterial tone by altering the activity of the pentose phosphate pathway (PPP), thereby leading to changes in the cellular [NADPH]/[NADP⁺] ratio. This would enable H₂O₂-independent regulation of redox networks, leading to effects on thiol oxidation which alter the function of proteins involved in controlling vascular tone. In PA, for example, hypoxia was shown to stimulate the PPP, thereby increasing the [NADPH]/[NADP⁺] ratio and as a result decreasing the activity of PKG1 α , which is redox-sensitive [22]. Somewhat paradoxically, several of these investigations found that activation of the PPP by hypoxia is due to a rise in cytoplasmic [H₂O₂], which acts through PKC δ to stimulate glucose-6-phosphate dehydrogenase (G-6-PD), the initial and rate-limiting step in the PPP. This rise in H₂O₂ was also proposed to stimulate contraction through both Ca²⁺-dependent and -independent pathways.

Although several of these studies investigated HPV in isolated perfused lungs or pulmonary artery rings from rats [447,547] or mice [483], most of this work was carried out using isolated bovine pulmonary artery (BPA) or lung homogenates, sometimes with comparative experiments in bovine coronary arteries (BCA), which relax to hypoxia. Importantly, rotenone, which has universally been found to block HPV in other species (Table 2), had no effect on HPV in BPA [458]. This would imply that mitochondrial ROS production plays no direct role in O₂ sensing in BPA, at least under the conditions used in these studies.

In most of these studies, hypoxia was imposed using PSS gassed with 5%CO₂/95%N₂, resulting in a PO₂ of 8–10 Torr. For studies of HPV, BPA rings were denuded of their endothelium to prevent hypoxic effects which might be mediated by NO and other endothelium-derived vasoactive factors, and were slightly pre-constricted, generally with 20–30 mM K⁺ but in some studies with U46619 or 5-HT, in order to enhance HPV. The authors generally adjusted the degree of pre-tone stimulus to maintain a similar amplitude of pre-constriction in the presence and absence of the interventions (e.g., blockers) which were being used to characterize mechanisms engaged by hypoxia or ROS.

8.1. Early Studies: HPV as the Loss of Tonic H₂O₂ and sGC-Mediated Vasorelaxation

A series of papers by Wolin's laboratory published over an approximately 10-year period starting in 1987 developed the concept that PA normally generate a basal level of superoxide (and therefore H₂O₂) due to the activity of an NAD(P)H oxidase [396,459,487,493,548–552]. This causes an ongoing stimulation of soluble guanylate cyclase (sGC) and therefore protein kinase G, creating a tonic vasodilating influence. The stimulation of sGC by H₂O₂ was proposed to be mediated by compound 1, a form of catalase which exists during its metabolism of H₂O₂ [493,548]. Vasodilation due to this mechanism is suppressed by hypoxia because there is less O₂ available for the formation of superoxide/H₂O₂, resulting in contraction (i.e., HPV). The development of this scheme is described

in a review by Wolin et al. [553]. The oxidase responsible for superoxide production in PA was initially seen as utilizing NADH. However, Mohazzab & Wolin [551] pointed out that its properties resembled those of the NADPH-dependent oxidoreductase which had been detected in neutrophils [554] and had been suggested to play a role in O₂ sensing in the carotid body [555]. It was eventually concluded [553] that it corresponded to the phagocytic NADPH oxidase and might be the same as the vascular oxidoreductase which others had shown utilized both NADH and NADPH to generate superoxide [556]. It soon became evident that there was a group of such oxidoreductases, christened the Nox family [557], and that they preferred to use NADPH rather than NAD as a substrate for superoxide production. In later work Wolin and colleagues concentrated on defining the role of NADPH rather than NADH in mediating the effects of hypoxia, stating that the [NADPH] in cells is likely to be 10–20 µM whereas [NADH] is ~1 µM [558].

Most of the subsequent investigations of HPV by this group focused on two interrelated themes. As described in Section 8.2, a series of papers published between 1999 and 2010, spearheaded by Sachin Gupte, examined how hypoxia-induced regulation of the PPP influences the redox state of the NADPH/NADP⁺ couple in such a way as to cause constriction of PA but relaxation of coronary arteries. Other work, published between 2010 and 2014, further examined how hypoxia, H₂O₂ and NADPH, acting through cellular redox networks, modulate PA tone by regulating the activity of the sGC/PKG axis (Section 8.3.2). In addition, Wolin's laboratory carried out pioneering experiments to examine the important possibility that HPV is influenced by the presence of extracellular H₂O₂ [482,483]

8.2. PPP Activity as a Determinant of the Effects of Hypoxia on Vascular Tone

An initial investigation of the role of the PPP in regulating vascular contraction [559] was based on the hypothesis that oxidation of the heme iron on sGC would inhibit its activity by preventing the binding of NO, and that NAD(P)H, by reducing the heme iron, might restore the responsiveness of sGC to NO, thereby promoting vasorelaxation. The results of the study provided support for this hypothesis, implying that the activity of the PPP, which generates cellular NADPH, could potentially regulate vascular tone of BPA via this mechanism.

A subsequent paper then considered two possible scenarios by which the activity of the PPP might influence HPV [547]. If NAD(P)H was indeed acting to augment the stimulation of sGC by NO, PPP activation should suppress HPV. Alternatively, based on evidence that NADP⁺ and GSSG activated the K⁺ current in PASMC [336], PPP-mediated reduction of NADP⁺ and/or GSSG might promote HPV by closing K⁺ channels and causing depolarization-induced Ca²⁺ influx. Their experiments showed that antagonists of glucose-6-phosphatase dehydrogenase (G-6-PD), which mediates the rate-limiting step of the PPP, suppressed HPV in isolated perfused rat lungs and relaxed contractions evoked by 30 mM K⁺ in both PA and aorta. These results supported the second hypothesis, implying that hypoxia-induced activation of the PPP could promote HPV by reducing the cytoplasmic NADPH: NAD(P)⁺ couple and thereby inhibiting K_v channel opening.

Gupte et al. [560] similarly observed that pharmacological blockade of G-6-PD caused relaxation of bovine coronary arteries (BCA) and that this was accompanied by a fall in the tissue contents of NADPH and GSH, whereas total NAD(P)⁺ and GSSG increased. The reducing agent DTT attenuated this relaxation, suggesting that it was due to an oxidizing effect of some kind. However, they also found that superoxide and H₂O₂ generation by BCA fell during PPP blockade, implying that this oxidation was not due to an increase in cellular [ROS]. Moreover, ebselen, which would be expected to diminish the concentrations of both ROS, did not suppress relaxation. They proposed that the fall in ROS production caused by PPP blockade occurred because there was less NADPH available to be used to produce superoxide. Although this study was in coronary rather than pulmonary

arteries and did not look at the effects of hypoxia, it was important for their subsequent studies of HPV because it showed that the PPP, presumably by producing NADPH, was able to influence vascular tone through ROS- and sGC-independent mechanisms, and that regulation of Ca^{2+} influx and release was more important in this respect than that of Kv channels.

Wolin's laboratory had previously shown [561] that hypoxia also relaxed BCA via a mechanism which appeared to be independent of ROS, since it was not affected by NBT (an antioxidant) and DPI (blocks Nox and the ETC). This suggested that the inhibition of the PPP might also be causing relaxation through the same ROS-independent mechanism. In accordance with this possibility, Gupte & Wolin [562] reported that hypoxia increased the $[\text{GSSG}]/[\text{GSH}]$ and $[\text{NADP}^+]/[\text{NADPH}]$ ratios in these arteries. These effects were associated with a decreased level of glucose-6-phosphate, which is produced by the first step of the PPP, suggesting that hypoxia was inhibiting the PPP. As with PPP blockade, the relaxation to hypoxia was attenuated by DTT pre-treatment. Interestingly, application to BCA of pyruvate (10 mM), which was used to promote the flux of glucose through the PPP by inhibiting phosphofructokinase [563], increased the arterial content of glucose-6-phosphate and NAD(P)H under both normoxic and hypoxic conditions. Pyruvate also prevented the increase in arterial GSSG and the vasorelaxation evoked by hypoxia. Based on these findings, the authors proposed that, due to a relative lack of G-6-PD in BCA, hypoxia causes a 'metabolic stress' leading to a reduced flux of glucose through the PPP. This causes a rise in the cellular $[\text{NADP}^+]/[\text{NADPH}]$ and $[\text{GSSG}]/[\text{GSH}]$ ratios, which act through redox networks to cause vasorelaxation by oxidizing thiol switches on a number of proteins controlling contraction (e.g., SERCA) [57].

Gupte et al. [558] examined the relationship between G-6-PD and cellular superoxide levels (assessed using 5 μM lucigenin) in BPA and BCA. They reported that mRNA expression of Nox1 was not present in either artery. On the other hand, Nox2 and Nox4 mRNA and protein were similarly expressed in the two types of arteries, although superoxide levels were 40–80% higher in BPA. Both the G-6-PD blocker 6-aminonicotinamide (6-AN) and the putative Nox antagonist apocynin decreased the lucigenin signal more in BPA than in BCA, suggesting that the greater level of superoxide production in BPA was due to a higher activity of Nox in these arteries. This in turn was driven by a relatively higher rate of NADPH formation by the PPP. This idea was supported by their observation that the level of NADPH and the expression and activity of G-6-PD was higher in BPA than in BCA. We suggest, however, that this conclusion should be interpreted with caution, since the $[\text{NADPH}]/[\text{NADP}^+]$ ratio was higher in BCA, suggesting that the PPP might be more active in these arteries.

Gupte et al. [447] then investigated the involvement of the PPP and NADPH in HPV recorded in perfused rat lungs. They found that acute hypoxia caused marked increases in the amounts of glucose-6-phosphate and NADPH in the lung. Blockers of G-6-PD abolished the hypoxia-induced rise in NADPH and also greatly reduced the amplitude of HPV. Similarly, hypoxia increased the $[\text{NADPH}]/[\text{NADP}^+]$ ratio by about 75% in isolated lobar PA, and this effect, along with HPV, was suppressed by the G-6-PD antagonists. In additional experiments, sGC activity was measured in homogenates prepared from isolated PA which had been subjected to hypoxia in the presence or absence of G-6-PD blockers or the sGC antagonist 1H-[1,2,4]oxadiazolo [4,3-a] quinoxalin-1-one (ODQ). It was found that hypoxia strongly increased the activity of sGC. This effect was also prevented by G-6-PD blockade, suggesting it was due to activation of the PPP. ODQ also prevented the increased sGC activity, but, in contrast to PPP blockade, enhanced HPV.

Based on these results, the authors proposed that hypoxia activates the PPP in PA, leading to an increased synthesis of NADPH, which is responsible for HPV. At the same time, the increased $[\text{NADPH}]$ also stimulates sGC (see also [559,564]), creating a

countervailing vasodilating influence which explained why HPV was potentiated by ODQ. However, this was outweighed by the pro-contraction response, which they speculated was due to Kv channel inhibition.

In summary, the results of these studies suggested that hypoxia increases and decreases NADPH production by the PPP in BPA and BCA, respectively, and that this difference is responsible for its opposite effects on vascular tone in these arteries. According to the model developed by the authors to explain their observations [565,566], hypoxia *inhibits* the PPP in BCA, leading to vasorelaxation consequent on a fall in cellular NADPH and a resulting oxidation of cellular thiol switches (e.g., on Kv channels and Ca²⁺ channels or pumps). Although ROS production decreases due to the fall in PO₂ and/or the suppression of the PPP, their evidence suggested that this does not exert an important effect on vascular tone.

In contrast, PPP activity in BPA is higher under basal conditions and is *increased* by hypoxia. This causes hypoxia-induced contraction (HPV) through two mechanisms. Firstly, the PPP generates sufficient NADPH under normoxic conditions to create an ongoing vasodilating influence dependent on superoxide/H₂O₂ produced by Nox. This vasodilating influence is greater in BPA than in BCA because the former has a higher expression and activity of G-6-PD and therefore has higher basal NADPH levels. Hypoxia removes this ongoing vasodilation by attenuating ROS production, causing contraction (i.e., HPV). Secondly, PPP activation by hypoxia leads to increased NADPH production, which promotes force development by causing the reduction of GSSG and/or protein thiol switches in such a way as to suppress relaxation or enhance contraction.

Gupte et al. [446] further investigated the role of the PPP in triggering HPV in BPA. Contraction mechanisms were examined in intact endothelium-denuded arteries and those which the plasma membrane had been permeabilized with α -toxin from *staphylococcus aureus* to enable the study of Ca²⁺ sensitization. They evaluated the role of G-6-PD in HPV by using PA which had been cultured with anti-G-6-PD or scrambled siRNA for two days. They also compared the responses to hypoxia in a mouse model in which expression of G-6-PD was halved due to a mutation in the 5' promoter region of its gene, and in mice that were heterozygous for this mutation and had normal levels of G-6-PD expression. They measured G-6-PD activity and the levels of pyridine nucleotides and other metabolic intermediates in homogenates made from frozen BPA.

The authors observed that the activation of the PPP by hypoxia previously detected in rat lung also occurred in BPA. This was shown by their observations that hypoxia increased the activity of G-6-PD as well as the [NADPH]/[NADP⁺] ratio and the NADPH content of these arteries. They also examined the effect of hypoxia on cellular ROS levels using MitoSox and DHE in arteries which were pre-constricted with PSS containing 30 mM K⁺, presumably to mimic the conditions used to study HPV. Hypoxia decreased the MitoSox signal, suggesting a fall in mitochondrial ROS levels, but increased the DHE signal, which they ascribed to cytoplasmic ROS. In light of previous evidence that the activity of G-6-PD is stimulated by H₂O₂, [567] they speculated that an increase in cytoplasmic ROS levels caused by hypoxia was responsible for activating the PPP, a concept which was subsequently supported by the subsequent observation that H₂O₂ could stimulate G-6-PD in BCA through PKC δ [568]. In addition, they found that hypoxia decreased the ATP/ADP ratio and increased the lactate/pyruvate ratio, indicating that glycolysis and oxidative phosphorylation were inhibited. These effects, plus a hypoxia-induced increase in glucose uptake [351], would be expected to increase the intracellular glucose concentration and shunt glucose into the PPP, further increasing its activity.

Curiously, however, they also found that hypoxia strongly increased the NAD⁺/NADH ratio. This effect is the opposite of what would be expected when NADH consumption by the ETC is inhibited, and runs counter to the effects of hypoxia on this

ratio observed by others in PASMC [351] and CBCC [187]. The implications of this apparent discrepancy were not discussed.

Importantly, they observed that decreasing the expression of G-6-PD strongly suppressed HPV. Furthermore, hypoxia increased contraction by stimulating both Ca^{2+} -dependent mechanisms and Ca^{2+} sensitization, and both responses were almost abolished in arteries treated with anti-G-6-PD siRNA. The former effect was studied by measuring hypoxia-induced increases in contractions of intact PA evoked by 30 mM K^+ PSS and 5-HT in the presence of extracellular Ca^{2+} . The latter effect manifested as a leftward shift of the Ca^{2+} vs. contraction curve in permeabilized PA and was also suggested by the increase in MLC_{20} phosphorylation which they detected in intact PA under Ca^{2+} -free conditions. Hypoxia also increased the phosphorylation of CPI-17. The hypoxia-induced phosphorylation of MLC_{20} but not CPI-17 was antagonized by 6-AN, suggesting that PPP-dependent Ca^{2+} sensitization was not due to CPI-17. Instead, the authors suggested that hypoxia was causing Ca^{2+} sensitization via rho kinase, although no direct evidence for this was presented. The involvement of the PPP in O_2 sensing in BPA, proposed by the authors based on these results and those of earlier studies, is illustrated in Figure 9.




Figure 9. The role of the PPP in HPV as described by Gupte et al. [439]. According to their model, hypoxia induces changes in glucose metabolism and cytoplasmic [ROS]. This causes an increased activity of the PPP, which, acting through NADPH, causes a further rise in [ROS]. This activates rho kinase and other pro-contractile pathways, evoking HPV. The figure is based closely on that presented in [446]. See the text for further details.

8.3. Loss of Basal H_2O_2 -Induced Stimulation of sGC and PKG as a Mechanism of HPV

8.3.1. Basal H_2O_2 Production and HPV

Ahmad et al. [482] tested the hypothesis that the removal by hypoxia of an ongoing H_2O_2 -dependent vasorelaxation contributes to HPV by using ebselen (100 μM) to mimic a hypoxia-induced fall in H_2O_2 . They found that the contraction of BPA to 20 mM K^+ , which they viewed as reflecting basal normoxic contractility, was enhanced by ebselen, while HPV was strongly depressed. Treatment of BPA with the CuZn SOD blocker DETCA exerted similar effects. Both ebselen and DETCA greatly decreased normoxic H_2O_2 levels (detected using a luminol/HRP assay). HPV was also markedly depressed in BPA which had been treated for 24 h with CoCl_2 , which increased tissue levels of H_2O_2 , putatively by enhancing the expression of extracellular SOD. This suggested that if produced extracellularly in sufficient quantities, H_2O_2 entry into PASMC would increase to the point where hypoxia was no longer able to depress its intracellular concentration enough to evoke HPV. This idea was supported by the observation that applying 1 mM catalase to remove extracellular H_2O_2 restored HPV in the CoCl_2 -treated arteries.

They then [458] examined the involvement of Nox2 and Nox4 in generating the H_2O_2 responsible for maintaining normoxic vasorelaxation in BPA. They found that pharmacological block or siRNA knockdown of Nox2 attenuated superoxide production but did not affect HPV or the contraction to 25 mM K^+ under normoxic conditions. Similarly, ETC block using rotenone (10 μM) or antimycin (10 μM) decreased superoxide production but had no effect on either type of contraction. In contrast, knockdown of Nox4, which also depressed superoxide production, halved the amplitude of HPV, although it too had no effect on the 25 mM K^+ contraction (i.e., knockdown of Nox4 was not affecting the basal contractility or vascular tone of the arteries). Based on these results, the authors concluded that Nox4 was responsible for generating H_2O_2 and maintaining a vasodilating influence which was removed by hypoxia, thereby causing or contributing to HPV.

However, this scheme would predict that interventions which diminish intracellular $[\text{H}_2\text{O}_2]$ even under normoxic conditions should increase basal contractility, and this conflicted with their finding that siRNA knockdown of Nox4 had no effect on the contraction to 25 mM K^+ . To explain this, they suggested that H_2O_2 produced by Nox4 had both dilating and constricting effects, and that following H_2O_2 depletion the dilating effect was lost immediately, whereas the constricting effect wore off more slowly. The immediate loss of the dilating effect would give rise to a rise in tension (e.g., HPV or the contraction caused by the antioxidant ebselen), whereas the eventual loss of the contractile effect would account for the fact that very prolonged H_2O_2 depletion associated with Nox4 knockout did not affect the response to 25 mM K^+ . They tested this concept by applying 25 mM K^+ before and after either a short- (35 min) or long-term (several hours) treatment with ebselen (again used to mimic the H_2O_2 -lowering effect of hypoxia) and found that short-term ebselen treatment produced the predicted enhancement of the high K^+ response, whereas long-term treatment did not. The authors suggested that the pro-contractile effect of ROS produced by Nox4 might be due to the stimulation of rho kinase.

Together, the results of these two papers [458,482] supported the hypothesis that HPV is due to the loss of vasorelaxation caused by H_2O_2 produced by Nox4.

8.3.2. HPV and Regulation of sGC and PKG by H_2O_2

The proposal by Wolin's laboratory that a fall in H_2O_2 production caused by hypoxia evokes or promotes HPV by inhibiting the activity of the sGC/PKG pathway grew out of their initial observations suggesting that H_2O_2 stimulates sGC through the action of compound C, a catalase intermediate which forms while it is metabolizing H_2O_2 [493,548,569,570]. The link between this mechanism and HPV was based on evidence [548]

that aminotriazole, which inactivates catalase by binding to compound C, caused a contraction and inhibited HPV in BPA. However, it is not clear from the results presented in this paper that the latter effect was statistically significant, and whereas aminotriazole raised basal PAP in perfused rat lung, it did not block HPV in this preparation [552]. Moreover, in perfused rabbit lung, where aminotriazole did block HPV, it caused an equivalent inhibition of the contractile responses to U46619 and angiotensin 2 [486]. The ability of compound C to block sGC appears not to have been confirmed by any other laboratories.

Wolin's laboratory subsequently explored other mechanisms by which hypoxia could regulate the sGC/PKG axis in BPA, reporting that hypoxia greatly enhanced the stimulation of cGMP production evoked by NO donors [571]. Nevertheless, sGC-mediated vasorelaxation was strongly depressed by hypoxia, although the NO-induced vasorelaxation was maintained by an sGC-independent mechanism, which their experiments suggested was the activation of the sarco/endoplasmic reticulum ATPase (SERCA), which pumps Ca^{2+} into the sarcoplasmic reticulum. This possibility was later supported by the demonstration that NO can stimulate SERCA in arteries via peroxynitrite-mediated S-glutathiolation of Cys674 [572]. The authors did not investigate or discuss how hypoxia was apparently diminishing the ability of cGMP to cause vasodilation. However, since PKG is the only known target of cGMP in vascular smooth muscle, its inhibition by hypoxia is a likely explanation.

They then showed that the oxidant diamide inhibited the activation of sGC by NO donors [564]. Blocking the PPP, which caused a fall in the level of NADPH and should therefore also lead to oxidation of thiols, had the same effect as diamide. This would predict that a fall in H_2O_2 and activation of the PPP caused by hypoxia should increase NO-induced stimulation of sGC, which was consistent with their earlier observation.

PKG exists as two isoforms, PKG1 and PKG2, with the former being predominantly expressed in the vasculature. Two splice variants of PKG1, PKG1 α and PKG1 β , are expressed in vascular smooth muscle. Both are homodimers, with the subunits held together by a leucine zipper interaction, and are activated by cGMP. In 2007, Burgoyne et al. [22] discovered that PKG1 α can also be activated in a cGMP-independent manner by the oxidative formation of a disulfide bond between the Cys42 residues of its two subunits. Although the inactive form of PKG1 α is often referred to as a monomer which dimerizes upon the formation of the disulfide bridge [507], this is incorrect; both active and inactive PKG1 α are dimers [573]. The monomeric form only exists under the denaturing conditions used for immunoblotting, which cause the dimers without the disulfide bond to come apart to form monomers, whereas dimers with a disulfide bridge remain intact; this allows the proportion of active and inactive forms of PKG1 α to be measured. We will use the term 'disulfide dimer complex' (DDC) [573] to refer to the form of PKG1 α activated by oxidation.

The finding that PKG1 α could be directly activated by DDC formation opened up the possibility that decreases in $[\text{H}_2\text{O}_2]$ caused by hypoxia could act on PKG1 α in PASMC to inhibit its vasodilating effect. This led Wolin's laboratory [502] to investigate the effect of exogenous H_2O_2 on DDC formation and the activity of PKG in BPA. They also examined whether H_2O_2 was still able to cause vasorelaxation in arteries in which the expression of sGC had been greatly diminished by a 48 h exposure to ODQ, an approach adapted from a protocol described in [574]. They found that application of 100 μM H_2O_2 to BPA significantly increased DDC formation and PKG activity (measured as the phosphorylation of the PKG1 α substrate VASP (vasodilator-stimulated phosphoprotein)). These effects were suppressed by pre-treating arteries with DTT, suggesting that they were due to PKG1 α oxidation. Depletion of sGC by prolonged ODQ treatment, which almost eliminated the dilating response induced by a NO donor, diminished the vasorelaxation evoked by exogenous H_2O_2 by ~50%, indicating that H_2O_2 was activating PKG1 α both directly and

indirectly, via sGC. Acute ODQ treatment, designed to prevent any effect of NO on sGC, did not block the response to H₂O₂, indicating that this was independent of NO.

Neo et al. [506] studied the relative contributions of sGC and PKG1 α to HPV. They found that, like H₂O₂, hypoxia decreased the DDC formation and activity of PKG α 1. Strikingly, siRNA knockdown of PKG1 α mimicked the acute effects of ebbsen observed previously [458], increasing the amplitude of the contraction induced by 25 mM K⁺ under normoxic conditions but decreasing the amplitude of HPV by ~50%. Additionally, treating BPA with DTT mimicked hypoxia by increasing the contraction to 25 mM K⁺ and diminishing both DDC formation and PKG1 α activity. DTT also abolished HPV.

Depletion of sGC also potentiated the contraction to 25 mM K⁺ under normoxic conditions while reducing the amplitude of HPV by ~60%. siRNA knockdown of the β -subunit of sGC in BPA, which reduced its expression by ~60%, caused similar effects and decreased H₂O₂-induced vasorelaxation and PKG1 α activity. siRNA knockdown of PKG1 α also suppressed H₂O₂-induced relaxation and PKG1 α activity and blocked HPV.

These findings supported the hypothesis that in BPA, hypoxia, by diminishing cellular H₂O₂ levels, causes a reduction of regulatory cysteine thiols on both sGC and PKG1 α , which suppresses their activities. This would attenuate ongoing normoxic vasodilation, thereby causing or augmenting HPV. In contrast to what was observed in BPA, hypoxia caused vasorelaxation in BCA, associated with increases in DDC formation and VASP phosphorylation. This presumably reflected an oxidizing shift in the NADP⁺/NADPH redox couple, which acted through redox networks to oxidize PKG1 α , rather than being due to H₂O₂ [560].

Neo et al. [575] investigated whether PPP-induced reduction of the NADP⁺/NADPH couple was influencing the activity of PKG1 α by regulating the redox state of antioxidant enzymes such as Trx-1 or Prx1 which could suppress its activation by reducing Cys42. In this case, it would be predicted that experimental interventions which interfered with the function of the PPP or of these enzymes would increase PKG1 α DDC formation, leading to its activation and a decrease in vascular contraction.

The authors described several observations which supported the idea that the PPP creates a pro-contractile influence in BPA, which is exerted via NADPH-dependent reduction (i.e., inhibition) of PKG1 α . Hypoxia caused an increase in the NADPH/NADP⁺ ratio, which was antagonized by the PPP blocker 6-AN. PPP blockade acutely relaxed high K⁺-pre-constricted PA, and increased PKG1 α DDC formation and VASP phosphorylation. In addition, the high K⁺ contraction under both normoxic and hypoxic conditions was smaller, and levels of PKG1 α DDC formation and VASP phosphorylation were higher, in PA in which G-6-PD had been knocked down by siRNA. Similar effects were seen following siRNA knockdown of Trx and TrxR-1. Knockdown of thioredoxin reductase also suppressed 6-AN-induced relaxation, DDC formation, and PKG1 α activity.

In contrast, knockdown of Prx1 had no significant effect on the amplitude of the high K⁺ contraction, PKG1 α DDC formation, PKG1 α activity, or 6-AN-induced relaxation. They also examined the effect of TrxR-1 and Prx1 knockdown on the responses of PA to the application of 100 μ M H₂O₂, which caused PKG1 α DDC formation and VASP phosphorylation. Knockdown of TrxR-1 enhanced peroxide-induced relaxation, DDC formation, and PKG1 α activity, whereas knockdown of Prx1 had the opposite effects.

The results supported the idea that the PPP exerts an inhibitory effect on the activity of PKG1 α , doing so by maintaining a high level of cellular NADPH, which acts through Trx-1 and TrxR-1 to reduce PKG1 α , thus suppressing its vasorelaxing effect. This inhibitory effect on the underlying vasorelaxing influence exerted by PKG is enhanced during hypoxia because the PPP is activated [446], promoting HPV. This concept was in line with the observation [576] that G6P-D, Trx-1, and PKG co-precipitated, suggesting that they form a complex.

The results also suggested that Prx1 is not involved in O_2 sensing in BPA. This was shown by the observation that Prx1 knockdown had no effect on the high K^+ contraction under normoxic conditions, which would mean that this antioxidant enzyme is not contributing to the normoxic vasorelaxation which is removed by hypoxia. The authors also observed that depletion of sGC had no effect on 6-AN-mediated relaxation under hypoxic conditions, implying that the PPP and NADPH were not directly regulating cGMP production. The authors therefore concluded that regulation of $PKG1\alpha$ DDC formation by NADPH might contribute to hypoxic responses associated with changes in the redox state of NADPH.

Figure 10 summarizes the proposed involvement of the sGC/PKG axis and hypoxia-induced removal of a tonic normoxic H_2O_2 -dependent vasodilation as a mechanism contributing to HPV.




Figure 10. Proposed role of $PKG1\alpha$ and sGC in HPV. According to this model, HPV is largely due to the removal of a basal vasodilating mechanism, which is caused by an ongoing activation of sGC and $PKG1\alpha$. Hypoxia removes this influence via two pathways. Firstly, by activating the PPP, it increases the NADPH/NADP⁺ ratio. The increase in NADPH causes the reduction of TrxR-1 and Trx-1, thereby reversing the resting oxidation and activation of $PKG1\alpha$, diminishing its vasorelaxing influence. Secondly, the fall in PO_2 decreases the ambient intracellular concentration of H_2O_2 , further decreasing the activation of $PKG1\alpha$ both directly by promoting its activity and indirectly by inhibiting sGC. This figure is largely based on information presented in [575].

8.4. Does the Presence of Extracellular H_2O_2 Affect HPV?

As described in Section 4, it is thought that the basal extracellular H_2O_2 concentration is in the range of 1–5 μM . This level of extracellular H_2O_2 is predicted to cause an inward leak, which would raise its intracellular concentration, especially adjacent to the cell membrane [249], and it is therefore conceivable that this could influence the effects of changes in mitochondrial H_2O_2 production on redox-sensitive HPV effector targets, especially those in the plasmalemma. In two studies, Wolin's laboratory examined the possible impact of extracellular H_2O_2 on HPV using catalase, which was applied at a low

concentration (1 μM) and is said to scavenge only extracellular H_2O_2 . These investigations, carried out in isolated PA from cow [482] and mouse [483], found no effect of catalase on HPV under control conditions. However, HPV in both arteries was suppressed by maneuvers which raised the extracellular $[\text{H}_2\text{O}_2]$, an effect which was partly reversed by catalase. Based on other results in their paper, Patel et al. [483] proposed that this was due to oxidation-dependent stimulation of $\text{PKG1}\alpha$, presumably exerted by an excessive inward leak of H_2O_2 . These results should be interpreted with caution since no evidence was presented that catalase was not penetrating into the cells (which it has been shown to do [358]). However, taken at face value, they support the possibility that the presence of extracellular H_2O_2 might influence HPV.

8.5. Summary and Critique: Does Activation of the PPP and the Withdrawal of H_2O_2 /PKG-Mediated Normoxic Vasodilation Cause HPV?

The picture of the involvement of ROS and cellular redox in HPV which emerges from the papers published on this subject by Wolin, Gupte, and colleagues is complex, but incorporates two proposed pathways, as shown in Figures 9 and 10:

1. Under normoxic conditions, the production of superoxide/ H_2O_2 by Nox4 is greater in PA than in systemic arteries because the PPP, being more active, generates more NADPH. The resulting higher level of H_2O_2 creates an ongoing vasodilating influence by stimulating $\text{PKG1}\alpha$, both directly, by oxidizing Cys42 on $\text{PKG1}\alpha$, and indirectly, by activating sGC. Hypoxia decreases the production of superoxide/ H_2O_2 by Nox4, thereby raising vascular tone by inhibiting this baseline vasorelaxation.
2. Hypoxia also suppresses the activity of $\text{PKG1}\alpha$ by stimulating G-6-PD and the PPP, causing a consequent increase in the NADPH/NADP⁺ ratio. This acts, at least in part, through Trx-1 and TrxR-1, to reduce $\text{PKG1}\alpha$, causing it to become less active, which promotes contraction. The activation of G-6-PD is proposed to result from an increase in cytoplasmic [ROS], which also promotes HPV by stimulating rho kinase and Ca^{2+} -dependent contractile mechanisms.

Below, we look critically at several aspects of these proposals.

8.5.1. Decreased Production of H_2O_2 by Nox as a Cause of HPV

The high $K_m(\text{O}_2)$ of Nox4 (Section 3.2.2), which allows its production of H_2O_2 to be sensitive to even small decreases in PO_2 , renders it an eminently feasible O_2 sensor for HPV. It has been shown, for example, that H_2O_2 production by Nox4 expressed in HEK293 cells falls by ~50% when the PO_2 is decreased from 150 to 80 Torr [68].

Nonetheless, the knockout of Nox4 had no effect on basal PAP or HPV [404] in mice, indicating that it does not regulate basal vascular PA tone and is not required for the response to hypoxia, at least in this species. On the other hand, sustained HPV was diminished by the knockout of $\text{p}^{22\text{phox}}$ [407], suggesting that another Nox isoform, probably Nox1, plays a role in HPV. However, $\text{p}^{22\text{phox}}$ knockout had no effect on basal PAP. This argues against the possibility that HPV in mice is due to the withdrawal of basal Nox-induced vasodilation, suggesting instead that Nox is contributing to sustained HPV by increasing its production of ROS.

These observations do not rule out the possibility that the loss of basal ROS production by Nox4 is important for HPV in species other than the mouse. Nonetheless, the involvement of decreased ROS production by Nox4 in HPV in cow can be questioned on several fronts. Most importantly, direct evidence that hypoxia decreases ROS levels in BPA is inconsistent. A hypoxia-induced fall in ROS in intact BPA was demonstrated in only a single study [549], which measured H_2O_2 using a high concentration of lucigenin with HRP. Lucigenin at this concentration is subject to redox cycling, making it a

suboptimal indicator for examining the effect of hypoxia on ROS levels (Section 7.4). Conversely, the investigation employing the intracellular indicator DHE reported that hypoxia increases ROS in BPA [446]. Although DHE also has its drawbacks, its suitability as a detector of cell oxidation has been vigorously defended [577].

Evidence [458] that the withdrawal of normoxic vasodilation caused by Nox4-generated H_2O_2 contributes to HPV in BPA can also be challenged. As described in this paper, siRNA knockout of Nox4 did not affect the contraction to 25 mM K^+ , which the authors used as an index of basal reactivity. This conflicted with their hypothesis that Nox4 exerts a tonic vasodilating effect on basal tone, which predicts that Nox4 knockout should increase the K^+ contraction. The authors explained this apparent discrepancy by proposing that ROS generation by Nox4 exerts opposing vasodilating and vasoconstricting effects, and that when ROS production is acutely interrupted by hypoxia, the dilating effect is immediately removed, whereas the constricting effect only disappears after a delay of several hours.

However, down-regulation of sGC and $\text{PKG1}\alpha$, the putative targets by which H_2O_2 produced by Nox4 was suggested to suppress basal tone, did increase the contraction to 25 mM K^+ [506]. This observation seemed to undermine the authors' explanation for the lack of an effect of Nox4 knockout on basal reactivity, which provides an important underpinning for their proposal that Nox4-derived H_2O_2 causes normoxic vasodilation. As to the possibility that the attenuation of superoxide/ H_2O_2 production by other Nox isoforms by hypoxia might contribute to HPV, this laboratory did not detect the presence of Nox1 in these arteries [558], and demonstrated that although blocking Nox2 with gp91-dsat diminished ROS levels in BPA, it had no effect on either basal reactivity or HPV [458]. We therefore believe that the proposal that a significant component of HPV in BPA is due to the loss of basal vasorelaxation associated with H_2O_2 production by Nox remains speculative.

8.5.2. Does a Fall in H_2O_2 During Hypoxia Cause HPV by Inhibiting sGC?

If HPV is partly caused by the removal of H_2O_2 -induced basal oxidative activation of sGC and/or an increase in PPP-derived NADPH [502,506,548,549], the following would be predicted: 1. A block of sGC or PKG should inhibit HPV and also mimic hypoxia in causing PA constriction. 2. Hypoxia should decrease the cGMP content of PA. 3. The activity of sGC should be increased by H_2O_2 or equivalent oxidant stimuli.

The first prediction is supported by an *in vivo* study of HPV in cats [578], which showed that the sGC blocker methylene blue (MB) increased baseline PAP and abolished the response to hypoxia. MB also increased baseline normoxic PAP in pigs and sheep, although its effect on HPV was not investigated [579]. Conversely, MB [580,581] and ODQ [582] had no effect on baseline normoxic PAP, and augmented HPV in an isolated perfused rat lung. In isolated rat PA, MB increased HPV, although this effect was absent in severe hypoxia [583]. Vermeersch and colleagues [584] compared HPV in wildtype mice and those in which the α_1 subunit of sGC $\alpha_1\beta_1$ isoform, the predominant form of sGC in the vasculature, had been rendered less active by deletion of exon 6 of the α_1 subunit gene. RVSP under normoxic conditions and during a 10 min hypoxic challenge was not different in the two groups and hypoxia had no effect on the lung cGMP content in the control mice. Also, although ODQ slightly reduced the amplitude of HPV in isolated perfused mouse lung, it did not increase normoxic PAP, going against the idea of sGC-mediated normoxic vasorelaxation [585]. Fouty et al. [582] found that the PKG inhibitor Rp-8-pCPT-cGMPs had no effect on basal PAP or HPV in isolated perfused rat lung. They verified that Rp-8-pCPT-cGMPs suppressed PKG activity by monitoring phosphorylation of the $\text{Ins}(1,4,5)\text{P}_3$ receptor, a PKG target. The differential effect of Rp-8-pCPT-cGMPs and ODQ on HPV in this study is difficult to explain, since protein kinase G is the only known target of cGMP

in vascular smooth muscle. Nevertheless, neither drug affected basal PAP. With regard to the second prediction, contrasting effects of hypoxia on cGMP levels in PA have been reported. In BPA, hypoxia decreased cGMP in endothelium-denuded BPA [548]. In rat PA, however, hypoxia decreased the cGMP content of intact but not endothelium-denuded PAs [586]. This suggested that hypoxia was suppressing the activity of sGC, but that this was secondary to a fall in endothelial NO release rather than being due a direct effect of hypoxia on sGC in the smooth muscle. Taken together, the results of these studies argue against a role for sGC in causing basal PA vasorelaxation in rodents, although this might occur in cats in a manner similar to that proposed for BPA.

The proposal that H₂O₂ production under normoxic conditions creates a vasodilating influence in BPA, which is in part due to the activation of sGC, and that this is decreased by hypoxia, rests on evidence that (1) exogenous H₂O₂ caused vasorelaxation and increased the cGMP content in endothelium-intact arteries [493], (2) hypoxia decreased the cGMP content of endothelium-denuded arteries [548], (3) depletion of cellular sGC β 1 using prolonged ODQ treatment increased basal reactivity (assessed as the contraction to 25 mM K⁺ under normoxic conditions), [502], (4) depletion of sGC β 1 using long-term ODQ treatment or siRNA knockdown inhibited vasorelaxation and activation of PKG (assessed using VASP phosphorylation) by exogenous H₂O₂ under hypoxic conditions [502,506], and (5) depletion of sGC using siRNA knockdown increased basal reactivity but decreased HPV [506].

Importantly, Neo et al. found that the vasorelaxation of BPA induced by the H₂O₂ occurred in endothelium-denuded arteries, and whereas it was decreased by downregulating sGC expression, it was not affected by acute treatment with ODQ [502]. Since ODQ would block stimulation of sGC due to any remaining NO, they concluded from these results that the activation of sGC by H₂O₂ must be NO-independent (indeed, this laboratory had previously presented evidence that NO-dependent activation of sGC was inhibited by oxidation [564]).

In this case, their model predicts that if hypoxia causes a fall in H₂O₂ production, it should decrease the cGMP concentration in endothelium-denuded arteries, an effect they had previously observed in BPA [548]. However, the cGMP content of endothelium-denuded rat PA was not affected by hypoxia [586], and in porcine PA, Ye et al. also found that hypoxia did not affect the cyclic GMP level in the presence of either ODQ or the eNOS antagonist nitro-L-arginine [587] and that reducing agents did not affect (DTT and GSH), or increased (L-cysteine and tris(2-carboxyethyl)phosphine), the cGMP content of nitro-L-arginine-treated arteries. Intriguingly, they also reported that sGC dimerization, which is required for its activation by NO [588], was present under basal conditions, and was decreased by both hypoxia and reductants. Accordingly, NO-dependent vasodilation was depressed by reductants. These observations contradict the evidence in BPA that oxidation depresses NO-dependent and increases NO-independent sGC activity but support the general concept that the reduction of sGC by hypoxia can cause its inhibition. However, since NO release falls during hypoxia [589], the extent to which the redox regulation of NO-dependent sGC activation could influence HPV is uncertain.

8.5.3. Does a Fall in H₂O₂ During Hypoxia Cause HPV by Directly Inhibiting PKG1 α ?

The proposal that a decrease in H₂O₂ levels and/or an increase in NADPH production by the PPP caused by hypoxia contributes to HPV by reversing oxidation-induced PKG1 α activation is consistent with extensive evidence showing that it exerts an important vasodilating influence in systemic arteries through its phosphorylation of multiple target proteins which control vascular tone [590]. It is also well established that PKG activity in arteries is increased by the oxidation of Cys42, leading to the formation of a disulfide bond in PKG1 α in mice [22,573,591], and that this creates an ongoing vasodilating influence

under baseline conditions, at least in systemic resistance vessels, as evidenced by the observation that blood pressure is raised in mice expressing a 'redox-dead' form of PKG1 α [592].

With regard to the involvement of PKG1 α in HPV, this hypothesis is supported by evidence, described in Section 8.3.2, for the following: 1. Exposure of BPA to 0.1 and 1 mM H₂O₂ increased PKG1 α DDC formation and phosphorylation of its target VASP, and these responses were reversed by DTT [502]. 2. Hypoxia caused contraction and decreased PKG DDC formation and VASP phosphorylation of endothelium-denuded BPA, whereas the opposite effects on PKG occurred in BCA, which relaxed to hypoxia. 3. DTT increased force in BPA under both aerobic and hypoxic conditions and diminished PKG1 α DDC formation and VASP phosphorylation. 4. siRNA knockdown of PKG1 α increased basal reactivity (i.e., the contraction to 25 mM K⁺ under normoxic conditions) but depressed HPV [506]. 5. Block of the PPP, which would reduce the supply of NADPH available to reduce and therefore inactivate PKG1 α , diminished the contraction evoked by 25 mM K⁺ under both normoxic and hypoxic conditions [575].

However, it is not clear whether the effect of the PPP on contraction, exerted through the redox regulation of PKG1 α [575], is specific to HPV. The authors proposed that the PPP was promoting hypoxic contraction of BPA by increasing the production of NADPH, which would transfer reducing equivalents via TrxR-1 and Trx-1 to PKG1 α , thereby suppressing its vasodilating activity (Figure 10). In support of this scheme, they found that interventions used to block the influence of the PPP on contraction (i.e., siRNA knockdown of G6PD, Trx-1, TrxR-1) decreased the contraction induced by 25 mM K⁺ under hypoxic conditions. Importantly, however, these interventions caused a comparable decrease in the high K⁺ contraction recorded under normoxic conditions, suggesting that the PPP similarly promotes contraction under both normoxic and hypoxic conditions rather than selectively enhancing HPV. Along similar lines, Chettimada et al., who also showed that blocking the PPP with 6-AN increased HPV, found that this caused equivalent increases in the activity of PKG under normoxic and hypoxic conditions [576]. Rather than demonstrating that the activation of the PPP acting via an NADPH/TrxR/Trx/PKG1 α pathway is a specific cause of HPV, these results suggest that NADPH produced by the PPP may promote contraction by inhibiting PKG1 α regardless of the oxygen concentration.

There is currently no information about the role of PKG1 α oxidation in acute HPV in other species, although there is evidence [593] that this is increased during chronic hypoxia and tends to ameliorate pulmonary hypertension in mice.

8.5.4. Is Activation of the PPP Important for HPV?

Although we have argued that the evidence that the PPP causes/promotes HPV by inhibiting PKG1 α is unconvincing, it is possible that stimulation of the PPP by hypoxia could trigger HPV via other contractile mechanisms, e.g., activation of RhoK [446], or inhibition of K_v channels by the interaction of NADPH with their b subunits [274]. Certainly, the evidence that hypoxia stimulates the PPP and causes contraction in BPA, whereas it inhibits the PPP and causes relaxation in BCA, offers support, albeit indirect, for the concept that it regulates the contractile response to low PO₂. Unfortunately, it is difficult to draw any conclusions about the involvement of the PPP in HPV because no other laboratories seem to have investigated this. Nevertheless, in agreement with the observation that the G-6-PD antagonist 6-AN blocked HPV in BPA, we have found that dehydroepiandrosterone, which also blocks this enzyme, has a similar effect in rat PA (Figure 11).




Figure 11. HPV was evoked by gassing small endothelium-intact PA from rats in a conventional myograph with 95% N/5%CO₂/0% O₂ as previously described; the PO₂ during hypoxia was shown to be 15–20 Torr under these conditions [10]. A low concentration of PGF_{2α} was applied before hypoxia to create a small degree of pre-tone; this increases the amplitude of HPV, particularly the transient initial Phase 1. Application of DHEA significantly ($p < 0.05$ by 2-way ANOVA and also paired T-test) depressed the amplitude of Phase 1 and also of the sustained Phase 2. In the absence of pre-tone (in which case phase 1 is not evident), DHEA also virtually abolished HPV in 7 arteries.

There is also evidence that the PPP in other types of cells is rapidly stimulated by oxidative stress [594] and several mechanisms by which the oxidation of thiols in glycolytic enzymes (e.g., glyceraldehyde-3-phosphate dehydrogenase, pyruvate kinase) can divert metabolic flux into the PPP have been defined (e.g., [595]. Supporting the involvement of G-6-PD and the PPP in regulating contraction, the vasoconstrictor U46619 caused an increase in G-6-PD activity in BCA, which was mediated by a ROS-dependent stimulation of PKC δ [568]. Also, complexes containing G-6-PD, Trx-1, and PKG can be detected in BPA homogenates [576], and L-type voltage-gated Ca²⁺ channels in BCA and human internal mammary artery smooth muscle cells were shown to be activated by a direct interaction with G-6-PD in caveoli [596]. Further investigation of the role of PPP activation in HPV, focused on determining the effect of hypoxia on the redox state of the NADP⁺:NADPH couple [597] and on identifying the effectors through which the PPP could induce HPV, would therefore be of great interest.

9. Conclusions

The proposal that cell redox mechanisms acting on cell thiols are responsible for O₂ sensing in HPV was first mooted by Kenneth Weir and colleagues in a remarkably prescient study published more than 40 years ago [283]. As we have described, experiments in the intervening time have led to the development of three main models designed to describe how hypoxia induces changes in PASMOC ROS production and cytoplasmic redox state which engage with specific HPV effector mechanisms. Although many of these studies utilized techniques which were innovative and state-of-the-art at the time they were carried out, the rapid pace of conceptual and technical development in the field of redox biology, especially over the past decade, has brought into question the interpretation of many of the observations which underpin these models. Largely because it is supported

by a larger evidence base, and one which draws more heavily on the use of newer techniques which are at present viewed as being more valid, we believe that the Mitochondrial ROS hypothesis currently provides the most convincing picture of how hypoxia acts through cell redox mechanisms to cause HPV.

Nevertheless, many interesting questions remain. The mechanisms by which hypoxia can increase mitochondrial ROS production remain uncertain, as do the identities of the HPV effectors which are linked to an increase in ROS, and the precise nature of the involvement of Cox4i2 in O₂ sensing in PASM. The possible involvement of NADH in HPV, possibly via the regulation of Kv or voltage-gated Ca²⁺ channels, is a long-standing but unresolved concept which has been given a renewed lease of life by recent observations in CBCC. Indeed, the question as to what extent common mechanisms are used by the various homeostatic sensors of acute hypoxia present in the body remains open. Recent evidence suggesting that hypoxia may exert complex effects on the cytoplasmic redox state (due to spatial and/or temporal compartmentalization?) rather than generating a simple up/down ROS signal is also worthy of note. In addition, concerns about the use of non-physiologic conditions and regarding whether cells were being exposed to O₂ concentrations which differed from those which were intended also suggest that the quest for the O₂-sensing mechanism in HPV should not be regarded as finished. It is noteworthy, for example, that the involvement in HPV of ROS-independent mechanisms which contribute to acute O₂ sensing in other types of cells has not been investigated.

Happily, progress in the development of increasingly sensitive/specific indicators for cellular H₂O₂ [598], peroxynitrite [599], free NADH ([600]), and NADPH ([100]), as well as probes for intracellular PO₂ ([601]), has made it more possible than ever to explore the involvement of cell redox mechanisms as O₂ sensors in PASM and other cells. It will be interesting to see where the research in this area takes us.

Funding: This research received no external funding.

Data Availability Statement: No new data were created or analyzed in this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gierhardt, M.; Pak, O.; Walmrath, D.; Seeger, W.; Grimminger, F.; Ghofrani, H.A.; Weissmann, N.; Hecker, M.; Sommer, N. Impairment of hypoxic pulmonary vasoconstriction in acute respiratory distress syndrome. *Eur. Respir. Rev.* **2021**, *30*, 210059. <https://doi.org/10.1183/16000617.0059-2021>.
2. Sommer, N.; Dietrich, A.; Schermuly, R.T.; Ghofrani, H.A.; Gudermaun, T.; Schulz, R.; Seeger, W.; Grimminger, F.; Weissmann, N. Regulation of hypoxic pulmonary vasoconstriction: Basic mechanisms. *Eur. Respir. J.* **2008**, *32*, 1639–1651. <https://doi.org/10.1183/09031936.00013908>.
3. Sylvester, J.T.; Shimoda, L.A.; Aaronson, P.I.; Ward, J.P. Hypoxic pulmonary vasoconstriction. *Physiol. Rev.* **2012**, *92*, 367–520. <https://doi.org/10.1152/physrev.00041.2010>.
4. Nagendran, J.; Stewart, K.; Hoskinson, M.; Archer, S.L. An anesthesiologist's guide to hypoxic pulmonary vasoconstriction: Implications for managing single-lung anesthesia and atelectasis. *Curr. Opin. Anaesthesiol.* **2006**, *19*, 34–43. <https://doi.org/10.1097/01.aco.0000192777.09527.9e>.
5. Dinunno, F.A. Skeletal muscle vasodilation during systemic hypoxia in humans. *J. Appl. Physiol.* **2016**, *120*, 216–225. <https://doi.org/10.1152/jappphysiol.00256.2015>.
6. Dorrington, K.L.; Clar, C.; Young, J.D.; Jonas, M.; Tansley, J.G.; Robbins, P.A. Time course of the human pulmonary vascular response to 8 hours of isocapnic hypoxia. *Am. J. Physiol.* **1997**, *273*, H1126–H1134. <https://doi.org/10.1152/ajpheart.1997.273.3.H1126>.
7. Talbot, N.P.; Balanos, G.M.; Dorrington, K.L.; Robbins, P.A. Two temporal components within the human pulmonary vascular response to approximately 2 h of isocapnic hypoxia. *J. Appl. Physiol.* **2005**, *98*, 1125–1139. <https://doi.org/10.1152/jappphysiol.00903.2004>.

8. Marshall, B.E.; Clarke, W.R.; Costarino, A.T.; Chen, L.; Miller, F.; Marshall, C. The dose-response relationship for hypoxic pulmonary vasoconstriction. *Respir. Physiol.* **1994**, *96*, 231–247. [https://doi.org/10.1016/0034-5687\(94\)90129-5](https://doi.org/10.1016/0034-5687(94)90129-5).
9. Fishman, A.P. Acute hypoxia and pulmonary vasoconstriction in humans: Uncovering the mechanism of the pressor response. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2004**, *287*, L893–L894. <https://doi.org/10.1152/classicessays.00004.2004>.
10. Robertson, T.P.; Hague, D.; Aaronson, P.I.; Ward, J.P. Voltage-independent calcium entry in hypoxic pulmonary vasoconstriction of intrapulmonary arteries of the rat. *J. Physiol.* **2000**, *525*, 669–680. <https://doi.org/10.1111/j.1469-7793.2000.t01-1-00669.x>.
11. Weissmann, N.; Dietrich, A.; Fuchs, B.; Kalwa, H.; Ay, M.; Dumitrascu, R.; Olschewski, A.; Storch, U.; Mederos y Schnitzler, M.; Ghofrani, H.A.; et al. Classical transient receptor potential channel 6 (TRPC6) is essential for hypoxic pulmonary vasoconstriction and alveolar gas exchange. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 19093–19098. <https://doi.org/10.1073/pnas.0606728103>.
12. Jain, P.P.; Hosokawa, S.; Xiong, M.; Babicheva, A.; Zhao, T.; Rodriguez, M.; Rahimi, S.; Pourhashemi, K.; Balistrieri, F.; Lai, N.; et al. Revisiting the mechanism of hypoxic pulmonary vasoconstriction using isolated perfused/ventilated mouse lung. *Pulm. Circ.* **2020**, *10*, 2045894020956592. <https://doi.org/10.1177/2045894020956592>.
13. Jabr, R.I.; Toland, H.; Gelband, C.H.; Wang, X.X.; Hume, J.R. Prominent role of intracellular Ca²⁺ release in hypoxic vasoconstriction of canine pulmonary artery. *Br. J. Pharmacol.* **1997**, *122*, 21–30. <https://doi.org/10.1038/sj.bjp.0701326>.
14. Weigand, L.; Foxson, J.; Wang, J.; Shimoda, L.A.; Sylvester, J.T. Inhibition of hypoxic pulmonary vasoconstriction by antagonists of store-operated Ca²⁺ and nonselective cation channels. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2005**, *289*, L5–L13. <https://doi.org/10.1152/ajplung.00044.2005>.
15. Robertson, T.P.; Dipp, M.; Ward, J.P.; Aaronson, P.I.; Evans, A.M. Inhibition of sustained hypoxic vasoconstriction by Y-27632 in isolated intrapulmonary arteries and perfused lung of the rat. *Br. J. Pharmacol.* **2000**, *131*, 5–9. <https://doi.org/10.1038/sj.bjp.0703537>.
16. Aaronson, P.I.; Robertson, T.P.; Ward, J.P. Endothelium-derived mediators and hypoxic pulmonary vasoconstriction. *Respir. Physiol. Neurobiol.* **2002**, *132*, 107–120. [https://doi.org/10.1016/s1569-9048\(02\)00053-8](https://doi.org/10.1016/s1569-9048(02)00053-8).
17. Archer, S.L.; Wu, X.C.; Thebaud, B.; Nsair, A.; Bonnet, S.; Tyrrell, B.; McMurtry, M.S.; Hashimoto, K.; Harry, G.; Michelakis, E.D. Preferential expression and function of voltage-gated, O₂-sensitive K⁺ channels in resistance pulmonary arteries explains regional heterogeneity in hypoxic pulmonary vasoconstriction: Ionic diversity in smooth muscle cells. *Circ. Res.* **2004**, *95*, 308–318. <https://doi.org/10.1161/01.RES.0000137173.42723.fb>.
18. Connolly, M.J.; Prieto-Lloret, J.; Becker, S.; Ward, J.P.; Aaronson, P.I. Hypoxic pulmonary vasoconstriction in the absence of pretone: Essential role for intracellular Ca²⁺ release. *J. Physiol.* **2013**, *591*, 4473–4498. <https://doi.org/10.1113/jphysiol.2013.253682>.
19. Moudgil, R.; Michelakis, E.D.; Archer, S.L. Hypoxic pulmonary vasoconstriction. *J. Appl. Physiol.* **2005**, *98*, 390–403. <https://doi.org/10.1152/japplphysiol.00733.2004>.
20. Wang, L.; Yin, J.; Nickles, H.T.; Ranke, H.; Tabuchi, A.; Hoffmann, J.; Tabeling, C.; Barbosa-Sicard, E.; Chanson, M.; Kwak, B.R.; et al. Hypoxic pulmonary vasoconstriction requires connexin 40-mediated endothelial signal conduction. *J. Clin. Investig.* **2012**, *122*, 4218–4230. <https://doi.org/10.1172/JCI59176>.
21. Weir, E.K.; Archer, S.L. The mechanism of acute hypoxic pulmonary vasoconstriction: The tale of two channels. *FASEB J.* **1995**, *9*, 183–189. <https://doi.org/10.1096/fasebj.9.2.7781921>.
22. Burgoyne, J.R.; Madhani, M.; Cuello, F.; Charles, R.L.; Brennan, J.P.; Schroder, E.; Browning, D.D.; Eaton, P. Cysteine redox sensor in PKGIa enables oxidant-induced activation. *Science* **2007**, *317*, 1393–1397. <https://doi.org/10.1126/science.1144318>.
23. Waypa, G.B.; Chandel, N.S.; Schumacker, P.T. Model for hypoxic pulmonary vasoconstriction involving mitochondrial oxygen sensing. *Circ. Res.* **2001**, *88*, 1259–1266. <https://doi.org/10.1161/hh1201.091960>.
24. Rathore, R.; Zheng, Y.M.; Niu, C.F.; Liu, Q.H.; Korde, A.; Ho, Y.S.; Wang, Y.X. Hypoxia activates NADPH oxidase to increase [ROS]_i and [Ca²⁺]_i through the mitochondrial ROS-PKCepsilon signaling axis in pulmonary artery smooth muscle cells. *Free Radic. Biol. Med.* **2008**, *45*, 1223–1231. <https://doi.org/10.1016/j.freeradbiomed.2008.06.012>.
25. Alruwaili, N.; Kandhi, S.; Sun, D.; Wolin, M.S. Metabolism and Redox in Pulmonary Vascular Physiology and Pathophysiology. *Antioxid. Redox Signal.* **2019**, *31*, 752–769. <https://doi.org/10.1089/ars.2018.7657>.
26. Moral-Sanz, J.; Lewis, S.A.; MacMillan, S.; Ross, F.A.; Thomson, A.; Viollet, B.; Foretz, M.; Moran, C.; Hardie, D.G.; Evans, A.M. The LKB1-AMPK-alpha1 signaling pathway triggers hypoxic pulmonary vasoconstriction downstream of mitochondria. *Sci. Signal.* **2018**, *11*, eaau0296. <https://doi.org/10.1126/scisignal.aau0296>.
27. Moral-Sanz, J.; Mahmoud, A.D.; Ross, F.A.; Eldstrom, J.; Fedida, D.; Hardie, D.G.; Evans, A.M. AMP-activated protein kinase inhibits Kv 1.5 channel currents of pulmonary arterial myocytes in response to hypoxia and inhibition of mitochondrial oxidative phosphorylation. *J. Physiol.* **2016**, *594*, 4901–4915. <https://doi.org/10.1113/JP272032>.

28. Olson, K.R.; Dombkowski, R.A.; Russell, M.J.; Doellman, M.M.; Head, S.K.; Whitfield, N.L.; Madden, J.A. Hydrogen sulfide as an oxygen sensor/transducer in vertebrate hypoxic vasoconstriction and hypoxic vasodilation. *J. Exp. Biol.* **2006**, *209*, 4011–4023. <https://doi.org/10.1242/jeb.02480>.
29. Olson, K.R.; Whitfield, N.L.; Bearden, S.E.; St Leger, J.; Nilson, E.; Gao, Y.; Madden, J.A. Hypoxic pulmonary vasodilation: A paradigm shift with a hydrogen sulfide mechanism. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2010**, *298*, R51–R60. <https://doi.org/10.1152/ajpregu.00576.2009>.
30. Olson, K.R. Are Reactive Sulfur Species the New Reactive Oxygen Species? *Antioxid. Redox Signal.* **2020**, *33*, 1125–1142. <https://doi.org/10.1089/ars.2020.8132>.
31. Olson, K.R. A Case for Hydrogen Sulfide Metabolism as an Oxygen Sensing Mechanism. *Antioxidants* **2021**, *10*, 1650. <https://doi.org/10.3390/antiox10111650>.
32. Grimmer, B.; Kuebler, W.M. The endothelium in hypoxic pulmonary vasoconstriction. *J. Appl. Physiol.* **2017**, *123*, 1635–1646. <https://doi.org/10.1152/japplphysiol.00120.2017>.
33. Nagaraj, C.; Tang, B.; Balint, Z.; Wygrecka, M.; Hrzenjak, A.; Kwapiszewska, G.; Stacher, E.; Lindenmann, J.; Weir, E.K.; Olschewski, H.; et al. Src tyrosine kinase is crucial for potassium channel function in human pulmonary arteries. *Eur. Respir. J.* **2013**, *41*, 85–95. <https://doi.org/10.1183/09031936.00211811>.
34. Riesco-Fagundo, A.M.; Perez-Garcia, M.T.; Gonzalez, C.; Lopez-Lopez, J.R. O₂ modulates large-conductance Ca²⁺-dependent K⁺ channels of rat chemoreceptor cells by a membrane-restricted and CO-sensitive mechanism. *Circ. Res.* **2001**, *89*, 430–436. <https://doi.org/10.1161/hh1701.095632>.
35. Chang, A.J.; Ortega, F.E.; Riegler, J.; Madison, D.V.; Krasnow, M.A. Oxygen regulation of breathing through an olfactory receptor activated by lactate. *Nature* **2015**, *527*, 240–244. <https://doi.org/10.1038/nature15721>.
36. McCartney, C.E.; McClafferty, H.; Huibant, J.M.; Rowan, E.G.; Shipston, M.J.; Rowe, I.C. A cysteine-rich motif confers hypoxia sensitivity to mammalian large conductance voltage- and Ca-activated K (BK) channel alpha-subunits. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 17870–17876. <https://doi.org/10.1073/pnas.0505270102>.
37. Sommer, N.; Strielkov, I.; Pak, O.; Weissmann, N. Oxygen sensing and signal transduction in hypoxic pulmonary vasoconstriction. *Eur. Respir. J.* **2016**, *47*, 288–303. <https://doi.org/10.1183/13993003.00945-2015>.
38. Dao, V.T.; Elbatreek, M.H.; Altenhofer, S.; Casas, A.I.; Pachado, M.P.; Neullens, C.T.; Knaus, U.G.; Schmidt, H. Isoform-selective NADPH oxidase inhibitor panel for pharmacological target validation. *Free Radic. Biol. Med.* **2020**, *148*, 60–69. <https://doi.org/10.1016/j.freeradbiomed.2019.12.038>.
39. Archer, S.L.; Gomberg-Maitland, M.; Maitland, M.L.; Rich, S.; Garcia, J.G.; Weir, E.K. Mitochondrial metabolism, redox signaling, and fusion: A mitochondria-ROS-HIF-1 α -Kv1.5 O₂-sensing pathway at the intersection of pulmonary hypertension and cancer. *Am. J. Physiol. Heart Circ. Physiol.* **2008**, *294*, H570–H578. <https://doi.org/10.1152/ajpheart.01324.2007>.
40. Nozik-Grayck, E.; Stenmark, K.R. Role of reactive oxygen species in chronic hypoxia-induced pulmonary hypertension and vascular remodeling. *Adv. Exp. Med. Biol.* **2007**, *618*, 101–112. https://doi.org/10.1007/978-0-387-75434-5_8.
41. Wang, Y.X.; Zheng, Y.M. ROS-dependent signaling mechanisms for hypoxic Ca²⁺ responses in pulmonary artery myocytes. *Antioxid. Redox Signal.* **2010**, *12*, 611–623. <https://doi.org/10.1089/ars.2009.2877>.
42. Resta, T.C.; Broughton, B.R.; Jernigan, N.L. Reactive oxygen species and RhoA signaling in vascular smooth muscle: Role in chronic hypoxia-induced pulmonary hypertension. *Adv. Exp. Med. Biol.* **2010**, *661*, 355–373. https://doi.org/10.1007/978-1-60761-500-2_23.
43. Veit, F.; Pak, O.; Brandes, R.P.; Weissmann, N. Hypoxia-dependent reactive oxygen species signaling in the pulmonary circulation: Focus on ion channels. *Antioxid. Redox Signal.* **2015**, *22*, 537–552. <https://doi.org/10.1089/ars.2014.6234>.
44. Jaitovich, A.; Jourdain, D. A Brief Overview of Nitric Oxide and Reactive Oxygen Species Signaling in Hypoxia-Induced Pulmonary Hypertension. *Adv. Exp. Med. Biol.* **2017**, *967*, 71–81. https://doi.org/10.1007/978-3-319-63245-2_6.
45. Huetsch, J.C.; Suresh, K.; Shimoda, L.A. Regulation of Smooth Muscle Cell Proliferation by NADPH Oxidases in Pulmonary Hypertension. *Antioxidants* **2019**, *8*, 56. <https://doi.org/10.3390/antiox8030056>.
46. Yan, S.; Resta, T.C.; Jernigan, N.L. Vasoconstrictor Mechanisms in Chronic Hypoxia-Induced Pulmonary Hypertension: Role of Oxidant Signaling. *Antioxidants* **2020**, *9*, 999. <https://doi.org/10.3390/antiox9100999>.
47. Reyes-Garcia, J.; Carbajal-Garcia, A.; Di Mise, A.; Zheng, Y.M.; Wang, X.; Wang, Y.X. Important Functions and Molecular Mechanisms of Mitochondrial Redox Signaling in Pulmonary Hypertension. *Antioxidants* **2022**, *11*, 473. <https://doi.org/10.3390/antiox11030473>.

48. Sies, H.; Belousov, V.V.; Chandel, N.S.; Davies, M.J.; Jones, D.P.; Mann, G.E.; Murphy, M.P.; Yamamoto, M.; Winterbourn, C. Defining roles of specific reactive oxygen species (ROS) in cell biology and physiology. *Nat. Rev. Mol. Cell. Biol.* **2022**, *23*, 499–515. <https://doi.org/10.1038/s41580-022-00456-z>.
49. Dunham-Snary, K.J.; Hong, Z.G.; Xiong, P.Y.; Del Paggio, J.C.; Herr, J.E.; Johri, A.M.; Archer, S.L. A mitochondrial redox oxygen sensor in the pulmonary vasculature and ductus arteriosus. *Pflugers Arch.* **2016**, *468*, 43–58. <https://doi.org/10.1007/s00424-015-1736-y>.
50. Smith, K.A.; Schumacker, P.T. Sensors and signals: The role of reactive oxygen species in hypoxic pulmonary vasoconstriction. *J. Physiol.* **2019**, *597*, 1033–1043. <https://doi.org/10.1113/JP275852>.
51. Pak, O.; Nolte, A.; Knoepp, F.; Giordano, L.; Pecina, P.; Huttemann, M.; Grossman, L.I.; Weissmann, N.; Sommer, N. Mitochondrial oxygen sensing of acute hypoxia in specialized cells—Is there a unifying mechanism? *Biochim. Biophys. Acta Bioenerg.* **2022**, *1863*, 148911. <https://doi.org/10.1016/j.bbabi.2022.148911>.
52. Archer, S.L.; Dunham-Snary, K.J.; Bentley, R.; Alizadeh, E.; Weir, E.K. Hypoxic Pulmonary Vasoconstriction: An Important Component of the Homeostatic Oxygen Sensing System. *Physiol. Res.* **2024**, *73*, S493–S510. <https://doi.org/10.33549/physiolres.935431>.
53. Archer, S.L.; Michelakis, E.D.; Thebaud, B.; Bonnet, S.; Moudgil, R.; Wu, X.C.; Weir, E.K. A central role for oxygen-sensitive K⁺ channels and mitochondria in the specialized oxygen-sensing system. *Novartis Found. Symp.* **2006**, *272*, 157–171, discussion 171–175, 214–217.
54. Villamor, E.; Moreno, L.; Mohammed, R.; Perez-Vizcaino, F.; Cogolludo, A. Reactive oxygen species as mediators of oxygen signaling during fetal-to-neonatal circulatory transition. *Free Radic. Biol. Med.* **2019**, *142*, 82–96. <https://doi.org/10.1016/j.freeradbiomed.2019.04.008>.
55. Waypa, G.B.; Schumacker, P.T. O₂ sensing in hypoxic pulmonary vasoconstriction: The mitochondrial door re-opens. *Respir. Physiol. Neurobiol.* **2002**, *132*, 81–91. [https://doi.org/10.1016/s1569-9048\(02\)00051-4](https://doi.org/10.1016/s1569-9048(02)00051-4).
56. Weir, E.K.; Olschewski, A. Role of ion channels in acute and chronic responses of the pulmonary vasculature to hypoxia. *Cardiovasc. Res.* **2006**, *71*, 630–641. <https://doi.org/10.1016/j.cardiores.2006.04.014>.
57. Wolin, M.S.; Gupte, S.A.; Mingone, C.J.; Neo, B.H.; Gao, Q.; Ahmad, M. Redox regulation of responses to hypoxia and NO-cGMP signaling in pulmonary vascular pathophysiology. *Ann. N. Y. Acad. Sci.* **2010**, *1203*, 126–132. <https://doi.org/10.1111/j.1749-6632.2010.05557.x>.
58. Keeley, T.P.; Mann, G.E. Defining Physiological Normoxia for Improved Translation of Cell Physiology to Animal Models and Humans. *Physiol. Rev.* **2019**, *99*, 161–234. <https://doi.org/10.1152/physrev.00041.2017>.
59. Warpsinski, G.; Smith, M.J.; Srivastava, S.; Keeley, T.P.; Siow, R.C.M.; Fraser, P.A.; Mann, G.E. Nrf2-regulated redox signaling in brain endothelial cells adapted to physiological oxygen levels: Consequences for sulforaphane mediated protection against hypoxia-reoxygenation. *Redox Biol.* **2020**, *37*, 101708. <https://doi.org/10.1016/j.redox.2020.101708>.
60. Smith, M.J.; Yang, F.; Griffiths, A.; Morrell, A.; Chapple, S.J.; Siow, R.C.M.; Stewart, T.; Maret, W.; Mann, G.E. Redox and metal profiles in human coronary endothelial and smooth muscle cells under hyperoxia, physiological normoxia and hypoxia: Effects of NRF2 signaling on intracellular zinc. *Redox Biol.* **2023**, *62*, 102712. <https://doi.org/10.1016/j.redox.2023.102712>.
61. Altun, H.Y.; Secilmis, M.; Yang, F.; Akgul Caglar, T.; Vatandaslar, E.; Toy, M.F.; Vilain, S.; Mann, G.E.; Ozturk, G.; Eroglu, E. Visualizing hydrogen peroxide and nitric oxide dynamics in endothelial cells using multispectral imaging under controlled oxygen conditions. *Free Radic. Biol. Med.* **2024**, *221*, 89–97. <https://doi.org/10.1016/j.freeradbiomed.2024.05.021>.
62. Alva, R.; Wiebe, J.E.; Stuart, J.A. Revisiting reactive oxygen species production in hypoxia. *Pflugers Arch.* **2024**, *476*, 1423–1444. <https://doi.org/10.1007/s00424-024-02986-1>.
63. Kamler, M.; Nowak, K.; Bock, M.; Herold, U.; Motsch, J.; Hagl, S.; Gebhard, M.M.; Jakob, H. Bronchial artery revascularization restores peribronchial tissue oxygenation after lung transplantation. *J. Heart Lung Transplant.* **2004**, *23*, 763–766. <https://doi.org/10.1016/j.healun.2003.07.016>.
64. Herold, U.; Jakob, H.; Kamler, M.; Thiele, R.; Tochtermann, U.; Weinmann, J.; Motsch, J.; Gebhard, M.M.; Hagl, S. Interruption of bronchial circulation leads to a severe decrease in peribronchial oxygen tension in standard lung transplantation technique. *Eur. J. Cardiothorac. Surg.* **1998**, *13*, 176–183. [https://doi.org/10.1016/s1010-7940\(97\)00314-x](https://doi.org/10.1016/s1010-7940(97)00314-x).
65. Rivera, B.K.; Naidu, S.K.; Subramanian, K.; Joseph, M.; Hou, H.; Khan, N.; Swartz, H.M.; Kuppusamy, P. Real-time, in vivo determination of dynamic changes in lung and heart tissue oxygenation using EPR oximetry. *Adv. Exp. Med. Biol.* **2014**, *812*, 81–86. https://doi.org/10.1007/978-1-4939-0620-8_11.
66. Marshall, C.; Marshall, B. Site and sensitivity for stimulation of hypoxic pulmonary vasoconstriction. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* **1983**, *55*, 711–716. <https://doi.org/10.1152/jappl.1983.55.3.711>.

67. Grayson, C.; Mailloux, R.J. Coenzyme Q(10) and nicotinamide nucleotide transhydrogenase: Sentinels for mitochondrial hydrogen peroxide signaling. *Free Radic. Biol. Med.* **2023**, *208*, 260–271. <https://doi.org/10.1016/j.freeradbiomed.2023.08.015>.
68. Nisimoto, Y.; Diebold, B.A.; Cosentino-Gomes, D.; Lambeth, J.D. Nox4: A hydrogen peroxide-generating oxygen sensor. *Biochemistry* **2014**, *53*, 5111–5120. <https://doi.org/10.1021/bi500331y>.
69. Knock, G.A. NADPH oxidase in the vasculature: Expression, regulation and signalling pathways; role in normal cardiovascular physiology and its dysregulation in hypertension. *Free Radic. Biol. Med.* **2019**, *145*, 385–427. <https://doi.org/10.1016/j.freeradbiomed.2019.09.029>.
70. Rogers, Z.J.; Flood, D.; Bencherif, S.A.; Taylor, C.T. Oxygen control in cell culture—Your cells may not be experiencing what you think! *Free Radic. Biol. Med.* **2025**, *226*, 279–287. <https://doi.org/10.1016/j.freeradbiomed.2024.11.036>.
71. Peniche Silva, C.J.; Liebsch, G.; Meier, R.J.; Gutbrod, M.S.; Balmayor, E.R.; van Griensven, M. A New Non-invasive Technique for Measuring 3D-Oxygen Gradients in Wells During Mammalian Cell Culture. *Front. Bioeng. Biotechnol.* **2020**, *8*, 595. <https://doi.org/10.3389/fbioe.2020.00595>.
72. Rogers, Z.J.; Colombani, T.; Khan, S.; Bhatt, K.; Nukovic, A.; Zhou, G.; Woolston, B.M.; Taylor, C.T.; Gilkes, D.M.; Slavov, N.; et al. Controlling Pericellular Oxygen Tension in Cell Culture Reveals Distinct Breast Cancer Responses to Low Oxygen Tensions. *Adv. Sci.* **2024**, *11*, e2402557. <https://doi.org/10.1002/advs.202402557>.
73. Ward, J.P. Oxygen sensors in context. *Biochim. Biophys. Acta* **2008**, *1777*, 1–14. <https://doi.org/10.1016/j.bbabi.2007.10.010>.
74. Malconian, M.K.; Rock, P.B.; Reeves, J.T.; Cymerman, A.; Houston, C.S. Operation Everest II: Gas tensions in expired air and arterial blood at extreme altitude. *Aviat. Space Environ. Med.* **1993**, *64*, 37–42.
75. Sutton, J.R.; Reeves, J.T.; Wagner, P.D.; Groves, B.M.; Cymerman, A.; Malconian, M.K.; Rock, P.B.; Young, P.M.; Walter, S.D.; Houston, C.S. Operation Everest II: Oxygen transport during exercise at extreme simulated altitude. *J. Appl. Physiol.* **1988**, *64*, 1309–1321. <https://doi.org/10.1152/jappl.1988.64.4.1309>.
76. West, J.B.; Hackett, P.H.; Maret, K.H.; Milledge, J.S.; Peters, R.M., Jr.; Pizzo, C.J.; Winslow, R.M. Pulmonary gas exchange on the summit of Mount Everest. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* **1983**, *55*, 678–687. <https://doi.org/10.1152/jappl.1983.55.3.678>.
77. Grocott, M.P.; Martin, D.S.; Levett, D.Z.; McMorrow, R.; Windsor, J.; Montgomery, H.E.; Caudwell Xtreme Everest Research, G. Arterial blood gases and oxygen content in climbers on Mount Everest. *N. Engl. J. Med.* **2009**, *360*, 140–149. <https://doi.org/10.1056/NEJMoa0801581>.
78. Kemp, P.J.; Telezhkin, V. Oxygen sensing by the carotid body: Is it all just rotten eggs? *Antioxid. Redox Signal.* **2014**, *20*, 794–804. <https://doi.org/10.1089/ars.2013.5377>.
79. Hawkins, C.L.; Davies, M.J. Detection, identification, and quantification of oxidative protein modifications. *J. Biol. Chem.* **2019**, *294*, 19683–19708. <https://doi.org/10.1074/jbc.REV119.006217>.
80. Sies, H.; Berndt, C.; Jones, D.P. Oxidative Stress. *Annu. Rev. Biochem.* **2017**, *86*, 715–748. <https://doi.org/10.1146/annurev-biochem-061516-045037>.
81. Lipinski, B. Hydroxyl radical and its scavengers in health and disease. *Oxid. Med. Cell. Longev.* **2011**, *2011*, 809696. <https://doi.org/10.1155/2011/809696>.
82. Bienert, G.P.; Moller, A.L.; Kristiansen, K.A.; Schulz, A.; Moller, I.M.; Schjoerring, J.K.; Jahn, T.P. Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J. Biol. Chem.* **2007**, *282*, 1183–1192. <https://doi.org/10.1074/jbc.M603761200>.
83. Pacher, P.; Beckman, J.S.; Liaudet, L. Nitric oxide and peroxynitrite in health and disease. *Physiol. Rev.* **2007**, *87*, 315–424. <https://doi.org/10.1152/physrev.00029.2006>.
84. Sies, H. Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: Oxidative eustress. *Redox Biol.* **2017**, *11*, 613–619. <https://doi.org/10.1016/j.redox.2016.12.035>.
85. Rios, E.J.; Fallon, M.; Wang, J.; Shimoda, L.A. Chronic hypoxia elevates intracellular pH and activates Na⁺/H⁺ exchange in pulmonary arterial smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2005**, *289*, L867–L874. <https://doi.org/10.1152/ajplung.00455.2004>.
86. Xiao, H.; Jedrychowski, M.P.; Schweppe, D.K.; Huttlin, E.L.; Yu, Q.; Heppner, D.E.; Li, J.; Long, J.; Mills, E.L.; Szpyt, J.; et al. A Quantitative Tissue-Specific Landscape of Protein Redox Regulation during Aging. *Cell* **2020**, *180*, 968–983.e924. <https://doi.org/10.1016/j.cell.2020.02.012>.
87. Ulrich, K.; Jakob, U. The role of thiols in antioxidant systems. *Free Radic. Biol. Med.* **2019**, *140*, 14–27. <https://doi.org/10.1016/j.freeradbiomed.2019.05.035>.

88. Salsbury, F.R., Jr.; Knutson, S.T.; Poole, L.B.; Fetrow, J.S. Functional site profiling and electrostatic analysis of cysteines modifiable to cysteine sulfenic acid. *Protein Sci.* **2008**, *17*, 299–312. <https://doi.org/10.1110/ps.073096508>.
89. Mailloux, R.J.; Grayson, C.; Koufos, O. Regulation of Mitochondrial Hydrogen Peroxide Availability by Protein S-glutathionylation. *Cells* **2022**, *12*, 107. <https://doi.org/10.3390/cells12010107>.
90. Burgoyne, J.R.; Mongue-Din, H.; Eaton, P.; Shah, A.M. Redox signaling in cardiac physiology and pathology. *Circ. Res.* **2012**, *111*, 1091–1106. <https://doi.org/10.1161/CIRCRESAHA.111.255216>.
91. Brennan, J.P.; Bardswell, S.C.; Burgoyne, J.R.; Fuller, W.; Schroder, E.; Wait, R.; Begum, S.; Kentish, J.C.; Eaton, P. Oxidant-induced activation of type I protein kinase A is mediated by RI subunit interprotein disulfide bond formation. *J. Biol. Chem.* **2006**, *281*, 21827–21836. <https://doi.org/10.1074/jbc.M603952200>.
92. Cuello, F.; Eaton, P. Cysteine-Based Redox Sensing and Its Role in Signaling by Cyclic Nucleotide-Dependent Kinases in the Cardiovascular System. *Annu. Rev. Physiol.* **2019**, *81*, 63–87. <https://doi.org/10.1146/annurev-physiol-020518-114417>.
93. Kettenhofen, N.J.; Wood, M.J. Formation, reactivity, and detection of protein sulfenic acids. *Chem. Res. Toxicol.* **2010**, *23*, 1633–1646. <https://doi.org/10.1021/tx100237w>.
94. Truong, T.H.; Carroll, K.S. Redox regulation of protein kinases. *Crit. Rev. Biochem. Mol. Biol.* **2013**, *48*, 332–356. <https://doi.org/10.3109/10409238.2013.790873>.
95. Ahsan, M.K.; Lekli, I.; Ray, D.; Yodoi, J.; Das, D.K. Redox regulation of cell survival by the thioredoxin superfamily: An implication of redox gene therapy in the heart. *Antioxid. Redox Signal.* **2009**, *11*, 2741–2758. <https://doi.org/10.1089/ars.2009.2683>.
96. Pillay, C.S.; Rohwer, J.M. Computational models as catalysts for investigating redoxin systems. *Essays Biochem.* **2024**, *68*, 27–39. <https://doi.org/10.1042/EBC20230036>.
97. Lubos, E.; Loscalzo, J.; Handy, D.E. Glutathione peroxidase-1 in health and disease: From molecular mechanisms to therapeutic opportunities. *Antioxid. Redox Signal.* **2011**, *15*, 1957–1997. <https://doi.org/10.1089/ars.2010.3586>.
98. Xiao, W.; Wang, R.S.; Handy, D.E.; Loscalzo, J. NAD(H) and NADP(H) Redox Couples and Cellular Energy Metabolism. *Antioxid. Redox Signal.* **2018**, *28*, 251–272. <https://doi.org/10.1089/ars.2017.7216>.
99. Winterbourn, C.C. Biological Production, Detection, and Fate of Hydrogen Peroxide. *Antioxid. Redox Signal.* **2018**, *29*, 541–551. <https://doi.org/10.1089/ars.2017.7425>.
100. Scherschel, M.; Niemeier, J.O.; Jacobs, L.; Hoffmann, M.D.A.; Diederich, A.; Bell, C.; Hohne, P.; Raetz, S.; Kroll, J.B.; Steinbeck, J.; et al. A family of NADPH/NADP(+) biosensors reveals in vivo dynamics of central redox metabolism across eukaryotes. *Nat. Commun.* **2024**, *15*, 10704. <https://doi.org/10.1038/s41467-024-55302-x>.
101. Gebicka, L.; Krych-Madej, J. The role of catalases in the prevention/promotion of oxidative stress. *J. Inorg. Biochem.* **2019**, *197*, 110699. <https://doi.org/10.1016/j.jinorgbio.2019.110699>.
102. Netto, L.E.; Antunes, F. The Roles of Peroxiredoxin and Thioredoxin in Hydrogen Peroxide Sensing and in Signal Transduction. *Mol. Cells* **2016**, *39*, 65–71. <https://doi.org/10.14348/molcells.2016.2349>.
103. Booth, D.M.; Enyedi, B.; Geiszt, M.; Varnai, P.; Hajnoczky, G. Redox Nanodomains Are Induced by and Control Calcium Signaling at the ER-Mitochondrial Interface. *Mol. Cell* **2016**, *63*, 240–248. <https://doi.org/10.1016/j.molcel.2016.05.040>.
104. Stocker, S.; Van Laer, K.; Mijuskovic, A.; Dick, T.P. The Conundrum of Hydrogen Peroxide Signaling and the Emerging Role of Peroxiredoxins as Redox Relay Hubs. *Antioxid. Redox Signal.* **2018**, *28*, 558–573. <https://doi.org/10.1089/ars.2017.7162>.
105. Sobotta, M.C.; Liou, W.; Stocker, S.; Talwar, D.; Oehler, M.; Ruppert, T.; Scharf, A.N.; Dick, T.P. Peroxiredoxin-2 and STAT3 form a redox relay for H₂O₂ signaling. *Nat. Chem. Biol.* **2015**, *11*, 64–70. <https://doi.org/10.1038/nchembio.1695>.
106. Portillo-Ledesma, S.; Randall, L.M.; Parsonage, D.; Dalla Rizza, J.; Karplus, P.A.; Poole, L.B.; Denicola, A.; Ferrer-Sueta, G. Differential Kinetics of Two-Cysteine Peroxiredoxin Disulfide Formation Reveal a Novel Model for Peroxide Sensing. *Biochemistry* **2018**, *57*, 3416–3424. <https://doi.org/10.1021/acs.biochem.8b00188>.
107. Trujillo, M.; Ferrer-Sueta, G.; Radi, R. Kinetic studies on peroxynitrite reduction by peroxiredoxins. *Methods Enzymol.* **2008**, *441*, 173–196. [https://doi.org/10.1016/S0076-6879\(08\)01210-X](https://doi.org/10.1016/S0076-6879(08)01210-X).
108. Rhee, S.G.; Woo, H.A. Multiple functions of 2-Cys peroxiredoxins, I and II, and their regulations via post-translational modifications. *Free Radic. Biol. Med.* **2020**, *152*, 107–115. <https://doi.org/10.1016/j.freeradbiomed.2020.02.028>.
109. van Dam, L.; Pages-Gallego, M.; Polderman, P.E.; van Es, R.M.; Burgering, B.M.T.; Vos, H.R.; Dansen, T.B. The Human 2-Cys Peroxiredoxins form Widespread, Cysteine-Dependent- and Isoform-Specific Protein-Protein Interactions. *Antioxidants* **2021**, *10*, 627. <https://doi.org/10.3390/antiox10040627>.
110. Mailloux, R.J.; Treberg, J.R. Protein S-glutathionylation links energy metabolism to redox signaling in mitochondria. *Redox Biol.* **2016**, *8*, 110–118. <https://doi.org/10.1016/j.redox.2015.12.010>.

111. Sies, H.; Jones, D.P. Reactive oxygen species (ROS) as pleiotropic physiological signalling agents. *Nat. Rev. Mol. Cell. Biol.* **2020**, *21*, 363–383. <https://doi.org/10.1038/s41580-020-0230-3>.
112. Chouchani, E.T.; Pell, V.R.; Gaude, E.; Aksentijevic, D.; Sundier, S.Y.; Robb, E.L.; Logan, A.; Nadtochiy, S.M.; Ord, E.N.J.; Smith, A.C.; et al. Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* **2014**, *515*, 431–435. <https://doi.org/10.1038/nature13909>.
113. Pelletier, M.; Lepow, T.S.; Billingham, L.K.; Murphy, M.P.; Siegel, R.M. New tricks from an old dog: Mitochondrial redox signaling in cellular inflammation. *Semin. Immunol.* **2012**, *24*, 384–392. <https://doi.org/10.1016/j.smim.2013.01.002>.
114. Starkov, A.A.; Andreyev, A.Y.; Zhang, S.F.; Starkova, N.N.; Korneeva, M.; Syromyatnikov, M.; Popov, V.N. Scavenging of H₂O₂ by mouse brain mitochondria. *J. Bioenerg. Biomembr.* **2014**, *46*, 471–477. <https://doi.org/10.1007/s10863-014-9581-9>.
115. Treberg, J.R.; Munro, D.; Banh, S.; Zacharias, P.; Sotiri, E. Differentiating between apparent and actual rates of H₂O₂ metabolism by isolated rat muscle mitochondria to test a simple model of mitochondria as regulators of H₂O₂ concentration. *Redox Biol.* **2015**, *5*, 216–224. <https://doi.org/10.1016/j.redox.2015.05.001>.
116. Dey, S.; Sidor, A.; O'Rourke, B. Compartment-specific Control of Reactive Oxygen Species Scavenging by Antioxidant Pathway Enzymes. *J. Biol. Chem.* **2016**, *291*, 11185–11197. <https://doi.org/10.1074/jbc.M116.726968>.
117. Andreyev, A.Y.; Kushnareva, Y.E.; Starkov, A.A. Mitochondrial metabolism of reactive oxygen species. *Biochemistry* **2005**, *70*, 200–214. <https://doi.org/10.1007/s10541-005-0102-7>.
118. Murphy, M.P. How mitochondria produce reactive oxygen species. *Biochem. J.* **2009**, *417*, 1–13. <https://doi.org/10.1042/BJ20081386>.
119. Droese, S.; Brandt, U.; Wittig, I. Mitochondrial respiratory chain complexes as sources and targets of thiol-based redox-regulation. *Biochim. Biophys. Acta* **2014**, *1844*, 1344–1354. <https://doi.org/10.1016/j.bbapap.2014.02.006>.
120. Andreyev, A.Y.; Kushnareva, Y.E.; Murphy, A.N.; Starkov, A.A. Mitochondrial ROS Metabolism: 10 Years Later. *Biochemistry* **2015**, *80*, 517–531. <https://doi.org/10.1134/S0006297915050028>.
121. Brand, M.D. Mitochondrial generation of superoxide and hydrogen peroxide as the source of mitochondrial redox signaling. *Free Radic. Biol. Med.* **2016**, *100*, 14–31. <https://doi.org/10.1016/j.freeradbiomed.2016.04.001>.
122. Hernansanz-Agustin, P.; Morales-Vidal, C.; Calvo, E.; Natale, P.; Marti-Mateos, Y.; Jaroszewicz, S.N.; Cabrera-Alarcon, J.L.; Acin-Perez, R.; Lopez-Montero, I.; Vazquez, J.; et al. A trans-mitochondrial sodium gradient controls membrane potential in mammalian mitochondria. *Cell* **2024**, *187*, 6599–6613. <https://doi.org/10.1016/j.cell.2024.08.045>.
123. Nicholls, D.G.; Ferguson, S.J. *Bioenergetics* 3, 3rd ed.; Academic Press: San Diego, CA, USA, 2002; p. xviii; 297p.
124. Gauthier, L.D.; Greenstein, J.L.; Cortassa, S.; O'Rourke, B.; Winslow, R.L. A computational model of reactive oxygen species and redox balance in cardiac mitochondria. *Biophys. J.* **2013**, *105*, 1045–1056. <https://doi.org/10.1016/j.bpj.2013.07.006>.
125. Castro, L.; Tortora, V.; Mansilla, S.; Radi, R. Aconitases: Non-redox Iron-Sulfur Proteins Sensitive to Reactive Species. *Acc. Chem. Res.* **2019**, *52*, 2609–2619. <https://doi.org/10.1021/acs.accounts.9b00150>.
126. Tahara, E.B.; Navarete, F.D.; Kowaltowski, A.J. Tissue-, substrate-, and site-specific characteristics of mitochondrial reactive oxygen species generation. *Free Radic. Biol. Med.* **2009**, *46*, 1283–1297. <https://doi.org/10.1016/j.freeradbiomed.2009.02.008>.
127. Hoffman, D.L.; Brookes, P.S. Oxygen sensitivity of mitochondrial reactive oxygen species generation depends on metabolic conditions. *J. Biol. Chem.* **2009**, *284*, 16236–16245. <https://doi.org/10.1074/jbc.M809512200>.
128. Balmaceda, V.; Komlodi, T.; Szibor, M.; Gnaiger, E.; Moore, A.L.; Fernandez-Vizarra, E.; Viscomi, C. The striking differences in the bioenergetics of brain and liver mitochondria are enhanced in mitochondrial disease. *Biochim. Biophys. Acta Mol. Basis Dis.* **2024**, *1870*, 167033. <https://doi.org/10.1016/j.bbadis.2024.167033>.
129. Slade, L.; Chalker, J.; Kuksal, N.; Young, A.; Gardiner, D.; Mailloux, R.J. Examination of the superoxide/hydrogen peroxide forming and quenching potential of mouse liver mitochondria. *Biochim. Biophys. Acta Gen. Subj.* **2017**, *1861*, 1960–1969. <https://doi.org/10.1016/j.bbagen.2017.05.010>.
130. Wong, H.S.; Benoit, B.; Brand, M.D. Mitochondrial and cytosolic sources of hydrogen peroxide in resting C2C12 myoblasts. *Free Radic. Biol. Med.* **2019**, *130*, 140–150. <https://doi.org/10.1016/j.freeradbiomed.2018.10.448>.
131. Chandel, N.S. Mitochondria. *Cold Spring Harb. Perspect. Biol.* **2021**, *13*, a040543. <https://doi.org/10.1101/cshperspect.a040543>.
132. Han, D.; Canali, R.; Rettori, D.; Kaplowitz, N. Effect of glutathione depletion on sites and topology of superoxide and hydrogen peroxide production in mitochondria. *Mol. Pharmacol.* **2003**, *64*, 1136–1144. <https://doi.org/10.1124/mol.64.5.1136>.
133. Hoehne, M.N.; Jacobs, L.; Lapacz, K.J.; Calabrese, G.; Murschall, L.M.; Marker, T.; Kaul, H.; Trifunovic, A.; Morgan, B.; Fricker, M.; et al. Spatial and temporal control of mitochondrial H₂O₂ release in intact human cells. *EMBO J.* **2022**, *41*, e109169. <https://doi.org/10.15252/embj.2021109169>.

134. Goncalves, R.L.S.; Watson, M.A.; Wong, H.S.; Orr, A.L.; Brand, M.D. The use of site-specific suppressors to measure the relative contributions of different mitochondrial sites to skeletal muscle superoxide and hydrogen peroxide production. *Redox Biol.* **2020**, *28*, 101341. <https://doi.org/10.1016/j.redox.2019.101341>.
135. Bleier, L.; Wittig, I.; Heide, H.; Steger, M.; Brandt, U.; Drose, S. Generator-specific targets of mitochondrial reactive oxygen species. *Free Radic. Biol. Med.* **2015**, *78*, 1–10. <https://doi.org/10.1016/j.freeradbiomed.2014.10.511>.
136. Gao, L.; Gonzalez-Rodriguez, P.; Ortega-Saenz, P.; Lopez-Barneo, J. Redox signaling in acute oxygen sensing. *Redox Biol.* **2017**, *12*, 908–915. <https://doi.org/10.1016/j.redox.2017.04.033>.
137. Dunham-Snary, K.J.; Wu, D.; Potus, F.; Sykes, E.A.; Mewburn, J.D.; Charles, R.L.; Eaton, P.; Sultanian, R.A.; Archer, S.L. Ndufs2, a Core Subunit of Mitochondrial Complex I, Is Essential for Acute Oxygen-Sensing and Hypoxic Pulmonary Vasoconstriction. *Circ. Res.* **2019**, *124*, 1727–1746. <https://doi.org/10.1161/CIRCRESAHA.118.314284>.
138. Paddenberg, R.; Ishaq, B.; Goldenberg, A.; Faulhammer, P.; Rose, F.; Weissmann, N.; Braun-Dullaeus, R.C.; Kummer, W. Essential role of complex II of the respiratory chain in hypoxia-induced ROS generation in the pulmonary vasculature. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2003**, *284*, L710–L719. <https://doi.org/10.1152/ajplung.00149.2002>.
139. Korde, A.S.; Yadav, V.R.; Zheng, Y.M.; Wang, Y.X. Primary role of mitochondrial Rieske iron-sulfur protein in hypoxic ROS production in pulmonary artery myocytes. *Free Radic. Biol. Med.* **2011**, *50*, 945–952. <https://doi.org/10.1016/j.freeradbiomed.2011.01.010>.
140. Waypa, G.B.; Marks, J.D.; Guzy, R.D.; Mungai, P.T.; Schriewer, J.M.; Dokic, D.; Ball, M.K.; Schumacker, P.T. Superoxide generated at mitochondrial complex III triggers acute responses to hypoxia in the pulmonary circulation. *Am. J. Respir. Crit. Care Med.* **2013**, *187*, 424–432. <https://doi.org/10.1164/rccm.201207-1294OC>.
141. Sommer, N.; Alebrahimdehkhordi, N.; Pak, O.; Knoepp, F.; Strielkov, I.; Scheibe, S.; Dufour, E.; Andjelkovic, A.; Sydykov, A.; Saraji, A.; et al. Bypassing mitochondrial complex III using alternative oxidase inhibits acute pulmonary oxygen sensing. *Sci. Adv.* **2020**, *6*, eaba0694. <https://doi.org/10.1126/sciadv.aba0694>.
142. Sommer, N.; Huttemann, M.; Pak, O.; Scheibe, S.; Knoepp, F.; Sinkler, C.; Malczyk, M.; Gierhardt, M.; Esfandiary, A.; Kraut, S.; et al. Mitochondrial Complex IV Subunit 4 Isoform 2 Is Essential for Acute Pulmonary Oxygen Sensing. *Circ. Res.* **2017**, *121*, 424–438. <https://doi.org/10.1161/CIRCRESAHA.116.310482>.
143. Gibbs, E.T.; Lerner, C.A.; Watson, M.A.; Wong, H.S.; Gerencser, A.A.; Brand, M.D. Site IQ in mitochondrial complex I generates S1QEL-sensitive superoxide/hydrogen peroxide in both the reverse and forward reactions. *Biochem. J.* **2023**, *480*, 363–384. <https://doi.org/10.1042/BCJ20220611>.
144. Bleier, L.; Drose, S. Superoxide generation by complex III: From mechanistic rationales to functional consequences. *Biochim. Biophys. Acta* **2013**, *1827*, 1320–1331. <https://doi.org/10.1016/j.bbabbio.2012.12.002>.
145. Guillaud, F.; Drose, S.; Kowald, A.; Brandt, U.; Klipp, E. Superoxide production by cytochrome bc1 complex: A mathematical model. *Biochim. Biophys. Acta* **2014**, *1837*, 1643–1652. <https://doi.org/10.1016/j.bbabbio.2014.05.358>.
146. Sarewicz, M.; Osyczka, A. Electronic connection between the quinone and cytochrome C redox pools and its role in regulation of mitochondrial electron transport and redox signaling. *Physiol. Rev.* **2015**, *95*, 219–243. <https://doi.org/10.1152/physrev.00006.2014>.
147. Rottenberg, H.; Covian, R.; Trumpower, B.L. Membrane potential greatly enhances superoxide generation by the cytochrome bc1 complex reconstituted into phospholipid vesicles. *J. Biol. Chem.* **2009**, *284*, 19203–19210. <https://doi.org/10.1074/jbc.M109.017376>.
148. Markevich, N.I.; Hoek, J.B. Computational modeling analysis of mitochondrial superoxide production under varying substrate conditions and upon inhibition of different segments of the electron transport chain. *Biochim. Biophys. Acta* **2015**, *1847*, 656–679. <https://doi.org/10.1016/j.bbabbio.2015.04.005>.
149. Quinlan, C.L.; Gerencser, A.A.; Treberg, J.R.; Brand, M.D. The mechanism of superoxide production by the antimycin-inhibited mitochondrial Q-cycle. *J. Biol. Chem.* **2011**, *286*, 31361–31372. <https://doi.org/10.1074/jbc.M111.267898>.
150. Tsubaki, M. Fourier-transform infrared study of cyanide binding to the Fea3-CuB binuclear site of bovine heart cytochrome c oxidase: Implication of the redox-linked conformational change at the binuclear site. *Biochemistry* **1993**, *32*, 164–173. <https://doi.org/10.1021/bi00052a022>.
151. Biscoe, T.J.; Duchon, M.R. Cellular basis of transduction in carotid chemoreceptors. *Am. J. Physiol.* **1990**, *258*, L271–L278. <https://doi.org/10.1152/ajplung.1990.258.6.L271>.
152. Wilson, D.F. Oxidative phosphorylation: Regulation and role in cellular and tissue metabolism. *J. Physiol.* **2017**, *595*, 7023–7038. <https://doi.org/10.1113/JP273839>.

153. Buckler, K.J.; Turner, P.J. Functional Properties of Mitochondria in the Type-1 Cell and Their Role in Oxygen Sensing. *Adv. Exp. Med. Biol.* **2015**, *860*, 69–80. https://doi.org/10.1007/978-3-319-18440-1_9.
154. Wilson, D.F.; Rumsey, W.L.; Green, T.J.; Vanderkooi, J.M. The oxygen dependence of mitochondrial oxidative phosphorylation measured by a new optical method for measuring oxygen concentration. *J. Biol. Chem.* **1988**, *263*, 2712–2718.
155. Sommer, N.; Pak, O.; Schorner, S.; Derfuss, T.; Krug, A.; Gnaiger, E.; Ghofrani, H.A.; Schermuly, R.T.; Huckstorf, C.; Seeger, W.; et al. Mitochondrial cytochrome redox states and respiration in acute pulmonary oxygen sensing. *Eur. Respir. J.* **2010**, *36*, 1056–1066. <https://doi.org/10.1183/09031936.00013809>.
156. Michelakis, E.D.; Hampl, V.; Nsair, A.; Wu, X.; Harry, G.; Haromy, A.; Gurtu, R.; Archer, S.L. Diversity in mitochondrial function explains differences in vascular oxygen sensing. *Circ. Res.* **2002**, *90*, 1307–1315. <https://doi.org/10.1161/01.res.0000024689.07590.c2>.
157. Roy, A.; Li, J.; Al-Mehdi, A.B.; Mokashi, A.; Lahiri, S. Effect of acute hypoxia on glomus cell Em and psi m as measured by fluorescence imaging. *J. Appl. Physiol.* **2002**, *93*, 1987–1998. <https://doi.org/10.1152/japplphysiol.00725.2001>.
158. Kurokawa, H.; Ito, H.; Inoue, M.; Tabata, K.; Sato, Y.; Yamagata, K.; Kizaka-Kondoh, S.; Kadonosono, T.; Yano, S.; Inoue, M.; et al. High resolution imaging of intracellular oxygen concentration by phosphorescence lifetime. *Sci. Rep.* **2015**, *5*, 10657. <https://doi.org/10.1038/srep10657>.
159. Gnaiger, E.; Steinlechner-Maran, R.; Mendez, G.; Eberl, T.; Margreiter, R. Control of mitochondrial and cellular respiration by oxygen. *J. Bioenerg. Biomembr.* **1995**, *27*, 583–596. <https://doi.org/10.1007/BF02111656>.
160. Gnaiger, E.; Lassnig, B.; Kuznetsov, A.; Rieger, G.; Margreiter, R. Mitochondrial oxygen affinity, respiratory flux control and excess capacity of cytochrome c oxidase. *J. Exp. Biol.* **1998**, *201*, 1129–1139. <https://doi.org/10.1242/jeb.201.8.1129>.
161. Brown, G.C.; Cooper, C.E. Nanomolar concentrations of nitric oxide reversibly inhibit synaptosomal respiration by competing with oxygen at cytochrome oxidase. *FEBS Lett.* **1994**, *356*, 295–298. [https://doi.org/10.1016/0014-5793\(94\)01290-3](https://doi.org/10.1016/0014-5793(94)01290-3).
162. Palacios-Callender, M.; Quintero, M.; Hollis, V.S.; Springett, R.J.; Moncada, S. Endogenous NO regulates superoxide production at low oxygen concentrations by modifying the redox state of cytochrome c oxidase. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 7630–7635. <https://doi.org/10.1073/pnas.0401723101>.
163. Edmunds, N.J.; Moncada, S.; Marshall, J.M. Does nitric oxide allow endothelial cells to sense hypoxia and mediate hypoxic vasodilatation? In vivo and in vitro studies. *J. Physiol.* **2003**, *546*, 521–527. <https://doi.org/10.1113/jphysiol.2002.023663>.
164. Quintero, M.; Colombo, S.L.; Godfrey, A.; Moncada, S. Mitochondria as signaling organelles in the vascular endothelium. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 5379–5384. <https://doi.org/10.1073/pnas.0601026103>.
165. Galkin, A.; Higgs, A.; Moncada, S. Nitric oxide and hypoxia. *Essays Biochem.* **2007**, *43*, 29–42. <https://doi.org/10.1042/BSE0430029>.
166. Fagan, K.A.; Tyler, R.C.; Sato, K.; Fouty, B.W.; Morris, K.G., Jr.; Huang, P.L.; McMurtry, I.F.; Rodman, D.M. Relative contributions of endothelial, inducible, and neuronal NOS to tone in the murine pulmonary circulation. *Am. J. Physiol.* **1999**, *277*, L472–L478. <https://doi.org/10.1152/ajplung.1999.277.3.L472>.
167. Leeman, M.; de Beyl, V.Z.; Delcroix, M.; Naeije, R. Effects of endogenous nitric oxide on pulmonary vascular tone in intact dogs. *Am. J. Physiol.* **1994**, *266*, H2343–H2347. <https://doi.org/10.1152/ajpheart.1994.266.6.H2343>.
168. Persson, M.G.; Gustafsson, L.E.; Wiklund, N.P.; Moncada, S.; Hedqvist, P. Endogenous nitric oxide as a probable modulator of pulmonary circulation and hypoxic pressor response in vivo. *Acta Physiol. Scand.* **1990**, *140*, 449–457. <https://doi.org/10.1111/j.1748-1716.1990.tb09021.x>.
169. Vaughan, D.J.; Brogan, T.V.; Kerr, M.E.; Deem, S.; Luchtel, D.L.; Swenson, E.R. Contributions of nitric oxide synthase isozymes to exhaled nitric oxide and hypoxic pulmonary vasoconstriction in rabbit lungs. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2003**, *284*, L834–L843. <https://doi.org/10.1152/ajplung.00341.2002>.
170. Blitzer, M.L.; Loh, E.; Roddy, M.A.; Stamler, J.S.; Creager, M.A. Endothelium-derived nitric oxide regulates systemic and pulmonary vascular resistance during acute hypoxia in humans. *J. Am. Coll. Cardiol.* **1996**, *28*, 591–596. [https://doi.org/10.1016/0735-1097\(96\)00218-5](https://doi.org/10.1016/0735-1097(96)00218-5).
171. Leach, R.M.; Robertson, T.P.; Twort, C.H.; Ward, J.P. Hypoxic vasoconstriction in rat pulmonary and mesenteric arteries. *Am. J. Physiol.* **1994**, *266*, L223–L231. <https://doi.org/10.1152/ajplung.1994.266.3.L223>.
172. Castello, P.R.; David, P.S.; McClure, T.; Crook, Z.; Poyton, R.O. Mitochondrial cytochrome oxidase produces nitric oxide under hypoxic conditions: Implications for oxygen sensing and hypoxic signaling in eukaryotes. *Cell Metab.* **2006**, *3*, 277–287. <https://doi.org/10.1016/j.cmet.2006.02.011>.
173. Palacios-Callender, M.; Hollis, V.; Mitchison, M.; Frakich, N.; Unitt, D.; Moncada, S. Cytochrome c oxidase regulates endogenous nitric oxide availability in respiring cells: A possible explanation for hypoxic vasodilation. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 18508–18513. <https://doi.org/10.1073/pnas.0709440104>.

174. Amdahl, M.B.; DeMartino, A.W.; Gladwin, M.T. Inorganic nitrite bioactivation and role in physiological signaling and therapeutics. *Biol. Chem.* **2019**, *401*, 201–211. <https://doi.org/10.1515/hsz-2019-0349>.
175. Olson, K.R.; DeLeon, E.R.; Gao, Y.; Hurley, K.; Sadauskas, V.; Batz, C.; Stoy, G.F. Thiosulfate: A readily accessible source of hydrogen sulfide in oxygen sensing. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2013**, *305*, R592–R603. <https://doi.org/10.1152/ajpregu.00421.2012>.
176. Petersen, L.C. The effect of inhibitors on the oxygen kinetics of cytochrome c oxidase. *Biochim. Biophys. Acta* **1977**, *460*, 299–307. [https://doi.org/10.1016/0005-2728\(77\)90216-x](https://doi.org/10.1016/0005-2728(77)90216-x).
177. Prieto-Lloret, J.; Snetkov, V.A.; Shaifta, Y.; Docio, I.; Connolly, M.J.; MacKay, C.E.; Knock, G.A.; Ward, J.P.T.; Aaronson, P.I. Role of reactive oxygen species and sulfide-quinone oxoreductase in hydrogen sulfide-induced contraction of rat pulmonary arteries. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2018**, *314*, L670–L685. <https://doi.org/10.1152/ajplung.00283.2016>.
178. Szabo, C.; Ransy, C.; Modis, K.; Andriamihaja, M.; Murghes, B.; Coletta, C.; Olah, G.; Yanagi, K.; Bouillaud, F. Regulation of mitochondrial bioenergetic function by hydrogen sulfide. Part I. Biochemical and physiological mechanisms. *Br. J. Pharmacol.* **2014**, *171*, 2099–2122. <https://doi.org/10.1111/bph.12369>.
179. Nicholson, R.A.; Roth, S.H.; Zhang, A.; Zheng, J.; Brookes, J.; Skrajny, B.; Bennington, R. Inhibition of respiratory and bioenergetic mechanisms by hydrogen sulfide in mammalian brain. *J. Toxicol. Environ. Health A* **1998**, *54*, 491–507. <https://doi.org/10.1080/009841098158773>.
180. Levitt, M.D.; Abdel-Rehim, M.S.; Furne, J. Free and acid-labile hydrogen sulfide concentrations in mouse tissues: Anomalous high free hydrogen sulfide in aortic tissue. *Antioxid. Redox Signal.* **2011**, *15*, 373–378. <https://doi.org/10.1089/ars.2010.3525>.
181. Wilson, D.F.; Harrison, D.K.; Vinogradov, A. Mitochondrial cytochrome c oxidase and control of energy metabolism: Measurements in suspensions of isolated mitochondria. *J. Appl. Physiol.* **2014**, *117*, 1424–1430. <https://doi.org/10.1152/japplphysiol.00736.2014>.
182. Harrison, D.K.; Fasching, M.; Fontana-Ayoub, M.; Gnaiger, E. Cytochrome redox states and respiratory control in mouse and beef heart mitochondria at steady-state levels of hypoxia. *J. Appl. Physiol.* **2015**, *119*, 1210–1218. <https://doi.org/10.1152/japplphysiol.00146.2015>.
183. Wilson, D.F.; Erecinska, M.; Drown, C.; Silver, I.A. Effect of oxygen tension on cellular energetics. *Am. J. Physiol.* **1977**, *233*, C135–C140. <https://doi.org/10.1152/ajpcell.1977.233.5.C135>.
184. Wilson, D.F.; Erecinska, M.; Drown, C.; Silver, I.A. The oxygen dependence of cellular energy metabolism. *Arch. Biochem. Biophys.* **1979**, *195*, 485–493. [https://doi.org/10.1016/0003-9861\(79\)90375-8](https://doi.org/10.1016/0003-9861(79)90375-8).
185. Pajuelo Reguera, D.; Cunatova, K.; Vrbacky, M.; Pecinova, A.; Houstek, J.; Mracek, T.; Pecina, P. Cytochrome c Oxidase Subunit 4 Isoform Exchange Results in Modulation of Oxygen Affinity. *Cells* **2020**, *9*, 443. <https://doi.org/10.3390/cells9020443>.
186. Huttemann, M.; Kadenbach, B.; Grossman, L.I. Mammalian subunit IV isoforms of cytochrome c oxidase. *Gene* **2001**, *267*, 111–123. [https://doi.org/10.1016/s0378-1119\(01\)00385-7](https://doi.org/10.1016/s0378-1119(01)00385-7).
187. Buckler, K.J.; Turner, P.J. Oxygen sensitivity of mitochondrial function in rat arterial chemoreceptor cells. *J. Physiol.* **2013**, *591*, 3549–3563. <https://doi.org/10.1113/jphysiol.2013.257741>.
188. Scialo, F.; Fernandez-Ayala, D.J.; Sanz, A. Role of Mitochondrial Reverse Electron Transport in ROS Signaling: Potential Roles in Health and Disease. *Front. Physiol.* **2017**, *8*, 428. <https://doi.org/10.3389/fphys.2017.00428>.
189. Treberg, J.R.; Quinlan, C.L.; Brand, M.D. Evidence for two sites of superoxide production by mitochondrial NADH-ubiquinone oxidoreductase (complex I). *J. Biol. Chem.* **2011**, *286*, 27103–27110. <https://doi.org/10.1074/jbc.M111.252502>.
190. Fernandez-Aguera, M.C.; Gao, L.; Gonzalez-Rodriguez, P.; Pintado, C.O.; Arias-Mayenco, I.; Garcia-Flores, P.; Garcia-Perganeda, A.; Pascual, A.; Ortega-Saenz, P.; Lopez-Barneo, J. Oxygen Sensing by Arterial Chemoreceptors Depends on Mitochondrial Complex I Signaling. *Cell Metab.* **2015**, *22*, 825–837. <https://doi.org/10.1016/j.cmet.2015.09.004>.
191. Arias-Mayenco, I.; Gonzalez-Rodriguez, P.; Torres-Torrel, H.; Gao, L.; Fernandez-Aguera, M.C.; Bonilla-Henao, V.; Ortega-Saenz, P.; Lopez-Barneo, J. Acute O₂ Sensing: Role of Coenzyme QH₂/Q Ratio and Mitochondrial ROS Compartmentalization. *Cell Metab.* **2018**, *28*, 145–158.e144. <https://doi.org/10.1016/j.cmet.2018.05.009>.
192. Swiderska, A.; Coney, A.M.; Alzahrani, A.A.; Aldossary, H.S.; Batis, N.; Ray, C.J.; Kumar, P.; Holmes, A.P. Mitochondrial Succinate Metabolism and Reactive Oxygen Species Are Important but Not Essential for Eliciting Carotid Body and Ventilatory Responses to Hypoxia in the Rat. *Antioxidants* **2021**, *10*, 840. <https://doi.org/10.3390/antiox10060840>.
193. Torres-Lopez, M.; Spiller, P.F.; Gao, L.; Garcia-Flores, P.; Murphy, M.P.; Ortega-Saenz, P.; Lopez-Barneo, J. Acute oxygen sensing by arterial chemoreceptors with a mutant mitochondrial complex I ND6 subunit lacking reverse electron transport. *FEBS Lett.* **2025**, *599*, 1122–1134. <https://doi.org/10.1002/1873-3468.70017>.

194. Hernansanz-Agustin, P.; Choya-Foces, C.; Carregal-Romero, S.; Ramos, E.; Oliva, T.; Villa-Pina, T.; Moreno, L.; Izquierdo-Alvarez, A.; Cabrera-Garcia, J.D.; Cortes, A.; et al. Na⁺ controls hypoxic signalling by the mitochondrial respiratory chain. *Nature* **2020**, *586*, 287–291. <https://doi.org/10.1038/s41586-020-2551-y>.
195. Hernansanz-Agustin, P.; Ramos, E.; Navarro, E.; Parada, E.; Sanchez-Lopez, N.; Pelaez-Aguado, L.; Cabrera-Garcia, J.D.; Tello, D.; Buendia, I.; Marina, A.; et al. Mitochondrial complex I deactivation is related to superoxide production in acute hypoxia. *Redox Biol.* **2017**, *12*, 1040–1051. <https://doi.org/10.1016/j.redox.2017.04.025>.
196. Vinogradov, A.D. Catalytic properties of the mitochondrial NADH-ubiquinone oxidoreductase (complex I) and the pseudo-reversible active/inactive enzyme transition. *Biochim. Biophys. Acta* **1998**, *1364*, 169–185. [https://doi.org/10.1016/s0005-2728\(98\)00026-7](https://doi.org/10.1016/s0005-2728(98)00026-7).
197. Roberts, P.G.; Hirst, J. The deactive form of respiratory complex I from mammalian mitochondria is a Na⁺/H⁺ antiporter. *J. Biol. Chem.* **2012**, *287*, 34743–34751. <https://doi.org/10.1074/jbc.M112.384560>.
198. Babot, M.; Birch, A.; Labarbuta, P.; Galkin, A. Characterisation of the active/de-active transition of mitochondrial complex I. *Biochim. Biophys. Acta* **2014**, *1837*, 1083–1092. <https://doi.org/10.1016/j.bbabo.2014.02.018>.
199. Babot, M.; Galkin, A. Molecular mechanism and physiological role of active-deactive transition of mitochondrial complex I. *Biochem. Soc. Trans.* **2013**, *41*, 1325–1330. <https://doi.org/10.1042/BST20130088>.
200. Maklashina, E.; Kotlyar, A.B.; Karliner, J.S.; Cecchini, G. Effect of oxygen on activation state of complex I and lack of oxaloacetate inhibition of complex II in Langendorff perfused rat heart. *FEBS Lett.* **2004**, *556*, 64–68. [https://doi.org/10.1016/s0014-5793\(03\)01369-3](https://doi.org/10.1016/s0014-5793(03)01369-3).
201. Chalmers, S.; Nicholls, D.G. The relationship between free and total calcium concentrations in the matrix of liver and brain mitochondria. *J. Biol. Chem.* **2003**, *278*, 19062–19070. <https://doi.org/10.1074/jbc.M212661200>.
202. Gherardi, G.; Monticelli, H.; Rizzuto, R.; Mammucari, C. The Mitochondrial Ca²⁺ Uptake and the Fine-Tuning of Aerobic Metabolism. *Front. Physiol.* **2020**, *11*, 554904. <https://doi.org/10.3389/fphys.2020.554904>.
203. Wescott, A.P.; Kao, J.P.Y.; Lederer, W.J.; Boyman, L. Voltage-energized Calcium-sensitive ATP Production by Mitochondria. *Nat. Metab.* **2019**, *1*, 975–984. <https://doi.org/10.1038/s42255-019-0126-8>.
204. Sena, L.A.; Li, S.; Jairaman, A.; Prakriya, M.; Ezponda, T.; Hildeman, D.A.; Wang, C.R.; Schumacker, P.T.; Licht, J.D.; Perlman, H.; et al. Mitochondria are required for antigen-specific T cell activation through reactive oxygen species signaling. *Immunity* **2013**, *38*, 225–236. <https://doi.org/10.1016/j.immuni.2012.10.020>.
205. Lee, I.; Bender, E.; Arnold, S.; Kadenbach, B. New control of mitochondrial membrane potential and ROS formation—A hypothesis. *Biol. Chem.* **2001**, *382*, 1629–1636. <https://doi.org/10.1515/BC.2001.198>.
206. Starkov, A.A.; Fiskum, G. Regulation of brain mitochondrial H₂O₂ production by membrane potential and NAD(P)H redox state. *J. Neurochem.* **2003**, *86*, 1101–1107. <https://doi.org/10.1046/j.1471-4159.2003.01908.x>.
207. Ramzan, R.; Vogt, S.; Kadenbach, B. Stress-mediated generation of deleterious ROS in healthy individuals—Role of cytochrome c oxidase. *J. Mol. Med.* **2020**, *98*, 651–657. <https://doi.org/10.1007/s00109-020-01905-y>.
208. Bertero, E.; Maack, C. Calcium Signaling and Reactive Oxygen Species in Mitochondria. *Circ. Res.* **2018**, *122*, 1460–1478. <https://doi.org/10.1161/CIRCRESAHA.118.310082>.
209. Duong, Q.V.; Hoffman, A.; Zhong, K.; Dessinger, M.J.; Zhang, Y.; Bazil, J.N. Calcium overload decreases net free radical emission in cardiac mitochondria. *Mitochondrion* **2020**, *51*, 126–139. <https://doi.org/10.1016/j.mito.2020.01.005>.
210. Grivennikova, V.G.; Kareyeva, A.V.; Vinogradov, A.D. Oxygen-dependence of mitochondrial ROS production as detected by Amplex Red assay. *Redox Biol.* **2018**, *17*, 192–199. <https://doi.org/10.1016/j.redox.2018.04.014>.
211. Stepanova, A.; Konrad, C.; Manfredi, G.; Springett, R.; Ten, V.; Galkin, A. The dependence of brain mitochondria reactive oxygen species production on oxygen level is linear, except when inhibited by antimycin A. *J. Neurochem.* **2019**, *148*, 731–745. <https://doi.org/10.1111/jnc.14654>.
212. Hoffman, D.L.; Salter, J.D.; Brookes, P.S. Response of mitochondrial reactive oxygen species generation to steady-state oxygen tension: Implications for hypoxic cell signaling. *Am. J. Physiol. Heart Circ. Physiol.* **2007**, *292*, H101–H108. <https://doi.org/10.1152/ajpheart.00699.2006>.
213. Wu, D.; Dasgupta, A.; Read, A.D.; Bentley, R.E.T.; Motamed, M.; Chen, K.H.; Al-Qazazi, R.; Mewburn, J.D.; Dunham-Snary, K.J.; Alizadeh, E.; et al. Oxygen sensing, mitochondrial biology and experimental therapeutics for pulmonary hypertension and cancer. *Free Radic. Biol. Med.* **2021**, *170*, 150–178. <https://doi.org/10.1016/j.freeradbiomed.2020.12.452>.
214. Weir, E.K.; Archer, S.L. The role of redox changes in oxygen sensing. *Respir. Physiol. Neurobiol.* **2010**, *174*, 182–191. <https://doi.org/10.1016/j.resp.2010.08.015>.

215. Genova, M.L.; Pich, M.M.; Bernacchia, A.; Bianchi, C.; Biondi, A.; Bovina, C.; Falasca, A.I.; Formiggini, G.; Castelli, G.P.; Lenaz, G. The mitochondrial production of reactive oxygen species in relation to aging and pathology. *Ann. N. Y. Acad. Sci.* **2004**, *1011*, 86–100. https://doi.org/10.1007/978-3-662-41088-2_10.
216. Read, A.D.; Bentley, R.E.; Archer, S.L.; Dunham-Snary, K.J. Mitochondrial iron-sulfur clusters: Structure, function, and an emerging role in vascular biology. *Redox Biol.* **2021**, *47*, 102164. <https://doi.org/10.1016/j.redox.2021.102164>.
217. Zickermann, V.; Wirth, C.; Nasiri, H.; Siegmund, K.; Schwalbe, H.; Hunte, C.; Brandt, U. Structural biology. Mechanistic insight from the crystal structure of mitochondrial complex I. *Science* **2015**, *347*, 44–49. <https://doi.org/10.1126/science.1259859>.
218. Treberg, J.R.; Braun, K.; Selseleh, P. Mitochondria can act as energy-sensing regulators of hydrogen peroxide availability. *Redox Biol.* **2019**, *20*, 483–488. <https://doi.org/10.1016/j.redox.2018.11.002>.
219. Cardoso, A.R.; Chausse, B.; da Cunha, F.M.; Luevano-Martinez, L.A.; Marazzi, T.B.; Pessoa, P.S.; Queliconi, B.B.; Kowaltowski, A.J. Mitochondrial compartmentalization of redox processes. *Free Radic. Biol. Med.* **2012**, *52*, 2201–2208. <https://doi.org/10.1016/j.freeradbiomed.2012.03.008>.
220. Secilmis, M.; Altun, H.Y.; Pilic, J.; Erdogan, Y.C.; Cokluk, Z.; Ata, B.N.; Sevimli, G.; Zaki, A.G.; Yigit, E.N.; Ozturk, G.; et al. A Co-Culture-Based Multiparametric Imaging Technique to Dissect Local H₂O₂ Signals with Targeted HyPer7. *Biosensors* **2021**, *11*, 338. <https://doi.org/10.3390/bios11090338>.
221. Lim, J.B.; Huang, B.K.; Deen, W.M.; Sikes, H.D. Analysis of the lifetime and spatial localization of hydrogen peroxide generated in the cytosol using a reduced kinetic model. *Free Radic. Biol. Med.* **2015**, *89*, 47–53. <https://doi.org/10.1016/j.freeradbiomed.2015.07.009>.
222. Archer, S.L.; Reeve, H.L.; Michelakis, E.; Puttagunta, L.; Waite, R.; Nelson, D.P.; Dinauer, M.C.; Weir, E.K. O₂ sensing is preserved in mice lacking the gp91 phox subunit of NADPH oxidase. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 7944–7949. <https://doi.org/10.1073/pnas.96.14.7944>.
223. Hilenski, L.L.; Clemus, R.E.; Quinn, M.T.; Lambeth, J.D.; Griendling, K.K. Distinct subcellular localizations of Nox1 and Nox4 in vascular smooth muscle cells. *Arterioscler. Thromb. Vasc. Biol.* **2004**, *24*, 677–683. <https://doi.org/10.1161/01.ATV.0000112024.13727.2c>.
224. Miller, E.W.; Dickinson, B.C.; Chang, C.J. Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 15681–15686. <https://doi.org/10.1073/pnas.1005776107>.
225. Oakley, F.D.; Abbott, D.; Li, Q.; Engelhardt, J.F. Signaling components of redox active endosomes: The redoxosomes. *Antioxid. Redox Signal.* **2009**, *11*, 1313–1333. <https://doi.org/10.1089/ars.2008.2363>.
226. Shaifita, Y.; Snetkov, V.A.; Prieto-Lloret, J.; Knock, G.A.; Smirnov, S.V.; Aaronson, P.I.; Ward, J.P. Sphingosylphosphorylcholine potentiates vasoreactivity and voltage-gated Ca²⁺ entry via NOX1 and reactive oxygen species. *Cardiovasc. Res.* **2015**, *106*, 121–130. <https://doi.org/10.1093/cvr/cvv029>.
227. Miyano, K.; Ueno, N.; Takeya, R.; Sumimoto, H. Direct involvement of the small GTPase Rac in activation of the superoxide-producing NADPH oxidase Nox1. *J. Biol. Chem.* **2006**, *281*, 21857–21868. <https://doi.org/10.1074/jbc.M513665200>.
228. Schroder, K.; Weissmann, N.; Brandes, R.P. Organizers and activators: Cytosolic Nox proteins impacting on vascular function. *Free Radic. Biol. Med.* **2017**, *109*, 22–32. <https://doi.org/10.1016/j.freeradbiomed.2017.03.017>.
229. Streeter, J.; Schickling, B.M.; Jiang, S.; Stanic, B.; Thiel, W.H.; Gakhar, L.; Houtman, J.C.; Miller, F.J., Jr. Phosphorylation of Nox1 regulates association with NoxA1 activation domain. *Circ. Res.* **2014**, *115*, 911–918. <https://doi.org/10.1161/CIRCRESAHA.115.304267>.
230. Niu, X.L.; Madamanchi, N.R.; Vendrov, A.E.; Tchivilev, I.; Rojas, M.; Madamanchi, C.; Brandes, R.P.; Krause, K.H.; Humphries, J.; Smith, A.; et al. Nox activator 1: A potential target for modulation of vascular reactive oxygen species in atherosclerotic arteries. *Circulation* **2010**, *121*, 549–559. <https://doi.org/10.1161/CIRCULATIONAHA.109.908319>.
231. Zhang, F.; Jin, S.; Yi, F.; Xia, M.; Dewey, W.L.; Li, P.L. Local production of O^{2•−} by NAD(P)H oxidase in the sarcoplasmic reticulum of coronary arterial myocytes: cADPR-mediated Ca²⁺ regulation. *Cell Signal.* **2008**, *20*, 637–644. <https://doi.org/10.1016/j.cellsig.2007.11.013>.
232. Vendrov, A.E.; Vendrov, K.C.; Smith, A.; Yuan, J.; Sumida, A.; Robidoux, J.; Runge, M.S.; Madamanchi, N.R. NOX4 NADPH Oxidase-Dependent Mitochondrial Oxidative Stress in Aging-Associated Cardiovascular Disease. *Antioxid. Redox Signal.* **2015**, *23*, 1389–1409. <https://doi.org/10.1089/ars.2014.6221>.
233. Anilkumar, N.; San Jose, G.; Sawyer, I.; Santos, C.X.; Sand, C.; Brewer, A.C.; Warren, D.; Shah, A.M. A 28-kDa splice variant of NADPH oxidase-4 is nuclear-localized and involved in redox signaling in vascular cells. *Arterioscler. Thromb. Vasc. Biol.* **2013**, *33*, e104–e112. <https://doi.org/10.1161/ATVBAHA.112.300956>.

234. Kracun, D.; Lopes, L.R.; Cifuentes-Pagano, E.; Pagano, P.J. NADPH oxidases: Redox regulation of cell homeostasis and disease. *Physiol. Rev.* **2025**, *105*, 1291–1428. <https://doi.org/10.1152/physrev.00034.2023>.
235. Sutliff, R.L.; Hilenski, L.L.; Amanso, A.M.; Parastatidis, I.; Dikalova, A.E.; Hansen, L.; Datla, S.R.; Long, J.S.; El-Ali, A.M.; Joseph, G.; et al. Polymerase delta interacting protein 2 sustains vascular structure and function. *Arterioscler. Thromb. Vasc. Biol.* **2013**, *33*, 2154–2161. <https://doi.org/10.1161/ATVBAHA.113.301913>.
236. Stone, J.R.; Yang, S. Hydrogen peroxide: A signaling messenger. *Antioxid. Redox Signal.* **2006**, *8*, 243–270. <https://doi.org/10.1089/ars.2006.8.243>.
237. Oshino, N.; Chance, B.; Sies, H.; Bucher, T. The role of H₂O₂ generation in perfused rat liver and the reaction of catalase compound I and hydrogen donors. *Arch. Biochem. Biophys.* **1973**, *154*, 117–131. [https://doi.org/10.1016/0003-9861\(73\)90040-4](https://doi.org/10.1016/0003-9861(73)90040-4).
238. Chance, B.; Sies, H.; Boveris, A. Hydroperoxide metabolism in mammalian organs. *Physiol. Rev.* **1979**, *59*, 527–605. <https://doi.org/10.1152/physrev.1979.59.3.527>.
239. Jones, D.P. Intracellular catalase function: Analysis of the catalatic activity by product formation in isolated liver cells. *Arch. Biochem. Biophys.* **1982**, *214*, 806–814. [https://doi.org/10.1016/0003-9861\(82\)90087-x](https://doi.org/10.1016/0003-9861(82)90087-x).
240. Lyublinskaya, O.; Antunes, F. Measuring intracellular concentration of hydrogen peroxide with the use of genetically encoded H₂O₂ biosensor HyPer. *Redox Biol.* **2019**, *24*, 101200. <https://doi.org/10.1016/j.redox.2019.101200>.
241. Bienert, G.P.; Chaumont, F. Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide. *Biochim. Biophys. Acta* **2014**, *1840*, 1596–1604. <https://doi.org/10.1016/j.bbagen.2013.09.017>.
242. Antunes, F.; Cadenas, E. Estimation of H₂O₂ gradients across biomembranes. *FEBS Lett.* **2000**, *475*, 121–126. [https://doi.org/10.1016/S0014-5793\(00\)01638-0](https://doi.org/10.1016/S0014-5793(00)01638-0).
243. Makino, N.; Mochizuki, Y.; Bannai, S.; Sugita, Y. Kinetic studies on the removal of extracellular hydrogen peroxide by cultured fibroblasts. *J. Biol. Chem.* **1994**, *269*, 1020–1025.
244. Wagner, B.A.; Witmer, J.R.; van't Erve, T.J.; Buettner, G.R. An Assay for the Rate of Removal of Extracellular Hydrogen Peroxide by Cells. *Redox Biol.* **2013**, *1*, 210–217. <https://doi.org/10.1016/j.redox.2013.01.011>.
245. Huang, B.K.; Sikes, H.D. Quantifying intracellular hydrogen peroxide perturbations in terms of concentration. *Redox Biol.* **2014**, *2*, 955–962. <https://doi.org/10.1016/j.redox.2014.08.001>.
246. Adimora, N.J.; Jones, D.P.; Kemp, M.L. A model of redox kinetics implicates the thiol proteome in cellular hydrogen peroxide responses. *Antioxid. Redox Signal.* **2010**, *13*, 731–743. <https://doi.org/10.1089/ars.2009.2968>.
247. Sobotta, M.C.; Barata, A.G.; Schmidt, U.; Mueller, S.; Millonig, G.; Dick, T.P. Exposing cells to H₂O₂: A quantitative comparison between continuous low-dose and one-time high-dose treatments. *Free Radic. Biol. Med.* **2013**, *60*, 325–335. <https://doi.org/10.1016/j.freeradbiomed.2013.02.017>.
248. Marinho, H.S.; Cyrne, L.; Cadenas, E.; Antunes, F. H₂O₂ delivery to cells: Steady-state versus bolus addition. *Methods Enzymol.* **2013**, *526*, 159–173. <https://doi.org/10.1016/B978-0-12-405883-5.00010-7>.
249. Lim, J.B.; Langford, T.F.; Huang, B.K.; Deen, W.M.; Sikes, H.D. A reaction-diffusion model of cytosolic hydrogen peroxide. *Free Radic. Biol. Med.* **2016**, *90*, 85–90. <https://doi.org/10.1016/j.freeradbiomed.2015.11.005>.
250. Jones, D.P. Radical-free biology of oxidative stress. *Am. J. Physiol. Cell Physiol.* **2008**, *295*, C849–C868. <https://doi.org/10.1152/ajp-cell.00283.2008>.
251. Pardo-Pena, K.; Yanez-Hernandez, A.; Medina-Ceja, L.; Morales-Villagran, A. Ellagic acid and allopurinol decrease H₂O₂ concentrations, epileptiform activity and astrogliosis after status epilepticus in the hippocampus of adult rats. *Exp. Brain Res.* **2022**, *240*, 1191–1203. <https://doi.org/10.1007/s00221-022-06323-9>.
252. Pardo-Pena, K.; Sanchez-Lira, A.; Salazar-Sanchez, J.C.; Morales-Villagran, A. A novel online fluorescence method for in-vivo measurement of hydrogen peroxide during oxidative stress produced in a temporal lobe epilepsy model. *Neuroreport* **2018**, *29*, 621–630. <https://doi.org/10.1097/WNR.0000000000001007>.
253. Pardo-Pena, K.; Lorea-Hernandez, J.J.; Camacho-Hernandez, N.P.; Ordaz, B.; Villasana-Salazar, B.; Morales-Villagran, A.; Pena-Ortega, F. Hydrogen peroxide extracellular concentration in the ventrolateral medulla and its increase in response to hypoxia in vitro: Possible role of microglia. *Brain Res.* **2018**, *1692*, 87–99. <https://doi.org/10.1016/j.brainres.2018.04.032>.
254. Puppulin, L.; Hosogi, S.; Sun, H.; Matsuo, K.; Inui, T.; Kumamoto, Y.; Suzaki, T.; Tanaka, H.; Marunaka, Y. Bioconjugation strategy for cell surface labelling with gold nanostructures designed for highly localized pH measurement. *Nat. Commun.* **2018**, *9*, 5278. <https://doi.org/10.1038/s41467-018-07726-5>.
255. Hosogi, S.; Marunaka, Y.; Ashihara, E.; Yamada, T.; Sumino, A.; Tanaka, H.; Puppulin, L. Plasma membrane anchored nanosensor for quantifying endogenous production of H₂O₂ in living cells. *Biosens. Bioelectron.* **2021**, *179*, 113077. <https://doi.org/10.1016/j.bios.2021.113077>.

256. Forman, H.J.; Bernardo, A.; Davies, K.J. What is the concentration of hydrogen peroxide in blood and plasma? *Arch. Biochem. Biophys.* **2016**, *603*, 48–53. <https://doi.org/10.1016/j.abb.2016.05.005>.
257. Sousa, T.; Gouveia, M.; Travasso, R.D.M.; Salvador, A. How abundant are superoxide and hydrogen peroxide in the vasculature lumen, how far can they reach? *Redox Biol.* **2022**, *58*, 102527. <https://doi.org/10.1016/j.redox.2022.102527>.
258. Bayer, S.B.; Maghzal, G.; Stocker, R.; Hampton, M.B.; Winterbourn, C.C. Neutrophil-mediated oxidation of erythrocyte peroxiredoxin 2 as a potential marker of oxidative stress in inflammation. *FASEB J.* **2013**, *27*, 3315–3322. <https://doi.org/10.1096/fj.13-227298>.
259. Ezerina, D.; Morgan, B.; Dick, T.P. Imaging dynamic redox processes with genetically encoded probes. *J. Mol. Cell. Cardiol.* **2014**, *73*, 43–49. <https://doi.org/10.1016/j.yjmcc.2013.12.023>.
260. Mishra, P.K.; Park, I.; Sharma, N.; Yoo, C.M.; Lee, H.Y.; Rhee, H.W. Enzymatic Recording of Local Hydrogen Peroxide Generation Using Genetically Encodable Enzyme. *Anal. Chem.* **2022**, *94*, 14869–14877. <https://doi.org/10.1021/acs.analchem.2c01966>.
261. Eid, M.; Barayeu, U.; Sulkova, K.; Aranda-Vallejo, C.; Dick, T.P. Using the heme peroxidase APEX2 to probe intracellular H₂O₂ flux and diffusion. *Nat. Commun.* **2024**, *15*, 1239. <https://doi.org/10.1038/s41467-024-45511-9>.
262. Yamdjeu, O.T.; Begerow, A.; Sommer, N.; Diener, M.; Weissmann, N.; Knoepp, F. H₂O₂ Sensitivity of K(v) Channels in Hypoxic Pulmonary Vasoconstriction: Experimental Conditions Matter. *Int. J. Mol. Sci.* **2025**, *26*, 6857. <https://doi.org/10.3390/ijms26146857>.
263. Mishina, N.M.; Tyurin-Kuzmin, P.A.; Markvicheva, K.N.; Vorotnikov, A.V.; Tkachuk, V.A.; Laketa, V.; Schultz, C.; Lukyanov, S.; Belousov, V.V. Does cellular hydrogen peroxide diffuse or act locally? *Antioxid. Redox Signal.* **2011**, *14*, 1–7. <https://doi.org/10.1089/ars.2010.3539>.
264. Mishina, N.M.; Bogdanova, Y.A.; Ermakova, Y.G.; Panova, A.S.; Kotova, D.A.; Bilan, D.S.; Steinhorn, B.; Arner, E.S.J.; Michel, T.; Belousov, V.V. Which Antioxidant System Shapes Intracellular H₂O₂ Gradients? *Antioxid. Redox Signal.* **2019**, *31*, 664–670. <https://doi.org/10.1089/ars.2018.7697>.
265. Pak, V.V.; Ezerina, D.; Lyublinskaya, O.G.; Pedre, B.; Tyurin-Kuzmin, P.A.; Mishina, N.M.; Thauvin, M.; Young, D.; Wahni, K.; Martinez Gache, S.A.; et al. Ultrasensitive Genetically Encoded Indicator for Hydrogen Peroxide Identifies Roles for the Oxidant in Cell Migration and Mitochondrial Function. *Cell Metab.* **2020**, *31*, 642–653.e646. <https://doi.org/10.1016/j.cmet.2020.02.003>.
266. Kritsiligkou, P.; Bosch, K.; Shen, T.K.; Meurer, M.; Knop, M.; Dick, T.P. Proteome-wide tagging with an H₂O₂ biosensor reveals highly localized and dynamic redox microenvironments. *Proc. Natl. Acad. Sci. USA* **2023**, *120*, e2314043120. <https://doi.org/10.1073/pnas.2314043120>.
267. Potekhina, E.S.; Bass, D.I.; Ezerina, D.; Fleckenstein, D.D.; Chebotarev, A.S.; Sysoeva, V.A.; Maltsev, D.I.; Pak, V.V.; Moshchenko, A.A.; Sokolov, A.I.; et al. A color-tailored fluorogenic sensor for hydrogen peroxide. *Nat. Chem. Biol.* **2025**, 1–12. <https://doi.org/10.1038/s41589-025-02036-6>.
268. Booth, D.M.; Varnai, P.; Joseph, S.K.; Hajnoczky, G. Oxidative bursts of single mitochondria mediate retrograde signaling toward the ER. *Mol. Cell* **2021**, *81*, 3866–3876.e3862. <https://doi.org/10.1016/j.molcel.2021.07.014>.
269. Joseph, S.K.; Young, M.P.; Alzayady, K.; Yule, D.I.; Ali, M.; Booth, D.M.; Hajnoczky, G. Redox regulation of type-I inositol trisphosphate receptors in intact mammalian cells. *J. Biol. Chem.* **2018**, *293*, 17464–17476. <https://doi.org/10.1074/jbc.RA118.005624>.
270. King, M.P.; Attardi, G. Isolation of human cell lines lacking mitochondrial DNA. *Methods Enzymol.* **1996**, *264*, 304–313. [https://doi.org/10.1016/s0076-6879\(96\)64029-4](https://doi.org/10.1016/s0076-6879(96)64029-4).
271. Firth, A.L.; Gordienko, D.V.; Yuill, K.H.; Smirnov, S.V. Cellular localization of mitochondria contributes to Kv channel-mediated regulation of cellular excitability in pulmonary but not mesenteric circulation. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2009**, *296*, L347–L360. <https://doi.org/10.1152/ajplung.90341.2008>.
272. Archer, S.L.; Huang, J.; Henry, T.; Peterson, D.; Weir, E.K. A redox-based O₂ sensor in rat pulmonary vasculature. *Circ. Res.* **1993**, *73*, 1100–1112. <https://doi.org/10.1161/01.res.73.6.1100>.
273. Weir, E.K. Does normoxic pulmonary vasodilatation rather than hypoxic vasoconstriction account for the pulmonary pressor response to hypoxia? *Lancet* **1978**, *1*, 476–477. [https://doi.org/10.1016/s0140-6736\(78\)90138-1](https://doi.org/10.1016/s0140-6736(78)90138-1).
274. Kilfoil, P.J.; Tipparaju, S.M.; Barski, O.A.; Bhatnagar, A. Regulation of ion channels by pyridine nucleotides. *Circ. Res.* **2013**, *112*, 721–741. <https://doi.org/10.1161/CIRCRESAHA.111.247940>.
275. Jones, D.P.; Sies, H. The Redox Code. *Antioxid. Redox Signal.* **2015**, *23*, 734–746. <https://doi.org/10.1089/ars.2015.6247>.
276. Dwenger, M.M.; Raph, S.M.; Reyzer, M.L.; Lisa Manier, M.; Riggs, D.W.; Wohl, Z.B.; Ohanyan, V.; Mack, G.; Pucci, T.; Moore, J.B.; et al. Pyridine nucleotide redox potential in coronary smooth muscle couples myocardial blood flow to cardiac metabolism. *Nat. Commun.* **2022**, *13*, 2051. <https://doi.org/10.1038/s41467-022-29745-z>.

277. Liu, S.Q.; Jin, H.; Zacarias, A.; Srivastava, S.; Bhatnagar, A. Binding of pyridine coenzymes to the beta-subunit of the voltage sensitive potassium channels. *Chem. Biol. Interact.* **2001**, *130*–132, 955–962. [https://doi.org/10.1016/s0009-2797\(00\)00248-9](https://doi.org/10.1016/s0009-2797(00)00248-9).
278. Gutscher, M.; Pauleau, A.L.; Marty, L.; Brach, T.; Wabnitz, G.H.; Samstag, Y.; Meyer, A.J.; Dick, T.P. Real-time imaging of the intracellular glutathione redox potential. *Nat. Methods* **2008**, *5*, 553–559. <https://doi.org/10.1038/nmeth.1212>.
279. Swain, L.; Kesemeyer, A.; Meyer-Roxlau, S.; Vettel, C.; Zieseniss, A.; Guntsch, A.; Jatho, A.; Becker, A.; Nanadikar, M.S.; Morgan, B.; et al. Redox Imaging Using Cardiac Myocyte-Specific Transgenic Biosensor Mice. *Circ. Res.* **2016**, *119*, 1004–1016. <https://doi.org/10.1161/CIRCRESAHA.116.309551>.
280. Weir, E.K.; Lopez-Barneo, J.; Buckler, K.J.; Archer, S.L. Acute oxygen-sensing mechanisms. *N. Engl. J. Med.* **2005**, *353*, 2042–2055. <https://doi.org/10.1056/NEJMr050002>.
281. Weir, E.K.; Will, J.A.; Lundquist, L.J.; Eaton, J.W.; Chesler, E. Diamide inhibits pulmonary vasoconstriction induced by hypoxia or prostaglandin F2 alpha. *Proc. Soc. Exp. Biol. Med.* **1983**, *173*, 96–103. <https://doi.org/10.3181/00379727-173-41615>.
282. Burghuber, O.; Mathias, M.M.; McMurtry, I.F.; Reeves, J.T.; Voelkel, N.F. Lung edema due to hydrogen peroxide is independent of cyclooxygenase products. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* **1984**, *56*, 900–905. <https://doi.org/10.1152/jappl.1984.56.4.900>.
283. Weir, E.K.; Eaton, J.W.; Chesler, E. Redox status and pulmonary vascular reactivity. *Chest* **1985**, *88*, 249S–252S. https://doi.org/10.1378/chest.88.4_supplement.249s.
284. Archer, S.L.; Will, J.A.; Weir, E.K. Redox status in the control of pulmonary vascular tone. *Herz* **1986**, *11*, 127–141.
285. Archer, S.L.; Peterson, D.; Nelson, D.P.; DeMaster, E.G.; Kelly, B.; Eaton, J.W.; Weir, E.K. Oxygen radicals and antioxidant enzymes alter pulmonary vascular reactivity in the rat lung. *J. Appl. Physiol.* **1989**, *66*, 102–111. <https://doi.org/10.1152/jappl.1989.66.1.102>.
286. McMurtry, I.F. Angiotensin is not required for hypoxic constriction in salt solution-perfused rat lungs. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* **1984**, *56*, 375–380. <https://doi.org/10.1152/jappl.1984.56.2.375>.
287. Archer, S.L.; Nelson, D.P.; Weir, E.K. Simultaneous measurement of O₂ radicals and pulmonary vascular reactivity in rat lung. *J. Appl. Physiol.* **1989**, *67*, 1903–1911. <https://doi.org/10.1152/jappl.1989.67.5.1903>.
288. Post, J.M.; Hume, J.R.; Archer, S.L.; Weir, E.K. Direct role for potassium channel inhibition in hypoxic pulmonary vasoconstriction. *Am. J. Physiol.* **1992**, *262*, C882–C890. <https://doi.org/10.1152/ajpcell.1992.262.4.C882>.
289. Rounds, S.; McMurtry, I.F. Inhibitors of oxidative ATP production cause transient vasoconstriction and block subsequent pressor responses in rat lungs. *Circ. Res.* **1981**, *48*, 393–400. <https://doi.org/10.1161/01.res.48.3.393>.
290. Leach, R.M.; Sheehan, D.W.; Chacko, V.P.; Sylvester, J.T. Effects of hypoxia on energy state and pH in resting pulmonary and femoral arterial smooth muscles. *Am. J. Physiol.* **1998**, *275*, L1051–L1060. <https://doi.org/10.1152/ajplung.1998.275.6.L1051>.
291. Leach, R.M.; Sheehan, D.W.; Chacko, V.P.; Sylvester, J.T. Energy state, pH, and vasomotor tone during hypoxia in precontracted pulmonary and femoral arteries. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2000**, *278*, L294–L304. <https://doi.org/10.1152/ajplung.2000.278.2.L294>.
292. Buescher, P.C.; Pearse, D.B.; Pillai, R.P.; Litt, M.C.; Mitchell, M.C.; Sylvester, J.T. Energy state and vasomotor tone in hypoxic pig lungs. *J. Appl. Physiol.* **1991**, *70*, 1874–1881. <https://doi.org/10.1152/jappl.1991.70.4.1874>.
293. Wu, W.; Platoshyn, O.; Firth, A.L.; Yuan, J.X. Hypoxia divergently regulates production of reactive oxygen species in human pulmonary and coronary artery smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2007**, *293*, L952–L959. <https://doi.org/10.1152/ajplung.00203.2007>.
294. Mehta, J.P.; Campian, J.L.; Guardiola, J.; Cabrera, J.A.; Weir, E.K.; Eaton, J.W. Generation of oxidants by hypoxic human pulmonary and coronary smooth-muscle cells. *Chest* **2008**, *133*, 1410–1414. <https://doi.org/10.1378/chest.07-2984>.
295. Divakaruni, A.S.; Jastroch, M. A practical guide for the analysis, standardization and interpretation of oxygen consumption measurements. *Nat. Metab.* **2022**, *4*, 978–994. <https://doi.org/10.1038/s42255-022-00619-4>.
296. Belousov, V.V.; Fradkov, A.F.; Lukyanov, K.A.; Staroverov, D.B.; Shakhbazov, K.S.; Tersikh, A.V.; Lukyanov, S. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat. Methods* **2006**, *3*, 281–286. <https://doi.org/10.1038/nmeth866>.
297. Brandt, U. Energy converting NADH:quinone oxidoreductase (complex I). *Annu. Rev. Biochem.* **2006**, *75*, 69–92. <https://doi.org/10.1146/annurev.biochem.75.103004.142539>.
298. Hameedi, M.A.; Grba, D.N.; Richardson, K.H.; Jones, A.J.Y.; Song, W.; Roessler, M.M.; Wright, J.J.; Hirst, J. A conserved arginine residue is critical for stabilizing the N₂ FeS cluster in mitochondrial complex I. *J. Biol. Chem.* **2021**, *296*, 100474. <https://doi.org/10.1016/j.jbc.2021.100474>.

299. Kashani-Poor, N.; Zwicker, K.; Kersch, S.; Brandt, U. A central functional role for the 49-kDa subunit within the catalytic core of mitochondrial complex I. *J. Biol. Chem.* **2001**, *276*, 24082–24087. <https://doi.org/10.1074/jbc.M102296200>.
300. Prieur, I.; Lunardi, J.; Dupuis, A. Evidence for a quinone binding site close to the interface between NUOD and NUOB subunits of Complex I. *Biochim. Biophys. Acta* **2001**, *1504*, 173–178. [https://doi.org/10.1016/s0005-2728\(01\)00158-x](https://doi.org/10.1016/s0005-2728(01)00158-x).
301. Read, A.D.; Bentley, R.E.T.; Martin, A.Y.; Mewburn, J.D.; Alizadeh, E.; Wu, D.; Lima, P.D.A.; Dunham-Snary, K.J.; Thebaud, B.; Sharp, W.; et al. Electron Leak From the Mitochondrial Electron Transport Chain Complex I at Site I(Q) Is Crucial for Oxygen Sensing in Rabbit and Human Ductus Arteriosus. *J. Am. Heart Assoc.* **2023**, *12*, e029131. <https://doi.org/10.1161/JAHA.122.029131>.
302. Huttemann, M.; Sommer, N.; Weissmann, N.; Grossman, L.I. Letter by Huttemann et al Regarding Article, “Ndufs2, a Core Subunit of Mitochondrial Complex I, Is Essential for Acute Oxygen-Sensing and Hypoxic Pulmonary Vasoconstriction”. *Circ. Res.* **2019**, *125*, e33–e34. <https://doi.org/10.1161/CIRCRESAHA.119.315815>.
303. Dunham-Snary, K.J.; Archer, S.L. Response by Dunham-Snary and Archer to Letter Regarding Article, “Ndufs2, a Core Subunit of Mitochondrial Complex I, Is Essential for Acute Oxygen-Sensing and Hypoxic Pulmonary Vasoconstriction”. *Circ. Res.* **2019**, *125*, e35–e36. <https://doi.org/10.1161/CIRCRESAHA.119.315826>.
304. Onukwufor, J.O.; Farooqi, M.A.; Vodickova, A.; Koren, S.A.; Baldzizhar, A.; Berry, B.J.; Beutner, G.; Porter, G.A., Jr.; Belousov, V.; Grossfield, A.; et al. A reversible mitochondrial complex I thiol switch mediates hypoxic avoidance behavior in *C. elegans*. *Nat. Commun.* **2022**, *13*, 2403. <https://doi.org/10.1038/s41467-022-30169-y>.
305. Bandara, A.B.; Drake, J.C.; James, C.C.; Smyth, J.W.; Brown, D.A. Complex I protein NDUFS2 is vital for growth, ROS generation, membrane integrity, apoptosis, and mitochondrial energetics. *Mitochondrion* **2021**, *58*, 160–168. <https://doi.org/10.1016/j.mito.2021.03.003>.
306. Sazanov, L.A. From the ‘black box’ to ‘domino effect’ mechanism: What have we learned from the structures of respiratory complex I. *Biochem. J.* **2023**, *480*, 319–333. <https://doi.org/10.1042/BCJ20210285>.
307. Pagniez-Mammeri, H.; Loublier, S.; Legrand, A.; Benit, P.; Rustin, P.; Slama, A. Mitochondrial complex I deficiency of nuclear origin I. Structural genes. *Mol. Genet. Metab.* **2012**, *105*, 163–172. <https://doi.org/10.1016/j.ymgme.2011.11.188>.
308. Burska, D.; Stiburek, L.; Krizova, J.; Vanisova, M.; Martinek, V.; Sladkova, J.; Zamecnik, J.; Honzik, T.; Zeman, J.; Hansikova, H.; et al. Homozygous missense mutation in UQCRC2 associated with severe encephalomyopathy, mitochondrial complex III assembly defect and activation of mitochondrial protein quality control. *Biochim. Biophys. Acta Mol. Basis Dis.* **2021**, *1867*, 166147. <https://doi.org/10.1016/j.bbadis.2021.166147>.
309. Iuso, A.; Scacco, S.; Piccoli, C.; Bellomo, F.; Petruzzella, V.; Trentadue, R.; Minuto, M.; Ripoli, M.; Capitanio, N.; Zeviani, M.; et al. Dysfunctions of cellular oxidative metabolism in patients with mutations in the NDUFS1 and NDUFS4 genes of complex I. *J. Biol. Chem.* **2006**, *281*, 10374–10380. <https://doi.org/10.1074/jbc.M513387200>.
310. McMurtry, I.F.; Davidson, A.B.; Reeves, J.T.; Grover, R.F. Inhibition of hypoxic pulmonary vasoconstriction by calcium antagonists in isolated rat lungs. *Circ. Res.* **1976**, *38*, 99–104. <https://doi.org/10.1161/01.res.38.2.99>.
311. Harder, D.R.; Madden, J.A.; Dawson, C. A membrane electrical mechanism for hypoxic vasoconstriction of small pulmonary arteries from cat. *Chest* **1985**, *88*, 233S–235S. https://doi.org/10.1378/chest.88.4_supplement.233s.
312. Hasunuma, K.; Rodman, D.M.; McMurtry, I.F. Effects of K⁺ channel blockers on vascular tone in the perfused rat lung. *Am. Rev. Respir. Dis.* **1991**, *144*, 884–887. <https://doi.org/10.1164/ajrccm/144.4.884>.
313. Yuan, X.J. Voltage-gated K⁺ currents regulate resting membrane potential and [Ca²⁺]_i in pulmonary arterial myocytes. *Circ. Res.* **1995**, *77*, 370–378. <https://doi.org/10.1161/01.res.77.2.370>.
314. Yuan, X.J.; Goldman, W.F.; Tod, M.L.; Rubin, L.J.; Blaustein, M.P. Hypoxia reduces potassium currents in cultured rat pulmonary but not mesenteric arterial myocytes. *Am. J. Physiol.* **1993**, *264*, L116–L123. <https://doi.org/10.1152/ajplung.1993.264.2.L116>.
315. Archer, S.L.; Huang, J.M.; Reeve, H.L.; Hampl, V.; Tolarova, S.; Michelakis, E.; Weir, E.K. Differential distribution of electrophysiologically distinct myocytes in conduit and resistance arteries determines their response to nitric oxide and hypoxia. *Circ. Res.* **1996**, *78*, 431–442. <https://doi.org/10.1161/01.res.78.3.431>.
316. Olschewski, A.; Hong, Z.; Nelson, D.P.; Weir, E.K. Graded response of K⁺ current, membrane potential, and [Ca²⁺]_i to hypoxia in pulmonary arterial smooth muscle. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2002**, *283*, L1143–L1150. <https://doi.org/10.1152/ajplung.00104.2002>.
317. Archer, S.L.; London, B.; Hampl, V.; Wu, X.; Nsair, A.; Puttagunta, L.; Hashimoto, K.; Waite, R.E.; Michelakis, E.D. Impairment of hypoxic pulmonary vasoconstriction in mice lacking the voltage-gated potassium channel Kv1.5. *FASEB J.* **2001**, *15*, 1801–1803. <https://doi.org/10.1096/fj.00-0649fje>.

318. Archer, S.L.; Souil, E.; Dinh-Xuan, A.T.; Schremmer, B.; Mercier, J.C.; El Yaagoubi, A.; Nguyen-Huu, L.; Reeve, H.L.; Hampl, V. Molecular identification of the role of voltage-gated K⁺ channels, Kv1.5 and Kv2.1, in hypoxic pulmonary vasoconstriction and control of resting membrane potential in rat pulmonary artery myocytes. *J. Clin. Investig.* **1998**, *101*, 2319–2330. <https://doi.org/10.1172/JCI333>.
319. Firth, A.L.; Platoshyn, O.; Brevnova, E.E.; Burg, E.D.; Powell, F.; Haddad, G.H.; Yuan, J.X. Hypoxia selectively inhibits KCNA5 channels in pulmonary artery smooth muscle cells. *Ann. N. Y. Acad. Sci.* **2009**, *1177*, 101–111. <https://doi.org/10.1111/j.1749-6632.2009.05040.x>.
320. Platoshyn, O.; Yu, Y.; Ko, E.A.; Remillard, C.V.; Yuan, J.X. Heterogeneity of hypoxia-mediated decrease in I(K(V)) and increase in [Ca²⁺]_(cyt) in pulmonary artery smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2007**, *293*, L402–L416. <https://doi.org/10.1152/ajplung.00391.2006>.
321. Olschewski, A.; Li, Y.; Tang, B.; Hanze, J.; Eul, B.; Bohle, R.M.; Wilhelm, J.; Morty, R.E.; Brau, M.E.; Weir, E.K.; et al. Impact of TASK-1 in human pulmonary artery smooth muscle cells. *Circ. Res.* **2006**, *98*, 1072–1080. <https://doi.org/10.1161/01.RES.0000219677.12988.e9>.
322. Fuchs, B.; Dietrich, A.; Gudermann, T.; Kalwa, H.; Grimminger, F.; Weissmann, N. The role of classical transient receptor potential channels in the regulation of hypoxic pulmonary vasoconstriction. *Adv. Exp. Med. Biol.* **2010**, *661*, 187–200. https://doi.org/10.1007/978-1-60761-500-2_12.
323. Olschewski, A.; Weir, E.K. Redox regulation of ion channels in the pulmonary circulation. *Antioxid. Redox Signal.* **2015**, *22*, 465–485. <https://doi.org/10.1089/ars.2014.5899>.
324. Rocher, A.; Aaronson, P.I. The Thirty-Fifth Anniversary of K⁺ Channels in O₂ sensing: What We Know and What We Don't Know. *Oxygen* **2024**, *4*, 53–89. <https://doi.org/10.3390/oxygen4010004>.
325. Firth, A.L.; Yuill, K.H.; Smirnov, S.V. Mitochondria-dependent regulation of Kv currents in rat pulmonary artery smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2008**, *295*, L61–L70. <https://doi.org/10.1152/ajplung.90243.2008>.
326. Searle, G.J.; Hartness, M.E.; Hoareau, R.; Peers, C.; Kemp, P.J. Lack of contribution of mitochondrial electron transport to acute O₂ sensing in model airway chemoreceptors. *Biochem. Biophys. Res. Commun.* **2002**, *291*, 332–337. <https://doi.org/10.1006/bbrc.2002.6428>.
327. Cogolludo, A.; Frazziano, G.; Cobeno, L.; Moreno, L.; Lodi, F.; Villamor, E.; Tamargo, J.; Perez-Vizcaino, F. Role of reactive oxygen species in Kv channel inhibition and vasoconstriction induced by TP receptor activation in rat pulmonary arteries. *Ann. N. Y. Acad. Sci.* **2006**, *1091*, 41–51. <https://doi.org/10.1196/annals.1378.053>.
328. Frazziano, G.; Moreno, L.; Moral-Sanz, J.; Menendez, C.; Escolano, L.; Gonzalez, C.; Villamor, E.; Alvarez-Sala, J.L.; Cogolludo, A.L.; Perez-Vizcaino, F. Neutral sphingomyelinase, NADPH oxidase and reactive oxygen species. Role in acute hypoxic pulmonary vasoconstriction. *J. Cell. Physiol.* **2011**, *226*, 2633–2640. <https://doi.org/10.1002/jcp.22611>.
329. Post, J.M.; Gelband, C.H.; Hume, J.R. [Ca²⁺]_i inhibition of K⁺ channels in canine pulmonary artery. Novel mechanism for hypoxia-induced membrane depolarization. *Circ. Res.* **1995**, *77*, 131–139. <https://doi.org/10.1161/01.res.77.1.131>.
330. Vandier, C.; Delpech, M.; Bonnet, P. Spontaneous transient outward currents and delayed rectifier K⁺ current: Effects of hypoxia. *Am. J. Physiol.* **1998**, *275*, L145–L154. <https://doi.org/10.1152/ajplung.1998.275.1.L145>.
331. Snetkov, V.A.; Smirnov, S.V.; Kua, J.; Aaronson, P.I.; Ward, J.P.; Knock, G.A. Superoxide differentially controls pulmonary and systemic vascular tone through multiple signalling pathways. *Cardiovasc. Res.* **2011**, *89*, 214–224. <https://doi.org/10.1093/cvr/cvq275>.
332. Svoboda, L.K.; Reddie, K.G.; Zhang, L.; Vesely, E.D.; Williams, E.S.; Schumacher, S.M.; O'Connell, R.P.; Shaw, R.; Day, S.M.; Anumonwo, J.M.; et al. Redox-sensitive sulfenic acid modification regulates surface expression of the cardiovascular voltage-gated potassium channel Kv1.5. *Circ. Res.* **2012**, *111*, 842–853. <https://doi.org/10.1161/CIRCRESAHA.111.263525>.
333. Raph, S.M.; Dwenger, M.M.; Hu, X.; Nystoriak, M.A. Basal NAD(H) redox state permits hydrogen peroxide-induced mesenteric artery dilatation. *J. Physiol.* **2023**, *601*, 2621–2634. <https://doi.org/10.1113/JP284195>.
334. Moreno, L.; Frazziano, G.; Cogolludo, A.; Cobeno, L.; Tamargo, J.; Perez-Vizcaino, F. Role of protein kinase C ζ and its adaptor protein p62 in voltage-gated potassium channel modulation in pulmonary arteries. *Mol. Pharmacol.* **2007**, *72*, 1301–1309. <https://doi.org/10.1124/mol.107.037002>.
335. Schumacher, S.M.; McEwen, D.P.; Zhang, L.; Arendt, K.L.; Van Genderen, K.M.; Martens, J.R. Antiarrhythmic drug-induced internalization of the atrial-specific k⁺ channel kv1.5. *Circ. Res.* **2009**, *104*, 1390–1398. <https://doi.org/10.1161/CIRCRESAHA.108.192773>.
336. Lee, S.; Park, M.; So, I.; Earm, Y.E. NADH and NAD modulates Ca²⁺-activated K⁺ channels in small pulmonary arterial smooth muscle cells of the rabbit. *Pflugers Arch.* **1994**, *427*, 378–380. <https://doi.org/10.1007/BF00374548>.

337. Reeve, H.L.; Weir, E.K.; Nelson, D.P.; Peterson, D.A.; Archer, S.L. Opposing effects of oxidants and antioxidants on K⁺ channel activity and tone in rat vascular tissue. *Exp. Physiol.* **1995**, *80*, 825–834. <https://doi.org/10.1113/expphysiol.1995.sp003890>.
338. Park, M.K.; Lee, S.H.; Ho, W.K.; Earm, Y.E. Redox agents as a link between hypoxia and the responses of ionic channels in rabbit pulmonary vascular smooth muscle. *Exp. Physiol.* **1995**, *80*, 835–842. <https://doi.org/10.1113/expphysiol.1995.sp003891>.
339. Park, M.K.; Bae, Y.M.; Lee, S.H.; Ho, W.K.; Earm, Y.E. Modulation of voltage-dependent K⁺ channel by redox potential in pulmonary and ear arterial smooth muscle cells of the rabbit. *Pflugers Arch.* **1997**, *434*, 764–771. <https://doi.org/10.1007/s004240050463>.
340. Olschewski, A.; Hong, Z.; Peterson, D.A.; Nelson, D.P.; Porter, V.A.; Weir, E.K. Opposite effects of redox status on membrane potential, cytosolic calcium, and tone in pulmonary arteries and ductus arteriosus. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2004**, *286*, L15–L22. <https://doi.org/10.1152/ajplung.00372.2002>.
341. Schach, C.; Xu, M.; Platoshyn, O.; Keller, S.H.; Yuan, J.X. Thiol oxidation causes pulmonary vasodilation by activating K⁺ channels and inhibiting store-operated Ca²⁺ channels. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2007**, *292*, L685–L698. <https://doi.org/10.1152/ajplung.00276.2006>.
342. Yuan, X.J.; Tod, M.L.; Rubin, L.J.; Blaustein, M.P. Deoxyglucose and reduced glutathione mimic effects of hypoxia on K⁺ and Ca²⁺ conductances in pulmonary artery cells. *Am. J. Physiol.* **1994**, *267*, L52–L63. <https://doi.org/10.1152/ajplung.1994.267.1.L52>.
343. Weir, E.K.; Hong, Z.; Porter, V.A.; Reeve, H.L. Redox signaling in oxygen sensing by vessels. *Respir. Physiol. Neurobiol.* **2002**, *132*, 121–130. [https://doi.org/10.1016/s1569-9048\(02\)00054-x](https://doi.org/10.1016/s1569-9048(02)00054-x).
344. Marshall, C.; Mamary, A.J.; Verhoeven, A.J.; Marshall, B.E. Pulmonary artery NADPH-oxidase is activated in hypoxic pulmonary vasoconstriction. *Am. J. Respir. Cell Mol. Biol.* **1996**, *15*, 633–644. <https://doi.org/10.1165/ajrcmb.15.5.8918370>.
345. Killilea, D.W.; Hester, R.; Balczon, R.; Babal, P.; Gillespie, M.N. Free radical production in hypoxic pulmonary artery smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2000**, *279*, L408–L412. <https://doi.org/10.1152/ajplung.2000.279.2.L408>.
346. Abdalla, S.; Will, J.A. Potentiation of the hypoxic contraction of guinea-pig isolated pulmonary arteries by two inhibitors of superoxide dismutase. *Gen. Pharmacol.* **1995**, *26*, 785–792. [https://doi.org/10.1016/0306-3623\(94\)00245-i](https://doi.org/10.1016/0306-3623(94)00245-i).
347. Duranteau, J.; Chandel, N.S.; Kulisz, A.; Shao, Z.; Schumacker, P.T. Intracellular signaling by reactive oxygen species during hypoxia in cardiomyocytes. *J. Biol. Chem.* **1998**, *273*, 11619–11624. <https://doi.org/10.1074/jbc.273.19.11619>.
348. Chandel, N.S.; Maltepe, E.; Goldwasser, E.; Mathieu, C.E.; Simon, M.C.; Schumacker, P.T. Mitochondrial reactive oxygen species trigger hypoxia-induced transcription. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 11715–11720. <https://doi.org/10.1073/pnas.95.20.11715>.
349. Turrens, J.F.; Alexandre, A.; Lehninger, A.L. Ubisemiquinone is the electron donor for superoxide formation by complex III of heart mitochondria. *Arch. Biochem. Biophys.* **1985**, *237*, 408–414. [https://doi.org/10.1016/0003-9861\(85\)90293-0](https://doi.org/10.1016/0003-9861(85)90293-0).
350. Heumuller, S.; Wind, S.; Barbosa-Sicard, E.; Schmidt, H.H.; Busse, R.; Schroder, K.; Brandes, R.P. Apocynin is not an inhibitor of vascular NADPH oxidases but an antioxidant. *Hypertension* **2008**, *51*, 211–217. <https://doi.org/10.1161/HYPERTENSIONAHA.107.100214>.
351. Leach, R.M.; Hill, H.M.; Snetkov, V.A.; Robertson, T.P.; Ward, J.P. Divergent roles of glycolysis and the mitochondrial electron transport chain in hypoxic pulmonary vasoconstriction of the rat: Identity of the hypoxic sensor. *J. Physiol.* **2001**, *536*, 211–224. <https://doi.org/10.1111/j.1469-7793.2001.00211.x>.
352. Robb, E.L.; Hall, A.R.; Prime, T.A.; Eaton, S.; Szibor, M.; Viscomi, C.; James, A.M.; Murphy, M.P. Control of mitochondrial superoxide production by reverse electron transport at complex I. *J. Biol. Chem.* **2018**, *293*, 9869–9879. <https://doi.org/10.1074/jbc.RA118.003647>.
353. Waypa, G.B.; Marks, J.D.; Mack, M.M.; Boriboun, C.; Mungai, P.T.; Schumacker, P.T. Mitochondrial reactive oxygen species trigger calcium increases during hypoxia in pulmonary arterial myocytes. *Circ. Res.* **2002**, *91*, 719–726. <https://doi.org/10.1161/01.res.0000036751.04896.f1>.
354. Weissmann, N.; Ebert, N.; Ahrens, M.; Ghofrani, H.A.; Schermuly, R.T.; Hanze, J.; Fink, L.; Rose, F.; Conzen, J.; Seeger, W.; et al. Effects of mitochondrial inhibitors and uncouplers on hypoxic vasoconstriction in rabbit lungs. *Am. J. Respir. Cell Mol. Biol.* **2003**, *29*, 721–732. <https://doi.org/10.1165/rcmb.2002-0217OC>.
355. Weissmann, N.; Tadic, A.; Hanze, J.; Rose, F.; Winterhalder, S.; Nollen, M.; Schermuly, R.T.; Ghofrani, H.A.; Seeger, W.; Grimminger, F. Hypoxic vasoconstriction in intact lungs: A role for NADPH oxidase-derived H₂O₂? *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2000**, *279*, L683–L690. <https://doi.org/10.1152/ajplung.2000.279.4.L683>.
356. Weissmann, N.; Winterhalder, S.; Nollen, M.; Voswinkel, R.; Quanz, K.; Ghofrani, H.A.; Schermuly, R.T.; Seeger, W.; Grimminger, F. NO and reactive oxygen species are involved in biphasic hypoxic vasoconstriction of isolated rabbit lungs. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2001**, *280*, L638–L645. <https://doi.org/10.1152/ajplung.2001.280.4.L638>.

357. Weissmann, N.; Zeller, S.; Schafer, R.U.; Turowski, C.; Ay, M.; Quanz, K.; Ghofrani, H.A.; Schermuly, R.T.; Fink, L.; Seeger, W.; et al. Impact of mitochondria and NADPH oxidases on acute and sustained hypoxic pulmonary vasoconstriction. *Am. J. Respir. Cell Mol. Biol.* **2006**, *34*, 505–513. <https://doi.org/10.1165/rcmb.2005-0337OC>.
358. Liu, J.Q.; Sham, J.S.; Shimoda, L.A.; Kuppusamy, P.; Sylvester, J.T. Hypoxic constriction and reactive oxygen species in porcine distal pulmonary arteries. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2003**, *285*, L322–L333. <https://doi.org/10.1152/ajplung.00337.2002>.
359. Dikalov, S.; Griendling, K.K.; Harrison, D.G. Measurement of reactive oxygen species in cardiovascular studies. *Hypertension* **2007**, *49*, 717–727. <https://doi.org/10.1161/01.HYP.0000258594.87211.6b>.
360. Guzy, R.D.; Hoyos, B.; Robin, E.; Chen, H.; Liu, L.; Mansfield, K.D.; Simon, M.C.; Hammerling, U.; Schumacker, P.T. Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing. *Cell Metab.* **2005**, *1*, 401–408. <https://doi.org/10.1016/j.cmet.2005.05.001>.
361. Waypa, G.B.; Guzy, R.; Mungai, P.T.; Mack, M.M.; Marks, J.D.; Roe, M.W.; Schumacker, P.T. Increases in mitochondrial reactive oxygen species trigger hypoxia-induced calcium responses in pulmonary artery smooth muscle cells. *Circ. Res.* **2006**, *99*, 970–978. <https://doi.org/10.1161/01.RES.0000247068.75808.3f>.
362. Schafer, F.Q.; Buettner, G.R. Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic. Biol. Med.* **2001**, *30*, 1191–1212. [https://doi.org/10.1016/s0891-5849\(01\)00480-4](https://doi.org/10.1016/s0891-5849(01)00480-4).
363. Sabharwal, S.S.; Waypa, G.B.; Marks, J.D.; Schumacker, P.T. Peroxiredoxin-5 targeted to the mitochondrial intermembrane space attenuates hypoxia-induced reactive oxygen species signalling. *Biochem. J.* **2013**, *456*, 337–346. <https://doi.org/10.1042/BJ20130740>.
364. Wang, Q.S.; Zheng, Y.M.; Dong, L.; Ho, Y.S.; Guo, Z.; Wang, Y.X. Role of mitochondrial reactive oxygen species in hypoxia-dependent increase in intracellular calcium in pulmonary artery myocytes. *Free Radic. Biol. Med.* **2007**, *42*, 642–653. <https://doi.org/10.1016/j.freeradbiomed.2006.12.008>.
365. Yang, Z.; Zhuan, B.; Yan, Y.; Jiang, S.; Wang, T. Roles of different mitochondrial electron transport chain complexes in hypoxia-induced pulmonary vasoconstriction. *Cell Biol. Int.* **2016**, *40*, 188–195. <https://doi.org/10.1002/cbin.10550>.
366. Waypa, G.B.; Marks, J.D.; Guzy, R.; Mungai, P.T.; Schriewer, J.; Dokic, D.; Schumacker, P.T. Hypoxia triggers subcellular compartmental redox signaling in vascular smooth muscle cells. *Circ. Res.* **2010**, *106*, 526–535. <https://doi.org/10.1161/CIRCRESAHA.109.206334>.
367. Meyer, A.J.; Dick, T.P. Fluorescent protein-based redox probes. *Antioxid. Redox Signal.* **2010**, *13*, 621–650. <https://doi.org/10.1089/ars.2009.2948>.
368. Desiredi, J.R.; Farrow, K.N.; Marks, J.D.; Waypa, G.B.; Schumacker, P.T. Hypoxia increases ROS signaling and cytosolic Ca²⁺ in pulmonary artery smooth muscle cells of mouse lungs slices. *Antioxid. Redox Signal.* **2010**, *12*, 595–602. <https://doi.org/10.1089/ars.2009.2862>.
369. Doctrow, S.R.; Huffman, K.; Marcus, C.B.; Musleh, W.; Bruce, A.; Baudry, M.; Malfroy, B. Salen-manganese complexes: Combined superoxide dismutase/catalase mimics with broad pharmacological efficacy. *Adv. Pharmacol.* **1997**, *38*, 247–269. [https://doi.org/10.1016/s1054-3589\(08\)60987-4](https://doi.org/10.1016/s1054-3589(08)60987-4).
370. Casteels, R.; Raeymaekers, L.; Suzuki, H.; Van Eldere, J. Tension response and ⁴⁵Ca release in vascular smooth muscle incubated in Ca-free solution. *Pflügers Arch.* **1981**, *392*, 139–145. <https://doi.org/10.1007/BF00581262>.
371. Paddenbergh, R.; Konig, P.; Faulhammer, P.; Goldenberg, A.; Pfeil, U.; Kummer, W. Hypoxic vasoconstriction of partial muscular intra-acinar pulmonary arteries in murine precision cut lung slices. *Respir. Res.* **2006**, *7*, 93. <https://doi.org/10.1186/1465-9921-7-93>.
372. Sanderson, M.J.; Bai, Y.; Perez-Zoghbi, J. Ca²⁺ oscillations regulate contraction of intrapulmonary smooth muscle cells. *Adv. Exp. Med. Biol.* **2010**, *661*, 77–96. https://doi.org/10.1007/978-1-60761-500-2_5.
373. Panzer, J.K.; Hiller, H.; Cohrs, C.M.; Almaca, J.; Enos, S.J.; Beery, M.; Cechin, S.; Drotar, D.M.; Weitz, J.R.; Santini, J.; et al. Pancreas tissue slices from organ donors enable in situ analysis of type 1 diabetes pathogenesis. *JCI Insight* **2020**, *5*, e134525. <https://doi.org/10.1172/jci.insight.134525>.
374. Truong, L.; Zheng, Y.M.; Wang, Y.X. Mitochondrial Rieske iron-sulfur protein in pulmonary artery smooth muscle: A key primary signaling molecule in pulmonary hypertension. *Arch. Biochem. Biophys.* **2020**, *683*, 108234. <https://doi.org/10.1016/j.abb.2019.108234>.
375. Song, T.; Zheng, Y.M.; Wang, Y.X. Cross Talk Between Mitochondrial Reactive Oxygen Species and Sarcoplasmic Reticulum Calcium in Pulmonary Arterial Smooth Muscle Cells. *Adv. Exp. Med. Biol.* **2017**, *967*, 289–298. https://doi.org/10.1007/978-3-319-63245-2_17.

376. Yang, Z.; Song, T.; Truong, L.; Reyes-Garcia, J.; Wang, L.; Zheng, Y.M.; Wang, Y.X. Important Role of Sarcoplasmic Reticulum Ca^{2+} Release via Ryanodine Receptor-2 Channel in Hypoxia-Induced Rieske Iron-Sulfur Protein-Mediated Mitochondrial Reactive Oxygen Species Generation in Pulmonary Artery Smooth Muscle Cells. *Antioxid. Redox Signal.* **2020**, *32*, 447–462. <https://doi.org/10.1089/ars.2018.7652>.
377. Anso, E.; Weinberg, S.E.; Diebold, L.P.; Thompson, B.J.; Malinge, S.; Schumacker, P.T.; Liu, X.; Zhang, Y.; Shao, Z.; Steadman, M.; et al. The mitochondrial respiratory chain is essential for haematopoietic stem cell function. *Nat. Cell Biol.* **2017**, *19*, 614–625. <https://doi.org/10.1038/ncb3529>.
378. Hughes, B.G.; Hekimi, S. A mild impairment of mitochondrial electron transport has sex-specific effects on lifespan and aging in mice. *PLoS ONE* **2011**, *6*, e26116. <https://doi.org/10.1371/journal.pone.0026116>.
379. Waypa, G.B.; Smith, K.A.; Mungai, P.T.; Dudley, V.J.; Helmin, K.A.; Singer, B.D.; Peek, C.B.; Bass, J.; Nelson, L.; Shah, S.J.; et al. Mitochondria regulate proliferation in adult cardiac myocytes. *J. Clin. Investig.* **2024**, *134*, e165482. <https://doi.org/10.1172/JCI165482>.
380. Horvath, A.; Horakova, E.; Dunajcikova, P.; Verner, Z.; Pravdova, E.; Slapetova, I.; Cuninkova, L.; Lukes, J. Downregulation of the nuclear-encoded subunits of the complexes III and IV disrupts their respective complexes but not complex I in procyclic *Trypanosoma brucei*. *Mol. Microbiol.* **2005**, *58*, 116–130. <https://doi.org/10.1111/j.1365-2958.2005.04813.x>.
381. Owens, K.M.; Kulawiec, M.; Desouki, M.M.; Vanniarajan, A.; Singh, K.K. Impaired OXPHOS complex III in breast cancer. *PLoS ONE* **2011**, *6*, e23846. <https://doi.org/10.1371/journal.pone.0023846>.
382. Kadenbach, B.; Huttemann, M. The subunit composition and function of mammalian cytochrome c oxidase. *Mitochondrion* **2015**, *24*, 64–76. <https://doi.org/10.1016/j.mito.2015.07.002>.
383. Zhou, T.; Chien, M.S.; Kaleem, S.; Matsunami, H. Single cell transcriptome analysis of mouse carotid body glomus cells. *J. Physiol.* **2016**, *594*, 4225–4251. <https://doi.org/10.1113/JP271936>.
384. Gao, L.; Bonilla-Henao, V.; Garcia-Flores, P.; Arias-Mayenco, I.; Ortega-Saenz, P.; Lopez-Barneo, J. Gene expression analyses reveal metabolic specifications in acute O_2 -sensing chemoreceptor cells. *J. Physiol.* **2017**, *595*, 6091–6120. <https://doi.org/10.1113/JP274684>.
385. Weissmann, N.; Kuzkaya, N.; Fuchs, B.; Tiyerili, V.; Schafer, R.U.; Schutte, H.; Ghofrani, H.A.; Schermuly, R.T.; Schudt, C.; Sydykov, A.; et al. Detection of reactive oxygen species in isolated, perfused lungs by electron spin resonance spectroscopy. *Respir. Res.* **2005**, *6*, 86. <https://doi.org/10.1186/1465-9921-6-86>.
386. Murphy, M.P.; Bayir, H.; Belousov, V.; Chang, C.J.; Davies, K.J.A.; Davies, M.J.; Dick, T.P.; Finkel, T.; Forman, H.J.; Janssen-Heininger, Y.; et al. Guidelines for measuring reactive oxygen species and oxidative damage in cells and in vivo. *Nat. Metab.* **2022**, *4*, 651–662. <https://doi.org/10.1038/s42255-022-00591-z>.
387. Dry, I.B.; Moore, A.L.; Day, D.A.; Wiskich, J.T. Regulation of alternative pathway activity in plant mitochondria: Nonlinear relationship between electron flux and the redox poise of the quinone pool. *Arch. Biochem. Biophys.* **1989**, *273*, 148–157. [https://doi.org/10.1016/0003-9861\(89\)90173-2](https://doi.org/10.1016/0003-9861(89)90173-2).
388. El-Khoury, R.; Dufour, E.; Rak, M.; Ramanantsoa, N.; Grandchamp, N.; Csaba, Z.; Duvillie, B.; Benit, P.; Gallego, J.; Gressens, P.; et al. Alternative oxidase expression in the mouse enables bypassing cytochrome c oxidase blockade and limits mitochondrial ROS overproduction. *PLoS Genet.* **2013**, *9*, e1003182. <https://doi.org/10.1371/journal.pgen.1003182>.
389. Maxwell, D.P.; Wang, Y.; McIntosh, L. The alternative oxidase lowers mitochondrial reactive oxygen production in plant cells. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 8271–8276. <https://doi.org/10.1073/pnas.96.14.8271>.
390. Moreno-Dominguez, A.; Ortega-Saenz, P.; Gao, L.; Colinas, O.; Garcia-Flores, P.; Bonilla-Henao, V.; Aragones, J.; Huttemann, M.; Grossman, L.I.; Weissmann, N.; et al. Acute O_2 sensing through HIF2 α -dependent expression of atypical cytochrome oxidase subunits in arterial chemoreceptors. *Sci. Signal.* **2020**, *13*, eaay9452. <https://doi.org/10.1126/scisignal.aay9452>.
391. Tanaka, H.; Nishimaru, K.; Aikawa, T.; Hirayama, W.; Tanaka, Y.; Shigenobu, K. Effect of SEA0400, a novel inhibitor of sodium-calcium exchanger, on myocardial ionic currents. *Br. J. Pharmacol.* **2002**, *135*, 1096–1100. <https://doi.org/10.1038/sj.bjp.0704574>.
392. Ruiz, A.; Alberdi, E.; Matute, C. CGP37157, an inhibitor of the mitochondrial $\text{Na}^+/\text{Ca}^{2+}$ exchanger, protects neurons from excitotoxicity by blocking voltage-gated Ca^{2+} channels. *Cell Death Dis.* **2014**, *5*, e1156. <https://doi.org/10.1038/cddis.2014.134>.
393. Friedman, S.M. The effects of external sodium substitution on cell sodium and potassium in vascular smooth muscle. *J. Physiol.* **1977**, *270*, 195–208. <https://doi.org/10.1113/jphysiol.1977.sp011946>.
394. Nita, I.I.; Hershfinkel, M.; Lewis, E.C.; Sekler, I. A crosstalk between Na^+ channels, Na^+/K^+ pump and mitochondrial Na^+ transporters controls glucose-dependent cytosolic and mitochondrial Na^+ signals. *Cell Calcium* **2015**, *57*, 69–75. <https://doi.org/10.1016/j.ceca.2014.12.007>.

395. Majander, A.; Finel, M.; Wikstrom, M. Diphenyleneiodonium inhibits reduction of iron-sulfur clusters in the mitochondrial NADH-ubiquinone oxidoreductase (Complex I). *J. Biol. Chem.* **1994**, *269*, 21037–21042.
396. Mohazzab, K.M.; Fayngersh, R.P.; Kaminski, P.M.; Wolin, M.S. Potential role of NADH oxidoreductase-derived reactive O₂ species in calf pulmonary arterial PO₂-elicited responses. *Am. J. Physiol.* **1995**, *269*, L637–L644. <https://doi.org/10.1152/ajplung.1995.269.5.L637>.
397. Grimminger, F.; Weissmann, N.; Spriestersbach, R.; Becker, E.; Rosseau, S.; Seeger, W. Effects of NADPH oxidase inhibitors on hypoxic vasoconstriction in buffer-perfused rabbit lungs. *Am. J. Physiol.* **1995**, *268*, L747–L752. <https://doi.org/10.1152/ajplung.1995.268.5.L747>.
398. Diatchuk, V.; Lotan, O.; Koshkin, V.; Wikstroem, P.; Pick, E. Inhibition of NADPH oxidase activation by 4-(2-aminoethyl)-benzenesulfonyl fluoride and related compounds. *J. Biol. Chem.* **1997**, *272*, 13292–13301. <https://doi.org/10.1074/jbc.272.20.13292>.
399. Lahiri, S.; Ehleben, W.; Acker, H. Chemoreceptor discharges and cytochrome redox changes of the rat carotid body: Role of heme ligands. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 9427–9432. <https://doi.org/10.1073/pnas.96.16.9427>.
400. Rathore, R.; Zheng, Y.M.; Li, X.Q.; Wang, Q.S.; Liu, Q.H.; Ginnan, R.; Singer, H.A.; Ho, Y.S.; Wang, Y.X. Mitochondrial ROS-PKCepsilon signaling axis is uniquely involved in hypoxic increase in [Ca²⁺]_i in pulmonary artery smooth muscle cells. *Biochem. Biophys. Res. Commun.* **2006**, *351*, 784–790. <https://doi.org/10.1016/j.bbrc.2006.10.116>.
401. Cogolludo, A.; Moreno, L.; Frazziano, G.; Moral-Sanz, J.; Menendez, C.; Castaneda, J.; Gonzalez, C.; Villamor, E.; Perez-Vizcaino, F. Activation of neutral sphingomyelinase is involved in acute hypoxic pulmonary vasoconstriction. *Cardiovasc. Res.* **2009**, *82*, 296–302. <https://doi.org/10.1093/cvr/cvn349>.
402. Moral-Sanz, J.; Gonzalez, T.; Menendez, C.; David, M.; Moreno, L.; Macias, A.; Cortijo, J.; Valenzuela, C.; Perez-Vizcaino, F.; Cogolludo, A. Ceramide inhibits Kv currents and contributes to TP-receptor-induced vasoconstriction in rat and human pulmonary arteries. *Am. J. Physiol. Cell Physiol.* **2011**, *301*, C186–C194. <https://doi.org/10.1152/ajpcell.00243.2010>.
403. Wu-Zhang, A.X.; Newton, A.C. Protein kinase C pharmacology: Refining the toolbox. *Biochem. J.* **2013**, *452*, 195–209. <https://doi.org/10.1042/BJ20130220>.
404. Veith, C.; Kraut, S.; Wilhelm, J.; Sommer, N.; Quanz, K.; Seeger, W.; Brandes, R.P.; Weissmann, N.; Schroder, K. NADPH oxidase 4 is not involved in hypoxia-induced pulmonary hypertension. *Pulm. Circ.* **2016**, *6*, 397–400. <https://doi.org/10.1086/687756>.
405. Murtaza, G.; Paddenberger, R.; Pfeil, U.; Goldenberg, A.; Mermer, P.; Kummer, W. Hypoxia-induced pulmonary vasoconstriction of intra-acinar arteries is impaired in NADPH oxidase 4 gene-deficient mice. *Pulm. Circ.* **2018**, *8*, 2045894018808240. <https://doi.org/10.1177/2045894018808240>.
406. Hales, C.A. The site and mechanism of oxygen sensing for the pulmonary vessels. *Chest* **1985**, *88*, 235S–240S. https://doi.org/10.1378/chest.88.4_supplement.235s.
407. Nagaraj, C.; Tabeling, C.; Nagy, B.M.; Jain, P.P.; Marsh, L.M.; Papp, R.; Pienn, M.; Witzernath, M.; Ghanim, B.; Klepetko, W.; et al. Hypoxic vascular response and ventilation/perfusion matching in end-stage COPD may depend on p22phox. *Eur. Respir. J.* **2017**, *50*, 1601651. <https://doi.org/10.1183/13993003.01651-2016>.
408. Littler, C.M.; Morris, K.G., Jr.; Fagan, K.A.; McMurtry, I.F.; Messing, R.O.; Dempsey, E.C. Protein kinase C-epsilon-null mice have decreased hypoxic pulmonary vasoconstriction. *Am. J. Physiol. Heart Circ. Physiol.* **2003**, *284*, H1321–H1331. <https://doi.org/10.1152/ajpheart.00795.2002>.
409. Robertson, T.P.; Aaronson, P.I.; Ward, J.P. Hypoxic vasoconstriction and intracellular Ca²⁺ in pulmonary arteries: Evidence for PKC-independent Ca²⁺ sensitization. *Am. J. Physiol.* **1995**, *268*, H301–H307. <https://doi.org/10.1152/ajpheart.1995.268.1.H301>.
410. Stuart, J.A.; Fonseca, J.; Moradi, F.; Cunningham, C.; Seliman, B.; Worsfold, C.R.; Dolan, S.; Abando, J.; Maddalena, L.A. How Supraphysiological Oxygen Levels in Standard Cell Culture Affect Oxygen-Consuming Reactions. *Oxid. Med. Cell. Longev.* **2018**, *2018*, 8238459. <https://doi.org/10.1155/2018/8238459>.
411. Wang, X.T.; McCullough, K.D.; Wang, X.J.; Carpenter, G.; Holbrook, N.J. Oxidative stress-induced phospholipase C-gamma 1 activation enhances cell survival. *J. Biol. Chem.* **2001**, *276*, 28364–28371. <https://doi.org/10.1074/jbc.M102693200>.
412. Strielkov, I.V.; Kizub, I.V.; Khromov, A.S.; Soloviev, A.I. Evidence for the role of phosphatidylcholine-specific phospholipase C in sustained hypoxic pulmonary vasoconstriction. *Vasc. Pharmacol.* **2013**, *58*, 292–298. <https://doi.org/10.1016/j.vph.2013.02.002>.
413. Liao, B.; Zheng, Y.M.; Yadav, V.R.; Korde, A.S.; Wang, Y.X. Hypoxia induces intracellular Ca²⁺ release by causing reactive oxygen species-mediated dissociation of FK506-binding protein 12.6 from ryanodine receptor 2 in pulmonary artery myocytes. *Antioxid. Redox Signal.* **2011**, *14*, 37–47. <https://doi.org/10.1089/ars.2009.3047>.
414. Chen, T.X.; Xu, X.Y.; Zhao, Z.; Zhao, F.Y.; Gao, Y.M.; Yan, X.H.; Wan, Y. Hydrogen peroxide is a critical regulator of the hypoxia-induced alterations of store-operated Ca(2+) entry into rat pulmonary arterial smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2017**, *312*, L477–L487. <https://doi.org/10.1152/ajplung.00138.2016>.

415. Lin, M.J.; Yang, X.R.; Cao, Y.N.; Sham, J.S. Hydrogen peroxide-induced Ca^{2+} mobilization in pulmonary arterial smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2007**, *292*, L1598–L1608. <https://doi.org/10.1152/ajplung.00323.2006>.
416. Park, J.M.; Do, V.Q.; Seo, Y.S.; Kim, H.J.; Nam, J.H.; Yin, M.Z.; Kim, H.J.; Kim, S.J.; Griendling, K.K.; Lee, M.Y. NADPH Oxidase 1 Mediates Acute Blood Pressure Response to Angiotensin II by Contributing to Calcium Influx in Vascular Smooth Muscle Cells. *Arterioscler. Thromb. Vasc. Biol.* **2022**, *42*, e117–e130. <https://doi.org/10.1161/ATVBAHA.121.317239>.
417. Maier, T.; Follmann, M.; Hessler, G.; Kleemann, H.W.; Hachtel, S.; Fuchs, B.; Weissmann, N.; Linz, W.; Schmidt, T.; Lohn, M.; et al. Discovery and pharmacological characterization of a novel potent inhibitor of diacylglycerol-sensitive TRPC cation channels. *Br. J. Pharmacol.* **2015**, *172*, 3650–3660. <https://doi.org/10.1111/bph.13151>.
418. Knock, G.A.; Ward, J.P. Redox regulation of protein kinases as a modulator of vascular function. *Antioxid. Redox Signal.* **2011**, *15*, 1531–1547. <https://doi.org/10.1089/ars.2010.3614>.
419. Knock, G.A.; Snetkov, V.A.; Shaifta, Y.; Drndarski, S.; Ward, J.P.; Aaronson, P.I. Role of src-family kinases in hypoxic vasoconstriction of rat pulmonary artery. *Cardiovasc. Res.* **2008**, *80*, 453–462. <https://doi.org/10.1093/cvr/cvn209>.
420. Knock, G.A.; Snetkov, V.A.; Shaifta, Y.; Connolly, M.; Drndarski, S.; Noah, A.; Pourmahram, G.E.; Becker, S.; Aaronson, P.I.; Ward, J.P. Superoxide constricts rat pulmonary arteries via Rho-kinase-mediated Ca^{2+} sensitization. *Free Radic. Biol. Med.* **2009**, *46*, 633–642. <https://doi.org/10.1016/j.freeradbiomed.2008.11.015>.
421. MacKay, C.E.; Shaifta, Y.; Snetkov, V.V.; Francois, A.A.; Ward, J.P.T.; Knock, G.A. ROS-dependent activation of RhoA/Rho-kinase in pulmonary artery: Role of Src-family kinases and ARHGEF1. *Free Radic. Biol. Med.* **2017**, *110*, 316–331. <https://doi.org/10.1016/j.freeradbiomed.2017.06.022>.
422. Forman, H.J.; Ursini, F.; Maiorino, M. An overview of mechanisms of redox signaling. *J. Mol. Cell. Cardiol.* **2014**, *73*, 2–9. <https://doi.org/10.1016/j.yjmcc.2014.01.018>.
423. Pak, O.; Scheibe, S.; Esfandiary, A.; Gierhardt, M.; Sydykov, A.; Logan, A.; Fysikopoulos, A.; Veit, F.; Hecker, M.; Kroschel, F.; et al. Impact of the mitochondria-targeted antioxidant MitoQ on hypoxia-induced pulmonary hypertension. *Eur. Respir. J.* **2018**, *51*, 1701024. <https://doi.org/10.1183/13993003.01024-2017>.
424. Jernigan, N.L.; Resta, T.C.; Gonzalez Bosc, L.V. Altered Redox Balance in the Development of Chronic Hypoxia-induced Pulmonary Hypertension. *Adv. Exp. Med. Biol.* **2017**, *967*, 83–103. https://doi.org/10.1007/978-3-319-63245-2_7.
425. Pourmahram, G.E.; Snetkov, V.A.; Shaifta, Y.; Drndarski, S.; Knock, G.A.; Aaronson, P.I.; Ward, J.P. Constriction of pulmonary artery by peroxide: Role of Ca^{2+} release and PKC. *Free Radic. Biol. Med.* **2008**, *45*, 1468–1476. <https://doi.org/10.1016/j.freeradbiomed.2008.08.020>.
426. Genet, N.; Billaud, M.; Rossignol, R.; Dubois, M.; Gillibert-Duplantier, J.; Isakson, B.E.; Marthan, R.; Savineau, J.P.; Guibert, C. Signaling Pathways Linked to Serotonin-Induced Superoxide Anion Production: A Physiological Role for Mitochondria in Pulmonary Arteries. *Front. Physiol.* **2017**, *8*, 76. <https://doi.org/10.3389/fphys.2017.00076>.
427. Poyton, R.O.; Castello, P.R.; Ball, K.A.; Woo, D.K.; Pan, N. Mitochondria and hypoxic signaling: A new view. *Ann. N. Y. Acad. Sci.* **2009**, *1177*, 48–56. <https://doi.org/10.1111/j.1749-6632.2009.05046.x>.
428. Martinez-Ruiz, A.; Cadenas, S.; Lamas, S. Nitric oxide signaling: Classical, less classical, and nonclassical mechanisms. *Free Radic. Biol. Med.* **2011**, *51*, 17–29. <https://doi.org/10.1016/j.freeradbiomed.2011.04.010>.
429. Viner, R.I.; Williams, T.D.; Schoneich, C. Peroxynitrite modification of protein thiols: Oxidation, nitrosylation, and S-glutathionylation of functionally important cysteine residue(s) in the sarcoplasmic reticulum Ca-ATPase. *Biochemistry* **1999**, *38*, 12408–12415. <https://doi.org/10.1021/bi9909445>.
430. Minetti, M.; Mallozzi, C.; Di Stasi, A.M. Peroxynitrite activates kinases of the src family and upregulates tyrosine phosphorylation signaling. *Free Radic. Biol. Med.* **2002**, *33*, 744–754. [https://doi.org/10.1016/s0891-5849\(02\)00891-2](https://doi.org/10.1016/s0891-5849(02)00891-2).
431. Lock, J.T.; Sinkins, W.G.; Schilling, W.P. Effect of protein S-glutathionylation on Ca^{2+} homeostasis in cultured aortic endothelial cells. *Am. J. Physiol. Heart Circ. Physiol.* **2011**, *300*, H493–H506. <https://doi.org/10.1152/ajpheart.01073.2010>.
432. Sun, J.; Yamaguchi, N.; Xu, L.; Eu, J.P.; Stamler, J.S.; Meissner, G. Regulation of the cardiac muscle ryanodine receptor by O_2 tension and S-nitrosoglutathione. *Biochemistry* **2008**, *47*, 13985–13990. <https://doi.org/10.1021/bi8012627>.
433. Flores-Tamez, V.; Escalante, B.; Rios, A. Peroxynitrite-Induced Intracellular Ca^{2+} Depression in Cardiac Myocytes: Role of Sarco/Endoplasmic Reticulum Ca^{2+} Pump. *Folia Biol.* **2019**, *65*, 237–245. <https://doi.org/10.14712/fb2019065050237>.
434. Evans, A.M.; Mustard, K.J.; Wyatt, C.N.; Peers, C.; Dipp, M.; Kumar, P.; Kinnear, N.P.; Hardie, D.G. Does AMP-activated protein kinase couple inhibition of mitochondrial oxidative phosphorylation by hypoxia to calcium signaling in O_2 -sensing cells? *J. Biol. Chem.* **2005**, *280*, 41504–41511. <https://doi.org/10.1074/jbc.M510040200>.

435. Evans, A.M.; Lewis, S.A.; Ogunbayo, O.A.; Moral-Sanz, J. Modulation of the LKB1-AMPK Signalling Pathway Underpins Hypoxic Pulmonary Vasoconstriction and Pulmonary Hypertension. *Adv. Exp. Med. Biol.* **2015**, *860*, 89–99. https://doi.org/10.1007/978-3-319-18440-1_11.
436. Emerling, B.M.; Weinberg, F.; Snyder, C.; Burgess, Z.; Mutlu, G.M.; Viollet, B.; Budinger, G.R.; Chandel, N.S. Hypoxic activation of AMPK is dependent on mitochondrial ROS but independent of an increase in AMP/ATP ratio. *Free Radic. Biol. Med.* **2009**, *46*, 1386–1391. <https://doi.org/10.1016/j.freeradbiomed.2009.02.019>.
437. Gusarova, G.A.; Trejo, H.E.; Dada, L.A.; Briva, A.; Welch, L.C.; Hamanaka, R.B.; Mutlu, G.M.; Chandel, N.S.; Prakriya, M.; Sznajder, J.I. Hypoxia leads to Na,K-ATPase downregulation via Ca⁽²⁺⁾ release-activated Ca⁽²⁺⁾ channels and AMPK activation. *Mol. Cell. Biol.* **2011**, *31*, 3546–3556. <https://doi.org/10.1128/MCB.05114-11>.
438. Hawley, S.A.; Ross, F.A.; Chevtzoff, C.; Green, K.A.; Evans, A.; Fogarty, S.; Towler, M.C.; Brown, L.J.; Ogunbayo, O.A.; Evans, A.M.; et al. Use of cells expressing gamma subunit variants to identify diverse mechanisms of AMPK activation. *Cell Metab.* **2010**, *11*, 554–565. <https://doi.org/10.1016/j.cmet.2010.04.001>.
439. Choi, S.L.; Kim, S.J.; Lee, K.T.; Kim, J.; Mu, J.; Birnbaum, M.J.; Soo Kim, S.; Ha, J. The regulation of AMP-activated protein kinase by H₂O₂. *Biochem. Biophys. Res. Commun.* **2001**, *287*, 92–97. <https://doi.org/10.1006/bbrc.2001.5544>.
440. Prieto-Lloret, J.; Aaronson, P.I. Hydrogen Sulfide as an O₂ Sensor: A Critical Analysis. *Adv. Exp. Med. Biol.* **2017**, *967*, 261–276. https://doi.org/10.1007/978-3-319-63245-2_15.
441. Prieto-Lloret, J.; Shaifta, Y.; Ward, J.P.; Aaronson, P.I. Hypoxic pulmonary vasoconstriction in isolated rat pulmonary arteries is not inhibited by antagonists of H₂S-synthesizing pathways. *J. Physiol.* **2015**, *593*, 385–401. <https://doi.org/10.1113/jphysiol.2014.277046>.
442. Iciek, M.; Bilska-Wilkosz, A.; Gorny, M. Sulfane sulfur—New findings on an old topic. *Acta Biochim. Pol.* **2019**, *66*, 533–544. https://doi.org/10.18388/abp.2019_2909.
443. Olson, K.R.; Gao, Y.; DeLeon, E.R.; Markel, T.A.; Drucker, N.; Boone, D.; Whiteman, M.; Steiger, A.K.; Pluth, M.D.; Tessier, C.R.; et al. Extended hypoxia-mediated H₂S production provides for long-term oxygen sensing. *Acta Physiol.* **2020**, *228*, e13368. <https://doi.org/10.1111/apha.13368>.
444. Gubern, M.; Andriamihaja, M.; Nubel, T.; Blachier, F.; Bouillaud, F. Sulfide, the first inorganic substrate for human cells. *FASEB J.* **2007**, *21*, 1699–1706. <https://doi.org/10.1096/fj.06-7407com>.
445. Skovgaard, N.; Olson, K.R. Hydrogen sulfide mediates hypoxic vasoconstriction through a production of mitochondrial ROS in trout gills. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2012**, *303*, R487–R494. <https://doi.org/10.1152/ajpregu.00151.2012>.
446. Gupte, R.S.; Rawat, D.K.; Chettimada, S.; Cioffi, D.L.; Wolin, M.S.; Gerthoffer, W.T.; McMurtry, I.F.; Gupte, S.A. Activation of glucose-6-phosphate dehydrogenase promotes acute hypoxic pulmonary artery contraction. *J. Biol. Chem.* **2010**, *285*, 19561–19571. <https://doi.org/10.1074/jbc.M109.092916>.
447. Gupte, S.A.; Okada, T.; McMurtry, I.F.; Oka, M. Role of pentose phosphate pathway-derived NADPH in hypoxic pulmonary vasoconstriction. *Pulm. Pharmacol. Ther.* **2006**, *19*, 303–309. <https://doi.org/10.1016/j.pupt.2005.08.002>.
448. Houtkooper, R.H.; Canto, C.; Wanders, R.J.; Auwerx, J. The secret life of NAD⁺: An old metabolite controlling new metabolic signaling pathways. *Endocr. Rev.* **2010**, *31*, 194–223. <https://doi.org/10.1210/er.2009-0026>.
449. Wilson, H.L.; Dipp, M.; Thomas, J.M.; Lad, C.; Galione, A.; Evans, A.M. Adp-ribosyl cyclase and cyclic ADP-ribose hydrolase act as a redox sensor. a primary role for cyclic ADP-ribose in hypoxic pulmonary vasoconstriction. *J. Biol. Chem.* **2001**, *276*, 11180–11188. <https://doi.org/10.1074/jbc.M004849200>.
450. Moreno-Dominguez, A.; Colinas, O.; Arias-Mayenco, I.; Cabeza, J.M.; Lopez-Ogayar, J.L.; Chandel, N.S.; Weissmann, N.; Sommer, N.; Pascual, A.; Lopez-Barneo, J. Hif1alpha-dependent mitochondrial acute O₂ sensing and signaling to myocyte Ca⁽²⁺⁾ channels mediate arterial hypoxic vasodilation. *Nat. Commun.* **2024**, *15*, 6649. <https://doi.org/10.1038/s41467-024-51023-3>.
451. Dwenger, M.M.; Ohanyan, V.; Navedo, M.F.; Nystoriak, M.A. Coronary microvascular Kv1 channels as regulatory sensors of intracellular pyridine nucleotide redox potential. *Microcirculation* **2018**, *25*, e12426. <https://doi.org/10.1111/micc.12426>.
452. Torres-Torrelo, H.; Ortega-Saenz, P.; Gao, L.; Lopez-Barneo, J. Lactate sensing mechanisms in arterial chemoreceptor cells. *Nat. Commun.* **2021**, *12*, 4166. <https://doi.org/10.1038/s41467-021-24444-7>.
453. Michelakis, E.D.; Rebeyka, I.; Wu, X.; Nsair, A.; Thebaud, B.; Hashimoto, K.; Dyck, J.R.; Haromy, A.; Harry, G.; Barr, A.; et al. O₂ sensing in the human ductus arteriosus: Regulation of voltage-gated K⁺ channels in smooth muscle cells by a mitochondrial redox sensor. *Circ. Res.* **2002**, *91*, 478–486. <https://doi.org/10.1161/01.res.0000035057.63303.d1>.
454. Hirst, J.; King, M.S.; Pryde, K.R. The production of reactive oxygen species by complex I. *Biochem. Soc. Trans.* **2008**, *36*, 976–980. <https://doi.org/10.1042/BST0360976>.

455. Quinlan, C.L.; Perevoshchikova, I.V.; Hey-Mogensen, M.; Orr, A.L.; Brand, M.D. Sites of reactive oxygen species generation by mitochondria oxidizing different substrates. *Redox Biol.* **2013**, *1*, 304–312. <https://doi.org/10.1016/j.redox.2013.04.005>.
456. Reeve, H.L.; Tolarova, S.; Nelson, D.P.; Archer, S.; Weir, E.K. Redox control of oxygen sensing in the rabbit ductus arteriosus. *J. Physiol.* **2001**, *533*, 253–261. <https://doi.org/10.1111/j.1469-7793.2001.0253b.x>.
457. Bonnet, S.; Michelakis, E.D.; Porter, C.J.; Andrade-Navarro, M.A.; Thebaud, B.; Bonnet, S.; Haromy, A.; Harry, G.; Moudgil, R.; McMurtry, M.S.; et al. An abnormal mitochondrial-hypoxia inducible factor-1 α -Kv channel pathway disrupts oxygen sensing and triggers pulmonary arterial hypertension in fawn hooded rats: Similarities to human pulmonary arterial hypertension. *Circulation* **2006**, *113*, 2630–2641. <https://doi.org/10.1161/CIRCULATIONAHA.105.609008>.
458. Ahmad, M.; Kelly, M.R.; Zhao, X.; Kandhi, S.; Wolin, M.S. Roles for Nox4 in the contractile response of bovine pulmonary arteries to hypoxia. *Am. J. Physiol. Heart Circ. Physiol.* **2010**, *298*, H1879–H1888. <https://doi.org/10.1152/ajpheart.01228.2009>.
459. Paky, A.; Michael, J.R.; Burke-Wolin, T.M.; Wolin, M.S.; Gurtner, G.H. Endogenous production of superoxide by rabbit lungs: Effects of hypoxia or metabolic inhibitors. *J. Appl. Physiol.* **1993**, *74*, 2868–2874. <https://doi.org/10.1152/jappl.1993.74.6.2868>.
460. Mathew, R.; Burke-Wolin, T.; Gewitz, M.H.; Wolin, M.S. O₂ and rat pulmonary artery tone: Effects of endothelium, Ca²⁺, cyanide, and monocalcium. *J. Appl. Physiol.* **1991**, *71*, 30–36. <https://doi.org/10.1152/jappl.1991.71.1.30>.
461. Young, T.A.; Cunningham, C.C.; Bailey, S.M. Reactive oxygen species production by the mitochondrial respiratory chain in isolated rat hepatocytes and liver mitochondria: Studies using myxothiazol. *Arch. Biochem. Biophys.* **2002**, *405*, 65–72. [https://doi.org/10.1016/s0003-9861\(02\)00338-7](https://doi.org/10.1016/s0003-9861(02)00338-7).
462. Starkov, A.A.; Fiskum, G. Myxothiazol induces H₂O₂ production from mitochondrial respiratory chain. *Biochem. Biophys. Res. Commun.* **2001**, *281*, 645–650. <https://doi.org/10.1006/bbrc.2001.4409>.
463. Watson, M.A.; Wong, H.S.; Brand, M.D. Use of S1QELs and S3QELs to link mitochondrial sites of superoxide and hydrogen peroxide generation to physiological and pathological outcomes. *Biochem. Soc. Trans.* **2019**, *47*, 1461–1469. <https://doi.org/10.1042/BST20190305>.
464. Kilbride, S.M.; Telford, J.E.; Davey, G.P. Complex I Controls Mitochondrial and Plasma Membrane Potentials in Nerve Terminals. *Neurochem. Res.* **2021**, *46*, 100–107. <https://doi.org/10.1007/s11064-020-02990-8>.
465. Cheung, J.Y.; Wang, J.; Zhang, X.Q.; Song, J.; Davidyock, J.M.; Prado, F.J.; Shanmughapriya, S.; Worth, A.M.; Madesh, M.; Judenharc-Haouzi, A.; et al. Methylene Blue Counteracts H₂S-Induced Cardiac Ion Channel Dysfunction and ATP Reduction. *Cardiovasc. Toxicol.* **2018**, *18*, 407–419. <https://doi.org/10.1007/s12012-018-9451-5>.
466. Cheranov, S.Y.; Jaggar, J.H. Mitochondrial modulation of Ca²⁺ sparks and transient K_{Ca} currents in smooth muscle cells of rat cerebral arteries. *J. Physiol.* **2004**, *556*, 755–771. <https://doi.org/10.1113/jphysiol.2003.059568>.
467. McCarron, J.G.; Olson, M.L.; Wilson, C.; Sandison, M.E.; Chalmers, S. Examining the role of mitochondria in Ca²⁺(⁺) signaling in native vascular smooth muscle. *Microcirculation* **2013**, *20*, 317–329. <https://doi.org/10.1111/micc.12039>.
468. Sward, K.; Dreja, K.; Lindqvist, A.; Persson, E.; Hellstrand, P. Influence of mitochondrial inhibition on global and local [Ca²⁺](I) in rat tail artery. *Circ. Res.* **2002**, *90*, 792–799. <https://doi.org/10.1161/01.res.0000015214.40360.84>.
469. Doege, K.; Heine, S.; Jensen, I.; Jelkmann, W.; Metzen, E. Inhibition of mitochondrial respiration elevates oxygen concentration but leaves regulation of hypoxia-inducible factor (HIF) intact. *Blood* **2005**, *106*, 2311–2317. <https://doi.org/10.1182/blood-2005-03-1138>.
470. Potter, M.; Badder, L.; Hoade, Y.; Johnston, I.G.; Morten, K.J. Monitoring Intracellular Oxygen Concentration: Implications for Hypoxia Studies and Real-Time Oxygen Monitoring. *Adv. Exp. Med. Biol.* **2016**, *876*, 257–263. https://doi.org/10.1007/978-1-4939-3023-4_32.
471. Hagen, T.; Taylor, C.T.; Lam, F.; Moncada, S. Redistribution of intracellular oxygen in hypoxia by nitric oxide: Effect on HIF1 α . *Science* **2003**, *302*, 1975–1978. <https://doi.org/10.1126/science.1088805>.
472. Wilcox, C.S.; Pearlman, A. Chemistry and antihypertensive effects of tempol and other nitroxides. *Pharmacol. Rev.* **2008**, *60*, 418–469. <https://doi.org/10.1124/pr.108.000240>.
473. Oyewole, A.O.; Birch-Machin, M.A. Mitochondria-targeted antioxidants. *FASEB J.* **2015**, *29*, 4766–4771. <https://doi.org/10.1096/fj.15-275404>.
474. Pedre, B.; Barayeu, U.; Ezerina, D.; Dick, T.P. The mechanism of action of N-acetylcysteine (NAC): The emerging role of H₂S and sulfane sulfur species. *Pharmacol. Ther.* **2021**, *228*, 107916. <https://doi.org/10.1016/j.pharmthera.2021.107916>.
475. Barbosa, N.V.; Nogueira, C.W.; Nogara, P.A.; de Bem, A.F.; Aschner, M.; Rocha, J.B.T. Organoselenium compounds as mimics of selenoproteins and thiol modifier agents. *Metallomics* **2017**, *9*, 1703–1734. <https://doi.org/10.1039/c7mt00083a>.

476. Olson, K.R.; Gao, Y.; Steiger, A.K.; Pluth, M.D.; Tessier, C.R.; Markel, T.A.; Boone, D.; Stahelin, R.V.; Batinic-Haberle, I.; Straubg, K.D. Effects of Manganese Porphyrins on Cellular Sulfur Metabolism. *Molecules* **2020**, *25*, 980. <https://doi.org/10.3390/molecules25040980>.
477. Batinic-Haberle, I.; Tovmasyan, A.; Spasojevic, I. An educational overview of the chemistry, biochemistry and therapeutic aspects of Mn porphyrins--From superoxide dismutation to H₂O₂-driven pathways. *Redox Biol.* **2015**, *5*, 43–65. <https://doi.org/10.1016/j.redox.2015.01.017>.
478. Silveira, L.R.; Pereira-Da-Silva, L.; Juel, C.; Hellsten, Y. Formation of hydrogen peroxide and nitric oxide in rat skeletal muscle cells during contractions. *Free Radic. Biol. Med.* **2003**, *35*, 455–464. [https://doi.org/10.1016/s0891-5849\(03\)00271-5](https://doi.org/10.1016/s0891-5849(03)00271-5).
479. Messent, M.; Griffiths, M.J.; Quinlan, G.J.; Gutteridge, J.M.; Evans, T.W. Ischaemia--reperfusion injury in the rat is modulated by superoxide generation and leads to an augmentation of the hypoxic pulmonary vascular response. *Clin. Sci.* **1996**, *90*, 47–54. <https://doi.org/10.1042/cs0900047>.
480. Shen, C.Y.; Lee, J.F.; Su, C.L.; Wang, D.; Chen, C.F. Hypoxia and reoxygenation of the lung tissues induced mRNA expressions of superoxide dismutase and catalase and interventions from different antioxidants. *Transplant. Proc.* **2008**, *40*, 2182–2184. <https://doi.org/10.1016/j.transproceed.2008.07.080>.
481. Francis, B.N.; Wilkins, M.R.; Zhao, L. Tetrahydrobiopterin and the regulation of hypoxic pulmonary vasoconstriction. *Eur. Respir. J.* **2010**, *36*, 323–330. <https://doi.org/10.1183/09031936.00188809>.
482. Ahmad, M.; Zhao, X.; Kelly, M.R.; Kandhi, S.; Perez, O.; Abraham, N.G.; Wolin, M.S. Heme oxygenase-1 induction modulates hypoxic pulmonary vasoconstriction through upregulation of ecSOD. *Am. J. Physiol. Heart Circ. Physiol.* **2009**, *297*, H1453–H1461. <https://doi.org/10.1152/ajpheart.00315.2009>.
483. Patel, D.; Alhawaj, R.; Wolin, M.S. Exposure of mice to chronic hypoxia attenuates pulmonary arterial contractile responses to acute hypoxia by increases in extracellular hydrogen peroxide. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2014**, *307*, R426–R433. <https://doi.org/10.1152/ajpregu.00257.2013>.
484. Bopp, C.; Auger, C.; Mebazaa, A.; Joshi, G.P.; Schini-Kerth, V.B.; Diemunsch, P. Urapidil, but not dihydropyridine calcium channel inhibitors, preserves the hypoxic pulmonary vasoconstriction: An experimental study in pig arteries. *Fundam. Clin. Pharmacol.* **2019**, *33*, 527–534. <https://doi.org/10.1111/fcp.12457>.
485. Demiryurek, A.T.; Wadsworth, R.M.; Kane, K.A. Pharmacological evidence for the role of mediators in hypoxia-induced vasoconstriction in sheep isolated intrapulmonary artery rings. *Eur. J. Pharmacol.* **1991**, *203*, 1–8. [https://doi.org/10.1016/0014-2999\(91\)90783-m](https://doi.org/10.1016/0014-2999(91)90783-m).
486. Weissmann, N.; Grimminger, F.; Voswinckel, R.; Conzen, J.; Seeger, W. Nitro blue tetrazolium inhibits but does not mimic hypoxic vasoconstriction in isolated rabbit lungs. *Am. J. Physiol.* **1998**, *274*, L721–L727. <https://doi.org/10.1152/ajplung.1998.274.5.L721>.
487. Mohazzab, K.M.; Wolin, M.S. Properties of a superoxide anion-generating microsomal NADH oxidoreductase, a potential pulmonary artery PO₂ sensor. *Am. J. Physiol.* **1994**, *267*, L823–L831. <https://doi.org/10.1152/ajplung.1994.267.6.L823>.
488. Du, W.; Frazier, M.; McMahon, T.J.; Eu, J.P. Redox activation of intracellular calcium release channels (ryanodine receptors) in the sustained phase of hypoxia-induced pulmonary vasoconstriction. *Chest* **2005**, *128*, 556S–558S. https://doi.org/10.1378/chest.128.6_suppl.556S.
489. Mansoori, S.; Moosavi, S.M.S.; Ketabchi, F. The Interaction between Trolox and 4,4'-diisothiocyanatostilbene-2,2'-disulfonic Acid on Hypoxic Pulmonary Vasoconstriction in the Isolated Rabbit Lung. *Iran. J. Med. Sci.* **2017**, *42*, 284–291.
490. Hodyc, D.; Snorek, M.; Brtnicky, T.; Herget, J. Superoxide dismutase mimetic tempol inhibits hypoxic pulmonary vasoconstriction in rats independently of nitric oxide production. *Exp. Physiol.* **2007**, *92*, 945–951. <https://doi.org/10.1113/expphysiol.2007.037135>.
491. Go, Y.M.; Chandler, J.D.; Jones, D.P. The cysteine proteome. *Free Radic. Biol. Med.* **2015**, *84*, 227–245. <https://doi.org/10.1016/j.freeradbiomed.2015.03.022>.
492. Meyrick, B.; Reid, L. Hypoxia-induced structural changes in the media and adventitia of the rat hilar pulmonary artery and their regression. *Am. J. Pathol.* **1980**, *100*, 151–178.
493. Burke, T.M.; Wolin, M.S. Hydrogen peroxide elicits pulmonary arterial relaxation and guanylate cyclase activation. *Am. J. Physiol.* **1987**, *252*, H721–H732. <https://doi.org/10.1152/ajpheart.1987.252.4.H721>.
494. Sheehan, D.W.; Giese, E.C.; Gugino, S.F.; Russell, J.A. Characterization and mechanisms of H₂O₂-induced contractions of pulmonary arteries. *Am. J. Physiol.* **1993**, *264*, H1542–H1547. <https://doi.org/10.1152/ajpheart.1993.264.5.H1542>.
495. Jin, N.; Rhoades, R.A. Activation of tyrosine kinases in H₂O₂-induced contraction in pulmonary artery. *Am. J. Physiol.* **1997**, *272*, H2686–H2692. <https://doi.org/10.1152/ajpheart.1997.272.6.H2686>.

496. Jones, R.D.; Thompson, J.S.; Morice, A.H. The effect of hydrogen peroxide on hypoxia, prostaglandin F₂ alpha and potassium chloride induced contractions in isolated rat pulmonary arteries. *Pulm. Pharmacol. Ther.* **1997**, *10*, 37–42. <https://doi.org/10.1006/pupt.1997.0071>.
497. Pelaez, N.J.; Braun, T.R.; Paul, R.J.; Meiss, R.A.; Packer, C.S. H₂O₂ mediates Ca²⁺- and MLC(20) phosphorylation-independent contraction in intact and permeabilized vascular muscle. *Am. J. Physiol. Heart Circ. Physiol.* **2000**, *279*, H1185–H1193. <https://doi.org/10.1152/ajpheart.2000.279.3.H1185>.
498. Kerkar, S.; Speyer, C.; Tyburski, J.; Steffes, C. Reactive oxygen metabolites induce a biphasic contractile response in microvascular lung pericytes. *J. Trauma* **2001**, *51*, 440–445. <https://doi.org/10.1097/00005373-200109000-00003>.
499. Nagi, M.N.; Mansour, M.A.; Al-Shabanah, O.A.; El-Kashef, H.A. Melatonin inhibits the contractile effect of vanadate in the isolated pulmonary arterial rings of rats: Possible role of hydrogen peroxide. *J. Biochem. Mol. Toxicol.* **2002**, *16*, 273–278. <https://doi.org/10.1002/jbt.10049>.
500. Oeckler, R.A.; Arcuino, E.; Ahmad, M.; Olson, S.C.; Wolin, M.S. Cytosolic NADH redox and thiol oxidation regulate pulmonary arterial force through ERK MAP kinase. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2005**, *288*, L1017–L1025. <https://doi.org/10.1152/ajplung.00223.2004>.
501. Maki, J.; Hirano, M.; Hoka, S.; Kanaide, H.; Hirano, K. Involvement of reactive oxygen species in thrombin-induced pulmonary vasoconstriction. *Am. J. Respir. Crit. Care Med.* **2010**, *182*, 1435–1444. <https://doi.org/10.1164/rccm.201002-0255OC>.
502. Neo, B.H.; Kandhi, S.; Wolin, M.S. Roles for soluble guanylate cyclase and a thiol oxidation-elicited subunit dimerization of protein kinase G in pulmonary artery relaxation to hydrogen peroxide. *Am. J. Physiol. Heart Circ. Physiol.* **2010**, *299*, H1235–H1241. <https://doi.org/10.1152/ajpheart.00513.2010>.
503. Zaloudikova, M.; Herget, J.; Vizek, M. The contractile response of isolated small pulmonary arteries induced by activated macrophages. *Physiol. Res.* **2014**, *63*, 267–270. <https://doi.org/10.33549/physiolres.932698>.
504. Lucchesi, P.A.; Belmadani, S.; Matrougui, K. Hydrogen peroxide acts as both vasodilator and vasoconstrictor in the control of perfused mouse mesenteric resistance arteries. *J. Hypertens.* **2005**, *23*, 571–579. <https://doi.org/10.1097/01.hjh.0000160214.40855.79>.
505. Thakali, K.; Davenport, L.; Fink, G.D.; Watts, S.W. Pleiotropic effects of hydrogen peroxide in arteries and veins from normotensive and hypertensive rats. *Hypertension* **2006**, *47*, 482–487. <https://doi.org/10.1161/01.HYP.0000201540.91234.8f>.
506. Neo, B.H.; Kandhi, S.; Wolin, M.S. Roles for redox mechanisms controlling protein kinase G in pulmonary and coronary artery responses to hypoxia. *Am. J. Physiol. Heart Circ. Physiol.* **2011**, *301*, H2295–H2304. <https://doi.org/10.1152/ajpheart.00624.2011>.
507. Zhang, D.X.; Borbouse, L.; Gebremedhin, D.; Mendoza, S.A.; Zinkevich, N.S.; Li, R.; Gutterman, D.D. H₂O₂-induced dilation in human coronary arterioles: Role of protein kinase G dimerization and large-conductance Ca²⁺-activated K⁺ channel activation. *Circ. Res.* **2012**, *110*, 471–480. <https://doi.org/10.1161/CIRCRESAHA.111.258871>.
508. Nishijima, Y.; Cao, S.; Chabowski, D.S.; Korishettar, A.; Ge, A.; Zheng, X.; Sparapani, R.; Gutterman, D.D.; Zhang, D.X. Contribution of K(V)1.5 Channel to Hydrogen Peroxide-Induced Human Arteriolar Dilation and Its Modulation by Coronary Artery Disease. *Circ. Res.* **2017**, *120*, 658–669. <https://doi.org/10.1161/CIRCRESAHA.116.309491>.
509. Gutterman, D.D.; Miura, H.; Liu, Y. Redox modulation of vascular tone: Focus of potassium channel mechanisms of dilation. *Arterioscler. Thromb. Vasc. Biol.* **2005**, *25*, 671–678. <https://doi.org/10.1161/01.ATV.0000158497.09626.3b>.
510. Matoba, T.; Shimokawa, H.; Kubota, H.; Morikawa, K.; Fujiki, T.; Kunihiro, I.; Mukai, Y.; Hirakawa, Y.; Takeshita, A. Hydrogen peroxide is an endothelium-derived hyperpolarizing factor in human mesenteric arteries. *Biochem. Biophys. Res. Commun.* **2002**, *290*, 909–913. <https://doi.org/10.1006/bbrc.2001.6278>.
511. Morgan, B.; Van Laer, K.; Owusu, T.N.; Ezerina, D.; Pastor-Flores, D.; Amponsah, P.S.; Tursch, A.; Dick, T.P. Real-time monitoring of basal H₂O₂ levels with peroxiredoxin-based probes. *Nat. Chem. Biol.* **2016**, *12*, 437–443. <https://doi.org/10.1038/nchem-bio.2067>.
512. Yang, W.; Block, E.R. Effect of hypoxia and reoxygenation on the formation and release of reactive oxygen species by porcine pulmonary artery endothelial cells. *J. Cell. Physiol.* **1995**, *164*, 414–423. <https://doi.org/10.1002/jcp.1041640222>.
513. Wojciak-Stothard, B.; Tsang, L.Y.; Haworth, S.G. Rac and Rho play opposing roles in the regulation of hypoxia/reoxygenation-induced permeability changes in pulmonary artery endothelial cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2005**, *288*, L749–L760. <https://doi.org/10.1152/ajplung.00361.2004>.
514. Munzel, T.; Afanas'ev, I.B.; Kleschyov, A.L.; Harrison, D.G. Detection of superoxide in vascular tissue. *Arterioscler. Thromb. Vasc. Biol.* **2002**, *22*, 1761–1768. <https://doi.org/10.1161/01.atv.0000034022.11764.ec>.
515. Wardman, P. Fluorescent and luminescent probes for measurement of oxidative and nitrosative species in cells and tissues: Progress, pitfalls, and prospects. *Free Radic. Biol. Med.* **2007**, *43*, 995–1022. <https://doi.org/10.1016/j.freeradbiomed.2007.06.026>.

516. Kalyanaraman, B.; Darley-Usmar, V.; Davies, K.J.; Dennery, P.A.; Forman, H.J.; Grisham, M.B.; Mann, G.E.; Moore, K.; Roberts, L.J., 2nd; Ischiropoulos, H. Measuring reactive oxygen and nitrogen species with fluorescent probes: Challenges and limitations. *Free Radic. Biol. Med.* **2012**, *52*, 1–6. <https://doi.org/10.1016/j.freeradbiomed.2011.09.030>.
517. Zielonka, J.; Kalyanaraman, B. Small-molecule luminescent probes for the detection of cellular oxidizing and nitrating species. *Free Radic. Biol. Med.* **2018**, *128*, 3–22. <https://doi.org/10.1016/j.freeradbiomed.2018.03.032>.
518. Li, Y.; Zhu, H.; Trush, M.A. Detection of mitochondria-derived reactive oxygen species production by the chemilumigenic probes lucigenin and luminol. *Biochim. Biophys. Acta* **1999**, *1428*, 1–12. [https://doi.org/10.1016/s0304-4165\(99\)00040-9](https://doi.org/10.1016/s0304-4165(99)00040-9).
519. Rembish, S.J.; Trush, M.A. Further evidence that lucigenin-derived chemiluminescence monitors mitochondrial superoxide generation in rat alveolar macrophages. *Free Radic. Biol. Med.* **1994**, *17*, 117–126. [https://doi.org/10.1016/0891-5849\(94\)90109-0](https://doi.org/10.1016/0891-5849(94)90109-0).
520. Janiszewski, M.; Souza, H.P.; Liu, X.; Pedro, M.A.; Zweier, J.L.; Laurindo, F.R. Overestimation of NADH-driven vascular oxidase activity due to lucigenin artifacts. *Free Radic. Biol. Med.* **2002**, *32*, 446–453. [https://doi.org/10.1016/s0891-5849\(01\)00828-0](https://doi.org/10.1016/s0891-5849(01)00828-0).
521. Budd, S.L.; Castilho, R.F.; Nicholls, D.G. Mitochondrial membrane potential and hydroethidine-monitored superoxide generation in cultured cerebellar granule cells. *FEBS Lett.* **1997**, *415*, 21–24. [https://doi.org/10.1016/s0014-5793\(97\)01088-0](https://doi.org/10.1016/s0014-5793(97)01088-0).
522. Adesina, S.E.; Kang, B.Y.; Bijli, K.M.; Ma, J.; Cheng, J.; Murphy, T.C.; Michael Hart, C.; Sutliff, R.L. Targeting mitochondrial reactive oxygen species to modulate hypoxia-induced pulmonary hypertension. *Free Radic. Biol. Med.* **2015**, *87*, 36–47. <https://doi.org/10.1016/j.freeradbiomed.2015.05.042>.
523. Sen, B.; Benoit, B.; Brand, M.D. Hypoxia decreases mitochondrial ROS production in cells. *Free Radic. Biol. Med.* **2024**, *224*, 1–8. <https://doi.org/10.1016/j.freeradbiomed.2024.08.016>.
524. Fang, J.; Wong, H.S.; Brand, M.D. Production of superoxide and hydrogen peroxide in the mitochondrial matrix is dominated by site I(Q) of complex I in diverse cell lines. *Redox Biol.* **2020**, *37*, 101722. <https://doi.org/10.1016/j.redox.2020.101722>.
525. Dooley, C.T.; Dore, T.M.; Hanson, G.T.; Jackson, W.C.; Remington, S.J.; Tsien, R.Y. Imaging dynamic redox changes in mammalian cells with green fluorescent protein indicators. *J. Biol. Chem.* **2004**, *279*, 22284–22293. <https://doi.org/10.1074/jbc.M312847200>.
526. Lukyanov, K.A.; Belousov, V.V. Genetically encoded fluorescent redox sensors. *Biochim. Biophys. Acta* **2014**, *1840*, 745–756. <https://doi.org/10.1016/j.bbagen.2013.05.030>.
527. Pouvreau, S. Genetically encoded reactive oxygen species (ROS) and redox indicators. *Biotechnol. J.* **2014**, *9*, 282–293. <https://doi.org/10.1002/biot.201300199>.
528. Nash, K.M.; Rockenbauer, A.; Villamena, F.A. Reactive nitrogen species reactivities with nitrones: Theoretical and experimental studies. *Chem. Res. Toxicol.* **2012**, *25*, 1581–1597. <https://doi.org/10.1021/tx200526y>.
529. Dikalov, S.I.; Polienko, Y.F.; Kirilyuk, I. Electron Paramagnetic Resonance Measurements of Reactive Oxygen Species by Cyclic Hydroxylamine Spin Probes. *Antioxid. Redox Signal.* **2018**, *28*, 1433–1443. <https://doi.org/10.1089/ars.2017.7396>.
530. Gonzalez, C.; Sanz-Alfayate, G.; Obeso, A.; Agapito, M.T. Role of glutathione redox state in oxygen sensing by carotid body chemoreceptor cells. *Methods Enzymol.* **2004**, *381*, 40–71. [https://doi.org/10.1016/S0076-6879\(04\)81003-6](https://doi.org/10.1016/S0076-6879(04)81003-6).
531. Dikalov, S.I.; Kirilyuk, I.A.; Voinov, M.; Grigor'ev, I.A. EPR detection of cellular and mitochondrial superoxide using cyclic hydroxylamines. *Free Radic. Res.* **2011**, *45*, 417–430. <https://doi.org/10.3109/10715762.2010.540242>.
532. Mills, E.L.; Kelly, B.; Logan, A.; Costa, A.S.H.; Varma, M.; Bryant, C.E.; Tourlomis, P.; Dabritz, J.H.M.; Gottlieb, E.; Latorre, I.; et al. Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria to Drive Inflammatory Macrophages. *Cell* **2016**, *167*, 457–470.e413. <https://doi.org/10.1016/j.cell.2016.08.064>.
533. Korshunov, S.S.; Skulachev, V.P.; Starkov, A.A. High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria. *FEBS Lett.* **1997**, *416*, 15–18. [https://doi.org/10.1016/s0014-5793\(97\)01159-9](https://doi.org/10.1016/s0014-5793(97)01159-9).
534. Brand, M.D. Riding the tiger—Physiological and pathological effects of superoxide and hydrogen peroxide generated in the mitochondrial matrix. *Crit. Rev. Biochem. Mol. Biol.* **2020**, *55*, 592–661. <https://doi.org/10.1080/10409238.2020.1828258>.
535. Bell, E.L.; Klimova, T.A.; Eisenbart, J.; Moraes, C.T.; Murphy, M.P.; Budinger, G.R.; Chandel, N.S. The Qo site of the mitochondrial complex III is required for the transduction of hypoxic signaling via reactive oxygen species production. *J. Cell Biol.* **2007**, *177*, 1029–1036. <https://doi.org/10.1083/jcb.200609074>.
536. Bell, E.L.; Klimova, T.A.; Eisenbart, J.; Schumacker, P.T.; Chandel, N.S. Mitochondrial reactive oxygen species trigger hypoxia-inducible factor-dependent extension of the replicative life span during hypoxia. *Mol. Cell. Biol.* **2007**, *27*, 5737–5745. <https://doi.org/10.1128/MCB.02265-06>.
537. Chandel, N.S.; Schumacker, P.T. Cellular oxygen sensing by mitochondria: Old questions, new insight. *J. Appl. Physiol.* **2000**, *88*, 1880–1889. <https://doi.org/10.1152/jappl.2000.88.5.1880>.

538. Dada, L.A.; Chandel, N.S.; Ridge, K.M.; Pedemonte, C.; Bertorello, A.M.; Sznajder, J.I. Hypoxia-induced endocytosis of Na,K-ATPase in alveolar epithelial cells is mediated by mitochondrial reactive oxygen species and PKC-zeta. *J. Clin. Investig.* **2003**, *111*, 1057–1064. <https://doi.org/10.1172/JCI16826>.
539. Hernansanz-Agustin, P.; Izquierdo-Alvarez, A.; Sanchez-Gomez, F.J.; Ramos, E.; Villa-Pina, T.; Lamas, S.; Bogdanova, A.; Martinez-Ruiz, A. Acute hypoxia produces a superoxide burst in cells. *Free Radic. Biol. Med.* **2014**, *71*, 146–156. <https://doi.org/10.1016/j.freeradbiomed.2014.03.011>.
540. Loor, G.; Kondapalli, J.; Iwase, H.; Chandel, N.S.; Waypa, G.B.; Guzy, R.D.; Vanden Hoek, T.L.; Schumacker, P.T. Mitochondrial oxidant stress triggers cell death in simulated ischemia-reperfusion. *Biochim. Biophys. Acta* **2011**, *1813*, 1382–1394. <https://doi.org/10.1016/j.bbamcr.2010.12.008>.
541. Mansfield, K.D.; Guzy, R.D.; Pan, Y.; Young, R.M.; Cash, T.P.; Schumacker, P.T.; Simon, M.C. Mitochondrial dysfunction resulting from loss of cytochrome c impairs cellular oxygen sensing and hypoxic HIF- α activation. *Cell Metab.* **2005**, *1*, 393–399. <https://doi.org/10.1016/j.cmet.2005.05.003>.
542. Mironov, S.L.; Langohr, K. Modulation of synaptic and channel activities in the respiratory network of the mice by NO/cGMP signalling pathways. *Brain Res.* **2007**, *1130*, 73–82. <https://doi.org/10.1016/j.brainres.2006.09.114>.
543. Na, N.; Chandel, N.S.; Litvan, J.; Ridge, K.M. Mitochondrial reactive oxygen species are required for hypoxia-induced degradation of keratin intermediate filaments. *FASEB J.* **2010**, *24*, 799–809. <https://doi.org/10.1096/fj.08-128967>.
544. Schafer, M.; Schafer, C.; Ewald, N.; Piper, H.M.; Noll, T. Role of redox signaling in the autonomous proliferative response of endothelial cells to hypoxia. *Circ. Res.* **2003**, *92*, 1010–1015. <https://doi.org/10.1161/01.RES.0000070882.81508.FC>.
545. Debski, D.; Smulik, R.; Zielonka, J.; Michalowski, B.; Jakubowska, M.; Debowska, K.; Adamus, J.; Marcinek, A.; Kalyanaraman, B.; Sikora, A. Mechanism of oxidative conversion of Amplex(R) Red to resorufin: Pulse radiolysis and enzymatic studies. *Free Radic. Biol. Med.* **2016**, *95*, 323–332. <https://doi.org/10.1016/j.freeradbiomed.2016.03.027>.
546. Ferrer-Sueta, G.; Radi, R. Chemical biology of peroxynitrite: Kinetics, diffusion, and radicals. *ACS Chem. Biol.* **2009**, *4*, 161–177. <https://doi.org/10.1021/cb800279q>.
547. Gupte, S.A.; Li, K.X.; Okada, T.; Sato, K.; Oka, M. Inhibitors of pentose phosphate pathway cause vasodilation: Involvement of voltage-gated potassium channels. *J. Pharmacol. Exp. Ther.* **2002**, *301*, 299–305. <https://doi.org/10.1124/jpet.301.1.299>.
548. Burke-Wolin, T.; Wolin, M.S. H₂O₂ and cGMP may function as an O₂ sensor in the pulmonary artery. *J. Appl. Physiol.* **1989**, *66*, 167–170. <https://doi.org/10.1152/jappl.1989.66.1.167>.
549. Burke-Wolin, T.M.; Wolin, M.S. Inhibition of cGMP-associated pulmonary arterial relaxation to H₂O₂ and O₂ by ethanol. *Am. J. Physiol.* **1990**, *258*, H1267–H1273. <https://doi.org/10.1152/ajpheart.1990.258.5.H1267>.
550. Omar, H.A.; Wolin, M.S. Endothelium-dependent and independent cGMP mechanisms appear to mediate O₂ responses in calf pulmonary resistance arteries. *Am. J. Physiol.* **1992**, *262*, L560–L565. <https://doi.org/10.1152/ajplung.1992.262.5.L560>.
551. Mohazzab, K.M.; Wolin, M.S. Sites of superoxide anion production detected by lucigenin in calf pulmonary artery smooth muscle. *Am. J. Physiol.* **1994**, *267*, L815–L822. <https://doi.org/10.1152/ajplung.1994.267.6.L815>.
552. Monaco, J.A.; Burke-Wolin, T. NO and H₂O₂ mechanisms of guanylate cyclase activation in oxygen-dependent responses of rat pulmonary circulation. *Am. J. Physiol.* **1995**, *268*, L546–L550. <https://doi.org/10.1152/ajplung.1995.268.4.L546>.
553. Wolin, M.S.; Burke-Wolin, T.M.; Mohazzab, H.K. Roles for NAD(P)H oxidases and reactive oxygen species in vascular oxygen sensing mechanisms. *Respir. Physiol.* **1999**, *115*, 229–238. [https://doi.org/10.1016/s0034-5687\(99\)00023-7](https://doi.org/10.1016/s0034-5687(99)00023-7).
554. Cross, A.R.; Jones, O.T. Enzymic mechanisms of superoxide production. *Biochim. Biophys. Acta* **1991**, *1057*, 281–298. [https://doi.org/10.1016/s0005-2728\(05\)80140-9](https://doi.org/10.1016/s0005-2728(05)80140-9).
555. Cross, A.R.; Henderson, L.; Jones, O.T.; Delpiano, M.A.; Hentschel, J.; Acker, H. Involvement of an NAD(P)H oxidase as a pO₂ sensor protein in the rat carotid body. *Biochem. J.* **1990**, *272*, 743–747. <https://doi.org/10.1042/bj2720743>.
556. Griendling, K.K.; Ushio-Fukai, M. NADH/NADPH Oxidase and Vascular Function. *Trends Cardiovasc. Med.* **1997**, *7*, 301–307. [https://doi.org/10.1016/S1050-1738\(97\)00088-1](https://doi.org/10.1016/S1050-1738(97)00088-1).
557. Lambeth, J.D.; Cheng, G.; Arnold, R.S.; Edens, W.A. Novel homologs of gp91phox. *Trends Biochem. Sci.* **2000**, *25*, 459–461. [https://doi.org/10.1016/s0968-0004\(00\)01658-3](https://doi.org/10.1016/s0968-0004(00)01658-3).
558. Gupte, S.A.; Kaminski, P.M.; Floyd, B.; Agarwal, R.; Ali, N.; Ahmad, M.; Edwards, J.; Wolin, M.S. Cytosolic NADPH may regulate differences in basal Nox oxidase-derived superoxide generation in bovine coronary and pulmonary arteries. *Am. J. Physiol. Heart Circ. Physiol.* **2005**, *288*, H13–H21. <https://doi.org/10.1152/ajpheart.00629.2004>.
559. Gupte, S.A.; Rupawalla, T.; Phillibert, D., Jr.; Wolin, M.S. NADPH and heme redox modulate pulmonary artery relaxation and guanylate cyclase activation by NO. *Am. J. Physiol.* **1999**, *277*, L1124–L1132. <https://doi.org/10.1152/ajplung.1999.277.6.L1124>.

560. Gupte, S.A.; Arshad, M.; Viola, S.; Kaminski, P.M.; Ungvari, Z.; Rabbani, G.; Koller, A.; Wolin, M.S. Pentose phosphate pathway coordinates multiple redox-controlled relaxing mechanisms in bovine coronary arteries. *Am. J. Physiol. Heart Circ. Physiol.* **2003**, *285*, H2316–H2326. <https://doi.org/10.1152/ajpheart.00229.2003>.
561. Mohazzab, H.K.; Kaminski, P.M.; Fayngersh, R.P.; Wolin, M.S. Oxygen-elicited responses in calf coronary arteries: Role of H₂O₂ production via NADH-derived superoxide. *Am. J. Physiol.* **1996**, *270*, H1044–H1053. <https://doi.org/10.1152/ajpheart.1996.270.3.H1044>.
562. Gupte, S.A.; Wolin, M.S. Hypoxia promotes relaxation of bovine coronary arteries through lowering cytosolic NADPH. *Am. J. Physiol. Heart Circ. Physiol.* **2006**, *290*, H2228–H2238. <https://doi.org/10.1152/ajpheart.00615.2005>.
563. Mallet, R.T.; Olivencia-Yurvati, A.H.; Bunger, R. Pyruvate enhancement of cardiac performance: Cellular mechanisms and clinical application. *Exp. Biol. Med.* **2018**, *243*, 198–210. <https://doi.org/10.1177/1535370217743919>.
564. Mingone, C.J.; Gupte, S.A.; Ali, N.; Oeckler, R.A.; Wolin, M.S. Thiol oxidation inhibits nitric oxide-mediated pulmonary artery relaxation and guanylate cyclase stimulation. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2006**, *290*, L549–L557. <https://doi.org/10.1152/ajplung.00331.2005>.
565. Wolin, M.S.; Ahmad, M.; Gupte, S.A. Oxidant and redox signaling in vascular oxygen sensing mechanisms: Basic concepts, current controversies, and potential importance of cytosolic NADPH. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2005**, *289*, L159–L173. <https://doi.org/10.1152/ajplung.00060.2005>.
566. Wolin, M.S. Reactive oxygen species and the control of vascular function. *Am. J. Physiol. Heart Circ. Physiol.* **2009**, *296*, H539–H549. <https://doi.org/10.1152/ajpheart.01167.2008>.
567. Pandolfi, P.P.; Sonati, F.; Rivi, R.; Mason, P.; Grosveld, F.; Luzzatto, L. Targeted disruption of the housekeeping gene encoding glucose 6-phosphate dehydrogenase (G6PD): G6PD is dispensable for pentose synthesis but essential for defense against oxidative stress. *EMBO J.* **1995**, *14*, 5209–5215. <https://doi.org/10.1002/j.1460-2075.1995.tb00205.x>.
568. Gupte, R.S.; Ata, H.; Rawat, D.; Abe, M.; Taylor, M.S.; Ochi, R.; Gupte, S.A. Glucose-6-phosphate dehydrogenase is a regulator of vascular smooth muscle contraction. *Antioxid. Redox Signal.* **2011**, *14*, 543–558. <https://doi.org/10.1089/ars.2010.3207>.
569. Neo, B.H.; Kandhi, S.; Ahmad, M.; Wolin, M.S. Redox regulation of guanylate cyclase and protein kinase G in vascular responses to hypoxia. *Respir. Physiol. Neurobiol.* **2010**, *174*, 259–264. <https://doi.org/10.1016/j.resp.2010.08.024>.
570. Wolin, M.S.; Burke, T.M. Hydrogen peroxide elicits activation of bovine pulmonary arterial soluble guanylate cyclase by a mechanism associated with its metabolism by catalase. *Biochem. Biophys. Res. Commun.* **1987**, *143*, 20–25. [https://doi.org/10.1016/0006-291x\(87\)90623-1](https://doi.org/10.1016/0006-291x(87)90623-1).
571. Mingone, C.J.; Gupte, S.A.; Iesaki, T.; Wolin, M.S. Hypoxia enhances a cGMP-independent nitric oxide relaxing mechanism in pulmonary arteries. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2003**, *285*, L296–L304. <https://doi.org/10.1152/ajplung.00362.2002>.
572. Adachi, T.; Weisbrod, R.M.; Pimentel, D.R.; Ying, J.; Sharov, V.S.; Schoneich, C.; Cohen, R.A. S-Glutathiolation by peroxynitrite activates SERCA during arterial relaxation by nitric oxide. *Nat. Med.* **2004**, *10*, 1200–1207. <https://doi.org/10.1038/nm1119>.
573. Burgoyne, J.R.; Oka, S.; Ale-Agha, N.; Eaton, P. Hydrogen peroxide sensing and signaling by protein kinases in the cardiovascular system. *Antioxid. Redox Signal.* **2013**, *18*, 1042–1052. <https://doi.org/10.1089/ars.2012.4817>.
574. Meurer, S.; Pioch, S.; Pabst, T.; Opitz, N.; Schmidt, P.M.; Beckhaus, T.; Wagner, K.; Matt, S.; Gegenbauer, K.; Geschka, S.; et al. Nitric oxide-independent vasodilator rescues heme-oxidized soluble guanylate cyclase from proteasomal degradation. *Circ. Res.* **2009**, *105*, 33–41. <https://doi.org/10.1161/CIRCRESAHA.109.198234>.
575. Neo, B.H.; Patel, D.; Kandhi, S.; Wolin, M.S. Roles for cytosolic NADPH redox in regulating pulmonary artery relaxation by thiol oxidation-elicited subunit dimerization of protein kinase G1alpha. *Am. J. Physiol. Heart Circ. Physiol.* **2013**, *305*, H330–H343. <https://doi.org/10.1152/ajpheart.01010.2011>.
576. Chettimada, S.; Rawat, D.K.; Dey, N.; Kobelja, R.; Simms, Z.; Wolin, M.S.; Lincoln, T.M.; Gupte, S.A. Glc-6-PD and PKG contribute to hypoxia-induced decrease in smooth muscle cell contractile phenotype proteins in pulmonary artery. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2012**, *303*, L64–L74. <https://doi.org/10.1152/ajplung.00002.2012>.
577. Daiber, A.; Oelze, M.; Steven, S.; Kroller-Schon, S.; Munzel, T. Taking up the cudgels for the traditional reactive oxygen and nitrogen species detection assays and their use in the cardiovascular system. *Redox Biol.* **2017**, *12*, 35–49. <https://doi.org/10.1016/j.redox.2017.02.001>.
578. Hyman, A.L.; Lipton, H.L.; Kadowitz, P.J. Methylene blue prevents hypoxic pulmonary vasoconstriction in cats. *Am. J. Physiol.* **1991**, *260*, H586–H592. <https://doi.org/10.1152/ajpheart.1991.260.2.H586>.
579. Cremona, G.; Wood, A.M.; Hall, L.W.; Bower, E.A.; Higenbottam, T. Effect of inhibitors of nitric oxide release and action on vascular tone in isolated lungs of pig, sheep, dog and man. *J. Physiol.* **1994**, *481*, 185–195. <https://doi.org/10.1113/jphysiol.1994.sp020429>.

580. Dumas, J.P.; Dumas, M.; Sgro, C.; Advenier, C.; Giudicelli, J.F. Effects of two K⁺ channel openers, aprikalim and pinacidil, on hypoxic pulmonary vasoconstriction. *Eur. J. Pharmacol.* **1994**, *263*, 17–23. [https://doi.org/10.1016/0014-2999\(94\)90518-5](https://doi.org/10.1016/0014-2999(94)90518-5).
581. Mazmanian, G.M.; Baudet, B.; Brink, C.; Cerrina, J.; Kirkiacharian, S.; Weiss, M. Methylene blue potentiates vascular reactivity in isolated rat lungs. *J. Appl. Physiol.* **1989**, *66*, 1040–1045. <https://doi.org/10.1152/jappl.1989.66.3.1040>.
582. Fouty, B.; Komalavilas, P.; Muramatsu, M.; Cohen, A.; McMurtry, I.F.; Lincoln, T.M.; Rodman, D.M. Protein kinase G is not essential to NO-cGMP modulation of basal tone in rat pulmonary circulation. *Am. J. Physiol.* **1998**, *274*, H672–H678. <https://doi.org/10.1152/ajpheart.1998.274.2.H672>.
583. Rodman, D.M.; Yamaguchi, T.; O'Brien, R.F.; McMurtry, I.F. Methylene blue enhances hypoxic contraction in isolated rat pulmonary arteries. *Chest* **1988**, *93*, 93S–94S. https://doi.org/10.1378/chest.93.3_supplement.93s.
584. Vermeersch, P.; Buys, E.; Pokreisz, P.; Marsboom, G.; Ichinose, F.; Sips, P.; Pellens, M.; Gillijns, H.; Swinnen, M.; Graveline, A.; et al. Soluble guanylate cyclase- α 1 deficiency selectively inhibits the pulmonary vasodilator response to nitric oxide and increases the pulmonary vascular remodeling response to chronic hypoxia. *Circulation* **2007**, *116*, 936–943. <https://doi.org/10.1161/CIRCULATIONAHA.106.677245>.
585. Spohr, F.; Busch, C.J.; Teschendorf, P.; Weimann, J. Selective inhibition of guanylate cyclase prevents impairment of hypoxic pulmonary vasoconstriction in endotoxemic mice. *J. Physiol. Pharmacol.* **2009**, *60*, 107–112.
586. Rodman, D.M.; Yamaguchi, T.; Hasunuma, K.; O'Brien, R.F.; McMurtry, I.F. Effects of hypoxia on endothelium-dependent relaxation of rat pulmonary artery. *Am. J. Physiol.* **1990**, *258*, L207–L214. <https://doi.org/10.1152/ajplung.1990.258.4.L207>.
587. Ye, L.; Liu, J.; Liu, H.; Ying, L.; Dou, D.; Chen, Z.; Xu, X.; Raj, J.U.; Gao, Y. Sulfhydryl-dependent dimerization of soluble guanylyl cyclase modulates the relaxation of porcine pulmonary arteries to nitric oxide. *Pflugers Arch.* **2013**, *465*, 333–341. <https://doi.org/10.1007/s00424-012-1176-x>.
588. Harteneck, C.; Koesling, D.; Soling, A.; Schultz, G.; Bohme, E. Expression of soluble guanylyl cyclase. Catalytic activity requires two enzyme subunits. *FEBS Lett.* **1990**, *272*, 221–223. [https://doi.org/10.1016/0014-5793\(90\)80489-6](https://doi.org/10.1016/0014-5793(90)80489-6).
589. Le Cras, T.D.; McMurtry, I.F. Nitric oxide production in the hypoxic lung. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **2001**, *280*, L575–L582. <https://doi.org/10.1152/ajplung.2001.280.4.L575>.
590. Manoury, B.; Idres, S.; Leblais, V.; Fischmeister, R. Ion channels as effectors of cyclic nucleotide pathways: Functional relevance for arterial tone regulation. *Pharmacol. Ther.* **2020**, *209*, 107499. <https://doi.org/10.1016/j.pharmthera.2020.107499>.
591. Rudyk, O.; Prysyazhna, O.; Burgoyne, J.R.; Eaton, P. Nitroglycerin fails to lower blood pressure in redox-dead Cys42Ser PKG1 α knock-in mouse. *Circulation* **2012**, *126*, 287–295. <https://doi.org/10.1161/CIRCULATIONAHA.112.101287>.
592. Prysyazhna, O.; Rudyk, O.; Eaton, P. Single atom substitution in mouse protein kinase G eliminates oxidant sensing to cause hypertension. *Nat. Med.* **2012**, *18*, 286–290. <https://doi.org/10.1038/nm.2603>.
593. Rudyk, O.; Rowan, A.; Prysyazhna, O.; Krasemann, S.; Hartmann, K.; Zhang, M.; Shah, A.M.; Ruppert, C.; Weiss, A.; Schermuly, R.T.; et al. Oxidation of PKG1 α mediates an endogenous adaptation to pulmonary hypertension. *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 13016–13025. <https://doi.org/10.1073/pnas.1904064116>.
594. Stincone, A.; Prigione, A.; Cramer, T.; Wamelink, M.M.; Campbell, K.; Cheung, E.; Olin-Sandoval, V.; Gruning, N.M.; Kruger, A.; Tauqeer Alam, M.; et al. The return of metabolism: Biochemistry and physiology of the pentose phosphate pathway. *Biol. Rev. Camb. Philos. Soc.* **2015**, *90*, 927–963. <https://doi.org/10.1111/brv.12140>.
595. Mullarky, E.; Cantley, L.C. Diverting Glycolysis to Combat Oxidative Stress. In *Innovative Medicine: Basic Research and Development*; Nakao, K., Minato, N., Uemoto, S., Eds.; Springer: Tokyo, Japan, 2015; pp. 3–23.
596. Gupte, R.; Dhagia, V.; Rocic, P.; Ochi, R.; Gupte, S.A. Glucose-6-phosphate dehydrogenase increases Ca⁽²⁺⁾ currents by interacting with Ca(v)1.2 and reducing intrinsic inactivation of the L-type calcium channel. *Am. J. Physiol. Heart Circ. Physiol.* **2020**, *319*, H144–H158. <https://doi.org/10.1152/ajpheart.00727.2019>.
597. Chang, H.H.; Bennett, A.M.; Cameron, W.D.; Floro, E.; Au, A.; McFaul, C.M.; Yip, C.M.; Rocheleau, J.V. Targeting Apolipoprotein NADP(+) to Image NADPH Generation in Pancreatic Beta-Cell Organelles. *ACS Sens.* **2022**, *7*, 3308–3317. <https://doi.org/10.1021/acssensors.2c01174>.
598. Eid, M.; Barayeu, U.; Dick, T.P. Chemogenetic detection and quantitation of H₂O₂ in living cells. *Nat. Protoc.* **2025**, 1–24. <https://doi.org/10.1038/s41596-025-01226-9>.
599. Han, H.H.; Ge, P.X.; Li, W.J.; Hu, X.L.; He, X.P. Recent Advancement in Fluorescent Probes for Peroxynitrite (ONOO⁽⁻⁾). *Sensors* **2025**, *25*, 3018. <https://doi.org/10.3390/s25103018>.

600. Hung, Y.P.; Albeck, J.G.; Tantama, M.; Yellen, G. Imaging cytosolic NADH-NAD(+) redox state with a genetically encoded fluorescent biosensor. *Cell Metab.* **2011**, *14*, 545–554. <https://doi.org/10.1016/j.cmet.2011.08.012>.
601. Zanetti, C.; Gaspar, R.D.L.; Zhdanov, A.V.; Maguire, N.M.; Joyce, S.A.; Collins, S.G.; Maguire, A.R.; Papkovsky, D.B. Heterosubstituted Derivatives of PtPFPP for O₂ Sensing and Cell Analysis: Structure-Activity Relationships. *Bioconjug. Chem.* **2022**, *33*, 2161–2169. <https://doi.org/10.1021/acs.bioconjchem.2c00400>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.