



Review

Chemical Constituents and Their Biological Activities from Genus *Styrax*

Ding-Ding Xia ^{1,2,†}, Xin-Yu Han ^{1,3,†}, Yu Zhang ^{1,*} and Na Zhang ^{4,*}

¹ Shanghai Frontiers Science Center for Chinese Medicine Chemical Biology, Institute of Interdisciplinary Integrative Medicine Research, Shanghai University of Traditional Chinese Medicine, No. 1200, Cailun Road, Shanghai 201203, China; 2103084016@stu.htu.edu.cn (D.-D.X.); 1710201016@vip.henu.edu.cn (X.-Y.H.)

² School of Chemistry and Chemical Engineering, Henan Normal University, 46 East of Construction Road, Xinxiang 453007, China

³ Institute of Pharmacy, Pharmacy College of Henan University, Jinming District, Kaifeng 475004, China

⁴ Department of Biology, Philipps University, Karl-von-Frisch-Straße 8, 35043 Marburg, Germany

* Correspondence: yzhang@shutcm.edu.cn (Y.Z.); na.zhang@biologie.uni-marburg.de (N.Z.)

† D.X. and X.H. contributed equally.

Abstract: Plants from the genus *Styrax* have been extensively used in folk medicines to treat diseases such as skin diseases and peptic ulcers and as an antiseptic and analgesic. Most *Styrax* species, especially *Styrax tonkinensis*, which is used as an expectorant, antiseptic, and analgesic in Chinese traditional medicine, could screen resin after external injury. *Styrax* is also used in folk medicines in Korea to treat sore throat, bronchitis, cough, expectoration, paralysis, laryngitis, and inflammation. Different parts of various *Styrax* species can be widely employed for ethnopharmacological applications. Moreover, for ethnopharmacological use, these parts of *Styrax* species can be applied in combination with other folk medicines. *Styrax* species consist of versatile natural compounds, with some of them exhibiting particularly excellent pharmacological activities, such as cytotoxic, acetylcholinesterase inhibitory, antioxidant, and antifungal activities. Altogether, these exciting results indicate that a comprehensive review of plants belonging to this genus is essential for helping researchers to continuously conduct an in-depth investigation. In this review, the traditional uses, phytochemistry, corresponding pharmacological activities, and structure–activity relationships of different *Styrax* species are clarified and critically discussed. More insights into potential opportunities for future research are carefully assessed.

Keywords: *Styrax* Genus; biological activities; constituents; lignans; terpenoid

Citation: Xia, D.-D.; Han, X.-Y.; Zhang, Y.; Zhang, N. Chemical Constituents and Their Biological Activities from Genus *Styrax*. *Pharmaceuticals* **2023**, *16*, 1043. <https://doi.org/10.3390/ph16071043>
Academic Editor: Shengbao Cai and Jiangbo He

Received: 13 June 2023

Revised: 14 July 2023

Accepted: 19 July 2023

Published: 22 July 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The genus *Styrax* has a widespread but dispersive distribution. It is found in East Asian, American, and Mediterranean regions. It is the largest genus of the Styracaceae family and contains approximately 130 species [1]. *Styrax* stands out from other genera in this family because it produces a resinous material known as benzoin resin. This resin is typically released when the bark is injured by sharp objects. It has been utilized in various regions across the globe for its aromatic properties, being commonly used in perfumes and cosmetics, and *Styrax* species have traditionally been used in herbal medicines for the treatment of various diseases [2]. Of note, many *Styrax* species, especially *S. tonkinensis*, which is used as an expectorant, antiseptic, and analgesic in Chinese traditional medicine, could screen resin after exterior injury [3,4]. As a folk medicine in Korea, *S. japonica* is used to treat cough, bronchitis, sore throat, inflammation, paralysis, laryngitis, and expectoration [5–7]. The resin from *Styrax*, mixed with other antibiotic substances and hardening material, is also indicated in Islamic medicine as working as a good dental restorative material [2]. The flower of *S. japonicus* sieb. et Zucc. is used in Chinese folk medicine to

relieve pain such as sore throat pain and toothache [8]. The leaves and roots of *Styrax suberifolium* are typically used as traditional medicines in China to cure rheumatic diseases [9].

Furthermore, the extensive investigation of pharmacologically active compounds derived from various *Styrax* species has been ongoing for several decades. While numerous *Styrax* species have been studied, *S. obassia* and *S. japonica* have emerged as the most extensively researched species, encompassing studies ranging from phytochemistry to comprehensive pharmacological investigations (**Figure 1**). An example of the pharmacological potential of *Styrax* species was the inhibitory effect of benzofurans extracted from *Styrax agrestis* A. Chev. on acetylcholinesterase (AChE) in vitro [10]. Triterpenoids isolated from the resin of *S. tonkinensis* (Pier.) Craib showed promising antiproliferative and differentiation effects on human leukemia HL-60 cells [4]. Additionally, the hydroalcoholic extracts of *S. camporum* Pohl demonstrated effectiveness in reducing chromosome and DNA damage [11]. Another notable finding was the promotion of estrogen biosynthesis by egonol gentiobioside and egonol gentiotrioside from *Styrax perkinsiae* through the action of aromatase [12].

Despite significant progress in discovering natural compounds from *Styrax* species and elucidating their potential pharmacological activities, there is still a need for a comprehensive and focused discussion of this rapidly growing research area. With our continuous interests in natural products discovery and pharmacological research [13–16], our aim is to provide researchers with a convenient and comprehensive resource that offers detailed and concise profiles of the *Styrax* genus. This review encompasses the examination of structural diversity and the pharmacological and biological significance and presents the exciting future research prospects in this field.




Figure 1. Research overview of the *Styrax* genus.

2. Results and Discussion

2.1. Chemical Constituents

2.1.1. Lignans

Lignans are the major constituents isolated from *Styrax* species. Most lignans are benzofuran, tetrahydrofuran, and furofuran lignans, and they are found in the stem nucleus of *S. perkinsiae*, *S. ferrugineus*, *S. macranthus*, *S. obassia*, *S. camporum*, *S. japonica*, and *S. officinalis* L. [2]. *S. perkinsiae* contains 16 norlignans including **1–14** and lignans **15** and **16** [17,18]. Norlignans (**2**, **4**, and **17–19**) from *S. ferrugineus* leaves were investigated and characterized [19]. Compounds **5** and **20–35** were isolated from the stem bark of *S. japonica* by several research groups [7,20–24]. Meanwhile, lignans **4**, **15**, and **16** were also afforded from *S. japonica* seeds [25]. Constituents of *S. obassia* were investigated, and norlignans (**2**, **3**, **6**, and **36**) were isolated [26]. Moreover, a series of reports revealed the presence of several benzofurans in *S. obassia* including **37–42** [27–31]. Six benzofuran derivatives comprising **4–7**, **14**, and **43–46** were afforded from the seeds of *S. macranthus* that grow in southwestern China [32,33]. Benzofurans **4**, **15**, **40**, and **47–49** were isolated from the hexane extract of the seeds of *S. officinalis* L. [34–37]. Thirteen compounds, **4**, **15**, **37**, **39–42**, and **50–55**, were obtained from the ethyl acetate (EtOAc) extract of the fruits of *S. agrestis* [10]. Moreover, compounds **4**, **17**, and **56** were isolated from *S. camporum*, and their protective activities were continuously assessed in vivo [11,38]. Bertanha et al. isolated benzofuran nor-neolignan derivatives **4**, **6**, **17**, **18**, and **57** from the aerial parts of *S. pohlii*. Several lignans including **58–65** were isolated from *S. perkinsiae* [39]. Seventeen phenylpropanoids were successfully isolated from the bark of *S. suberifolius*, including ten benzofuran derivatives (**4–5**, and **66–73**), two dihydrofuran derivatives (**23** and **65**), two new neolignans (**74** and **75**), and three benzalcohols (**76–78**) [40]. Eight lignans (**79–85**) were isolated from the leaves of *S. tonkinensis* (Pierre) Craib ex Hartw [41]. Two new phenylpropanoids (**86** and **87**) were isolated from the resin of *S. tonkinensis* (Pierre) Craib ex Hartw by Fang's groups [42]. Two lignans (**88** and **89**) and five nor-lignan-type benzofurans, including **4** and **90–93**, were separated from *S. argentifolius* by Son's group [43].

2.1.2. Terpenoids

Terpenoids were also obtained from the *Styrax* genus as one of its major constituents. It should be noted that a vast majority of the terpenoids isolated from the *Styrax* genus were pentacyclic triterpenoids. To date, these molecules were only found in four species of the *Styrax* genus. Compounds **94–102** were isolated from the stem bark of *S. japonica* Sieb. et Zucc. by several research groups [20,22,44,45]. A phytochemical investigation on the fresh fruits of *S. japonica* Sieb. et Zucc. was also conducted, and four new triterpenoid glycosides including jegosaponins A–D (**103–106**) were found [46]. Furthermore, *S. japonica* Sieb. et Zucc. continued to be investigated by Kwon's group, and **107–110** and taraxerol (**94**) were isolated [47]. In addition to the plants themselves, triterpenoids were also found from the resin of *S. tonkinensis* (Pier.) Craib containing **111–119** [4,6]. A pentacyclic triterpenoid (**120**), three triterpenoid saponins *styrax*-saponins A–C (**121–123**), and deacylsaponin (**124**) were also obtained from *S. officinalis* L. [48,49]. Moreover, several monoterpenes, such as α -terpineol, linalool, and geraniol, were isolated from the benzoe resin of *S. officinalis* L. [50]. Recently, two cinnamyl esters and seven pentacyclic triterpene acids (**119**, **125–130**) were separated and characterized from *S. tonkinensis* (Pierre) Craib ex Hartw [42,51]. A triterpenoid (**131**) was obtained from *S. argentifolius* very recently [43].

2.1.3. Aromatic Compounds

Aromatic compounds, as a small proportion, were reported in the *Styrax* genus as well. In the species of *S. tonkinensis* (Pier.) Craib, seven aromatic compounds including **132–139** were reported [52,53]. Moreover, Kim and coworkers found **140** and **141** from the stem bark of *S. japonica* (SJ) [54]. In *S. perkinsiae* Rhed., **142** was separated [39]. Recently, a

new epicatechin glucopyranoside, **143**, and three mononuclear phenolic acid esters, **144–146**, were isolated from the bark of *S. suberifolius* Hook [40].

2.1.4. Steroids

Luo and coworkers reported that three steroids including stigmasterol (**147**), styraxosides A (**148**), and daucosterol (**149**) were obtained from the seeds of *S. macranthus* Perk [32]. Another Steroid named β -sitosterol (**150**) was reported in *S. perkinsiae* Rehder [17]. A sterol, **151**, was separated from *S. argentifolius* H.L. Li by Son's group [43].

2.1.5. Others

In addition to the commonly isolated products from the genus *Styrax*, other types of natural products were also reported with relatively limited numbers. For example, in 1973, a preliminary result regarding the seeds of *S. officinalis* L. showed that the oil content amounts to 50% [55]. Moreover, flavonoids are not frequently reported in the *Styrax* genus according to literature studies. Only four flavonoids including **152–155** were isolated from the aerial parts of *S. pohlii* A. DC. and the leaves of *S. camporum* Pohl [56]. Later, two new polyketones, **156–157**, were isolated from *Styrax camporum* Pohl. [57]. Recently, two bioactive saponins, Jegosaponin A and B (**158–159**), were extracted and subsequently identified from *S. japonica* Siebold et al. Zuccarini [58].

2.2. Chemical Constituents Biological Activities

2.2.1. Cytotoxic Activity

S. perkinsiae was investigated, and the cytotoxic activity of the compounds isolated from this species was tested through the colorimetric chemosensitivity assay with SRB. (Figure 2). Interestingly, **11** and **14** revealed cytotoxic activities in vitro against two breast cancer cell lines, MCF-7 (IC_{50} = 5.5 and 15.0 μ g/mL, respectively) and MDA-MB-231 (IC_{50} = 3.81 and 13.71 μ g/mL, respectively) [17].

Later, the cytotoxic activities of lignans isolated from *S. camporum* against three cell lines, namely, HeLa (human cervix carcinoma), C6 (rat glioma), and Hep-2 (larynx epidermoid carcinoma), were analyzed using the standard MTT. Compound **4** showed strong cytotoxic activities against the Hep-2 (IC_{50} = 3.6 μ g/mL) and C6 (IC_{50} = 3.2 μ g/mL) cell lines. Compound **17** exhibited significant cytotoxic activities against the HeLa (IC_{50} = 5.3 μ g/mL) and C6 (IC_{50} = 4.9 μ g/mL) cell lines. Compound **56** exhibited moderate cytotoxic activities against the Hep-2 (IC_{50} = 28.0 μ g/mL), HeLa (IC_{50} = 31.7 μ g/mL), and C6 (IC_{50} = 10.7 μ g/mL) cell lines. Moreover, when combined, **4** and **17** exhibited higher cytotoxic activities than the hydroalcoholic extract or either of the lignans alone, with the lowest IC_{50} being 13.3 μ g/mL [38,59].

Seven compounds isolated from *S. obassia* were screened for their cytotoxic activities against the HeLa, HL-60, and MCF-7 cell lines. Among them, compounds **3** and **5** exhibited significant antitumor properties. Compound **3** exhibited cytotoxicity against the HeLa (IC_{50} = 23.3 μ g/mL), HL-60 (IC_{50} = 16.8 μ g/mL), and MCF-7 cells (IC_{50} = 53.5 μ g/mL). Meanwhile, compound **5** exhibited cytotoxicity against HeLa (IC_{50} = 23.3 μ g/mL), HL-60 (IC_{50} = 47.8 μ g/mL), and MCF-7 cells (IC_{50} = 27.9 μ g/mL) [60].




Figure 2. Natural compounds with cytotoxic activity.

Through the Cell Counting Kit-8 (CCK-8) test in vitro, compounds **86** and **87** were tested for their cytotoxic activities against five tumor cell lines (PC-3, MCF-7, A549, HeLa, and HepG-2). Among them, the cytotoxic effect of compound **86** was observed against the MCF-7 and HeLa cell lines (IC_{50} = 26.75 and 45.16 μ M, respectively), which was better or similar to that of the positive control cisplatin (IC_{50} = 40.95 and 47.36 μ M, respectively). Compound **86** exhibited moderate cytotoxicity against the PC-3 and HepG-2 cell lines. The other biomolecule, **87**, displayed moderate cytotoxicity against MCF-7 cells (IC_{50} = 57.1 μ M) [42].

Son's group assessed the cytotoxicity and α -glucosidase inhibitory activity of isolated compounds from *S. argentifolius*. They suggested that the activities of triterpenoid **131** and norlignan-type benzofurans (**4** and **91–93**) are superior to those of others including sterol **153** and lignans **88** and **89**. The better activities of benzofurans (**4** and **91–93**) were postulated to be an effect of the substitutions at the side chain of carbon C-5. Among them, compound **4** exhibited potential cytotoxicity against three cancer cell lines, namely, Lu (IC_{50} = 21.50 μ g/mL), KB (IC_{50} = 22.11 μ g/mL), and HepG-2 (IC_{50} = 18.15 μ g/mL) [43].

2.2.2. Antibacterial and Antifungal Activity

Initially, the extract of *S. ferrugineus* exhibited antifungal and antibacterial activities against *Candida albicans*, *Cladosporium sphaerospermum*, and *Staphylococcus aureus*. To identify the potential biomolecules from this species that exhibit antifungal and antibacterial activities, the isolated lignans were tested (**Figure 3**). Among them, lignans **4** and **17** exhibited antifungal and antibacterial activities against *S. aureus* (MIC = 10 μ g/mL and 20 μ g/mL, respectively), *C. albicans* (MIC = 10 μ g/mL and 12 μ g/mL, respectively), and *C. sphaerospermum* (MIC = 5 μ g/mL and 10 μ g/mL, respectively), whereas the other three

natural products (**5**, **18**, and **19**) only inhibited *C. albicans* (MIC = 15 µg/mL, 20 µg/mL, and 15 µg/mL, respectively) and *S. aureus* (MIC = 20 µg/mL, 20 µg/mL, and 20 µg/mL, respectively) [19].

To exploit the antibacterial activity of the aerial parts of *S. pohlii*, different fractions, especially those extracted using *n*-hexane, EtOAc, *n*-BuOH, and methanol, were evaluated against *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *S. pyogenes*, *Streptococcus pneumoniae*, and *Klebsiella pneumoniae*. The broth microdilution method was used for measuring the minimum inhibitory concentration (MIC). Among the fractions, the *n*-hexane fraction exhibited excellent antibacterial activity against Gram-positive *S. pneumoniae* (MIC = 200 µg/mL). The MIC values of compounds **4** and **17** (400.0 µg/mL) against *P. aeruginosa* and *S. pneumoniae* were the best [61].




Figure 3. Natural compounds with antibacterial and antifungal activities.

By conducting the radial growth-inhibition experiment, the antifungal activities of compounds from the bark of *S. suberifolius* against three plants' fungal pathogen, namely, *Phomopsis cytospora*, *Fusarium oxysporum*, and *Alternaria Solani*, was exhibited. Compounds **144**, **145**, and **146** exhibited selective suppressive activities against the tested fungi. Notably, compound **146** was a significantly effective inhibitor of *Phomopsis cytospora* at 100.0 µg/mL, with an inhibition rate of 86.72% [40].

2.2.3. Antiproliferative and Differentiation Effects

In 2006, Wang's group found that triterpenoids (**111–120**) isolated from *S. tonkinensis* inhibit HL-60 cell growth ($IC_{50} = 8.9–99.4 \mu M$). Of note, oleanolic acid **119** acted as the most effective antiproliferative agent ($IC_{50} = 8.9 \mu g/mL$) (**Figure 4**). Compound **113** exhibited the lowest growth-inhibitory effect. According to the NBT-reduction assay, compound **113** induced HL-60 cell differentiation, as measured in [4].




Figure 4. Natural compounds with antiproliferative and differentiation properties.

2.2.4. Anti-Complement Activity

Egonol (**4**), masutakeside I (**10**), styraxlignolide A (**28**), and styraxoside B (**101**) isolated from *S. japonica* could inhibit the hemolytic activity of the complement system ($IC_{50} = 33, 166, 123$, and $65 \mu M$, respectively) (**Figure 5**). This finding strongly suggested that the methyl enedioxy group of lignans has a vital role in inhibiting the hemolytic activity of human serum against erythrocytes [22].




Figure 5. Natural compounds with anti-complement activity.

2.2.5. Anti-Complement Activity

Natural products isolated from *S. japonica* were tested for in vitro antioxidant activities through the DPPH radical scavenging test. Among them, **30–33** exhibited weak DPPH radical scavenging activities ($IC_{50} = 380, 278, 194$, and $260 \mu M$, respectively) (**Figure 6**) [7]. Moreover, Oliveira et al. reported that the hydroalcoholic extract of *S. camporum* could concentration-dependently scavenge DPPH radicals; a maximum scavenging activity of 85% was observed at $30.0 \mu g/mL$ [11].




Figure 6. Natural compounds with antioxidant activity.

2.2.6. Induction of Apoptosis

Lee and Lim revealed that the ethanol extract of *S. japonica* Siebold et al., Zuccarini (SJSZ) induced programmed cell death (apoptosis) in HepG2 cells under the experimental condition (75.0 µg/mL of SJSZ for 4 h treatment). The results indicated that the ethanol extract of SJSZ (75 µg/mL) stimulates an increase in the number of iROS, Ca²⁺, and the apoptotic-related factors in HepG2 cells [62].

2.2.7. Induction of Apoptosis

In 2002, a nonradioactive assay was established for measuring aromatase activity by using human ovarian granulosa KGN cells. Lignans **6** and **7** exhibited approximately 1.62- and 1.95-fold increases, respectively, in 17 β-estradiol biosynthesis at 10 µM, and significantly improved 17 β-estradiol biosynthesis by approximately 1.53- and 1.71-fold, respectively, in 3T3-L1 preadipocyte cells (**Figure 7**). Moreover, egonol gentiotrioside increased serum estrogen levels in ovariectomized rats. These results suggested that these two lignans induce estrogen biosynthesis through the allosteric regulation of aromatase activity [12].




Figure 7. Natural compounds exhibiting estrogen-promoting activity.

2.2.8. Acetylcholinesterase Inhibitors and Structure–Activity Relationships

In 2011, Liu et al. screened their library of plant extracts through a high-throughput assay. They found that the EtOAc extract of *S. agrestis* fruits exhibited significant inhibitory activity against AChE. They proved that two active subfractions were responsible for this inhibition and further isolated 13 compounds from the EtOAc extract. Later, they examined the selectivity and inhibitory potency of benzofurans on hAChE, BChE, and EeAChE by using the improved Ellman's colorimetric method (**Figure 8**). Some egonol derivatives were synthesized through chemical modifications to clearly understand the structure–activity relationships. According to the results, the inhibition ratio affects the bulkiness and length of the alkyl ester group. In particular, compounds **50–53** exhibited inhibitory activity against AChE (IC₅₀ = 1.4–3.1 µM). Compound **50** at 100.0 µM displayed obvious inhibition of Aβ aggregation (77.6%). Liu et al.'s SAR (Structure-Activity Relationships) studies indicated that compounds exhibiting anti-AChE activity are observed with the incorporation of alkyl chains consisting of more than three carbon units, the furan ring, and the ester group. Molecular docking studies proposed a binding site for this class of compound on AChE and identified multiple key residues at the peripheral site that are crucial for mediating the inhibitory effect [10]. The anti-AChE and antifungal activities of two novel polyketides, **156** and **157**, were also tested through TLC bioautographic assays. The results indicated that compound **156** could inhibit AChE activity [57].




Figure 8. Natural compounds as potential acetylcholinesterase inhibitors.

2.2.9. Inhibitory Effect on Interleukin

Lee and Lim separated a glycoprotein with an approximate molecular mass of 38 kDa from *S. japonica*. Subsequently, an immunoblot analysis and RT-PCR were conducted to evaluate ERK, JNK, and NF- κ B activities and the levels of inflammation-related factors (COX-2, inducible nitric oxide synthase (iNOS), and interleukin (IL)-1 β) in Cr-induced BNL CL2 cells. The SJSZ glycoprotein (50.0 $\mu\text{g/mL}$) inhibited the expression of ERK, NF- κ B, JNK, iNOS, IL-1 β , and COX-2 [63].

With further investigation of the SJSZ glycoprotein (38 kDa), Lee and Kim proved that this glycoprotein modulates IFN- γ , IL-2, and IL-12 expression in cyclophosphamide (CTX)-induced Balb/c mice. The glycoprotein counteracted the CTX-induced immunosuppressive effects. It effectively restored the spleen and thymus weights to normal levels and enhanced the phagocytic activity of peritoneal macrophages in response to CTX. Furthermore, the SJSZ glycoprotein exerted regulatory effects on the proliferation of T and B lymphocytes, cytotoxicity of NK cells, and production of key cytokines (IFN- γ , IL-2, and IL-12). Additionally, it improved the activity of antioxidant enzymes (e.g., SOD, CAT, and GPx) [64].

2.2.10. Matrix Metalloproteinase's Activity

Some triterpenoids (**95–97**, **100**) were isolated and further tested the Matrix Metalloproteinases (MMPs)' activity of the methylene chloride soluble fraction of a methanol extract from the stems of *S. japonica*. Among them, **95** and **100** displayed effective cytotoxic activities against human dermal fibroblasts (IC₅₀ = 20.0 and 1.12 μM , respectively) (**Figure 9**). In addition, **96** and **97** exhibited no cytotoxicity for the same cells at the test dose (0.01–1 μM). However, **96** dose-dependently reduced UV-induced MMP-1 protein levels to normal levels by 73.1% at 0.01 μM [65]. In a dose-dependent manner, **96** effectively downregulated MMP-1 protein expression, whereas it upregulated type-1 procollagen protein expression in the UV-irradiated cultured human skin fibroblasts of an elderly person [66].




Figure 9. Natural compounds with matrix metalloproteinase's activity.

Styrax japonoside B (**26**) exerted inhibitory activity against MMP-1 and prevented UV-induced changes in MMP-1 expression. At 10 μ M, the treatment led to a significant dose-dependent reduction in MMP-1 protein expression, with an average decrease of 62.1% compared with the vehicle-treated control cells. The findings suggested that the glycoprotein can potentially be used as a potent antimetastatic agent. This glycoprotein exerts its effect by suppressing MMP-9 enzymatic activity through the NF- κ B and AP-1 signaling pathways [67].

Two cinnamyl esters (**86** and **87**) and seven pentacyclic triterpene acids (**119** and **125–130**) in *Styrax* are the key components that inhibit hCES1A activity. These seven pentacyclic triterpene acids in the two active sites of *Styrax* exert a significant inhibitory effect on hCES1A (IC_{50} = 41–478 nM). Among them, epibetulinic acid (**129**) (IC_{50} = 0.041 μ g/mL), oleanonic acid (**125**) (IC_{50} = 0.49 μ g/mL), and betulonic acid (**126**) (IC_{50} = 1.48 μ g/mL) exhibited the strongest inhibitory activity against hCES1A [42,51].

2.2.11. Antiasthmatic, Antiulcer, and Anti-Inflammatory Activities

In a murine asthma model, homoegonol (**17**) exerted significant effects in reducing inflammatory cell infiltration and Th2 cytokine production in the bronchoalveolar lavage fluid. It also attenuated airway hyperresponsiveness, decreased serum IgE levels, and downregulated iNOS and MMP-9 expression. Thus, compound **17** exhibited the potential to effectively suppress OVA challenge-induced asthmatic responses (Figure 10).




Figure 10. Natural compounds with antiasthmatic, antiulcer, and anti-inflammatory activities. The EtOAc fraction of *S. camporum* displayed antiulcer activity. It reduced the ulcer area and gastric secretion volume and increased the number of collagen fibers [37,68].

In 2005, the extracted fractions of *S. pohlii* aerial parts, including the EtOAc fraction, ethanolic extract, and hexane fraction, were evaluated for their inhibitory activities against COX-1 and COX-2. The isolated products were further assessed against COX-1 and COX-2. The results revealed that all crude fractions and isolated products induced weak-to-

moderate COX-1 and COX-2 inhibition. Among them, **57** exerted mild COX-1 inhibition, of 35.7% at 30 μ M [69].

2.2.12. Other Activity

Through micronucleus and comet assays, Oliveira demonstrated that different doses (250, 500, and 1000 mg/kg body weight) of the *S. camporum* extract's compounds **4** and **17** had no genotoxic effect in Swiss mice. Moreover, they were effective in reducing doxorubicin- and methanesulfonate-induced DNA and chromosomal damage [11].

Braguine [69] investigated the EtOAc fractions of *S. camporum* and *S. pohlii* and isolated and identified compounds **152–155**. Upon biological evaluation, they found that the EtOAc fractions, as well as compounds **152** and **155**, could separate coupled *Schistosoma mansoni* adult worms. Additionally, compound **155** killed adult schistosomes in vitro. This research group also observed that homoeogonol and homoeogonol glucoside exhibited the best results against *S. mansoni* adult worms [70].

In vitro assessments were conducted to determine the protein tyrosine phosphatase 1B (PTP1B)'s inhibitory activities of compounds from *S. japonica* stem bark. Among the isolated compounds, **108** and **109** had the highest inhibitory activities (IC_{50} = 7.8 and 9.3 μ M, respectively) [45].

By downregulating NF- κ B–DNA binding activity, styraxoside A (**148**) derived from *S. japonica* exerted inhibitory effects on the expression of LPS-induced iNOS, COX-2, tumor necrosis factor- α , and IL-1 β [45].

Jegosaponins A and B (**158** and **159**, respectively) exhibited potent hemolytic activity in sheep defibrillation (IC_{50} = 2.1 and 20.2 μ g/mL, respectively) and could improve the performance of PC-3 cells and zebrafish embryos through the identification of a membrane nonpermeable DRAQ7, which is a fluorescent nucleus staining dye [61] (**Figure 11**).




Figure 11. Natural compounds with other properties.

3. Materials and Methods

Through the search of a variety of online libraries such as Wiley Online Library, PubMed, Scifinder Web, ACS, and Web of Science, a summary of the newly discovered chemicals isolated from the genus *styrax* and their related biological activities in recent decades was provided. All species names were checked using <http://www.theplantlist.org> (accessed on 10 May 2023).

4. Conclusions

In summary, the *Styrax* genus comprises 130 species, and most of the species are extensively used as traditional medicines (**Appendix A**), particularly in China and Korea. *Styrax* can be easily collected because of its extensive distribution. All the species of the *Styrax* family, which were reported regarding the aspects of phytochemistry and pharmacology, were comprehensively summarized. In total, 159 compounds (**Appendix B**), including lignans, terpenoids, steroids, etc., were isolated from various species. The biological activities of those isolated compounds were subsequently investigated, exhibiting broad bioactivities such as cytotoxic activity, antioxidant activity, antifungal activity, apoptotic activity, anti-inflammation activity, anti-complement activity and so on. Chemical and pharmacological studies on the *Styrax* genus also proved that its main constituents are lignans and terpenoids. Moreover, several bioactive molecules exhibiting strong pharmacological activities were also isolated from *Styrax* (**Appendix C**).

Of note, information about the structure–activity relationships of most bioactive compounds is insufficient due to the lack of derivatives. Therefore, the exploitation of the versatility of the potentially bioactive natural compounds obtained from this genus is in great demand. Moreover, some species used in traditional medicines are still untapped such as *S. suberifolius*, which is used as a cure for rheumatic arthritis, whereas the modern physiochemical and pharmacological investigations are missing. Furthermore, in-depth pharmacological studies, especially in vivo studies, of the isolated biomolecules should be conducted in the future.

Author Contributions: Conceptualization, D.-D.X. and X.-Y.H.; methodology, Y.Z.; formal analysis, N.Z.; investigation, D.-D.X. and X.-Y.H.; resources, Y.Z.; data curation, D.-D.X.; writing—original draft preparation, D.-D.X. and X.-Y.H.; writing—review and editing, D.-D.X., X.-Y.H., and Y.Z.; visualization, N.Z.; supervision, N.Z.; project administration, Y.Z.; funding acquisition, Y.Z. All authors have read and agreed to the published version of the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Chenguang Program of the Shanghai Education Development Foundation and the Shanghai Municipal Education Commission (22CGA51 to Y.Z.). And the start-up funding from Shanghai University of Traditional Chinese Medicine to Y.Z.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We are extremely appreciative of Wei-Dong Zhang and Jin-Xin Wang from the Second Military Medical University for the helpful discussion.

Conflicts of Interest: The authors declare no conflicts of interest.

Appendix A

Table A2. Genus *Styrax* and their traditional use.

Genus <i>Styrax</i>	Traditional Use
<i>S. perkinsiae</i>	Estrogen synthesis promotion

<i>S. obassia</i>	Anticancer activities
<i>S. japonica</i>	Cough, bronchitis, sore throat, inflammation, paralysis, laryngitis, and expectoration treatment
<i>S. pohlii</i>	/
<i>S. camporum</i>	Antimicrobial, anticancer, antifungal, hypolipidemic, and immunosuppressive activities
<i>S. macranthus</i>	/
<i>S. officinalis</i>	Antiseptic and anti-respiratory disease
<i>S. argentifolius</i>	/
<i>S. ferrugineus</i>	/
<i>S. agrestis</i>	/
<i>S. tonkinensis</i>	Expectorant, antiseptic, and analgesic activities
<i>S. suberifolius</i>	Rheumatic disease cure
<i>S. ramirezii</i>	/

Appendix B

Table A1. Chemical constituents of plants from the genus *Styrax*.

No.	Compound Class and Name	Source	Ref.
Lignans			
1	5-(2-Propen-1-one)-7-me-thoxy-2-(3,4-methylenedioxyphenyl) benzofuran	<i>S. perkinsiae</i>	[18]
2	1''-Hydroxyegonol gentiobioside	<i>S. perkinsiae</i>	[18]
		<i>S. obassia</i>	[2]
		<i>S. ferrugineus</i>	[19]
3	Obassioside B	<i>S. perkinsiae</i>	[18]
		<i>S. obassia</i>	[2]
		<i>S. obassia</i>	[2]
		<i>S. obassia</i>	[61]
4	Egonol	<i>S. perkinsiae</i>	[18]
		<i>S. japonica</i>	[21]
			[25]
		<i>S. obassia</i>	[28]
			[31]
		<i>S. agrestis</i>	[10]
		<i>S. ferrugineus</i>	[19]
		<i>S. pohlii</i>	[71]
		<i>S. camporum</i>	[11]
		<i>S. macranthus</i>	[32]
5	Egonol glucoside	<i>S. officinalis</i> L.	[35]
		<i>S. argentifolius</i>	[43]
		<i>S. perkinsiae</i>	[18]
		<i>S. japonica</i>	[20]
		<i>S. suberifolius</i>	[55]
		<i>S. macranthus</i>	[32]
		<i>S. obassia</i>	[28]
6	Egonol gentiobioside		[61]
		<i>S. perkinsiae</i>	[17] [18]

		<i>S. macranthus</i>	[32]
		<i>S. obassia</i>	[2]
		<i>S. pohlii</i>	[71]
		<i>S. perkinsiae</i>	[17]
7	Egonol gentiotrioside	<i>S. perkinsiae</i>	[18]
		<i>S. macranthus</i>	[32]
		<i>S. perkinsiae</i>	[18]
8	Masutakeside I (sutakeside I)	<i>S. japonica</i>	[22]
		<i>S. obassia</i>	[27]
9	<i>trans</i> -5-(3-Hydroxypropyl)-7-methoxy-2-[3-methoxy-4-hydroxyphenyl]-benzofuran-5-yl]benzofuran	<i>S. perkinsiae</i>	[17]
10	(<i>E</i>)-5-(2-Formylvinyl)-7-methoxy-2-(3,4-methylenedioxyphenyl)benzofuran	<i>S. perkinsiae</i>	[17]
11	5-(3-Butanoyloxypropyl)-7-methoxy-2-(3,4-methylene dioxyphenyl)benzofuran	<i>S. perkinsiae</i>	[17]
12	5-(3-Hydroxypropyl)-7-hydroxy-2-(3,4-methylene- dioxyphenyl) benzofuran	<i>S. perkinsiae</i>	[17]
		<i>S. macranthus</i>	[32]
12	5-(3-Hydroxypropyl)-7-hydroxy-2-(3,4-methylene- dioxyphenyl) benzofuran	<i>S. perkinsiae</i>	[17]
13	Egonol acetate	<i>S. macranthus</i>	[32]
		<i>S. perkinsiae</i>	[17]
		<i>S. japonica</i>	[25]
		<i>S. obassia</i>	[28,29,31]
		<i>S. agrestis</i>	[10]
		<i>S. officinalis</i> L.	[37]
14	Demethoxy egonol acetate	<i>S. perkinsiae</i>	[17]
		<i>S. japonica</i>	[25]
15	Styraxlignolide B	<i>S. perkinsiae</i>	[18]
		<i>S. japonica</i>	[7]
16	Styraxjaponoside C	<i>S. perkinsiae</i>	[18]
		<i>S. japonica</i>	[7]
17	Nor-lignans 5-(3''-hydroxypropyl)-7-methoxy-2-(3',4'-methylenedioxyphenyl) benzofuran	<i>S. ferrugineus</i>	[19]
		<i>S. camporum</i>	[11]
		<i>S. pohlii</i>	[71]
18	5-[3''-(β -D-Glucopyranosyloxy)propyl]-7-methoxy-2-(3',4'-methylenedioxyphenyl) benzofuran	<i>S. ferrugineus</i>	[19]
		<i>S. pohlii</i>	[71]
19	Dihydrodehydrodiconiferyl alcohol	<i>S. ferrugineus</i>	[19]
20	Styraxjaponoside C	<i>S. japonica</i>	[21]
21	Arctiin	<i>S. japonica</i>	[21]
22	Matairesinoside	<i>S. japonica</i>	[20,21]
23	Pinoresinol-4-O- β -D-glucopyranoside	<i>S. japonica</i>	[21]
24	2 <i>R</i> -(4'-hydroxy-3'-methoxyphenyl)-6 <i>R</i> -(3'',4''-methylenedioxyphenyl)-8-oxo-3,7-dioxabicyclo [3.3.0]octane 4'-hydroxyl	<i>S. japonica</i>	[7]
25	Styraxjaponoside A	<i>S. japonica</i>	[20]
26	Styraxjaponoside B	<i>S. japonica</i>	[20]

27	Dihydrodehydrodiconiferyl alcohol 9-O-glucoside	<i>S. japonica</i>	[20]
28	Styraxlignolide A	<i>S. japonica</i>	[22]
29	Styraxlignolide B	<i>S. japonica</i>	[7]
30	Styraxlignolide C	<i>S. japonica</i>	[7]
31	Styraxlignolide D	<i>S. japonica</i>	[7]
32	Styrax lignolides F	<i>S. japonica</i>	[7,24]
33	(-)-Pinoresinol glucoside	<i>S. japonica</i>	[7]
34	Styrlignan A	<i>S. japonica</i>	[23]
35	1R,2R,5S,6R-2-(4'-Hydroxy-3'-methoxyphenyl)-6-(3'',4''-dimethoxyphenyl)-3,7-dioxabicyclo-[3.3.0]octane-4'-O- β -D-glucopyranoside	<i>S. japonica</i>	[24]
36	Obassioside A	<i>S. japonica</i>	[26]
37	Methyl 3-[7-methoxy-2-(3',4'-methylene-dioxyphenyl)-5-benzofuranyl]-propionate	<i>S. obassia</i>	[31]
38	Methyl 3-[2-(3',4'-methylenedioxyphenyl)-5-benzofuranyl]-propionate	<i>S. obassia</i>	[31]
39	5-(3''-Propanoyloxypropyl)-7-methoxy-2-(3',4'-methylenedioxyphenyl)-benzofuran	<i>S. obassia</i> <i>S. agrestis</i>	[28] [10]
40	Egonol-2-methylbutanoate	<i>S. obassia</i>	[31]
41	7-Demethoxyegonol-2-methylbutanoate	<i>S. obassia</i>	[5]
42	Egonol propanoate	<i>S. obassia</i> <i>S. agrestis</i>	[29] [10]
43	3-[7-Methoxy-2-(3,4-methylenedioxyphenyl)benzofuran-5-yl]propyl-3-[7-methoxy-2-(3,4-methylenedioxyphenyl)-benzofuran-5-yl]propanoate	<i>S. macranthus</i>	[32]
44	Demethoxy egonol gentiobioside	<i>S. macranthus</i>	[32]
45	7-Methoxy-2-(3,4-methylenedioxyphenyl)benzofuran-5-carbaldehyde	<i>S. macranthus</i>	[32]
46	Demethoxy egonol	<i>S. macranthus</i> <i>S. obassia</i> <i>S. japonica</i>	[32] [28] [23]
47	5-3''-(2-Methylbutanoyloxy)propyl]-7-methoxy-2-(3',4'-dimethoxyphenyl)benzofuran	<i>S. officinalis</i> L.	[23]
48	5-(3''Benzoyloxypropyl)-7-methoxy-2-(3',4'-methylenedioxyphenyl)-benzofuran	<i>S. officinalis</i> L.	[36]
49	4-[3''-(1c-methylbutanoyloxy)propyl]-2-methoxy-(3',4'-methylenedioxyphenyl)-1a,5b-dihydrobenzo-[3,4]-cyclobutaoxirene	<i>S. officinalis</i> L.	[67]
50	Egonol-9(Z),12(Z) linoleate	<i>S. agrestis</i>	[10]
51	7-Demethoxyegonol-9(Z),12(Z) linoleate	<i>S. agrestis</i>	[10]

52	7-Demethoxyegonol oleate	<i>S. agrestis</i>	[10]
53	Egonol oleate	<i>S. agrestis</i>	[10]
54	7-Demethoxyegonol acetate	<i>S. agrestis</i>	[10]
55	Egonol-2-methylpropanoate	<i>S. agrestis</i>	[10]
56	(±)Syringaresinol	<i>S. camporum</i>	[38]
57	Homoegonol gentiobioside	<i>S. pohlii</i>	[71]
58	Lariciresinol 4-O-β-D-glucoside	<i>S. perkinsiae</i>	[39]
59	(-)-Secoisolariciresinol 4-O-β-D-Glucopyranoside	<i>S. perkinsiae</i>	[39]
60	Lariciresinol 4'-O-β-D-glucoside	<i>S. perkinsiae</i>	[39]
61	Lanicepside A	<i>S. perkinsiae</i>	[39]
62	Solariciresinol 4-O-β-D-glucopyranoside	<i>S. perkinsiae</i>	[39]
63	(+)-Lariciresinol 9-O-β-D-glucopyranoside	<i>S. perkinsiae</i>	[39]
64	2R,3S-Dihydrodehydrodiconiferyl alcohol 4'-O-β-D-glucopyranoside	<i>S. perkinsiae</i>	[39]
65	Pinoresinol	<i>S. perkinsiae</i>	[39]
66	Homoegonol glucoside	<i>S. suberifolius</i>	[55]
67	2-(4-Hydroxy-3-methoxyphenyl)-5-(3-hydroxypropyl)-7-methoxybenzofuran	<i>S. suberifolius</i>	[55]
68	2-(3-Hydroxy-4-methoxyphenyl)-7-methoxy-5benzofuranpropanol	<i>S. suberifolius</i>	[55]
69	(+)-Cedrusin	<i>S. suberifolius</i>	[55]
70	(-)-(7R,8S)-Dihydrodehydrodiconiferyl alcohol	<i>S. suberifolius</i>	[55]
71	(-)-(7R,8S)-Dihydrodehydrodiconiferyl alcohol 4-O-β-D-glucopyranoside	<i>S. suberifolius</i>	[55]
72	(-)-(7S,8R)-Dihydrodehydrodiconiferyl alcohol 4-O-β-D-glucopyranoside	<i>S. suberifolius</i>	[55]
73	(+)-(7S,8R)-Dihydrodehydrodiconiferyl alcohol	<i>S. suberifolius</i>	[55]
74	(+)-(7S,8R)-Erythro-4,7,9,9'-tetrahydroxy-3,3'-dimethoxy-8-O-4'-neolignan	<i>S. suberifolius</i>	[55]
75	(-)-Symplocosneolignan A	<i>S. suberifolius</i>	[55]
76	(-)-7-O-Ethylguaiacylglycerol (15)	<i>S. suberifolius</i>	[55]
77	2[4-(3-Hydroxypropyl)-2-methoxyphenoxy]-1,3-propanediol	<i>S. suberifolius</i>	[55]
78	Dihydroconiferyl alcohol	<i>S. suberifolius</i>	[55]
79	3,3-Bis(3,4-dihydro-6-methoxy-2H-1-benzopyran	<i>S. tonkinensis</i>	[41]
80	Rac-(8α,8'β)-4,4'-dihydroxy-3,3'-dimethoxylignan-9,9'-diyl diacetate	<i>S. tonkinensis</i>	[41]
81	(-)-Secoisolariciresinol	<i>S. tonkinensis</i>	[41]
82	4,4'-Dihydroxy-3,3'-dimethoxy-9-ethoxy-9,9'-epoxylignan	<i>S. tonkinensis</i>	[41]
83	(2S,3R,4R)-4-[1-Ethoxy-1-(4-hydroxy-3-methoxyphenyl)methyl-2(4-hydroxy-3-methoxyphenyl)-3-hydroxymethyl-tetrahydrofuran	<i>S. tonkinensis</i>	[41]
84	(-)-Neo-olivil-(9-O-9'')-seco-isolariciresinol	<i>S. tonkinensis</i>	[41]

85	Isolariciresinol	<i>S. tonkinensis</i>	[41]
86	Stytonkinol A	<i>S. tonkinensis</i>	[42]
87	Stytonkinol B	<i>S. tonkinensis</i>	[42]
88	Styraxin	<i>S. argentifolius</i>	[43]
89	Vladinol D	<i>S. argentifolius</i>	[43]
90	5-Carboxy7-methoxy-2-(3',4'-methylenedioxyphenyl)benzofuran	<i>S. argentifolius</i>	[43]
91	5-((E)-2-Carboxyvinyl)-7-methoxy-2-(3',4'-methylenedioxyphenyl)benzofuran	<i>S. argentifolius</i>	[43]
92	(-)-Machicendiol	<i>S. argentifolius</i>	[43]
93	Machicendona	<i>S. argentifolius</i>	[43]

Terpenoids







94	Taraxerol	<i>S. japonica</i>	[7] [45]
95	Oleanolic aldehyde acetate	<i>S. japonica</i>	[22,54]
96	Erythrodil-3-acetate	<i>S. japonica</i>	[22,54]
97	Euphorginol	<i>S. japonica</i>	[22,54]
98	3 β -Acetoxylean-12-en-28-acid	<i>S. japonica</i>	[45]
99	3 β -Acetoxylean-17 β -hydroxy-28-norolean-12-ene	<i>S. japonica</i>	[45]
100	Anhydrosophoradiol-3-acetate	<i>S. japonica</i>	[22,54]
101	Styraxosides B	<i>S. japonica</i>	[44]
102	Camellenodiol	<i>S. japonica</i>	[44]
103	Jegosaponins A	<i>S. japonica</i>	[46]
104	Jegosaponins B	<i>S. japonica</i>	[46]
105	Jegosaponins C	<i>S. japonica</i>	[46]
106	Jegosaponins D	<i>S. japonica</i>	[46]
107	3 β -Acetoxylean-28-hydroxyolean-12-ene	<i>S. japonica</i>	[47]
108	3 β -Acetoxylean-12-en-28-acid	<i>S. japonica</i>	[47]
109	3 β -Acetoxylean-12-en-28-aldehyde	<i>S. japonica</i>	[47]
110	3 β -Acetoxylean-17 β -hydroxy-28-norolean-12-ene	<i>S. japonica</i>	[47]
111	6 β -Hydroxy-3-oxo-11 α ,12 α -epoxyolean-28,13 β -olide	<i>S. tonkinensis</i>	[4]
112	3 β ,6 β -Dihydroxy-11 α ,12 α -epoxyolean-28,13 β -olide	<i>S. tonkinensis</i>	[4]
113	3 β ,6 β -Dihydroxy-11-oxo-olean-12-en-28-oic acid	<i>S. tonkinensis</i>	[4]
114	3 β -Hydroxy-12-oxo-13HR-olean-28,19 β -olide	<i>S. tonkinensis</i>	[4]
115	19 β -Hydroxy-3-oxo-olean-12-en-28-oic acid	<i>S. tonkinensis</i>	[4]
116	6 β -Hydroxy-3-oxo-olean-12-en-28-oic acid	<i>S. tonkinensis</i>	[4]
117	Sumaresinolic acid	<i>S. tonkinensis</i>	[4,51]
118	Siaresinolic acid	<i>S. tonkinensis</i>	[4,51]
119	Oleanolic acid	<i>S. tonkinensis</i>	[4,51]
120	21-Benzoylbarringtonol C	<i>S. officinalis</i> L.	[4]
121	Styrax-saponin A	<i>S. officinalis</i> L.	[49]
122	Styrax-saponin B	<i>S. officinalis</i> L.	[49]











123	Styrax-saponin C	<i>S. officinalis</i> L.	[49]
124	Deacylsaponin	<i>S. officinalis</i> L.	[49]
125	Oleanonic acid	<i>S. tonkinensis</i>	[42] [51]
126	Betulonic acid	<i>S. tonkinensis</i>	[42] [51]
127	Corosolic acid	<i>S. tonkinensis</i>	[42] [51]
128	Maslinic acid	<i>S. tonkinensis</i>	[42] [51]
129	Epibetulinic acid	<i>S. tonkinensis</i>	[42]
130	Betulinic acid	<i>S. tonkinensis</i>	[42] [51]
131	2 α ,3 α ,24-trihydroxy-urs-12-en-28-oic acid	<i>S. argentifolius</i>	[43]
Aromatic Compounds			
132	<i>trans</i> -(Tetrahydro-2-(4-hydroxy-3-methoxyphenyl)-5-oxofuran-3-yl)methylbenzoate	<i>S. tonkinensis</i>	[52]
133	3-(4-Hydroxy-3-methoxyphenyl)-2-oxopropylbenzoate	<i>S. tonkinensis</i>	[52]
134	4-(<i>E</i>)-3-Ethoxyprop-1-enyl)-2-methoxyphenol	<i>S. tonkinensis</i>	[52]
135	Benzoic acid	<i>S. tonkinensis</i>	[52]
136	Vanillin	<i>S. tonkinensis</i>	[52]
137	Dehydrodivanillin	<i>S. tonkinensis</i>	[52]
138	Vanillic acid	<i>S. tonkinensis</i>	[52]
139	Coniferyl aldehyde	<i>S. tonkinensis</i>	[52]
140	Methylsyringin	<i>S. japonica</i>	[21]
141	Syringin	<i>S. japonica</i>	[7,21]
142	Isotachioside	<i>S. perkinsiae</i>	[39]
143	(2R,3R)-3,7,4'-Trihydroxy-5,3'-dimethoxyflavan 7-O- β -D-glucopyranoside	<i>S. suberifolius</i>	[55]
144	Methyl orsellinate	<i>S. suberifolius</i>	[55]
145	Ethyl orsellinate	<i>S. suberifolius</i>	[55]
146	Methyl β -orcinolcarboxylate	<i>S. suberifolius</i>	[55]
Steroids			
147	Stigmasterol	<i>S. macranthus</i>	[32]
148	Styraxosides A	<i>S. japonica</i>	[32]
149	Daucosterol	<i>S. macranthus</i>	[32]
150	β -Sitosterol	<i>S. perkinsiae</i>	[17]
151	(20R)-24Ethylcholest-5,22-dien-7-one	<i>S. argentifolius</i>	[43]
Others			
152	Kaempferol-3-O-(2'',4''-di-O-(<i>E</i>)-p-coumaroyl)- β -D-glucopyranoside)	<i>S. pohlii</i> <i>S. camporum</i>	[57] [57]
153	Kaempferol-3-O-(2'',6''-di-O-(<i>E</i>)-p-coumaroyl)- β -D-glucopyranoside	<i>S. pohlii</i> <i>S. camporum</i>	[57] [57]
154	Quercetin	<i>S. pohlii</i>	[57]













155	Kaempferol	<i>S. camporum</i>	[57]
		<i>S. pohlii</i>	[57]
		<i>S. camporum</i>	[58]
156	Koninginin T		
157	Koninginin U	<i>S. pohlii</i>	[64]
		<i>S. pohlii</i>	[64]
158	Jegosaponins A	<i>S. japonica</i>	[59]
159	Jegosaponins B	<i>S. japonica</i>	[59]












Appendix C





Table A3. Structural information of the active molecule.

No.	Structure	Phytochemistry	Pharmacological Activity
3		<i>S. perkinsiae</i> <i>S. obassia</i>	MCF-7 (IC ₅₀ = 53.5 µg/mL) Hela (IC ₅₀ = 23.3 µg/mL) HL-60 (IC ₅₀ = 16.8 µg/mL)
4		<i>S. perkinsiae</i> <i>S. japonica</i> <i>S. obassia</i> <i>S. agrestis</i> <i>S. ferrugineus</i> <i>S. pohlii</i> <i>S. camporum</i> <i>S. macranthus</i> <i>S. officinalis</i> <i>S. argentifolius</i>	Hep-2 (IC ₅₀ = 3.6 µg/mL) C6 (IC ₅₀ = 3.2 µg/mL) KB (IC ₅₀ = 22.11 µg/mL) HepG2 (IC ₅₀ = 18.15 µg/mL) Lu (IC ₅₀ = 21.50 µg/mL) Hemolytic activity (IC ₅₀ = 33 µg/mL) <i>C. sphaerospermum</i> (MIC = 5 mg/mL) <i>C. albicans</i> (MIC = 10 µg/mL) <i>S. aureus</i> (MIC = 10 µg/mL) ATCC 27853 and ATCC 6305 (MIC = 400 mg/mL) COX-1 (inhibition rate = 35.7% at 30 µM) Reduce doxorubicin- and methanesulfonate-induced DNA and chromosomal damage
5		<i>S. perkinsiae</i> <i>S. japonica</i> <i>S. suberifolius</i> <i>S. macranthus</i> <i>S. obassia</i>	MCF-7 (IC ₅₀ = 27.9 µg/mL) Hela (IC ₅₀ = 23.3 µg/mL) <i>S. aureus</i> (MIC = 20 µg/mL) <i>C. albicans</i> (MIC = 15 µg/mL) HL-60 (IC ₅₀ = 47.8 µg/mL)
6		<i>S. perkinsiae</i> <i>S. macranthus</i> <i>S. obassia</i> <i>S. pohlii</i>	Estrogen biosynthesis
7		<i>S. perkinsiae</i> <i>S. macranthus</i>	Estrogen biosynthesis
10		<i>S. perkinsiae</i>	Hemolytic activity (IC ₅₀ = 166 µg/mL)

11		<i>S. perkinsiae</i>	MCF-7 (IC ₅₀ = 5.5 µg/mL) MDA-MB-231 (IC ₅₀ = 3.81 µg/mL)
14		<i>S. perkinsiae</i> <i>S. japonica</i>	MCF-7 (IC ₅₀ = 15.0 µg/mL) MDA-MB-231 (IC ₅₀ = 13.71 µg/mL)
17		<i>S. ferrugineus</i> <i>S. camporum</i> <i>S. pohlii</i>	HeLa (IC ₅₀ = 5.3 µg/mL) C6 (IC ₅₀ = 4.9 µg/mL) <i>C. sphaerospermum</i> (MIC = 10 µg/mL) <i>C. albicans</i> (MIC = 12 µg/mL) <i>S. aureus</i> (MIC = 10 µg/mL) Th2 cytokines, iNOS, MMP-9 Reduce doxorubicin- and methanesulfonate-induced DNA and chromosomal damage
18		<i>S. ferrugineus</i> <i>S. pohlii</i>	<i>S. aureus</i> (MIC = 20 µg/mL) <i>C. albicans</i> (MIC = 20 µg/mL)
19		<i>S. ferrugineus</i>	<i>S. aureus</i> (MIC = 20 µg/mL) <i>C. albicans</i> (MIC = 15 µg/mL)
26		<i>S. japonica</i>	MMP-1 (inhibition rate = 62.1% at 10 µM)
28		<i>S. japonica</i>	Hemolytic activity (IC ₅₀ = 123 µg/mL)
30		<i>S. japonica</i>	DPPH (IC ₅₀ = 380 µM)
31		<i>S. japonica</i>	DPPH (IC ₅₀ = 278 µM)
32		<i>S. japonica</i>	DPPH (IC ₅₀ = 194 µM)
33		<i>S. japonica</i>	DPPH (IC ₅₀ = 260 µM)

50		<i>S. agrestis</i>	EeAChE ($IC_{50} = 1.4 \mu\text{g/mL}$) hAChE ($IC_{50} = 1.7 \mu\text{g/mL}$)
51		<i>S. agrestis</i>	EeAChE ($IC_{50} = 2.0 \mu\text{g/mL}$) hAChE ($IC_{50} = 2.7 \mu\text{g/mL}$)
52		<i>S. agrestis</i>	EeAChE ($IC_{50} = 1.4 \mu\text{g/mL}$) hAChE ($IC_{50} = 1.8 \mu\text{g/mL}$)
53		<i>S. agrestis</i>	EeAChE ($IC_{50} = 2.2 \mu\text{g/mL}$) hAChE ($IC_{50} = 3.1 \mu\text{g/mL}$)
86		<i>S. tonkinensis</i>	Hela ($IC_{50} = 26.75 \mu\text{M}$) MCF-7 ($IC_{50} = 45.16 \mu\text{M}$)
87		<i>S. tonkinensis</i>	MCF-7 ($IC_{50} = 57.1 \mu\text{M}$) Hemolytic activity ($IC_{50} = 65 \mu\text{g/mL}$)
92		<i>S. argentifolius</i>	KB ($IC_{50} = 96.01 \mu\text{g/mL}$) HepG2 ($IC_{50} = 86.60 \mu\text{g/mL}$) Lu ($IC_{50} = 106.86 \mu\text{g/mL}$)
95		<i>S. japonica</i>	Human dermal fibroblasts ($IC_{50} = 20 \mu\text{M}$)
96		<i>S. japonica</i>	MMP-1 (inhibition rate = 73.1% at $0.01 \mu\text{M}$)
100		<i>S. japonica</i>	Human dermal fibroblasts ($IC_{50} = 1.12 \mu\text{M}$)
101		<i>S. tonkinensis</i>	Hemolytic activity ($IC_{50} = 65 \mu\text{g/mL}$)
108		<i>S. japonica</i>	PTP1B ($IC_{50} = 7.8 \mu\text{g/mL}$)
109		<i>S. japonica</i>	PTP1B ($IC_{50} = 9.3 \mu\text{g/mL}$)
112		<i>S. tonkinensis</i>	HL-60 ($IC_{50} = 27.5 \mu\text{g/mL}$)

116		<i>S. tonkinensis</i>	HL-60 (IC ₅₀ = 14.2 µg/mL)
118		<i>S. tonkinensis</i>	HL-60 (IC ₅₀ = 29.0 µg/mL)
119		<i>S. tonkinensis</i>	HL-60 (IC ₅₀ = 8.9 µg/mL)
125		<i>S. tonkinensis</i>	hCES1A (IC ₅₀ = 0.49 mg/mL)
126		<i>S. tonkinensis</i>	hCES1A (IC ₅₀ = 1.48 µg/mL)
129		<i>S. tonkinensis</i>	hCES1A (IC ₅₀ = 0.041 µg/mL)
144		<i>S. suberifolius</i>	<i>A. solani</i> (inhibition rate = 58.41% at 100 µg/mL) <i>F. oxysporum</i> (inhibition rate = 67.39% at 100 µg/mL)
145		<i>S. suberifolius</i>	<i>A. solani</i> (inhibition rate = 59.31% at 100 µg/mL) <i>F. oxysporum</i> (inhibition rate = 45.65% at 100 µg/mL)
146		<i>S. suberifolius</i>	<i>F. oxysporum</i> (inhibition rate = 61.41% at 100 µg/mL) <i>P. cytospora</i> (inhibition rate = 86.72% at 100 µg/mL)
149		<i>S. macranthus</i>	iNOS, COX-2, TNF-α, and IL-1β
152		<i>S. pohlii</i> <i>S. camporum</i>	Separate coupled <i>Schistosoma mansoni</i> adult worms

155		<i>S. pohlii</i> <i>S. camporum</i>	Separate coupled <i>Schistosoma mansonii</i> adult worms Kill adult schistosomes
156		<i>S. pohlii</i>	AChE
158		<i>S. japonica</i>	Hemolytic activity (IC ₅₀ = 2.1 µg/mL)
159		<i>S. japonica</i>	Hemolytic activity (IC ₅₀ = 20.2 µg/mL)

Abbreviations

MCF-7	Metastatic breast adenocarcinoma cell line
MDA-MB-231	Human breast cancer cell line
Hep-2	Human larynx carcinoma cell line
HeLa	Human cervix carcinoma cell lines
C6	Rat glioma cell lines
HL-60	Human leukemia cell line
A549	Human lung cancer cell line
PC-3	Human prostatic cancer cell line
Lu	Lung cancer cell lines
CCK-8	Cell Counting Kit-8
ATCC 6305	<i>Streptococcus pneumoniae</i>
ATCC 19615	<i>Streptococcus pyogenes</i>
ATCC 10211	<i>Hemophilus influenzae</i>
ATCC 27853	<i>Pseudomonas aeruginosa</i>
ATCC 10031	<i>Klebsiella pneumoniae</i>
MIC	Minimum inhibitory concentration
DPPH	DPPH radical
HepG2	Hepatocellular carcinoma cell line
KGN	Human ovarian granulosa cells
AChE	Acetylcholinesterase
EeAChE	<i>Electrophorus electricus</i> AChE
hAChE	Human AChE
BChE	Butyrylcholinesterase
CTX	Cyclophosphamide
IFN-γ	Interferon-γ
hCES1A	Human carboxylesterase 1A1
MMP-1	Matrix metalloproteinase-1
PTP1B	The protein tyrosine phosphatase 1B
iNOS	Nitric oxide synthase
TNF-α	Tumor necrosis factor-α
IL-1β	Interleukin-1β
COX-1	Cyclooxygenase-1
COX-2	Cyclooxygenase-2

NF- κ BNuclear factor- κ B

References

1. Fritsch, P.W.; Almeda, F.; Renner, S.S.; Martins, A.B.; Cruz, B.C. Phylogeny and circumscription of the near-endemic Brazilian tribe Microlicieae (Melastomataceae). *Am. J. Bot.* **2004**, *91*, 1105–1114.
2. Pauletti, P.M.; Teles, H.L.; Silva, D.H.S.; Araújo, A.R.; Bolzani, V.S. The Styracaceae. *Rev. Bras. Farmacogn.* **2006**, *16*, 576–590.
3. Jiang, S. *Dictionary of Chinese Materia Medica*; Shanghai Scientific and Technical Publishing House: Shanghai, China, **1997**, p. 1723–1725.
4. Wang, F.; Hua, H.; Pei, Y.; Chen, D.; Jing, K.Y. Triterpenoids from the Resin of *Styrax tonkinensis* and Their Antiproliferative and Differentiation Effects in Human Leukemia HL-60 Cells. *J. Nat. Prod.* **2006**, *69*, 807–810.
5. Lee, T.B. *Illustrated Flora of Korea*; Hwang Mun Sa: Seoul, Republic of Korea, **1982**; p. 614.
6. Kitagawa, I.; Imakura, Y.; Hayashi, T.
7. 6. Kitagawa, I.; Imakura, Y.; Hayashi, T.; Yosioka, I. Structure of deacyl-jegosaponin, a common deacyl derivative of jegosaponin isolated from pericarps of *Styrax japonica* Sieb. et Zucc. *Chem. Pharm. Bull.* **1974**, *22*, 1675–1677. Min, B.S.; Na, M.K.; Oh, S.R.; Ahn, K.S.; Jeong, G.S.; Li, G.; Lee, S.K.; Joung, H.; Lee, H.K. New Furofuran and Butyrolactone Lignans with Antioxidant Activity from the Stem Bark of *Styrax japonica*. *J. Nat. Prod.* **2004**, *67*, 1980–1984.
8. He, L.; Zhou, Y.; Wan, G.J.; Wang, W.C.; Zhang, N.; Yao, L. Antinociceptive effects of flower extracts and the active fraction from *Styrax japonicus*. *J. Ethnopharmacol.* **2022**, *284*, 114779–114788.
9. Xie, Z.W. *Compilation of Chinese Herbal Medicine*; People's Medical Publishing House: Beijing, China, **1975**.
10. Liu, J.; Dumontet, V.; Simonin, A. Benzofurans from *styrax agrestis* as acetylcholinesterase inhibitors: Structure–activity relationships and molecular modeling studies. *J. Nat. Prod.* **2011**, *74*, 2081–2088.
11. Francielli de Oliveira, P.; Furtado, R.A.; Acésio, N.O.; Leandro, L.F.; Montanheiro, G.; de Padua, F.C.; Correa, M.B.; Braguini, C.G.; Pauletti, P.M.; Tavares, D.C. In Vivo Protective Activity of *Styrax camporum* Hydroalcoholic Extract against Genotoxicity Induced by Doxorubicin and Methyl Methanesulfonate in the Micronucleus and Comet Assays. *Planta. Med.* **2012**, *78*, 1899–1905.
12. Lu, D.; Yang, L.; Li, Q.L.; Gao, X.P.; Wang, F.; Zhang, G.L. Egonol gentiobioside and egonol gentiotrioside from *Styrax perkinsiae* promote the biosynthesis of estrogen by aromatase. *Eur. J. Pharmacol.* **2012**, *691*, 275–282.
13. Zhang, Y.; Zhao, X.C.; Xie, Y.G.; Fan, C.; Huang, Y.Y.; Yan, S.K.; Zhang, Y.; Jin, H.Z.; Zhang, W.D. Eight new γ -lactam alkaloids from the roots of the *Hemerocallis minor* Mill. *Fitoterapia* **2017**, *118*, 80–86.
14. Xie, Q.; Wu, G.Z.; Yang, N.; Shen, Y.H.; Tang, J.; Zhang, W.D. Delavatine A, an unusual isoquinoline alkaloid exerts anti-inflammation on LPS-induced proinflammatory cytokines production by suppressing NF- κ B activation in BV-2 microglia. *Biochem. Biophys. Res. Commun.* **2018**, *502*, 202–208.
15. Zhao, J.; Zhao, Q.; Lu, J.Z.; Ye, D.; Mu, S.; Yang, X.D.; Zhang, W.D.; Ma, B.L. Natural Nano-Drug Delivery System in Coptidis Rhizoma Extract with Modified Berberine Hydrochloride Pharmacokinetics. *Int. J. Nanomed.* **2021**, *16*, 6297–6311.
16. Yuan, W.L.; Huang, Z.H.; Li, B.; Xiao, S.J.; Shen, X.P.; He, Y.Q.; Zhang, W.D.; Shen, Y.H. Trisecocycloartane triterpene dilactones from the cones of *Pseudolarix amabilis*. *Phytochem. Lett.* **2021**, *45*, 88–92.
17. Li, Q.; Guo, Y.; He, X.; Li, G. Bifunctional Cu(II)-containing PDA-PEI copolymer dots: Demonstration of a dual-mode platform for colorimetric-fluorescent detection of glyphosate in the environment. *Talanta* **2023**, *265*, 124865.
18. Chen, Q.F.; Chen, X.Z.; Li, G.Y.; Wang, C.; Zhang, G.L. Two new 2-phenylbenzofurans from the bark of *Styrax perkinsiae*. *Chin. J. Nat. Med.* **2012**, *10*, 92–97.
19. Pauletti, P.M.; Araújo, A.R.; Young, M.C.M.; Giesbrecht, A.M.; Bolzani, V.D.S. nor-Lignans from the leaves of *Styrax ferrugineus* (Styracaceae) with antibacterial and antifungal activity. *Phytochemistry* **2000**, *55*, 597–601.
20. Kim, M.; Moon, H.; Chung, J.H.; Moon, Y.H.; Hahm, K.; Woo, E. Matrix metalloproteinase-1 inhibitor from the stem bark of *Styrax japonica* S. et Z. *Chem. pharm. Bull.* **2004**, *52*, 1466–1469.
21. Kim, M.; Moon, H.T.; Lee, D.G.; Woo, E.R. A new lignan glycoside from the stem bark of *Styrax japonica* S. et Z. *Arch. Pharm. Res.* **2007**, *30*, 425–430.
22. Min, B.; Oh, S.; Ahn, K. Anti-complement activity of norlignans and terpenes from the stem bark of *Styrax japonica*. *Planta Med.* **2004**, *70*, 1210–1215.
23. Wu, N.; Wang, L.; Chzn, Y.; Liao, Z.; Yang, G.Y.; Hu, Q.F. Lignans from the Stem of *Styrax japonica*. *Asian J. Chem.* **2011**, *23*, 931–932.
24. KwanKwonwon, Y.; UnKim, S. Phytochemical studies on seeds of *Styrax japonica* Sieb. et Zucc. *J. Appl. Biol. Chem.* **2002**, *45*, 28–30.
25. Hung, T.M.; Lee, M.K.; Min, B.; Kim, J.C.; Choi, J.S.; Lee, H.K. Two Lignans from the stem bark of *Styrax japonica*. *J. Korean Soc. Appl. Biol.* **2009**, *52*, 560–563.
26. Kinoshita, T.; Haga, Y.; Narimatsu, S. New nor-neolignan glycosides from *Styrax obassia* (Styracaceae). *Heterocycles* **2005**, *65*, 1471–1480.
27. Takanashi, M.; Takizawa, Y. New Egonol Glycoside Having D-Xylose, from Immature Seeds of *Styrax obassia* Sieb. et Zucc. *J. Oleo Sci.* **2002**, *51*, 423–426.

28. Takanashi, M.; Takizawa, Y. New benzofurans related to egonol from immature seeds of *Styrax obassia*. *Phytochemistry* **1988**, *27*, 1224–1226.
29. Takanashi, M.; Takizawa, Y.; Mitsunashi, T. Determination of Mercury by Direct Heating of Mercury-Adsorbed Ion-Exchange Resin in Electrothermal Atomic Absorption Spectrometry. *Chem. Lett.* **1974**, *8*, 869–871.
30. Park, S.Y.; Lee, H.; Lee, O.; Kang, H.; Choi, D.H.; Paik, K.H.; Khan, M. Benzofurans from the Seeds of *Styrax obassia*. *Bull. Korean Chem. Soc.* **2007**, *28*, 1874.
31. Lee, H.J.; Park, S.Y.; Lee, O.K.; Jo, H.J.; Kang, H.Y.; Choi, D.H.; Paik, K.H.; Khan, M. Benzofurans and sterol from the seeds of *Styrax obassia*. *Chem. Nat. Compd.* **2008**, *44*, 435–439.
32. Luo, Y.; He, Z.; Li, H. 2-Aryl benzofurans and their derivatives from seeds of *Styrax macranthus*. *Fitoterapia* **2007**, *78*, 211–214.
33. Schneiders, G.E.; Stevenson, R. Synthesis of (+)-machicendiol. *J. Org. Chem.* **1979**, *44*, 4710–4711.
34. Anil, H. Four benzofurans glycosides from *Styrax officinalis*. *Phytochemistry* **1980**, *19*, 2784–2786.
35. Akgul, Y.Y.; Anil, H. Benzofuran from seeds of *Styrax officinalis*. *Fitoterapia* **2003**, *74*, 743–745.
36. Akgul, Y.Y.; Anil, H. Benzofurans and another constituent from seeds of *Styrax officinalis*. *Phytochemistry* **2003**, *63*, 939–943.
37. Bacchi, E.M.; Sertie, J.A.; Villa, N.; Katz, H. Antiulcer action and toxicity of *Styrax camporum* and *Caesalpinia ferrea*. *Planta. Med.* **1995**, *61*, 204–207.
38. Teles, H.L.; Hemerly, J.P.; Pauletti, P.M. Cytotoxic lignans from the stems of *Styrax camporum* (Styracaceae). *Nat. Prod. Res.* **2005**, *19*, 319–323.
39. Zhang, P.Z.; Zhang, Y.M. Tetrahydrofuranoid Lignans from Stem Bark of *Styrax perkinsiae*. *Chin. J. Experimental Tradit. Med. Formulae* **2015**, *21*, 91–94.
40. Liu, B.L.; Ding, W.B.; Huang, S.Y.; Sun, W.H.; Li, Y.Z. Chemotaxonomic significance of phenylpropanoids from *Styrax suberifolius* Hook. Et Arn. *Biochem. Syst. Ecol.* **2018**, *78*, 35–38.
41. Hu, W.L.; Li, Z.L.; Chen, Q.J.; Sun, Y.W.; Zhai, S.; Lu, F.; Zhang, C.F. Triterpenes and lignans from the leaves of *Styrax tonkinensis*. *Biochem. Systemat. Ecol.* **2019**, *86*, 103891.
42. Wang, F.; Zhang, L.; Zhang, Q.; Chen, A.; Wang, S.M.; Fang, Z.F. Two new phenylpropanoids from the resin of *Styrax tonkinensis* (Pierre) Craib ex Hartw. *J. Nat. Med.* **2020**, *74*, 819–824.
43. Tra, N.T.; Tuyen, N.V.; Cuong, P.V.; Ha, N.T.T.; Anh, L.T.T.; Son, T.N. Chemical constituents from the leaves of *Styrax argentifolius* H.L. Li and their biological activities. *Phytochem. Lett.* **2021**, *41*, 70–73.
44. Moon, H.; Kim, M.; Woo, E.; Chung, J.H. Triterpenoid from *Styrax japonica* SIEB. et ZUCC, and its effects on the expression of matrix metalloproteinases-1 and type 1 procollagen caused by ultraviolet irradiated cultured primary human skin fibroblasts. *Biol. Pharm. Bull.* **2005**, *28*, 2003–2006.
45. Yun, K.; Min, B.; Kim, J.; Lee, K.T. Styraxoside A isolated from the stem bark of *Styrax japonica* inhibits lipopolysaccharide-induced expression of inducible nitric oxide synthase and cyclooxygenase-2 in RAW 264.7 cells by suppressing nuclear factor-kappa B activation. *Biol. Pharm. Bull.* **2007**, *30*, 139–144.
46. Yoshikawa, K.; Hirai, H.; Tanaka, M. Antisweet Natural Products. XV. Structures of Jegosaponins AD from *Styrax japonica* SIEB. et Zucc. *Chem. Pharm. Bull.* **2000**, *48*, 1093–1096.
47. Kwon, J.H.; Chang, M.J.; Seo, H.W.; Lee, J.H.; Min, B.S.; Na, M.; Kim, J.C.; Woo, M.H.; Choi, J.S.; Lee, H.K. Triterpenoids and a sterol from the stem-bark of *Styrax japonica* and their protein tyrosine phosphatase 1B inhibitory activities. *Phytother. Res.* **2008**, *22*, 1303–1306.
48. Anil, H. 21-Benzoyl-barringtonol C, a sapogenin from *Styrax officinalis*. *Phytochemistry* **1979**, *18*, 1760–1761.
49. Yayla, Y.; Alankuş-Çalışkan, Ö.; Anıl, H.; BBates, R.; Stessman, C.; Kane, V. Saponins from *Styrax officinalis*. *Fitoterapia* **2002**, *73*, 320–326.
50. Tayoub, G.; Schwob, I.; Bessiere, J.; Masotti, V.; Rabier, J.; Ruzzier, M.; Viano, J. Composition of volatile oils of *Styrax* (*Styrax officinalis* L.) leaves at different phenological stages. *Biochem. Syst. Ecol.* **2006**, *34*, 705–709.
51. Wang, L.; Guan, X.Q.; He, R.J.; Huo, P.C.; Qin, W.W.; Cui, L.T.; Hu, Q.; Hou, J.; Mamat, M.; Jia, S.N.; et al. Discovery and characterization of pentacyclic triterpenoid acids in *Styrax* as potent and reversible pancreatic lipase inhibitors. *J. Funct. Foods* **2020**, *74*, 104159.
52. Wang, F.; Hua, H.; Bian, X.; Pei, Y.H.; Jing, K.Y. Two new aromatic compounds from the resin of *Styrax tonkinensis* (Pier.) Craib. *J. Asian Nat. Prod. Res.* **2006**, *8*, 137–141.
53. Wang, F.; Fang, Z.F. Chemical Constituents from Resin of *Styrax tonkinensis*. *Chin. J. Exp. Tradi. Med. Formulae* **2012**, *18*, 17.
54. Kim, M.; Lee, H.; Hahm, K.; Moon, Y.H.; Woo, E. Pentacyclic triterpenoids and their cytotoxicity from the stem bark of *Styrax japonica* S. et Z. *Arch Pharm. Res.* **2004**, *27*, 283–286.
55. Vardar, Y.; Oflas, S. Preliminary studies on the *Styrax* oil. *Qual. Plant. Mater. Veg.* **1973**, *22*, 145–148.
56. Braguine, C.G.; Bertanha, C.S.; Gonçalves, U.O.; Magalhães, L.G.; Rodrigues, V.; Gimenez, V.M.M. Schistosomicidal evaluation of flavonoids from two species of *Styrax* against *Schistosoma mansoni* adult worms. *Pharm. Biol.* **2012**, *50*, 925–929.
57. Biasetto, C.R.; Somensio, A.; Sordia, R.; Chaplab, V.M.; Ebrahimi, S.N.; Silvad, G.H.; Telese, H.L.; Bolzania, V.S.; Youngf, M.C.M.; Pfenning, L.H.; et al. The new koniginins T-U from *Phomopsis stipata*, an endophytic fungus isolated from *Styrax camporum* Pohl. *Phytochem. Lett.* **2020**, *36*, 106–110.
58. Nishimura, M.; Fuchino, H.; Takayanagi, K.; Kawakami, H.; Nakayama, H.; Kawahara, H.; Shimada, Y. Toxicity of Jegosaponins A and B from *Styrax japonica* Siebold et al., Zuccarini in Prostate Cancer Cells and Zebrafish Embryos Resulting from Increased Membrane Permeability. *Int. J. Mol. Sci.* **2021**, *22*, 6354.

59. Francielli de Oliveira, P.; Damasceno, J.L.; Bertanha, C.S.; Bertanha, C.S.; Araujo, A.R.B.; Pauletti, P.M.; Tavares, D.C. Study of the cytotoxic activity of *Styrax camporum* extract and its chemical markers, egonol and homoegonol. *Cytotechnology* **2016**, *68*, 1597–1602.
60. Cao, T.Q.; Lee, B.M.; Jung, Y.W.; Nguyen, V.T.; Kim, J.A.; Min, B.S. Cytotoxic Activity of Compounds from *Styrax obassia*. *Nat. Prod. Commun.* **2017**, *12*, 259–260.
61. Bertanha, C.S.; Utrera, S.H.; Gimenez, V.M.M.; Groppo, M.; Silva, M.; Cunha, W.R.; Martins, C.H.G.; Januário, A.H.; Pauletti, P.M. Antibacterial evaluation of *Styrax pohlii* and isolated compounds. *Braz. J. Med. Biol. Res.* **2013**, *49*, 653–658.
62. Lee, J.; Lim, K. Apoptotic activity of ethanol extract from *Styrax Japonica* Siebold et al Zuccarini in HepG2 cells. *J. Ethnopharmacol.* **2010**, *131*, 210–215.
63. Lee, J.; Lim, K. Inhibitory effect of *Styrax Japonica* Siebold et al., Zuccarini glycoprotein (38 kDa) on interleukin-1 β and induction proteins in chromium (VI)-treated BNL CL. 2 cells. *Mol. Cell Biochem.* **2012**, *367*, 103–111.
64. Lee, J.; Lim, K. SJSZ glycoprotein (38 kDa) modulates expression of IL-2, IL-12, and IFN- γ in cyclophosphamide-induced Balb/c. *Inflamm. Res.* **2012**, *61*, 1319–1328.
65. Schreiber, F.G.; Stevenson, R. Synthesis of benzofuran *Styrax* extractives. *J. Chem. Soc. Perkin Trans.* **1976**, *1*, 1514–1518.
66. Moon, H.; Lee, J.; Chung, J.H. The effect of erythrodil-3-acetate on the expressions of matrix metalloproteinase-1 and type-1 procollagen caused by ultraviolet irradiated cultured primary old aged human skin fibroblasts. *Phytomedicine* **2006**, *13*, 707–711.
67. Lee, J.; Lim, K. Inhibitory effect of phyto glycoprotein (38 kDa) on expression of matrix metalloproteinase-9 in 12-O-tetradecanoylphorbol-13-acetate-treated HepG2 cells. *Naunyn Schmiedeberg's Arch. pharmacol.* **2011**, *384*, 185–196.
68. Bacchi, E.M.; Sertie, J.A. Antiulcer action of *Styrax camporum* and *Caesalpinia ferrea* in rats. *Planta Med.* **1994**, *60*, 118–120.
69. Bertanha, C.S.; Braguine, C.G.; Moraes, A.G.; Gimenez, V.M.; Groppo, M.; Silva, M.; Cunha, W.R.; Januário, A.H.; Pauletti, P.M. Cyclooxygenase inhibitory properties of nor-neolignans from *Styrax pohlii*. *Nat. Prod. Res.* **2012**, *26*, 2323–2329.
70. Bertanha, C.S.; Soares, E.; Gonçalves, U.O.; Francielli de Oliveira, P.; Magalhães, L.G.; Rodrigues, V.; Tavares, D.C.; Andrade e Silva, M.L.; Cunha, W.R.; Januário, A.H.; et al. In vitro Antischistosomal and Cytotoxic Activities of Norneolignans from *Styrax pohlii* A. DC. *J. Herbs Spices Med. Plants* **2014**, *20*, 394–401.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.