

Article

Influence of Land Use on the C and N Status of a C₄ Invasive Grass in a Semi-Arid Region: Implications for Biomonitoring

Edison A. Díaz-Álvarez ¹ and Erick de la Barrera ^{2,*}

¹ Instituto de Investigaciones Forestales, Universidad Veracruzana, Parque Ecológico “El Haya”, Carretera Antigua a Coatepec-Coapexpan, Xalapa, Veracruz 91070, Mexico; edisondiaz@uv.mx

² Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional Autónoma de México, Antigua Carretera a Pátzcuaro 8701, Col. Ex-Hacienda de San José del Cerro, Morelia Michoacán 58190, Mexico

* Correspondence: delabarrera@unam.mx

Abstract: Biomonitoring of atmospheric pollution is an increasingly accepted practice. However, most existing biomonitors are usually epiphytic species from mesic environments. This work assessed the suitability of buffelgrass (*Cenchrus ciliaris*), an invasive C₄ grass in northwestern Mexico, as a biomonitor, by means of the spatial distribution of the carbon and nitrogen content and isotopic signatures for grass samples collected from urban, agricultural, and natural areas throughout the state of Sonora. We found the highest tissue carbon content of 45.6% (on a dry weight basis) and highest nitrogen content of 3.31% for buffelgrass from the Yaqui Valley. We also found the lowest $\delta^{13}\text{C}$ of -15.9‰ , and the highest $\delta^{15}\text{N}$ of 16.7‰ in the same region. In contrast, the lowest carbon and nitrogen content of 39.4 and 1.49% were found for Bahía de Kino and Río Sonora mountains, respectively. The lowest $\delta^{15}\text{N}$ of 2.18‰ and the highest $\delta^{13}\text{C}$ of -13.7‰ were measured for two remote locations. These results show the influence that pollutant emissions, including agriculture and transportation, have on elemental and isotopic composition of vegetation. Buffelgrass is most adequate for tracking carbon and nitrogen emissions in arid environments and for determining alterations on nitrogen soil reactions, as a first approximation for saturation.

Keywords: biomonitoring; carbon emissions; environmental pollution; global ecology; nitrogen deposition; stable isotopes



Citation: Díaz-Álvarez, E.A.; de la Barrera, E. Influence of Land Use on the C and N Status of a C₄ Invasive Grass in a Semi-Arid Region: Implications for Biomonitoring. *Plants* **2021**, *10*, 942. <https://doi.org/10.3390/plants10050942>

Academic Editor: Giedrė Kacienė

Received: 13 March 2021

Accepted: 4 May 2021

Published: 9 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Anthropic activities, mainly those related to urbanization and agriculture, cause the release and subsequent deposition of substantial amounts of pollutants. Carbon and nitrogen compounds are two groups of pollutants of special concern given their noxious environmental effects, the better-known of which are climate change and nitrogen deposition [1–3]. Atmospheric pollution not only has significant impacts on natural environments, since it is considered one of the leading drivers of biodiversity loss, but it also is a matter of human health, as air pollution causes premature mortality and various diseases that have become a global burden [4–6].

Various methods can be used for monitoring atmospheric pollution, with the gold standard being the deployment of automated monitoring stations that record the atmospheric concentrations of various atmospheric pollutants such as ozone, sulfur dioxide, and heavy metals in a continuous and reliable manner [7]. However, because multiple point measurements are required to monitor vast areas, the deployment of numerous monitoring stations is necessary, which can represent excessive costs of operation, maintenance, and space. A more economical alternative is the use of passive collectors, such as those that are commonly used to quantify nitrogen deposition. This methodology is effective in tracking atmospheric pollutants in large areas, but has the disadvantage of requiring that

the collectors be replaced every few weeks and that the measurements are sensitive to variations of temperature, air movement, and humidity [8,9].

Another alternative is the use of biomonitors, i.e., plants that occur naturally in regions of interest, which have been shown to be particularly suited for assessing environmental pollution by carbon and nitrogen emissions [10,11]. Biomonitors further allow the integration of different environmental factors, including water availability, air temperature, and atmospheric pollution, as well as the respective changes in such parameters in both natural and modified environments [8,12,13]. Of particular promise are the so-called atmospheric biomonitors, plants whose mineral nutrition depends mostly on deposited or gaseous atmospheric sources. Alas, such epiphytic species tend to occur predominantly in mesic environments and tend to be less abundant in arid and semi-arid regions, thus limiting the widespread use of this “biotechnology”. To address this limitation, the elemental and isotopic composition of grass biomonitors have recently been shown to detect modifications to the nitrogen cycle in natural and modified environments, including within arid regions [8,12–15]. Stable isotopes have been deemed as a powerful analytical tool that allows an integral assessment of various biogeochemical and ecological processes [16]. In particular, as the carbon and nitrogen move through different planetary compartments, they are subject to different reactions and their consequent fractionations. This also applies to pollutants emitted by different human activities. For example, while -8‰ is a typical $\delta^{13}\text{C}$ value for the air in natural environments, the carbon emissions from coal, gasoline, diesel, and natural gas combustion are substantially more negative, ranging from -42 to -25‰ , a signal that is picked up by photosynthetic biomonitors [11,17–21]. For nitrogen emissions, oxidized compounds emitted by vehicular and industrial activities tend to have positive $\delta^{15}\text{N}$ values, contrasting with the very negative values measured for reduced compounds such as those emitted by agriculture and animal husbandry [8,10,15].

We assessed the potential of *Cenchrus ciliaris* (common name is buffelgrass) as a candidate for biomonitoring carbon and nitrogen pollution in a semi-arid region from northwestern Mexico, where this plant is abundant, and the environmental conditions disallow the widespread occurrence of epiphytes. Indeed, with annual precipitations as low as 140 mm, this plant is most able to germinate, develop, and spread effectively throughout the study region, reasons for which it was introduced to the southern United States in the 1940s and to northwestern Mexico in the 1950s for forage and fodder production [22–25]. Buffelgrass has since escaped and invaded vast areas, displacing numerous local species [26–28]. In fact, this African grass has become one of the most important invasive plants worldwide, as a result of its phenological and ecophysiological characteristics, including rapid growth and tolerance to drought, which stems from its C_4 photosynthesis [29].

Even though the human population in the state of Sonora amounts to 2 million and that productive activities presumably represent the emission of large amounts of atmospheric pollutants, such that their monitoring is required by the Mexican environmental regulation, automated monitoring stations have been deployed in Hermosillo, the state capital, and Ciudad Obregón, but they are not operational [30–32]. Monitoring air quality in the rest of the state is irregular, at best, and limited to sporadic measurements of SO_2 and particulate matter [32]. For these reasons, taking advantage of the regional abundance of this invasive grass, we measured its carbon and nitrogen elemental and isotopic composition by means of sampling leaves and seeds of this plant from 34 sites with various land uses in Sonora, Mexico (Figure 1).




Figure 1. Study area in Sonora, Mexico. Green circles indicate the sites where buffelgrass was collected. Numbers indicate the collection site from small (sites 13 and 21), medium (9, 33 and 11) and large urban areas (1–4, 15, 17–19), agricultural (5–8, 10, 14, 16, and 24–30) and rural sites (20, 22, 23, 31, 32, and 34). Image data: Google Earth; date: 20 April 2020.

2. Results

2.1. Environmental Characteristics

The mean monthly temperature averaged 23.5 ± 0.15 °C among the 34 collecting sites from Sonora, Mexico, with a minimum temperature of the coldest month of 7.5 ± 0.25 °C and a maximum temperature of the warmest month of 38.0 ± 0.15 °C (Table 1). The annual precipitation was 287 ± 20 mm throughout the study region. The soil at the study sites was mostly alluvial in origin, with a sandy clay loam texture and an electrical conductivity of 0.71 ± 0.14 mmhos/cm. The primary soil type for three quarters of the sites was either a cambisol, a luvisol, or a vertisol. Although a considerable variation was found in the study region, a Principal Components Analysis revealed that the environment was essentially indistinguishable among the various land-use classes considered (Figure 2; Table 2).

Table 1. Climate and soil attributes from the study region in Sonora, Mexico. Data are means \pm 1 S.E. (n = 34 sites).

Parameter	Value
Mean temperature (°C)	23.5 ± 0.1
Minimum temperature (°C)	7.5 ± 0.2
Maximum temperature (°C)	38.0 ± 0.1
Annual precipitation (mm)	287 ± 20
Primary soil type	Cambisol (26% of sites), luvisol (26%), vertisol (24%), leptosol (12%), regosol (6%), phaeozem (6%)
Geological origin	Alluvial (85%), sedimentary (12%), igneous (3%)
Clay content (%)	32.2 ± 3.0
Silt content (%)	22.5 ± 2.0
Sand content (%)	45.3 ± 3.6
Electrical conductivity (mmho/cm)	0.71 ± 0.14
pH	8.45 ± 0.14




Figure 2. Principal Components Analysis for climate and soil characteristics from 34 collection sites with different land use. Data are plotted along the two main components from Table 2 for sites with agricultural (circles), large (squares), medium (triangles), and small (open circles) urban, or rural (diamonds) use. The numbers are the site identity from Figure 1. Ellipses indicate the 95% confidence interval for each land-use class. Percentages along the axes indicate the variance explained by each component.

Table 2. Principal Components Analysis for climate and soil attributes from 34 sites in Sonora, Mexico. Data are the component loadings for each attribute considered and the percent of the variance explained by each component individually and cumulative. For each component, the *p*-values are from a Kruskal–Wallis test for the axis values for each land use class.

	Principal Component								
	1	2	3	4	5	6	7	8	9
Sand	0.32	0.39	−0.28	0.14	−0.35	−0.28	0.29	−0.04	0.59
Silt	0.03	0.38	0.37	−0.55	0.47	0.19	0.05	0.07	0.39
Clay	−0.29	−0.54	0.03	0.19	0.04	0.13	−0.27	−0.01	0.70
Minimum temperature	−0.51	0.21	0.06	−0.13	−0.13	−0.27	−0.09	−0.76	−0.00
Maximum temperature	0.13	−0.39	−0.63	−0.44	0.32	−0.06	0.27	−0.24	−0.00
Mean Temperature	−0.49	0.03	−0.13	−0.24	−0.02	−0.58	−0.01	0.58	0.00
Precipitation	−0.35	0.30	−0.47	−0.15	−0.29	0.65	−0.12	0.14	0.00
Electrical conductivity	0.40	0.03	−0.13	−0.31	−0.16	−0.18	−0.82	−0.02	−0.00
pH	−0.07	0.35	−0.35	0.50	0.65	−0.09	−0.27	−0.02	−0.00
Individual variance	0.41	0.20	0.13	0.12	0.07	0.04	0.03	0.00	0.00
Cumulative variance	0.41	0.61	0.74	0.86	0.93	0.97	1.00	1.00	1.00
<i>p</i> -value	0.328	0.812	0.036	0.232	0.812	0.079	0.814	0.085	1.00

2.2. Elemental and Isotopic Composition for Buffelgrass

Over the study region the leaf carbon content of buffelgrass ranged from 39.4% (dry weight basis), for plants from site No. 21 located near the coast, to 45.6%, for plants from site No. 10 within the Yaqui Valley, averaging $42.1 \pm 0.16\%$ throughout the study region (Figure 3A). The lowest seed carbon content of 42.8% was also measured for the plants collected at site No. 21, while the highest seed carbon content of 51.4% was found for site no.31 in a natural protected area (Figure 3B). For medium urban sites, carbon content of leaves was higher than for rural and small urban sites. The carbon content for seeds averaged $47.5 \pm 0.89\%$ and it was significantly higher than for the leaves ($p = 0.001$; Tables 3 and A1; Figure 4). Land use had a significant effect on the carbon content of

C. ciliaris as significant differences were found among the five land uses ($p = 0.001$). Seeds from small urban sites had a lower carbon content than those from rural and medium urban sites. The interaction organ \times land use was also significant ($p = 0.002$; Table 3; Figure 4A).




Figure 3. Spatial distribution for the elemental composition for buffelgrass in Sonora, Mexico. Carbon content (C%, dry weight) for (A) leaves and (B) seeds, nitrogen content (N%, dry weight) for (C) leaves and (D) seeds, and the C:N ratio for (E) leaves and (F) fascicles. Image data: Google Earth; date: 20 April 2020.

Table 3. Two-way PERMANOVA for carbon and nitrogen parameters of buffelgrass growing in Sonora, Mexico. *p*-values in **bold** indicate significant differences.

	d.f.	C Content (% Dry Weight)		N Content (% Dry Weight)		C:N Ratio		$\delta^{13}\text{C}$ (‰)		$\delta^{15}\text{N}$ (‰)	
		F	<i>p</i>	F	<i>p</i>	F	<i>p</i>	F	<i>p</i>	F	<i>p</i>
Organ	1	463.2	0.001	172.3	0.001	276.98	0.001	94.641	0.001	0.741	0.416
Land use	4	6.261	0.001	5.109	0.002	4.529	0.001	4.892	0.002	12.89	0.001
Organ \times Land use	4	4.805	0.002	2.285	0.057	2.918	0.015	4.037	0.003	2.954	0.012




Figure 4. Elemental and isotopic carbon and nitrogen composition for leaves and seeds of *C. ciliaris* collected from sites with different land use. (A) Carbon and nitrogen content (dry mass basis) for leaves (open symbols) and seeds (closed symbols) collected from small (circles), medium (inverted triangles), and large cities (squares), agricultural lands (diamonds), and rural sites (triangles). (B) Carbon and nitrogen isotopic signatures for leaves and seeds of *C. ciliaris* collected from sites with different land use. Data are shown as mean \pm S.E. See underlying data in Appendix A.

The lowest leaf nitrogen content was 1.49%, found for the plants collected at site no.16, a remote rural location close to the Río Sonora mountains. Conversely, the highest nitrogen content of 3.31% was found for plants growing at site no.14, also a rural location but, in this case, close to a secondary road and an agricultural field (Figure 3C). On average, the leaf N content reached $2.52 \pm 0.05\%$ and it was significantly lower, $1.60 \pm 0.04\%$, than for the seeds ($p = 0.001$; Table 3). The leaves from agricultural sites had the highest nitrogen content, small urban sites had the lowest. The lowest nitrogen content for seeds was found for plants from site no. 12 that reached 0.99%, while the highest content was 2.35% for site no. 11 in a rural area (Figure 3D). Land use had a significant effect on the nitrogen content of *C. ciliaris* ($p = 0.002$). For example, this parameter was higher for seeds from medium urban sites than for those from rural sites. However, the interaction between organ \times land use was not significant ($p = 0.057$; Table 3; Figure 4A).

The C:N ratio averaged 17.98 ± 0.49 for leaves, ranging from 11.9 at site no. 29 to 28.9 at site no. 16 (Figure 3E). Significant differences were observed among land uses ($p = 0.002$; Table 3). C:N ratio for leaves from large urban was higher than for agriculture. For the seeds, the C:N ratio averaged 31.85 ± 0.96 , with the highest value of 49.6 found for the plants from site no. 31, and the lowest value in site no.29, which reached 20.5 (Figure 3F). Significant differences were also observed for the C:N ratio between leaves and seeds ($p = 0.001$; Table 3). In addition, the C:N ratios among land uses were different ($p = 0.001$), and the interaction between organ \times land use for *C. ciliaris* was significant ($p = 0.015$; Table 3). Seeds collected from rural sites had the highest C:N ratios, while the lowest were from agricultural sites.

The $\delta^{13}\text{C}$ values for the leaves ranged from -15.9‰ for plants growing in site no. 17, in the city of Hermosillo to -13.7‰ for plants collected in site no. 29, near a road and a wheat field, the land use had a significant effect on this parameter ($p = 0.002$; Table 3; Figure 5A). The $\delta^{13}\text{C}$ values for the leaves from small urban sites and the lowest was found in large urban sites (Figure 4). The average $\delta^{13}\text{C}$ value for seeds was $-14.3 \pm 0.30\text{‰}$. The lowest $\delta^{13}\text{C}$ value for seeds reached -17.2‰ for plants growing at site no. 31 in a natural protected area, while the highest value of -13.9‰ was measured for plants growing at site no. 13 in a small urban center, significant differences were observed among land uses ($p = 0.002$; Table 3; Figure 5B). The $\delta^{13}\text{C}$ values for *C. ciliaris* were significant for the organs ($p = 0.001$), also, the interaction between organ \times land use was significant ($p = 0.003$; Figure 4B; Table 3). In addition, the seeds from small urban sites were more enriched than those from rural sites (Figure 4).

The $\delta^{15}\text{N}$ values for leaves ranged from 2.18 to 16.7‰ , respectively for plants collected near a highway in an isolated area (site no. 12) and a site where urban and cultivated fields join (site no. 9). Significant differences were found among the different classes of land use ($p = 0.001$; Table 3; Figures 4B and 5C). The $\delta^{15}\text{N}$ values for the leaves of buffelgrass averaged $8.37 \pm 0.20\text{‰}$ and $8.77 \pm 0.26\text{‰}$ for the seeds, throughout the study area. The lowest $\delta^{15}\text{N}$ value was found for leaves from small urban sites, in contrast agriculture and rural sites had the highest $\delta^{15}\text{N}$ value (Figure 4). The lowest value of 2.71‰ was found for seeds at the most remote point in site no. 12, the highest of 16.4‰ was found near a highway in site no. 10 (Figures 4B and 5D). No differences were found for the $\delta^{15}\text{N}$ values between leaves and seeds ($p = 0.416$; Table 3). On the contrary, the interaction between organ \times land use was significant ($p = 0.012$; Table 3). For seeds, the $\delta^{15}\text{N}$ values were highest for medium urban and lowest for large urban sites (Figure 4).




Figure 5. Spatial distribution for buffelgrass isotopic composition in Sonora, Mexico. Carbon isotopic composition ($\delta^{13}\text{C}$, ‰) for (A) leaves and (B) seeds and nitrogen isotopic composition ($\delta^{15}\text{N}$, ‰) for (C) leaves and (D) seeds. Image data: Google Earth; date: 20 April 2020.

3. Discussion

Plant carbon content is influenced by multiple environmental factors, including water and nutrient availability, the prevailing CO_2 concentration where plants grow [33]. Buffelgrass can become very successful in allocating biomass to its organs under adverse environmental conditions, owing to physiological traits such as the use of C_4 photosynthesis, which improves their water use efficiency [34]. Indeed, the grass can maintain its maximum photosynthetic gas exchange rates under water limitations [35]. These physiological advantages are further enhanced under elevated CO_2 concentrations, which are locally present in urban and agricultural environments, effectively stimulating biomass and carbon accumulation by this species [22,36–38]. The carbon isotopic composition of C_4 plants generally responds to water availability, with discrimination against ^{13}C increasing with aridity [39–41]. Given that the underlying environmental characteristics of the collection sites were indistinguishable among land-use classes, the differences that we observed in the isotopic signatures of buffelgrass were assumed to be, at least in part, caused by local differences in gaseous carbon concentrations and sources. This has been especially documented in urban environments, where the intensive use of fossil fuel for vehicles and industry increases the local concentrations of anthropogenic CO and CO_2 , whose distinctive isotopic signatures can be tracked by the vegetation growing in these environments [11,17].

The lowest carbon content, which we found for plants collected from a small coastal locality that is mainly dedicated to tourism (Site no. 21), was accompanied by the most positive $\delta^{13}\text{C}$ values of the study area, similar to the isotopic signatures for other C_4 grasses in natural environments [42]. In turn, the highest carbon content was found at various

sites within the Yaqui Valley, dominated by agricultural activities that presumably lead to a locally high CO₂ concentration, not only from the use of agricultural machinery and the transportation of supplies, products, and workers, but also from biogenic emissions released by agricultural soil nutrient management, such as the use of synthetic fertilizers and, in many cases, the burning of crop residues [43–45]. The carbon compounds that are released from biomass burning also have isotopic signatures that are considerably more negative than those from “clean” environments, leaving a distinctive signature on the exposed grasses [18,21,46]. Additionally, the highest nitrogen emissions from the biomass burning typical of anthropic environments have a significant effect on the carbon content, as the increased nitrogen availability can increase photosynthetic rates [13,47].

The nitrogen concentration of plants growing in anthropic environments is strongly related to the amount of foliar uptake of reactive nitrogen species and to the rate of nitrogen deposition on the soil surface [13,48]. Plants growing close to a significant source of emissions such as traffic, industries, cultivated fields, or animal husbandry, have been found to increase considerably their nitrogen concentration, changing, from 0.8% in rural to 3.6% (dry weight) in highly urbanized areas [8,10,13]. Buffelgrass followed this pattern, with the lowest nitrogen concentration measured for individuals from a remote site with scant human activity, where nitrogenous emissions are presumably lower than for other sites included in this study, and the highest concentration found for plants collected near agricultural fields or major roads. In this respect, anthropogenic compounds, such as nitrogen oxides (NO_x) and ammonia (NH₃) are easily assimilated by plants via gas exchange (dry deposition). These compounds can also be dissolved into water to be deposited onto the soil by rain (wet deposition), where they can be taken up by the roots, further increasing the plant nitrogen concentration [13,14,49].

The nitrogen isotopic composition of the vegetation depends on different factors, with the regional distribution of human activity being one of the most important [14,50,51]. For example, plants growing in rural and natural sites, where either the nitrogen deposition rates are low (< 5 Kg N ha⁻¹ year⁻¹) or the nitrogenous emissions are low, tend to have δ¹⁵N values that are negative but close to zero [52]. In these areas, the main sources for nitrogen are lightning and biological fixation by legumes, which do not discriminate between isotopes. On the other hand, soil type and its particular isotope signature also play a large role in plant δ¹⁵N values, this is particularly important for soils in natural regions. However, when the nitrogen emissions and deposition increase, for example, as a consequence of traffic or agriculture, the importance of soil type and their original δ¹⁵N values for plants isotope compositions becomes reduced [14,53]. We observed this effect for the different land uses comprising our study area. In particular, as the human activity increased δ¹⁵N values became more positive. In addition, in natural regions the microbial and vegetation N cycling are closely coupled, i.e., no important modifications because land use or atmospheric deposition are presented and N cycling is “in equilibrium”, for these regions the leaching and gaseous N losses are negligible, thus the isotopic fractionation is low, and the δ¹⁵N values are generally negative close to zero [54–57].

The δ¹⁵N values that we measured for buffelgrass were consistently positive, regardless of the site of collection. For plants growing in urban environments, this has been attributed to the interaction of two main factors. First, urban vegetation is exposed to gaseous NO_x which are taken up by direct stomatal absorption, these pollutants emitted by industry and cars are usually enriched in ¹⁵N, with δ¹⁵N values that can reach up to 10‰, which explains in part the isotopic signature that we found for buffelgrass in those environments [10,14,47,49,51]. The NO_x can be dissolved in water as nitrate (NO₃⁻) and deposited either by rain, fog, or snow onto the soil and reach the urban vegetation, their δ¹⁵N values are generally positive [54,58]. An additional source of nitrogen is reduced gaseous NH₃ or dissolved NH₄⁺, compounds that tend to be ¹⁵N depleted [54]. However, reduced nitrogen compounds do not abound in cities, thus representing a small proportion of the nitrogen sources for urban vegetation [10,54]. Second, urban vegetation, especially ruderal plants such as buffelgrass, have positive δ¹⁵N values because they root on modified

soils, in which the nitrogen reactions are altered leading to a ^{15}N enrichment as discussed below [44,57–62]. Therefore, the combination of the constant stomatal uptake of oxidized nitrogen by leaves and roots, coupled with rooting on modified soils contributed to the observed positive $\delta^{15}\text{N}$ values for buffelgrass in the urban environments [14]. As it occurs for plants from urban environments, those growing near roads and highways and even from isolated sites are exposed to the same process, but to a lesser extent. This explains why the $\delta^{15}\text{N}$ values for buffelgrass changed in the human activity gradient from very positive in urban to less positive closer to zero in isolated regions where emissions are low.

For agricultural areas, the nitrogenous fertilizer applied to crops can be taken up by plants, released as a gas, retained in the soil, or leached to the groundwater. For the case of the Yaqui Valley in our study region, one third of the applied nitrogen ends up in harvested crops, the other two thirds are released to the environment [63,64]. Not only for agricultural lands, but also for areas with other land uses, increasing rates of deposition result in soil nitrogen losses, which produces the ^{15}N enrichment of the available nitrogen in soil for plants [14,53,65–68]. For example, N_2O emissions are produced in agricultural fields after fertilization and in ornamental lawns when the incomplete transformation of NH_4^+ to nitrate, or during the incomplete transformation of NO_3^- to N_2 occurs, nitrification or denitrification, respectively. N_2O emissions can reach $\delta^{15}\text{N}$ values as negative as -48.2‰ [68,69]. NH_3 volatilization is another form of nitrogen loss from agricultural and other soils exposed to increasing rates of nitrogen inputs; it has $\delta^{15}\text{N}$ values that reach -56‰ [70,71].

Both N_2O and NH_3 can be directly absorbed by plants growing near agricultural fields, which would result in very negative $\delta^{15}\text{N}$ values [50,58]. Another potential N source with negative $\delta^{15}\text{N}$ values is NO_3^- , which is massively used in the study region for crop fertilization, such that it is lixiviated in high quantities, e.g., 8200 Mg N/yr just in the Yaqui Valley, most of which reaches the groundwater and is exported to the Gulf of California [63,72], similar to that which occurs in other agricultural lands [64]. However, such isotopic signals were not detected here. Considering that fertilizers are applied directly to the crops and we collected in the vicinity, but not directly in the agricultural plots, the positive $\delta^{15}\text{N}$ values that we measured for buffelgrass in agricultural sites may have multiple origins. For instance, when enough nitrogen accumulates in the soil, as a consequence of high rates of nitrogen deposition, gaseous reactive species of nitrogen (NO_x or NH_3) have been shown not to have a significant effect on the $\delta^{15}\text{N}$ value of grasses regardless of their local atmospheric concentration [13,50]. Additionally, the ^{15}N enrichment that results from N loss is common in disturbed soils, such as those where we collected buffelgrass [13,53,65–67,73]. In addition, the high rates of deposition of NO_x -derived compounds that are prevalent in many areas with anthropogenic activities may help explain the positive $\delta^{15}\text{N}$ values found in the study area [74,75].

The isotopic differences that we found among the various land-use classes may be influenced, at least in part, by existing differences in the isotopic fractionation rates of soil processes involved in nitrogen loss [64–66], which in turn respond to the amount and type of available nitrogen. The signal of the nitrogen remaining in the soil is picked up by the plants. However, higher rates of nitrogen deposition do not necessarily result in higher carbon or nitrogen content in plant tissues because negative feedback mechanisms can reduce or even halt further assimilation of nitrogen beyond species-specific physiological thresholds, which have been documented for various epiphytic orchids, terrestrial herbs, and trees [8,47,66,67,76]. Whether the amount of nitrogen available in the study region surpassed such a threshold for buffelgrass should be considered in future work by determining both the nitrogen-tolerance of this species and conducting measurements of the prevailing forms of nitrogen available in the various atmospheric and edaphic compartments. Our results, however, provide some indication of the regions in Sonora that are more susceptible of reaching such a nitrogen saturation threshold, including the Yaqui Valley, Coast of Hermosillo, and the two large cities of the region.

As the geographical expansion of buffelgrass continues in Mexico, owing to its tolerance to high temperatures and low water availability, its use as a biomonitor of nitrogen deposition can be a silver lining for this biological invasion as the grass' elemental and isotopic composition appears to respond directly to the intensity of anthropogenic disturbance under different land uses.

4. Materials and Methods

4.1. Study Region

The study was conducted in the state of Sonora, located in northwestern Mexico, along the Gulf of California (Figure 1). Collection sites were selected by the spontaneous occurrence of buffelgrass in this heterogeneous landscape. Each site was assigned to one of five classes of land use along a gradient of intensity of human activity [77]: (1) urban settlements from 1 to 49,999 habitants (Small urban), (2) urban settlements from 50,000 to 399,000 (Medium urban), (3) settlements more than 400,000 (Large urban), (4) agriculture and, (5) rural landscape. The study area included two important agricultural regions, the Yaqui Valley (6 sites), accredited to be the birthplace of the Green Revolution, comprises ca. 230,000 ha of irrigated cropland, mainly for the production of wheat and maize [74]. The coast of Hermosillo (10 sites), with 1.5 million ha, also under irrigation, is mainly dedicated to the production of various horticultural crops, such as table grapes and watermelon. The study region also contained two major cities, Hermosillo (population 812,229) and Ciudad Obregón (433,050), as well as various mid-sized settlements [28,78].

Climate and soil characteristics were determined for each collection site. In particular, the minimum temperature of the coldest month, the maximum temperature of the warmest month, the mean monthly temperature, and the annual precipitation were obtained for the 1961–2001 normals, through climatic spline models for each site [79]. In turn, the soil's geological origin, primary soil type, clay, silt, and sand content, electrical conductivity, and pH were obtained for the nearest reference point from the National soil profile database [80].

4.2. Sampling of Plant Material

Tissue samples of the invasive exotic grass *Cenchrus ciliaris* L. 1771, which is amply distributed in the study region [22,23], were collected from 34 sites from Sonora, Mexico, in March 2017. In particular, fully developed green leaves and mature seeds (fascicles were left intact) were collected at each site from each one of 5 individuals growing at least 5 m apart. The samples were placed in paper envelopes and kept in a shaded cardboard container during its transport to the laboratory, which usually occurred within 2 days.

4.3. Elemental and Isotopic Analysis

The plant materials were dried at 60 °C in a gravity convection oven until reaching constant weight, usually within 48 h. The dried tissues were ground to a fine powder in a ball mill, wrapped into tin capsules (Costech Analytical, Inc. Valencia, CA, USA), and weighed with a microbalance (resolution of 0.01 mg, Sartorius, Göttingen, Germany). For each sample, the carbon and nitrogen content, as well as the corresponding isotopic composition, were determined at the Stable Isotope Facility University of Wyoming (Laramie, WY, USA). The analyses were conducted with a Costech 4010 elemental analyzer (Costech Analytical Inc., Valencia, CA, USA) attached to a continuous flow isotope ratio mass spectrometer (Finnigan Delta Plus XP, Thermo Electron Corp, Waltham, MA, USA). Carbon and nitrogen isotope ratios, reported in parts per thousand, were calculated relative to V-PDB or atmospheric air by means of glutamic acid and alfalfa secondary standards (accessions: 36-UWSIF-Glutamic 1, 39-UWSIF-Glutamic 2, and UWSIF05). The analytical precision for the $\delta^{13}\text{C}$ was $0.3 \pm 0.02\text{‰}$ and $0.4 \pm 0.03\text{‰}$ (SD) for $\delta^{15}\text{N}$. The natural abundances of ^{13}C and ^{15}N were calculated as:

$$\delta^{13}\text{C} (\text{‰ versus V-PDB}) = (R_{\text{sample}}/R_{\text{standard}} - 1) \times 1000$$

$$\delta^{15}\text{N} (\text{‰ versus air}) = (R_{\text{sample}}/R_{\text{standard}} - 1) \times 1000$$

where R is the ratio of $^{13}\text{C}/^{12}\text{C}$ for carbon and $^{15}\text{N}/^{14}\text{N}$ for nitrogen isotope abundance for a given sample [81,82].

4.4. Statistical Analyses

To determine whether an underlying grouping of soil and climate characteristics existed for the five land-use categories, we conducted a Principal Components Analysis, excluding the categorical attributes of soil geological origin and primary type, followed by a Kruskal–Wallis test by ranks for the values of the collection sites along each principal component.

Differences among land use and organs for the plant carbon and nitrogen contents, the C:N ratio, as well as for the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values, were determined by means of a non-parametric two-way PERMANOVA (factors were organ and land use class). Pairwise comparisons were made by means of the function pairwise.adonis 2 from the ‘Vegan’ package in R ($p < 0.05$). Analyses were conducted in R (version 3.5.3, R Core Team, R foundation for Statistical Computing, Vienna, Austria), except for the Principal Components Analysis that was run on ClustVis [83].

4.5. Geostatistical Analysis

The ordinary Kriging method [84] within ArcGIS 10.5 (Esri, Redlands, California, USA), was used to determine the spatial distribution of plant carbon and nitrogen content, the C:N ratio, and the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values based on the interpolation of the point data measured for buffelgrass.

Author Contributions: Conceptualization, methodology, investigation: E.A.D.-Á., E.d.l.B.; formal analysis, writing—original draft preparation: E.A.D.-Á.; writing—review and editing, funding acquisition: E.d.l.B. All authors have read and agreed to the published version of the manuscript.

Funding: Research and APC were funded by the Dirección General de Asuntos del Personal Académico, Universidad Nacional Autónoma de México, grants PAPIIT IN205616 and IN211519.

Data Availability Statement: Data are available from the corresponding author on reasonable request.

Acknowledgments: We thank useful discussions with E.A. Yépez, as well as his generous hospitality in Ciudad Obregón, Sonora. We also thank the Huatabampo Police Department for releasing us while collecting within the Yaqui Valley.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Underlying data on the elemental and isotopic carbon and nitrogen status for leaves and seeds of *C. ciliaris* collected from sites with different land use.

Land-Use Class	Carbon Content (% Dry Mass)		Nitrogen Content (% Dry Mass)		C:N (Adimensional)		$\delta^{13}\text{C}$ (‰)		$\delta^{15}\text{N}$ (‰)	
	Leaves	Seeds	Leaves	Seeds	Leaves	Seeds	Leaves	Seeds	Leaves	Seeds
Small urban	41.13 ± 0.64	44.31 ± 0.54	2.31 ± 0.20	1.75 ± 0.22	18.96 ± 1.43	29.05 ± 3.31	−14.11 ± 0.12	−14.13 ± 0.14	6.56 ± 0.42	9.32 ± 0.52
Medium urban	43.31 ± 0.53	48.48 ± 0.58	2.71 ± 0.20	1.90 ± 0.17	17.30 ± 1.33	29.81 ± 3.41	−14.65 ± 0.16	−15.51 ± 0.33	12.23 ± 1.04	11.40 ± 0.96
Large urban	41.95 ± 0.30	47.15 ± 0.32	2.32 ± 0.09	1.73 ± 0.08	19.34 ± 0.91	29.39 ± 1.27	−14.67 ± 0.17	−15.58 ± 0.23	7.33 ± 0.24	8.01 ± 0.26
Agriculture	42.17 ± 0.25	47.15 ± 0.32	2.86 ± 0.07	1.81 ± 0.06	15.66 ± 0.57	27.68 ± 0.85	−14.28 ± 0.06	−15.39 ± 0.15	8.67 ± 0.29	8.57 ± 0.35
Rural	41.12 ± 0.19	48.91 ± 0.61	2.48 ± 0.10	1.51 ± 0.08	17.29 ± 0.74	35.38 ± 2.47	−14.14 ± 0.08	−16.12 ± 0.33	8.92 ± 0.30	8.85 ± 0.33

References

- Foley, J.A.; DeFries, R.; Asner, G.P.; Barford, C.; Bonan, G.; Carpenter, S.R.; Chapin, F.S.; Coe, M.T.; Daily, G.C.; Gibbs, H.K.; et al. Global Consequences of Land Use. *Science* **2005**, *309*, 570–574. [[CrossRef](#)]
- Houghton, R.A.; House, J.I.; Pongratz, J.; van der Werf, G.R.; DeFries, R.S.; Hansen, M.C.; Ramankutty, N. Carbon emissions from land use and land-cover change. *Biogeosciences* **2012**, *9*, 5125–5142. [[CrossRef](#)]
- Gasser, T.; Ciais, P. A theoretical framework for the net land-to-atmosphere CO₂ flux and its implications in the definition of “emissions from land-use change”. *Earth Syst. Dyn.* **2013**, *4*, 171–186. [[CrossRef](#)]
- Sala, O.E.; Chapin, F.S., III; Armesto, J.J.; Berlow, E.; Bloomfield, J.; Dirzo, R.; Huber-Sanwald, E.; Huenneke, L.F.; Jackson, R.B.; Kinzing, A.; et al. Global biodiversity scenarios for the year 2100. *Science* **2000**, *287*, 1770–1774. [[CrossRef](#)]
- Cohen, A.J.; Brauer, M.; Burnett, R. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: An analysis of data from the Global Burden of Diseases Study 2015. *Lancet* **2017**, *389*, 1907–1918. [[CrossRef](#)]
- Landrigan, P.J.; Fuller, R.; Acosta, N.J.R. The Lancet Commission on pollution and health. *Lancet* **2018**, *391*, 462–512. [[CrossRef](#)]
- Xie, X.; Semanjski, I.; Gautama, S.; Tsiligianni, E.; Deligiannis, N.; Rajan, R.; Pasveer, F.; Philips, W. A Review of Urban Air Pollution Monitoring and Exposure Assessment Methods. *ISPRS Int. J. Geo. Inf.* **2017**, *6*, 389. [[CrossRef](#)]
- Díaz-Álvarez, E.A.; Lindig-Cisneros, R.; de la Barrera, E. Biomonitoring of atmospheric nitrogen deposition: Potential uses and limitations. *Conserv. Physiol.* **2018**, *6*, coy011. [[CrossRef](#)] [[PubMed](#)]
- Górecki, T.; Namieśnik, J. Passive sampling. *TrAC Trends Analyt. Chem.* **2002**, *21*, 276–291. [[CrossRef](#)]
- Díaz-Álvarez, E.A.; de la Barrera, E. Characterization of nitrogen deposition in a megalopolis by means of atmospheric biomonitoring. *Sci. Rep.* **2018**, *8*, 13569. [[CrossRef](#)]
- Díaz-Álvarez, E.A.; de la Barrera, E. Isotopic biomonitoring of anthropic carbon emissions in a megalopolis. *PeerJ* **2020**, *8*, e9283. [[CrossRef](#)] [[PubMed](#)]
- Lichtfouse, E.; Lichtfouse, M.; Jaffrézic, A. $\delta^{13}\text{C}$ Values of Grasses as a Novel Indicator of Pollution by Fossil-Fuel-Derived Greenhouse Gas CO₂ in Urban Areas. *Environ. Sci. Technol.* **2003**, *37*, 87–89. [[CrossRef](#)] [[PubMed](#)]
- Wang, W.; Pataki, D.E. Spatial patterns of plant isotope tracers in the Los Angeles urban region. *Landsc. Ecol.* **2010**, *25*, 35–52. [[CrossRef](#)]
- Wang, W.; Pataki, D.E. Drivers of spatial variability in urban plant and soil isotopic composition in the Los Angeles basin. *Plant Soil* **2011**, *350*, 323–338. [[CrossRef](#)]
- Redling, K.; Elliot, E.; Bain, D.; Sherwell, J. Highway contributions to reactive nitrogen deposition: Tracing the fate of vehicular NO_x using stable isotopes and plant biomonitoring. *Biogeochem.* **2013**, *116*, 261–274. [[CrossRef](#)]
- Sulzman, E.W. Stable Isotope Chemistry and Measurement: A Primer. In *Stable Isotopes in Ecology and Environmental Science*, 2nd ed.; Michener, R., Lajtha, K., Eds.; Blackwell Publishing Ltd.: Malden, MA, USA, 2007; pp. 1–121.
- Maricle, B.R.; Caudle, K.L.; Adler, P.B. Influence of water Availability on Photosynthesis, Water Potential, Leaf $\delta^{13}\text{C}$, and Phenology in Dominant C₄ Grasses In Kansas, USA. *Trans. Kans. Acad. Sci.* **2015**, *118*, 173–193. [[CrossRef](#)]
- Williams, D.G.; Gempko, V.; Fravolini, A.; Leavitt, S.W.; Wall, G.W.; Kimball, B.A.; Pinter, P.J., Jr.; LaMorte, R.; Ottman, M. Carbon isotope discrimination by Sorghum bicolor under CO₂ enrichment and drought. *New Phytol.* **2001**, *150*, 285–293. [[CrossRef](#)]
- Pataki, D.E.; Bowling, D.R.; Ehleringer, J.R. Seasonal cycle of carbon dioxide and its isotopic composition in an urban atmosphere: Anthropogenic and biogenic effects. *J. Geophys. Res.* **2003**, *108*, 4735. [[CrossRef](#)]
- Widory, D.; Javoy, M. The carbon isotope composition of atmospheric CO₂ in Paris. *Earth Planet. Sci. Lett.* **2003**, *215*, 289–298. [[CrossRef](#)]
- Semmens, C.; Ketler, R.; Schwendenmann, L.; Nesic, Z.; Christen, A. *Isotopic Composition of CO₂ in Gasoline, Diesel and Natural Gas Combustion Exhaust in Vancouver, BC, Canada*; Technical Report; The University of British Columbia: Columbia, BC, Canada, 2014; pp. 1–12.
- Molina-Freaner, F.; Gámez, R.C.; Tinoco-Ojanguren, C.; Castellanos, A.E. Vine species diversity across environmental gradients in northwestern México. *Biodivers. Conserv.* **2004**, *13*, 1853–1874. [[CrossRef](#)]
- Cox, J.R.; Martin-R., M.H.; Ibarra-F., F.A.; Fourie, J.H.; Rethman, N.F.G.; Wilcos, D.G. The influence of climate and soils on the distribution of four African grasses. *J. Range Manag.* **1988**, *41*, 127–139. [[CrossRef](#)]
- Ibarra-Flores, F.; Cox, J.R.; Martin, M.; Crawl, T.A.; Call, C.A. Predicting buffelgrass survival across a geographical and environmental gradient. *J. Range Manag.* **1995**, *48*, 53–59. [[CrossRef](#)]
- Tinoco-Ojanguren, C.; Reyes-Ortega, I.; Sánchez-Coronado, M.E.; Molina-Freaner, F.; Orozco-Segovia, A. Germination of an invasive *Cenchrus ciliaris* L. (buffel grass) population of the Sonoran Desert under various environmental conditions. *S. Afr. J. Bot.* **2016**, *104*, 112–117. [[CrossRef](#)]
- Arriaga, L.; Castellanos, A.E.; Moreno, E.; Alarcón, J. Potential ecological distribution of alien invasive species and risk assessment: A case study for buffel grass in arid regions of Mexico. *Conserv. Biol.* **2004**, *18*, 1504–1514. [[CrossRef](#)]
- de la Barrera, E. Recent invasion of buffel grass (*Cenchrus ciliaris*) of a natural protected area from the southern Sonoran Desert. *Rev. Mex. Biodivers.* **2008**, *79*, 385–392.
- Marshall, V.M.; Lewis, M.M.; Ostendorf, B. Buffel grass (*Cenchrus ciliaris*) as an invader and threat to biodiversity in arid environments: A review. *J. Arid Environ.* **2012**, *78*, 1–12. [[CrossRef](#)]
- Williams, D.G.; Baruch, Z. African grass invasion in the Americas: Ecosystem consequences and the role of ecophysiology. *Biol. Invasions* **2000**, *2*, 123–140. [[CrossRef](#)]

30. SEMARNAT. *Norma Oficial Mexicana NOM-156-SEMARNAT-2012, Establecimiento y Operación de Sistemas de Monitoreo de la Calidad del Aire*; Diario Oficial de la Federación: México D.F., Mexico, 2012.
31. INEGI. Cuéntame, Información por Entidad. 2020. Available online: <http://www.cuentame.inegi.org.mx/monografias/informacion/Son/Poblacion/default.aspx?tema=ME&e=26> (accessed on 23 March 2020).
32. INECC, Instituto Nacional de Ecología y Cambio Climático. *Informe Nacional de Calidad del Aire 2018, México*; Coordinación General de Contaminación y Salud Ambiental, Dirección de Investigación de Calidad del Aire y Contaminantes Climáticos: Ciudad de México, Mexico, 2019; pp. 1–395.
33. Ma, S.; He, F.; Tian, D.; Zou, D.; Yan, Z.; Yang, Y.; Zhou, T.; Huang, K.; Shen, H.; Fang, J. Variations and determinants of carbon content in plants: A global synthesis. *Biogeosciences* **2018**, *15*, 693–702. [[CrossRef](#)]
34. Castellanos-Villegas, A.E.; Yanes, G.; Valdez-Zamudio, D. Drought-tolerant exotic buffel-grass and desertification. In *Weeds Across Borders*; Tellman, B., Ed.; Proceedings of the North American Conference; The Arizona-Sonora Desert Museum: Tucson, Arizona, 2002; pp. 99–112.
35. Bhatt, R.K.; Baig, M.J.; Tiwari, H.S. Growth, biomass production, and assimilatory characters in *Cenchrus ciliaris* L. Under elevated CO₂ condition. *Photosynthetica* **2007**, *45*, 296–298. [[CrossRef](#)]
36. Williams, D.G.; Black, R.A. Response to drought for a native and introduced Hawaiian grass. *Oecologia* **1994**, *97*, 512–519. [[CrossRef](#)]
37. Ripley, B.S.; Abraham, T.I.; Osborne, C.P. Consequences of C₄ photosynthesis for the partitioning of growth: A test using C₃ and C₄ subspecies of *Alloterospis semialata* under nitrogen-limitation. *J. Exp. Bot.* **2007**, *59*, 1705–1714. [[CrossRef](#)]
38. Buchmann, N.; Brooks, J.R.; Rapp, K.D.; Ehleringer, J.R. Carbon isotope composition of C₄ grasses is influenced by light and water supply. *Plant Cell Environ.* **1996**, *19*, 392–402. [[CrossRef](#)]
39. Millar, N.; Urrea, A.; Kahmark, K.; Shcherbak, I.; Robertson, G.P.; Ortiz-Monasterio, I. Nitrous oxide (N₂O) flux responds exponentially to nitrogen fertilizer in irrigated wheat in the Yaqui Valley, Mexico. *Agric. Ecosyst. Environ.* **2018**, *261*, 125–132. [[CrossRef](#)]
40. Cogley, L.A.E.; Pataki, D.E. Vehicle emissions and fertilizer impact the leaf chemistry of urban trees in Salt Lake Valley, UT. *Environ. Pollut.* **2019**, *254*, 112984. [[CrossRef](#)] [[PubMed](#)]
41. Naus, S.; Röckmann, T.; Popa, M.E. The isotopic composition of CO in vehicle exhaust. *Atmospheric Environ.* **2018**, *177*, 132–142. [[CrossRef](#)]
42. Winter, K.; Troughton, J.H.; Card, K.A. δ¹³C values of grass species collected in the northern Sahara desert. *Oecologia* **1976**, *25*, 115–123. [[CrossRef](#)]
43. Börjesson, P.I.I. Emissions of CO₂ from biomass production and transportation in agriculture and forestry. *Energy Convers. Manag.* **1996**, *37*, 1235–1240. [[CrossRef](#)]
44. FAO. *Agriculture, Forestry and Other Land Use Emissions by Sources and Removals by Sinks. 1990–2011 Analysis*; FAO Statistics Division, Working Paper Series; FAO: Rome, Italy, 2014; pp. 1–89.
45. Liu, X.; Zhang, S.; Bae, J. The nexus of renewable energy-agriculture-environment in BRICS. *Appl. Energy* **2017**, *204*, 489–496. [[CrossRef](#)]
46. Gitz, V.; Ciais, P. Future Expansion Of Agriculture and Pasture Acts to Amplify Atmospheric CO₂ Levels in Response to Fossil-Fuel and Land-Use Change Emissions. *Clim. Change* **2004**, *67*, 161–184. [[CrossRef](#)]
47. Díaz-Álvarez, E.A.; Lindig-Cisneros, R.; de la Barrera, E. Responses to simulated nitrogen deposition by the neotropical epiphytic orchid *Laelia speciosa*. *PeerJ* **2015**, *3*, e1021. [[CrossRef](#)] [[PubMed](#)]
48. Harrison, A.F.; Schulze, E.D.; Gebauer, G.; Bruckner, G. Canopy Uptake and Utilization of Atmospheric Pollutant Nitrogen. In *Carbon and Nitrogen Cycling in European Forest Ecosystems*; Schulze, E.D., Ed.; Springer: Berlin, Germany, 2000; pp. 171–184.
49. Ammann, M.; Siegwolf, R.; Pichlmayer, F.; Suter, M.; Saurer, M.; Brunold, C. Estimating the uptake of traffic-derived NO₂ from ¹⁵N abundance in Norway spruce needles. *Oecologia* **1999**, *118*, 124–131. [[CrossRef](#)]
50. Vallano, D.; Sparks, J. Quantifying foliar uptake of gaseous nitrogen dioxide using enriched foliar δ¹⁵N values. *New Phytol.* **2008**, *177*, 946–955. [[CrossRef](#)]
51. Jung, K.; Gebauer, G.; Gehre, M.; Hofmann, D.; Weißflog, L.; Schüttrmann, G. Anthropogenic impacts on natural nitrogen isotope variations in pinus sylvestris stands in an industrially polluted area. *Environ. Pollut.* **1997**, *97*, 175–181. [[CrossRef](#)]
52. Craine, J.M.; Brookshire, E.N.J.; Cramer, M.D.; Hasselquist, N.J.; Koba, K.; Marin-Spiota, E.; Wang, L. Ecological interpretations of nitrogen isotope ratios of terrestrial plants and soils. *Plant Soil* **2015**, *396*, 1–26. [[CrossRef](#)]
53. Xu, Y.; Xiao, H.; Wu, D. Traffic-related dustfall and NO_x, but not NH₃, seriously affect nitrogen isotopic compositions in soil and plant tissues near the roadside. *Environ. Pollut.* **2019**, *249*, 655–665. [[CrossRef](#)] [[PubMed](#)]
54. Hoering, T. The isotopic composition of the ammonia and the nitrate ion in rain. *Geochim. Cosmochim. Acta* **1957**, *12*, 97–102. [[CrossRef](#)]
55. Gebauer, G.; Dietrich, P. Nitrogen isotope ratios in different compartments of a mixed stand of spruce, larch and beech trees and of understorey vegetation including fungi. *Isot. Environ. Health Stud.* **1993**, *29*, 35–44. [[CrossRef](#)]
56. Gebauer, G.; Giesemann, A.; Schulze, E.-D.; Jäger, H.-J. Isotope ratios and concentrations of sulfur and nitrogen in needles and soils of *Picea abies* stands as influenced by atmospheric deposition of sulfur and nitrogen compounds. *Plant Soil* **1994**, *164*, 267–281. [[CrossRef](#)]
57. Aber, J.; McDowell, W.; Nadelhoffer, K.; Magill, A.; Berntson, G.; Kamakea, M.; McNulty, S.; Currie, W.; Rustad, L.; Fernandez, I. Nitrogen saturation in temperate forest ecosystems. *Bioscience* **1998**, *48*, 921–934. [[CrossRef](#)]

58. Stewart, G.; Aidar, M.P.; Joly, C.A.; Schmidt, S. Impact of point source pollution on nitrogen isotope signatures ($\delta^{15}\text{N}$) of vegetation in SE Brazil. *Oecologia* **2002**, *131*, 468–472. [CrossRef] [PubMed]
59. Eugster, W.; Haeni, M. Nutrients or Pollutants? Nitrogen Deposition to European Forests. In *Climate Change, Air Pollution and Global Challenges—Understanding and Perspectives from Forest Research*; Matyssek, R., Clarke, N., Cudlin, P., Mikkelsen, T.N., Tuovinen, J.-P., Wieser, G., Paoletti, E., Eds.; Elsevier: Amsterdam, The Netherlands, 2013; Volume 13, pp. 37–56.
60. Zech, M.; Bimüller, C.; Hemp, A.; Samimi, C.; Broesike, C.; Hörold, C.; Zech, W. Human and climate impact on ^{15}N natural abundance of plants and soils in high-mountain ecosystems: A short review and two examples from the Eastern Pamirs and Mt. Kilimanjaro. *Isot. Environ. Health Stud.* **2011**, *47*, 286–296. [CrossRef] [PubMed]
61. Díaz-Álvarez, E.A.; Reyes-García, C.; de la Barrera, E. A $\delta^{15}\text{N}$ assessment of nitrogen deposition for the endangered epiphytic orchid *Laelia speciosa* from a city and an oak forest in Mexico. *J. Plant Res.* **2016**, *129*, 863–872. [CrossRef] [PubMed]
62. Kenkel, J.A.; Sisk, T.D.; Hultine, K.R.; Sesnie, S.E.; Bowker, M.A.; Johnson, N.C. Indicators of vehicular emission inputs into semi-arid roadside ecosystems. *J. Arid Environ.* **2016**, *134*, 150–159. [CrossRef]
63. Ahrens, T.; Harrison, J.; Berman, M.; Jewett, P.; Matson, P.A. Nitrogen in the Yaqui Valley: Sources, transfers, and consequences. In *Seeds of Sustainability, Lessons from the Birthplace of the Green Revolution*; Matson, P.A., Ed.; Island Press: Washington, DC, USA, 2012; pp. 171–195.
64. Parra Suárez, S.; Peiffer, S.; Gebauer, G. Origin and fate of nitrate runoff in an agricultural catchment: Haeon, South Korea—Comparison of two extremely different monsoon seasons. *Sci. Total Environ.* **2019**, *648*, 66–79. [CrossRef] [PubMed]
65. Hogberg, P.; Johannisson, C. ^{15}N abundance of forests is correlated with losses of nitrogen. *Plant Soil* **1993**, *157*, 147–150. [CrossRef]
66. Pardo, L.H.; Templer, P.H.; Goodale, C.L.; Duke, S.; Groffman, P.M.; Adams, M.B.; Boeckx, P.; Boggs, J.; Campbell, J.; Colman, B.; et al. Regional assessment of N saturation using foliar and root. *Biogeochemistry* **2006**, *80*, 143–171. [CrossRef]
67. Emmett, B.A. Nitrogen saturation of terrestrial ecosystems: Some recent findings and their implications for our conceptual framework. *Water Air Soil Pollut. Focus* **2007**, *7*, 99–109. [CrossRef]
68. Panek, J.A.; Matson, P.A.; Ortiz-Monasterio, I.; Brooks, P. Distinguishing nitrification and denitrification sources of N_2O in a Mexican wheat system using ^{15}N . *Ecol. Appl.* **2000**, *10*, 506–514.
69. Townsend-Small, A.; Pataki, D.E.; Czimeczik, C.I.; Tyler, S.C. Nitrous oxide emissions and isotopic composition in urban and agricultural systems in southern California. *J. Geophys. Res.* **2011**, *116*, G01013. [CrossRef]
70. Felix, J.D.; Elliott, E.M. Isotopic composition of passively collected nitrogen dioxide emissions: Vehicle, soil and livestock source signatures. *Atmos. Environ.* **2014**, *92*, 359–366. [CrossRef]
71. Felix, J.D.; Elliot, E.M.; Gay, D.A. Spatial and temporal patterns of nitrogen isotopic composition of ammonia at US ammonia monitoring network sites. *Atmos. Environ.* **2017**, *150*, 434–442. [CrossRef]
72. Ahrens, T.D.; Beman, J.M.; Harrison, J.A.; Jewett, P.K.; Matson, P.A. A synthesis of nitrogen transformations and transfers from land to the sea in the Yaqui Valley agricultural region of northwest Mexico. *Water Resour. Res.* **2008**, *44*, W00A05. [CrossRef]
73. Billy, C.; Billen, G.; Sebilo, M.; Birgand, F.; Tournebize, J. Nitrogen isotopic composition of leached nitrate and soil organic matter as an indicator of denitrification in a sloping drained agricultural plot and adjacent uncultivated riparian buffer strips. *Soil Biol. Biochem.* **2012**, *42*, 108–117. [CrossRef]
74. Payne, R.J.; Dise, N.B.; Field, C.D.; Dore, A.J.; Caporn, S.J.; Stevens, C.J. Nitrogen deposition and plant biodiversity: Past, present, and future. *Front. Ecol. Environ.* **2017**, *15*, 431–436. [CrossRef]
75. Matson, P.; Luers, A.M.; Seto, K.C.; Naylor, R.L.; Ortiz-Monasterio, I. People, Land Use, and Environment in the Yaqui Valley, Sonora, Mexico. In *Land Use, and Environment: Research Directions*; Entwisle, B., Stern, P.C., Eds.; National Research Council: Washington, DC, USA, 2005; pp. 238–264.
76. Díaz-Álvarez, E.A.; de la Barrera, E.; Barrios-Hernández, E.Y.; Arróniz-Crespo, M. Morphophysiological screening of potential organisms for biomonitoring nitrogen deposition. *Ecol. Indic.* **2020**, *108*, 105729. [CrossRef]
77. Forman, R.T.T.; Godron, M. *Landscape Ecology*; Wiley: New York, NY, USA, 1986; pp. 1–619.
78. Adams, A.S.; Vázquez, J.L.M.; Ley, A.N.L. Agricultura y manejo sustentable del acuífero de la Costa de Hermosillo. *Rec. Soc.* **2012**, *3*, 155–179.
79. Research on Forest Climate Change: Potential Effects of Global Warming on Forests and Plant Climate Relationships in Western North America and Mexico. Available online: <http://charcoal.cnre.vt.edu/climate/> (accessed on 9 April 2021).
80. Instituto Nacional de Geografía y Estadística. Conjunto de datos de perfiles de suelos. Escala 1:250,000. Serie II (Continuo nacional), Mexico. Available online: <https://www.inegi.org.mx/temas/edafologia/#Descargas> (accessed on 9 April 2021).
81. Ehleringer, J.R.; Osmond, B.O. Stable isotopes. In *Plant Physiological Ecology*; Pearcy, R.W., Ehleringer, J.R., Mooney, H.A., Rundel, P.W., Eds.; Springer: Dordrecht, The Netherlands; Berlin, Germany, 1989; pp. 281–300.
82. Evans, R. Physiological mechanisms influencing plant nitrogen isotope composition. *Trends Plant Sci.* **2001**, *6*, 121–126. [CrossRef]
83. Metsalu, T.; Vilo, J. Clustvis: A web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap. *Nucleic Acids Res.* **2015**, *43*, W566–W570. [CrossRef]
84. Cressie, N. Spatial prediction and ordinary Kriging. *Math. Geol.* **1988**, *20*, 405–421. [CrossRef]