

Extended Abstract

A New Approach to the Synthesis of Functionalized Bicyclo[3.2.1]octanes [†]

Arif Idris Ismiev ¹, Gunay Zakhid Mamedova ¹, Victor V. Dotsenko ^{2,3,4,*}, Nikolai A. Aksenov ³, Inna V. Aksenova ³ and Abel Mamedali Magerramov ¹

¹ Department of Chemistry, Baku State University, Akademik Zahid Xəlilov küçəsi-23, Baku 1148, Azerbaijan; arif_ismiev@mail.ru (A.I.I.); gunka479@mail.ru (G.Z.M.); rector@bsu.az (A.M.M.)

² Department of Chemistry and High Technologies, Kuban State University, 149 Stavropolskaya Str., 350040 Krasnodar, Russia

³ Department of Chemistry, North Caucasus Federal University, 1a Pushkina Str., 355009 Stavropol, Russia; radioanimation@rambler.ru (N.A.A.); inna-aksenova00@rambler.ru (I.V.A.)

⁴ ChemEx Lab, Vladimir Dal' Lugansk National University, 20A Molodezhny, 91034 Lugansk, Russia

* Correspondence: victor_dotsenko_@mail.ru


[†] Presented at the 22nd International Electronic Conference on Synthetic Organic Chemistry, 15 November–15 December 2018; Available Online: <https://sciforum.net/conference/ecsoc-22>.

Published: 14 November 2018

Abstract: The reaction of furfural with secondary amines and isopropyl cyanoacetate leads to the formation of the previously unknown diisopropyl 8-(dialkylamino)-2,4-dicyano-3-(2-furyl)-6-oxobicyclo[3.2.1]octane-2,4-dicarboxylates. The structures of the products were confirmed by NMR and X-ray analysis.


Keywords: furfural; isopropyl cyanoacetate; Stenhouse salt; Nazarov reaction; carbocyclization; bicyclo[3.2.1]octane

Recently, a number of papers [1–6] dealing with the development of various approaches to the synthesis of bicyclo[3.2.1]octanes have appeared (for reviews, see [7–9]). First, this bicyclic system is of interest due to its biological activity—inhibitors of dopamine and serotonin transfer [10] were found among bicyclo[3.2.1]octanes. They can also be used in the synthesis of tricyclic terpenoids [11]. The bicyclo[3.2.1]octane motif is the structural core unit of various natural terpene compounds and alkaloids [8]. In continuation of our studies in the chemistry of active methylene nitriles [12–16], we would like to propose a new approach to the synthesis of bicyclo[3.2.1]octanes based on an elegant cascade reaction between furfural and isopropyl cyanoacetate in the presence of secondary amines. We found that the reaction of furfural with secondary amines and cyanoacetic ester leads to the formation of bicyclooctanes **1a,b** (Scheme 1) in moderate yields (37%–45%).



Scheme 1. Synthesis of compounds **1**

The proposed mechanism firstly involves the formation of the deprotonated Stenhouse salt **2** from a secondary amine and furfural (Scheme 2). It should be noted that when furfural and cyanoacetic ester were put into the reaction simultaneously, the reaction led to the formation of the expected Knoevenagel products, 3-(2-furyl)-2-cyanoacrylates. We suppose that the intermediate **2** further undergoes spontaneous Nazarov-type 4π -electrocyclization [17] to give 2,3-diaminocyclopentenones **3**. The formation of **3** from furfural and amines is well known and described in recent literature (e.g., [18–25], see also the review by Piancatelli et al. [26]). When furfural and isopropyl cyanoacetate were added to the reaction mixture, 3-(2-furyl)-2-cyanoacrylate **4** and Michael adduct **5** were formed sequentially. Under basic conditions, compound **5** undergoes the Michael addition to cyclopentenone **3**, followed by carbocyclization of adduct **6**, probably through the nucleophilic substitution of a secondary amine (Scheme 2).



Scheme 2. Probable mechanism of the formation of **1**.

The structures of compounds **1a,b** were confirmed by means of IR and NMR $^1\text{H}/^{13}\text{C}$ data, as well as X-ray diffraction analysis for compound **1a** (Figure 1).

Conclusions

In summary, we have discovered a new cascade reaction, opening a new opportunity for the synthesis of functionalized bicyclo[3.2.1]octanes from available reagents under one-pot conditions. The mechanistic details, scope, and limitations of the reaction will be reported elsewhere.

Experimental

IR spectra were recorded on a Varian 3600 FT-IR Excalibur Series spectrometer in KBr pellets. NMR spectra were recorded on a Bruker AC-300 (300 MHz for ^1H , 75 MHz for ^{13}C) in CDCl_3 . The selected experimental procedure (synthesis of **1a**) is given. Microanalysis was obtained using a Carlo Erba 1106 analyzer.

A mixture of 0.83 mL (0.01 mol) of furfural and 0.02 mol of Et_2NH in 96% EtOH (10 mL) was stirred for 2 h. Then, a mixture of 0.83 mL (0.01 mol) of furfural and 2.51 mL (0.02 mol) of isopropyl cyanoacetate was added to the solution under vigorous stirring. The reaction mass was maintained for 24 h, and the precipitated crystals were filtered off.

Beige crystals, yield 37%, mp 154 °C (EtOH). IR, ν , cm^{-1} : 2240 ($\text{C}\equiv\text{N}$), 1750 ($\text{C}=\text{O}$), 1654 ($\text{CO}_2\text{Pr-i}$). NMR ^1H (300 MHz, CDCl_3), δ , ppm: 1.02 t (6H, 2 CH_2CH_3 , 3J 7.0 Hz), 1.21–1.30 m (12H, 2 $\text{CH}(\text{CH}_3)_2$),

2.53–2.54 m (2H, H-7), 2.69 q (4H, 2 CH_2CH_3 , 3J 7.0 Hz), 2.92 br.s (1H, H-8), 3.12–3.14 m (1H, H-1), 3.97 s (1H, H-5), 4.29 s (1H, H-3), 5.01–5.12 m (2H, 2 $\text{CH}(\text{CH}_3)_2$), 6.34 dd (1H, H-4_{furyl}, $^3J_{\text{H}(4)-\text{H}(3)}$ 3.1 Hz, $^3J_{\text{H}(4)-\text{H}(5)}$ 1.7 Hz), 6.67 d (1H, H-3_{furyl}, $^3J_{\text{H}(4)-\text{H}(3)}$ 3.1 Hz), 7.35–7.36 m (1H, H-5_{furyl}). NMR ^{13}C (75 MHz, CDCl_3), δ , ppm: 11.1 (2 CH_2CH_3), 21.14 ($\text{CH}(\text{CH}_3)_2$), 21.18 ($\text{CH}(\text{CH}_3)_2$), 21.23 ($\text{CH}(\text{CH}_3)_2$), 21.27 ($\text{CH}(\text{CH}_3)_2$), 35.6 (C-3), 38.0 (C-1), 42.1 (2 CH_2CH_3), 42.5 (C-8), 50.4 (C-4 or C-2), 51.8 (C-2 or C-4), 54.5 (C-7), 63.2 (C-5), 72.5 ($\text{CH}(\text{CH}_3)_2$), 72.9 ($\text{CH}(\text{CH}_3)_2$), 110.79 (C-4_{furyl}), 110.81 (C-3_{furyl}), 115.3 (C \equiv N), 116.2 (C \equiv N), 142.8 (C-5_{furyl}), 146.6 (C-1_{furyl}), 163.4 ($\text{CO}_2\text{Pr-}i$), 165.1 ($\text{CO}_2\text{Pr-}i$), 206.9 (C=O). Found, %: C 64.62; H 6.97; N 8.63. $\text{C}_{26}\text{H}_{33}\text{N}_3\text{O}_6$. Calculated, %: C 64.58; H 6.88; N 8.69.




Figure 1. The structure of diisopropyl 2,4-dicyano-8-(diethylamino)-3-(2-furyl)-6-oxobicyclo[3.2.1]octane-2,4-dicarboxylate (**1a**) (X-ray data).

References

1. Lefranc, A.; Gremaud, L.; Alexakis, A. Construction of Bicyclo[3.2.1]octanes with Four Stereogenic Centers by Organocatalytic Domino Michael/Aldol Reaction. *Org. Lett.* **2014**, *16*, 5242–5245. doi:10.1021/o1502171h
2. Rueping, M.; Kuenkel, A.; Tato, F.; Bats, J.W. Asymmetric organocatalytic domino Michael/aldol reactions: enantioselective synthesis of chiral cycloheptanones, tetrahydrochromenones, and polyfunctionalized bicyclo[3.2.1]octanes. *Angew. Chem. Int. Ed.* **2009**, *48*, 3699–3702. doi:10.1002/anie.200900754
3. Tan, B.; Lu, Y.; Zeng, X.; Chua, P.J.; Zhong, G. *Org. Lett.* **2010**, *12*, 2682–2685. doi: 10.1021/o11007795
4. Zhao, J.; Yang, S.; Xie, X.; Li, X.; Liu, Y.J. Facile Domino Access to Chiral Bicyclo[3.2.1]octanes and Discovery of a New Catalytic Activation Mode. *Org. Chem.* **2018**, *83*, 1287–1297. doi:10.1021/acs.joc.7b02816
5. He, C.; Bai, Z.; Hu, J.; Wang, B.; Xie, H.; Yu, L.; Ding, H. A divergent [5+2] cascade approach to bicyclo[3.2.1]octanes: facile synthesis of ent-kaurene and cedrene-type skeletons. *Chem. Commun.* **2017**, *53*, 8435–8438. doi:10.1039/c7cc04292b
6. Takatori, K.; Ota, S.; Tendo, K.; Matsunaga, K.; Nagasawa, K.; Watanabe, S.; Kishida, A.; Kogen, H.; Nagaoka, H. Synthesis of Methylenebicyclo[3.2.1]octanol by a Sm(II)-Induced 1,2-Rearrangement Reaction with Ring Expansion of Methylenebicyclo[4.2.0]octanone. *Org. Lett.* **2017**, *19*, 3763–3766. doi:10.1021/acs.orglett.7b01604
7. Filippini, M.H.; Rodriguez, J. Synthesis of Functionalized Bicyclo[3.2.1]octanes and Their Multiple Uses in Organic Chemistry. *Chem. Rev.* **1999**, *99*, 27–76. doi:10.1021/cr970029u

8. Presset, M.; Coquerel, Y.; Rodriguez, J. Syntheses and Applications of Functionalized Bicyclo[3.2.1]octanes: Thirteen Years of Progress. *Chem. Rev.* **2013**, *113*, 525–595. doi: 10.1021/cr200364p
9. Presset, M.; Coquerel, Y.; Rodriguez, J. Single-Step Metallocatalyzed and Organocatalyzed Enantioselective Construction of Bicyclo[3.2.1]octanes. *Chem. Cat. Chem.* **2012**, *4*, 172–174. doi:10.1002/cctc.201100357
10. Meltzer, P.C.; Blundell, P.; Yong, Y.F.; Chen, Z.; George, C.; Gonzalez, M.D.; Madras, B.K. 2-Carbomethoxy-3-aryl-8-bicyclo[3.2.1]octanes: Potent Non-Nitrogen Inhibitors of Monoamine Transporters. *J. Med. Chem.* **2000**, *43*, 2982–2991. doi:10.1021/jm000191g.
11. Peter, C.; Geoffroy, P.; Miesch, M. Intramolecular Morita–Baylis–Hillman reaction as a strategy for the construction of tricyclic sesquiterpene cores. *Org. Biomol. Chem.* **2018**, *16*, 1381–1389. doi:10.1039/c7ob03124f
12. Dotsenko, V.V.; Ismiev, A.I.; Khrustaleva, A.N.; Frolov, K.A.; Krivokolysko, S.G.; Chigorina, E.A.; Snizhko, A.P.; Gromenko, V.M.; Bushmarinov, I.S.; Askerov, R.K.; et al. Synthesis, structure, and reactions of (4-aryl-3-cyano-6-oxopiperidin-2-ylidene)malononitriles. *Chem. Heterocycl. Compds.* **2016**, *52*, 473–483. doi:10.1007/s10593-016-1918-3
13. Hajiyeva, K.; Ismiev, A.; Franz, M.; Schmidtman, M.; Martens, J.; Maharramov, A. Simple one-pot multicomponent approach to polyfunctionalized 2-amino-1-methyl-6-(methylthio)-5-nitro-4-aryl-1,4-dihydropyridine-3-carbonitriles. *Synth. Commun.* **2017**, *47*, 2031–2035. doi:10.1080/00397911.2017.1359845
14. Dotsenko, V.V.; Frolov, K.A.; Pekhtereva, T.M.; Papaianina, O.S.; Suykov, S.Yu., Krivokolysko, S.G. Design and Synthesis of Pyrido[2,1-b][1,3,5]thiadiazine Library via Uncatalyzed Mannich-Type Reaction. *ACS Comb. Sci.* **2014**, *16*, 543–550. doi:10.1021/co5000807
15. Dotsenko, V.V.; Krivokolysko, S.G.; Litvinov, V.P. 1-(Cyanoacetyl)-3,5-dimethylpyrazole as Active Methylene Compound in Hantzsch-type Pyridine Synthesis: A Convenient and Highly Effective Approach to 3,5-Dicyano-4-(het)aryl-6-oxo-1,4,5,6-tetrahydropyridine-2-thiolates. *Monatsh. Chem.* **2007**, *138*, 607–613. doi:10.1007/s00706-007-0664-8
16. Dotsenko, V.V.; Krivokolysko, S.G.; Polovinko, V.V.; Litvinov, V.P. On the regioselectivity of the reaction of cyanothioacetamide with 2-acetylcyclohexanone, 2-acetylcyclopentanone, and 2-acetyl-1-(morpholin-4-yl)-1-cycloalkenes. *Chem. Heterocycl. Compds.* **2012**, *48*, 309–319. doi:10.1007/s10593-012-0991-5
17. Tius, M.A. Some New Nazarov Chemistry. *Eur. J. Org. Chem.* **2005**, *2005*, 2193–2206. doi:10.1002/ejoc.200500005
18. Griffiths, K.; Gallop, C.W.D.; Abdul-Sada, A.; Vargas, A.; Navarro, O.; Kostakis, G.E. Heteronuclear 3 d/Dy^{III} Coordination Clusters as Catalysts in a Domino Reaction. *Chem. Eur. J.* **2015**, *21*, 6358–6361. doi:10.1002/chem.201500505
19. Ramesh, D.; Srikanth Reddy, T.; Narasimhulu, M.; Rajaram, S.; Suryakiran, N.; Chinni Mahesh, K.; Venkateswarlu, Y. Efficient and Rapid Stereoselective Synthesis of trans-4,5-Diaminocyclopent-2-enones by Acidic Ionic Liquid under Solvent-free Conditions. *Chem. Lett.* **2009**, *38*, 586–587. doi:10.1246/cl.2009.586.
20. Procopio, A.; Costanzo, P.; Curini, M.; Nardi, M.; Oliverio, M.; Sindona, G. Erbium(III) Chloride in Ethyl Lactate as a Smart Ecofriendly System for Efficient and Rapid Stereoselective Synthesis of trans-4,5-Diaminocyclopent-2-enones. *ACS Sust. Chem. Eng.* **2013**, *1*, 541–544. doi:10.1021/sc4000219
21. Nardi, M.; Costanzo, P.; De Nino, A.; Di Gioia, M.L.; Olivito, F.; Sindona, G.; Procopio, A. Water excellent solvent for the synthesis of bifunctionalized cyclopentenones from furfural. *Green Chem.* **2017**, *19*, 5403–5411. doi:10.1039/C7GC02303K
22. Griffiths, K.; Kumar, P.; Mattock, J.D.; Abdul-Sada, A.; Pitak, M.B.; Coles, S.J.; Navarro, O.; Vargas, A.; Kostakis, G.E. Efficient Ni^{II}/Ln^{III}2 Electrocyclization Catalysts for the Synthesis of trans-4,5-Diaminocyclopent-2-enones from 2-Furaldehyde and Primary or Secondary Amines. *Inorg. Chem.* **2016**, *55*, 6988–6994. doi:10.1021/acs.inorgchem.6b00720
23. Liu, J.; Yu, J.; Zhu, M.; Li, J.; Zheng, X.; Wang, L. Novel Role of p-Toluenesulfonamide in the Preparation of 4,5-Diaminocyclopent-2-enones. *Synthesis* **2013**, *45*, 2165–2170. doi:10.1055/s-0033-1339180
24. Estevão, M.S.; Afonso, C.A.M. Synthesis of trans-4,5-diaminocyclopent-2-enones from furfural catalyzed by Er(III) immobilized on silica. *Tetrahedron Lett.* **2017**, *58*, 302–304. doi:10.1016/j.tetlet.2016.12.002

25. Li, S.-W.; Batey, R.A. Mild lanthanide(III) catalyzed formation of 4,5-diaminocyclopent-2-enones from 2-furaldehyde and secondary amines: A domino condensation/ring-opening/electrocyclization process. *Chem. Commun.* **2007**, 3759–3761. DOI: 10.1039/b709337n
26. Piancatelli, G.; D'Auria, M.; D'Onofrio, F. Synthesis of 1,4-Dicarbonyl Compounds and Cyclopentenones from Furans. *Synthesis* **1994**, 1994, 867–889. doi:10.1055/s-1994-25591



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).