








Article

Pregestational Stress Representing a Maternal Depression Model and Prenatally Applied Antidepressant Mirtazapine Modulate Hippocampal Excitability in Offspring

Lucia Dubiel-Hoppanova ^{1,†} , Alzbeta Filipova ^{1,†} , Stanislava Bukatova ², Katarina Ondacova ¹ ,
Matus Tomko ¹ , Bohumila Jurkovicova-Tarabova ¹ , Michal Dubovicky ², Eliyahu Dremencov ^{1,*} 
and Lubica Lacinova ^{1,*} 

- ¹ Centre of Biosciences, Institute of Molecular Physiology and Genetics, Slovak Academy of Sciences, 840 05 Bratislava, Slovakia; lucia.hoppanova@savba.sk (L.D.-H.); alzbeta.idunkova@savba.sk (A.F.); katarina.ondacova@savba.sk (K.O.); matus.tomko@savba.sk (M.T.); bohumila.tarabova@savba.sk (B.J.-T.)
- ² Centre of Experimental Medicine, Institute of Experimental Pharmacology and Toxicology, Slovak Academy of Sciences, 841 04 Bratislava, Slovakia; stanislava.bukatova@savba.sk (S.B.); michal.dubovicky@savba.sk (M.D.)
- * Correspondence: eliyahu.dremencov@savba.sk (E.D.); lubica.lacinova@savba.sk (L.L.)
- † These authors contributed equally to this work.

Abstract: Maternal depression negatively affects the neurodevelopment of offspring, but its pharmacological treatment during gestation remains controversial. This study reports the consequences of maternal depression and/or prenatal antidepressant treatment with mirtazapine on offspring early neurodevelopment via an animal model of maternal depression induced by pregestational chronic unpredictable stress (CUS). Offspring from four groups were studied: nonstressed vehicle-treated dams, nonstressed mirtazapine-treated dams, stressed vehicle-treated dams, and stressed mirtazapine-treated dams. The hippocampal excitability of offspring was examined in primary hippocampal cultures established on the first postnatal day, reflecting mostly prenatal development, and in hippocampal slices prepared on postnatal days 11–13, reflecting an early postnatal development. The pregestational CUS modeling of maternal depression moderately suppressed offspring hippocampal excitability in primary cultures but facilitated it in slices. Mirtazapine administered to CUS-exposed dams partly rectified the changes observed in primary cultures of pups from untreated dams and, more prominently, in slices. Mirtazapine itself negatively affected the hippocampal excitability of nonstressed dam offspring in primary culture, and this effect was diminished in slices. Since altered hippocampal neurotransmission might be responsible, at least in part, for the neuropsychopathologies frequently observed in the offspring of depressed mothers, and mirtazapine was able to partly relieve such changes, this treatment may be also beneficial during the prenatal and perinatal periods.

Keywords: pregestational stress; maternal depression; prenatal antidepressant treatment; hippocampal excitability



Academic Editor: Nilesh Patel

Received: 14 February 2025

Revised: 6 March 2025

Accepted: 22 March 2025

Published: 31 March 2025

Citation: Dubiel-Hoppanova, L.; Filipova, A.; Bukatova, S.; Ondacova, K.; Tomko, M.; Jurkovicova-Tarabova, B.; Dubovicky, M.; Dremencov, E.; Lacinova, L. Pregestational Stress Representing a Maternal Depression Model and Prenatally Applied Antidepressant Mirtazapine Modulate Hippocampal Excitability in Offspring. *Sci. Pharm.* **2025**, *93*, 17. <https://doi.org/10.3390/scipharm93020017>

Copyright: © 2025 by the authors. Published by MDPI on behalf of the Österreichische Pharmazeutische Gesellschaft. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Maternal depression is a general term referring to a spectrum of depressive conditions that typically occur during pregnancy and up to 12 months postpartum [1]. It is also a critical period of fetal and child development, and maternal disorders can have far-reaching impacts on offspring [2,3]. Maternal depression is a known risk factor for preterm birth, low

birth weight, and intrauterine growth restriction [4,5]. These complications are reported to be the leading cause of death in children under five years of age [6,7].

Many studies have examined the effects of prenatal maternal stress, anxiety, and depression on child development, impacting cognitive, behavioral, social, mood, language, and attachment outcomes [8–10]. Children of depressed mothers also face higher rates of cognitive, social, and mood disorders later in life [11,12]. Distinguishing the effects of untreated prenatal depression from those of other factors is challenging. Animal models, with their controlled environments, help us better understand the adverse effects of maternal stress and depression on offspring development.

The most frequently used antidepressant drugs are the second-generation antidepressants such as selective serotonin (5-HT) reuptake inhibitors (SSRIs) [13,14]. However, concerns have been raised about their use during pregnancy and the postpartum period because SSRIs easily cross the placenta and are present in breast milk, and fetuses and infants may have difficulty metabolizing SSRIs efficiently [15–17]. SSRIs may inadvertently affect serotonin signaling in offspring, leading to doubts about whether prescribing SSRIs to pregnant and postpartum women is safe for offspring. Several studies suggest that prenatal SSRI exposure has developmental consequences (enhancement of anxiety states) in both rats and human offspring [18,19].

Mirtazapine, a third-generation antidepressant, according to a recent meta-study, belongs to a small group of the most effective antidepressants [20] and is recommended for use during pregnancy. Mirtazapine is an antagonist of α_2 -adrenergic inhibitory, 5-HT₂, and 5-HT₃ receptors [21]. However, insufficient data are available to support its safe use during pregnancy [22].

In this study, we focused on the possible effects of maternal depression and its pharmacological treatment during gestation on the neuronal development and behavior of offspring in an early stage of postnatal development (up to 2 weeks). As it is involved in learning memory, cognitive function, and emotion regulation, the hippocampus is among the key brain regions affected by the pathogenesis of depression [23–25]. Therefore, we focused on possible alterations in hippocampal excitability.

As an animal model of maternal depression, we have used pregestational exposure to chronic unpredictable stress (CUS) [17,26,27]. We had previously reported that the CUS dams showed behavioral features resembling the symptoms of depression, as well as impaired hippocampal neurogenesis. Furthermore, we have shown that the behavioral and neuroanatomical abnormalities in the CUS dams were reversed by chronic antidepressant treatment [28]. CUS dams therefore possess constructive and predictive validity as a model of prenatal depression. In this work, we tested whether the previously observed alterations are reflected in altered hippocampal excitability. We compared the hippocampal excitability of offspring from four experimental groups: (i) dams not exposed to CUS and treated with vehicle; (ii) dams not exposed to CUS and treated with mirtazapine; (iii) dams exposed to CUS and treated with vehicle; and (iv) dams exposed to CUS and treated with mirtazapine.

The primary goal of our study was to test the hypothesis that the maternal stress and/or alteration in maternal monoamine neurotransmission during gestation affect the intrinsic properties of the embryonal glutamate neurons, or their progenitors. This alteration in the intrinsic properties of the embryonal neurons may subsequently affect their maturation and functioning in *in vivo* or *in vitro* conditions.

2. Results

2.1. Effects of Pregestational Stress and Prenatal Mirtazapine on the Electrophysiological Characteristics of Cultivated Pyramidal Neurons Isolated from Newborn Offspring Hippocampi

The resting membrane potentials (V_{rest}) and input resistance (R_{inp}) reflect the expression of voltage-gated ion channels in respective cells. V_{rest} and R_{inp} of the cultivated neurons isolated from the hippocampi of the newborn offspring of the dams exposed to pregestational CUS, prenatal mirtazapine treatment, or their combination are shown in Figure 1A. The Two-way ANOVA revealed a significant effect of CUS and perinatal mirtazapine on V_{rest} ($F_{1,280} = 38.85, p < 0.001$, and $F_{1,280} = 4.04, p = 0.045$, respectively) and R_{inp} ($F_{1,279} = 11.57, p < 0.001$, and $F_{1,279} = 10.02, p = 0.002$, respectively). The interaction between these factors for V_{rest} and R_{inp} was not statistically significant.

Observed differences were diminished during 11–13 days of postnatal development, as shown in Figure 1C,D. According to the two-way ANOVA results, maternal stress and mirtazapine treatment did not significantly affect V_{rest} or R_{inp} . The interactions between these factors were also not statistically significant.




Figure 1. Resting membrane potential (A,C) and input resistance (B,D) of cultivated neurons (A,B) and neurons in slices (C,D) isolated from the hippocampi of offspring of dams that experienced pregestational chronic unpredictable stress (CUS), prenatal mirtazapine treatment, or their combination. n (the number of individual cells/number of litters in the group) = 55/5 in NO CUS+VEH, 75/5 in NO CUS+MIR, 82/5 in CUS+VEH and 69/5 CUS+MIR in primary cultures, 54/5 in NO CUS+VEH, 63/5 in NO CUS+MIR, 38/5 in CUS+VEH and 39/5 CUS+MIR in acute slices; * $p < 0.05$, and *** $p < 0.001$, Bonferroni post hoc test.

The waveform of a single action potential (AP) activated by a brief depolarizing current pulse was characterized by the threshold (Figure 2A) and peak (Figure 2B) voltages, latency time (Figure 2C), width of action potential (Figure 2D), rise time (Figure 2E), and minimal current needed to evoke a single action potential (Figure 2F). CUS depressed excitability by increasing the voltage threshold for AP generation. Treatment by mirtazapine facilitated it by decreasing both the voltage threshold and minimal current amplitude needed for an activation of an AP.

For the monitored parameters, a significant effect of CUS on the peak voltage of the AP ($F_{1,278} = 9.35, p = 0.002$; two-way ANOVA) and its rise time ($F_{1,278} = 9.67, p < 0.002$; two-way ANOVA) was detected; a significant effect of maternal mirtazapine treatment on the threshold ($F_{1,278} = 7.99, p = 0.005$; two-way ANOVA) and peak ($F_{1,278} = 11.28, p < 0.001$; two-way ANOVA) voltages and latency times was detected ($F_{1,278} = 4.13, p = 0.04$; two-way ANOVA); and a significant interaction between these factors concerning the voltage threshold ($F_{1,278} = 49.59, p < 0.001$; two-way ANOVA), rise time ($F_{1,278} = 13.27, p < 0.001$;

two-way ANOVA), and the minimal current needed to induce a single action potential was detected ($F_{1,278} = 14.23, p < 0.001$; two-way ANOVA).




Figure 2. Single action potential threshold (A) and peak (B) voltages, latency time (C), width of action potential (D), rise time (E), and the minimal current needed to induce a single action potential (F) for cultivated neurons isolated from the hippocampi of newborn offspring of dams that experienced pregestational chronic unpredictable stress (CUS), prenatal mirtazapine treatment, or their combination. n (the number of individual cells/number of litters in the group) = 55/5 in NO CUS+VEH, 75/5 in NO CUS+MIR, 81/5 in CUS+VEH, and 68/5 in CUS+MIR; * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$, Bonferroni post hoc test.

The parameters characterizing the AP series (three or more APs generated during a depolarizing current pulse) activated by the second suprathreshold current pulse were analyzed in the same experimental groups are shown in Figure 3.




Figure 3. The mean number of action potentials within the depolarization current-induced series (A), latency time (B) and threshold voltage (C) of the first action potential, peak voltage within the series of action potentials (D), and minimal current needed to induce the series (E) for cultivated neurons isolated from the hippocampi of newborn offspring of dams that experienced pregestational chronic unpredictable stress (CUS), prenatal mirtazapine treatment, or their combination. n (the number of individual cells/number of litters in the group) = 48/5 in NO CUS+VEH, 60/5 in NO CUS+MIR, 68/5 in CUS+VEH, and 50/5 in CUS+MIR; * $p < 0.05$ and *** $p < 0.001$, Bonferroni post hoc test. Panel (F) shows representative example of an AP recording.

The two-way ANOVA revealed a statistically significant effect of maternal mirtazapine on the first AP latency time ($F_{1,225} = 4.93, p = 0.03$; Figure 3B) and a significant interaction between maternal stress and mirtazapine with respect to the number of APs within the

series ($F_{1,225} = 4.46, p = 0.04$; Figure 3A), first AP threshold voltage ($F_{1,225} = 33.65, p < 0.001$; Figure 3C), and minimal current required to induce the series ($F_{1,225} = 7.52, p = 0.007$; Figure 3E). Again, CUS suppressed excitability by increasing both the voltage threshold for the generation of the first AP and minimal current amplitude needed for the generation of the first AP series. Mirtazapine rectified both changes.

Figure 4 illustrates the effects of maternal stress, mirtazapine, and their combination on the percentage of neurons in primary cultures with AP series observed during depolarizing current pulses with amplitudes between 100 and 400 pA. A significant interaction effect between maternal stress and mirtazapine was detected during depolarizing pulses of 300 ($F_{1,19} = 4.77, p = 0.04$), 350 ($F_{1,19} = 5.14, p = 0.04$), and 400 pA ($F_{1,19} = 6.44, p = 0.02$). CUS increased excitability at higher depolarizing pulses and mirtazapine treatment suppressed this effect.




Figure 4. Percent of neurons with AP series after depolarization pulse application in primary cultures established from the hippocampi of newborn offspring (A); = 48/5 in NO CUS+VEH, 60/5 in NO CUS+MIR, 68/5 in CUS+VEH, and 50/5 in CUS+MIR) or in slices isolated from the hippocampi of 11–13-day-old offspring (B); = 53/5 in NO CUS+VEH, 63/5 in NO CUS+MIR, 40/5 in CUS+VEH, and 40/5 in CUS+MIR) of dams that experienced pregestational chronic unpredictable stress (CUS), prenatal mirtazapine treatment, or their combination. I *, significant interaction between maternal stress and mirtazapine, $p < 0.05$, S * and S ***, significant effect of maternal stress, $p < 0.05$ and $p < 0.001$, respectively, two-way ANOVA; * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$, between-group comparison, Bonferroni post hoc test.

2.2. Effects of Pregestational Stress and Prenatal Mirtazapine on the Electrophysiological Characteristics of Pyramidal Neurons in Hippocampal Slices Isolated from 11–13-Day-Old Offspring

Individual AP parameters of the pyramidal neurons in the slices isolated from the hippocampi of 11–13-day-old offspring of the dams that experienced pregestational stress, prenatal mirtazapine treatment, or their combination are shown in Figure 5. The two-way ANOVA revealed a significant effect of maternal stress on the threshold voltage ($F_{1,189} = 89.92, p < 0.001$), latency time ($F_{1,189} = 17.75, p < 0.001$), width of action potential ($F_{1,189} = 28.15, p < 0.001$), rise time ($F_{1,189} = 59.21, p < 0.001$), and minimal current needed to induce a series of action potentials ($F_{1,189} = 28.05, p < 0.001$). With respect to the width of the action potential, there was also a significant interaction effect between maternal stress and mirtazapine treatment ($F_{1,189} = 4.53, p = 0.04$).

Figure 6 shows the mean number of action potentials within the depolarization current-induced series (A), latency time (B) and threshold voltage (C) of the first AP, maximal peak voltage within the series of APs (D), and minimal current needed to induce the series (E).

We observed a significant effect of maternal stress on the latency time ($F_{1,195} = 9.73, p < 0.002$; two-way ANOVA) and threshold voltage ($F_{1,195} = 79.40, p < 0.001$; two-way ANOVA) of the first AP and the minimal current needed to induce the series ($F_{1,195} = 25.42, p < 0.001$; two-way ANOVA), a significant effect of maternal mirtazapine on the peak voltage within the series of APs ($F_{1,195} = 4.49, p < 0.04$; two-way ANOVA) and the minimal current needed to induce the series ($F_{1,195} = 9.06, p < 0.003$; two-way ANOVA), and significant

interactions between these factors with respect to the peak voltage ($F_{1,195} = 5.47, p < 0.02$; two-way ANOVA).




Figure 5. Single action potential threshold (A) and peak (B) voltages, latency time (C), width of action potential (D), rise time (E), and the minimal current needed to induce a single action potential (F) for the neurons in the slices isolated from the hippocampi of 11–13-day-old offspring of the dams that experienced pregestational chronic unpredictable stress (CUS), prenatal mirtazapine treatment, or their combination. n (the number of individual cells/number of litters in the group) = 50/5 in NO CUS+VEH, 63/5 in NO CUS+MIR, 38/5 in CUS+VEH, and 39/5 in CUS+MIR; * $p < 0.05$ and *** $p < 0.001$, Bonferroni post hoc test.




Figure 6. The mean number of action potentials within the depolarization current-induced series (A), latency time (B) and threshold voltage (C) of the first action potential, peak voltage within the series of action potentials (D), and the minimal current needed to induce the series (E) for neurons in the slices isolated from the hippocampi of 11–13-day-old offspring of dams that experienced pregestational chronic unpredictable stress (CUS), prenatal mirtazapine treatment, or their combination. n (the number of individual cells/number of litters in the group) = 53/5 in NO CUS+VEH, 63/5 in NO CUS+MIR, 40/5 in CUS+VEH, and 40/5 in CUS+MIR; * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$, Bonferroni post hoc test. Panel (F) shows a representative example of the original AP recordings.

The effects of maternal stress, mirtazapine, and their combination on the percentage of neurons in hippocampal slices with AP series observed during depolarizing current pulses with amplitudes between 100 and 400 pA are shown above in Figure 4B. A significant effect of maternal stress was detected during depolarizing pulses of 250 ($F_{1,18} = 5.79, p = 0.03$), 300 ($F_{1,18} = 26.30, p < 0.001$), 350 ($F_{1,18} = 21.86, p < 0.001$), and 400 pA ($F_{1,18} = 8.53, p = 0.01$).

2.3. Spontaneous Activity of Pyramidal Neurons in Hippocampal Slices Isolated from 11–13-Day-Old Offspring

The spontaneous activity of hippocampal neurons was measured only in the slices. On PNDs 11–13, of all the measured cells, 25% of the neurons in the offspring of nonstressed vehicle-treated dams, 22% of the neurons in the nonstressed mirtazapine-treated group, 35% of the neurons in the CUS-exposed vehicle-treated group, and up to 54% of the neurons in the CUS-exposed mirtazapine-treated group manifested spontaneous activity. The parameters of spontaneous activity, i.e., the average number of fired action potentials, the mean instantaneous frequency, and the firing frequency of action potentials, were evaluated, and no significant differences were found.

3. Discussion

The effects of depression on offspring development and how antidepressant treatment during pregnancy influences such changes are still open questions. Animal models offer the possibility to precisely control the pregestational and gestational states of the mother and pharmacotherapy. The offspring of treated mothers can be monitored under defined conditions for a longer period of time. Moreover, it is possible to combine behavioral, molecular, and electrophysiological analyses in the same settings [27].

In this study, we used a well-established pregestational chronic unpredictable stress (CUS) rat model of maternal depression [21,29]. As a tested antidepressant, a nonselective alpha 2-adrenoceptor antagonist, mirtazapine, which has antidepressant activity in major depression, was used. In this model, we recently reported behavioral changes in offspring at juvenile and adolescent ages [21].

CUS and/or mirtazapine significantly affected several parameters of hippocampal excitability in cultured primary neurons, whereas another set of parameters was not changed. CUS suppresses the excitability, as manifested in hyperpolarized resting membrane potential, increased the voltage threshold for an action potential (AP), and increased the minimal depolarizing current pulse amplitude for activation of both single AP and AP series. Similar effects were caused by mirtazapine when it was administered to dams not exposed to CUS: the resting membrane potential was hyperpolarized, the AP threshold and latency were increased, the minimal depolarizing current pulse amplitude for the activation of a single AP was increased, and the AP rise time was prolonged. Mirtazapine administered to CUS-exposed dams had a positive effect on excitability in selected CUS-affected parameters: it lowered the voltage threshold for the generation of a single AP and AP series and lowered the minimal depolarizing current pulse amplitude for the activation of a single AP. Taken together, the administration of mirtazapine to control dams had minor effects on the hippocampal excitability of offspring. Pregestational CUS, however, suppressed the hippocampal excitability of offspring to a moderate extent. The treatment of CUS-exposed dams with mirtazapine partly compensated for the observed changes in hippocampal excitability.

Approximately two weeks of maturation of the hippocampus under native conditions resulted in a different pattern of excitability alterations observed in acute slices. CUS mostly facilitates excitability by lowering both the threshold for AP activation and the minimal current amplitude needed to activate a single AP and AP series, shortening the AP latency and accelerating the AP rise time. The negative effects of mirtazapine administration to healthy control offspring almost disappeared during the first weeks of postnatal development. It was restricted to lowering the minimal current amplitude needed to activate a single AP and AP series (facilitation of excitability) and lowering the AP peak voltage in the AP series (negative effect). Mirtazapine administered to CUS-exposed dams facilitated the excitability of offspring in a majority of the analyzed parameters. The effect

of CUS was opposite to that observed in primary cultures but remained minor to moderate. Mirtazapine treatment did not compensate for these changes.

While numerous studies have analyzed the behavioral and/or metabolic effects of maternal stress on offspring, alterations in neuronal excitability measured at the level of individual neurons are less frequently investigated. Grigoryan and Segal [30] analyzed the excitability of hippocampal neurons in a primary culture established from the offspring of dams exposed to stress during gestation on the day of birth. They reported no differences in the resting membrane potential, input resistance, or AP threshold voltage between the stressed and nonstressed groups. Nevertheless, they reported reduced GABAergic inhibition, which resulted in a greater frequency of network activity in cultured neurons [30]. In the prenatally stressed group, a reduction in inhibitory tone was also confirmed in hippocampal slices of 2–3-week-old rats from the same cohort. These findings support our findings that maternal stress may facilitate the hippocampal excitability of offspring in early life, whereas they contrast with our findings in primary cultures. This difference may arise from a different stress paradigm and the time of its application. The stress paradigm applied during gestation might have had a stronger effect at the time of birth. Another study in primary hippocampal culture [27] reported a negative effect of pregestational stress on AP generation, which aligns with the findings of this study. Furthermore, maternal CUS decreased the expression of 5-HT and serotonin transporters. Moreover, it increased the 5-HT level in the hippocampi of 20-day-old embryos [31], which is also in line with the decreased excitability of hippocampal neurons just after birth observed in this work.

Experiments on acute slices reflect not only two weeks of neuronal development in a native environment but also altered developmental interactions with mothers and siblings. Indeed, the results we obtained from the slices differ from those obtained from the primary cultures. The effect of the administration of mirtazapine to nonstressed dams was markedly reduced in slices and restricted to only two parameters. The significant change in the resting membrane potential observed in cultures disappeared, and several parameters of excitability were increased in offspring from CUS-exposed mothers. These results parallel those of several other studies in which a pregestational maternal stress model was used. The antidepressant venlafaxine decreased the number of mature hippocampal neurons in juvenile (31 days) offspring of the control but not CUS dams, and this effect disappeared in adolescent (47 days) and adult (75 days) offspring [32]. In adult male rats, pregestational maternal stress reduces neurogenesis in hippocampal granule cells, and perinatal treatment with another antidepressant, fluoxetine, reverses this effect [33].

Pregestational stress decreased the metabolism of the inhibitory neurotransmitter GABA in the hippocampus of 30-day-old offspring, as documented by MRI scans [34], in line with a model used by Grigoryan and Segal [30] and in line with the increased excitability observed in our model in slices. The same results were obtained in our model by Makova et al. [35].

The effects of mirtazapine reported in this study may also be partly explained by its action as an antagonist of 5-HT₂ and 5-HT₃ receptors. While the activation of 5-HT₂, 5-HT₃, or 5-HT₄ receptors results in the depolarization of the membrane potential and increased neuronal excitability [36–38], the activation of 5-HT_{1A} receptors causes hyperpolarization and decreased excitability [36,39,40].

In the present study, we observed spontaneous activity in hippocampal neurons from all four offspring groups, with no significant differences in quantitative parameters of AP firing between the groups. However, the percentage of spontaneously active neurons was increased in CUS group. In our earlier work [27], we reported increased levels of spontaneous activity in neurons from primary hippocampal cultures in the stressed group.

Our findings can be interpreted in terms of altered GABAergic and glutamatergic systems, which enhance spontaneous action potential firing [8].

The primary limitation of the present study is the fact that the offspring in each group (those of nonstressed vehicle treated dams, nonstressed mirtazapine treated dams, stressed vehicle treated dams, and stressed mirtazapine treated dams) were assessed as a whole, without distinguishing between males and females.

4. Materials and Methods

4.1. Animals

Female and male Wistar rats (initial weight 200–220 g; 2–3 months old) were obtained from the Department of Toxicology and Laboratory Animals Breeding, Centre of Experimental Medicine of the Slovak Academy of Sciences, Dobra Voda, Slovak Republic. All the animals were housed (38 × 59 × 25 cm³ large cages) under standard laboratory conditions (temperature: 22 ± 2 °C, humidity: 55 ± 10%) with a 12 h light/12 h dark cycle (lights on at 7.00 a.m.). Pelleted food and tap water were available ad libitum. The State Veterinary and Food Administration of the Slovak Republic approved all the experimental procedures (Approval No. 3334-5/2020-220 from 11 December 2020). The rats were handled according to the Guide for the Care and Use of Laboratory Animals (N.R.C., 1996) and the European Communities Council Directive of September 22, 2010 (2010/63/EU, 74). In accordance with the Directive, the animal environment was enriched, e.g., by providing hiding places and tissue paper for exploration and nest building.

4.2. Experimental Design

The timing of the experiments is shown in Figure 7.




Figure 7. Schedule of the experiment. G—gestation day; PND—postnatal day; DIV—days in vitro.

4.3. Chronic Unpredictable Stress Schedule

After one week of adaptation, the female rats ($n = 20$) were randomly allocated to one of the following two groups: no chronic unpredictable stress (NO CUS; $n = 10$) and chronic unpredictable stress (CUS; $n = 10$). Animals from the stressed group were randomly exposed to 6 distinct stressors for three consecutive weeks: (1) exposure to damp bedding (12 h); (2) food deprivation (24 h); (3) predator stress—clothing with cat odor (10 h); (4) water deprivation (12 h); (5) cage decline at a 45° angle (6 h); and (6) Strobe light flicker lights (12 h). After the stress procedure, female rats from both groups were mated with males (3:1). At gestational day 0 (G0), each pregnant rat was housed separately. The G0 day was detected by the presence of sperm in vaginal smears.

4.4. Mirtazapine Treatment

On the tenth day of gestation, the dams from each group (NO CUS and CUS) were randomly divided into two sub-groups: vehicle (VEH) and mirtazapine (MIR), with five animals in each sub-group: NO CUS+VEH, NO CUS+MIR, CUS+VEH, and CUS+MIR. Mirtazapine (Mikrochem Trade s.r.o.; $M = 265.36 \text{ g/mol}$) was dissolved in citric acid and administered orally via a wafer biscuit (size of 1 cm^3) to pregnant rats at a clinically relevant dose of 10 mg/kg/day (Figure 7). The dose used in this experiment was chosen on the basis of previous results [21]. The female rats in the control groups received a 1 cm^3 wafer biscuit once daily, which was filled with vehicle.

4.5. Primary Hippocampal Cell Culture

Primary hippocampal cultures were prepared on PND 0 according to the methods of Reichova et al. [41]. For each primary culture preparation, three newborn pups were randomly selected from each litter of each dam from the groups NO CUS+VEH, NO CUS+MIR, CUS+VEH, and CUS+MIR. The animals were euthanized by cutting off the head using a special guillotine designed for killing laboratory rodents. The brains were dissected in ice-cold Hank's balanced salt solution (HBSS; Gibco, Jenks, OK, USA) supplemented with 1% antibiotic-antimycotic (Sigma-Aldrich, Darmstadt, Germany) and 0.3 M HEPES (Sigma-Aldrich, Darmstadt, Germany) under a stereomicroscope (Optika Microscopes, Ponteranica, Italy). Hippocampal tissues were dissociated for 20 min at 37°C with an enzymatic solution containing HBSS, 0.1% trypsin (Gibco, Jenks, OK, USA), and 0.1 mg/mL DNase I (Roche, Mannheim, Germany). The cells were subsequently resuspended and plated on 24-well plates containing coverslips (Sarstedt, Nümbrecht, Germany) coated with $10 \mu\text{g/mL}$ poly-D-lysine (Sigma-Aldrich, Darmstadt, Germany) at a density of $0.8\text{--}1 \times 10^5 \text{ cells/mL}$ in RPMI medium (Gibco, Jenks, OK, USA) containing 10% fetal bovine serum (Sigma-Aldrich, Darmstadt, Germany). After 3 h of cell culture at 37°C in a 5% CO_2 incubator, the medium was replaced with selective growth medium (Neurobasal A; Gibco, Jenks, OK, USA) supplemented with 1% antibiotic-antimycotic (Sigma-Aldrich, Darmstadt, Germany), 1% glutamine (Sigma-Aldrich, Darmstadt, Germany), and 2% B27 supplement (Gibco, Jenks, OK, USA). This was assigned as day 1 in vitro (DIV 1). At DIV5, 50% of the fresh growth medium was replaced. Cultured hippocampal neurons with pyramidal-like morphology were used for electrophysiology experiments at DIV 8–10. The excitability of at least 30 neurons were recorded from each culture. The exact numbers of the neurons/recordings from the hippocampal cultures are provided in the legends to the figures.

4.6. Acute Hippocampal Slices

Acute hippocampal slices were prepared on PNDs 11–13. Three were randomly selected from each litter of each dam from the groups NO CUS+VEH, NO CUS+MIR, CUS+VEH, and CUS+MIR. The animals were euthanized by cutting off the head using a special guillotine designed for killing laboratory rodents. The brains were cut on a vibratome (Campden Instruments, Loughborough, England) into $300 \mu\text{m}$ thick slices in ice-cold artificial cerebrospinal fluid (ACSF) containing the following (in mM): 125 NaCl, 3 KCl, 25 NaHCO_3 , 1.25 NaH_2PO_4 , 2 CaCl_2 , 1 MgCl_2 , and 10 D-glucose (all chemicals were purchased from Sigma-Aldrich, Darmstadt, Germany). ACSF was oxygenated with 95% O_2 and 5% CO_2 , pH 7.4. Before recording, the slices were incubated for 60 min at 37°C and then for 30 min at room temperature in oxygenated ACSF solution. The exact numbers of the neurons/recordings from the hippocampal slices are provided in the legends to the figures.

4.7. Electrophysiological Experiments

All electrophysiology measurements were performed via the whole-cell current clamp configuration of the patch-clamp technique. The signals were recorded via a HEKA EPC10 amplifier (HEKA Electronics, Reutlingen, Germany) controlled by Patchmaster v2x73.3 (HEKA Electronics, Reutlingen, Germany) software. The data were analyzed via Fitmaster v2x73.3 (HEKA Electronics, Reutlingen, Germany) and Origin 2022 Academic (OriginLab Co., Northampton, MA, USA) software. The parameters of an action potential (AP) series were extracted from the experimental traces with custom code via the open-source Electrophysiological Feature Extraction Library (eFEL, <https://github.com/BlueBrain/eFEL>, accessed on 25 March 2025).

Patch electrodes were pulled from borosilicate glass with 3–4 M Ω resistance. The extracellular solution in primary hippocampal cultures contained the following (in mM): 130 NaCl; 3 KCl; 10 HEPES; 2 CaCl₂; 1 MgCl₂; 10 D-glucose; pH 7.4 (with NaOH), and the intracellular solution contained the following (in mM): 120 K-gluconate; 20 KCl; 2 MgCl₂; 2 Na₂ATP; 0.25 Na₂GTP; 10 HEPES; pH 7.4 (with KOH) (all chemicals from Sigma–Aldrich, Darmstadt, Germany). The extracellular solution for acute neuronal slices was ASCF solution oxygenated with 95% O₂ and 5% CO₂, pH 7.4, and the intracellular solution was the same as that for the primary hippocampal cultures. The osmolality of an intracellular solution was approximately 290–300 mOsm/L (Osmomat 030-Gonotec, Berlin, Germany). The osmolality of an extracellular solution was adjusted by adding sucrose to a final osmolality, which was 2–3 mOsm/L lower than the osmolality of the corresponding intracellular solution.

The actual resting membrane potential (V_{rest}) was read immediately after establishing the whole-cell configuration and switching to current-clamp mode. During all other experiments, the resting membrane potential was held at -70 mV. The input resistance (R_{inp}) was measured by a series of 250 ms long hyperpolarizing current pulses with the amplitude decreasing from -70 mV with a step of -25 pA. The membrane potential at the end of each current pulse was read and the R_{inp} value was calculated from the fit of the voltage–current relationship by the Ohm equation. Single APs were activated by a series of 5 ms long depolarizing current pulses with the amplitude increasing with a step of $+50$ pA. A series of APs were activated by a series of 300 ms long depolarizing current pulses with the amplitude increasing with a step of $+50$ pA. Spontaneous activity was measured for 5 min from each neuron's resting membrane potential.

4.8. Statistical Analyses

The statistical analyses were performed via GraphPad Prism 8.3 software. Individual recordings from individual neurons were used as data points for the statistical analysis. Outliers were identified and excluded via GraphPad PRISM's ROUT method ($Q = 1\%$). Neurons, considered as an outlier according to one parameter, were excluded from the assessment of all other characteristics. The data were analyzed by a two-way analysis of variance (ANOVA) with treatment (mirtazapine) and conditions (stress vs. nonstress) as the main factors, followed by the Bonferroni post hoc test. The results are expressed as the mean \pm standard error of the mean (SEM), and a value of $p \leq 0.05$ was considered statistically significant.

5. Conclusions

To summarize, we have shown that in our animal model, the treatment of control nonstressed dams with mirtazapine during pregnancy moderately affects the hippocampal excitability of offspring. Prenatal CUS results in moderately suppressed hippocampal excitability in offspring at birth, and this effect reverses to moderate facilitation during

early postnatal development. Treatment with mirtazapine during pregnancy partly partially reduced the effects of CUS, and this partial reduction becomes more prominent during postnatal development. Therefore, treatment with mirtazapine may be also beneficial during the prenatal and perinatal periods.

Author Contributions: Conceptualization, M.D. and L.L.; methodology, L.D.-H., A.F., S.B., B.J.-T. and K.O., software, M.T.; validation, M.D., E.D. and L.L.; formal analysis, L.D.-H., A.F., M.T. and E.D.; investigation, L.D.-H., A.F., S.B., K.O. and BJT; resources, M.D., E.D. and L.L.; data curation, L.D.-H., A.F., S.B. and K.O.; writing—original draft preparation, L.D.-H. and K.O.; writing—review and editing, L.D.-H., A.F., S.B., K.O., M.T., B.J.-T., M.D., E.D. and L.L.; visualization, E.D. and L.L.; supervision, M.D. and L.L.; project administration, M.D. and L.L.; funding acquisition, M.D., E.D. and L.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the Slovak Research and Development Agency under Contracts No. APVV-19-0435, APVV-20-0202, and APVV-22-0061 and by the Research Grant Agency of the Slovak Academy of Sciences and Ministry of Education, Science, Research and Sport of the Slovak Republic via the grants No. VEGA 2/0057/22, VEGA 2/0081/22, VEGA 2/0133/23.

Institutional Review Board Statement: The State Veterinary and Food Administration of the Slovak Republic approved all experimental procedures with animals (Approval No. 3334-5/2020-220 from 11 December 2020).

Informed Consent Statement: Not applicable.

Data Availability Statement: The data that support the findings of this study are available from the corresponding authors upon reasonable request.

Acknowledgments: The authors thank Alexandra Reichová, for practical advice and recommendations on preparing primary hippocampal cultures.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

5-HT	Serotonin
ANOVA	Analysis of variance
AP	Action potential
CUS	Chronic unpredictable stress
MIR	Mirtazapine
MDPI	Multidisciplinary Digital Publishing Institute
R _{in} p	Input resistance
SSRIs	Selective serotonin reuptake inhibitors
VEH	Vehicle
V _{rest}	Resting membrane potentials
WHO	World Health Organization

References

1. Shidhaye, P.; Giri, P. Maternal depression: A hidden burden in developing countries. *Ann. Med. Health Sci. Res.* **2014**, *4*, 463–465. [[PubMed](#)]
2. Barker, E.D.; Jaffee, S.R.; Uher, R.; Maughan, B. The contribution of prenatal and postnatal maternal anxiety and depression to child maladjustment. *Depress. Anxiety* **2011**, *28*, 696–702. [[PubMed](#)]
3. Stein, A.; Pearson, R.M.; Goodman, S.H.; Rapa, E.; Rahman, A.; McCallum, M.; Howard, L.M.; Pariante, C.M. Effects of perinatal mental disorders on the fetus and child. *Lancet* **2014**, *384*, 1800–1819. [[CrossRef](#)] [[PubMed](#)]
4. Gelaye, B.; Rondon, M.B.; Araya, R.; Williams, M.A. Epidemiology of maternal depression, risk factors, and child outcomes in low-income and middle-income countries. *Lancet Psychiatry* **2016**, *3*, 973–982.

5. Accortt, E.E.; Cheadle, A.C.; Dunkel Schetter, C. Prenatal depression and adverse birth outcomes: An updated systematic review. *Matern. Child. Health J.* **2015**, *19*, 1306–1337.
6. Silveira, M.F.; Victora, C.G.; Horta, B.L.; da Silva, B.G.C.; Matijasevich, A.; Barros, F.C.; Pelotas Cohorts Study, G. Low birthweight and preterm birth: Trends and inequalities in four population-based birth cohorts in Pelotas, Brazil, 1982–2015. *Int. J. Epidemiol.* **2019**, *48* (Suppl. S1), i46–i53.
7. Eshete, A.; Alemu, A.; Zerfu, T.A. Magnitude and Risk of Dying among Low Birth Weight Neonates in Rural Ethiopia: A Community-Based Cross-Sectional Study. *Int. J. Pediatr.* **2019**, *2019*, 9034952. [[CrossRef](#)]
8. Ahun, M.N.; Cote, S.M. Maternal depressive symptoms and early childhood cognitive development: A review of putative environmental mediators. *Arch. Womens Ment. Health* **2019**, *22*, 15–24.
9. Kingston, D.; Kehler, H.; Austin, M.P.; Mughal, M.K.; Wajid, A.; Vermeyden, L.; Benzies, K.; Brown, S.; Stuart, S.; Giallo, R. Trajectories of maternal depressive symptoms during pregnancy and the first 12 months postpartum and child externalizing and internalizing behavior at three years. *PLoS ONE* **2018**, *13*, e0195365.
10. Rogers, A.; Obst, S.; Teague, S.J.; Rossen, L.; Spry, E.A.; Macdonald, J.A.; Sunderland, M.; Olsson, C.A.; Youssef, G.; Hutchinson, D. Association Between Maternal Perinatal Depression and Anxiety and Child and Adolescent Development: A Meta-analysis. *JAMA Pediatr.* **2020**, *174*, 1082–1092.
11. Braun, K.; Bock, J.; Wainstock, T.; Matas, E.; Gaisler-Salomon, I.; Fegert, J.; Ziegenhain, U.; Segal, M. Experience-induced transgenerational (re-)programming of neuronal structure and functions: Impact of stress prior and during pregnancy. *Neurosci. Biobehav. Rev.* **2020**, *117*, 281–296. [[CrossRef](#)]
12. Grundwald, N.J.; Brunton, P.J. Prenatal stress programs neuroendocrine stress responses and affective behaviors in second generation rats in a sex-dependent manner. *Psychoneuroendocrinology* **2015**, *62*, 204–216. [[CrossRef](#)] [[PubMed](#)]
13. Benard-Laribiere, A.; Pambrun, E.; Sutter-Dallay, A.L.; Gautier, S.; Hurault-Delarue, C.; Damase-Michel, C.; Lacroix, I.; Begaud, B.; Pariente, A. Patterns of antidepressant use during pregnancy: A nationwide population-based cohort study. *Br. J. Clin. Pharmacol.* **2018**, *84*, 1764–1775. [[CrossRef](#)] [[PubMed](#)]
14. Molenaar, N.M.; Kamperman, A.M.; Boyce, P.; Bergink, V. Guidelines on treatment of perinatal depression with antidepressants: An international review. *Aust. N. Z. J. Psychiatry* **2018**, *52*, 320–327. [[CrossRef](#)] [[PubMed](#)]
15. Kim, J.; Riggs, K.W.; Misri, S.; Kent, N.; Oberlander, T.F.; Grunau, R.E.; Fitzgerald, C.; Rurak, D.W. Stereoselective disposition of fluoxetine and norfluoxetine during pregnancy and breast-feeding. *Br. J. Clin. Pharmacol.* **2006**, *61*, 155–163. [[CrossRef](#)]
16. Rampono, J.; Simmer, K.; Ilett, K.F.; Hackett, L.P.; Doherty, D.A.; Elliot, R.; Kok, C.H.; Coenen, A.; Forman, T. Placental transfer of SSRI and SNRI antidepressants and effects on the neonate. *Pharmacopsychiatry* **2009**, *42*, 95–100. [[CrossRef](#)]
17. Dubovicky, M.; Belovicova, K.; Csatlósova, K.; Bogi, E. Risks of using SSRI/SNRI antidepressants during pregnancy and lactation. *Interdiscip. Toxicol.* **2017**, *10*, 30–34. [[CrossRef](#)]
18. Gemmel, M.; Hazlett, M.; Bogi, E.; De Lacalle, S.; Hill, L.A.; Kokras, N.; Hammond, G.L.; Dalla, C.; Charlier, T.D.; Pawluski, J.L. Perinatal fluoxetine effects on social play, the HPA system, and hippocampal plasticity in pre-adolescent male and female rats: Interactions with pre-gestational maternal stress. *Psychoneuroendocrinology* **2017**, *84*, 159–171. [[CrossRef](#)]
19. Millard, S.J.; Weston-Green, K.; Newell, K.A. The effects of maternal antidepressant use on offspring behaviour and brain development: Implications for risk of neurodevelopmental disorders. *Neurosci. Biobehav. Rev.* **2017**, *80*, 743–765. [[CrossRef](#)]
20. Cipriani, A.; Furukawa, T.A.; Salanti, G.; Chaimani, A.; Atkinson, L.Z.; Ogawa, Y.; Leucht, S.; Ruhe, H.G.; Turner, E.H.; Higgins, J.P.T.; et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *Lancet* **2018**, *391*, 1357–1366. [[CrossRef](#)]
21. Viñas-Noguera, M.; Csatlósova, K.; Šimončíčová, E.; Bögi, E.; Ujházy, E.; Dubovický, M.; Belovičová, K. Sex- and age- dependent effect of pre-gestational chronic stress and mirtazapine treatment on neurobehavioral development of Wistar rat offspring. *PLoS ONE* **2022**, *17*, e0255546.
22. Smit, M.; Dolman, K.M.; Honig, A. Mirtazapine in pregnancy and lactation—A systematic review. *Eur. Neuropsychopharmacol.* **2016**, *26*, 126–135. [[CrossRef](#)] [[PubMed](#)]
23. Idunkova, A.; Lacinova, L.; Dubiel-Hoppanova, L. Stress, depression, and hippocampus: From biochemistry to electrophysiology. *Gen. Physiol. Biophys.* **2023**, *42*, 107–122.
24. Castanheira, L.; Silva, C.; Cheniaux, E.; Telles-Correia, D. Neuroimaging Correlates of Depression—Implications to Clinical Practice. *Front. Psychiatry* **2019**, *10*, 703.
25. Aleksandrova, L.R.; Wang, Y.T.; Phillips, A.G. Evaluation of the Wistar-Kyoto rat model of depression and the role of synaptic plasticity in depression and antidepressant response. *Neurosci. Biobehav. Rev.* **2019**, *105*, 1–23.
26. Bogi, E.; Belovicova, K.; Csatlósova, K.; Dubovicky, M. Animal models of maternal depression for monitoring neurodevelopmental changes occurring in dams and offspring. *Interdiscip. Toxicol.* **2017**, *10*, 35–39.
27. Bogi, E.; Belovicova, K.; Moravcikova, L.; Csatlósova, K.; Dremencov, E.; Lacinova, L.; Dubovicky, M. Pre-gestational stress impacts excitability of hippocampal cells in vitro and is associated with neurobehavioral alterations during adulthood. *Behav. Brain Res.* **2019**, *375*, 112131.

28. Belovicova, K.; Bogi, E.; Koprdoва, R.; Ujhazy, E.; Mach, M.; Dubovicky, M. Effects of venlafaxine and chronic unpredictable stress on behavior and hippocampal neurogenesis of rat dams. *Neuro Endocrinol. Lett.* **2017**, *38*, 19–26.
29. Czarzasta, K.; Bogacki-Rychlik, W.; Segiet-Swiecicka, A.; Kruszewska, J.; Malik, J.; Skital, V.; Kasarello, K.; Wrzesien, R.; Bialy, M.; Sajdel-Sulkowska, E.M. Gender differences in short- vs. long-term impact of maternal depression following pre-gestational chronic mild stress. *Exp. Neurol.* **2022**, *353*, 114059. [[CrossRef](#)]
30. Grigoryan, G.; Segal, M. Prenatal stress affects network properties of rat hippocampal neurons. *Biol. Psychiatry* **2013**, *73*, 1095–1102.
31. Huang, Y.; Xu, H.; Li, H.; Yang, H.; Chen, Y.; Shi, X. Pre-gestational stress reduces the ratio of 5-HIAA to 5-HT and the expression of 5-HT1A receptor and serotonin transporter in the brain of foetal rat. *BMC Neurosci.* **2012**, *13*, 22.
32. Belovicova, K.; Simoncicova, E.; Noguera, M.V.; Dubovicky, M.; Bogi, E. Long-term effects of pre-gestational stress and perinatal venlafaxine treatment on neurobehavioral development of female offspring. *Behav. Brain Res.* **2021**, *398*, 112944. [[PubMed](#)]
33. Gemmel, M.; De Lacalle, S.; Mort, S.C.; Hill, L.A.; Charlier, T.D.; Pawluski, J.L. Perinatal fluoxetine has enduring sexually differentiated effects on neurobehavioral outcomes related to social behaviors. *Neuropharmacology* **2019**, *144*, 70–81. [[PubMed](#)]
34. Huang, Y.; Shen, Z.; Hu, L.; Xia, F.; Li, Y.; Zhuang, J.; Chen, P.; Huang, Q. Exposure of mother rats to chronic unpredictable stress before pregnancy alters the metabolism of gamma-aminobutyric acid and glutamate in the right hippocampus of offspring in early adolescence in a sexually dimorphic manner. *Psychiatry Res.* **2016**, *246*, 236–245. [[PubMed](#)]
35. Makova, M.; Kasparova, S.; Tvrdik, T.; Noguera, M.; Belovicova, K.; Csatlosova, K.; Dubovicky, M. Mirtazapine modulates Glutamate and GABA levels in the animal model of maternal depression. MRI and (1)H MRS study in female rats. *Behav. Brain Res.* **2023**, *442*, 114296.
36. Xiang, Z.; Prince, D.A. Heterogeneous actions of serotonin on interneurons in rat visual cortex. *J. Neurophysiol.* **2003**, *89*, 1278–1287.
37. McMahon, L.L.; Kauer, J.A. Hippocampal interneurons are excited via serotonin-gated ion channels. *J. Neurophysiol.* **1997**, *78*, 2493–2502.
38. Ropert, N.; Guy, N. Serotonin facilitates GABAergic transmission in the CA1 region of rat hippocampus in vitro. *J. Physiol.* **1991**, *441*, 121–136.
39. Beck, S.G.; Choi, K.C.; List, T.J. Comparison of 5-hydroxytryptamine1A-mediated hyperpolarization in CA1 and CA3 hippocampal pyramidal cells. *J. Pharmacol. Exp. Ther.* **1992**, *263*, 350–359.
40. Tanaka, E.; North, R.A. Actions of 5-hydroxytryptamine on neurons of the rat cingulate cortex. *J. Neurophysiol.* **1993**, *69*, 1749–1757.
41. Reichova, A.; Schaller, F.; Bukatova, S.; Bacova, Z.; Muscatelli, F.; Bakos, J. The impact of oxytocin on neurite outgrowth and synaptic proteins in Magel2-deficient mice. *Dev. Neurobiol.* **2021**, *81*, 366–388.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.