

Article

Characterization of Chemical Constituents of *Oxytropis microphylla* (Pall.) DC. by Ultra-High-Performance Liquid Chromatography Coupled with Quadrupole-Time-of Flight Tandem Mass Spectrometry

Mengqi Xia ^{1,2,†}, Min Yao ^{1,3,†}, Junmao Li ¹, Jianjian Zhang ^{1,2}, Yayun Yu ^{1,2}, Shilin Yang ¹, Guoyue Zhong ^{1,4}, Na Pei ^{1,5}, Hui Ouyang ^{1,6,*} and Yulin Feng ^{1,6,*}

¹ School of pharmacy, Jiangxi University of Traditional Chinese Medicine, Nanchang 330006, China

² Jiangxi Bencao Tiangong Technology Co., Ltd., Nanchang 330006, China

³ Jiangxi Institute for Drug Control, Nanchang 330029, China

⁴ Research Center of Natural Resources of Chinese Medicinal Materials and Ethnic Medicine, Jiangxi University of Traditional Chinese Medicine, Nanchang 330004, China

⁵ School of Pharmacy, Yichun University, Yichun 336000, China

⁶ State Key Laboratory of Innovative Drug and Efficient Energy-Saving Pharmaceutical Equipment, Nanchang 330006, China

* Correspondence: 20070982@jxutcm.edu.cn (H.O.); 20081002@jxutcm.edu.cn (Y.F.); Tel.: +0791-87119632 (Y.F.)

† These authors contributed equally to this work.

Citation: Xia, M.; Yao, M.; Li, J.; Zhang, J.; Yu, Y.; Yang, S.; Zhong, G.; Pei, N.; Ouyang, H.; Feng, Y. Characterization of Chemical Constituents of *Oxytropis microphylla* (Pall.) DC. by Ultra-High-Performance Liquid Chromatography Coupled with Quadrupole-Time-of Flight Tandem Mass Spectrometry. *Separations* **2022**, *9*, 297.

<https://doi.org/10.3390/separations9100297>

Academic Editor: Victoria Samanidou

Received: 26 August 2022

Accepted: 30 September 2022

Published: 9 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: *Oxytropis microphylla* (Pall.) DC. is a traditional Tibetan medicine used as an external preparation for clearing heat and detoxification, healing sore muscles, astringent vein hemostasis, defecation, and treating plague, constipation, anthrax, and swollen and painful furuncles. It remains a challenge to comprehensively analyze and identify the chemical constituents of *Oxytropis microphylla* (Pall.) DC. In this study, a new analytical method using a combination of ultra-high-performance liquid chromatography–mass spectrometry (UHPLC-MS) and effective data mining techniques was established to identify the chemical constituents of *Oxytropis microphylla*. A total of 127 compounds were identified in *O. microphylla* extract, including 92 flavonoids, 15 indole alkaloids, and 20 others. After the oral administration of the extract to rats, 22 metabolites were identified in the plasma. The primary in vivo metabolic reactions that occurred after the administration of *O. microphylla* extract were glucuronidation and sulfation. Therefore, we successfully devised a high-efficiency method to distinguish compounds and used it as a source of post-study to identify the active biological components of *O. microphylla* extract.

Keywords: *Oxytropis microphylla* (Pall.) DC.; HPLC-TOF-MS; data mining strategy; chemical profiling; metabolites

1. Introduction

More than 140 species and 3 varieties of *Oxytropis* (Family: Leguminosae) are found in China. *Oxytropis* plants were used extensively in the west owing to their outstanding efficiency [1–3]. *Oxytropis microphylla* (Pall.) DC. (OMDC) is a perennial herb, locally identified as the “king of herbs” on account of its analgesic and outstanding anti-inflammatory effects. It is found in the valleys, hillsides, and meadows of the Qinghai–Tibet plateau at an altitude of 2700–4300 m [4–6]. In ancient books, OMDC has been recorded for its efficiency as an analgesic and for anti-inflammatory, detoxification, promotion of blood circulation, and heat-clearing effects [7,8]. Several bioactive compounds have been separated from *Oxytropis* using phytochemical methods. Flavones are the major active ingredients of *Oxytropis* that are known to exert anti-inflammatory, analgesic, ultraviolet damage-

protective, and antitumor effects, and to enhance immune cofunction [9,10]. Currently, only a few reports exist in the literature on the systematic characterization of the chemical components of this plant, which do not effectively explain its active components [11]. Therefore, there is an urgent requirement to establish a scientific method to identify the chemical constituents of OMDC.

To better understand the characteristics of OMDC, it is necessary to establish a robust identification method to analyze the chemical constituents of the plant [12,13]. The current technology used to analyze the potential active components of traditional Chinese medicine is UHPLC-QTOF-MS, which can predict the elemental composition of the tested compound and accurately determine its quality [13–16]. Its high resolution, efficiency, and structure recognition ability can detect hundreds or thousands of chemical components in complex samples and rapidly characterize the extracts used in traditional Chinese medicine [17–20]. Therefore, UHPLC-QTOF-MS is widely used in the qualitative analysis of medicinal materials.

To the best of our knowledge, only a few reports have used UHPLC-QTOF-MS to identify compounds in the crude extract of OMDC, and there is no comprehensive or systematic explanation to analyze their chemical components. Therefore, there is an urgent requirement to establish a reliable method to identify the chemical components of OMDC at micro-concentration levels and to summarize MS fragmentation behaviors. To address this problem, we established a UHPLC-QTOF-MS method for the identification, classification, and systematic investigation of the chemical components of OMDC.

2. Materials and Methods

2.1. Chemicals and Reagents

Acetonitrile and formic acid used for sample processing were sourced from Fisher Scientific (Fair Lawn, NJ, USA) and Sigma Aldrich (Sigma Aldrich, St. Louis, MO, USA), respectively. Pure distilled water was obtained using a Millipore Alpha-Q water purification system. All 6 reference standards having a purity > 98% (2',4'-dihydroxychalcone, 7-hydroxyflavanone, pinocembrin, quercetin, apigenin, luteolin) were purchased from Sichuan Weikeqi Biological Technology Co., Ltd. (Chengdu, China).

2.2. Animal Experiments

Three male Sprague-Dawley rats (weighing 180 ± 20 g) were obtained from the Hunan Laike Jingda Experimental Animal Co. Ltd., Changsha, China. All rats were housed in a room (a temperature of 20°C and humidity of 50%) and were provided access to food and water ad libitum. OMDC extract was suspended in 0.5% CMC-Na and administered orally to rats at a dose of 150 mg/kg body weight. After drug administration, blood was collected from the inner canthus vein at fixed intervals (1, 2, 4, 6, 12 h). Blood samples were centrifuged at 5000 rpm for 10 min to obtain the serum. The animal experiments were approved by the Laboratory Animal Ethics Committee of Jiangxi University of Traditional Chinese Medicine (approval No. SYXK2017-0004).

2.3. Sample Preparation

OMDC was obtained from Tibet Province in 2020. The sample was identified by Professor Guo-yue Zhong and Ming Yao (Jiangxi University of Traditional Chinese Medicine). The crude extract of OMDC was prepared using the medicinal powder. Briefly, 1 g of OMDC powder was mixed with 50 mL methanol and extracted for 30 min using ultrasonication in a water bath maintained at room temperature ($20\text{--}30^{\circ}\text{C}$). Extracts were filtered through a $0.22\text{-}\mu\text{M}$ microporous membrane filter and $3.0\text{ }\mu\text{L}$ of the filtered sample was used for UHPLC-QTOF-MS.

All reference materials were dissolved in methanol to yield the respective stock solutions, which were stored at 4°C . After mixing all the reference stock solutions, they were

diluted with methanol to a concentration of 10.0 $\mu\text{g}/\text{mL}$, and 3.0 μL of the solution was used for UHPLC-QTOF-MS.

The plasma of the 3 rats from each time point was mixed. Plasma samples were added to 3 times their volume of methanol, vortexed for 2 min, and centrifuged at 12,000 rpm for 10 min. This supernatant was dried under a nitrogen stream at 35 $^{\circ}\text{C}$. The residue was redissolved in 100 μL of 50% methanol and vortexed for 2 min. The solution was centrifuged at 12,000 rpm for 10 min, and 5 μL of the supernatant was injected for UHPLC-QTOF-MS.

2.4. UHPLC-QTOF-MS/MS

A Shimadzu system (Kyoto, Japan) was used for separation. The other systems included a CTO-30AC column oven, a DGU-20A3 degasser, an LC-3AD solvent delivery system, a CBM-20A controller, and a SIL-30ACXR auto-sampler. A Welch UHPLC AQ-C18 (100 mm \times 2.1 mm, 1.8 μm) was used at a temperature of 40 $^{\circ}\text{C}$. The mobile phase was composed of 0.1% formic acid in water (solvent A) and acetonitrile (solvent B) and the flow rate was 0.3 mL/min. The elution conditions were as follows: 0.1–2.0 min, 5–10% B; 2.0–10.0 min, 10–14% B; 10.0–13.0 min, 14–20% B; 13.0–30.0 min, 20–40% B; 30.0–37.0 min, 40–50% B; 37.0–45.0 min, 50–65% B; 45.0–50.0 min, 65–95% B; 50.0–53.0 min, 95% B; 53.0–58.0 min, 5% B.

UHPLC-QTOF-MS/MS analyses were performed in the negative electrospray ionization mode using a Triple TOF™ 5600+ system with a Duo Spray source (AB SCIEX, Foster City, CA, USA). The mass spectrometry conditions were as follows: ion spray voltage, -4500 V; ion source temperature, 550 $^{\circ}\text{C}$; curtaingas, 25 psi; nebulizer gas (GS1), 55 psi; heater gas (GS2), 55 psi; and decluster potential (DP), -100 V. The mass ranges of TOF-MS and TOF MS/MS experiments were all set at 50–1250 Da.

3. Results and Discussion

3.1. Screening and Identification of Chalcone

In the mass spectrometry of protonated chalcones, phCO^+ and $\text{ph}'\text{CH}=\text{CHCO}^+$ were observed as the major fragments. Compound **82** was obtained as a quasi-molecular ion at m/z 239.0713, which was in accordance with the chemical formula of $\text{C}_{15}\text{H}_{12}\text{O}_3$ (Figures 1–3; Table 1). The characteristic ions at m/z 135.0090 [$\text{M}-\text{H}-\text{C}_8\text{H}_8$] $^-$, 197.0600 [$\text{M}-\text{H}-\text{C}_2\text{H}_2\text{O}$] $^-$, 148.0169 [$\text{M}-\text{H}-\text{C}_7\text{H}_7$] $^-$, and 109.0308 [$\text{M}-\text{H}-\text{C}_9\text{H}_6\text{O}$] $^-$ were yielded by C-C bond. Thus, the compound was identified as 2',4'-dihydroxychalcone by retention time of the standard. Compound **58** was inferred by sifting using the DPI acquired at m/z 239.0712 and the NL at 162 Da ($\text{C}_6\text{H}_{10}\text{O}_5$). The chemical formula was computed as $\text{C}_{21}\text{H}_{22}\text{O}_8$ based on the exact mass precursor ion. This compound was determined to be 2',4'-dihydroxychalcone-gluco-side. Compound **43** has two more OH radicals than 2',4'-dihydroxychalcone. The molecular fragment had m/z 253.0507 [$\text{M}-\text{H}-\text{H}_2\text{O}$] $^-$, 151.0038 [$\text{M}-\text{H}-\text{C}_8\text{H}_8\text{O}$] $^-$, 125.0244 [$\text{M}-\text{H}-\text{C}_9\text{H}_6\text{O}_2$] $^-$, 119.0508 [$\text{M}-\text{H}-\text{C}_7\text{H}_4\text{O}_4$] $^-$, 93.0354 [$\text{M}-\text{H}-\text{C}_9\text{H}_6\text{O}_4$] $^-$ and it was structurally similar naringenin chalcone that is reported in the literature [21]. Compound **80** had a mass of 2 Da (2H) more than 2',4'-dihydroxychalcone and was identified as 2',4'-dihydroxydihydro-chalcone [22]. The deprotonation molecular ion of compound **59** ($\text{C}_{15}\text{H}_{12}\text{O}_4$) was discovered at m/z 256.0735. The fragment ions (m/z 237.056 [$\text{M}-\text{H}-\text{H}_2\text{O}$] $^-$, 135.0093 [$\text{M}-\text{H}-\text{C}_8\text{H}_8\text{O}$] $^-$, 119.0512 [$\text{M}-\text{H}-\text{C}_7\text{H}_4\text{O}_3$] $^-$, and 91.0204 [$\text{M}-\text{H}-\text{H}_2\text{O}-\text{C}_9\text{H}_7\text{O}_2$] $^-$) were consistent with isoliquiritigenin reported in the literature [23]. Compounds **81** had a mass of 28 Da (C_2H_2), more than isoliquiritigenin. This generated fragment ions at m/z 225.0916 [$\text{M}-\text{H}-\text{CO}_2$] $^-$, 148.0151 [$\text{M}-\text{H}-\text{CH}_3-\text{C}_7\text{H}_6\text{O}$] $^-$, 119.0507 [$\text{M}-\text{H}-\text{C}_8\text{H}_6\text{O}_3$] $^-$, which was substantially identical with those reported previously, and was tentatively identified as 2',4-dihydroxy-4'-methoxy chalcone (Figure 2).




Figure 1. Representative chromatograms from the analysis of OMDC extracts. (A) and (B): extracted ion chromatogram (EIC) of reference standards; (C): positive ion mode; (D): negative ion mode) total ion chromatogram (TIC) of OMDC extract.




Figure 2. Structures of compounds identified from OMDC.




Figure 3. Proposed fragmentation pathways of 2',4'-dihydroxychalcone.

Table 1. Chromatographic and mass spectrometric data (negative ion) of compounds identified from OMDC using UHPLC-QTOF-MS/MS.

No.	Formula	[M-H] ⁻	Error (ppm)	tR (min)	MS/MS (Characteristic Product Ions)	Identity	Intensity
1	C ₁₅ H ₁₂ O ₅	271.0612	-0.9	8.17	243.0656, 227.0728, 185.0579, 164.0089, 136.0160, 109.0292, 91.0215, 65.0417	2',3,4,4'-Tetrahydroxy-chalcone	7488
2	C ₁₅ H ₁₂ O ₆	287.0561	-0.7	12.52	269.0428, 259.0615, 243.0683, 203.0330, 173.0608, 151.0031, 125.0255	(-)-dihydrokaempferol	3122
3	C ₁₅ H ₁₄ O ₅	273.0769	-4	12.87	255.0650, 243.0622, 109.0287, 253.0493, 243.0659, 227.0566, 13.0819,	2,4,2',5' -Tetrahydrodihydrochalcone	1984
4	C ₁₅ H ₁₂ O ₅	271.0612	-0.7	12.94	164.0107, 136.0168, 109.0303, 91.0195, 67.0190	2',3,5,4'-Tetrahydroxy-chalcone	21,430
5	C ₁₅ H ₁₀ O ₅	269.0456	-0.9	13.75	241.0515, 225.0545, 197.0611, 181.0653, 135.0096, 91.0205	2',5,7-Trihydroxyflavone	5705
6	C ₂₁ H ₂₂ O ₁₀	433.11402	3.2	13.84	271.0590, 165.0192	Naringenin-glucose	1139
7	C ₁₅ H ₁₀ O ₇	301.0354	-0.6	14.08	283.0248, 255.0286, 215.0340, 151.0003, 145.9294	Quercetin; 2-(3,4-Dihydroxyphenyl)-3,5,7-trihydroxy-4H-chromen-4-one	3244
8	C ₁₅ H ₁₄ O ₅	273.0769	-1.4	14.12	227.0768, 167.0349, 149.0244, 137.0246, 123.0453, 109.0302	2',4',6',4'-Tetrahydroxydihydrochalcone	3608
9	C ₁₅ H ₁₂ O ₆	287.0561	-0.5	14.79	269.0435, 259.0605, 243.0667, 201.0534, 177.0558, 151.0031, 125.0242, 83.0133, 63.0250	Dihydrokaempferol; 3,5,7-trihydroxy-2-(4-hydroxyphenyl)-2,3-dihydrochromen-4-one	6907
10	C ₁₇ H ₁₆ O ₆	315.08741	-1.2	14.83	300.0646, 257.0497, 178.9552, 149.0271	3',5-Dihydroxy-4',7-Dimethoxyflavanone	1362

11	C ₁₆ H ₁₄ O ₆	301.07176	-0.7	15.1	286.0473, 283.0595, 271.0616, 255.0702, 191.0359, 179.0344, 176.0093, 164.9257, 151.0394, 135.0081, 121.0295	3',5,7-Trihydroxy-4'-Methoxyflavanone	4211
12	C ₁₅ H ₁₀ O ₅	269.0456	-0.7	15.26	251.0317, 241.0492, 225.0556, 181.0643, 149.0225, 135.0082, 91.0232	7,8,3'-Trihydroxyflavone	8460
13	C ₂₁ H ₂₂ O ₁₀	433.11402	0.7	15.42	271.0601, 165.0179	Naringenin-glucose	1470
14	C ₁₅ H ₁₄ O ₅	273.0769	-0.9	15.62	255.0670, 227.0683, 167.0347, 148.0172, 2,4,2',5'-Tetrahydrodihydrochalcone 137.0131, 123.0455, 109.0286	5233	
15	C ₁₇ H ₁₆ O ₆	315.08741	-0.8	16.23	300.0618, 257.0438, 178.9567, 149.0228, 121.0309	4',6-Dihydroxy-5,7-Dimethoxyflavanone	1794
16	C ₁₆ H ₁₄ O ₆	301.07176	-0.9	16.39	283.0612, 271.0613, 225.0554, 179.0352, 151.0418, 136.0136, 121.0305, 93.0353	7-methoxy-3',4',5-trihydroxyflavanone	4732
17	C ₁₅ H ₁₀ O ₆	285.0405	-0.8	16.9	257.0449, 217.0506, 199.0390, 175.0396, 149.0229, 133.0292, 83.0140, 65.0201	Luteolin; 3',4',5,7-Tetrahydroxyflavone	42,838
18	C ₁₅ H ₁₂ O ₄	255.0663	-1.4	17.65	135.0088, 119.0504, 91.0198	Liquiritigenin; 4',7-dihydroxyflavanone	114,322
19	C ₁₅ H ₁₀ O ₆	285.0405	-0.8	17.71	257.0504, 271.0504, 199.0396, 175.0391, 2',4',5,7-Tetrahydroxyisoflavone 149.0239, 133.0293, 83.0159, 65.0047	56,144	
20	C ₁₆ H ₁₄ O ₆	301.07176	-1.3	18.06	191.0353, 176.0103, 164.9285, 148.0148, 109.0297, 67.0208	3',4',7-Trihydroxy-5-methoxyflavanone	2937
21	C ₉ H ₈ O ₂	147.04515	-0.8	18.3	103.0556, 77.0391, 61.9902	Cinnamic acid	8302
22	C ₁₆ H ₁₂ O ₅	283.0612	-0.5	18.42	268.0377, 239.0341, 215.0326, 211.0400, 195.0450, 184.0531, 147.0451, 112.9849, 61.9902	6,7-dihydroxy-5-methoxyflavone	9831
23	C ₁₅ H ₁₂ O ₄	255.0663	-1.3	18.64	237.0560, 209.0608, 199.0761, 167.0861, 149.0247, 135.0090, 109.0301, 91.0206	6,7-dihydroxyflavanone	225,821
24	C ₁₆ H ₁₂ O ₅	283.0612	-1	18.73	268.0376, 239.0345, 211.0394, 184.0519, 148.0168, 135.0089	5,7-dihydroxy-4'-Methoxy isoflavones	91,281
25	C ₁₆ H ₁₆ O ₅	287.0925	-1.4	18.91	272.0683, 257.0439, 163.0405, 150.0321, 135.0439, 121.0306, 109.0280, 91.0580	7,2',3'-Trihydroxy-4'-methoxyisoflavane	4783
26	C ₂₁ H ₂₂ O ₉	417.1191	0.6	18.96	255.0659, 177.0147, 151.0057	Pinocembrin-glucoside	4369
27	C ₁₆ H ₁₂ O ₅	283.0612	-0.4	19.22	268.0375, 239.0344, 211.0399, 184.0526, 148.0168, 135.0089	4',7-Dihydroxy-2'-methoxyisoflavan	131,630
28	C ₂₁ H ₂₂ O ₉	417.1191	0.4	19.47	255.0669, 177.0191, 151.0029	Pinocembrin-glucoside	4277
29	C ₁₆ H ₁₆ O ₅	287.0925	-0.4	19.48	272.0690, 257.0435, 163.0397, 150.0317, 135.0463, 121.0292, 109.0279, 91.0615	7,2',3'-Trihydroxy-6-Methoxy isoflavane	5798
30	C ₁₅ H ₁₂ O ₄	255.0663	-1.1	19.52	237.0610, 209.0610, 199.0765, 165.0709, 135.0089, 109.0302, 91.0202	3',6-dihydroxyflavanone	279,373
31	C ₁₆ H ₁₄ O ₄	269.0819	-1	19.75	253.0522, 163.0401, 148.0178, 135.0084, 7-hydroxy-4'-methoxy dihydroflavone 119.0499, 109.0299, 91.0224	7,3',4'-trihydroxy-5-methoxy flavone	36,522
32	C ₁₆ H ₁₂ O ₆	299.0561	-1	19.96	109.0297, 65.0433	7,3',4'-trihydroxy-5-methoxy flavone	13,371
33	C ₁₆ H ₁₄ O ₄	269.0819	-0.6	20.26	253.0499, 225.0541, 163.0397, 148.0158, 135.0085, 119.0508, 109.0294, 91.0188	2',4'-Dihydroxy-2-methoxychalcone	46,597
34	C ₂₁ H ₂₀ O ₁₀	431.09837	0.1	20.32	269.0456, 226.9665, 158.9784, 140.9788	Apigenin-glucoside	1697
35	C ₂₁ H ₂₂ O ₉	417.1191	0.7	20.33	255.0658, 145.0644	Pinocembrin-glucoside	9326
36	C ₁₆ H ₁₂ O ₆	299.0561	-0.4	20.45	109.0299, 65.0411	2',4',5-Trihydroxy-7-methoxyisoflavanone	14,610
37	C ₂₁ H ₂₀ O ₁₀	431.09837	1.5	20.62	269.0439, 223.0971, 140.9806	Apigenin-glucoside	1657
38	C ₂₁ H ₂₂ O ₉	417.1191	0.9	20.76	255.0659, 211.0790, 171.0418, 151.0031	Pinocembrin-glucoside	10,565

39	C ₁₆ H ₁₆ O ₅	287.0925	-0.7	20.83	257.0818, 239.0698, 224.0470, 136.0171, 109.0297, 91.0210 253.0506, 225.0542, 215.0712, 197.0600,	7,2',3'-Trihydroxy-4'-methoxyisoflavane	11,412
40	C ₁₅ H ₁₂ O ₅	271.0612	-0.4	20.84	177.0190, 161.0607, 151.0033, 119.0504, 107.0144, 93.0352	Naringenin; 4',5,7-trihydroxyflavanone	196,476
41	C ₁₆ H ₁₂ O ₅	283.0612	-0.8	21.16	268.0372, 239.0360, 211.0406, 184.0529, 146.9650, 135.0069, 61.9916	3'-methoxy-5,7-dihydroxyflavone	9696
42	C ₁₆ H ₁₆ O ₅	287.0925	-0.7	21.26	257.0820, 239.0712, 224.0484, 136.0170, 109.0298, 91.0204 253.0507, 225.0548, 215.0706, 197.0598,	7,2',3'-Trihydroxy-4'-methoxyisoflavane Naringenin chalcone;	13,087
43	C ₁₅ H ₁₂ O ₅	271.0612	-0.7	21.32	185.0599, 177.0191, 161.0605, 151.0038, 119.0508, 107.0143, 93.0354, 63.0268	2',4,4',6'-tetrahydroxy-chalcone	300,571
44	C ₁₆ H ₁₂ O ₅	283.0612	-0.7	21.53	268.0375, 239.0348, 211.0406, 195.0449, 6,4'-dihydroxy-7-methoxyflavone 184.0515, 146.9653, 135.0073, 61.9892	6,4'-dihydroxy-7-methoxyflavone	11,665
45	C ₁₅ H ₁₀ O ₅	269.0456	-1	21.83	225.0554, 201.0550, 181.0660, 151.0031, 149.0242, 117.0347, 107.0154, 87.0472	Apigenin; 4',5,7-Trihydroxyflavone	48,246
46	C ₁₇ H ₁₆ O ₆	315.08741	-0.2	21.84	297.0389, 269.0453, 109.0294, 65.0404	5,4'-dihydroxy-7,3'-dimethoxy-flavanone	6428
47	C ₁₆ H ₁₂ O ₇	315.051	0.2	22.08	297.0397, 269.0450, 254.0218, 226.0249, 165.0199, 109.0293, 65.0390	Rhamnetin	4246
48	C ₁₅ H ₁₂ O ₅	271.0612	-0.8	22.1	253.0502, 227.0706, 185.0606, 151.0034, 143.0499, 107.0138, 83.0146, 65.0044	7,3',5'-trihydroxyflavanone	46,118
49	C ₁₇ H ₁₆ O ₆	315.08741	-0.6	22.14	297.0401, 269.0448, 254.0230, 226.0184, 3',4'-Dihydroxy-6,7-methoxyflavanone 198.0315, 165.0202, 109.0293, 65.0433	3',4'-Dihydroxy-6,7-methoxyflavanone	6592
50	C ₁₅ H ₁₂ O ₅	271.0612	-0.8	22.43	253.0500, 227.0706, 185.0601, 151.0030, 143.0495, 107.0143, 83.0157, 65.0053	3,7,4'-Trihydroxyflavanone	55,627
51	C ₁₆ H ₁₂ O ₆	299.0561	-0.5	23.26	284.0323, 255.0261, 227.0338, 211.0398, 5,7,2'-trihydroxy-4'-methoxy flavone 148.0180, 91.0200	5,7,2'-trihydroxy-4'-methoxy flavone	8908
52	C ₁₅ H ₁₄ O ₄	257.0819	-1.1	23.56	163.0400, 151.0405, 135.0088, 107.0511, 93.0358, 65.0414	2',4',4'-Trihydroxydihydrochalcone	218,604
53	C ₁₅ H ₁₀ O ₃	237.0557	-1.2	23.69	208.0531, 193.0658, 180.0573, 165.0721, 132.0213, 91.0209	7-Hydroxyflavone	52,893
54	C ₁₆ H ₁₂ O ₄	267.0663	-0.3	24.19	252.0418, 223.0387, 196.0520, 168.0583, 7-methoxy-4'-hydroxyisoflavone 135.0085, 117.0344, 91.0192	7-methoxy-4'-hydroxyisoflavone	26,052
55	C ₁₅ H ₁₀ O ₃	237.0557	-2.4	24.54	208.0515, 193.0641, 180.0579, 135.0094, 117.0348, 91.0268	6-Hydroxyflavone	10,396
56	C ₁₆ H ₁₄ O ₅	285.07685	-1	24.56	267.0644, 163.0393, 135.0452, 121.0306, Vestitone; 2',7-dihydroxy-4'-methoxyisoflavanone 109.03222, 91.0673	2',7-dihydroxy-4'-methoxyisoflavanone	5450
57	C ₁₆ H ₁₀ O ₅	281.0456	-0.9	24.73	253.0504, 224.0477, 209.0602, 195.0446, 167.0493, 135.0088, 117.0337, 91.0201	Pseudobaptigenin	158,719
58	C ₂₁ H ₂₂ O ₈	401.1242	0.8	24.76	239.0712, 197.0607, 135.0087, 112.9851	2',4'-Dihydroxychalcone-glucose	13,190
59	C ₁₅ H ₁₂ O ₄	255.0663	-1.2	25.06	135.0093, 119.0512, 91.0204	Isoliquiritigenin; 2',4,4'-Trihydroxychalcone	518,486
60	C ₁₆ H ₁₂ O ₆	299.0561	-0.9	25.39	284.0315, 256.0351, 165.0191, 149.9952, 121.0293, 65.0262	3'-Methoxy-4',5,7-trihydroxyflavanone	39,790
61	C ₁₆ H ₁₂ O ₄	267.0663	-1.2	25.45	252.0424, 223.0391, 208.0522, 195.0442, 167.0485, 132.0214, 91.0181	Formononetin; 7-Hydroxy-4'-methoxyisoflavanone	101,086
62	C ₁₅ H ₁₂ O ₃	239.0714	-2.3	25.85	197.0603, 169.0658, 148.0164, 135.0090, 109.0302, 91.0207	7-Hydroxyflavanone	1,785,720

63	C ₁₆ H ₁₄ O ₄	269.0819	-0.8	25.9	225.0899, 148.0184, 119.0510 237.0560, 209.0605, 193.0657, 169.0661,	4,4'-dihydroxy-2'-methoxychalcone (2R)-Pinocembrin; (2R)-5,7-dihydroxy-2-phenyl-2,3-dihydrochromen-4-one	20,639
64	C ₁₅ H ₁₂ O ₄	255.0663	-1.1	26.12	145.0294, 135.0088, 119.0503, 109.0302, 91.0201		202,511
65	C ₁₆ H ₁₂ O ₆	299.0561	-0.7	26.31	284.0326, 255.03040, 227.0347, 211.0384, 148.0152,	3',4',7-Trihydroxy-5-methoxyflavone	12,184
66	C ₁₅ H ₁₀ O ₄	253.0506	-0.8	26.78	223.0384, 208.0526, 195.0450, 180.0582, 152.0633, 132.0422, 116.9286, 92.0318	4',7-Dihydroxyisoflavone	26,456
67	C ₁₇ H ₁₈ O ₅	301.1082	-0.9	27.07	286.0853, 271.0602, 179.0728, 164.0482, 7,3 '-dihydroxy-2', 4 '-dimethoxy isoflavane		5886
68	C ₁₆ H ₈ O ₆	295.02481	-0.8	27.88	267.0292, 266.0218, 239.0352, 211.0407, 195.0285, 158.9768, 114.9876	Medicagol	2425
69	C ₁₅ H ₁₀ O ₄	253.0506	-0.9	29.23	225.0561, 209.0610, 181.0659, 143.0504, 119.0504, 107.0147, 63.0270	Chrysin; 5,7-Dihydroxyflavone	872,015
70	C ₁₆ H ₁₄ O ₅	285.07685	-0.1	29.5	269.0472, 165.0191, 119.0505, 97.0290, 89.0041, 65.0129	Sakuranetin, 4',5-Dihydroxy-7-methoxyflavanone	52,887
71	C ₁₅ H ₁₂ O ₄	255.0663	-1.3	29.68	227.0708, 213.0549, 211.0762, 185.0603, 171.0447, 169.0655, 151.0035, 145.0659, 107.0146, 83.0160, 65.0068	Pinocembrin; 5,7-Dihydroxyflavanone	2,838,626
72	C ₁₅ H ₁₂ O ₅	271.0612	-2	29.72	253.0504, 243.0659, 227.0706, 185.0603, 173.0606, 152.0107, 124.0162, 95.0139, 65.0050	7,3',4'-Trihydroxyflavanone	45,968
73	C ₁₅ H ₁₀ O ₃	237.0557	-0.3	29.79	208.0519, 193.0671, 180.0560, 165.0724, 135.0106, 107.0156, 91.0193, 65.0103	5-Hydroxyflavone	13,845
74	C ₁₅ H ₁₂ O ₃	239.0714	-1.7	30.26	211.0796, 197.0601, 195.0805, 169.0653, 148.0164, 135.0089, 109.0305, 91.0203, 65.0057	2',5'-Dihydroxychalcone	74,066
75	C ₁₅ H ₁₀ O ₅	269.0456	-0.1	30.44	252.0426, 239.0369, 224.0454, 200.8814, 169.0643,	3',4',5-Trihydroxyflavon	7107
76	C ₁₆ H ₁₄ O ₅	285.07685	-1.9	30.57	267.0656, 145.0293, 139.0398, 124.0168, 96.0229	3',4'-Dihydroxy-5'-methoxyflavanone	63,776
77	C ₁₅ H ₁₄ O ₄	257.0819	-2.2	30.73	239.0721, 213.0925, 195.0814, 173.0610, 151.0039, 122.0377, 107.0153, 81.0367, 65.0043	2',4',6'-Trihydroxydihydrochalcone	127,231
78	C ₁₆ H ₈ O ₆	295.02481	-0.8	30.79	267.0293, 266.0216, 237.0185, 211.0393, 135.0100	Medicagol; 7-Hydroxy-11,12-methylenedioxycoumestan	23,276
79	C ₁₆ H ₁₂ O ₆	299.0561	-0.6	30.99	284.0320, 271.0610, 256.0367, 240.0415, 178.0270, 165.0190, 151.0033, 122.0014, 93.0355, 65.0054	Rhamnocitrin; 3,4',5-Trihydroxy-7-methoxyflavone	64,943
80	C ₁₅ H ₁₄ O ₃	241.087	-1.2	33.52	223.0762, 197.0970, 150.0321, 135.0091, 122.0380, 109.0307, 91.0209, 65.0436	2',4'-Dihydroxydihydrochalcone	2,684,260
81	C ₁₆ H ₁₄ O ₄	269.0819	-1.2	33.69	225.0916, 148.0151, 119.0507, 93.0337	2',4-dihydroxy-4'-Methoxy chalcone	108,605
82	C ₁₅ H ₁₂ O ₃	239.0714	-1.4	33.95	211.0761, 197.0600, 195.0812, 169.0662, 148.0169, 135.0090, 109.0308, 91.0210, 65.0074	2',4'-Dihydroxychalcone	3,251,005
83	C ₁₅ H ₁₀ O ₅	269.0456	-0.5	35.22	241.0498, 225.0544, 197.0596, 181.0645	3',4',7-Trihydroxyflavone	9165

84	$C_{16}H_{14}O_4$	269.0819	-1.2	35.86	254.0574, 226.0620, 198.0652, 165.0185, 5-hydroxy-7-methoxy di- 149.9948, 122.0016, 65.0080 hydroflavone	5383
85	$C_{16}H_{16}O_4$	271.09758	-0.2	36.11	256.0734, 165.0207, 152.0110, 139.0396, 4,5-Dihydroxy-2-Methoxy 124.0166 dihydrochalcone	7714
86	$C_{16}H_{16}O_4$	271.09758	-1.4	36.66	256.0752, 253.0872, 165.0202, 151.0116, 4,5-Dihydroxy-2-Methoxy 139.0392, 124.0158 dihydrochalcone	7147
87	$C_{16}H_{14}O_4$	269.0819	-0.9	38.02	254.0580, 226.0628, 177.0184, 165.0196, 5-hydroxy-7-methoxy di- 149.9946, 122.0012, 65.0046 hydroflavone	40,090
88	$C_{16}H_{14}O_5$	285.07685	-0.9	38.02	270.0547, 165.0202, 145.0286, 139.0408, 3',4'-Dihydroxy-5'-meth- 93.0353 oxyflavanone	3772
89	$C_{20}H_{20}O_4$	323.1289	-0.7	39.75	255.0667, 203.0706, 159.0805, 119.0506, isobavachin; 4',7-dihy- 93.0361 droxy-8-prenylflavanone	25,579
90	$C_{16}H_{14}O_5$	285.07685	-0.9	40.66	267.0642, 241.0865, 176.0101, 148.0137, 3',4'-Dihydroxy-6-meth- 109.0191 oxyflavanone	3395
91	$C_{20}H_{20}O_4$	323.1289	-0.4	43.34	305.1204, 277.1659, 255.0628, 219.0698, 4,2',4'-trihydroxy-3'-iso- 186.9312 pentenyl chalcone	2040
92	$C_{16}H_{14}O_6$	301.07176	-0.8	52.85	283.0601, 257.0847, 192.0055, 173.0594, 4',5,7-trihydroxy-3'-meth- 164.0103 oxyflavanone	9203

3.2. Screening and Identification of Flavone

The fragmentation of $A^{1,3-}$ and $B^{1,4-}$ was formed by Retro-Diels-Alder (RDA) reaction on the C ring of flavone (Figure 4) and flavonol aglycones. Furthermore, isoflavone aglycones appeared specific fragmentations of $B^{0,3-}$ and $A^{1,3-}$, which derived from the C ring's breakage (Figure 4). The reason may be that the various conjugated systems belong to flavone and isoflavone aglycones influenced the fragmentation behavior of C ring. They also tend to lose some small neutral molecules, for example CO, CHO, C_3O_2 , CO_2 , C_2H_2O , etc. The ion fragments of some compounds are listed in this work.




Figure 4. RDA collision of flavone and isoflavone aglycones at C ring (A, B, and C stand for different six-membered rings).

Compound 53 had a deprotonated molecular ion at m/z 238.0629 and its molecular formula was determined to be $C_{15}H_{10}O_3$. This compound generated fragment ions at m/z 208.0531, 193.0658, 180.0573, 165.0721, and 91.0209, and was tentatively identified as 7-hydroxyflavone (Figure 3) [22]. The molecular formulae for 69, 45, and 17 were $C_{15}H_{10}O_4$, $C_{15}H_{10}O_5$, and $C_{15}H_{10}O_6$, respectively, and the compounds were identified as chrysins, apigenin, and luteolin, respectively, by DPLs (chrysins: m/z 225.0561 [$M-H-CO$] $^-$, 209.0610 [$M-H-CO_2$] $^-$, 151.0034 [$M-H-C_8H_6$] $^-$, 143.0504 [$M-H-C_6H_6O_2$] $^-$, 107.0147 [$M-H-C_9H_6O_2$] $^-$; apigenin: m/z 225.0554 [$M-H-CO_2$] $^-$, 151.003 [$M-H-C_8H_6O$] $^-$, 117.0347 [$M-H-C_7H_4O_4$] $^-$, 107.0154 [$M-H-C_9H_6O_3$] $^-$; luteolin: m/z 257.0048 [$M-H-H_2O$] $^-$, 241.0504 [$M-H-CO_2$] $^-$, 217.0504 [$M-H-CO_2-C_2H_2$] $^-$, 149.0239 [$M-H-C_7H_4O_3$] $^-$, 133.0293 [$M-H-C_7H_4O_4$] $^-$) and NL [44 Da (CO_2) and

68 Da (C_4H_4O]), the results were consistent with those reported in the literature [24,25]. (Compound **61**) The quasi-molecular ions acquired at m/z 267.0670 were 14 Da (CH_2) more than chrysin and the compound was determined to be formononetin via the DPIs (252.0424 [$M-H-CH_3$] $^-$, 223.0391 [$M-H-CO_2$] $^-$, 195.0442 [$M-H-CO_2-CO$] $^-$, 132.0214 [$M-H-C_7H_3O_3$] $^-$, 91.0181 [$M-H-CH_3-C_9H_5O_3$] $^+$) [26]. Compound **24** was screened in OMDC extracts based on the precursor ions at m/z 283.0609 and the chemical formula was computed as $C_{16}H_{12}O_5$. The fragment ions (m/z 268.0375 [$M-H-CH_3$] $^-$, 239.0344 [$M-H-CO_2$] $^-$, 184.0526 [$M-H-C_4H_4O_3$] $^-$, 148.0526 [$M-H-C_7H_3O_3$] $^-$, 135.0089 [$M-H-C_9H_8O_2$] $^-$) of these compounds were the same as those reported in the literature [4,27]. Compound **24** was, therefore, tentatively deduced to be 5,7-dihydroxy-4'-methoxy isoflavones (Figures 3 and 5). The molecular formula for compound **51** was determined to be $C_{16}H_{12}O_6$ based on the accurate mass of the quasi-molecular ions at m/z 299.0558. This organic compound was confirmed to be 5,7,2'-trihydroxy-4'-methoxy flavone (Figure 3) by the DPI m/z 284.0323, 255.0261, 227.0338, 211.0398, 148.0180 [2]. Compound **57** was determined to be pseudobaptigenin based on the parent ions at m/z 281.0455, and its molecular formula was deduced as $C_{16}H_{10}O_5$. It generated fragment ions at m/z 253.0504 [$M-H-CO_2$] $^-$, 225.0557 [$M-H-CO_2-CO$] $^-$, 135.0088 [$M-H-C_9H_6O_2$] $^-$, and 91.0201 [$M-H-C_{10}H_6O_4$] $^-$, and its structure was consistent with that reported in the literature [28,29].




Figure 5. Proposed fragmentation pathways of 5,7-dihydroxy-4'-methoxy isoflavones.

Compounds **71** and **18** yielded parent ions at m/z 255.0656, corresponding to the formula $C_{15}H_{12}O_4$. Compound **71** was determined to be pinocembrin based on its chromatographic data and comparison with the fragment ions with standard pinocembrin. Compounds **26**, **28**, and **38** were determined to be pinocembrin derivatives based on their DPIs acquired at m/z 211.0790, 171.0418, and 151.0031. These compounds were tentatively classified as pinocembrin-glucoside due to the DPIs and NL 162 Da ($C_6H_{10}O_5$). Compound **18** and isoliquiritigenin were structural isomers. It was found to be liquiritigenin by the DPIs acquired at m/z 135.0093 [$M-H-C_8H_8O$] $^-$, 119.0512 [$M-H-C_7H_4O_3$] $^-$, and 91.0204 [$M-H-C_9H_8O_3$] $^-$. These compounds were similar, as reported in the literature [30]. Two compounds (**31** and **84**) had a deprotonated molecular ion at m/z 269.0815 in the MS data, which matched with the chemical formula $C_{16}H_{14}O_4$. Based on the DPIs (m/z 163.0397 [$M-H-C_6H_5O$] $^-$, 148.0151 [$M-H-C_6H_3O_2$] $^-$, 135.0085 [$M-H-C_8H_8O$] $^-$, 119.0507 [$M-H-C_7H_4O_3$] $^-$, and 91.0188 [$M-H-C_9H_8O_3$] $^-$) of the compound **31** was identified as 7-hydroxy-4'-methoxy dihydroflavone (Figure 6) [31], and compound **84** was identified as 5-hydroxy-7-methoxy dihydroflavone by DPIs (m/z 165.0196, 149.9946, 122.0012, and 65.0046) [32]. Compound **89** had a mass of 68 Da (C_5H_8), more than pinocembrin and its chemical formula was calculated to be $C_{20}H_{20}O_4$. This compound was provisionally designated as isobavachin based on DPIs (m/z 255.0667 [$M-H-C_5H_8$] $^-$, 203.0706 [$M-H-C_6H_3O_2$] $^-$, 159.0805 [$M-H-C_9H_8O_3$] $^-$, 119.0506 [$M-H-C_5H_8-C_7H_3O_3$] $^-$, and 93.0361 [$M-H-C_{14}H_{14}O_3$] $^-$) [33].




Figure 6. Proposed fragmentation pathways of 7-hydroxy-4'-methoxy dihydroflavone.

3.3. Screening and Identification of Flavonol

The molecular formula of compound **7** was calculated as $C_{15}H_{10}O_7$, and the ion fragments that were obtained were mainly m/z 283.0248 [$M-H-H_2O$] $^-$, 255.0286 [$M-H-H_2O-CO$] $^-$, and 151.0003 [$M-H-C_8H_6O_3$] $^-$. Based on the results that were consistent with those reported in the literature, the compound was identified as quercetin [34]. Compound **79** had a deprotonated molecular ion at m/z 299.0565 and its molecular formula was calculated as $C_{16}H_{12}O_6$. These fragments (m/z 284.0320 [$M-H-CH_3$] $^-$, 271.0610 [$M-H-CO$] $^-$, 165.0190 [$M-H-C_8H_6O_2$] $^-$, 151.0033 [$M-H-CH_3-C_8H_6O_2$] $^-$, and 93.0355 [$M-H-C_{10}H_6O_5$] $^-$) were determined to be those of rhamnocitrin (Figures 3 and 7), which was consistent with that reported in the literature [35]. Compound **9** was identified as dihydrokaempferol (Figure 8) by DPLs (m/z 259.0605 [$M-H-CO$] $^-$, 269.00435 [$M-H-H_2O$] $^-$, 177.0558 [$M-H-H_2O-C_6H_4O$] $^-$, and 151.0031 [$M-H-H_2O-C_8H_6O$] $^-$) and its molecular formula was $C_{15}H_{12}O_6$.




Figure 7. Proposed fragmentation pathways of rhamnocitrin.




Figure 8. Proposed fragmentation pathways of dihydrokaempferol.

3.4. Screening and Identification of Isoflavane

The molecular ion peak of compound **67** at *m/z* 301.1048 was determined based on quantitative data and its molecular formula was estimated to be C₁₇H₁₈O₅. The compound was confirmed as 5,7-dimethoxy-2',4'-dihydroxy isoflavan (Figures 3 and 9) by the DPIS at 286.0853[M-H-CH₃]⁻, 271.0602[M-H-CH₃-CH₃]⁻, 135.0451[M-H-C₉H₁₀O₃]⁻, and 109.0293[M-H-C₁₁H₁₂O₃]⁻ (Figure 2), and further confirmed by comparison with fracture modes reported in the literature [4,36]. The parent ion of Compound **39** was identified as *m/z* 287.0912 (C₁₆H₁₆O₅) and it had a mass of 14 Da (CH₂) less than that of compound **67**. The compound was provisionally identified as 7,2',3'-trihydroxy-4'-methoxy isoflavan (Figure 3) by DPIS (*m/z* 257.0820 [M-H-CH₃O]⁻, 239.0712 [M-H-CH₃O-H₂O]⁻, 136.0170[M-H-C₈H₇O₃]⁻, and 109.0298[M-H-C₁₀H₁₀O₃]⁻) [36].




Figure 9. Proposed fragmentation pathways of 5,7-dimethoxy-2',4'-dihydroxy isoflavan.

3.5. Screening and Identification of Indole Alkaloids

Compound **112** was an isomer with a quasi-molecular ion m/z 238.0867, which was consistent with the chemical formula of $C_{15}H_{11}NO_2$. It was determined to be 3-hydroxy-N-benzoyl indole based on the resulting DPIS (m/z 165.0705, 105.0342, 77.0397) [36]. The deprotonated molecular ions at m/z 254.0821 ($C_{15}H_{11}NO_3$) for two compounds (**93** and **98**) were 16 Da (O) more than that of 3-hydroxy-N-benzoyl indole. The structure of Compound **93** was deduced from the DPIS (165.0695, 121.0288, 107.0498) and speculated as 3-hydroxy-N-(3-hydroxybenzoyl) indole. Based on the DPIS at m/z 135.0449, Compound **98** was tentatively identified as 3-hydroxy-N-(p-hydroxybenzoyl) indole [36]. Compounds **108** had a mass of 30 Da (CH_2O) more than that of 3-hydroxy-N-benzoyl indole and was identified as 3-methoxy-N-(p-hydroxybenzoyl) indole by the fragmentation ions (m/z 253.0742, 121.0291, 93.0347) [36]. The precursor ion of Compound **118** was identified as m/z 322.1456, and had a mass of 84 Da (C_5H_8O) more than that of 3-hydroxy-N-benzoyl indole. It was provisionally identified as 3-hydroxy-N-(3-isopentenyl-4-hydroxybenzoyl) (Figure 10 and Table 2) indole on the basis of DPIS (m/z 266.0825 [$M+H-C_4H_7$]⁺, 238.0865 [$M+H-C_4H_8-CO$]⁺, 189.0913, 133.0289).




Figure 10. Proposed fragmentation pathways of 3-hydroxy-N-(3-isopentenyl-4-hydroxybenzoyl) indoles.

Table 2. Chromatographic and mass spectrometric data (positive ion) of compounds identified from OMDC using UHPLC-QTOF-MS/MS.

No.	Formula	$[M+H]^-$	Error (ppm)	tR (min)	MS/MS (Characteristic Product Ions)	Identity	Identity
93	$C_{15}H_{11}NO_3$	254.0812	0.5	21.29	236.0703, 208.0760, 165.0695, 121.0288, 107.0498, 93.0677, 65.0383	3-Hydroxy-N-(3-hydroxybenzoyl) indole	6243

94	C ₁₅ H ₂₆ O ₂	239.2006	-0.8	22.19	221.1896, 203.1798, 147.1171, 135.1166, 133.1010, 107.0862, 95.0863, 81.0710 236.0713, 208.0754, 165.0694, 121.0289, 107.0493, 93.0342, 65.0396 269.0690, 226.0628, 150.0299, 121.0280, 269.0685, 150.0309, 120.0442 236.0716, 208.0768, 181.0657, 165.0705, 135.0449, 121.0291, 107.0499, 93.0341, 65.0399 122.0602, 105.0342, 103.0553, 77.0399 236.0714, 209.0585, 181.0654, 165.0708, 153.0689, 135.0453, 121.0286, 107.0482, 93.0338, 65.0401 509.1637, 424.1442, 385.1068, 373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	3-Methyl-5-(1,3,3-trimethyl-7-oxabi-cyclo [2.2.1] hept-2-yl)-pent-1-en-3-ol	18,924
95	C ₁₅ H ₁₁ NO ₃	254.0812	0.4	22.85	165.0694, 121.0289, 107.0493, 93.0342, 65.0396 269.0690, 226.0628, 150.0299, 121.0280, 269.0685, 150.0309, 120.0442 236.0716, 208.0768, 181.0657, 165.0705, 135.0449, 121.0291, 107.0499, 93.0341, 65.0399 122.0602, 105.0342, 103.0553, 77.0399 236.0714, 209.0585, 181.0654, 165.0708, 153.0689, 135.0453, 121.0286, 107.0482, 93.0338, 65.0401 509.1637, 424.1442, 385.1068, 373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	3-Hydroxy-N-(3-hydroxybenzoyl) indole	33,513
96	C ₁₆ H ₁₃ NO ₄	284.0917	0.7	22.91	226.0628, 150.0299, 121.0280, 269.0685, 150.0309, 120.0442 236.0716, 208.0768, 181.0657, 165.0705, 135.0449, 121.0291, 107.0499, 93.0341, 65.0399 122.0602, 105.0342, 103.0553, 77.0399 236.0714, 209.0585, 181.0654, 165.0708, 153.0689, 135.0453, 121.0286, 107.0482, 93.0338, 65.0401 509.1637, 424.1442, 385.1068, 373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	3-Methoxy-4-hydroxy-n-(p-hydroxybenzoyl) indole	9202
97	C ₁₆ H ₁₃ NO ₄	284.0917	0.8	23.79	165.0705, 135.0449, 121.0291, 107.0499, 93.0341, 65.0399 122.0602, 105.0342, 103.0553, 77.0399 236.0714, 209.0585, 181.0654, 165.0708, 153.0689, 135.0453, 121.0286, 107.0482, 93.0338, 65.0401 509.1637, 424.1442, 385.1068, 373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	3-Methoxy-4-hydroxy-n-(p-hydroxybenzoyl) indole	165,711
98	C ₁₅ H ₁₁ NO ₃	254.0812	0.9	24.9	165.0705, 135.0449, 121.0291, 107.0499, 93.0341, 65.0399 122.0602, 105.0342, 103.0553, 77.0399 236.0714, 209.0585, 181.0654, 165.0708, 153.0689, 135.0453, 121.0286, 107.0482, 93.0338, 65.0401 509.1637, 424.1442, 385.1068, 373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	3-Hydroxy-N-(p-hydroxybenzoyl) indole	251,175
99	C ₁₅ H ₁₅ NO	226.1226	0.7	25.7	122.0602, 105.0342, 103.0553, 77.0399 236.0714, 209.0585, 181.0654, 165.0708, 153.0689, 135.0453, 121.0286, 107.0482, 93.0338, 65.0401 509.1637, 424.1442, 385.1068, 373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	N-Benzoyl-phenylethylamine	1,428,756
100	C ₁₅ H ₁₁ NO ₃	254.0812	0.9	26.75	122.0602, 105.0342, 103.0553, 77.0399 236.0714, 209.0585, 181.0654, 165.0708, 153.0689, 135.0453, 121.0286, 107.0482, 93.0338, 65.0401 509.1637, 424.1442, 385.1068, 373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	3-Hydroxy-N-(p-hydroxybenzoyl) indole	50,895
101	C ₃₂ H ₃₂ O ₉	561.2119	-1.3	27.96	373.1072, 259.0970, 167.0699, 123.0433, 107.0498 221.1925, 203.1806,	(3R)-Proptero-B-(α ,6)-(-)-isomucronulatol	4620
102	C ₁₅ H ₂₆ O ₂	239.2006	0.6	28.48	221.1925, 203.1806,	3-Methyl-5-(2,2,4-trimethylcyclohexanol-3-yl)-pent-1-ene-3-ol	18,924

103	$C_{16}H_{13}NO_4$	284.0917	1	28.57	161.1315, 147.1172, 119.0849, 95.0860, 81.0722 269.0685, 137.0245, 91.0530 131.0490, 122.0962, 105.0699, 103.0545, 79.0548 148.0760, 131.0492, 105.0699, 103.0543, 79.0550, 77.0395 220.0760, 165.0700, 121.0291, 91.0548 220.0758, 165.0698, 121.0289, 93.0336, 65.0405, 253.0742, 225.0792, 210.0675, 149.0602, 134.0366, 121.0291, 107.0496, 93.0347, 65.0397 253.0745, 150.0318, 121.0298, 105.0342, 77.0398 439.3568, 421.3458, 376.1913, 245.1907, 233.1914, 185.1312, 163.1505, 147.1145, 109.0989, 81.0724	3-Methoxy-4-hydroxy-n-(p-hydroxybenzoyl) indole	7924
104	$C_{17}H_{19}NO$	254.1539	0.8	29.84	N-Hydrocinnamoyl-2-phenylethylamine	113,125	
105	$C_{17}H_{17}NO$	252.1383	1.1	30.75	N-Cinnamoyl-2-phenylethylamine	5,724,222	
106	$C_{15}H_{11}NO_2$	238.0863	0.8	30.84	N-p-Hydroxybenzoyl indole	369,512	
107	$C_{15}H_{11}NO_2$	238.0863	0	31.83	N-p-Hydroxybenzoyl indole	52,164	
108	$C_{16}H_{13}NO_3$	268.0968	0.7	32.1	3-Methoxy-N-(p-hydroxybenzoyl) indole	520,562	
109	$C_{16}H_{13}NO_3$	268.0968	1.1	32.63	3-Methoxy-hydroxy-n-benzoyl indole	503,247	
110	$C_{30}H_{48}O_3$	457.3676	-1	33.09	Soyasapogenol E	13,542	
111	$C_{16}H_{13}NO_3$	268.0968	0.5	33.3	3-Methoxy-hydroxy-n-benzoyl indole	29,002	

112	$C_{15}H_{11}NO_2$	238.0863	1	33.57	150.0320, 134.0367, 121.0285, 105.0349, 93.0333, 77.0394, 65.0390 221.0604, 220.0762, 193.0655,	3-Hydroxy-N-benzoyl indole	1,270,624
113	$C_{16}H_{14}O_3$	255.1016	1.6	34.48	165.0705, 135.0442, 105.0342, 77.0397 240.0793, 209.0972, 194.0735, 177.0560,	2'-Hydroxy-4'-methoxychalcone	2,708,762
114	$C_{30}H_{46}O_4$	471.3469	-1	35.03	165.0707, 151.0395, 131.0500, 103.0553, 95.0504, 77.0400 453.3366, 435.3233, 395.2965, 199.1518, 173.1328, 159.1160, 145.0998, 97.0654 439.3575, 421.3461, 409.3465, 381.3164, 309.2588, 259.2056,	3,22,24-Trihydroxy- γ -lactone-olean-12-en-29-oic acid	15,709
115	$C_{30}H_{48}O_3$	457.3676	-0.7	35.18	245.1891, 233.1904, 205.1586, 145.1007, 135.1165, 119.0863, 81.0704 253.0736, 165.0544, 150.0313, 137.121.0291, 105.0341, 77.0393	Melilotigenin C	64,956
116	$C_{16}H_{13}NO_3$	268.0968	0.8	35.67	3-Methoxy-hydroxy-n-benzoyl indole	59,907	
117	$C_{20}H_{19}NO_3$	322.1438	1	35.79	266.0819, 238.0874,	3-Hydroxy-N-(3-isopentenyl-4-hydroxybenzoyl) indole	11,031

118	$C_{20}H_{19}NO_3$	322.1438	0.7	37.49	211.0753, 133.0322, 121.00286 266.0825, 254.0814, 211.0757, 189.0913, 165.0703, 133.0289, 105.0336, 77.0389 240.0783, 209.0972, 194.0725, 177.0551,	3-Hydroxy-N-(3-isopentenyl-4-hydroxybenzoyl) indole	261,939
119	$C_{16}H_{14}O_3$	255.1016	0.9	41.5	165.0699, 151.0392, 131.0492, 103.0545, 95.0498, 77.0385 453.3360, 435.3254, 423.3261, 395.2930, 287.2016, 259.1681, 247.1705, 201.1631, 189.1624, 147.1173, 109.1043, 95.0854 421.3458, 399.2707, 297.2570,	2'-Hydroxy-4'-methoxychalcone	129,251
120	$C_{30}H_{46}O_4$	471.3469	-0.6	42.6	215.1794, 173.1332, 135.1171, 109.1006 387.2866,	Unknown	17,953
121	$C_{30}H_{48}O_3$	457.3676	-0.9	45.42	325.1404, 233.1514, 148.0852 441.3717, 423.3615, 355.1895,	Unknown	54,664
122	$C_{30}H_{46}O_4$	471.3469	-5.7	45.93	335.2005, 247.2064, 203.1779, 163.0758,	24-Hydroxy-3-oxoolean-12-en-29-oic acid	52,681
123	$C_{30}H_{50}O_3$	459.3833	-0.5	46.13		Soyasapogenol B	8837

124	$\text{N}_{19}\text{H}_{21}\text{NO}_4$	366.2127	-1.7	46.43	131.0486, 81.0691 244.1758, 145.0256, 121.0996, 85.0651 457.2366, 321.1122,	3-Methoxyaegeline	64,028
125	$\text{C}_{30}\text{H}_{50}\text{O}_4$	475.3782	-0.5	46.52	267.0679, 219.1767, 179.0354 441.3741, 423.3649, 323.1291, 311.1293,	Wistariasapogenol B	12,829
126	$\text{C}_{30}\text{H}_{50}\text{O}_3$	459.3833	-0.2	47.04	219.0657, 203.1810, 161.1330, 135, 1163, 123.1181, 95.0873 441.3746, 323.1950, 311.1284,	Unknown	25,684
127	$\text{C}_{30}\text{H}_{50}\text{O}_3$	459.3833	-0.4	48.77	293.1184, 201.1654, 179.0698, 109.1012, 107.0856	Unknown	18,190

3.6. Identification of 22 Metabolites (M1–M22) in Mice

Since drug metabolism largely determines the pharmacokinetic characteristics and bioavailability of most drugs, in order to better understand the metabolic pathway of OMDC, *in vivo* metabolite analysis was carried out. To analyze the *in vivo* metabolites, a megadose of OMDC solution (150 mg/kg body weight) was administered orally to rats. Plasma samples from three animals at different sampling sites (1, 2, 4, 6, and 12 h) were mixed and gathered for LC/MS. In total, 22 metabolites from plasma were tentatively identified (Table 3) based on fragment ions. These compounds were metabolized by glucuronidation and sulfation (Figures 11 and 12).

Table 3. *In vivo* metabolites after the administration of OMDC to rats.

No.	Formula	[M-H] ⁻	Error (ppm)	tR (min)	MS/MS (Characteristic Product Ions)	Identity
M1	$\text{C}_{27}\text{H}_{28}\text{O}_{16}$	607.13046	3.3	15.94	431.0981, 255.0655	Di-glucuronidation of 71
M2	$\text{C}_{22}\text{H}_{20}\text{O}_{11}$	459.09329	0.4	16.3	283.0594, 268.0382	Glucuronidation of 24
M3	$\text{C}_{21}\text{H}_{20}\text{O}_{11}$	447.09329	-48	16.42	271.0603, 151.0036, 119.0506	Glucuronidation of 43
M4	$\text{C}_{27}\text{H}_{28}\text{O}_{15}$	591.13554	2.7	16.67	415.1037, 239.0711	Di-glucuronidation of 82
M5	$\text{C}_{27}\text{H}_{30}\text{O}_{15}$	593.15119	4.8	17.05	417.1143, 241.0861	Di-glucuronidation of 80
M6	$\text{C}_{21}\text{H}_{18}\text{O}_{10}$	429.08272	3.5	17.36	253.0505, 195.0454	Glucuronidation of 69
M7	$\text{C}_{21}\text{H}_{20}\text{O}_9$	415.10346	1.4	19.78	239.0709, 197.0611, 135.0085	Glucuronidation of 82
M8	$\text{C}_{21}\text{H}_{20}\text{O}_{12}\text{S}$	495.06027	5.2	21.37	319.0283, 239.0682	Sulfation and glucuronidation of 82
M9	$\text{C}_{21}\text{H}_{20}\text{O}_{13}\text{S}$	511.05519	1.1	22.12	431.0999, 255.0656	Sulfation and glucuronidation of 71

M10	$C_{21}H_{20}O_{10}$	431.09837	1.9	23.03	255.0667, 213.0552, 211.0769, 171.0448	Glucuronidation of 71
M11	$C_{22}H_{20}O_{12}$	475.0882	-61.9	23.27	299.0558, 284.0336, 255.0281, 227.0366	Glucuronidation of 51
M12	$C_{15}H_{12}O_8S$	351.01801	0.4	23.58	271.0618, 177.0213, 151.0031, 119.0517, 107.0156	Sulfation of 43
M13	$C_{22}H_{22}O_{10}$	445.11402	0.9	25.99	269.0817, 254.0572, 226.0632, 165.0196	Glucuronidation of 84
M14	$C_{21}H_{22}O_9$	417.11911	0.2	27.33	241.0865, 197.0986, 150.0330	Glucuronidation of 80
M15	$C_{16}H_{12}O_8S$	363.01801	1	27.59	283.0616, 268.0378	Sulfation of 24
M16	$C_{16}H_{10}O_8S$	361.00236	-1	28.34	281.0453, 253.0508,	Sulfation of 57
M17	$C_{16}H_{12}O_9S$	379.01293	0	29.92	299.0560, 284.0319, 165.0181	Sulfation of 51
M18	$C_{15}H_{12}O_6S$	319.02818	0	30.36	239.0707, 197.0598, 148.0168, 135.0084	Sulfation of 82
M19	$C_{16}H_{14}O_7S$	349.03875	-0.7	32.49	269.0813, 165.0194, 149.9945	Sulfation of 84
M20	$C_{15}H_{12}O_7S$	335.0231	-0.5	35.08	255.0653, 213.0538, 171.0463, 151.0000, 145.0642, 107.0139	Sulfation of 71
M21	$C_{15}H_{14}O_7S$	337.03875	0.3	40.05	257.0818, 239.0714, 151.0031	Sulfation of 52
M22	$C_{15}H_{14}O_6S$	321.04383	-0.8	41.09	241.0864, 197.0967, 150.0318, 135.0087	Sulfation of 80




Figure 11. Structures of identified metabolites of drugs.

Eight glucuronidated, 9 sulfated, 3 both glucuronidated and glucuronidated, and 2 both glucuronidated and sulfated metabolites were identified. M10 showed $[M-H]^-$ at m/z 431.0984 and its molecular formula was predicted as $C_{21}H_{20}O_{10}$. The fragment ion was formed by the neutral loss of 176.0326 Da ($C_6H_8O_6$) in the MS^2 spectra; thus, it was determined to be pinocembrin glucuronide. M20 showed $[M-H]^-$ at m/z 335.0231 and its molecular formula was predicted to be $C_{15}H_{12}O_7S$. Based on the fragment ion at m/z 255.0650, 171.0463, and 151.0000, the molecule was determined to be pinocembrin sulfate. M1 showed $[M-H]^-$ at m/z 607.1304 and its chemical formula was $C_{27}H_{28}O_{16}$. The fragment ion at m/z 255.0650, forecasted as $C_{15}H_{12}O_4$, was generated by the loss of two 176.0326 Da ($C_6H_8O_6$). Subsequently, the molecule was provisionally determined to be pinocembrin di-glucuronide. M9 showed $[M-H]^-$ at m/z 511.0552 and its chemical formula was $C_{21}H_{20}O_{13}S$. It was generated by a loss of 79.95 Da (SO_3) and 176.0326 Da ($C_6H_8O_6$). Hence, the metabolite was provisionally confirmed as pinocembrin glucuronide sulfate. Four

corresponding metabolites (M4, M7, M8, and M18) of 2',4'-dihydroxychalcone were detected (Figure 11). Other metabolites are also derived from flavonoids. As the major chemical components and major in vivo metabolites of OMDC, flavonoids may be crucial for the pharmacological effects of OMDC.




Figure 12. The EICs of 22 metabolites in rat plasma after administration of OMDC.

4. Conclusions

A rapid method using UHPLC-QTOF-MS was established for the isolation and authentication of the chemical constituents of OMDC. The compounds in OMDC extract were identified by NL and DPI sifting schema. Our results indicated that a total of 127 compounds, including 92 flavonoids, 15 indole alkaloids, and 20 others, could be identified in the OMDC extract. After the oral administration of OMDC extract to rats, 22 different compounds were found in the plasma, which appeared to be flavonoid metabolites. The primary in vivo metabolic reactions undergone by OMDC were glucuronidation and sulfation. This UHPLC-MS method is the first of its kind to determine the chemical constituents of OMDC in the positive and negative ion modes. Our findings revealed that the combination of UHPLC-MS and effective data mining is a logical, practical, and systematic method for the characterization of the chemical constituents and metabolites of OMDC.

Author Contributions: Conceptualization: Y.F. and H.O.; methodology: J.L.; validation: J.Z., N.P. and Y.Y.; formal analysis: M.X. and M.Y.; investigation: M.X. and M.Y.; writing—original draft preparation: M.X. and M.Y.; writing—review and editing: M.X., M.Y., H.O. and J.L.; supervision: S.Y. and G.Z.; project administration: Y.F. and H.O.; funding acquisition: Y.F. and H.O. All authors have read and agreed to the published version of the manuscript.

Funding: The study was supported by the National Key R and D Program of China (2019YFC1712302; Yulin Feng); Jiangxi University of Traditional Chinese Medicine 1050 youth talent project and Jiangxi University of Chinese Medicine Science and Technology Innovation Team Development Program (Hui Ouyang).

Institutional Review Board Statement: The animal experiments were approved by the Laboratory Animal Ethics Committee of Jiangxi University of Traditional Chinese Medicine (approval No. SYXK2017-0004).

Informed Consent Statement: Not applicable.

Data Availability Statement: All data related to the manuscript are available in the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Chen, C.; Liu, F.Q.; Zhao, J.Y.; Chen, T.; Li, Y.L.; Zhang, D.J. Efficient Separation of Five Flavonoids from *Oxytropis falcata* Bunge by High-Speed Counter-Current Chromatography and their Anticancer Activity. *Acta Chromatogr.* **2020**, *32*, 189–193.
2. Wang, W.D.; Dang, J.; Shao, Y.; Wang, Q.L.; Mei, L.J.; Tao, Y.D. Flavonoids from the poisonous plant *Oxytropis falcata*. *Chem. Nat. Compd.* **2019**, *55*, 6.
3. Chen, Z.P.; Qu, M.M.; Chen, H.X.; Liu, D.; Xiao, Y.Y.; Chen, J.; Lu, T.L.; Cai, B.C. The studies of anti-inflammatory and analgesic activities and pharmacokinetics of *Oxytropis falcata* Bunge extraction after transdermal administration in rats. *Fitoterapia* **2011**, *82*, 426–433.
4. Chen, W.H.; Wang, R.; Shi, Y.P. Flavonoids in the Poisonous Plant *Oxytropis falcata*. *J. Nat. Prod.* **2010**, *73*, 1398–1403.
5. Jiang, H.; Zhan, W.Q.; Liua, X.; Jiang, S.X. Antioxidant activities of extracts and flavonoid compounds from *Oxytropis falcata* Bunge. *Nat. Prod. Res.* **2008**, *22*, 1650–1656.
6. Wang, D.; Tang, W.; Yang, G.M.; Cai, B.C. Anti-inflammatory, Antioxidant and Cytotoxic Activities of Flavonoids from *Oxytropis falcata* Bunge. *Chin. J. Nat. Med.* **2010**, *8*, 461–465.
7. Gu, Q.L.; Huang, C.L.; Zhang, L.; Jiang, H. A Preliminary Study on Anti-inflammatory Effects of Rhamnocitrin from *Oxytropis falcata* Bunge. *Chin. J. Inf. Tradit. Chin. Med.* **2014**, *12*, 48–50.
8. Zhang, X.J.; Liu, W.X.; Wei, P.; Kelsang, N.; Que, S.; Zhang, L.; Zhang, Q.Y. Subacute toxicity and chemical analysis of Tibetan medicine *Oxytropis falcata*. *China J. Chin. Mater. Med.* **2014**, *39*, 1157–1162.
9. Jiang, H.; Hu, J.R.; Zhan, W.Q.; Liu, X. Screening for fractions of *Oxytropis falcata* Bunge with antibacterial activity. *Nat. Prod. Res.* **2009**, *23*, 953–959.
10. Yang, G.M.; Wang, D.; Tang, W.; Chen, X.; Fan, L.Q.; Zhang, F.F.; Yang, H.; Cai, B.C. Anti-inflammatory and Antioxidant Activities of *Oxytropis falcata* Fractions and Its Possible Anti-inflammatory Mechanism. *Chin. J. Nat. Med.* **2010**, *8*, 285–292.
11. Yang, L.X.; Wang, Z.C.; Jiang, L.G.; Sun, E.; Fan, Q.; Liu, T.H. Total Flavonoids Extracted from *Oxytropis falcata* Bunge Improve Insulin Resistance through Regulation on the IKK β /NF- κ B Inflammatory Pathway. *Hindawi Evid. Based Complement. Altern. Med.* **2017**, *2017*, 2405124.
12. Guo, Y.; Zhan, B.Y.; Peng, Y.F.; Chang, L.C.; Li, Z.Q.; Zhang, X.X.; Zhang, D.J. Mechanism of Action of Flavonoids of *Oxytropis falcata* on the Alleviation of Myocardial Ischemia-Reperfusion Injury. *Molecules* **2022**, *27*, 1706.
13. Wang, Y.; Liu, S.; Hu, Y.; Li, P.; Wan, J.B. Current state of the art of mass spectrometry-based metabolomics studies—A review focusing on wide coverage, high throughput and easy identification. *RSC Adv.* **2015**, *5*, 78728–78737.
14. Li, J.M.; Wen, Q.; Feng, Y.L.; Zhang, J.; Luo, Y.; Tan, T. Characterization of the multiple chemical components of Glechomae Herba using ultra high performance liquid chromatography coupled to quadrupole-time-of-flight tandem mass spectrometry with diagnostic ion filtering strategy. *J. Sep. Sci.* **2019**, *42*, 1312–1322.
15. Rodriguez, A.M.; Gurny, R.; Veuthey, J.L.; Guillarme, D. Coupling ultra-high-pressure liquid chromatography with mass spectrometry: Constraints and possible applications. *J. Chromatogr. A* **2013**, *1292*, 2–18.
16. Garran, T.A.; Ji, R.F.; Chen, J.L.; Xie, D.M.; Guo, L.P.; Huang, L.Q.; Lai, C.J.S. Elucidation of metabolite isomers of *Leonurus japonicus* and *Leonurus cardiaca* using discriminating metabolite isomerism strategy based on ultra-high performance liquid chromatography tandem quadrupole time-of-flight mass spectrometry. *J. Chromatogr. A* **2019**, *1598*, 141–153.
17. Guillarme, D.; Schappeler, J.; Rudaz, S.; Venthey, J.L. Coupling ultrahigh-pressure liquid chromatography with mass spectrometry. *TrAC Trend Anal. Chem.* **2010**, *29*, 15–27.
18. Ouyang, H.; Li, J.M.; Wu, B.; Zhang, X.Y.; Li, Y.; Yang, S.L.; He, M.Z.; Feng, Y.L. A robust platform based on ultra-high performance liquid chromatography quadrupole time of flight tandem mass spectrometry with a two-step data mining strategy in

the investigation, classification, and identification of chlorogenic acids in *Ainsliaea fragrans* Champ. *J. Chromatogr. A* **2017**, *1502*, 38–50.

19. He, M.Z.; Jia, J.; Li, J.M.; Wu, B.; Huang, W.P.; Liu, M.; Li, Y.; Yang, S.L.; Ouyang, H.; Feng, Y.L. Application of characteristic ion filtering with ultra-high performance liquid chromatography quadrupole time of flight tandem mass spectrometry for rapid detection and identification of chemical profiling in *Eucommia ulmoides* Oliv. *J. Chromatogr. A* **2018**, *1554*, 81–91.

20. Xue, Z.Z.; Lai, C.J.S.; Kang, L.P.; Kotani, A.; Hakamata, H.; Jing, Z.X.; Li, H.; Wang, W.H.; Yang, B. Profiling and isomer recognition of phenylethanoid glycosides from *Magnolia officinalis* based on diagnostic/holistic fragment ions analysis coupled with chemometrics. *J. Chromatogr. A* **2019**, *1611*, 460583.

21. Oshimura, M.Y.; Sano, A.T.; Amei, J.U.C.K.; Bata, A.K.O. Identification and Quantification of Metabolites of Orally Administered Naringenin Chalcone in Rats. *J. Agric. Food Chem.* **2009**, *57*, 6432–6437.

22. Norbo, K.; Padro, Q.K.; Zhang, X.J.; Samnor; Liang, H.; Zhang, Q.Y. High-performance liquid chromatographic fingerprint analysis of *Oxytropis falcata* Bunge and *Oxytropis chiliophylla* Royle. *J. Chin. Pharm. Sci.* **2014**, *23*, 783–789.

23. Shen, L.; Cong, W.J.; Lin, X.; Hong, Y.L.; Hu, R.H.; Feng, Y.; Xu, D.S.; Ruan, K.F. Characterization Using LC/MS of the Absorption Compounds and Metabolites in Rat Plasma after Oral Administration of a Single or Mixed Decoction of Shaoyao and Gancao. *Chem. Pharm. Bull.* **2012**, *60*, 712–721.

24. Zhang, J.Q.; Wang, Y.; Li, H.L.; Wen, Q.; Yin, H.; Zeng, N.K.; Lai, W.Y.; Wei, N.; Cheng, S.Q.; Kang, S.L.; et al. Simultaneous quantification of seventeen bioactive components in rhizome and aerial parts of *Alpinia officinarum* Hance using LC-MS/MS. *Anal. Methods* **2015**, *7*, 4919–4926.

25. Tian, D.; Yang, Y.; Yu, M.; Han, Z.Z.; Wei, M.; Zhang, H.W.; Jia, H.M.; Zou, Z.M. Anti-inflammatory chemical constituents of *Flos Chrysanthemi Indici* determined by UPLC-MS/MS integrated with network pharmacology. *Food Funct.* **2020**, *11*, 6340–6351.

26. Nóra, G.; András, D.; Szilvia, L.; Szabolcs, B.; László, K. Characterization and identification of isoflavanoid glycosides in the root of Spiny restharrow (*Ononis spinosa* L.) by HPLC-QTOF-MS, HPLC-MS/MS and NMR. *J. Pharm. Biomed.* **2016**, *123*, 74–81.

27. Wang, H.; Wang, J.Y.; Wu, X.P.; Shen, H.Y.; Luo, Y.; Dai, H.F.; Mei, W.L. HPLC-ESI-MS Analysis of Flavonoids Obtained from Tissue Culture of *Dracaena cambodiana*. *Chem. Res. Chin. Univ.* **2015**, *31*, 38–43.

28. Nóra, G.; András, D.; László, K.; Szabolcs, B. Separation and characterization of homopipericolic acid isoflavanoid ester derivatives isolated from *Ononis spinosa* L. root. *J. Chromatogr. B* **2018**, *1091*, 21–28.

29. Yu, D.D.; Liang, X.R. Characterization and Identification of Isoflavonoids in the Roots of *Millettia speciosa* Champ. by UPLC-Q-TOF-MS/MS. *Curr. Pharm. Anal.* **2019**, *15*, 580–591.

30. Wang, Y.; Xu, C.H.; Wang, P.; Lin, X.Y.; Yang, Y.; Li, D.H.; Li, H.F.; Wu, X.Z.; Liu, H.B. Pharmacokinetic comparisons of different combinations of Shaoyao-Gancao-Decoction in rats: Simultaneous determination of ten active constituents by HPLC-MS/MS. *J. Chromatogr. B* **2013**, *932*, 76–87.

31. Xie, S.; Zhao, T.; Zhang, Z.W.; Meng, J.F. Reduction of Dihydrokaempferol by *Vitis vinifera* Dihydroflavonol 4-Reductase to Produce Orange Pelargonidin-type Anthocyanins. *J. Agric. Food Chem.* **2018**, *66*, 3524–3532.

32. Tsimogiannis, D.; Samiotaki, M.; Panayotou, G.; Oreopoulou, V. Characterization of Flavonoid Subgroups and Hydroxy Substitution by HPLC-MS/MS. *Molecules* **2007**, *12*, 593–606.

33. Tang, X.Y.; Dai, Z.Q.; Wu, Q.C.; Zeng, J.X.; Gao, M.X.; Xiao, H.H.; Yao, Z.H.; Dai, Y.; Yao, X. Simultaneous determination of multiple components in rat plasma and pharmacokinetic studies at a pharmacodynamic dose of Xian-Ling-Gu-Bao capsule by UPLC-MS/MS. *J. Pharm. Biomed. Anal.* **2020**, *177*, 112836.

34. Lee, J.H.; Lee, S.J.; Park, S.; Kim, H.K.; Jeong, W.Y.; Choi, J.Y.; Sung, N.J.; Lee, W.S.; Lim, C.S.; Kim, G.S.; et al. Characterisation of flavonoids in *Orostachys japonicus* A. Berger using HPLC-MS/MS: Contribution to the overall antioxidant effect. *Food Chem.* **2011**, *124*, 1627–1633.

35. Yao, Y.F.; Lin, C.Z.; Liu, F.L.; Zhang, R.J.; Zhang, Q.Y.; Huang, T.; Zou, Y.S.; Wang, M.Q.; Zhu, C.C. Identification and Pharmacokinetic Studies on Complanatuside and Its Major Metabolites in Rats by UHPLC-Q-TOF-MS/MS and LC-MS/MS. *Molecules* **2019**, *24*, 71.

36. Chen, W.H. Investigation on the Chemical Constituents of *Tridax procumbens* and *Oxytropis falcata* and Total Synthesis of A New Pterocarpan. Ph.D. Thesis, Lanzhou University, Lanzhou, China, 2009.