

Review

Bridging Magnetic Field Agriculture and UAV-Based Precision Monitoring: An Integrated Dual-Stream Evidence Synthesis and Conceptual Framework for Field-Scale Validation

George Papadopoulos ^{1,2,*} , Evgenia Georgiou ³, Antonia Oikonomou ⁴ , Spyros Fountas ² and Dimitrios Bilalis ¹ 

¹ Laboratory of Agronomy, Department of Crop Science, Agricultural University of Athens, 11855 Athens, Greece; bilalis@aau.gr

² Laboratory of Agricultural Engineering, Department of Natural Resources Management & Agricultural Engineering, School of Environment and Agricultural Engineering, Agricultural University of Athens, 11855 Athens, Greece; sfountas@aau.gr

³ Laboratory of Soil Science and Agricultural Chemistry, Agricultural University of Athens, 11855 Athens, Greece; e.georgiou@aau.gr

⁴ Laboratory of Mineralogy and Geology, Department of Natural Resources and Agricultural Engineering, School of Environment and Agricultural Engineering, Agricultural University of Athens, 11855 Athens, Greece; aoikonomou@aau.gr

* Correspondence: gpapadopoulos@aau.gr

Abstract

Magnetic field (MF) technologies have been explored in agriculture since the 1930s, with research activity increasing markedly since 2016. However, they have not achieved mainstream adoption, partly because no MF-specific validated methodology exists for evaluating their effects under realistic field conditions. Unmanned Aerial Vehicle (UAV)-based multispectral sensing represents a potential pathway to address this limitation by providing spatially explicit, non-destructive estimates of key canopy physiological variables at field scale, thereby enabling, for the first time, the systematic evaluation and validation of MF treatment responses under open-field conditions. To realise this potential, however, a common evidential basis must first be established by identifying crop physiological variables that are both consistently modulated by MF treatments and reliably detectable by UAV remote sensing. This study addressed this challenge through a dual-stream evidence synthesis of 216 peer-reviewed publications, comprising 102 studies on MF treatments in agricultural crops and 114 studies on UAV-based multispectral monitoring. Evidence from both research domains was synthesised to identify physiological variables that are simultaneously responsive to MF treatments and detectable through UAV remote sensing. Five direct bridge variables were identified: chlorophyll content, nitrogen use efficiency (NUE)/nitrogen assimilation, above-ground biomass (AGB), leaf area index (LAI), and yield. Chlorophyll content emerged as the strongest bridge variable, combining consistent MF responsiveness with UAV estimation accuracies of up to $R^2 = 0.90$. Based on these findings, a conceptual framework was developed linking MF treatments, UAV-derived vegetation indices, ground-truth measurements, and machine-learning approaches for field-scale validation. The review revealed a complete absence of integration between the two research domains within the reviewed corpus, despite their strong biological and methodological compatibility. The proposed framework is conceptual and remains to be experimentally validated; it provides the first operational pathway for evaluating MF technologies under realistic farming conditions and may support future research on sustainable and digitally enabled crop production systems.



Academic Editor: Roberto Mancinelli

Received: 17 June 2026

Revised: 30 July 2026

Accepted: 4 August 2026

Published: 6 August 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and

conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

Keywords: magnetic field treatments; UAV multispectral sensing; vegetation indices; precision agriculture; nitrogen use efficiency; sustainable agriculture; evidence synthesis; conceptual framework

1. Introduction

Global food systems are under increasing pressure to expand production while simultaneously reducing their environmental footprint: global food production must increase by an estimated 60% by 2050 to meet rising demand, even as agriculture already accounts for 10–12% of global greenhouse gas emissions [1]. This challenge is particularly evident in the context of nitrogen (N) fertilisation, whose global losses to the environment are disproportionately large relative to other agricultural inputs, contributing significantly to both eutrophication and greenhouse gas emissions. The EU Farm to Fork Strategy [2], a key component of the European Green Deal, aims to reduce nutrient losses by 50% and fertiliser use by at least 20% by 2030 [2,3]. These targets reflect growing recognition that decades of intensive nitrogen application have contributed to widespread aquatic eutrophication, nitrate contamination, and biodiversity loss. However, in conventional farming systems, nitrogen use efficiency (NUE), defined as the proportion of applied N assimilated by the crop, routinely remains below 50%, meaning that more than half of the applied nitrogen is lost to the environment [4]. Improving NUE without compromising productivity requires not only agronomic optimisation but also the development of complementary physical technologies capable of enhancing plant nitrogen assimilation without additional chemical inputs. Within this context, the application of magnetic fields (MFs) to seeds, irrigation water, or growing crops has re-emerged as a promising, although still insufficiently field-validated, area of agronomic research. A plausible physiological basis for linking MF treatment to NUE is its reported effect on nitrate reductase (NR), the rate-limiting enzyme of nitrogen assimilation, suggesting a potential mechanism through which nitrogen utilisation may be enhanced without increasing fertiliser inputs. At the same time, the available evidence remains heterogeneous, with treatment protocols, field strengths, experimental conditions, and reported effect sizes varying considerably among studies, while validation under commercial field conditions remains limited.

Magnetic field treatments in agriculture are generally applied through three principal approaches. The first is magnetopriming, in which seeds are exposed to a controlled static or pulsed MF before sowing. The second is magnetised water irrigation (MWI), where irrigation water passes through a permanent magnetic device installed on the supply line. The third involves direct and continuous exposure of growing crops through in-field electromagnetic systems. Although these approaches are often grouped together, they differ substantially in exposure geometry, field strength, exposure duration, and, in the case of MWI, in their effects on water physicochemistry, and they consequently act on different primary biological targets. They share only a broad high-level rationale: the interaction of externally applied magnetic flux, whether acting directly on plant tissue or indirectly through treated water, with cellular processes involved in ion transport, enzyme activity, and reactive oxygen species (ROS) signalling [5]. The specific pathways underlying each delivery mode remain largely proposed rather than definitively established, and are an active area of investigation [4]. Across these delivery modes, investigated field intensities span weak static fields (0–100 μ T) through milliTesla- and Tesla-level fields to extremely-low-frequency (ELF) configurations, with reported responses varying according to crop species, developmental stage, and exposure duration [5]. At the cellular level, MF treatments alter ion fluxes, enzyme activity, and gene expression, improving water and

solute transport while enhancing chlorophyll and carotenoid synthesis and increasing the quantum efficiency of Photosystem II [6,7].

The agronomic potential of MF technology is supported by a growing body of evidence. In controlled germination experiments, magnetopriming of flax (*Linum usitatissimum* L.) at 350 mT for 100–120 min resulted in 100% germination, a 26.81-fold increase in seedling vigour index I, and a 2.63-fold increase in catalase activity compared with untreated controls [8]. The benefits of MF treatment extend beyond germination and persist through subsequent crop development. Field experiments on common bean (*Phaseolus vulgaris* L.) under MF treatment recorded increases of 13%, 21%, and 26% in chlorophyll a, chlorophyll b, and carotenoid content, respectively [9]. Of particular relevance to sustainable agriculture, MF treatments have been associated with improved NUE through the upregulation of NR activity, a key enzyme in plant nitrogen assimilation [6]. An evidence synthesis of magnetopriming effects on photosynthetic performance confirmed that static magnetic field (SMF) treatments consistently improve Photosystem II (PSII) efficiency, photosynthetic pigment concentrations, and leaf gas exchange parameters across multiple crop species [6]. These examples illustrate the upper range of reported responses and should not be read as typical. MF treatment effects are not universal: null, weak, and inconsistent responses are also reported across crop species and treatment configurations, and existing reviews emphasise substantial heterogeneity in outcomes alongside variability in experimental protocols and proposed mechanisms [6,9]. These physiological benefits are also attracting commercial interest: magnetic water treatment devices are already marketed to farmers, and magnetically treated water, including brackish water, is increasingly used in water-limited and saline environments where conventional irrigation is costly [7]. This uptake, however, is outpacing independent field validation: adopters do not have an established, non-destructive method to confirm whether a treatment yields a measurable physiological benefit under their own conditions.

Despite this convergence of scientific evidence and commercial adoption, an important methodological gap continues to limit the broader validation and agronomic credibility of MF technologies. Achieving such validation requires a field-scale monitoring approach capable of objectively assessing treatment responses under realistic agricultural conditions. Much MF research has been conducted under controlled laboratory or greenhouse conditions, including pot experiments, small uniform plots, and simplified monoculture systems in which spatial variability, microclimatic gradients, and field heterogeneity are intentionally minimised [5,7]. Even where field experiments exist, treatment responses have typically been assessed only through destructive or point-based measurement rather than spatially explicit, hectare-scale monitoring. Consequently, the translation of these findings to open-field agriculture at hectare scale remains largely unexplored. Farmers deploying MF technologies currently have no MF-specific validated field-monitoring methodology for determining whether treatments produce measurable physiological responses under their specific field conditions, at which spatial locations those responses occur, or at which growth stages they are most pronounced. Inconsistencies among experimental protocols, exposure conditions, and proposed mechanisms of action further hinder the interpretation and comparison of results [7,10], limitations that are compounded by the absence of non-destructive monitoring tools capable of assessing treatment effects at field scale. It is precisely this absence of field-scale monitoring capacity that Unmanned Aerial Vehicle (UAV)-based multispectral remote sensing is positioned to fill.

UAV-based multispectral remote sensing has meanwhile matured into a widely deployed precision agriculture technology capable of estimating canopy physiological status non-destructively across entire fields [11]. Estimation accuracy, however, depends on several factors, including lighting conditions, phenological stage, sensor type, and radiometric calibration. The critical observation for the present argument is not that UAV technology exists,

but that the specific variables it measures with greatest accuracy, chlorophyll content, nitrogen nutrition index (NNI), above-ground biomass (AGB), and LAI, are precisely the variables that MF treatments consistently improve. Normalised difference red-edge index (NDRE)-based indices have achieved R^2 values as high as 0.90 for chlorophyll content estimation, and NNI mapping from UAV imagery has been validated across wheat, rice, and maize at field scale. These reported accuracies are therefore conditional rather than universal and should not be assumed to transfer directly across platforms, crops, and environments [10]. This overlap enables a closed monitoring cycle: MF treatment induces a physiological response, the canopy spectral signature shifts, and a UAV detects the change. However, spectral changes alone cannot be attributed uniquely to MF treatment because they may also arise from soil heterogeneity, nutrient availability, irrigation, pest or disease pressure, phenology, or microclimatic variation. Consequently, UAV observations must be embedded within an appropriate experimental design, where treated and control plots are analysed using machine-learning approaches to distinguish treatment-induced responses from background variability. Framed in this way, the monitoring cycle provides exactly what MF research has long lacked at field scale: spatially explicit evidence of whether, where, and when measurable physiological responses occur. Exploiting this opportunity, however, requires first identifying which variables both systems share, the common ground that makes integrated monitoring possible. Identifying these shared variables is the primary objective of the present review.

This review establishes the evidential basis for that monitoring cycle by synthesising evidence from two previously unconnected domains: MF-induced physiological responses in agricultural crops (Stream A) and UAV multispectral detection of those same variables at field scale (Stream B). The proposed framework translates this synthesis into a concrete operational protocol. MF treatment parameters are selected based on Stream A evidence, while UAV flight plans and vegetation indices are matched to the identified bridge variables. Machine-learning models are then used to distinguish treatment-induced spectral signatures from background field variability. In doing so, the review responds directly to the Farm to Fork Strategy's emphasis on digital precision fertilisation and sustainable agricultural practices [2,12]. By identifying the existing research gap, integrating evidence across disciplines, and proposing a standardised protocol for future multi-site field experiments, this review aims to support the transition of MF technologies from controlled experimental settings to practical field-scale applications. The framework presented here is therefore intended as a conceptual, hypothesis-generating foundation that synthesises evidence from two previously independent research domains to define a pathway for future field-scale validation, rather than to claim that MF treatment effects have already been experimentally validated under commercial farming conditions.

2. Materials and Methods

2.1. Study Design and Rationale

This paper is structured in two complementary components: a dual-stream evidence synthesis and a conceptual framework construction. In the first component, two distinct literatures, the physiological effects of MF treatments on crops (Stream A) and UAV-based multispectral estimation of crop physiological variables (Stream B), were each screened and synthesised following PRISMA 2020 principles [13], applied as a structured methodological framework for transparent and reproducible study selection rather than as a formal evidence synthesis protocol. In the second component, the synthesised evidence from both streams was combined through an interpretive cross-domain variable-mapping procedure to construct the proposed monitoring framework. This second step involves interpretive judgement that extends beyond standard PRISMA evidence synthesis, and is described separately in Section 2.6. Rather than adopting a conventional single-domain approach, the study was designed to address a cross-

disciplinary research gap: the absence of a validated methodology for monitoring the in-field physiological responses of crops to MF treatments at agronomically relevant scales.

To develop a scientifically grounded monitoring framework, evidence was synthesised from two previously unconnected research domains. The first documents the physiological effects of MF treatments on agricultural crops, while the second demonstrates the capability of UAV-based multispectral remote sensing to estimate the same physiological variables non-destructively at field scale. The review focuses specifically on UAV multispectral sensing because it represents the most widely adopted, commercially mature, and operationally scalable remote sensing platform currently available for precision agriculture applications. Although other sensing approaches, including hyperspectral, thermal, fluorescence, and proximal sensing technologies, may provide complementary information, multispectral UAV systems currently offer the greatest potential for practical field-scale deployment. It should be noted that operational field-scale deployment ultimately situates UAV-derived measurements within a broader digital-agriculture ecosystem, including GIS-based soil and terrain data, climate layers, irrigation records, and geospatial decision-support platforms. The present framework is concerned specifically with establishing whether MF treatment responses are detectable by UAV multispectral sensing, a precondition for their subsequent integration into such decision-support systems; that integration is identified as a direction for future work rather than developed here.

The two streams therefore serve complementary roles within a single evidentiary framework rather than constituting independent reviews. Stream A identifies the physiological responses most consistently associated with MF treatments, whereas Stream B evaluates the capacity of UAV multispectral sensing to detect and quantify those responses under field conditions. The integration of evidence from both streams forms the basis of the conceptual monitoring framework proposed in this study.

As a Review applying PRISMA 2020 screening principles rather than a formal systematic review, the study was not prospectively registered; prospective registration in PROSPERO is intended for systematic reviews of health-related interventions and is not applicable to a cross-disciplinary methodological synthesis of this kind. The complete search strings (Table 1), screening criteria (Table 2), and extraction fields (Table 3) are reported in full to support transparency and reproducibility. The PRISMA-style flow diagram summarising the screening process for both streams is presented in Figure 1.




Figure 1. PRISMA 2020 flow diagram of the dual-stream evidence synthesis. Stream A (MF treatments, blue) and Stream B (UAV multispectral sensing, amber).

2.2. Research Questions

Two complementary research questions structured the literature search and data extraction:

RQ1 (Stream A): Which physiological variables are consistently and significantly modulated by MF treatments in agricultural crops, and at what experimental scales have these effects been documented?

RQ2 (Stream B): Can UAV-mounted multispectral sensors, through validated vegetation indices, detect and quantify the same physiological variables identified in RQ1, and with what level of reported accuracy?

The synthesis of answers to RQ1 and RQ2 directly informs the construction of the proposed monitoring framework: variables identified in Stream A as MF-responsive are cross-referenced against variables for which Stream B demonstrates remote-sensing detectability, yielding a set of index-variable pairs that form the operational basis of the framework.

2.3. Search Strategy

A systematic literature search was conducted in Scopus (www.scopus.com) in May 2026. Scopus was selected for its comprehensive, concurrent indexing of both the plant-science and remote-sensing/engineering literatures relevant to this review, providing a single, consistent indexing baseline against which the non-overlap of the two literatures could be established. Two independent queries were constructed, one per stream, using Boolean operators and field-code restrictions (TITLE-ABS-KEY). Search terms were selected to maximise precision rather than recall. This was a deliberate design choice: the aim was not exhaustive coverage of the precision-agriculture literature, but focused identification of two specific bodies of work, MF treatment effects and UAV multispectral estimation of the corresponding physiological variables, and determination of whether they intersect. Establishing this non-intersection depends on precise identification of each literature rather than on maximal retrieval; the present work is therefore a targeted evidence synthesis rather than a fully comprehensive systematic review. Literatures addressing the downstream integration of crop-monitoring data into geospatial and decision-support systems fell outside this scope, and their relevance to operational deployment is acknowledged in Section 4.5. The full queries are presented in Table 1. Stream A was restricted to publications from 2016 onwards to capture the period of significant growth in magnetopriming and magnetized water irrigation research. Stream B was restricted to publications from 2019 onwards to reflect the maturation of affordable UAV multispectral platforms (e.g., DJI Phantom 4 Multispectral, MicaSense RedEdge) in precision agriculture applications.

Table 1. Scopus search queries for Stream A (MF treatments) and Stream B (UAV remote sensing), with associated research questions.

Stream	Research Question	Scopus Query (TITLE-ABS-KEY)—Abbreviated
Stream A MF Treatments	Which physiological variables do MF treatments consistently modulate?	TITLE-ABS-KEY (("static magnetic field" OR "pulsed magnetic field" OR "electro-magnetic field" OR "magnetopriming" OR "magnetized water" OR "magnetically treated water" OR "magnetic water treatment") AND ("seed germination" OR "plant growth" OR "chlorophyll" OR "nitrogen use efficiency" OR "NUE" OR "biomass" OR "photosyn-thesis" OR "nitrogen uptake" OR "crop yield" OR "antioxidant enzyme" OR "reactive oxygen species") AND ("agriculture" OR "agronomy" OR "crop" OR "wheat" OR "maize" OR "soybean" OR "tomato" OR "lettuce" OR "rice" OR "sunflower" OR "barley")) AND NOT TITLE-ABS-KEY ("cancer" OR "tumor" OR "MRI" OR "medical" OR "animal" OR "livestock" OR "poultry" OR "fish" OR "scale inhibition" OR "pipe" OR "industrial" OR "wastewater treatment" OR "heavy metal removal") AND PUBYEAR > 2015 AND DOCTYPE (ar OR re) AND LANGUAGE (english)

Table 1. Cont.

Stream	Research Question	Scopus Query (TITLE-ABS-KEY)—Abbreviated
Stream B UAV Sensing	Can UAV multispectral indices detect those same variables with validated accuracy?	TITLE-ABS-KEY ((“UAV” OR “unmanned aerial vehicle” OR “drone” OR “UAS”) AND (“multispectral” OR “NDVI” OR “NDRE” OR “vegetation index”) AND (“chloro-phyll” OR “nitrogen use efficiency” OR “NUE” OR “nitrogen nutrition index” OR “NNI” OR “leaf nitrogen content” OR “nitrogen status” OR “photosynthetic pigment” OR “bio-mass estimation”) AND (“wheat” OR “maize” OR “rice” OR “soybean” OR “barley” OR “cereal” OR “field crop”)) AND NOT TITLE-ABS-KEY (“satellite” OR “Sentinel” OR “Landsat” OR “MODIS” OR “hyperspectral” OR “greenhouse” OR “indoor” OR “forest” OR “vineyard” OR “fruit tree”) AND PUBYEAR > 2018 AND DOCTYPE (ar OR re) AND LANGUAGE (english)

Details of Table: UAS: Unmanned Aerial System; NDVI: Normalised Difference Vegetation Index; MODIS: Moderate Resolution Imaging Spectroradiometer.

2.4. Study Selection and Screening

Following the database search, retrieved records were screened following the four-phase PRISMA 2020 structure: Identification, Screening, Eligibility, and Inclusion. First-pass screening was conducted at the title and abstract level applying the exclusion criteria in Table 2. Prior to screening, duplicate records were identified and removed. Title and abstract screening was conducted independently by two authors, and any disagreements were resolved through discussion until consensus was reached; records that remained ambiguous after discussion were retained and carried forward to full-text assessment rather than excluded. Papers were retained in cases of ambiguity and subjected to full-text eligibility review. Second-pass full-text screening applied additional exclusion criteria focused on methodological relevance to the framework objective: papers were excluded if they measured only molecular or intracellular endpoints with no canopy-level physiological outcome detectable by remote sensing (Stream A), or if the reported spectral data lacked quantified accuracy metrics and did not estimate a variable relevant to MF treatment response (Stream B).

Table 2. Inclusion and exclusion criteria applied at title/abstract and full-text screening stages.

Stream	Category	Stream A—Exclude If:	Stream B—Exclude If:
Both	Publication type	Conference abstract, book chapter, or non-peer-reviewed	Conference abstract, book chapter, or non-peer-reviewed
A only	Wrong organism	Animals, humans, bacteria, cell cultures, trees, ornamentals	—
A only	Wrong domain	Water purification, scale inhibition, medical, veterinary, industrial, post-harvest preservation	—
B only	Wrong platform	—	Satellite-only (Sentinel, Landsat, MODIS); manned aircraft; ground sensors with no UAV component
B only	Wrong sensor	—	RGB-only (no NIR band); hyperspectral; thermal-only imaging

Table 2. Cont.

Stream	Category	Stream A—Exclude If:	Stream B—Exclude If:
B only	Wrong target variable	—	Disease/pest/weed detection; soil mapping without canopy measurement; fruit quality; no N/chlorophyll/biomass link
B only	No quantified accuracy	—	No R ² , RMSE, or correlation metric between VI and physiological variable

2.5. Data Extraction

Data extraction was performed independently for each stream using a standardized extraction sheet developed a priori (Table 3). Extraction was designed specifically to serve the framework construction objective: fields were selected to generate the evidence base required to match MF-responsive physiological variables (Stream A) against UAV-detectable physiological variables (Stream B). Extraction was therefore purposive rather than exhaustive: fields were restricted to those required for cross-domain variable mapping and accuracy assessment, and were recorded from the title, abstract, and indexed metadata of each record. Study-level methodological parameters not required for the bridge mapping, such as replication, plot design, and sensor-calibration detail, were not extracted, and their standardised reporting is instead recommended for future primary studies. The critical operational field in each stream is the ‘framework bridge flag’: for Stream A, this records whether the physiological outcome measured is, in principle, detectable by UAV multispectral imaging; for Stream B, it records whether the target variable estimated corresponds to a variable consistently modulated by MF treatment. Papers flagged YES in both streams at the same variable form the direct evidentiary basis for each node in the proposed framework.

Table 3. Data extraction fields for Stream A (MF treatment papers) and Stream B (UAV remote sensing papers).

Field	Stream A—MF Papers	Stream B—UAV Papers
Technology type	SMF/PEMF/magnetized water/direct exposure	UAV platform (DJI Phantom/Matrice/Parrot/other)
Physical parameters	MF intensity (mT/μT/T); exposure duration; application mode	Sensor type; spectral bands available
Application stage	Pre-sowing/seedling/vegetative/reproductive/full season	Growth stage(s) at which UAV flights conducted
Target variable	Chlorophyll/NUE/NNI/biomass/N uptake/PSII/yield	Chlorophyll/NUE/NNI/leaf N/biomass/LAI/PSII/yield
Quantified accuracy	% change from control (direction and magnitude)	R ² and/or RMSE; ML algorithm if used
Framework bridge flag	Does the paper measure any variable detectable by UAV? (YES/NO/PARTIAL)	Does the paper estimate a variable directly modulated by MF? (YES/NO)

2.6. Framework Construction Approach

The dual-stream evidence synthesis was designed not only to characterise the existing literature in each domain but to identify where it can be operationally connected. Following extraction of all Stream A and Stream B data, a cross-domain variable mapping procedure was applied to determine which physiological variables simultaneously satisfy three evidence-based inclusion criteria: (i) the variable is reported with a consistent positive directional effect in at least two independent Stream A studies using MF treatment (a criterion applied to identify candidate variables for framework construction, not as a claim regarding the overall consistency or reliability of MF treatment effects, which vary across the literature); (ii) at least one Stream B study has demonstrated that the same variable can be estimated from UAV multispectral imagery at canopy scale with a reported $R^2 \geq 0.60$. This threshold corresponds to the conventional interpretation of the coefficient of determination in agronomic and remote-sensing model evaluation, in which values in the approximate range 0.5–0.7 are commonly regarded as indicating moderate predictive performance appropriate for exploratory rather than operational monitoring. It is acknowledged that R^2 characterises the strength of association rather than the magnitude of prediction error; reported error metrics and validation design are therefore considered alongside R^2 in the synthesis, and the threshold is applied as a screening criterion for candidate bridge variables rather than as a sufficient guarantee of operational accuracy; and (iii) the biological quantity is physically measurable at canopy scale by passive optical remote sensing, without requiring contact instrumentation or destructive sampling.

Variables meeting all three criteria were assigned a bridge status of YES (direct connection confirmed). Variables meeting criterion (i) but only partially meeting (ii) or failing (iii), specifically, variables not directly estimable by passive multispectral UAV but accessible via an indirect proxy or an emerging active sensing technique, were assigned a PARTIAL flag. Variables present in Stream A but absent from Stream B at the required accuracy threshold were excluded from the bridge table and recorded as monitoring gaps. Antioxidant enzyme activity (SOD, CAT, APX), for example, is consistently reported as a significant MF treatment response in Stream A but has no validated canopy-scale UAV detection method in Stream B and was therefore excluded.

The confirmed bridge connections formed the basis for a three-stage conceptual framework for scalable agronomic validation of MF treatments. The framework proceeds as follows. **Stage 1 (Treatment Design)** uses the Stream A evidence base to define experimental parameters, crop species, MF treatment type, intensity, and application timing, informed by the physiological responses most consistently documented for the target crop. **Stage 2 (UAV Monitoring Protocol)** specifies the UAV multispectral flight plan and the vegetation index (VI) corresponding to each confirmed bridge variable: NDRE for chlorophyll content and NNI, SAVI combined with DSM-derived canopy height for AGB, NDVI and NDRE with PROSAIL radiative transfer model inversion for LAI, and multi-temporal NDVI with machine learning for yield, generating spatially explicit canopy maps of treatment response at each critical growth stage. **Stage 3 (ML Validation)** applies machine learning architectures validated in Stream B to distinguish treatment-related spectral patterns, relative to control plots, from background spatial variability. This variability is attributable to environmental covariates including soil texture and moisture, terrain (slope and elevation), microclimate, rainfall and temperature, and irrigation infrastructure and management zones; where available, GIS-based layers describing these covariates provide the spatial context against which treatment-attributable canopy variation is isolated. Accounting for these covariates at the validation stage is essential to attributing canopy response to the MF intervention rather than to confounding environmental factors. Within the proposed framework, this approach can support a more robust interpretation of canopy responses

observed in MF-treated fields while reducing the influence of potential confounding factors. The cross-domain variable mapping that underpins this framework, together with the supporting evidence from both streams, is presented and discussed in Section 3.3.

2.7. Quality Assessment

Given the cross-disciplinary and framework-oriented nature of this review, formal risk-of-bias assessment tools designed for clinical studies (e.g., GRADE, Cochrane RoB) were not applicable. Methodological quality was assessed using two criteria applied during full-text screening: (i) whether the study reported quantified outcomes (% change from control, R^2 , RMSE) rather than only qualitative descriptions, and (ii) whether the experimental design included a true control group. A third criterion, field relevance, distinguished studies conducted under field or open-field conditions from those conducted in laboratory or controlled-environment settings, reflecting the framework's specific concern with field-scale validation. Together, these criteria distinguish studies providing quantified, control-referenced evidence under agronomically relevant conditions from those providing only qualitative, uncontrolled, or controlled-environment observations. A formal multi-criterion risk-of-bias instrument was not applied, consistent with the targeted, framework-oriented scope of the review and with quality-assessment practice in comparable technology reviews. Papers meeting both criteria were prioritized in framework construction; papers meeting only one were included in the narrative synthesis but flagged [14].

3. Results

3.1. Stream A: Systematic Evidence on Magnetic Field Treatment Effects in Agricultural Crops

3.1.1. Screening Outcomes and Study Characteristics

The dual-stream PRISMA screening process is summarised in Table 4 and Figure 1. For Stream A, the Scopus query returned 172 records. First-pass title and abstract screening excluded 49 papers: 23 fell outside the agricultural MF domain (food processing, pseudoscience, non-MF electromagnetic applications, or engineering papers); 15 studied non-target organisms (tree crops, medicinal herbs, model plants, fungi, or human cell lines); and 11 addressed wrong outcome variables (heavy metal phytoremediation, herbicide management, or soil engineering). Second-pass full-text screening of the remaining 123 papers excluded a further 21 papers: 11 reported exclusively molecular or cellular endpoints with no canopy-level physiological outcome; 4 used treatments confounded by a second physical agent with no MF-only control; and 6 were excluded for other methodological or crop-context reasons, including hydroponic indoor systems, germination- or water-quality-only outcomes, and off-topic crop contexts. The final Stream A corpus comprised 102 included papers.

Table 4. PRISMA screening outcomes for Stream A (MF treatments) and Stream B (UAV remote sensing).

PRISMA Phase	Stream A—MF	Stream B—UAV
Records identified via Scopus	172	170
Excluded at 1st screening (title/abstract)	49	55
Retained after 1st screening	123	115
Excluded at 2nd screening (full abstract/text eligibility review)	21	1
FINAL INCLUDED for synthesis	102	114

Of the 102 papers, 93 were original research articles and 9 were review articles. Of the 93 original research articles, 89 provided extractable treatment data and were therefore

used for the descriptive characterisation of study scale, crop distribution, and MF treatment categories. Among the 89 empirical studies, study scale distribution reflects the lab-to-field gap central to this paper: 23 papers (26%) used purely laboratory or controlled-chamber conditions; 21 papers (24%) employed pot or greenhouse experiments; and 45 papers (51%) reported results from small field plots or open-field conditions. Although just over half of the empirical corpus (45 of 89 studies, 51%) was therefore conducted under field or open-field conditions, indicating that field research on MF treatment is not itself lacking, all of these field studies relied exclusively on destructive or point-based measurements. None employed spatially explicit, remotely monitored, or multi-site approaches at hectare scale, the specific form of field validation that the framework proposed here is designed to enable. Soybean (*Glycine max* L.) was the most studied species (16 papers), followed by wheat (*Triticum* spp., 13 papers), tomato (*Solanum lycopersicum* L., 10 papers), maize (*Zea mays* L., 10 papers), lettuce (*Lactuca sativa* L., 5 papers), and rice (*Oryza sativa* L., 2 papers). By MF treatment type, calculated from the 89 non-review empirical papers and recorded as non-mutually exclusive categories, MWI was reported in 47% of studies ($n = 42$), SMF magnetopriming in 36% ($n = 32$), pulsed electromagnetic field PEMF treatments in 16% ($n = 14$), and combined or direct in-season exposure in 7% ($n = 6$). Of the 102 papers, 52 (51%) received a framework bridge flag of YES and 50 (49%) received a PARTIAL flag. Critically, within the included corpus, none of the 102 Stream A papers employed remote sensing of any kind to monitor treatment response, a finding that directly quantifies the research gap this review addresses. This result describes the reviewed studies rather than asserting the global non-existence of such work; its robustness to the search strategy is considered in Section 4.6. The full quantitative breakdown, including document type, study scale, crop species distribution, MF treatment-type frequencies, physiological variables measured, and framework bridge-flag classification, is presented in Table 5.

Table 5. Stream A study characteristics.

Category/Variable	<i>n</i>	%	Denominator
Document Type			
Original research article	93	91.2%	102
Review article	9	8.8%	102
Study Scale (89 empirical)			
Laboratory/controlled chamber only	23	25.8%	89
Pot/greenhouse only	21	23.6%	89
Field plot/open field	45	50.6%	89
Crop Species			
Soybean (<i>Glycine max</i>)	16	15.7%	102
Wheat (<i>Triticum</i> spp.)	13	12.7%	102
Tomato (<i>Solanum lycopersicum</i>)	10	9.8%	102
Maize (<i>Zea mays</i>)	10	9.8%	102
Lettuce (<i>Lactuca sativa</i>)	5	4.9%	102
Rice (<i>Oryza sativa</i>)	2	2.0%	102
Other/multiple species	39	38.2%	102
MF Treatment Type (89 empirical) a			
Magnetised water irrigation (MWI)	42	47.2%	89
Static MF/magnetopriming (SMF)	32	36.0%	89
Pulsed electromagnetic field (PEMF)	14	15.7%	89
Combined/direct in-season exposure	6	6.7%	89

Table 5. *Cont.*

Category/Variable	<i>n</i>	%	Denominator
Physiological Variable Measured			
Chlorophyll/photosynthetic pigments	43	42.2%	102
Grain/fruit yield	47	46.1%	102
PSII maximum quantum efficiency (Fv/Fm)	23	22.5%	102
Nitrogen metabolism/NUE/nitrate reductase (NR)	15	14.7%	102
Water use efficiency (WUE)	11	10.8%	102
Above-ground biomass (AGB)	10	9.8%	102
Leaf area index (LAI)	6	5.9%	102
Framework Bridge Flag			
YES—directly UAV-detectable	52	51.0%	102
PARTIAL—proxy/indirect only	50	49.0%	102
Remote Sensing Used			
Yes	0	0.0%	102
No—destructive/point measurement only	102	100%	102

Table details: Denominator: base population for each percentage (102 = full corpus; 89 = empirical studies only). Bold text within the table body indicates category headings.

3.1.2. MF Treatment Types and Application Modes

Three principal MF delivery modes are represented in the Stream A corpus: magnetopriming, magnetized water irrigation (MWI), and direct in-season MF exposure. These approaches differ in their mode of application and duration of exposure but share the common objective of modulating plant physiological processes through MF intervention, albeit through delivery-mode-specific and largely proposed mechanisms. These approaches differ in their mode of application and duration of exposure but share the common objective of modulating plant physiological processes through MF intervention.

Magnetopriming involves exposing seeds to a static or pulsed MF prior to sowing, typically for 30–120 min at field intensities ranging from 50 mT to 1000 mT [5,6,15]. As a pre-sowing treatment, magnetopriming is applied only once before crop establishment. The proposed mode of action includes modifications in seed membrane permeability, enzyme activity, and water absorption kinetics, resulting in enhanced germination, accelerated emergence, and improved seedling vigour. Importantly, several studies report carry-over effects that extend beyond early development, including improvements in chlorophyll content and nitrogen metabolism that persist through later growth stages.

Magnetized water irrigation (MWI) involves passing irrigation water through a device containing permanent magnets before application to the crop [16–18]. The treatment is proposed to alter physicochemical water properties, particularly hydrogen bonding angle, surface tension, and dissolved oxygen content. Unlike magnetopriming, MWI can be applied continuously throughout the growing season and is currently the only MF treatment mode with established commercial deployment at farm scale. Commercial systems are available for integration into drip and sprinkler irrigation infrastructure, making MWI the most operationally mature MF technology represented in the corpus.

Direct in-season MF exposure applies continuous or periodic electromagnetic fields to actively growing crops through dedicated in-field equipment [19–21]. Although it is the least frequently studied approach in the reviewed literature, largely because of its infrastructure requirements, it provides the most direct means of influencing physiological processes

during critical developmental stages such as anthesis and grain filling. As a result, it offers the potential for targeted intervention during periods of high physiological sensitivity.

3.1.3. Synthesis of MF Effects on UAV-Detectable Physiological Variables Chlorophyll Content and Photosynthetic Pigments

Enhanced chlorophyll accumulation is the most consistently reported MF effect across the Stream A corpus, appearing in 43 of the 102 included papers and spanning all three treatment modes as well as all major crop species. This response is therefore not restricted to a particular crop or MF application strategy but represents a broadly documented physiological outcome of MF treatment.

Evidence of enhanced chlorophyll content has been reported across multiple crop species and treatment configurations. SMF magnetopriming of soybean improved Fv/Fm together with chlorophyll a and chlorophyll b concentrations under both UV-B stress conditions [22,23] and natural growing conditions [24]. Similarly, magnetopriming of tomato seeds enhanced chlorophyll content and leaf area during vegetative development [25], while magnetized water irrigation increased chlorophyll a, chlorophyll b, and carotenoid concentrations in tomato [26,27]. In addition, PEMF treatment of durum wheat during the growing season produced significant improvements in chlorophyll content alongside increases in yield and enzyme activity [28]. Field-scale magnetized water experiments have also reported enhanced chlorophyll pigment concentrations in bean [29] and tomato [26], including increases in chlorophyll a, chlorophyll b, and carotenoids.

The mechanistic basis for these responses has been documented across multiple crop species. Proposed mechanisms include enhanced Mg^{2+} availability for chlorophyll synthesis, reduced photo-oxidative damage, and improved thylakoid membrane integrity, all of which contribute to greater photosynthetic capacity and pigment accumulation [5–7]. From a framework perspective, chlorophyll content represents the strongest bridge variable identified in this review. Among all physiological variables evaluated, chlorophyll exhibits the highest reported UAV estimation accuracy in the Stream B corpus, with NDRE-based models achieving R^2 values of up to 0.90 [30]. This combination of consistent MF responsiveness and highly accurate remote sensing detectability establishes chlorophyll content as the primary and most robust connection between MF treatment physiology and field-scale UAV monitoring. It should be emphasised that chlorophyll content, although the most responsive and most detectable bridge variable, is not specific to MF treatment: canopy chlorophyll variation can also arise from nitrogen availability, water status, disease, cultivar differences, and growth stage. UAV-detected chlorophyll change therefore indicates a canopy response requiring interpretation, not MF causation per se. Robust use of chlorophyll as a validation variable consequently requires paired ground-truth measurement, SPAD readings or extractive chlorophyll determination, together with randomised spatial blocking of treated and control plots, so that treatment-attributable variation can be separated from background agronomic and environmental sources.

Nitrogen Assimilation and Nitrogen Use Efficiency

A total of 15 papers in the Stream A corpus report direct effects of MF treatment on plant nitrogen metabolism, making nitrogen assimilation one of the most consistently documented physiological responses relevant to sustainable crop production. Across these studies, improvements in nitrogen uptake, assimilation, and utilisation have been reported under both SMF and magnetized water treatments, indicating that the effect is not restricted to a single MF application strategy. The primary mechanistic pathway involves the upregulation of NR activity, the enzyme catalysing the first and rate-limiting step of nitrate reduction ($NO_3^- \rightarrow NO_2^-$), thereby enabling subsequent ammonium assimilation

into amino acids. Enhanced NR activity under SMF and magnetopriming conditions has been reported across multiple studies [31,32], providing a physiological basis for the observed improvements in nitrogen-related parameters.

Representative evidence has been documented across several crop species and experimental conditions. SMF treatment in wheat significantly enhanced NR activity, nitrogen assimilation efficiency, and canopy chlorophyll content (CCC), with effects confirmed under both controlled and open-field conditions [33]. These improvements were, where reported, generally observed under comparable agronomic management conditions, suggesting that the observed enhancements in nitrogen assimilation and utilisation were attributable to the MF treatment rather than differences in nutrient supply. In legumes, magnetized water irrigation combined with root MF exposure increased nodule formation and biological nitrogen fixation in bean crops [34]. Similarly, magnetized water irrigation improved nitrogen absorption in rice seedlings grown under saline-alkaline stress conditions [35]. Additional improvements in nitrogen-related parameters, including nitrogen content, protein accumulation, and water and radiation use efficiency, have been reported in chickpea under both SMF magnetopriming [36] and magnetized water irrigation [37], as well as in winter wheat under magnetized water irrigation [38,39].

From a framework perspective, nitrogen assimilation represents one of the most important bridge variables identified in this review. The biochemical responses documented in Stream A are directly linked to canopy-level indicators of crop nitrogen status that can be estimated non-destructively using UAV multispectral imagery. In particular, NNI mapping, validated extensively in Stream B, provides a practical means of monitoring the spatial expression of MF-induced improvements in nitrogen assimilation and NUE under field conditions. These nitrogen-related quantities operate at different biological scales and should be distinguished. Nitrate reductase activity is an enzymatic response measured at the tissue level and is not itself observable by remote sensing. Leaf and canopy nitrogen content and the NNI are canopy-scale indicators of crop nitrogen status, estimable from UAV multispectral imagery. Nitrogen use efficiency, by contrast, is an efficiency ratio of yield or biomass relative to nitrogen input, and therefore requires agronomic input–output accounting that spectral estimation alone cannot provide. UAV-based NNI estimation thus offers a canopy-scale indicator consistent with the nitrogen-assimilation improvements documented in Stream A, corroborating rather than confirming the underlying biochemical mechanism such as NR upregulation.

Above-Ground Biomass and Yield

Improvements in biomass accumulation and final yield represent some of the most agronomically significant outcomes reported in the Stream A corpus. Field-scale yield increases are documented in 47 papers, while explicit measurements of AGB are reported in approximately 10 studies. Evidence of positive biomass and yield responses spans a wide range of crop species, including wheat [38–43], maize [21,44–46], sunflower [20,47,48], and various legume species [49–52].

Yield improvements have been reported across all major MF treatment types and under diverse environmental conditions. The strongest responses are generally associated with magnetized water irrigation, particularly under water-stressed conditions where improvements in water uptake and physiological efficiency may confer a greater agronomic advantage. In field-grown soybean, magnetopriming produced significant yield gains accompanied by enhanced gas exchange and PSII efficiency [50]. Similarly, PEMF seed treatment in buckwheat and maize resulted in measurable increases in photosynthetic performance, biomass accumulation, and final yield in replicated field trials [53,54].

These findings suggest that biomass and yield improvements are not isolated responses but rather downstream manifestations of the enhanced physiological performance documented throughout the Stream A literature, including improvements in chlorophyll content, photosynthetic activity, and nitrogen assimilation. From a framework perspective, AGB represents a particularly valuable bridge variable because it can be estimated reliably at field scale using UAV multispectral imagery. In the Stream B corpus, SAVI and DSM-integrated approaches consistently achieve AGB estimation accuracies ranging from $R^2 = 0.61$ to 0.85 [55,56], providing a practical means of assessing MF-induced growth responses under real farming conditions. Because yield integrates the cumulative effects of many processes across the entire season, it is inherently more confounded than physiological indicators such as chlorophyll or LAI, and is correspondingly estimated with lower accuracy by UAV imagery ($R^2 = 0.33$ – 0.61 ; Section 3.2.2). Yield is therefore best treated as a final validation endpoint rather than as an early diagnostic indicator: chlorophyll, nitrogen status, and LAI provide the earlier and more mechanistically informative signals, while yield confirms whether those responses ultimately translate into productivity. Robust use of yield as a validation variable requires harvest-based ground truth, for example, plot-level yield maps from calibrated combine yield monitors, rather than reliance on spectral estimation alone.

Leaf Area Index and Canopy Development

Increases in LAI and canopy development are reported less frequently than chlorophyll or biomass responses, appearing in a limited but biologically plausible subset of six studies in the Stream A corpus. While too few to establish consistency in a statistical sense, these reports are mechanistically coherent with the improvements in leaf expansion and canopy development documented for other bridge variables. These responses are generally associated with improved leaf expansion, enhanced stomatal conductance, and accelerated canopy closure, all of which contribute to greater light interception and photosynthetic capacity.

Representative examples include magnetized water irrigation of bean [57] and magnetopriming of maize [44], both of which resulted in significant increases in leaf area and LAI under field conditions. In addition, downstream improvements in radiation use efficiency associated with enhanced canopy development have been documented in chickpea and sunflower under SMF treatment [20,36]. Although the number of studies is smaller than for other physiological variables, the consistency of the reported responses suggests that canopy structural development is a recurring outcome of MF treatment.

From a framework perspective, LAI represents an important bridge variable because it can be estimated directly from UAV multispectral imagery with validated accuracy across multiple crop species. Studies included in Stream B report LAI estimation accuracies ranging from $R^2 = 0.70$ to 0.88 using NDVI-, NDRE-, and PROSAIL-based approaches (see Section 3.2.2). Furthermore, UAV-derived LAI maps can be generated at sub-metre spatial resolutions, reaching ground sampling distances as low as 0.015 m [58]. This combination of biological relevance and reliable remote sensing detectability makes LAI a valuable component of the proposed monitoring framework.

Implications of MF Treatment Modes for UAV Monitoring

Although the principal physiological bridge variables identified in this review are common across the different MF treatment modes, their temporal expression under field conditions is expected to vary with treatment timing, from the single pre-sowing application of magnetopriming to the cumulative, season-long exposure of magnetized water irrigation (Section 3.1.2). Accordingly, while the physiological targets monitored by UAV

multispectral sensing remain similar, the timing and frequency of image acquisition should be adapted to the expected response dynamics of each treatment mode. However, the current evidence does not yet permit the definition of treatment-specific UAV monitoring protocols, highlighting an important direction for future field-scale validation studies.

3.2. Stream B: UAV Multispectral Remote Sensing for Physiological Variable Estimation

3.2.1. Screening Outcomes and Study Characteristics

The Stream B Scopus query returned 170 records. First-pass title and abstract screening excluded 55 papers across three categories: 23 studies lacked an operational UAV monitoring component, relying instead on satellite remote sensing platforms (e.g., Sentinel-2, Landsat, MODIS), ground-based proximal or handheld spectroscopy, or general radiative transfer modelling without drone-acquired imagery; 13 studies used sensor configurations incompatible with the multispectral framework, predominantly RGB-only cameras or thermal-only sensors without near-infrared spectral bands; and 19 studies targeted application domains with no link to N status, chlorophyll, or biomass estimation, including crop disease and pest detection, herbicide tolerance screening, evapotranspiration mapping, grain maturity identification, radiometric correction methodology, and UAV spraying system design. Second-pass full-text screening excluded one additional paper [59] whose primary output was a flight-time methodology study without any estimation of agronomic physiological variables. The final Stream B corpus comprised 114 papers from which all data were extracted.

The corpus spans publications from 2019 to 2026, reflecting the rapid growth of UAV multispectral applications following the commercialisation of affordable consumer-grade platforms. The most commonly studied crops are wheat (*Triticum aestivum* and *T. durum*, 40 papers, 35%), rice (*Oryza sativa*, approximately 30 papers, 26%), maize (*Zea mays*, approximately 28 papers, 25%), and soybean (*Glycine max*, approximately 10 papers, 9%). UAV platforms are dominated by multirotor DJI systems, including the DJI Phantom 4 Multispectral (P4M) [60,61], alongside fixed-wing platforms such as the senseFly eBee equipped with the Parrot Sequoia+ multispectral sensor [62]. Sensor configurations are predominantly 5-band multispectral (Blue, Green, Red, Red-Edge, NIR), with some studies combining multispectral with RGB imagery for feature fusion [63–66]; a subset of studies augmented spectral data with LiDAR for explicit 3D canopy structure mapping [67–69], while several others derived canopy height estimates from UAV photogrammetry-based digital surface models (DSMs) without LiDAR [55,56]. Of 114 papers, 111 (97%) received a framework bridge flag of YES, confirming that the estimated variable corresponds to a physiological parameter consistently modulated by MF treatment in Stream A. Zero of the 114 Stream B papers studied magnetically treated crops, confirming the complete absence of cross-domain application that this review addresses. This finding does not derive from the search terms: the Stream B query contained no magnetic-treatment terms and its exclusion criteria did not filter them out, so a UAV study monitoring a magnetically treated crop would have satisfied all inclusion criteria and been retrieved. Each retrieved paper was instead assessed for magnetic treatment during data extraction (the framework bridge flag), so the absence is established by the content of the included studies rather than by query design. The full quantitative breakdown, including crop species distribution, primary target variables, analytical approach, and framework bridge-flag classification, is presented in Table 6.

Table 6. Stream B study characteristics.

Category/Variable	<i>n</i>	%	Denominator
Document Type			
Original research article	113	99.1%	114
Review article	1	0.9%	114
Crop Species			
Wheat (<i>Triticum</i> spp.)	40	35.1%	114
Rice (<i>Oryza sativa</i>)	30	26.3%	114
Maize (<i>Zea mays</i>)	28	24.6%	114
Soybean (<i>Glycine max</i>)	10	8.8%	114
Other/multi-crop	10	8.8%	114
Primary Target Variable (non-excl.)			
Chlorophyll/SPAD/LCC/CCC	43	37.7%	114
NNI/NUE/N status	42	36.8%	114
Above-ground biomass (AGB)	30	26.3%	114
Grain/fruit/seed yield	35	30.7%	114
Leaf area index (LAI)	15	13.2%	114
Canopy height/phenology	6	5.3%	114
Stomatal/WUE/CWSI	5	4.4%	114
Analytical Approach			
Machine learning (beyond regression)	36	31.6%	114
Framework Bridge Flag			
YES—detects MF-modulated variable	111	97.4%	114
PARTIAL—indirect/proxy	3	2.6%	114
MF-TREATED CROPS STUDIED	0	0.0%	114

Table details: Denominator = 114 (full Stream B corpus) for all rows. Bold text indicates category headings within the table.

3.2.2. Vegetation Index Performance for MF-Relevant Variables

The Stream B literature consistently evaluated a common set of physiological variables using UAV multispectral sensing, although the level of evidence and estimation performance varied among variables. To facilitate comparison across the reviewed studies, Table 7 summarizes the principal physiological bridge variables, the approximate number of supporting studies, the dominant vegetation indices and analytical approaches, the highest reported estimation accuracies, and their relevance within the proposed conceptual framework. The subsequent sections discuss these variables in greater detail.

Table 7. Summary of UAV multispectral performance for the principal physiological bridge variables identified in Stream B.

Physiological Variable	No of Studies	Dominant VI	Best Reported Performance	Framework Significance
Chlorophyll content	43	NDRE; GNDVI; ML/Radiative Transfer Model (RTM)	R ² up to 0.90	Strongest bridge variable linking MF-induced chlorophyll enhancement with UAV monitoring
Nitrogen Nutrition Index/NUE	41	NDRE; ML ensembles	High accuracy across crops	Direct indicator of MF-induced improvements in nitrogen assimilation

Table 7. *Cont.*

Physiological Variable	No of Studies	Dominant VI	Best Reported Performance	Framework Significance
AGB	29	NDRE + canopy height (DSM)	Improved when spectral + structural features combined	Integrates cumulative crop growth responses
LAI	15	PROSAIL; RTM; ML	High accuracy across seasons	Indicator of canopy development
Yield	33	Multispectral + ML	$R^2 = 0.33\text{--}0.61$	Final agronomic outcome of MF-induced physiological responses

Chlorophyll Content Estimation

Chlorophyll estimation from UAV multispectral imagery is the most extensively validated application identified in the Stream B corpus, appearing as the primary target variable in 43 studies. Across a wide range of crop species and management conditions, chlorophyll content consistently emerges as one of the physiological variables that can be estimated with the highest accuracy using UAV-derived vegetation indices.

Among the evaluated indices, the normalized difference red-edge index (NDRE) consistently outperforms NDVI and other broadband vegetation indices due to its sensitivity to chlorophyll absorption in the red-edge spectral region (700–730 nm). A principal reason for this advantage is that NDVI tends to saturate at high canopy chlorophyll and biomass, losing discriminatory sensitivity in the range most relevant to well-developed crop canopies, whereas red-edge indices such as NDRE remain responsive across this range. This advantage is, however, contingent on sensor configuration: not all UAV multispectral sensors include a red-edge band, and the applicability of NDRE-based monitoring within the proposed framework therefore depends on the deployment of a sensor with an appropriate red-edge channel. Across multiple cereal and field crop systems, NDRE-based approaches achieved the highest reported accuracies, reaching $R^2 = 0.90$ for chlorophyll estimation in triticale and sunflower under varying nitrogen amendment levels [30]. Similarly, two-year UAV monitoring of winter wheat demonstrated stable and reliable chlorophyll estimation across growing seasons [70].

Alternative vegetation indices also demonstrated strong performance under specific conditions. The green normalized difference VI (GNDVI) was identified as the most effective green-band index for chlorophyll estimation in smallholder maize systems subjected to varying nitrogen fertilisation regimes [71]. These findings suggest that although NDRE is generally the most robust indicator, alternative indices may provide advantages depending on crop type, management practices, and sensor configuration.

Recent studies have further improved chlorophyll estimation through the integration of machine learning and physically based modelling approaches. Physics-informed transfer learning frameworks combining UAV multispectral imagery with radiative transfer models (RTM) achieved high accuracy for leaf chlorophyll content (LCC) and CCC estimation in winter wheat [72], while improving model transferability across environments. Similarly, a double-layer physical-statistical modelling approach successfully estimated canopy photosynthetic pigments, including chlorophyll a, chlorophyll b, and carotenoids, together with canopy nitrogen content in winter wheat [73].

Collectively, these findings establish chlorophyll content as the most robust and consistently validated physiological variable within the Stream B corpus. Combined with the strong evidence for MF-induced chlorophyll enhancement identified in Stream A, this high level of remote sensing detectability reinforces chlorophyll as the strongest bridge variable linking MF treatment responses with field-scale UAV monitoring.

Nitrogen Nutrition Index and NUE Estimation

The NNI, defined as the ratio of actual crop nitrogen concentration to the critical nitrogen concentration at a given biomass level, is one of the most agronomically relevant indicators of crop nitrogen status and is directly linked to NUE. Reflecting its importance for nutrient management and precision agriculture, 42 papers in the Stream B corpus address the estimation of NNI or NUE using UAV multispectral imagery. Across these studies, NDRE emerged as the dominant VI for nitrogen status assessment. Its sensitivity to changes in leaf nitrogen content within the red-edge spectral region enables the detection of nitrogen-related variation before visible deficiency symptoms become apparent. As a result, NDRE-based approaches have become a central component of UAV-assisted nitrogen management strategies.

The capability of UAV systems to estimate NNI and NUE has been demonstrated across a wide range of crops, environmental conditions, and management scenarios. High-accuracy NNI estimation was reported in rice at the critical early panicle initiation stage using a combination of spectral, texture, and discrete wavelet transform (DWT) features evaluated across three growing seasons and ten varieties [74]. UAV-based NNI estimation has also been successfully applied in smallholder winter wheat systems at the village scale using an eBee fixed-wing platform equipped with a Parrot Sequoia+ multispectral sensor, demonstrating applicability beyond large experimental research stations [62]. Direct UAV-based NUE phenotyping has been reported across full growing seasons in rice [75], spring wheat [76], wheat under varying soil mineral nitrogen conditions [77], and winter wheat [61,78–81].

Recent studies further demonstrate the value of advanced analytical approaches for improving nitrogen status estimation. Stacking ensemble machine learning models integrating UAV-derived spectral information with environmental and management variables achieved enhanced NNI prediction accuracy across diverse rice-growing environments [82]. These developments illustrate the increasing maturity of UAV-based nitrogen monitoring and its capacity to support field-scale assessment of crop nitrogen dynamics under heterogeneous conditions.

From a framework perspective, NNI and NUE represent the most operationally valuable bridge variables identified in this review. The improvements in nitrogen assimilation and NR activity documented in Stream A can be translated into spatially explicit indicators of crop nitrogen status through UAV-derived NNI mapping. Consequently, UAV multispectral sensing provides a practical and non-destructive approach for evaluating whether MF-induced improvements in nitrogen metabolism are expressed at canopy scale under real field conditions. It should be noted that UAV-based estimation addresses crop nitrogen status (NNI, canopy/leaf nitrogen content) rather than a complete measure of NUE. Where the framework contributes to precision nutrient management, it does so by providing spatially explicit nitrogen-status information; complete evaluation of NUE requires this to be combined with agronomic yield and nitrogen-input records.

Above-Ground Biomass and LAI Estimation

Above-ground biomass estimation from UAV multispectral imagery is addressed in 30 papers within the Stream B corpus. Unlike chlorophyll estimation, which can often

be achieved using a single VI, accurate biomass estimation generally requires the integration of both spectral and structural crop information. Consequently, the best-performing approaches combine vegetation indices with canopy height metrics derived from DSMs generated through UAV photogrammetry, consistently improving prediction accuracy compared with VI-only regression approaches [56,83].

Several studies demonstrate the effectiveness of this integrated approach across diverse crop systems and management conditions. The photosynthetic accumulation model (PAM), which combines vegetation indices and canopy height information, achieved improved AGB estimation performance in rice across multiple cultivars and growing seasons [55]. Biomass estimation has also been successfully evaluated under varying irrigation and nitrogen management regimes in corn [84] and in maize-based smallholder farming systems [85]. In addition, soil-adjusted vegetation indices such as SAVI have been incorporated to minimise the influence of soil background effects under partial canopy cover conditions commonly encountered in grain crops [86].

Leaf area index estimation is addressed in approximately 15 papers and represents a complementary measure of canopy development and crop growth. Several studies demonstrate that UAV multispectral imagery can provide reliable LAI retrievals through both empirical and physically based approaches. Among these, PROSAIL radiative transfer model inversion enabled physically interpretable LAI estimation at a spatial resolution of 0.015 m in maize [58]. Similarly, physics-informed transfer learning frameworks estimated LAI together with LCC and CCC with high accuracy across multiple growing seasons in winter wheat [72].

From a framework perspective, both AGB and LAI represent valuable indicators of the cumulative physiological effects of MF treatments. Whereas chlorophyll and NNI provide insight into specific physiological processes, biomass accumulation and canopy development integrate the combined effects of enhanced photosynthetic performance, nitrogen assimilation, and crop growth over time. The demonstrated capability of UAV systems to estimate these variables at field scale therefore provides an important mechanism for evaluating whether MF-induced physiological improvements translate into measurable agronomic outcomes under real production conditions.

Yield Prediction

Yield prediction is addressed in 35 papers within the Stream B corpus and represents one of the most practically relevant applications of UAV multispectral remote sensing. Unlike chlorophyll content, NNI, or LAI, yield is not a directly measurable physiological variable but rather the cumulative outcome of multiple interacting processes occurring throughout the growing season. Consequently, yield prediction presents a greater modelling challenge and typically exhibits lower prediction accuracies than intermediate physiological indicators.

Despite this complexity, UAV-based yield estimation has been successfully demonstrated across several cereal systems using commercially available platforms. Consumer-grade DJI Phantom 4 Multispectral (P4M) systems have been validated for grain yield prediction in both barley [60] and winter wheat [61], highlighting the accessibility of the approach for both research and practical applications. Reported grain yield prediction accuracies range from $R^2 = 0.33$ to 0.61 in barley breeding trials employing nonparametric modelling across multiple measurement dates [60].

Although these accuracies are lower than those reported for chlorophyll content, NNI, or LAI estimation, they remain sufficient for detecting treatment-level differences at field scale. From a framework perspective, this is particularly important because yield represents the ultimate agronomic outcome of the physiological responses induced by

MF treatments. As such, UAV-based yield prediction provides a means of evaluating whether improvements in chlorophyll content, nitrogen assimilation, biomass accumulation, and canopy development ultimately translate into measurable productivity gains under field conditions.

3.2.3. Machine Learning Integration for Predictive Validation

A total of 36 papers in the Stream B corpus employ machine learning (ML) approaches beyond conventional regression techniques for the estimation of physiological variables from UAV-derived spectral information. The reported methods span a broad range of algorithms, including Random Forest (RF), support vector machines (SVM), convolutional neural networks (CNN), deep learning architectures, and ensemble learning approaches. Collectively, these studies demonstrate the increasing role of ML in improving prediction accuracy, handling complex non-linear relationships, and integrating multiple sources of agronomic information.

Several studies highlight the advantages of advanced ML frameworks over traditional modelling approaches. A CNN-LSTM-XGBoost hybrid architecture applied to multimodal UAV imagery achieved robust performance for field-scale nitrogen stress classification while providing interpretable, feature-level insights into nitrogen status decisions [87]. Similarly, PSO-optimised ensemble learning combined with spectral feature selection significantly outperformed individual models for chlorophyll content estimation in rice under varying nitrogen application regimes [88]. Ensemble learning approaches have also improved the estimation of maize LCC and fractional vegetation cover (FVC) across multiple growth stages, addressing variations in model performance throughout the growing season [89].

The increasing adoption of ML-based approaches reflects a broader shift from simple VI regression towards data-driven predictive modelling capable of integrating spectral, structural, environmental, and management-related information. This development is particularly relevant for field-scale applications, where crop responses are influenced by multiple interacting factors and where traditional linear relationships may not fully capture observed variability. A related challenge for multi-site field deployment is that predictive performance can degrade substantially when models trained on one field, sensor, or season are applied to another; recent deep learning architectures address this directly, from bi-directional domain adaptation networks that separate domain-invariant and domain-specific spectral features for cross-temporal classification [90], to parameter-efficient fine-tuning frameworks that adapt pretrained models to new spectral modalities and unseen regions or seasons by updating only a small fraction of model parameters [91].

From a framework perspective, these ML architectures provide a potential mechanism for strengthening the interpretation of UAV-derived physiological responses in MF-treated fields. Once baseline spatial maps of chlorophyll content, nitrogen status, biomass, or canopy development have been generated, ML models may help identify treatment-related spectral patterns while accounting for background variability associated with soil heterogeneity, management history, and microclimatic conditions. Although these approaches have not yet been specifically validated for MF-treated cropping systems, the evidence presented in Stream B suggests that they offer a promising pathway for future field-scale assessment and validation of MF treatment responses. A methodological caveat applies to the accuracies reported across this literature. In spatially structured crop datasets, random partitioning into training and test sets can allow information to leak between them, for instance, when neighbouring plots or repeated flights over the same field are split across the partition, inflating apparent performance through spatial autocorrelation. Accuracies obtained under random cross-validation should therefore be regarded as potentially op-

timistic relative to genuinely independent data. Spatially and temporally independent validation, such as spatial cross-validation or hold-out by site or growing season, gives a more realistic indication of the transferable accuracy achievable on new fields, as would be required for MF treatment validation.

3.3. Cross-Domain Evidence Synthesis: The Framework Bridge Table

The central analytical output of this dual-stream review is the cross-domain variable mapping presented in Table 8. The purpose of this mapping is not simply to list variables reported in the two corpora, but to identify which MF-modulated physiological responses can realistically be monitored at field scale using UAV-based multispectral sensing. Each row therefore represents either a confirmed or partially confirmed bridge connection between Stream A and Stream B.

A bridge connection was classified as direct when the same physiological variable was consistently modulated by MF treatment in Stream A and could be estimated non-destructively at canopy scale by UAV multispectral imaging with validated accuracy in Stream B. These connections were assigned a YES flag. Variables were classified as partial when the MF-responsive trait could not be directly measured by standard passive multispectral UAV sensors but could potentially be inferred through a proxy variable or an emerging sensing approach. These connections were assigned a PARTIAL flag. In this way, the bridge table provides the operational basis for the proposed monitoring framework by distinguishing variables that are immediately suitable for field-scale validation from those requiring additional methodological development.

Table 8. Cross-domain bridge table: MF-modulated physiological variables (Stream A) mapped against UAV-detectable variables with best-performing VI and reported accuracy (Stream B). Bridge status indicates whether the same variable is directly detectable (YES) or via proxy/indirect method (PARTIAL).

MF-Modulated Variable	Stream A (<i>n</i> Papers)	MF Effect—Direction & Evidence	Best UAV Index	Best R ²	Bridge	Major Confounders	Required Ground-Truth	Recommended Validation Approach
Chlorophyll content (Chl a + b, carotenoids)	43	Consistent improvement across SMF, PEMF, and MWI in wheat, maize, soybean, tomato, lettuce, and cotton. Mechanism: Mg ²⁺ channel activation → enhanced chlorophyll synthesis; reduced photo-oxidative damage to thylakoid membranes [22–24,26–28,33,46]	NDRE, GNDVI	0.90 [30]	YES	Nitrogen availability, water status, disease, cultivar, growth stage	SPAD/extractive chlorophyll assay	Paired ground-truth + randomised spatial blocking
Nitrogen status (NNI/leaf N)—indicator of N assimilation	15	SMF and magnetopriming upregulate NR (NR: NO ₃ [−] → NO ₂ [−]), the first and rate-limiting step of N reduction. NUE and canopy N content improvements in soybean, wheat, chickpea, rice field experiments [31–35]	NDRE → NNI model	≥0.85 [62,74,82]	YES	Water status, cultivar, growth stage; enzyme mechanism not directly observed	Leaf N (Kjeldahl/Dumas), plant N% for NNI	NNI mapping + agronomic N-input accounting
AGB	10	Positive AGB and canopy growth across all MF types in field experiments—wheat, maize, sunflower, legumes. Enhanced dry matter via photosynthate translocation [20,21,39,40,44,45]	SAVI; NDVI + DSM (canopy height)	0.61–0.85 [55,56]	YES	Soil background, canopy structure, planting density	Destructive biomass harvest	VI + DSM canopy-height fusion; harvest calibration
LAI	6	Limited but biologically plausible evidence: MF and magnetopriming associated with increased leaf expansion via improved stomatal conductance and cell division. Field-scale LAI increases reported in bean and maize only [44,46,51,57]	NDVI, NDRE; PROSAIL RTM inversion	0.70–0.88 [58,72]	YES	Growth stage, cultivar canopy architecture	Ground LAI (ceptometer/LAI-2200)	Multi-temporal UAV + ground LAI calibration
Grain/fruit yield	47	Aggregate downstream outcome of improved photosynthesis, NUE, and biomass. Documented across wheat, maize, soybean, chickpea, and bean in open-field trials [28,37,39,40,46–48,50]	NDVI at heading/grain-fill; RF/ANN (multi-stage)	0.33–0.61 [60,61]	YES	Highly integrative—weather, management, all upstream factors	Plot-level harvest/calibrated combine yield maps	Final endpoint only; plot yield maps

Table 8. Cont.

MF-Modulated Variable	Stream A (n Papers)	MF Effect—Direction & Evidence	Best UAV Index	Best R ²	Bridge	Major Confounders	Required Ground-Truth	Recommended Validation Approach
PSII maximum quantum efficiency (Fv/Fm)	23	SMF preserves PSII reaction centres; consistent Fv/Fm improvements in magnetopriming studies under stress [22–24,46,50]	Indirect: GNDVI (Chl proxy); chlorophyll fluorescence + UAV emerging [92]	Indirect	PARTIAL	Not passively detectable; requires active fluorometry	PAM fluorometry (point measurement)	Not currently field-deployable; flag as monitoring gap
WUE	11	MWI reduces surface tension → improved water uptake and stomatal regulation. Documented in chickpea, watermelon, sunflower, cotton under deficit irrigation [36,48,93]	CWSI (thermal + MS); stomatal conductance via PROSAIL [94–96]	Indirect	PARTIAL	Requires thermal/water-status data; not multispectral-detectable	Gas exchange/sap flow/thermal CWSI	Requires thermal payload; monitoring gap

Five of the seven mapped variables received a YES bridge flag, confirming direct methodological connections between MF treatment outcomes and UAV monitoring capability. Chlorophyll content represents the strongest bridge node. It is the most consistently reported MF-responsive variable in Stream A, with 43 papers documenting improvements in chlorophyll a, chlorophyll b, and carotenoid concentrations across all MF treatment modes and several major crop species [22–28]. At the same time, Stream B demonstrates very high UAV-based detectability for chlorophyll, with NDRE-based models reaching $R^2 = 0.90$ in a multi-crop, multi-season field study [30]. This makes chlorophyll the most robust entry point for field-scale validation of MF treatment responses, because it combines strong biological responsiveness with high remote sensing accuracy.

Nitrogen nutrition index and NUE form the second critical bridge node and are particularly important from a sustainability perspective. Stream A provides evidence that MF treatments can influence nitrogen metabolism through NR upregulation and improvements in canopy nitrogen status [31–33]. Stream B, in turn, demonstrates that NNI can be estimated using NDRE-based UAV approaches across wheat, rice, and maize systems [62,74,82]. This bridge matters because it links the biochemical mechanism most relevant to NUE improvement with a practical field-scale monitoring method. It therefore provides a direct pathway for assessing whether MF treatments contribute to more efficient nitrogen use under real agronomic conditions.

Above-ground biomass, LAI, and yield complete the five confirmed direct bridge connections [39–41,44–46,55,56,58,60,61,83]. These variables are important because they represent progressively more integrated expressions of crop performance. AGB and LAI reflect the cumulative effects of improved photosynthesis, nitrogen assimilation, and canopy development, while yield represents the final agronomic outcome of these physiological processes. Although UAV-based yield prediction shows lower accuracy than chlorophyll or LAI estimation, with reported values ranging from $R^2 = 0.33$ to 0.61 , it remains relevant for detecting treatment-level productivity differences at field scale. In contrast, LAI estimation reaches higher accuracies of $R^2 = 0.70$ – 0.88 , supporting its use as a reliable canopy development indicator within the proposed framework.

Collectively, the five confirmed bridge variables represent different levels of crop response to MF treatment. Chlorophyll content and nitrogen assimilation reflect primary physiological processes directly associated with photosynthetic performance and nutrient metabolism. Above-ground biomass and LAI represent structural canopy responses that integrate these physiological effects over time, while yield constitutes the final agronomic outcome of their cumulative expression. Together, these variables provide a hierarchical framework for evaluating MF treatment effects from mechanism to productivity under field conditions. In parallel, PSII efficiency and WUE remain physiologically important partial bridge variables, but their direct field-scale estimation requires sensing approaches beyond standard passive multispectral UAV platforms.

A further consideration governs the practical utility of each bridge connection: whether the magnitude of the MF-induced change exceeds the estimation error of the corresponding UAV retrieval. The bridge variables are not equivalent in this respect. Reported MF effects on chlorophyll content (e.g., increases of 13–26% in pigment concentrations [9]) are large relative to the residual error implied by NDRE-based estimation at $R^2 = 0.90$ [30], indicating that treatment-level contrasts should be resolvable from single-pass imagery. For AGB ($R^2 = 0.61$ – 0.85 [55,56]) and particularly yield ($R^2 = 0.33$ – 0.61), the margin between treatment effect and retrieval error is considerably narrower, implying that replicated plots and multi-temporal acquisition are required. Chlorophyll therefore represents the recommended entry point for field-scale validation, a design constraint on the framework rather than an objection to it.

Two variables received a PARTIAL bridge flag. F_v/F_m , documented in 23 Stream A papers [22–24,46,50], is physiologically important because it reflects the functional status of Photosystem II and is frequently improved by MF treatment, particularly under stress conditions. However, direct F_v/F_m measurement requires active pulse-amplitude modulation fluorometry (PAMF), which is not available through standard passive multispectral UAV platforms. Only one Stream B paper linked chlorophyll fluorescence measurements with UAV canopy imagery [92], leaving direct field-scale spatial mapping of F_v/F_m unvalidated. Similarly, WUE, reported in 11 Stream A papers [36,38,48,93], can currently be estimated only indirectly through thermal-derived Crop Water Stress Index (CWSI) or PROSAIL-coupled stomatal conductance modelling [94–96]. These approaches require sensor configurations beyond standard five-band multispectral UAV payloads. Therefore, PSII efficiency and WUE remain promising but methodologically less mature bridge variables.

The most important finding of this synthesis is not contained within either corpus independently but emerges from their comparison. Within the corpus identified by this review, none of the 102 Stream A papers used remote sensing to monitor MF treatment responses in the field, and none of the 114 Stream B papers examined magnetically treated crops. Within the corpus identified by the present search strategy, no study directly integrated MF crop treatments with UAV multispectral monitoring. The bridge identified in Table 8 therefore represents, within this evidence base, a first-time connection between two established bodies of agricultural research rather than a simple methodological refinement. This absence of direct overlap confirms the central research gap addressed by the review and provides the rationale for the conceptual monitoring framework proposed in the following section.

4. Discussion

4.1. Bridging Two Previously Disconnected Research Domains

This review pursued three objectives. The first was to characterise the physiological variables consistently modulated by MF treatments (Stream A). The second was to establish which of those variables can be estimated at field scale through UAV multispectral sensing (Stream B). The third was to determine, through cross-domain mapping, which variables can serve as bridge variables within an operational monitoring framework. The discussion that follows evaluates the extent to which each of these objectives was met. The central finding of this review is neither the evidence for MF treatment effects on crop physiology nor the established capability of UAV multispectral sensing to detect those effects [5,7,11]. The central finding is that 102 papers documenting MF-induced physiological responses and 114 papers demonstrating UAV-based detection of those same physiological variables were identified, published and cited across the same ten-year period (2016–2026) without any documented intersection between them. None of the Stream A papers used remote sensing of any kind to monitor treatment responses in the field, and none of the Stream B papers studied crops subjected to MF treatment. This complete absence of cross-domain integration, across a combined corpus of 216 peer-reviewed papers spanning 2016–2026, is the observation on which the scientific novelty of this work rests.

Perhaps the most telling indicator of the depth of this separation is the directionality of the bridge connections identified in Table 8. When the 114 Stream B papers were systematically assessed against the physiological variables most consistently modulated by MF treatments, 111 (97.4%) were found to be studying variables that Stream A confirms are MF-responsive, principally chlorophyll content, NNI, AGB, and LAI. The two domains were, in effect, investigating the same biological phenomena: one through experimental intervention and contact measurement, the other through spatially resolved, non-destructive

remote sensing. The separation therefore appears to reflect disciplinary fragmentation and limited methodological integration rather than a lack of subject matter compatibility.

The crop species studied in the two corpora reinforce this observation. Wheat (*Triticum* spp.) featured in 13 Stream A papers and 40 Stream B papers. Maize (*Zea mays*) appeared in 10 Stream A and 28 Stream B studies, while soybean (*Glycine max*) was examined in 16 Stream A and 10 Stream B papers. In each case, the same crop species was being investigated by both research communities during the same publication period, under conditions that would in principle have supported methodological integration. Nevertheless, no evidence of such integration was identified.

Several factors may plausibly have contributed to this structural separation. While a formal bibliometric analysis lies beyond the scope of this review, some indication is available within the corpus data itself. The most likely contributing factor is disciplinary fragmentation. Stream A papers were concentrated in journals such as *Bioelectromagnetics*, *Plants*, *Horticulturae*, *Plant Signaling & Behavior*, and the *International Journal of Molecular Sciences*, reflecting a focus on plant physiology, agronomy, and biophysics. In contrast, Stream B papers were concentrated in journals such as *Remote Sensing*, *Precision Agriculture*, *Computers and Electronics in Agriculture*, and *Smart Agricultural Technology*, reflecting expertise in remote sensing, engineering, and digital agriculture. These research areas have largely evolved within different disciplinary communities, which appears to have limited opportunities for methodological cross-referencing.

A second factor that may have contributed is the measurement paradigm established within MF research. By the time UAV multispectral platforms became widely accessible, MF studies had already adopted a standard toolkit based on contact and destructive measurements, including SPAD meters, portable fluorimeters, gas-exchange analysers, and gravimetric biomass determination. Consequently, few studies in the MF literature drew upon or referenced the rapidly expanding UAV sensing literature, as confirmed by the complete absence of remote sensing applications identified in Stream A. A third possible factor is the mechanistic heterogeneity that has characterised MF research. Variability in experimental protocols and proposed mechanisms across studies [7,10] may have limited engagement from the precision agriculture community, where methodological reproducibility and operational transferability are often prioritised [11]. A formal bibliometric or citation-network analysis would be required to confirm the relative contribution of these factors, and represents a useful direction for future investigation.

The consequences of this separation have been substantial. Of the 89 empirical Stream A studies, 45 (51%) were conducted under open-field or field-plot conditions, yet all relied exclusively on contact-based or destructive measurements to assess treatment outcomes. As a result, the spatial distribution of physiological responses, within-field variability in treatment expression, and temporal dynamics across growth stages remained largely unexplored throughout the field-scale MF literature, not because these dimensions were scientifically unimportant, but because the monitoring methodology capable of capturing them at agronomic scale had matured in a parallel literature that MF researchers had not incorporated. The 45 Stream A field experiments therefore represent a body of research in which spatial expression, temporal trajectory, and within-field heterogeneity of MF treatment response have received comparatively limited attention.

From the perspective of digital agriculture, the findings carry an equally significant implication. The 111 Stream B papers identified as directly relevant to MF treatment monitoring had already validated spectral indices, estimation approaches, and species-specific protocols for variables known to respond to MF treatment. This accumulated methodological evidence provides a readily available foundation for the design of UAV-based MF validation trials. Spectral indices validated extensively in wheat, maize, and rice systems, such as

NDRE for chlorophyll content and NNI estimation, SAVI and DSM-integrated approaches for AGB assessment, and PROSAIL-based methods for LAI retrieval, could be applied to MF treatment experiments without requiring fundamental methodological redevelopment.

Beyond its specific implications for MF research and UAV precision agriculture, the findings of this review suggest that similar opportunities for cross-domain integration may exist elsewhere in agricultural research, where technological and biological disciplines often evolve in parallel despite addressing closely related agronomic questions. The dual-stream PRISMA approach adopted here, combining independent systematic evidence synthesis with structured cross-domain variable mapping, may therefore provide a useful methodology for identifying and formalising such connections in future studies.

4.2. From Physiological Responses to Field-Scale Validation

The bridge variables identified in Table 8 describe more than a technical correspondence between MF-responsive traits and UAV-detectable indicators. They represent a biologically coherent sequence of crop responses, progressing from primary physiological processes to canopy development and finally to agronomic performance. This hierarchy is important because it reflects how treatment effects are expected to emerge in the crop: first through changes in photosynthetic and nutritional processes, then through canopy growth, and ultimately through biomass accumulation and yield formation. Importantly, it also provides a mechanistic framework for interpreting MF treatment responses under field conditions. If changes are detected only at the physiological level but fail to propagate to canopy development or yield, this may indicate that the treatment effect is transient or environmentally constrained. Conversely, consistent responses across all three biological levels would provide stronger evidence that MF treatments generate agronomically meaningful benefits.

The identification of bridge variables across these biological scales is particularly important because it transforms the challenge of MF validation from a single end-point assessment into a process-based monitoring strategy. Rather than evaluating MF treatments solely through final yield measurements, researchers can monitor the progression of treatment responses throughout the growing season, thereby improving both mechanistic understanding and field-scale validation (Figure 2).

At the physiological level, chlorophyll content, nitrogen assimilation, NUE, and PSII efficiency represent some of the earliest and most mechanistically informative indicators of MF treatment response. These variables are directly associated with the plant's capacity to capture light, assimilate carbon, and use nitrogen efficiently. Leaf nitrogen, chlorophyll content, photosynthetic capacity, and radiation use efficiency are closely linked to biomass accumulation and crop productivity in the broader crop physiology literature [97,98], and these relationships underpin the biological coherence of the proposed bridge connections. In the context of this review, this is particularly important because the most consistently reported MF responses in Stream A correspond to the physiological variables most reliably estimated by UAV multispectral sensing in Stream B.

The second level of the response pathway is represented by structural canopy traits, particularly LAI and AGB. These variables integrate physiological changes over time and provide a more spatially explicit expression of crop growth. Both LAI and AGB are widely recognised as key indicators of crop development, reflecting canopy expansion, radiation interception capacity, and overall crop productivity potential [99]. While chlorophyll and nitrogen status indicate whether physiological activity has been altered, LAI and biomass show whether these changes are translated into canopy expansion, light interception, and dry matter production. This intermediate level is essential for field validation because it connects early physiological responses with whole-crop performance. It also fits the

practical strengths of UAV monitoring, which can repeatedly map canopy development and biomass-related indicators across entire field plots without destructive sampling.

Yield represents the final agronomic level of the hierarchy. Unlike chlorophyll, NNI, LAI, or biomass, yield is not a direct physiological trait but the cumulative outcome of many interacting processes occurring across the growing season. For this reason, yield prediction is expected to show lower accuracy than intermediate indicators. However, this does not reduce its relevance within the framework. Instead, it clarifies its role: yield should be treated as the final validation endpoint, while chlorophyll, nitrogen status, LAI, and biomass provide earlier diagnostic indicators explaining how and when treatment effects develop.

This hierarchical organisation strengthens the proposed framework because it avoids reliance on any single variable as proof of MF treatment effectiveness. A yield-only assessment would be too late and too integrative to explain treatment mechanisms, while a chlorophyll-only assessment would be insufficient to demonstrate agronomic relevance. By combining physiological, structural, and agronomic indicators, the framework enables MF treatment effects to be evaluated across multiple biological scales. This creates a more robust validation pathway, where early physiological responses can be linked to canopy development and then examined in relation to final productivity.

Therefore, the value of UAV-based monitoring in this context is not only its capacity to estimate individual variables, but its ability to follow the progression of MF-induced responses through time and space. This is particularly relevant for MF technologies, where treatment effects may depend on crop species, application method, growth stage, and environmental conditions. A multi-level monitoring strategy can help determine whether observed physiological responses remain localised, whether they develop into measurable canopy-level changes, and whether they ultimately contribute to agronomic performance under field conditions.




Figure 2. Process-based validation pathway for MF treatment effects, tracked by UAV monitoring from physiological responses to canopy development and final agronomic outcome.

4.3. The Proposed Framework as a Pathway Towards Scalable Validation of MF Technologies

One of the most persistent challenges in MF agriculture research is not the demonstration of treatment effects, but their validation under realistic farming conditions. Although the Stream A corpus provides substantial evidence that MF treatments can influence chlorophyll content, nitrogen metabolism, biomass accumulation, and yield-related traits, the majority of studies have relied on measurement approaches that inherently constrain experimental scale and monitoring intensity. Among the 89 empirical studies included in this review, 23 were conducted under laboratory or controlled-environment conditions

and a further 21 under greenhouse or pot-based conditions. Even among the 45 studies performed under field conditions, treatment responses were assessed exclusively through contact-based or destructive measurements. Consequently, most evidence regarding MF effectiveness remains derived from observations collected at individual plants, small plots, or limited sampling locations rather than from continuous field-scale assessments.

This methodological limitation has important implications for the interpretation and transferability of existing findings. Agricultural fields are inherently heterogeneous environments characterised by spatial variability in soil properties, nutrient availability, water status, microclimate, and crop development. Under such conditions, treatment responses may not be spatially uniform. However, traditional monitoring approaches such as SPAD measurements, chlorophyll extraction, biomass harvesting, gas exchange analysis, and fluorometry provide information only at discrete sampling points. While these methods remain essential for physiological validation, they cannot readily characterise the spatial distribution of treatment effects across entire fields or capture how responses evolve through time at agronomically relevant scales.

The framework proposed in this review addresses this scalability bottleneck by shifting the focus from point-based assessment to spatially explicit monitoring. Rather than replacing conventional physiological measurements, UAV multispectral sensing complements them by extending observations from individual sampling locations to complete field parcels at agronomically meaningful spatial resolution. This capability is particularly relevant because the variables most consistently affected by MF treatments, including chlorophyll content, nitrogen status, LAI, biomass, and yield-related traits, correspond closely to the variables most extensively validated for UAV-based estimation in Stream B. The framework therefore builds on an existing evidence base rather than requiring the development of entirely new sensing methodologies.

An additional advantage of the proposed approach lies in its temporal dimension. Many MF studies evaluate treatment responses at a limited number of growth stages or rely on end-of-season assessments. Such measurements provide valuable information regarding treatment outcomes but reveal little about how responses emerge, persist, or diminish during crop development. By enabling repeated observations throughout the growing season, UAV-based monitoring creates opportunities to investigate treatment dynamics rather than treatment outcomes alone. This distinction is important because MF-induced responses may vary according to crop species, environmental conditions, developmental stage, and treatment method. Monitoring these dynamics could improve understanding of when treatment effects are strongest and which physiological pathways are most closely associated with subsequent canopy development and productivity.

The framework may also contribute to improving the reproducibility and comparability of future MF studies. One of the recurring challenges identified in the MF literature is the diversity of treatment protocols, field strengths, exposure durations, crop species, and measured response variables. This heterogeneity complicates comparisons among studies and makes meta-analysis difficult. The use of a common monitoring approach based on validated UAV-derived indicators could facilitate more standardised assessment of treatment responses across sites, crops, and growing conditions. While such standardisation would not eliminate biological variability, it could improve the consistency with which treatment outcomes are quantified and reported.

Importantly, the framework should not be viewed as evidence that MF technologies are agronomically effective under all conditions. Rather, its value lies in providing a practical pathway through which such effectiveness can be evaluated at field scale. By integrating physiological indicators, canopy-level measurements, and agronomic outcomes within a single monitoring strategy, the framework offers a means of testing whether

responses observed under controlled conditions can be reproduced under the spatial and environmental complexity of commercial agricultural systems. In this respect, the framework should be understood as a hypothesis-driven pathway for field-scale validation rather than an experimentally verified monitoring system, a distinction that the limitations discussed in Section 4.6 address directly.

4.4. Implications for Sustainable Agriculture and Precision Nutrient Management

The methodological integration proposed in this review has direct relevance to sustainable agriculture and precision nutrient management, two domains explicitly addressed by current EU agricultural policy. Current agricultural policy increasingly emphasises the need to improve resource-use efficiency while reducing the environmental impacts associated with crop production. Within the European Union, the Farm to Fork Strategy identifies nutrient losses, excessive fertiliser use, and declining environmental sustainability as major challenges requiring the adoption of innovative and evidence-based agricultural practices [2,100].

Among the physiological responses most consistently associated with MF treatments in Stream A, improvements in chlorophyll content, nitrogen assimilation, NR activity, and NUE are particularly relevant in this context. These responses suggest a potential capacity of MF technologies to influence crop nutrient utilisation and physiological performance without requiring additional chemical inputs. However, despite the promising evidence reported across controlled-environment and field studies, the absence of a standardised field-scale monitoring methodology has limited the ability to evaluate whether these benefits remain consistent under commercial farming conditions.

The framework proposed in this review addresses this challenge by linking MF-responsive physiological variables with UAV-derived indicators already established within precision agriculture. In doing so, it creates a pathway through which MF technologies can be assessed using the same monitoring approaches currently employed for nitrogen management, crop status assessment, and decision support. Rather than introducing a completely new monitoring paradigm, the framework builds upon existing digital agriculture infrastructure and validated multispectral sensing workflows that are already widely used in research and commercial farming [101].

This connection is particularly relevant for precision nutrient management. Nitrogen use efficiency is widely recognised as a critical indicator of both economic and environmental sustainability because inefficient nitrogen use contributes simultaneously to production costs, greenhouse gas emissions, nitrate leaching, and water quality degradation [12,102]. UAV-based estimation of chlorophyll content and nitrogen nutrition status has become an established approach for supporting site-specific nutrient management and reducing unnecessary fertiliser inputs [103]. The identification of direct bridge connections between MF-induced improvements in nitrogen-related processes and UAV-derived NNI estimation therefore creates opportunities to evaluate whether MF technologies could contribute to more efficient nutrient use under field conditions.

Realising these sustainability benefits in practice requires that UAV-derived physiological information be translated into actionable, farmer-facing decisions. Spatially explicit maps of chlorophyll content, nitrogen status, and biomass can be aggregated into management-zone delineations and delivered to growers through geospatial decision-support and WebGIS platforms, which convert canopy-scale measurements into site-specific nutrient-management recommendations [104]. Integrating the monitoring framework proposed here with such decision-support infrastructure would close the loop between physiological detection and management action and represents the operational pathway through which UAV-based MF validation could contribute to the nutrient-efficiency objectives of

the Farm to Fork Strategy. The development of this decision-support layer is identified as a priority for future work.

More broadly, the framework aligns with the principles of sustainable intensification and digital agriculture. Sustainable intensification seeks to increase agricultural productivity while minimising environmental impacts and maintaining ecosystem services [105]. This is closely aligned with the broader transition toward climate neutrality in agriculture, in which the optimisation of nutrient and irrigation-water use is a central requirement [106]. Achieving this objective requires technologies that can be evaluated using robust, transparent, and scalable monitoring systems. By integrating physiological indicators, canopy development metrics, and agronomic outcomes within a single monitoring pathway, the proposed framework provides a mechanism through which the performance of MF technologies can be assessed objectively across diverse production environments.

Realising this potential, however, depends on weighing the practical advantages of the approach against its costs. Its principal advantages are that it requires no additional chemical input, is non-destructive, and can be deployed at field scale using commercially available multispectral platforms, allowing treatment responses to be monitored at a spatial resolution that conventional field scouting cannot match. These benefits are offset by the additional instrumentation and data-processing costs relative to conventional monitoring, and by the current absence of formal cost-benefit evidence for MF treatment itself (Section 4.6). A balanced consideration of these factors, the resource-efficiency gains on one side, the monitoring and treatment costs on the other, will ultimately determine the contribution of the dual MF-UAV approach to the transition toward climate-neutral agriculture.

Importantly, the framework should not be interpreted as evidence that MF technologies are inherently sustainable or agronomically beneficial. If future field validation confirms the physiological and agronomic benefits reported in controlled studies, the proposed framework could provide the monitoring infrastructure required to evaluate MF technologies under practical farming conditions. Such evidence would be essential before any conclusions could be drawn regarding their contribution to sustainable crop production, nutrient management, or future precision agriculture strategies.

Ultimately, the value of the framework lies not only in its potential application to MF technologies but also in its demonstration of how digital agriculture tools can accelerate the evaluation of emerging agronomic innovations. As agriculture increasingly moves towards data-driven and sustainability-oriented management systems, approaches capable of linking physiological processes with field-scale monitoring will become increasingly important for translating experimental findings into practical agricultural solutions.

4.5. Future Research Priorities

The most immediate research priority emerging from this review is the implementation of dedicated field experiments integrating MF treatments with UAV multispectral monitoring. While the bridge connections identified in this study demonstrate that several MF-responsive physiological variables can be estimated using established UAV methodologies, no study identified within the literature corpus included in the present review combined MF treatments with UAV multispectral monitoring within a single experimental framework. Consequently, the proposed framework remains a hypothesis-driven pathway requiring empirical validation under real agronomic conditions. Such experiments should include both conventional physiological measurements and UAV-derived indicators to assess whether treatment responses observed at plant level can also be detected consistently at canopy scale. Additionally, given that Stream A identified three distinct MF delivery modes, magnetized water irrigation (MWI, 47% of empirical papers), SMF magnetopriming (36%), and PEMF treatment (16%), future field experiments should evaluate whether

the strength and spatial pattern of canopy-level responses detectable by UAV monitoring differ systematically among these treatment types. MWI, as the only mode with established commercial deployment at farm scale, represents the highest-priority candidate for field-scale UAV validation trials. Such trials should additionally compare the three MF delivery modes, MWI, static-field magnetopriming, and pulsed electromagnetic treatment, to establish whether the strength and temporal profile of canopy-level responses detectable by UAV differ systematically among them (Section 3.1.3).

A second priority is the evaluation of the framework across multiple crops, management systems, and environmental conditions. The current evidence base is dominated by soybean, wheat, tomato, and maize, while several economically important crop species remain underrepresented. Furthermore, environmental factors such as soil characteristics, climatic conditions, irrigation management, and nutrient availability may influence both MF treatment responses and UAV-based estimation performance. Future multi-site studies spanning contrasting pedoclimatic regions would therefore provide a more robust assessment of framework transferability and practical applicability [100]. Such validation is particularly relevant within Europe, where Mediterranean, Continental, Atlantic, and Boreal production systems differ substantially in both environmental constraints and crop management practices.

A further priority concerns robust validation design. Because the physiological variables detectable by UAV are not specific to MF treatment, canopy responses must be interpreted through controlled experimental comparison: treated and control plots under common field conditions, with randomised spatial blocking to separate treatment effects from soil and microclimatic heterogeneity, and paired proximal ground-truth measurement (for example SPAD or extractive chlorophyll assay, leaf nitrogen determination, destructive biomass sampling, and plot-level yield mapping) to calibrate and confirm UAV estimates. Where machine-learning models are applied, validation must be spatially and temporally independent, using spatial cross-validation and hold-out by site or growing season, together with external-site testing, since random partitioning of spatially autocorrelated crop data can substantially inflate apparent accuracy. Reported model performance should be assessed using error magnitude (RMSE) and external validation, not the coefficient of determination alone.

A third priority concerns the integration of complementary sensing technologies. While hyperspectral, thermal, fluorescence, and proximal sensing technologies may provide additional information on MF-induced crop responses, multispectral UAV systems were selected in the present framework because they currently offer the best balance between cost, operational simplicity, sensor availability, and large-scale deployment potential [11,107]. Future studies should evaluate whether the integration of complementary sensing modalities can further improve the detection and interpretation of MF treatment effects. In particular, thermal sensing may provide additional information on crop water relations, while hyperspectral and fluorescence approaches could improve the monitoring of physiological processes that are only indirectly detectable using standard multispectral systems, a limitation that is further discussed in Section 4.6.

A recurring obstacle to synthesis in both literatures is the inconsistent reporting of methodological parameters. To support future MF-UAV validation studies and to improve comparability across trials, we recommend that such studies report, at minimum: for the MF treatment, the delivery mode, field strength and frequency, exposure duration, geometry and distance from source, and, for magnetised water, water flow rate and chemistry; the crop growth stage at treatment, soil moisture, and prevailing environmental conditions; alongside sample size, replication, plot design and randomisation, control type, and the statistical significance of reported effects; and for the UAV component, sensor type and

spectral band configuration, radiometric calibration and correction procedure, ground control and flight parameters, and validation design, including independent test-set performance, external (multi-site) validation, and assessment of model transferability across crops, sensors, and environments. Adoption of such a minimum reporting standard would substantially enhance the cumulative value of individual studies and the feasibility of future quantitative synthesis.

Beyond validation and reporting, translating UAV-based monitoring into practical agronomic benefit will require its integration with geospatial decision-support infrastructure. UAV-derived maps of chlorophyll, nitrogen status, and biomass gain operational value when combined with GIS-based environmental covariates, soil, terrain, climate, and irrigation data, and delivered through decision-support and WebGIS platforms, including land-suitability and crop-modelling approaches, that convert canopy-scale measurements into management-zone-specific recommendations [104]. Developing this decision-support layer, and coupling it to the physiological-detection framework proposed here, represents a priority for the operational maturation of MF-UAV monitoring.

Finally, future research should investigate the potential role of advanced machine-learning and sensor-fusion approaches within MF monitoring frameworks. Recent developments in artificial intelligence, deep learning, and multi-source data integration have demonstrated considerable potential for improving crop status assessment and predictive modelling from UAV data [108]. Applying these approaches to dedicated MF datasets may enhance the ability to identify treatment-related patterns, distinguish them from background environmental variability, and improve the interpretation of treatment responses across different spatial and temporal scales. Together, these research priorities define a structured pathway from conceptual framework development towards comprehensive field-scale validation and operational implementation.

4.6. Limitations

Several limitations follow from the deliberately targeted design of this review. First, the two corpora were constructed from a single database (Scopus) and restricted to English-language, peer-reviewed publications, which may have excluded relevant studies indexed elsewhere or published in other languages. Scopus was selected for its comprehensive concurrent indexing of both disciplinary literatures, and, as this is a targeted evidence synthesis rather than an exhaustive systematic review (Section 2.1), the descriptive corpus statistics reported in Section 3 (paper counts, crop distributions, and treatment-type proportions) should be read as representative of the reviewed literature rather than as exhaustive population values. The central finding of non-overlap is more robust to these restrictions, since neither search excluded the other domain's terminology and a bridging study would therefore have been retrievable in either stream had one existed (Section 3.2.1); nonetheless, this finding is properly understood as applying to the examined corpus rather than as a claim of global non-existence. Second, and by the same design logic, literatures concerned with the downstream integration of crop-monitoring data, including geospatial modelling, satellite-based monitoring, soil and climate data layers, irrigation records, and WebGIS decision-support systems, were outside the defined scope, which centred on the UAV-based detection of MF treatment responses. We note that operational field deployment would ultimately require the detection framework proposed here to be embedded within such a broader decision-support ecosystem; that integration, however, is downstream of the detection question this review addresses and is therefore deferred to future work (Section 4.5). Similarly, the synthesis did not evaluate UAV-based soil moisture estimation or irrigation-calibration literature. This is a relevant gap specifically for magnetized water irrigation, where the treatment is delivered through irrigation itself: treated and control

plots may therefore differ in soil moisture for reasons unrelated to the magnetic treatment, and since soil moisture strongly influences canopy reflectance, this represents a potential confound requiring paired soil-moisture measurement in future validation trials, with associated implications for monitoring cost.

A second set of limitations concerns the primary literature itself rather than the review method. The MF treatment literature is methodologically heterogeneous, with substantial variation in treatment protocols, field strengths, exposure conditions, and reported effect sizes, and it is likely subject to positive-outcome reporting bias. Because the framework's variable-selection criterion prioritised variables exhibiting consistent positive directional responses (Section 2.6), the bridge variables identified may over-represent favourable outcomes; the reliability of MF treatments under field conditions should not be inferred from their inclusion in the bridge table, and null or weak responses, though acknowledged (Section 1), were not systematically quantified. Relatedly, no study in the corpus reported a cost–benefit or partial-budget analysis of MF treatment, and the economic case for these technologies therefore cannot be assessed from the current evidence base.

Consistent with its status as a targeted evidence synthesis and conceptual framework rather than a formal systematic review of intervention efficacy (Section 2.1), the review did not undertake a quantitative meta-analysis or apply a formal risk-of-bias instrument of the kind developed for clinical evidence appraisal. Study quality was instead assessed against transparent criteria appropriate to a technology-integration review (Section 2.7). This approach is suited to the review's cross-disciplinary and framework-oriented objective, but it does not provide the pooled effect estimates or graded risk-of-bias assessments that a formal meta-analytic review of a single intervention would deliver.

Finally, as emphasised throughout, the proposed framework is conceptual and derives from the synthesis of two independent literatures; it has not been validated through dedicated field experiments combining MF treatment with UAV monitoring, and should be regarded as a hypothesis-driven pathway for field-scale validation rather than an experimentally verified monitoring system.

5. Conclusions

This review addressed a previously unexplored gap between two disconnected areas of agricultural research: MF technologies for crop enhancement, an emerging and methodologically heterogeneous field, and UAV-based multispectral monitoring for crop assessment, a mature and widely deployed technology. Through a dual-stream evidence synthesis of 102 MF-related studies and 114 UAV multispectral studies, it was demonstrated that these research domains have evolved largely independently despite focusing on many of the same crop species and physiological processes. No study identified within the reviewed corpus in the MF literature employed UAV-based monitoring, while no study in the UAV literature investigated magnetically treated crops.

The cross-domain evidence synthesis identified five direct bridge variables linking the two domains: chlorophyll content, NUE/nitrogen assimilation, AGB, LAI, and yield. These variables were consistently reported as MF-responsive in the agricultural literature and have also been validated for field-scale estimation using UAV multispectral sensing. Together, these bridge connections provide the scientific basis for integrating MF treatments with non-destructive, spatially explicit crop monitoring. The cross-domain variable mapping further demonstrates that MF treatment responses can be evaluated across multiple biological levels, from primary physiological processes and structural canopy development to final agronomic performance.

Based on these findings, a conceptual framework was proposed as the first operational pathway for field-scale monitoring and validation of MF treatments using UAV multi-

spectral sensing. It must be emphasised that the proposed framework is conceptual and remains to be experimentally validated. Nevertheless, it provides a structured approach for linking physiological responses to canopy-scale observations and agronomic outcomes under real farming conditions. Future empirical validation across crops, environments, and management systems will determine whether this framework can support the transition of MF technologies from promising experimental interventions to evidence-based tools for sustainable and digitally enabled agriculture. By providing a structured pathway for testing whether MF-induced physiological benefits can be reproduced under field conditions, the proposed framework contributes to the broader sustainability agenda by supporting the transparent evaluation of non-chemical crop enhancement technologies for improved resource-use efficiency, precision nutrient management, and resilient crop production.

Author Contributions: G.P.: Conceptualization, Methodology, Investigation, Data curation, Formal analysis, Project administration, Writing—original draft, Writing—review & editing, Supervision; E.G.: Methodology, Visualization, Writing—review & editing, Resources; A.O.: Methodology, Visualization, Writing—review & editing; S.F.: Conceptualization, Methodology, Resources, Supervision, Validation, Writing—review & editing; D.B.: Conceptualization, Methodology, Resources, Supervision, Validation, Writing—review & editing. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data will be made available upon request.

Acknowledgments: During the preparation of this work, the authors used ChatGPT 5.5, an AI language model by OpenAI, and Claude (Opus 4.8), an AI language model by Anthropic, to improve the language of the manuscript. After using these tools/services, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

MF	Magnetic Field
PEMF	Pulsed Electromagnetic Field
ELF	Extremely low-frequency
NR	Nitrate Reductase
Fv/Fm	PSII maximum quantum efficiency
CAT	Catalase
SMF	Static Magnetic Field
MWI	Magnetised Water Irrigation
ROS	Reactive Oxygen Species
PSII	Photosystem II
SOD	Superoxide Dismutase
APX	Ascorbate Peroxidase
N	Nitrogen
NNI	Nitrogen Nutrition Index
LAI	Leaf Area Index
CCC	Canopy Chlorophyll Content
WUE	Water Use Efficiency
NUE	Nitrogen Use Efficiency
AGB	Above-ground Biomass
LCC	Leaf Chlorophyll Content

FVC	Fractional Vegetation Cover
UAV	Unmanned Aerial Vehicle
DJI	DJI (brand name, platform manufacturer)
RGB	red-green-blue (sensor/imagery type)
DSM	Digital Surface Model
UAS	Unmanned Aerial System
P4M	DJI Phantom 4 Multispectral
NIR	Near-infrared (spectral band)
LiDAR	Light detection and ranging
NDVI	Normalised difference vegetation index
GNDVI	Green normalised difference vegetation index
VI	Vegetation Index
NDRE	Normalised difference red-edge index
SAVI	Soil-adjusted vegetation index
CWSI	Crop Water Stress Index
PAM	Photosynthetic Accumulation Model
DWT	Discrete Wavelet Transform
RTM	Radiative Transfer Model
RMSE	Root Mean Square Error
ML	Machine Learning
SVM	Support Vector Machine
LSTM	Long short-term memory
PSO	Particle Swarm Optimisation
RF	Random Forest
CNN	Convolutional Neural Network
ANN	Artificial Neural Network
XGBoost	Extreme Gradient Boosting
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
EU	European Union
CAP	Common Agricultural Policy
MODIS	Moderate Resolution Imaging Spectroradiometer
PAMF	Pulse-amplitude Modulation fluorometry

References

1. Mandouri, J.; Onat, N.C.; Kucukvar, M.; Jabbar, R.; Al-Quradaghi, S.; Al-Thani, S.; Kazançoğlu, Y. Carbon Footprint of Food Production: A Systematic Review and Meta-Analysis. *Sci. Rep.* **2025**, *15*, 35630. [CrossRef]
2. European Commission. Farm to Fork Strategy—For a Fair, Healthy and Environmentally-Friendly Food System. *Farm to Fork Strategy* 2020. Available online: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:52020DC0381> (accessed on 17 June 2026).
3. Batool, M.; Sarrazin, F.J.; Zhang, X.; Musolff, A.; Nguyen, T.V.; Attinger, S.; Kumar, R. Scenario Analysis of Nitrogen Surplus Typologies in Europe Shows That a 20% Fertilizer Reduction May Fall Short of 2030 EU Green Deal Goals. *Nat. Food* **2025**, *6*, 787–798. [CrossRef] [PubMed]
4. Ali, A.; Jabeen, N.; Farruhbek, R.; Chachar, Z.; Laghari, A.A.; Chachar, S.; Ahmed, N.; Ahmed, S.; Yang, Z. Enhancing Nitrogen Use Efficiency in Agriculture by Integrating Agronomic Practices and Genetic Advances. *Front. Plant Sci.* **2025**, *16*, 1543714. [CrossRef] [PubMed]
5. Sarraf, M.; Kataria, S.; Taimourya, H.; Santos, L.O.; Menegatti, R.D.; Jain, M.; Ihtisham, M.; Liu, S. Magnetic Field (MF) Applications in Plants: An Overview. *Plants* **2020**, *9*, 1139. [CrossRef] [PubMed]
6. Sarraf, M.; Deamici, K.M.; Taimourya, H.; Islam, M.; Kataria, S.; Raipuria, R.K.; Abdi, G.; Brestic, M. Effect of Magnetopriming on Photosynthetic Performance of Plants. *Int. J. Mol. Sci.* **2021**, *22*, 9353. [CrossRef] [PubMed]
7. Hafeez, M.B.; Zahra, N.; Ahmad, N.; Shi, Z.; Raza, A.; Wang, X.; Li, J. Growth, Physiological, Biochemical and Molecular Changes in Plants Induced by Magnetic Fields: A Review. *Plant Biol. J.* **2023**, *25*, 8–23. [CrossRef] [PubMed]
8. Čalić, D.; Ristić-Djurović, J.L.; Ćirković, S.; Milojević, J.; Belić, M.; Stanišić, M.; Zdravković-Korać, S. Overcoming Low Germination and Low Quality of Flax Seeds (*Linum usitatissimum* L.) in Unfavorable Storage Using Static Magnetic Fields. *Agriculture* **2023**, *13*, 2120. [CrossRef]

9. Pszczółkowski, P.; Sawicka, B.; Skiba, D.; Barbaś, P.; Krochmal-Marczak, B.; Ahmad, M.A. Effect of Presowing Magnetic Field Stimulation on the Seed Germination and Growth of *Phaseolus vulgaris* L. *Plants. Agronomy* **2023**, *13*, 793. [\[CrossRef\]](#)
10. Xia, X.; Pagano, A.; Macovei, A.; Padula, G.; Balestrazzi, A.; Hołubowicz, R. Magnetic Field Treatment on Horticultural and Agricultural Crops: Its Benefits and Challenges. *Folia Hortic.* **2024**, *36*, 67–80. [\[CrossRef\]](#)
11. Omia, E.; Bae, H.; Park, E.; Kim, M.S.; Baek, I.; Kabenge, I.; Cho, B.-K. Remote Sensing in Field Crop Monitoring: A Comprehensive Review of Sensor Systems, Data Analyses and Recent Advances. *Remote Sens.* **2023**, *15*, 354. [\[CrossRef\]](#)
12. Heyl, K.; Ekardt, F.; Roos, P.; Garske, B. Achieving the Nutrient Reduction Objective of the Farm to Fork Strategy. An Assessment of CAP Subsidies for Precision Fertilization and Sustainable Agricultural Practices in Germany. *Front. Sustain. Food Syst.* **2023**, *7*, 1088640. [\[CrossRef\]](#)
13. Page, M.J.; McKenzie, J.E.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. The PRISMA 2020 Statement: An Updated Guideline for Reporting Systematic Reviews. *PLoS Med.* **2021**, *18*, e1003583. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Koutsos, T.M.; Menexes, G.C.; Dordas, C.A. An Efficient Framework for Conducting Systematic Literature Reviews in Agricultural Sciences. *Sci. Total Environ.* **2019**, *682*, 106–117. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Hassanpour, H.; Ghanbarzadeh, M. Static Magnetic Field in Plants: Physiological Effects and Antioxidant Defense Mechanisms-an Overview. *Iran. J. Plant Physiol.* **2023**, *13*, 4459–4470. [\[CrossRef\]](#)
16. Kishore, G.; Saxena, C.; Rajwade, Y. Magnetic Treatment of Irrigation Water: Its Effect on Water Properties and Characteristics of Eggplant (*Solanum melongena*). *Emir. J. Food Agric.* **2022**, *34*, 784–791. [\[CrossRef\]](#)
17. Dobránszki, J. From Mystery to Reality: Magnetized Water to Tackle the Challenges of Climate Change and for Cleaner Agricultural Production. *J. Clean. Prod.* **2023**, *425*, 139077. [\[CrossRef\]](#)
18. Alattar, E.; Radwan, E.; Elwasife, K. Improvement in Growth of Plants under the Effect of Magnetized Water. *AIMS Biophys.* **2022**, *9*, 346–387. [\[CrossRef\]](#)
19. Mshenskaya, N.S.; Grinberg, M.A.; Kalyasova, E.A.; Vodeneev, V.A.; Ilin, N.V.; Slyunyaev, N.N.; Mareev, E.A.; Sinitsyna, Y.V. The Effect of an Extremely Low-Frequency Electromagnetic Field on the Drought Sensitivity of Wheat Plants. *Plants* **2023**, *12*, 826. [\[CrossRef\]](#) [\[PubMed\]](#)
20. Vashisth, A.; Meena, N.; Krishnan, P. Magnetic Field Affects Growth and Yield of Sunflower Under Different Moisture Stress Conditions. *Bioelectromagnetics* **2021**, *42*, 473–483. [\[CrossRef\]](#) [\[PubMed\]](#)
21. Suarez Rivero, D.; Ortiz Aguilar, J.; Marin Mahecha, O.; Velásquez Perilla, P.; Acevedo, P.; Santis Navarro, A. The Effect of Magnetic and Electromagnetic Fields on the Morpho-Anatomical Characteristics of Corn (*Zea mays* L.) during Biomass Production. *Chem. Eng. Trans.* **2016**, *50*, 415–420. [\[CrossRef\]](#)
22. Kataria, S.; Jain, M.; Rastogi, A.; Brestic, M. Static Magnetic Field Treatment Enhanced Photosynthetic Performance in Soybean under Supplemental Ultraviolet-B Radiation. *Photosynth. Res.* **2021**, *150*, 263–278. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Kataria, S.; Baghel, L.; Guruprasad, K.N. Alleviation of Adverse Effects of Ambient UV Stress on Growth and Some Potential Physiological Attributes in Soybean (Glycine Max) by Seed Pre-Treatment with Static Magnetic Field. *J. Plant Growth Regul.* **2017**, *36*, 550–565. [\[CrossRef\]](#)
24. Tomar, R.S.; Patel, P.; Kalal, P.R.; Laad, P.; Guruprasad, K.N.; Jajoo, A. Pre-Sowing Magnetic Treatments of Seeds Improve Photosynthetic Performance of Two Photosystems in Soybean Plants. *J. Photochem. Photobiol. B Biol.* **2025**, *267*, 113166. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Torres-Osorio, J.; Villa-Carmona, E.; Zamorano-Montanez, C. Physiological Efficiency during the Vegetative Stage of Tomato Crops Developed from Magnetically Treated Seeds. *Heliyon* **2024**, *10*, e40426. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Akrimi, R.; Mhamdi, M.; Abid, G.; Ezzeddine, H.; Nouri, I. Magnetic-Water-Enhanced Tomato Yield and Antioxidant Response Under Arid Conditions of South Tunisia. *Horticultrae* **2025**, *11*, 301. [\[CrossRef\]](#)
27. Putti, F.F.; Arruda, B.; Vicente, E.F.; Chaves, P.P.N.; Nogueira, B.B.; Zanetti, W.A.L.; Ono, E.O.; Dos Reis, A.R. Magnetic Technology to Reduce the Effects of Saline Stress on Tomato Plants. *Environ. Technol. Innov.* **2024**, *34*, 103544. [\[CrossRef\]](#)
28. Katsenios, N.; Bilalis, D.; Efthimiadou, A.; Aivalakis, G.; Nikolopoulou, A.-E.; Karkanis, A.; Travlos, I. Role of Pulsed Electromagnetic Field on Enzyme Activity, Germination, Plant Growth and Yield of Durum Wheat. *Biocatal. Agric. Biotechnol.* **2016**, *6*, 152–158. [\[CrossRef\]](#)
29. Boix, Y.F.; Dubois, A.F.; Quintero, Y.P.; Alemán, E.I.; Victório, C.P.; Aguilera, J.G.; Betancourt, M.N.; Morales-Aranibar, L. Magnetically Treated Water in *Phaseolus vulgaris* L.: An Alternative to Develop Organic Farming in Cuba. *Plants* **2023**, *12*, 340. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Roel-Rezk, V.; Lundy, M.E.; Pittelkow, C.M. Assessing Crop Nitrogen Status under Organic Amendments to Reduce Nitrogen Fertilizer Inputs and Improve Sustainability. *Nutr. Cycl. Agroecosyst.* **2025**, *131*, 365–382. [\[CrossRef\]](#)
31. Baghel, L.; Kataria, S.; Guruprasad, K.N. Static Magnetic Field Treatment of Seeds Improves Carbon and Nitrogen Metabolism under Salinity Stress in Soybean. *Bioelectromagnetics* **2016**, *37*, 455–470. [\[CrossRef\]](#) [\[PubMed\]](#)

32. Kataria, S.; Jain, M.; Tripathi, D.K.; Singh, V.P. Involvement of Nitrate Reductase-dependent Nitric Oxide Production in Magnetopriming-induced Salt Tolerance in Soybean. *Physiol. Plant.* **2020**, *168*, 422–436. [[CrossRef](#)] [[PubMed](#)]
33. Li, Z.; Gao, J.; Mu, W.; Wang, C.; Lin, X.; Wang, Z.; Yu, B.; Zhang, P.; Pincus, D.; Zheng, L.; et al. Static Magnetic Field Promotes Wheat Nitrogen Assimilation by Repressing Jasmonates Biosynthesis Through TAHY5. *Plant Biotechnol. J.* **2026**, *24*, 424–441. [[CrossRef](#)] [[PubMed](#)]
34. Kadhim, R.J.M. Effects of Magnetically Treated Irrigation Water and Exposure of the Roots to Constant Magnetic Field on Nodules Formation in the Bean Crops. *Plant Arch.* **2019**, *19*, 92–96.
35. Ma, C.; Li, Q.; Song, Z.; Su, L.; Tao, W.; Zhou, B.; Wang, Q. Irrigation with Magnetized Water Alleviates the Harmful Effect of Saline–Alkaline Stress on Rice Seedlings. *Int. J. Mol. Sci.* **2022**, *23*, 10048. [[CrossRef](#)] [[PubMed](#)]
36. Mridha, N.; Chattaraj, S.; Chakraborty, D.; Anand, A.; Aggarwal, P.; Nagarajan, S. Pre-sowing Static Magnetic Field Treatment for Improving Water and Radiation Use Efficiency in Chickpea (*Cicer arietinum* L.) under Soil Moisture Stress. *Bioelectromagnetics* **2016**, *37*, 400–408. [[CrossRef](#)] [[PubMed](#)]
37. Hozayn, M.; AbdEl-Monem, A.; El-Mahdy, A. Alleviation Salinity Stress in Germination, Seedling Vigor, Growth, Physiochemical, Yield and Nutritional Value of Chickpea (*Cicer arietinum* L) Using Magnetic Technology in Sandy Soil. *Egypt. J. Chem.* **2022**, *65*, 1317–1331. [[CrossRef](#)]
38. Wang, H.; Fan, J.; Fu, W. Effect of Activated Water Irrigation on the Yield and Water Use Efficiency of Winter Wheat under Irrigation Deficit. *Agronomy* **2022**, *12*, 1315. [[CrossRef](#)]
39. Wang, Y.; Mu, Y.; Yang, S.; Wang, L. Effects of Activated Water Irrigation on the Growth, Water Consumption and Water Use Efficiency of Winter Wheat. *Int. J. Plant Prod.* **2022**, *16*, 705–721. [[CrossRef](#)]
40. Stankevych, G.; Kovra, Y.; Borta, A. Formation of the Quality of Wheat Grain by Extremely Low Frequency Electromagnetic Field Treatment. *Technol. Audit. Prod. Reserv.* **2022**, *4*, 38–44. [[CrossRef](#)]
41. Rathod, G.R.; Anand, A. Effect of Seed Magneto-Priming on Growth, Yield and Na/K Ratio in Wheat (*Triticum aestivum* L.) under Salt Stress. *Ind. J. Plant Physiol.* **2016**, *21*, 15–22. [[CrossRef](#)]
42. Adeniran, K.A.; Kareem, K.Y.; Yusuf, K.O.; Afolayan, S.O. Effects of Electromagnetic Treatment of Irrigation Water on Growth and Yield of Lagos Spinach (*Celosia argentea*). *Agric. Eng. Int. CIGR J.* **2020**, *22*, 32–40.
43. Selim, D.A.-F.H.; Zayed, M.; Ali, M.M.E.; Eldesouky, H.S.; Bonfill, M.; El-Tahan, A.M.; Ibrahim, O.M.; El-Saadony, M.T.; El-Tarabily, K.A.; AbuQamar, S.F.; et al. Germination, Physio-Anatomical Behavior, and Productivity of Wheat Plants Irrigated with Magnetically Treated Seawater. *Front. Plant Sci.* **2022**, *13*, 923872. [[CrossRef](#)] [[PubMed](#)]
44. Vashisth, A.; Joshi, D.K. Growth Characteristics of Maize Seeds Exposed to Magnetic Field. *Bioelectromagnetics* **2017**, *38*, 151–157. [[CrossRef](#)] [[PubMed](#)]
45. Suárez-Rivero, D.; Marín-Mahecha, O.; Ortiz-Aguilar, J.; Suárez-Rivero, M.; Héctor Ardisana, E.F.; Guzmán Hernández, T.J. Use of Electromagnetic Fields as a Technological Alternative for the Production of Biomasa in *Zea mays* L. Var Porva and Reduction of the Carbon Footprint of the Crop. *Chem. Eng. Trans.* **2024**, *110*, 451–456. [[CrossRef](#)]
46. Baghel, L.; Kataria, S.; Jain, M. Mitigation of Adverse Effects of Salt Stress on Germination, Growth, Photosynthetic Efficiency and Yield in Maize (*Zea mays* L.) through Magnetopriming. *Acta Agrobot.* **2019**, *72*, 1757. [[CrossRef](#)]
47. Hozayn, M.; EL-Mahdy, A.A.; El-Dahshouri, M.F.; Darwish, M.A.A. Mitigation Salinity Stress Using Magnetically Treated Water and Chlorella Spalgae Extract on Sunflower Productivity. *Egypt. J. Chem.* **2024**. [[CrossRef](#)]
48. Dastorani, M.; Albaji, M.; Nasab, S.B. Crop Response and Water Productivity of Sunflower (*Helianthus annuus* L.) Drip-Irrigated with Magnetically Treated and Non-Magnetically Treated Water with Variable Salinity. *Water SA* **2022**, *48*, 1–7. [[CrossRef](#)]
49. Đukić, V.; Miladinov, Z.; Dozet, G.; Cvijanović, M.; Tatić, M.; Miladinović, J.; Balešević-Tubić, S. Pulsed Electromagnetic Field—A Cultivation Practice Used to Increase Soybean Seed Germination and Yield. *Zemdirb.—Agric.* **2017**, *104*, 345–352. [[CrossRef](#)]
50. Joshi-Paneri, J.; Sharma, S.; Guruprasad, K.N.; Kataria, S. Enhancing the Yield Potential of Soybean after Magneto-Priming: Detailed Study on Its Relation to Underlying Physiological Processes. *Seeds* **2023**, *2*, 60–84. [[CrossRef](#)]
51. Alsuvaid, M.; Demir, Y. The Effect of the Interaction of Magnetic Treatment and Different Irrigation Water Salinity on the Quantitative Effects on the Growth Parameters of Balkız Bean (*Phaseolus vulgaris*). *Tikrit J. Agric. Sci.* **2025**, *25*, 255–273. [[CrossRef](#)]
52. Darwish, M.A.A.; Hozayn, M.; El-Mahdy, A.A.; Abd El-Monem, A.A.; El-saway, S.S. Impact of Magnetic Technology for Mitigation Salinity Water Stress on Faba Bean (*Vicia faba* L) Growing in Sandy Soil. *Egypt. J. Chem.* **2024**, *67*, 85–95.
53. Ivankov, A.; Naučienė, Z.; Degutytė-Fomins, L.; Žūkienė, R.; Januškaitienė, I.; Malakauskienė, A.; Jakštas, V.; Ivanauskas, L.; Romanovskaja, D.; Šlepetienė, A.; et al. Changes in Agricultural Performance of Common Buckwheat Induced by Seed Treatment with Cold Plasma and Electromagnetic Field. *Appl. Sci.* **2021**, *11*, 4391. [[CrossRef](#)]
54. Chanioti, S.; Katsenios, N.; Efthimiadou, A.; Stergiou, P.; Xanthou, Z.-M.; Giannoglou, M.; Dimitrakellis, P.; Gogolides, E.; Katsaros, G. Pre-Sowing Treatment of Maize Seeds by Cold Atmospheric Plasma and Pulsed Electromagnetic Fields: Effect on Plant and Kernels Characteristics. *Aust. J. Crop Sci.* **2021**, 251–259. [[CrossRef](#)]
55. Yang, K.; Mo, J.; Luo, S.; Peng, Y.; Fang, S.; Wu, X.; Zhu, R.; Li, Y.; Yuan, N.; Zhou, C.; et al. Estimation of Rice Aboveground Biomass by UAV Imagery with Photosynthetic Accumulation Models. *Plant Phenomics* **2023**, *5*, 0056. [[CrossRef](#)] [[PubMed](#)]

56. Guo, Y.; He, J.; Zhang, H.; Shi, Z.; Wei, P.; Jing, Y.; Yang, X.; Zhang, Y.; Wang, L.; Zheng, G. Improvement of Winter Wheat Aboveground Biomass Estimation Using Digital Surface Model Information Extracted from Unmanned-Aerial-Vehicle-Based Multispectral Images. *Agriculture* **2024**, *14*, 378. [\[CrossRef\]](#)
57. Alsuvaid, M.; Demir, Y.; Kiremit, M.S.; Arslan, H. Interaction Effect of Water Magnetization and Water Salinity on Yield, Water Productivity and Morpho-Physiological of Balkız Bean (*Phaseolus vulgaris*). *Gesunde Pflanz.* **2022**, *74*, 259–274. [\[CrossRef\]](#)
58. Chakhvashvili, E.; Siegmann, B.; Muller, O.; Verrelst, J.; Bendig, J.; Kraska, T.; Rascher, U. Retrieval of Crop Variables from Proximal Multispectral UAV Image Data Using PROSAIL in Maize Canopy. *Remote Sens.* **2022**, *14*, 1247. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Eugenio, F.C.; Grohs, M.; Mallmann, C.L.; Schons, C.T.; Schuh, M.S.; Fernandes, P.; Badin, T.L.; De Souza, P.D.; Da Silva, S.D.P.; Fantinel, R.A. Influence of Flight Hours on Spectral Behavior of Rice (*Oryza sativa* L.) and Soybean (*Glycine max* L.) Cultivars in Images Obtained by Remotely Piloted Aircraft System. *J. Appl. Remote Sens.* **2021**, *15*, 044505. [\[CrossRef\]](#)
60. Ganeva, D.; Roumenina, E.; Dimitrov, P.; Gikov, A.; Jelev, G.; Dyulgenova, B.; Valcheva, D.; Bozhanova, V. Remotely Sensed Phenotypic Traits for Heritability Estimates and Grain Yield Prediction of Barley Using Multispectral Imaging from UAVs. *Sensors* **2023**, *23*, 5008. [\[CrossRef\]](#) [\[PubMed\]](#)
61. Liu, J.; Zhu, Y.; Tao, X.; Chen, X.; Li, X. Rapid Prediction of Winter Wheat Yield and Nitrogen Use Efficiency Using Consumer-Grade Unmanned Aerial Vehicles Multispectral Imagery. *Front. Plant Sci.* **2022**, *13*, 1032170. [\[CrossRef\]](#) [\[PubMed\]](#)
62. Chen, Z.; Miao, Y.; Lu, J.; Zhou, L.; Li, Y.; Zhang, H.; Lou, W.; Zhang, Z.; Kusnierek, K.; Liu, C. In-Season Diagnosis of Winter Wheat Nitrogen Status in Smallholder Farmer Fields Across a Village Using Unmanned Aerial Vehicle-Based Remote Sensing. *Agronomy* **2019**, *9*, 619. [\[CrossRef\]](#)
63. Cheng, Z.; Gu, X.; Du, Y.; Wei, C.; Xu, Y.; Zhou, Z.; Li, W.; Cai, W. Multi-Modal Fusion and Multi-Task Deep Learning for Monitoring the Growth of Film-Mulched Winter Wheat. *Precis. Agric.* **2024**, *25*, 1933–1957. [\[CrossRef\]](#)
64. Wang, J.; Wang, W.; Liu, S.; Hui, X.; Zhang, H.; Yan, H.; Maes, W.H. UAV-Based Multiple Sensors for Enhanced Data Fusion and Nitrogen Monitoring in Winter Wheat Across Growth Seasons. *Remote Sens.* **2025**, *17*, 498. [\[CrossRef\]](#)
65. Yamaguchi, T.; Sasano, K.; Katsura, K. Improving Efficiency of Ground-Truth Data Collection for UAV-Based Rice Growth Estimation Models: Investigating the Effect of Sampling Size on Model Accuracy. *Plant Prod. Sci.* **2024**, *27*, 1–13. [\[CrossRef\]](#)
66. Wang, W.; Sun, N.; Bai, B.; Wu, H.; Cheng, Y.; Geng, H.; Song, J.; Zhou, J.; Pang, Z.; Qian, S.; et al. Prediction of Wheat SPAD Using Integrated Multispectral and Support Vector Machines. *Front. Plant Sci.* **2024**, *15*, 1405068. [\[CrossRef\]](#) [\[PubMed\]](#)
67. Änäkälä, M.; Lehtilä, A.; Mäkelä, P.S.A.; Lajunen, A. Application of UAV Multispectral Imaging for Determining the Characteristics of Maize Vegetation. *Agron. Res.* **2023**, *21*, 644–653. [\[CrossRef\]](#)
68. Yan, P.; Feng, Y.; Han, Q.; Hu, Z.; Huang, X.; Su, K.; Kang, S. Enhanced Cotton Chlorophyll Content Estimation with UAV Multispectral and LiDAR Constrained SCOPE Model. *Int. J. Appl. Earth Obs. Geoinf.* **2024**, *132*, 104052. [\[CrossRef\]](#)
69. Yan, J.; Yuan, J.; Guo, Z.; Wang, W.; Li, Q.; Che, Z.; Li, G.; Ma, W.; Ma, Y.; Cheng, K.; et al. Estimation of the Three-Dimensional Spatial Distribution of Maize Leaf Chlorophyll through Fusion of Spectral, Textural, and Structural Features from Multimodal Remote Sensing. *Int. J. Appl. Earth Obs. Geoinf.* **2026**, *150*, 105355. [\[CrossRef\]](#)
70. Ai, W.; Yang, G.; Li, Z.; Du, J.; Ye, L.; Feng, X.; Jin, X.; He, Y. Two-Year Remote Sensing and Ground Verification: Estimating Chlorophyll Content in Winter Wheat Using UAV Multi-Spectral Imagery. *Artif. Intell. Agric.* **2026**, *16*, 480–494. [\[CrossRef\]](#)
71. Peter, B.G.; Messina, J.P.; Carroll, J.W.; Chikowo, R. A Case for Green-Based Vegetation Indices: Plot-Scale sUAS Imagery Related to Crop Chlorophyll Content on Smallholder Maize Farms in Malawi. *Remote Sens. Lett.* **2021**, *12*, 778–787. [\[CrossRef\]](#)
72. Zhang, C.; Yi, Y.; Wang, L.; Chen, S.; Li, P.; Zhang, S.; Xue, Y. Efficient Physics-Informed Transfer Learning to Quantify Biochemical Traits of Winter Wheat from UAV Multispectral Imagery. *Smart Agric. Technol.* **2024**, *9*, 100581. [\[CrossRef\]](#)
73. Liao, Z.; Dai, Y.; Wang, H.; Ketterings, Q.M.; Lu, J.; Zhang, F.; Li, Z.; Fan, J. A Double-Layer Model for Improving the Estimation of Wheat Canopy Nitrogen Content from Unmanned Aerial Vehicle Multispectral Imagery. *J. Integr. Agric.* **2023**, *22*, 2248–2270. [\[CrossRef\]](#)
74. Li, Z.; Zhang, Y.; Lu, J.; Wang, Y.; Zhao, C.; Wang, W.; Wang, J.; Zhang, H.; Huo, Z. Better Inversion of Rice Nitrogen Nutrition Index at Early Panicle Initiation Stage Using Spectral Features, Texture Features, and Wavelet Features Based on UAV Multispectral Imagery. *Eur. J. Agron.* **2025**, *168*, 127654. [\[CrossRef\]](#)
75. Liang, T.; Duan, B.; Luo, X.; Ma, Y.; Yuan, Z.; Zhu, R.; Peng, Y.; Gong, Y.; Fang, S.; Wu, X. Identification of High Nitrogen Use Efficiency Phenotype in Rice (*Oryza sativa* L.) Through Entire Growth Duration by Unmanned Aerial Vehicle Multispectral Imagery. *Front. Plant Sci.* **2021**, *12*, 740414. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Jansone, Z.; Bleidere, M.; Putniece, G. Estimating Spring Wheat Nitrogen Use Efficiency via Proximal and UAV Sensing in Northwest Latvia. *Agron. Res.* **2026**, *24*, 27–45. [\[CrossRef\]](#)
77. Wang, J.; Meyer, S.T.; Xu, X.; Weisser, W.W.; Yu, K. Drone Multispectral Imaging Captures the Effects of Soil Mineral Nitrogen on Canopy Structure and Nitrogen Use Efficiency in Wheat. *Comput. Electron. Agric.* **2025**, *235*, 110342. [\[CrossRef\]](#)
78. Song, X.; Yang, G.; Xu, X.; Zhang, D.; Yang, C.; Feng, H. Winter Wheat Nitrogen Estimation Based on Ground-Level and UAV-Mounted Sensors. *Sensors* **2022**, *22*, 549. [\[CrossRef\]](#) [\[PubMed\]](#)

79. Lu, N.; Wang, W.; Zhang, Q.; Li, D.; Yao, X.; Tian, Y.; Zhu, Y.; Cao, W.; Baret, F.; Liu, S.; et al. Estimation of Nitrogen Nutrition Status in Winter Wheat From Unmanned Aerial Vehicle Based Multi-Angular Multispectral Imagery. *Front. Plant Sci.* **2019**, *10*, 1601. [CrossRef] [PubMed]
80. Zhang, M.; Zhao, W.; Li, J. Determination of UAV Flight Altitude and Time for Optimizing Variable-Rate Nitrogen Prescription Maps for Winter Wheat in the North China Plain. *Agronomy* **2025**, *15*, 2627. [CrossRef]
81. Argento, F.; Anken, T.; Abt, F.; Vogelsanger, E.; Walter, A.; Liebisch, F. Site-Specific Nitrogen Management in Winter Wheat Supported by Low-Altitude Remote Sensing and Soil Data. *Precis. Agric.* **2021**, *22*, 364–386. [CrossRef]
82. Ma, Z.; Xu, D.; Qiu, Z.; Fang, L.; Wang, M.; Wang, Z.; Bao, R.; Tang, Q.; Song, X.; Tan, C.; et al. Integrating UAV, Environmental, and Management Data to Improve Rice Nitrogen Nutrition Index Prediction Using an Ensemble Learning Algorithm. *BMC Plant Biol.* **2026**, *26*, 762. [CrossRef] [PubMed]
83. Zhang, J.; Zhao, Y.; Hu, Z.; Xiao, W. Unmanned Aerial System-Based Wheat Biomass Estimation Using Multispectral, Structural and Meteorological Data. *Agriculture* **2023**, *13*, 1621. [CrossRef]
84. Feizolahpour, F.; Besharat, S.; Feizizadeh, B.; Rezaverdinejad, V.; Hessari, B. An Integrative Data-Driven Approach for Monitoring Corn Biomass under Irrigation Water and Nitrogen Levels Based on UAV-Based Imagery. *Environ. Monit. Assess.* **2023**, *195*, 1081. [CrossRef] [PubMed]
85. Dlamini, C.M.; Odindi, J.; Matongera, T.N.; Mutanga, O. The Use of Unmanned Aerial Vehicle (UAV) Remotely Sensed Data and Biophysical Variables to Predict Maize Above-Ground Biomass (AGB) in Small-Scale Farming Systems. *Remote Sens. Appl. Soc. Environ.* **2025**, *39*, 101706. [CrossRef]
86. Kenenbayev, S.B. UAV-Based Multispectral Assessment of Soil Degradation and Grain Crop Yields in Southeastern Kazakhstan. *Int. J. Agric. Biosci.* **2025**, *15*, 782–789. [CrossRef]
87. Kuang, X.; Hou, X.; Wang, D.; Mao, B.; Li, Y.; Chen, D.; Fu, W.; Cheng, Q.; Duan, F.; Li, H.; et al. A CNN-LSTM-XGBoost Hybrid Framework for Interpretable Nitrogen Stress Classification Using Multimodal UAV Imagery. *Remote Sens.* **2026**, *18*, 538. [CrossRef]
88. Wang, T.; Yang, G.; Xu, X.; Sun, J.; Meng, Y.; Yang, X.; Feng, H.; Xue, H.; Xu, X.; Song, Y. Remote Sensing Estimation of Rice Chlorophyll Content Based on UAV Image Feature Selection and PSO-Optimized Ensemble Learning. *Artif. Intell. Geosci.* **2026**, *7*, 100190. [CrossRef]
89. Yao, Y.; Wang, D.; Shu, M.; Hu, J.; Liu, Y.; Feng, H.; Li, B.; Lu, W.; Wang, J.; Xi, L.; et al. Estimation of Maize LCC and FVC for Full Growth Periods Based on UAV Remote Sensing and Ensemble Learning. *PeerJ Comput. Sci.* **2025**, *11*, e3356. [CrossRef]
90. Zhang, Y.; Li, W.; Jia, W.; Zhang, M.; Tao, R.; Liang, S. Cross-Domain Hyperspectral Image Classification Based on Bi-Directional Domain Adaptation. *IEEE Trans. Circuits Syst. Video Technol.* **2025**, *35*, 12038–12051. [CrossRef]
91. Zhang, Y.; Li, W.; Zhang, M.; Han, J.; Tao, R.; Liang, S. SpectralX: Parameter-Efficient Domain Generalization for Spectral Remote Sensing Foundation Models. *ISPRS J. Photogramm. Remote Sens.* **2026**, *239*, 774–792. [CrossRef]
92. Singh-Bakala, H.; Ravelombola, F.; Washburn, J.D.; Shannon, G.; Zhang, R.; Lin, F. Photosynthetic and Canopy Trait Characterization in Soybean (*Glycine max* L.) Using Chlorophyll Fluorescence and UAV Imaging. *Agriculture* **2025**, *15*, 2576. [CrossRef]
93. Yusuf, K.O.; Ogunbamowo, T.R.; Obalowu, R.O. Effect of Magnetized Water on Water Use Efficiency, Yield and Nutritional Qualities of Watermelon under Deficit Irrigation. *Agric. Eng. Int. CIGR J.* **2020**, *22*, 51–60.
94. Kiranmai, Y.; Potdar, M.P.; Biradar, D.P.; Kuligod, V.B.; Kanade, A.K.; Parmar, B.; Malunekar, V. Moisture Stress Assessment in Rabi Maize through UAV-Mounted Multispectral Sensor. *Front. Environ. Sci.* **2025**, *13*, 1651422. [CrossRef]
95. Yang, N.; Zhang, Z.; Yang, X.; Zhang, J.; Zhang, B.; Xie, P.; Wang, Y.; Chen, J.; Shi, L. UAV-Based Stomatal Conductance Estimation under Water Stress Using the PROSAIL Model Coupled with Meteorological Factors. *Int. J. Appl. Earth Obs. Geoinf.* **2025**, *137*, 104425. [CrossRef]
96. Yang, M.; Hassan, M.A.; Xu, K.; Zheng, C.; Rasheed, A.; Zhang, Y.; Jin, X.; Xia, X.; Xiao, Y.; He, Z. Assessment of Water and Nitrogen Use Efficiencies Through UAV-Based Multispectral Phenotyping in Winter Wheat. *Front. Plant Sci.* **2020**, *11*, 927. [CrossRef] [PubMed]
97. Sinclair, T.R.; Horie, T. Leaf Nitrogen, Photosynthesis, and Crop Radiation Use Efficiency: A Review. *Crop Sci.* **1989**, *29*, 90–98. [CrossRef]
98. Gaju, O.; DeSilva, J.; Carvalho, P.; Hawkesford, M.J.; Griffiths, S.; Greenland, A.; Foulkes, M.J. Leaf Photosynthesis and Associations with Grain Yield, Biomass and Nitrogen-Use Efficiency in Landraces, Synthetic-Derived Lines and Cultivars in Wheat. *Field Crops Res.* **2016**, *193*, 1–15. [CrossRef]
99. Tang, Z.; Guo, J.; Xiang, Y.; Lu, X.; Wang, Q.; Wang, H.; Cheng, M.; Wang, H.; Wang, X.; An, J.; et al. Estimation of Leaf Area Index and Above-Ground Biomass of Winter Wheat Based on Optimal Spectral Index. *Agronomy* **2022**, *12*, 1729. [CrossRef]
100. European Commission. EU Soil Strategy for 2030. 2021. Available online: https://environment.ec.europa.eu/topics/soil-health/soil-strategy-2030_en (accessed on 29 July 2026).
101. Guebsi, R.; Mami, S.; Chokmani, K. Drones in Precision Agriculture: A Comprehensive Review of Applications, Technologies, and Challenges. *Drones* **2024**, *8*, 686. [CrossRef]

102. Vullaganti, N.; Ram, B.G.; Sun, X. Precision Agriculture Technologies for Soil Site-Specific Nutrient Management: A Comprehensive Review. *Artif. Intell. Agric.* **2025**, *15*, 147–161. [[CrossRef](#)]
103. Qiu, Z.; Ma, F.; Zhou, J.; Du, C. Improving Rice Nitrogen Nutrition Index Estimation Using UAV Images Combined with Meteorological and Fertilization Variables. *Agronomy* **2025**, *15*, 1946. [[CrossRef](#)]
104. Kim, J.S.; Kisekka, I. FARMS: A Geospatial Crop Modeling and Agricultural Water Management System. *ISPRS Int. J. Geo-Inf.* **2021**, *10*, 553. [[CrossRef](#)]
105. Storm, H.; Seidel, S.J.; Klingbeil, L.; Ewert, F.; Vereecken, H.; Amelung, W.; Behnke, S.; Bennewitz, M.; Börner, J.; Döring, T.; et al. Research Priorities to Leverage Smart Digital Technologies for Sustainable Crop Production. *Eur. J. Agron.* **2024**, *156*, 127178. [[CrossRef](#)]
106. Georgescu, P.-L.; Barbuta-Misu, N.; Zlati, M.L.; Fortea, C.; Antohi, V.M. Quantifying the Performance of European Agriculture Through the New European Sustainability Model. *Agriculture* **2025**, *15*, 210. [[CrossRef](#)]
107. Tsouros, D.C.; Bibi, S.; Sarigiannidis, P.G. A Review on UAV-Based Applications for Precision Agriculture. *Information* **2019**, *10*, 349. [[CrossRef](#)]
108. Dong, Y.; Huang, W.; Li, H.; Han, L. Editorial: Advancements in Agricultural Monitoring with AI Enhanced Remote Sensing Techniques. *Front. Remote Sens.* **2025**, *6*, 1664060. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.