

Phthalate Exposure: From Quantification to Risk Assessment

Lidia Caporossi ^{1,*} and Maria Marino ^{2,*}

¹ INAIL, Department of Occupational and Environmental Medicine, Epidemiology and Hygiene, Via Fontana Candida 1, 00078 Monteporzio Catone, Italy

² Department of Science, University of Rome Tre, Viale G. Marconi, 446, 00146 Rome, Italy

* Correspondence: l.caporossi@inail.it (L.C.); maria.marino@uniroma3.it (M.M.)

Phthalates (di-esters of phthalic acid) are a group of synthetic organic compounds present in the environment because of their wide use in a multitude of important industrial products (e.g., cosmetics, medical devices, plastics, and food packages), mainly as plasticizers to improve mechanical properties such as flexibility, transparency, durability, and longevity of polyvinyl chloride. The widespread occurrence of phthalates in the environment as well as their endocrine-disrupting and reproductive effects contribute to their recognition as substances of high concern [1–4]. In this Special Issue of *Toxics* entitled “Phthalate Exposure: From Quantification to Risk Assessment”, we collected both reviews and articles to identify the levels of these substances in the environment and in other sources of human exposure (e.g., occupational), their toxic effects, and their action mechanisms. Special attention was paid to analytical methods able to quantify low concentrations of phthalates in different complex matrixes, particularly in human biomonitoring.

Exposure to phthalates occurs through contaminated food ingestion; through skin absorption, i.e., in the case of cosmetic creams; or through inhalation [2]. A particular type of exposure is that which occurs by migration of phthalates from medical devices made of plastic, such as drip bags or tubes for therapies. The wide presence of these substances in living environments as well as in specific workplaces has allowed us to highlight their generalized exposure, with registered dosages in the general population, through surveys of biomonitoring in both the United States and Europe [5,6]. Women of childbearing age seem to have higher urinary phthalate levels than men, probably due to the use of personal care products [7]. The paper by Caporossi and colleagues [8] provided evidence of a relationship between the possibility of pregnancy loss, and exposure to diethylhexyl phthalate (DEHP) and diethyl phthalate (DEP). The presence of significantly higher values of certain phthalates, DEP metabolite in particular, especially in women with recurrent pregnancy loss and idiopathic infertility, suggests a possible involvement of these compounds as competing factors in reproductive issues [8]. These data require specific attention due to phthalate being widely used and to the absence of particular law restriction in Europe for DEP. Furthermore, studies carried out on male populations suggested that even for seminal fluid quality, phthalate exposure could be deleterious [9]. In particular, exposure to DEP seems to be linked with a reduction in semen concentration. Moreover, this type of exposure could also lead to increased oxidative stress, even in seminal fluid, because the production of oxidative stress is one of the possible mechanisms of endocrine dysfunction owing to chemicals, probably linked to an interference in antioxidant protective mechanisms [10,11]. Indeed, some researchers proposed reactive oxygen species (ROS) as possible biomarkers of toxicity [12]. The paper by Pignini and collaborators [13] documented the correlation between phthalates and ROS in two hundred and seventy-four couples undergoing an assisted reproduction technology treatment. In particular, 8-oxo-7,8-dihydro-2-deoxyguanosine (8-oxo-dGuo) was found to be associated with higher levels of DEHP, DnBP, and diisobutylphthalate

Citation: Caporossi, L.; Marino, M. Phthalate Exposure: From Quantification to Risk Assessment. *Toxics* **2022**, *10*, 330. <https://doi.org/10.3390/toxics10060330>

Received: 9 June 2022

Accepted: 15 June 2022

Published: 16 June 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

(DiBP) [13,14], and 8-oxo-7,8-guanosine (8-oxo-Guo) emerged as a really sensitive marker of oxidative stress in this type of exposure [15]. Of particular interest was the studies by Jenkins and colleagues carried out in the Neonatal Intensive Care Units [16], where the possibility of the exposure of an extremely fragile population to the pollutants released from medical devices was tested.

Zebrafish is a model widely applied in toxicology to estimated possible developmental risks for humans and for exposure to endocrine disrupter toxics [17,18]. The paper by Kwang and colleagues [19] collected data obtained from acute and chronic exposure of zebrafish to DEHP. This experimental model allowed them to study the toxic effects of phthalates starting from the embryonic stage to different developmental stages, thus obtaining a model that could facilitate risk assessments of DEHP exposure. Closely related to phthalate toxic effects is the paper by Liu and collaborators [20], who developed a 3D-QSAR model based on the investigation of the phthalates effects on nuclear factor erythroid 2 related factor 2 (NRF2), a nuclear transcription factor that regulates oxidative stress and plays a crucial role in liver detoxification. The experiments performed in human hepatocyte (HepG2) cells could help predict the toxicity of new analogues and provide a reference for the design of less toxic phthalates PAEs in the future.

The mechanisms at the root of the endocrine disruptor effects of phthalates has been addressed in the paper by Fiocchetti and colleagues [21]. These authors demonstrated that DEP could induce estrogen receptor α subtype (ER α) phosphorylation and degradation even without direct binding. In this way, DEP modulates estrogenic signaling through the indirect induction of changes in conformation in ER α and the corresponding functional outcomes and transduction mechanisms [21].

Finally, the validation and use of an analytical method useful for specific determination, for animals [22], for humans, and for studying the possible route of exposure are core questions for obtaining valuable findings. The analysis of phthalates represents a challenge for numerous researchers due to their, usually, low concentrations in different matrixes (food, human urine, different materials such as gloves [23], animal blood [22], and environmental matrixes). Furthermore, a specific complexity is linked to possible cross-contamination in the laboratory during the steps of quantification owing to plastic materials being used for laboratory analysis. During the years, for the pretreatment of samples, an evolution focused on quickness, simplicity, low sample handling, and the optimization of the use of solvents and reagents was documented. Both solid phase extraction (SPE) and liquid/liquid extract (LLE) were developed for the extraction of phthalates from different matrixes. Regarding analytical methods, the chromatographic techniques were the most commonly used, usually gas chromatography or high performance liquid chromatography (HPLC) coupled with mass spectrometry (MS) or triple quadrupole mass spectrometry (MS/MS) [23,24].

In Europe, the HBM4EU biomonitoring project was implemented [25] to understand background levels of phthalates in the general population due to daily environmental exposure. Urinary phthalate levels were analyzed in the recruited subjects. Within this project, reported in the paper by Mol and colleagues [26], the analytical, validation, and quality aspects of the data were crucial, and therefore, inter-laboratory evaluations were conducted to ensure the quality of the data produced.

In conclusion, the adverse effects of phthalates on animals and humans were highlighted in this Special Issue. In particular, the reprotoxicity of phthalates clearly emerged. Given the prevalence of infertility in the human population, estimated at about 15%, investigating any possible risk factors is significant. Furthermore, the industrial production of phthalates occurs, again, worldwide at very consistent levels; this call for further studies to promote healthy workplaces and to reduce possible environmental contamination. Because of the daily use of personal care products, the communication of potential health risks for consumers represents a specific target for future actions.

Funding: this research received no external funding.

Conflicts of Interest: authors declare no conflict of interest.

References

- Wang, Y.; Zhu, H.; Kannan, K. A Review of Biomonitoring of Phthalate Exposures. *Toxics* **2019**, *7*, 21.
- Latini, G. Monitoring phthalate exposure in humans. *Clin. Chim. Acta* **2005**, *361*, 20–29.
- Mallow, E.B.; Fox, M.A. Phthalates and critically ill neonates: Device-related exposures and non-endocrine toxic risks. *J. Perinatol.* **2014**, *34*, 892–897.
- Yu, K.; Li, G.; Feng, W.; Liu, L.; Zhang, J.; Wu, W.; Xu, L.; Yan, Y. Chlorpyrifos is Estrogenic and Alters Embryonic Hatching, Cell Proliferation and Apoptosis in Zebrafish. *Chem. Biol. Interact.* **2015**, *239*, 26–33.
- Blount, B.C.; Silva, M.J.; Caudill, S.P.; Needham, L.L.; Pirkle, J.L.; Sampson, E.J.; Lucier, G.W.; Jackson, R.J.; Brock, J.W. Levels of seven urinary phthalate metabolites in a human reference population. *Environ. Health Perspect.* **2000**, *108*, 979–982.
- Koch, H.M.; Rossbach, B.; Drexler, H.; Angerer, J. Internal exposure of the general population to DEHP and other phthalates determination of secondary and primary phthalate monoester metabolites in urine. *Environ. Res.* **2003**, *93*, 177–185.
- Huang, P.C.; Liao, K.W.; Chang, J.W.; Chan, S.H.; Lee, C.C. Characterization of phthalates exposure and risk for cosmetics and perfume sales clerks. *Environ. Pollut.* **2018**, *233*, 577–587.
- Caporossi, L.; Viganò, P.; Paci, E.; Capanna, S.; Alteri, A.; Campo, G.; Pignini, D.; De Rosa, M.; Tranfo, G.; Papaleo, B. Female Reproductive Health and Exposure to Phthalates and Bisphenol A: A Cross Sectional Study. *Toxics* **2021**, *9*, 299.
- Caporossi, L.; Alteri, A.; Campo, G.; Paci, E.; Tranfo, G.; Capanna, S.; Papaleo, E.; Pignini, D.; Viganò, P.; Papaleo, B. Cross sectional study on exposure to BPA and phthalates and semen parameters in men attending a fertility center. *Int. J. Environ. Res. Public Health* **2020**, *17*, 489.
- Minamiyama, Y.; Ichikawa, H.; Takemura, S.; Kusunoki, H.; Naito, Y.; Yoshikawa, T. Generation of reactive oxygen species in sperms of rats as an earlier marker for evaluating the toxicity of endocrine-disrupting chemicals. *Free Radic. Res.* **2010**, *44*, 1398–1406.
- Botelho, G.G.K.; Bufalo, A.C.; Boareto, A.C.; Muller, J.C.; Morais, R.N.; Martino-Andrade, A.J.; Lemos, K.R.; Dalsenter, P.R. Vitamin C and Resveratrol Supplementation to Rat Dams Treated with Di(2-ethylhexyl)phthalate: Impact on Reproductive and Oxidative Stress End Points in Male Offspring. *Arch. Environ. Contam. Toxicol.* **2009**, *57*, 785–793.
- Devine, P.J.; Perreault, S.D.; Luderer, U. Roles of Reactive Oxygen Species and Antioxidants in Ovarian Toxicity. *Biol. Reprod.* **2012**, *86*, 1–10.
- Pignini, D.; Caporossi, L.; Paci, E.; Capanna, S.; Viganò, P.; Alteri, A.; Rabellotti, E.; Buonauro, F.; Papaleo, B.; Tranfo, G. Phthalate Exposure and Biomarkers of Oxidation of Nucleic Acids: Results on Couples Attending a Fertility Center. *Toxics* **2022**, *10*, 61.
- Guo, Y.; Weck, J.; Sundaram, R.; Goldstone, A.E.; Louis, G.B.; Kannan, K. Urinary Concentrations of Phthalates in Couples Planning Pregnancy and Its Association with 8-Hydroxy-20-deoxyguanosine, a Biomarker of Oxidative Stress: Longitudinal Investigation of Fertility and the Environment Study. *Environ. Sci. Technol.* **2014**, *48*, 9804–9811.
- Osburn, W.O.; Kensler, T.W. Nrf2 signaling: An adaptive response pathway for protection against environmental toxic insults. *Mutat. Res.* **2008**, *659*, 31–39.
- Jenkins, R.; Ondusko, D.; Montrose, L.; Forbush, R.; Rozansky, D. Phthalate Exposures in the Neonatal Intensive Care Unit. *Toxics* **2021**, *9*, 90. <https://doi.org/10.3390/toxics9050090>.
- Moreman, J.; Lee, O.; Trznadel, M.; David, A.; Kudoh, T.; Tyler, C.R. Acute Toxicity, Teratogenic, and Estrogenic Effects of Bisphenol A and Its Alternative Replacements Bisphenol S, Bisphenol F, and Bisphenol AF in Zebrafish Embryo-Larvae. *Environ. Sci. Technol.* **2017**, *51*, 12796–12805.
- Love, D.R.; Pichler, F.B.; Dodd, A.; Copp, B.R.; Greenwood, D.R. Technology for high-throughput screens: The present and future using zebrafish. *Curr. Opin. Biotechnol.* **2004**, *15*, 564–571.
- Kwan, W.S.; Roy, V.A.L.; Yu, K.N. Review on Toxic Effects of Di(2-ethylhexyl) Phthalate on Zebrafish Embryos. *Toxics* **2021**, *9*, 193.
- Liu, H.; Huang, H.; Xiao, X.; Zhao, Z.; Liu, C. Effects of Phthalate Esters (PAEs) on Cell Viability and Nrf2 of HepG2 and 3D-QSAR Studies. *Toxics* **2021**, *9*, 134. <https://doi.org/10.3390/toxics9060134>.
- Fiocchetti, M.; Bastari, G.; Cipolletti, M.; Leone, S.; Acconcia, F.; Marino, M. The Peculiar Estrogenicity of Diethyl Phthalate: Modulation of Estrogen Receptor α Activities in the Proliferation of Breast Cancer Cells. *Toxics* **2021**, *9*, 237.
- Notardonato, I.; Di Fiore, C.; Iannone, A.; Russo, M.V.; Blasi, M.F.; Favero, G.; Mattei, D.; Protano, C.; Vitali, M.; Avino, P. Fast and Reliable Determination of Phthalic Acid Esters in the Blood of Marine Turtles by Means of Solid Phase Extraction Coupled with Gas Chromatography-Ion Trap/Mass Spectrometry. *Toxics* **2021**, *9*, 279.
- Poitou, K.; Rogez-Florent, T.; Lecoq, M.; Danel, C.; Regnault, R.; Vérité, P.; Monteil, C.; Foulon, C. Analysis of Phthalates and Alternative Plasticizers in Gloves by as Chromatography–Mass Spectrometry and Liquid Chromatography–UV Detection: A Comparative Study. *Toxics* **2021**, *9*, 200.

24. Luís, C.; Algarra, M.; Câmara, J.S.; Perestrelo, R. Comprehensive Insight from Phthalates Occurrence: From Health Outcomes to Emerging Analytical Approaches. *Toxics* **2021**, *9*, 157.
25. Gillesa, L.; Govartsa, E.; Rambaud, L.; Vogel, N.; Castaño, A.; Esteban López, M.; Rodriguez, L.M.; Koppen, G.; Remy, S.; Vrijheid, M.; et al. HBM4EU combines and harmonises human biomonitoring data across the EU, building on existing capacity—The HBM4EU survey. *Int. J. Hyg. Environ. Health* **2021**, *237*, 113809.
26. Mol, H.G.J.; Elbers, I.; Pålme, C.; Bury, D.; Göen, T.; López, M.E.; Nübler, S.; Vaccher, V.; Antignac, J.-P.; Dvoráková, D.; et al. Proficiency and Interlaboratory Variability in the Determination of Phthalate and DINCH Biomarkers in Human Urine: Results from the HBM4EU Project. *Toxics* **2022**, *10*, 57.