

Review

Identification and Detection of Aflatoxigenic *Aspergillus flavus* and Aflatoxins: Current Technologies and Future Perspectives

Md Mostafa Masud ¹, Sumyya Waliullah ², Abdus Sobhan ¹ and Emran Ali ^{1,*} ¹ Department of Agriculture, Alcorn State University, Lorman, MS 39096, USA² Department of Biological Sciences, Alcorn State University, Lorman, MS 39096, USA

* Correspondence: eali@alcorn.edu

Abstract

Aspergillus flavus is a major fungal species that contaminates a wide range of crops and produces aflatoxins under favorable environmental conditions. Several approaches have been developed for aflatoxigenic *A. flavus* identification and aflatoxins detection. This review summarizes the existing methods for the identification of aflatoxigenic *A. flavus*, along with recent progress in aflatoxins detection methods. It also highlights limitations and possible factors associated with the accurate identification of aflatoxigenic isolates. Particularly, this review underscores the utilization of advanced analytical or detection methods along with existing identification techniques to accurately identify aflatoxigenic *A. flavus*.

Keywords: aflatoxigenic *Aspergillus flavus*; aflatoxins detection; identification; biosynthesis gene cluster; aflatoxins

Key Contribution: 1. This paper highlights current molecular approaches for identifying aflatoxigenic *A. flavus*, along with recent quantification techniques for aflatoxins detection. 2. Evaluates conventional and advanced molecular techniques for identifying aflatoxigenic *A. flavus*, highlighting their strengths and limitations. 3. Analyzes all the possible factors influencing aflatoxin production and discusses their implications for the reliable identification of aflatoxigenic *A. flavus*. 4. Explores the integration of both identification and detection approaches for accurate identification of aflatoxigenic *A. flavus* isolates.

1. Literature Search Strategy

A comprehensive search was performed across different electronic databases such as Google Scholar, PubMed, Web of Science, and ScienceDirect. The search included the published literature between 2007 and 2024, with a focus on existing methods for identification of aflatoxigenic *A. flavus*. Also, the literature included recent detection methods for aflatoxin quantification, particularly highlighting published articles between 2018 and 2025. Boolean operators were utilized to find the articles in the following terms: (“*Aspergillus flavus*” OR “*A. flavus*”) AND (aflatoxigenic OR toxigenic) AND (identification), (“*Aspergillus flavus*” OR “*A. flavus*”) AND (“aflatoxigenic” OR “toxigenic”) AND (“identification”) AND (“PCR” OR “RT-PCR” OR “RT-qPCR” OR “real-time” OR “LAMP”), ((“*Aspergillus flavus*” OR “*A. flavus*”) AND “toxigenic” AND “molecular” AND “detection”), (“aflatoxin detection” OR “prediction” OR “mycotoxin detection”) AND (“biosensor” OR “artificial intelligence” OR “machine learning”), (“mycotoxin detection” AND “recent advances” OR “recent progress”).



Received: 26 June 2026

Revised: 29 July 2026

Accepted: 31 July 2026

Published: 4 August 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

2. Introduction

Aspergillus flavus is a major fungal contaminant of food and feed commodities, posing significant risks to both human and animal health. As an opportunistic and airborne pathogen, it readily infects agricultural products, particularly grains and nuts, under warm and humid environmental conditions. Consequently, *A. flavus* has emerged as a global concern, especially in tropical, subtropical, and Mediterranean regions where climatic conditions favor fungal growth and toxin production [1].

This pathogen can contaminate crops at multiple stages of the food chain, including pre-harvest, post-harvest, storage, and transportation [2,3]. A key concern associated with *A. flavus* is its ability to produce aflatoxins (AFs), a group of highly toxic secondary metabolites that compromise food safety and quality [4]. Among them, aflatoxin B1 (AFB1) is the most potent and is classified as a Group 1 carcinogen by the International Agency for Research on Cancer (IARC) [5]. In addition to AFB1, *A. flavus* produces AFB2 and other secondary metabolites such as cyclopiazonic acid. In contrast, AFM1 is not directly produced by the fungus but is a hydroxylated metabolite of AFB1 formed in animals such as poultry and livestock and commonly detected in milk following ingestion of contaminated feed [6].

AFs contamination is associated with substantial economic losses worldwide. In the United States alone, annual losses in the corn industry due to AFs contamination range from millions to billions of dollars [7]. Beyond economic impacts, AFs pose serious health risks, including hepatotoxicity, immunosuppression, and hepatocellular carcinoma, particularly in individuals co-infected with hepatitis B virus [6,8–18]. Therefore, effective monitoring and accurate detection of aflatoxigenic *A. flavus* are critical for ensuring food safety and mitigating public health risks.

Traditionally, identification of *A. flavus* has relied on morphological and cultural characteristics, including colony morphology, pigmentation, and microscopic features [19,20]. However, these methods are labor-intensive, time-consuming, and often lack the sensitivity and specificity required to distinguish between aflatoxigenic and atoxigenic strains [21,22]. Moreover, there are no reliable morphological markers that can differentiate aflatoxigenic isolates from non-aflatoxigenic ones.

To overcome these limitations, molecular approaches have been widely adopted for the detection and characterization of aflatoxigenic *A. flavus*. These methods typically target genes within the AFs biosynthesis gene cluster, which is responsible for toxin production. Although molecular techniques such as PCR and sequencing offer higher sensitivity and specificity [21,23–25], challenges remain due to genetic variability, the absence or mutation of specific genes, and the inconsistent relationship between gene presence, gene expression, and actual AFs production [26–29].

This review mainly focuses on the traditional and existing methods that have been widely used for the identification of aflatoxigenic *A. flavus* and highlights the potential limitations associated with this method. Even though an isolate of *A. flavus* is identified through morphological and molecular methods, it cannot produce AFs in real time. Therefore, this review also highlights how the identification technique could be well adapted along with the AFs detection technique to accurately predict the aflatoxigenic *A. flavus* isolates. Moreover, several factors need to be considered while measuring AFs detection because they could influence the final concentration of AFs accumulation in food and feed crops.

Therefore, a comprehensive understanding of existing detection methods, along with their limitations, is essential for developing reliable and efficient diagnostic tools. It also emphasizes the need for integrated, multi-method strategies to enhance the accuracy and reliability of detection in both laboratory and field settings.

3. Traditional Approaches/Cultural Techniques

Morphological identification is one of the earliest and most widely used approaches for detecting fungal pathogens, including *A. flavus*. This method relies on observable characteristics such as colony morphology, growth pattern, pigment production, hyphal structure, and conidial formation on culture media [30–34]. Based on these traits, *A. flavus* isolates can be preliminarily identified by culturing on different types of media, each designed to support specific growth or analytical objectives (Table 1).

Different culture media are selected depending on the purpose of the study, including isolation, identification, toxin production assessment, and gene expression analysis. As summarized in Table 1, media such as *A. flavus* and *parasiticus* agar (AFPA), Aflatoxins-producing ability medium (APA), coconut-based media (CMA, CEA, CCA), Czapek yeast extract agar (CYA), and yeast extract sucrose (YES) media are widely used for culturing *Aspergillus* section *Flavi* under varying experimental conditions [35].

Despite its widespread use, morphological and cultural identification has several limitations. These methods require considerable time for media preparation, fungal growth, and microscopic examination. Additionally, accurate identification often requires substantial mycological expertise. More importantly, morphological approaches lack reliability due to significant intraspecific and interspecific variability among *Aspergillus* species [36]. As a result, distinguishing aflatoxigenic and non-aflatoxigenic strains based solely on morphology is not feasible.

To address these limitations, several culture-based techniques have been developed to identify aflatoxigenic and non-aflatoxigenic isolates based on their ability to produce AFs. These include the ammonia vapor test and fluorescence-based detection under UV light, which are discussed below.

Table 1. Different types of media that have been utilized by several researchers for culturing *Aspergillus* section *Flavi*.

Media	Application	References
AFPA	Selective media for the visual detection of <i>A. flavus</i> and <i>A. parasiticus</i> , Visual differentiation of aflatoxigenic and non-aflatoxigenic <i>A. flavus</i>	[16,37,38]
CYA, YES, CAM, APA, CCA, CMA, AFPA	Useful for growing aflatoxigenic <i>A. flavus</i> and determining AFs production capability.	[39–43]
DRBC, MRBA	Isolation and identification of <i>Aspergillus</i> spp. from soil	[16,44,45]
YESD	Better visual identification of aflatoxigenic <i>A. flavus</i> strains	[38]
MEA, PDA, YES, CYA	Visualization of the differences in morphological characteristics in <i>A. flavus</i> and <i>A. oryzae</i> , such as colony color, growth rate, sporulation, and formation of conidial chain, vesicle formation, and shape of conidia	[46]
PDA	Widely used media for visualization of macroscopic and microscopic structures of <i>A. flavus</i> , Culture-based detection methods based on color change, such as the Ammonia Vapor test and Fluorescence-based detection.	[42,47–49]
PDB and Modified PDB	Widely used for the higher growth of <i>Aspergillus</i> spp. to perform DNA extraction and measure AFs production.	[50–54]
MACM	Differences in AFB1 production by <i>A. flavus</i> through several sampling time periods and <i>A. flavus</i> growth over time and its dynamics.	[55]
PNA	To observe the interaction of climate change-related abiotic factors (temperature and water stress) on growth, gene expression, and AFB1 production	[56]

Note: AFPA: *A. flavus* and *parasiticus* agar; APA: Aflatoxins producing ability medium; CMA: Coconut milk agar; CEA: coconut extract agar; CCA: coconut cream agar; PDA: Potato Dextrose Agar; PDB: Potato Dextrose Broth; CYA: Czapek yeast extract agar; YESD: Yeast extract sucrose agar supplemented with methyl β -cyclodextrin and sodium desoxycholate; MACM: Maize Agar Culture Medium; PNA: Pistachio Nut Agar; YES: Yeast extract sucrose broth; MRBA: Modified Rose Bengal Agar Media; DRBC: Dichloran rose bengal chloramphenicol agar.

3.1. Ammonia Vapor Test

The ammonia vapor test is a conventional, rapid, and cost-effective method used to differentiate aflatoxigenic and non-aflatoxigenic *A. flavus* isolates. This technique is particularly useful for preliminary screening of non-aflatoxigenic strains for potential use as biocontrol agents.

In this method, *A. flavus* isolates are cultured on suitable agar media and exposed to ammonium hydroxide vapor by placing a drop of ammonia solution on the lid of an inverted Petri dish. After exposure, color changes in the fungal colonies are observed, which are associated with AFs production (Figure 1). The color variation is attributed to the interaction between ammonia vapor and AFs-related pigments, including averantin, norsolorinic acid, averufin, versicolorin A, versicolorin C, nidurufin, and versicolorin A hemiacetal [57]. These pigments act as pH-sensitive indicators, changing color in response to increased pH conditions.




Figure 1. Differentiation of aflatoxigenic and non-aflatoxigenic *A. flavus* isolates using the ammonia vapor test. Exposure to ammonium hydroxide vapor induces color changes in colonies due to pH-sensitive AFs-associated pigments, enabling visual discrimination of aflatoxigenic isolates.

Several studies have demonstrated the application of this method. For example, ref. [19] evaluated 25 *A. flavus* isolates and successfully differentiated aflatoxigenic and non-aflatoxigenic strains based on color variation. Similarly, ref. [20] reported that aflatoxigenic isolates produced a reddish-pink coloration on coconut agar medium (CAM) following ammonia treatment. The dichlorvos-ammonia (DV-AM) method has also been employed, where isolates grown on YES-DOC-CP-DV medium exhibited a yellow-to-red color change upon ammonia exposure [58].

Despite its advantages, the ammonia vapor test has several limitations. It is a qualitative method that does not provide quantitative information on AFs levels. Additionally, results may vary depending on culture age, media composition, pigment availability, and environmental conditions [56]. Therefore, this method is not standardized and should be used primarily as a preliminary screening tool rather than a definitive diagnostic approach.

3.2. Fluorescent-Based Detection

Fluorescence-based detection is another widely used culture-dependent method for identifying aflatoxigenic *A. flavus*. This technique relies on the emission of blue to blue-green fluorescence when fungal cultures are exposed to UV light (365 nm) on both solid and liquid media such as potato dextrose agar (PDA), coconut agar medium (CAM), and aflatoxins-producing ability (APA) medium [59].

In this method, fungal colonies are incubated and then observed under UV light. Aflatoxigenic isolates typically produce fluorescence, whereas non-aflatoxigenic isolates do not [57,60] (Figure 2). For example, ref. [16] demonstrated fluorescence-based detection using coconut cream agar (CCA), where enhanced visualization was achieved through the addition of chloramphenicol and incubation under controlled conditions.




Figure 2. Fluorescence-based detection of aflatoxigenic *A. flavus* isolates under UV light (365 nm). Aflatoxigenic strains exhibit blue to blue-green fluorescence due to AFs production, whereas non-aflatoxigenic isolates show little or no fluorescence.

The intensity of fluorescence can be enhanced using additives such as cyclodextrins and iodine, which improve AFs visualization [61]. Comparative studies have shown varying detection efficiencies; for instance, ref. [62] reported that 62% of isolates tested positive using UV fluorescence, whereas 54% were identified using the ammonia vapor test.

Although fluorescence-based detection is relatively simple and cost-effective, it also has limitations. Fluorescence intensity can vary depending on media composition, environmental conditions, and fungal growth stage. Furthermore, these methods do not provide quantitative data and may lead to misclassification of isolates.

3.3. Limitations of Cultural Methods and Need for Molecular Approaches

Overall, cultural methods are advantageous due to their low cost and simplicity; however, they lack sensitivity, specificity, and reproducibility. AFs production is influenced by multiple factors, including nutrient availability, pH, temperature, oxidative stress, and environmental conditions [63]. For example, low-nutrient media may reduce fungal growth and subsequently limit AFs production [63,64].

Therefore, while cultural techniques are useful for preliminary screening, they are insufficient for accurate identification of aflatoxigenic *A. flavus*. This limitation highlights the need for molecular approaches, which provide greater precision by targeting genes involved in AFs biosynthesis.

4. AFs Biosynthesis Gene Cluster

The AFs biosynthesis gene cluster is a complex genomic region approximately 70 kb in size, located near the telomeric region of chromosome 3 in *A. flavus* [63,65]. This cluster contains a large number of structural and regulatory genes that collectively encode enzymes involved in the AFs biosynthesis pathway (Table 2).

At least 27 enzymatic reactions are involved in AFs production, forming a highly coordinated polyketide pathway [66]. The expression of these genes is primarily regulated by two key transcriptional regulators, *aflR* and *aflS* [63,67]. Among these, *aflR* acts as a major transcriptional activator belonging to the *Gal4* family and plays a central role in regulating the expression of downstream biosynthetic genes [68]. Overexpression of *aflR* has been shown to enhance transcriptional activity and promote AFs production [69].

The *aflS* gene functions as a co-activator that enhances the regulatory activity of *aflR*. Interestingly, studies have shown that *aflS* expression is more closely associated with AFs production than *aflR*, as *aflS* expression decreases under conditions where AFs production is reduced, even when *aflR* expression remains stable [70]. This suggests that *aflS* may play a critical role in fine-tuning the regulation of AFs biosynthesis.

In addition to these regulatory genes, several transcription factors such as bZIP proteins (e.g., *atfA* and *atfB*) also influence AFs production by responding to oxidative stress and environmental signals. These factors bind to gene promoters and regulate transcriptional responses under stress conditions.

4.1. Stages of AFs Biosynthesis

The AFs biosynthesis pathway can be divided into four major stages based on the formation of intermediate compounds (Figure 3):




Figure 3. Schematic representation of the AFs biosynthesis pathway in *A. flavus*, illustrating key intermediate compounds and associated genes involved in AFB1 production.

4.1.1. Stage 1: Formation of Norsolorinic Acid (NOR)

The initial stage involves the conversion of acetate into norsolorinic acid (NOR), the first stable intermediate. This process is mediated by genes such as *aflA*, *aflB*, *aflC*, and *hypC* [71,72].

4.1.2. Stage 2: Conversion to Versicolorin A (VERA)

In this stage, NOR is converted into versicolorin A through a series of enzymatic reactions involving genes such as *aflD*, *aflG*, *aflH*, *aflK*, *aflI*, *aflV*, *aflW*, *aflJ*, and *aflL* [65,73–81].

4.1.3. Stage 3: Formation of Sterigmatocystin (ST)

Versicolorin A is further converted into sterigmatocystin through the action of genes such as *aflM*, *aflN*, *aflY*, *aflX*, and *aflO* [79,82–84].

4.1.4. Stage 4: Final Conversion to Aflatoxin B1 (AFB1)

The final stage involves the conversion of sterigmatocystin into aflatoxin B1 through genes such as *aflP*, *aflQ*, *hypB*, *aflE*, and *hypE* [24,85–87].

Among these stages, genes involved in the final steps of the pathway are considered particularly important, as they directly contribute to AFs production.

4.2. Regulation and Environmental Influence

AFs biosynthesis is not solely determined by gene presence but is strongly influenced by environmental and physiological factors. Variables such as temperature, CO₂ concentration, nutrient availability, oxidative stress, and developmental stage significantly affect gene expression and toxin production [88,89].

For example, nutrient-rich conditions promote AFs biosynthesis; whereas, nutrient limitation suppresses production. Similarly, oxidative stress plays a key role in regulating gene expression through transcription factors such as *atfA* and *atfB*.

4.3. Implications for Detection

Although the presence of AFs biosynthesis genes is often used as a marker for identifying aflatoxigenic isolates, it does not guarantee toxin production. The complexity of gene regulation, combined with environmental influences, makes it difficult to accurately predict AFs production based solely on gene presence.

Therefore, effective identification of aflatoxigenic *A. flavus* requires an integrated approach that considers gene presence, gene expression, and environmental factors. This complexity underscores the importance of combining molecular and analytical methods for reliable identification.

Table 2. AFs biosynthesis genes, their encoded enzymes, and associated functions. This table is adapted from [63,65,81,87,90].

Updated and Previous Name	Encoded Enzyme and Function	Stage of AFs Biosynthesis
<i>aflA (fas-2)</i>	Fatty acid synthase α ; conversion of hexanoate units into a polyketide structure	Conversion of acetate to norsolorinic acid (NOR)
<i>aflB (fas-1)</i>	Fatty acid synthase β ; conversion of hexanoate units into a polyketide structure	Conversion of acetate to NOR
<i>aflC (pksA)</i>	Polyketide synthase; synthesis of polyketide skeleton and conversion into norsolorinic acid anthrone (NAA)	Conversion of acetate to NOR
<i>hypC (hypB1)</i>	Anthrone oxidase; conversion of NAA into NOR	Conversion of acetate to NOR
<i>aflD (nor-1)</i>	Norsolorinic acid ketoreductase; conversion of NOR to averantin (AVN)	Conversion of NOR to versicolorin A
<i>aflG (avnA)</i>	Cytochrome P450 monooxygenase; hydroxylation of averantin into 5'-hydroxyaverantin (HAVN)	Conversion of NOR to versicolorin A
<i>aflH (adhA)</i>	Alcohol dehydrogenase; conversion of HAVN into 5'-oxoaverantin (OAVN)	Conversion of NOR to versicolorin A
<i>aflK (vbs)</i>	Cyclase/VERB synthase; transformation of versiconal into versicolorin B and conversion of intermediates to versicolorin B	Conversion of NOR to versicolorin A
<i>aflI (avfA)</i>	Averufin oxidase; conversion of averufin (AVF) into versiconal hemiacetal acetate (VHA)	Conversion of NOR to versicolorin A
<i>aflV (cypX)</i>	Cytochrome P450 monooxygenase; conversion of AVF into hydroxyversicolorone (HVN) and conversion of HVN into VHA	Conversion of NOR to versicolorin A
<i>aflJ (estA)</i>	Esterase; conversion of VHA into versiconal (VAL)	Conversion of NOR to versicolorin A
<i>aflL (verB)</i>	Cytochrome P450 monooxygenase/desaturase; conversion of versicolorin B into versicolorin A (VERA)	Conversion of NOR to versicolorin A
<i>aflM (ver-1)</i>	Ketoreductase; conversion of versicolorin A into demethylsterigmatocystin (DMST)	Conversion of versicolorin A to sterigmatocystin
<i>aflN (verA)</i>	Cytochrome P450 monooxygenase; conversion of versicolorin A into intermediate compounds	Conversion of versicolorin A to sterigmatocystin
<i>aflY (hypA)</i>	Hypothetical protein; involved in conversion of VERA to DMST	Conversion of versicolorin A to sterigmatocystin
<i>aflX (ordB)</i>	Oxidoreductase; catalyzes oxidative decarboxylation and ring closure reactions	Conversion of versicolorin A to sterigmatocystin
<i>aflO (omtB)</i>	O-methyltransferase; conversion of DMST into sterigmatocystin (ST)	Conversion of versicolorin A to sterigmatocystin
<i>aflP (omtA)</i>	O-methyltransferase; conversion of sterigmatocystin into O-methylsterigmatocystin (OMST)	Conversion of sterigmatocystin to AFB1
<i>aflQ (ordA)</i>	Cytochrome P450 monooxygenase; conversion of OMST to AFB1 and G1	Conversion of sterigmatocystin to AFB1
<i>aflE (norA)</i>	Aryl alcohol dehydrogenase; involved in late-stage AFs biosynthesis	Conversion of sterigmatocystin to AFB1
<i>hypE (aflLa)</i>	Hypothetical protein; involved in final steps of AFs production	Final stage of AFs biosynthesis

Note: NOR = norsolorinic acid; AVN = averantin; HAVN = hydroxyaverantin; OAVN = oxoaverantin; AVF = averufin; VHA = versiconal hemiacetal acetate; VAL = versiconal; VERA = versicolorin A; DMST = demethylsterigmatocystin; ST = sterigmatocystin; OMST = O-methylsterigmatocystin.

5. Existing Molecular Methods and Potential Drawbacks

Molecular detection approaches have become indispensable tools for identifying aflatoxigenic *A. flavus*, particularly due to the limitations associated with conventional morphological and cultural methods. As previously discussed, traditional diagnostic techniques, including visual symptom assessment, culturing, and biochemical assays, are often labor-intensive, time-consuming, and lack sufficient sensitivity and specificity [91].

To address these challenges, molecular techniques such as PCR, real-time PCR (qPCR), reverse transcription PCR (RT-PCR), multiplex PCR, loop-mediated isothermal amplification (LAMP) have been widely adopted [92,93] (Table 3). These approaches enable rapid, sensitive, and specific detection of *A. flavus*, even in complex environmental and food matrices. However, despite these advancements, accurate classification of aflatoxigenic isolates

remains difficult due to variability in gene presence, gene expression, and environmental influences on AFs biosynthesis.

Table 3. Comparison of different methods for accurate identification of aflatoxigenic *A. flavus*.

Methods/Techniques	Analysis Time	Cost	Sensitivity and Detection of Limit (LOD)	Specificity	Limitations	Field Applicability	Target Entity/Outcomes
Traditional/Cultural Techniques							
Ammonia Vapor Test	4–7 days	Low	Low		False negative	Low	Qualitative measurement of toxin production
Fluorescence-based detection	4–7 days	Moderate	Low		False negative	Low	Qualitative measurement of toxin production
Molecular Methods							
Standard PCR	3–5 h	Moderate to high	High	high	False positive/false negative	None	Only identify the presence of gene
RT-PCR	4–6 h	High	High	Very high	False negative	None	Determine gene expression of AFs genes
RT-qPCR	4–6 h	High	Very high	Very high	False positive/negative	None	Gene expression quantification of AFs genes
LAMP	30–60 min	Low to moderate	High	High	False positive/negative	High	Only identify the presence of gene
Analytical/Detection Methods							
Traditional (HPLC, LC-MS/MS, GC-MS, TLC, ELISA)	1–2 days	High to very high	Very high to excellent	Very high	False positive/negative	Low	Quantify Mycotoxins
Biosensor-based detection	15–45 min	Low	Excellent	Very high to excellent	False positive/negative	Very high	Quantify Mycotoxins

As shown in Table 4, multiple molecular methods are currently employed to detect aflatoxigenic *A. flavus* by targeting genes from the AFs biosynthesis gene cluster. While some molecular approaches improve identification efficiency, their ability to accurately classify isolates as aflatoxigenic or non-aflatoxigenic depends on integrating multiple layers of information, including gene presence, expression, and toxin quantification. Also, as shown in Table 3, no single method is sufficient for accurate identification of aflatoxigenic *A. flavus* because solely molecular detection could not identify aflatoxigenic isolates. Rather, integration of molecular and quantification methods provides better insights into aflatoxigenic *A. flavus* isolates.

Table 4. Use of several AFs biosynthesis genes and molecular methods to identify aflatoxigenic *A. flavus* in different crops, food and feed.

Molecular Method	Targeted Genes/Regions	Sample Source	Identification	References
Conventional PCR	Biosynthetic Pathway genes (<i>aflR</i> , <i>aflS</i> , <i>aflD</i> , <i>aflP</i> , <i>aflR</i> , <i>aflD</i> , <i>aflP</i> , <i>aflR</i> , <i>aflM</i> , <i>aflR</i> , <i>aflS</i> , <i>aflQ</i> , <i>aflP</i> , <i>aflD</i> , <i>aflM</i> , <i>aflO</i>)	Food, poultry feed, grains, stored grains, nuts, peanuts, spices	aflatoxigenic <i>A. flavus</i>	[20,94–101]
	Barcoding regions (ITS1, ITS4, Species-specific primer, <i>CaM</i>)	Chicken liver		[102–104]
PCR-RFLP	<i>aflP</i>	Peanut	aflatoxigenic <i>A. flavus</i>	[105]

Table 4. Cont.

Molecular Method	Targeted Genes/Regions	Sample Source	Identification	References
Multiplex PCR/Rt-PCR/qPCR	Biosynthetic Pathway Genes (<i>aflD</i> , <i>aflM</i> , <i>aflP</i> , <i>aflQ</i> , <i>aflD</i> , <i>aflO</i> , <i>aflQ</i> , <i>aflG</i> , <i>aflH</i> , <i>aflI</i> , <i>aflK</i> , <i>aflT</i> , <i>aflR</i> , <i>aflS</i> , <i>laeA</i> , <i>lipA</i>) and Housekeeping genes (<i>calmodulin</i> , β - <i>tub</i>)	Peanuts, soil and stored cultures, Brazil nuts, corn, almonds, hazelnuts, coffee, and pistachio nuts	aflatoxigenic <i>A. flavus</i>	[23,24,51,64,69,106–113]
LAMP Assay	Biosynthetic Pathway and Housekeeping genes (<i>aflP</i> , <i>aflD</i> , <i>aflT</i> and <i>acl1</i>)	Food samples, brazil nuts, spices, dried fruit, nuts and coffee	aflatoxigenic <i>A. flavus</i>	[30,114–117]

5.1. PCR

Polymerase chain reaction (PCR) is one of the most widely used molecular techniques for detecting *A. flavus*, based on the amplification of specific DNA fragments. The amplified products are typically visualized using gel electrophoresis with fluorescent dyes [118]. PCR offers several advantages, including rapid detection, high sensitivity, and the ability to analyze samples without extensive culturing.

Early applications of PCR in fungal identification focused on ribosomal DNA (rDNA) regions, particularly the internal transcribed spacer (ITS), which is highly variable and useful for species-level identification [119–121]. ITS-based assays have been successfully used to detect *A. flavus* and differentiate it from closely related species such as *A. parasiticus* [27,122,123]. In addition, protein-coding genes such as *calmodulin* (*CaM*) and β -*tubulin* (*BenA*) provide improved phylogenetic resolution within *Aspergillus* section Flavi [124,125].

For detecting aflatoxigenic strains, PCR assays often target genes within the AFs biosynthesis gene cluster, including regulatory genes (*aflR*, *aflT*) and structural genes (*aflD*, *aflM*, *aflP*) [108,126]. Several studies have demonstrated that combinations of these genes can be used to differentiate aflatoxigenic and non-aflatoxigenic isolates [20,127]. However, PCR-based methods have significant limitations. They primarily detect gene presence and do not provide information on gene expression or functional activity. For instance, the presence of *aflR* alone is insufficient to confirm AFs production, as it regulates multiple metabolic pathways [128].

Additionally, PCR assays are prone to false-positive and false-negative results due to primer mismatches, mutations within target genes, DNA quality, and low template concentration. Despite extensive research using various biosynthetic genes, no standardized gene set has been established for definitive identification of aflatoxigenic *A. flavus* [129,130]. Therefore, PCR alone is insufficient for accurate identification and must be complemented with other molecular or analytical techniques.

5.2. Multiplex PCR

Multiplex PCR enables simultaneous amplification of multiple target genes in a single reaction, providing a more comprehensive and efficient detection strategy compared to conventional PCR [113]. This method is particularly useful for targeting multiple genes from different stages of the AFs biosynthesis pathway, thereby improving detection reliability.

Several studies have utilized multiplex PCR to detect aflatoxigenic fungi using gene sets such as *nor-1*, *ver-1*, *omt-A*, and *apa-2* [108]. Similarly, multiplex PCR has been applied to detect multiple mycotoxigenic fungi across different genera, including *Aspergillus*, *Fusarium*, and *Penicillium* [113]. Other studies have targeted gene combinations such as

omtA, *omtB*, *ver-1*, and *aflR* for detecting aflatoxigenic *A. flavus* isolates from environmental samples [107].

Despite its advantages, multiplex PCR has limitations related to primer interactions, which can reduce amplification efficiency and sensitivity [106]. More importantly, similar to conventional PCR, multiplex PCR does not account for gene expression or toxin production. Therefore, although it improves detection coverage, it cannot definitively identify isolates as aflatoxigenic.

5.3. Reverse Transcriptase Polymerase Chain Reaction (RT-PCR)

RT-PCR is a powerful technique used to evaluate gene expression by converting RNA into complementary DNA (cDNA) prior to amplification. This approach provides insights into gene activity rather than mere gene presence, making it particularly valuable for studying AFs biosynthesis pathways.

Several studies have reported correlations between the expression of specific genes and AFs production. For example, *aflQ* and *aflO* genes have consistently shown strong positive correlations with AFs accumulation [23,111,113]. Similarly, *aflD* has been identified as an early-stage gene involved in AFs biosynthesis [112,131].

However, gene expression patterns are highly variable and influenced by environmental conditions, growth media, incubation time, and nutrient availability [89,124,132]. In some cases, isolates expressing multiple biosynthetic genes fail to produce detectable AFs levels, resulting in false-positive interpretations [112]. Furthermore, genetic mutations and regulatory differences can affect transcriptional activity, complicating interpretation [124].

These inconsistencies highlight that although RT-PCR provides valuable insights into gene expression, it cannot independently confirm AFs production. Therefore, it should be integrated with quantitative and analytical methods for accurate detection.

5.4. Quantitative PCR (qPCR)

Quantitative PCR (qPCR), also known as real-time PCR, allows simultaneous detection and quantification of DNA targets using fluorescent dyes or probes. This technique is widely used for detecting *A. flavus* in food, feed, and environmental samples due to its high sensitivity and specificity [133–136].

Several studies have demonstrated the effectiveness of qPCR in detecting AFs biosynthesis genes such as *omt-1*, *aflT*, and *aflR*, with detection limits as low as nanogram to femtogram levels [134,137,138].

However, qPCR shares similar limitations with conventional PCR, as it primarily detects gene presence rather than functional toxin production. Mutations within the AFs biosynthesis gene cluster and variability in gene expression can lead to inaccurate identification [135]. Therefore, integrating qPCR with RT-PCR and toxin quantification methods enhances detection reliability.

5.5. LAMP (Loop-Mediated Isothermal Amplification) Assay

Loop-mediated isothermal amplification (LAMP) is a rapid and sensitive DNA amplification technique that operates under constant temperature conditions [139]. This eliminates the need for thermal cycling equipment, making it suitable for field-based applications.

Several LAMP assays have been developed targeting genes such as *aflP*, *aflT*, *aflD*, and ITS regions for detecting aflatoxigenic *A. flavus* [30,106,116,117]. These assays demonstrate high sensitivity, with detection limits reaching picogram or femtogram levels [30].

Despite these advantages, LAMP assays may exhibit non-specific amplification and cross-reactivity with closely related species [115,116]. Additionally, false negatives can occur due to low DNA concentration or sample degradation [115]. Therefore, LAMP is most effective when used in combination with other molecular and analytical methods.

6. Recent Detection Techniques Mostly Focused on Aflatoxins Quantification but Limited to the Identification of Aflatoxigenic *A. flavus*

There are several methods already established for detecting AFs contamination in food and feed commodities, such as gas chromatography (GC), thin-layer chromatography (TLC), high-performance liquid chromatography (HPLC), fluorescence polarization assays, enzyme-linked immunosorbent assay (ELISA), and other microbiological methods used in a lab setting. These methods may provide accurate results; however, most of them need a well-organized laboratory, skilled personnel, and sophisticated methodologies [140]. Moreover, these techniques are usually expensive and need substantial time, particularly to detect AFs for large-scale screening, and are hard to incorporate into production or storage systems to monitor AFs detection or contamination. To minimize these limitations, some of the advanced techniques have been utilized to detect AFs contamination, which are rapid, sensitive, specific, and innovative.

6.1. Biosensor-Based Detections

A biosensor is a modern tool that is used to detect microorganisms and metabolites by merging biological recognition elements (DNA, enzymes, and antibodies) with a signal transducer. Those recognition elements usually bind to the specific target and develop a reaction, which is then converted through a transducer from a biological interaction to measurable signals (Electrical, Optical, and Physical). Biosensors have been widely used to detect mycotoxins in several foods and feeds. Depending on the bioreceptor and transducer, several types of biosensor systems have been utilized, focusing on different principles and functions such as aptamers, enzymes, nucleic acids, microorganisms, and protein-receptor based technique [141,142].

6.1.1. Enzymatic Sensor

This type of biosensor mainly operates on the association between substrate detection and enzyme inhibition [143,144]. Enzymatic biosensors can be widely incorporated in different transducers such as calorimetric, optical, and electrochemical. However, the most commonly used transducer is electrochemical [145]. This type of biosensor has wide application in different fields such as food, agricultural, environmental, pharmaceutical, and medical industries [146]. One of the main reasons behind this is the rapid and cost-effective method along with its analytical performance, which is listed in the study of [147].

For example, this technique was performed for the detection of tyramine, which was cross-linked with glutaraldehyde inside calcium orthophosphate, and then an amperometric technique was performed to measure the changes in the electrochemical signal of o-dopaquinone. Also, citrinin mycotoxin in rice samples was measured through the use of an amperometric biosensor utilizing horseradish peroxidase [147,148]. Furthermore, several types of enzymes have been used in terms of enzymatic inhibition, such as urease, cholinesterase, and glucose oxidase. For example, a commonly used enzyme, Acetylcholinesterase (AChE), is used because of its susceptibility to mycotoxins. Another study also supported this statement that AFB1 was detected by the use of AChE enzymatic activity due to the inhibitory effect between AFB1 and AChE [149]. Moreover, another study revealed that the use of cholinesterase in the biosensor technique is useful for the detection of AFB1 [150]. However, the utilization of nanomaterials in the development of biosensors improves the performance metrics such as stability and sensitivity. The sensitivity can be improved by the methods of inhibition, enzyme amount, and incubation time. Due to these advantages, it could be adapted for on-site monitoring and large-scale screening of different compounds.

6.1.2. Immunosensor

Immunosensors are termed as a bioanalytical device based on analyte-antibody interactions, where a specific antibody is used to detect a specific mycotoxin. In this technique, three primary groups have been developed, such as surface plasmon resonance, colorimetric sensors, and electrochemical sensors.

A label-free optical biosensor has been developed by Xu et al., (2013), which is based on gold nanorods and able to detect aflatoxin B1 [151]. The optical nano-biosensor provides highly selective, simple, and sensitive detection due to its strong ionic strength and reducing false results through the competitive dispersion principle [151]. AFB1 could be detected at a very low limit of detection (0.2 ng/mL) compared to the traditional ELISA method. For example, Surface plasmon resonance (SPR) is an optical detection technique that measures the total reflection of light upon a thin gold film, such as Au or Ag [152]. In one study, the sensitivity of the SPR was also higher compared to the ELISA method for AFB1 detection using the same immunoreagents. The sensitivity of the SPR was 3.0–49 ng/mL, while the sensitivity of the ELISA was 12–25,000 ng/mL [153]. Also, electrochemical immunosensors are based on the principle of the competitive ELISA, in which an electrochemical transducer allows direct oxidation-reduction reactions [154]. The detection of AFB1 was observed through the electrochemical immunosensor, where the limit of detection was 0.15 ng/mL in a solution. Apart from this, this method is also utilized to detect OTA with higher specificity [155].

6.1.3. Nucleic Acid Sensor or Aptasensors

Aptasensors are termed as biosensors, where aptamers (usually single-stranded RNA or DNA consisting of 2–60 nucleotides) particularly bind with the target molecule according to the lock-and-key theory [156,157]. Different immobilization strategies can influence aptasensor development, such as interactions between graphene oxide-modified interfaces and DNA base aptamers [158], binding thiolated aptamers to gold materials (Au-based materials) or CdTe quantum dots (QDs) [159], and linking aptamers to surface carboxylic groups [160].

So far, several aptasensors have been developed against different types of mycotoxins such as AFB1, OTA, fumonisin, ZEN, and T-2 toxin [158,161–166]. Moreover, mycotoxins, such as ZEN, OTA, and FB, in corn samples were detected based on multicolor quantum dot nanobeads with a detection limit of 5, 20, and 10 ng/mL, respectively [167]. An ultra-sensitive RT-qPCR-based aptasensor has been developed that can detect AFB1 in foods and feeds with high sensitivity (25 fg/mL), specificity, and low cost [168]. Also, an enzyme-free sensor has been developed by Chen et al. to detect aflatoxin B1 by utilizing a catalytic DNA circuit. The circuit is developed by two components such as gold nanoparticles and three hairpin DNA probes, which recognize specific AFB1 [169]. The advantages of the aptasensors are that they can detect mycotoxin contamination through a high-throughput technique, which could be adapted in the field for large-scale on-site detection. Also, these aptasensors are low-cost, sensitive, portable, and rapid compared to the traditional techniques, HPLC-MS and HPLC with fluorescence detection (FLD). Despite several advantages, aptasensor-based detection has some limitations, such as the proper selection of aptamers, a time-consuming process, and distinguishing and identifying similar mycotoxins (AFB1, AFB2, AFM1, and so on). Therefore, more research is required for the development of a specific aptamer and a multi-mycotoxin analysis scheme. Even though a significant advancement has been developed, signal amplification techniques and aptamer probe stability should be increased.

6.1.4. Solid-State Gas Sensor

It is another type of rapid and advanced technique that could be useful for the indirect detection of mycotoxins by utilizing volatile organic compounds (VOCs) emitted from the aflatoxigenic fungi [170,171]. Several aflatoxigenic species, such as *F. verticillioides*, *A. flavus*, and *P. verrucosum*, which release specific types of VOCs during their infection and colonization of different food and feed, could be utilized as reliable biomarkers for detecting the presence of mycotoxins [172].

The advantage of this method is that it provides a non-destructive, early method for detecting mycotoxins in different facilities, such as silos, storage facilities, and processing environments, offering an alternative to conventional analytical methods [49,173]. Also, mycotoxins can be detected even before significant toxin accumulation and visible fungal growth [171]. The main part of the principle of this technique is the interaction between the surface of the sensing layer and VOC molecules [174,175]. When aflatoxigenic fungi release VOCs, they interact with adsorbed O₂ species on the surface of the sensing layer and ultimately release electrons. Then, the electrical signal is measured in proportion to the VOC concentration [176,177]. To improve the performance of the gas sensor, nanoparticles could be integrated, which increases the adsorption sites and surface area [178]. In addition, metal doping and specific coatings also increase the specificity of VOCs [179,180]. This technique is also associated with some limitations, such as interference from environmental factors and non-specific VOCs, which could interfere with the accuracy, reproducibility, and sensitivity of the detection [181].

6.2. Optical-Based Detections

Among the several techniques mentioned above, optical-based methods are a rapid and non-destructive method for AFs detection, which is based on fluorescence spectroscopy (FS), hyperspectral imaging (HSI), Raman spectroscopy, and near-infrared spectroscopy (NIRS).

6.2.1. Fluorescence Spectroscopy (FS)

FS is termed an analytical technique that has been utilized in several disciplines to investigate the functions, structure, and reactivity of synthetic polymers, small molecules, other biological molecules, and proteins. Mainly, three processes are involved in the fluorescence process, such as excitation, conversion, and finally emission. The absorption of visible and ultraviolet light by a substrate or molecule is termed a fluorophore, which actually absorbs energy at a fixed wavelength and emits the light by liberating energy at a longer wavelength [182].

FS techniques have been widely used to detect mycotoxin contamination in several food and feed commodities [183]. There has been a link between AFs contamination, *A. flavus* infection, and bright-greenish-yellow fluorescence (BGYF), which was first reported in cottonseeds in 1969. After that, BGYF has been widely used as an indicator of AFs contamination [184]. A study performed by Wu et al. developed a self-built laser-induced fluorescence spectroscopy (LIFS) system integrated with three detection probes, which can classify AFB1-contaminated and non-contaminated pistachio kernels at a limit of detection of 5 µg kg⁻¹ and a classification accuracy of more than 91.0% [185]. Previously, studies reported the detection of AFs through the traditional FS methods [184,186]; however, this method has some limitations, such as the production of kojic acids, which influences the fluorescence signal for AFs detection [187] and can lead to false positive results [188]. Apart from that, the FS method relies on one specific wavelength, which is utilized on the sample. Compared to this technique, a new technique has been developed, named Fluorescence Fingerprinting (FF), which is a method for detecting a specific emission pattern across

several excitation wavelengths [189]. The FF method provides more sensitive and precise quantification compared to the FS [189], which is also a rapid and robust method of quantification of AFs [190].

6.2.2. Raman Spectroscopy

Raman Spectroscopy is a non-destructive technique based on the scattered light incident, which interacts with the chemical bonds within a sample. The Raman spectra can be obtained by detecting the Raman shifts and it has been widely used for the detection of food-borne pathogens in both the food and agri-industries due to its rapid, specific, and sensitive method [191–193].

Several studies have performed Raman spectroscopy techniques to differentiate bacterial species from different sources with an accuracy ranging from approximately 87% to 100% [194–196]. It has also been useful to detect bacteria in different survival stages, such as the viable but non-culturable state and biofilm formation state, which are crucial in terms of food safety and public health issues [197,198]. Besides bacteria, this technique is also utilized in non-bacterial pathogens such as viruses and fungi [199,200]. Apart from detecting microbes, it is also useful for detecting hazards and mycotoxin contamination [201–205]. To minimize the limitations of the traditional Raman spectroscopy method, a new analytical technique has been developed utilizing new nanomaterials such as copper, silver and gold called surface-enhanced Raman spectroscopy (SERS) [206]. Due to its characteristic spectroscopic fingerprint and superfast data-processing technique, SERS is useful for the direct and rapid detection of mycotoxins. Even though several advancements have been made in the SERS technique [207–209], there are still limitations regarding large-scale application, utilization of proper and specific binding reagents, point of need screening, and improvement of the SERS data analysis. Also, due to the low probability and the low intensity of the Raman scattering, it is difficult to detect molecules that are below the ppm level [210].

6.2.3. Near-Infrared Spectroscopy (NIRS)

Near-infrared spectroscopy (NIRS) relies on the interactions between the molecules in the samples and the absorbed light, which result in vibration and molecular overtone associated with several functional groups such as N-H, O-H, and C-H [211]. This technique has been widely used in the agricultural field, particularly as a promising approach for AFs detection and aflatoxigenic fungi due to low cost and high efficiency [212–214].

A diode array NIRS has been used in a study to separate moldy dried figs using spectroscopy and observed that the rate of separation between healthy and moldy figs was more than 88% by using partial least squares regression (PLSR) [214]. The classification accuracy can also be as high as 97% in terms of the detection of fungus infection [215]. To improve the detection technique of NIRS, a new technique has evolved named as Fourier transform near-infrared spectroscopy (FT-NIRS), which improves the accuracy, specificity, reproducibility, and wavelength resolution [216]. This improved method of detection can differentiate the contaminated and non-contaminated samples with an accuracy of more than 90% while utilizing a logarithmic linear classifier (LOGLC) and linear discriminant classifier (LDC) [217]. This accuracy rate was also further increased up to 100% with LDC to classify healthy and infected samples [218].

6.2.4. Hyperspectral Imaging (HSI)

A hyperspectral imaging technique is a relatively new technique that combines imaging and spectroscopic methods to provide spatial and spectral information of the samples being tested [219]. Therefore, compared to the traditional spectroscopic technique, it provides chemical imaging of the samples, which is useful for fungal infection and the

detection of AFs contamination. There are three approaches that are involved with this technique to acquire 3-D hypercubes, such as area scan, point scan, and line scan methods.

It is widely used in several fields such as agriculture, food safety, and quality [220–225]. One of the main advantages of this technique is its integration into a wide range of applications, such as total viable counts (TVC), Psychrotrophic plate counts (PPC), detection of bacteria, fungal infection, and microbial toxin detection [183]. Several studies have been reported to utilize this technique to determine the *Fusarium* and *Aspergillus*-infected cereal grains, along with measuring the level of AFs and fumonisin contamination [226–230]. Moreover, this method can also be adapted to monitor the early detection of mycotoxin contamination and fungal infection [231]. It has the potential to detect several microbial contaminants from different sources; however, it has some limitations. Some of the factors need to be evaluated further to enhance the efficacy and reproducibility of this method, such as minimizing redundant data during analysis, inaccuracy of the detection for homogeneous samples, and selecting the most effective wavelength. Moreover, integration of the exact reference value for the accuracy and effectiveness of the calibration model will improve better prediction.

7. Integration of Artificial Intelligence and Multi-Omics Approach in Aflatoxins Research

In the era of artificial intelligence, different advanced techniques have been integrated nowadays to monitor AFs risk prediction and AFs detection. For AFs detection, several ML-based models can be applied, such as Support Vector Machines, Random Forest, Gradient Boosting, and the classification of contamination risk using environmental datasets [232,233]. For example, Long Short-Term Memory (LSTM) and Convolutional Neural Networks (CNNs) can analyze time series and image-based datasets effectively up to 90% [234]. For the real-time prediction of AFs contamination, Hybrid AI-based models incorporate the factors related to fungal growth and colonization with sensor inputs, which will ultimately increase the prediction and interpretation [235,236]. Most of the machine learning models have a higher level of accuracy, ranging from 80 to 90% in terms of accurate classification of contaminated vs un-contaminated samples, forecasting contamination trends, and early contamination risk in storage conditions.

In addition, multi-omics-based approaches have also been developed recently in AFs research. In terms of metabolomics, several studies have been performed to identify the aflatoxigenic fungi, such as *A. flavus*, based on the chemical markers utilizing through gene expression [237–239]. Also, studies utilized metabolomics to determine the correlation between fungal infection and AFs accumulation [240,241]. Regarding the genomics approach, several genomics tools have been widely used for aflatoxigenic *A. flavus* [242–244] with a focus on whole-genome sequencing, AFs biosynthesis, and gene expression profiles [245–247]. Transcriptomics is one of the advanced fields in the recent era [248]. Transcriptomics research enables researchers to study gene expression at the transcriptional level along with gene expression regulation, functions of gene products, protein modification, and evolutionary changes [249,250]. For example, a study reported the use of RNA sequencing to identify gene functions related to *A. flavus* infection and AFs production [251,252]. Also, another study revealed how genes are expressed in different stages of infection, such as early or late stages [252]. Moreover, a transcriptomic study revealed how oxidative stress influences AFs accumulation [253]. Proteomics is another novel field that has been reviewed for AFs research [254]. Proteomic research in the mycotoxin field provides a clear understanding of cellular functions, protein profiling, and the effects of biotic factors. Also, it provides valuable information about plant responses against pathogen infestations and mycotoxin accumulation.

8. Conclusions and Future Perspectives

Over the years, both traditional (morphological and cultural) and molecular approaches have been developed for the detection of aflatoxigenic *A. flavus*. However, accurate identification remains challenging due to the complex nature of AFs biosynthesis, which is governed by multiple genes within the AFs biosynthesis gene cluster and influenced by gene expression dynamics and environmental conditions.

This review highlights that reliance on the presence of gene (single-gene or even multi-gene) identification strategies is often insufficient to definitively classify an isolate as aflatoxigenic. The presence or transcription of specific genes does not always correlate with actual AFs production, as toxin biosynthesis depends on a combination of genetic, physiological, and environmental factors. Therefore, not all *A. flavus* isolates possessing key biosynthetic genes can be considered aflatoxigenic.

As illustrated in Figure 4, accurate identification requires careful consideration of multiple interacting factors, including the selection of appropriate molecular targets across different stages of the AFs biosynthesis pathway, optimization of gene expression analysis, proper culture conditions, and the integration of reliable quantification methods. Additionally, methodological factors such as the use of multi-gene detection strategies, timing of sample preparation, and selection of suitable media further influence detection accuracy.




Figure 4. Key factors influencing accurate identification of aflatoxigenic *A. flavus*.

Looking forward, future research should focus on developing standardized, multi-target identification platforms that integrate molecular, biochemical, and environmental parameters along with advanced analytical or detection methods (Figures 4 and 5). The incorporation of advanced molecular techniques, along with emerging tools such as biosensors and portable sequencing technologies, offers promising opportunities for rapid and on-site detection. Furthermore, integrating omics-based approaches, including genomics and transcriptomics, can improve the understanding of gene regulation and enhance the predictive accuracy of AFs production and aflatoxigenic *A. flavus* identification. With advances in machine learning and deep learning algorithms, techniques such as AI-based predictive modeling, CRISPR-Cas, HIS, and NIR could provide a framework for identifying aflatoxigenic isolates and assessing their AFs production.




Figure 5. Future perspective on identification of aflatoxigenic *A. flavus* and AFs detection.

In conclusion, the future of detecting aflatoxigenic *A. flavus* lies in the development of integrated, multidisciplinary strategies that align with the complex factors outlined in Figure 4 and the utilization of techniques depicted in Figure 5. Such approaches will significantly improve detection reliability and support effective food safety monitoring and sustainable disease management practices.

Author Contributions: Conceptualization, E.A. and S.W.; investigation, E.A.; resources, E.A. and S.W.; writing—original draft preparation, M.M.M.; writing—review and editing, M.M.M., E.A., S.W. and A.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the United States Department of Agriculture (USDA); the National Institute of Food and Agriculture (NIFA) Capacity Building Grants Program (Award No. 2023-38821-40835); and Mississippi INBRE through the National Institutes of Health (Grant No. P20GM103476).

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviation is used in this manuscript:

AFs Aflatoxins

References

1. Medina, A.; Mohale, S.; Samsudin, N.I.P.; Rodriguez-Sixtos, A.; Rodriguez, A.; Magan, N. Biocontrol of Mycotoxins: Dynamics and Mechanisms of Action. *Curr. Opin. Food Sci.* **2017**, *17*, 41–48. [[CrossRef](#)]
2. Guo, B.; Chen, Z.Y.; Lee, R.D.; Scully, B.T. Drought Stress and Preharvest Aflatoxin Contamination in Agricultural Commodity: Genetics, Genomics and Proteomics. *J. Integr. Plant Biol.* **2008**, *50*, 1281–1291. [[CrossRef](#)]
3. Passone, M.A.; Ruffino, M.; Ponzio, V.; Resnik, S.; Etcheverry, M.G. Postharvest Control of Peanut *Aspergillus Section Flavi* Populations by a Formulation of Food-Grade Antioxidants. *Int. J. Food Microbiol.* **2009**, *131*, 211–217. [[CrossRef](#)] [[PubMed](#)]
4. Wang, P.; Qu, T.; Zhao, X.; Ge, Y.; Chen, Y. An Efficient Droplet Digital PCR Approach for Detection DNA at Low Concentrations of Toxigenic Fungi in Food Products. *Food Anal. Methods* **2022**, *1*, 1695–1706. [[CrossRef](#)]
5. IARC. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins. *IARC Monogr. Eval. Carcinog. Risks Hum.* **1993**, *56*, 259–362. [[CrossRef](#)]

6. Güç, İ.; Yalçın, E.; Çavuşoğlu, K.; Acar, A. Toxicity Mechanisms of Aflatoxin M1 Assisted with Molecular Docking and the Toxicity-Limiting Role of Trans-Resveratrol. *Sci. Rep.* **2022**, *12*, 14471. [[CrossRef](#)] [[PubMed](#)]
7. Mitchell, N.J.; Bowers, E.; Hurburgh, C.; Wu, F. Potential Economic Losses to the US Corn Industry from Aflatoxin Contamination. *Food Addit. Contam.—Part A Chem. Anal. Control Expo. Risk Assess.* **2016**, *33*, 540–550. [[CrossRef](#)] [[PubMed](#)]
8. Lengur, E.R.A.; Daten, H.; Nadut, A.; Lalong, P.R.F. Identification of Molds Producing Aflatoxin from Raw Materials of Free-Range Chicken Feed in East Nusa Tenggara, Indonesia. *Biodiversitas* **2024**, *25*, 1588–1595. [[CrossRef](#)]
9. Algammal, A.M.; Elsayed, M.E.; Hashem, H.R.; Ramadan, H.; Sheraba, N.S.; El-Diasty, E.M.; Abbas, S.M.; Hetta, H.F. Molecular and HPLC-Based Approaches for Detection of Aflatoxin B1 and Ochratoxin A Released from Toxicogenic *Aspergillus* Species in Processed Meat. *BMC Microbiol.* **2021**, *21*, 82. [[CrossRef](#)] [[PubMed](#)]
10. Gong, Y.Y.; Watson, S.; Routledge, M.N. Aflatoxin Exposure and Associated Human Health Effects, a Review of Epidemiological Studies. *Food Saf.* **2016**, *4*, 14–27. [[CrossRef](#)] [[PubMed](#)]
11. Kumar, P.; Mahato, D.K.; Kamle, M.; Mohanta, T.K.; Kang, S.G. Aflatoxins: A Global Concern for Food Safety, Human Health and Their Management. *Front. Microbiol.* **2017**, *7*, 2170. [[CrossRef](#)] [[PubMed](#)]
12. Moral, J.; Garcia-Lopez, M.T.; Camiletti, B.X.; Jaime, R.; Michailides, T.J.; Bandyopadhyay, R.; Ortega-Beltran, A. Present Status and Perspective on the Future Use of Aflatoxin Biocontrol Products. *Agronomy* **2020**, *10*, 491. [[CrossRef](#)]
13. Mehrzad, J.; Devriendt, B.; Baert, K.; Cox, E. Aflatoxin B1 Interferes with the Antigen-Presenting Capacity of Porcine Dendritic Cells. *Toxicol. Vitro* **2014**, *28*, 531–537. [[CrossRef](#)] [[PubMed](#)]
14. Bruix, J.; Merle, P.; Granito, A.; Huang, Y.-H.; Bodoky, G.; Yokosuka, O.; Rosmorduc, O.; Breder, V.; Gerolami, R.; Masi, G.; et al. LBA-03 Efficacy and Safety of Regorafenib versus Placebo in Patients with Hepatocellular Carcinoma (HCC) Progressing on Sorafenib: Results of the International, Randomized Phase 3 RESORCE Trial. *Ann. Oncol.* **2016**, *27*, ii140. [[CrossRef](#)]
15. Yoshida, H.; Mamada, Y.; Taniai, N.; Uchida, E. Spontaneous Ruptured Hepatocellular Carcinoma. *Hepatol. Res.* **2016**, *46*, 13–21. [[CrossRef](#)] [[PubMed](#)]
16. Khan, R.; Ghazali, F.M.; Mahyudin, N.A.; Samsudin, N.I.P. Morphological Characterization and Determination of Aflatoxigenic and Non-Aflatoxigenic *Aspergillus Flavus* Isolated from Sweet Corn Kernels and Soil in Malaysia. *Agriculture* **2020**, *10*, 450. [[CrossRef](#)]
17. McGlynn, K.A.; London, W.T. Epidemiology and Natural History of Hepatocellular Carcinoma. *Best Pract. Res. Clin. Gastroenterol.* **2005**, *19*, 3–23. [[CrossRef](#)] [[PubMed](#)]
18. Benkerroum, N. Mycotoxins in Dairy Products: A Review. *Int. Dairy J.* **2016**, *62*, 63–75. [[CrossRef](#)]
19. Kumar, S.; Shekhar, M.; Khan, A.A.; Sharma, P. A Rapid Technique for Detection of Toxicogenic and Nontoxicogenic Strains of *Aspergillus Flavus* from Maize Grains. *Indian Phytopathol.* **2007**, *60*, 31–34.
20. Alkhursan, R.N.; Abbas, B.A.; Khudor, M.H. PCR-Based Detection of Aflatoxigenic Strains of *Aspergillus Flavus* Isolated from Poultry Feed. *IOP Conf. Ser. Earth Environ. Sci.* **2021**, 761. [[CrossRef](#)]
21. Sardiñas, N.; Vázquez, C.; Gil-Serna, J.; González-Jaén, M.T.; Patiño, B. Specific Detection of *Aspergillus Parasiticus* in Wheat Flour Using a Highly Sensitive PCR Assay. *Food Addit. Contam.—Part A* **2010**, *27*, 853–858. [[CrossRef](#)] [[PubMed](#)]
22. Abbas, H.K.; Zablutowicz, R.M.; Weaver, M.A.; Horn, B.W.; Xie, W.; Shier, W.T. Comparison of Cultural and Analytical Methods for Determination of Aflatoxin Production by Mississippi Delta *Aspergillus* Isolates. *Can. J. Microbiol.* **2004**, *50*, 193–199. [[CrossRef](#)] [[PubMed](#)]
23. Mahmoud, M.A. Detection of *Aspergillus Flavus* in Stored Peanuts Using Real-Time PCR and the Expression of Aflatoxin Genes in Toxicogenic and Atoxigenic *A. Flavus* Isolates. *Foodborne Pathog. Dis.* **2015**, *12*, 289–296. [[CrossRef](#)] [[PubMed](#)]
24. Scherm, B.; Palomba, M.; Serra, D.; Marcello, A.; Migheli, Q. Detection of Transcripts of the Aflatoxin Genes AflD, AflO, and AflP by Reverse Transcription-Polymerase Chain Reaction Allows Differentiation of Aflatoxin-Producing and Non-Producing Isolates of *Aspergillus Flavus* and *Aspergillus Parasiticus*. *Int. J. Food Microbiol.* **2005**, *98*, 201–210. [[CrossRef](#)] [[PubMed](#)]
25. Vegi, A.; Wolf-Hall, C.E. Multiplex Real-Time PCR Method for Detection and Quantification of Mycotoxigenic Fungi Belonging to Three Different Genera. *J. Food Sci.* **2013**, *78*, M70–M76. [[CrossRef](#)] [[PubMed](#)]
26. Criseo, G.; Bagnara, A.; Bisignano, G. Differentiation of aflatoxin-Producing and Non-Producing Strains of *Aspergillus flavus* Group. *Lett. Appl. Microbiol.* **2001**, *33*, 291–295. [[CrossRef](#)] [[PubMed](#)]
27. González-Salgado, A.; González-Jaén, T.; Vázquez, C.; Patiño, B. Highly Sensitive Pcr-Based Detection Method Specific for *Aspergillus Flavus* in Wheat Flour. *Food Addit. Contam.—Part A Chem. Anal. Control Expo. Risk Assess.* **2008**, *25*, 758–764. [[CrossRef](#)] [[PubMed](#)]
28. Mayer, Z.; Färber, P.; Geisen, R. Monitoring the Production of Aflatoxin B1 in Wheat by Measuring the Concentration of Nor-1 mRNA. *Appl. Environ. Microbiol.* **2003**, *69*, 1154–1158. [[CrossRef](#)] [[PubMed](#)]
29. Sweeney, M.J.; Pàmies, P.; Dobson, A.D.W. The Use of Reverse Transcription-Polymerase Chain Reaction (RT-PCR) for Monitoring Aflatoxin Production in *Aspergillus Parasiticus* 439. *Int. J. Food Microbiol.* **2000**, *56*, 97–103. [[CrossRef](#)] [[PubMed](#)]
30. Liu, P.; Li, B.; Yin, R.; Weng, Q.; Chen, Q. Development and Evaluation of ITS- and AflP-Based LAMP Assays for Rapid Detection of *Aspergillus Flavus* in Food Samples. *Can. J. Microbiol.* **2014**, *60*, 579–584. [[CrossRef](#)] [[PubMed](#)]

31. Weaver, M.A.; Callicott, K.A.; Mehl, H.L.; Opoku, J.; Park, L.C.; Fields, K.S.; Mandel, J.R. Characterization of the *Aspergillus Flavus* Population from Highly Aflatoxin-Contaminated Corn in the United States. *Toxins* **2022**, *14*, 755. [[CrossRef](#)] [[PubMed](#)]
32. Pallavi, R.M.V.; Ramana, D.; Sashidhar, R.B. Short Note: A Simple Test Tube Screening Method for Identifying Aflatoxin-Producing Strains of *Aspergillus* Sp. Using Coconut Milk Agar Medium. *World J. Microbiol. Biotechnol.* **1997**, *13*, 713–714. [[CrossRef](#)]
33. Wei, D.; Zhou, L.; Selvaraj, J.N.; Zhang, C.; Xing, F.; Zhao, Y.; Wang, Y.; Liu, Y. Molecular Characterization of Atoxigenic *Aspergillus Flavus* Isolates Collected in China. *J. Microbiol.* **2014**, *52*, 559–565. [[CrossRef](#)] [[PubMed](#)]
34. Cotty, P.J. Fungi. *Mycopathologia* **1994**, *125*, 157–162. [[PubMed](#)]
35. Shabeer, S.; Asad, S.; Jamal, A.; Ali, A. Aflatoxin Contamination, Its Impact and Management Strategies: An Updated Review. *Toxins* **2022**, *14*, 307. [[CrossRef](#)] [[PubMed](#)]
36. Klich, M.A.; Pitt, J.I. Differentiation of *Aspergillus Flavus* from *A. Parasiticus* and Other Closely Related Species. *Trans. Br. Mycol. Soc.* **1988**, *91*, 99–108. [[CrossRef](#)]
37. Zhang, W.; Chang, X.; Wu, Z.; Dou, J.; Yin, Y.; Sun, C.; Wu, W. Rapid Isolation of Non-Aflatoxigenic *Aspergillus Flavus* Strains. *World Mycotoxin J.* **2020**, *13*, 277–286. [[CrossRef](#)]
38. Pitt, J.I.; Hocking, A.D.; Glenn, D.R. An Improved Medium for the Detection of *Aspergillus Flavus* and *A. Parasiticus*. *J. Appl. Bacteriol.* **1983**, *54*, 109–114. [[CrossRef](#)] [[PubMed](#)]
39. Fente, C.A.; Ordaz, J.J.; Vázquez, B.I.; Franco, C.M.; Cepeda, A. New Additive for Culture Media for Rapid Identification of Aflatoxin-Producing *Aspergillus* Strains. *Appl. Environ. Microbiol.* **2001**, *67*, 4858–4862. [[CrossRef](#)] [[PubMed](#)]
40. Dyer, S.K.; Mccammon, S. Related Species on Coconut Cream Agar. *J. Appl. Bacteriol.* **1994**, *76*, 75–78. [[CrossRef](#)] [[PubMed](#)]
41. Mamo, F.T.; Selvaraj, J.N.; Wang, Y.; Liu, Y. Recent Developments in the Screening of Atoxigenic *Aspergillus Flavus* towards Aflatoxin Biocontrol. *J. Appl. Environ. Microbiol.* **2017**, *5*, 20–30. [[CrossRef](#)]
42. Jaimez Ordaz, J.; Fente, C.A.; Vázquez, B.I.; Franco, C.M.; Cepeda, A. Development of a Method for Direct Visual Determination of Aflatoxin Production by Colonies of the *Aspergillus Flavus* Group. *Int. J. Food Microbiol.* **2003**, *83*, 219–225. [[CrossRef](#)] [[PubMed](#)]
43. Liu, J.; Sun, L.; Zhang, N.; Zhang, J.; Guo, J.; Li, C.; Rajput, S.A.; Qi, D. Effects of Nutrients in Substrates of Different Grains on Aflatoxin B1 Production by *Aspergillus Flavus*. *Biomed Res. Int.* **2016**, *2016*, 7232858. [[CrossRef](#)] [[PubMed](#)]
44. Njoroge, S.M.C.; Kanenga, K.; Siambi, M.; Waliyar, F.; Monyo, E.S. Identification and Toxigenicity of *Aspergillus* Spp. from Soils Planted to Peanuts in Eastern Zambia. *Peanut Sci.* **2016**, *43*, 148–156. [[CrossRef](#)]
45. Katati, B.; Kovács, S.; Njapau, H.; Kachapulula, P.W.; Zwaan, B.J.; van Diepeningen, A.D.; Schoustra, S.E. Maize *Aspergillus* Section Flavi Isolate Diversity May Be Distinct from That of Soil and Subsequently the Source of Aflatoxin Contamination. *Mycotoxin Res.* **2024**, *40*, 351–367. [[CrossRef](#)] [[PubMed](#)]
46. Suleiman, W.B. A Multi-Aspect Analysis of Two Analogous *Aspergillus* Spp. Belonging to Section Flavi: *Aspergillus Flavus* and *Aspergillus Oryzae*. *BMC Microbiol.* **2023**, *23*, 71. [[CrossRef](#)] [[PubMed](#)]
47. Zhou, L.; Wei, D.D.; Selvaraj, J.N.; Shang, B.; Zhang, C.S.; Xing, F.G.; Zhao, Y.J.; Wang, Y.; Liu, Y. A Strain of *Aspergillus Flavus* from China Shows Potential as a Biocontrol Agent for Aflatoxin Contamination. *Biocontrol Sci. Technol.* **2015**, *25*, 583–592. [[CrossRef](#)]
48. Liang, M.; Cai, X.; Gao, Y.; Yan, H.; Fu, J.; Tang, X.; Zhang, Q.; Li, P. A Versatile Nanozyme Integrated Colorimetric and Photothermal Lateral Flow Immunoassay for Highly Sensitive and Reliable *Aspergillus Flavus* Detection. *Biosens. Bioelectron.* **2022**, *213*, 114435. [[CrossRef](#)] [[PubMed](#)]
49. Okechukwu, V.O.; Kappo, A.P.; Njobeh, P.B.; Mamo, M.A. Morphed Aflaxotin Concentration Produced by *Aspergillus Flavus* Strain VKMN22 on Maize Grains Inoculated on Agar Culture. *Food Chem. Mol. Sci.* **2024**, *8*, 100197. [[CrossRef](#)] [[PubMed](#)]
50. Meyers, D.M.; Obrian, G.; Du, W.L.; Bhatnagar, D.; Payne, G.A. Characterization of AflJ, a Gene Required for Conversion of Pathway Intermediates to Aflatoxin. *Appl. Environ. Microbiol.* **1998**, *64*, 3713–3717. [[CrossRef](#)] [[PubMed](#)]
51. Okayo, R.O.; Andika, D.O.; Dida, M.M.; K’otuto, G.O.; Gichimu, B.M. Morphological and Molecular Characterization of Toxigenic *Aspergillus Flavus* from Groundnut Kernels in Kenya. *Int. J. Microbiol.* **2020**, *2020*. [[CrossRef](#)] [[PubMed](#)]
52. Palumbo, J.D.; Sarreal, S.B.L.; Kim, J.H. Simultaneous Detection of Mycotoxigenic *Aspergillus* Species of Sections Circumdati and Flavi Using Multiplex Digital PCR. *Lett. Appl. Microbiol.* **2023**, *76*, ovad142. [[CrossRef](#)] [[PubMed](#)]
53. Geng, Q.; Hu, J.; Xu, P.; Sun, T.; Qiu, H.; Wang, S.; Song, F.; Shen, L.; Li, Y.; Liu, M.; et al. The Autophagy-Related Protein ATG8 Orchestrates Asexual Development and AFB1 Biosynthesis in *Aspergillus Flavus*. *J. Fungi* **2024**, *10*, 349. [[CrossRef](#)] [[PubMed](#)]
54. Rao, K.R.; Vipin, A.V.; Venkateswaran, G. Mechanism of Inhibition of Aflatoxin Synthesis by Non-Aflatoxigenic Strains of *Aspergillus Flavus*. *Microb. Pathog.* **2020**, 104280. [[CrossRef](#)] [[PubMed](#)]
55. Guo, X.; Jia, B.; Zhang, H.; Ni, X.; Zhuang, H.; Lu, Y.; Wang, W. Evaluation of *Aspergillus Flavus* Growth and Detection of Aflatoxin B1 Content on Maize Agar Culture Medium Using Vis/NIR Hyperspectral Imaging. *Agriculture* **2023**, *13*, 237. [[CrossRef](#)]
56. Baazeem, A.; Rodriguez, A.; Medina, A.; Magan, N. Impacts of Climate Change Interacting Abiotic Factors on Growth, AflD and AflR Gene Expression and Aflatoxin B1 Production by *Aspergillus Flavus* Strains in Vitro and on Pistachio Nuts. *Toxins* **2021**, *13*, 385. [[CrossRef](#)] [[PubMed](#)]

57. Abbas, H.K.; Shier, W.T.; Horn, B.W.; Weaver, M.A. Cultural Methods for Aflatoxin Detection. *J. Toxicol.—Toxin Rev.* **2004**, *23*, 295–315. [\[CrossRef\]](#)
58. Guo, Z.; Zhang, J.; Wang, H.; Dong, H.; Li, S.; Shao, X.; Huang, J.; Yin, X.; Zhang, Q.; Guo, Y.; et al. Enhanced Detection of *Aspergillus Flavus* in Peanut Kernels Using a Multi-Scale Attention Transformer (MSAT): Advancements in Food Safety and Contamination Analysis. *Int. J. Food Microbiol.* **2024**, *423*, 110831. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Alkhersan, R.N.; Khudor, M.H.; Abbas, B.A. Rapid Detection of Aflatoxigenic Producing Strains of *Aspergillus Flavus* from Poultry Feeds by UV Light and Ammonia. *Basrah J. Vet. Res.* **2016**, *14*, 169–178.
60. Rodrigues, P.; Soares, C.M.G.; Kozakiewicz, Z.; Paterson, R.R.M.; Lima, N.; Venâncio, A. Identification and Characterization of *Aspergillus Flavus* and Aflatoxins. In *Microbiology Book Series—Communicating, Current Research and Educational Topics and Trends in Applied, Microbiology*; Méndez-Vilas, A., Ed.; Formatex: Badajoz, Spain, 2007; Volume 2, pp. 527–534.
61. Ouadhene, M.A.; Callicott, K.A.; Ortega-Beltran, A.; Mehl, H.L.; Cotty, P.J.; Battilani, P. Structure of *Aspergillus Flavus* Populations Associated with Maize in Greece, Spain, and Serbia: Implications for Aflatoxin Biocontrol on a Regional Scale. *Environ. Microbiol. Rep.* **2024**, *16*, e13249. [\[CrossRef\]](#) [\[PubMed\]](#)
62. Agbetiamah, D.; Ortega-beltran, A.; Awuah, R.T.; Atehnkeng, J. Field Efficacy of Two Atoxigenic Biocontrol Products for Mitigation of Aflatoxin Contamination in Maize and Groundnut in Ghana. *Biol. Control* **2020**, *150*, 104351. [\[CrossRef\]](#) [\[PubMed\]](#)
63. Accinelli, C.; Abbas, H.K.; Little, N.S.; Kotowicz, J.K.; Shier, W.T. Biological Control of Aflatoxin Production in Corn Using Non-Aflatoxigenic *Aspergillus Flavus* Administered as a Bioplastic-Based Seed Coating. *Crop Prot.* **2018**, *107*, 87–92. [\[CrossRef\]](#)
64. Passone, M.A.; Rosso, L.C.; Ciancio, A.; Etcheverry, M. Detection and Quantification of *Aspergillus Section Flavi* Spp. in Stored Peanuts by Real-Time PCR of nor-1 Gene, and Effects of Storage Conditions on Aflatoxin Production. *Int. J. Food Microbiol.* **2010**, *138*, 276–281. [\[CrossRef\]](#) [\[PubMed\]](#)
65. Yu, J.; Chang, P.K.; Ehrlich, K.C.; Cary, J.W.; Bhatnagar, D.; Cleveland, T.E.; Payne, G.A.; Linz, J.E.; Woloshuk, C.P.; Bennett, J.W. Clustered Pathway Genes in Aflatoxin Biosynthesis. *Appl. Environ. Microbiol.* **2004**, *70*, 1253–1262. [\[CrossRef\]](#) [\[PubMed\]](#)
66. Bennett, J.W.; Moore, G.G. Mycotoxins. *Encycl. Microbiol.* **2019**, 267–273. [\[CrossRef\]](#)
67. Price, M.S.; Yu, J.; Nierman, W.C.; Kim, H.; Pritchard, B.; Jacobus, C.A.; Bhatnagar, D.; Cleveland, T.E.; Payne, G.A. The Aflatoxin Pathway Regulator AflR Induces Gene Transcription inside and Outside of the Aflatoxin Biosynthetic Cluster. *FEMS Microbiol. Lett.* **2006**, *255*, 275–279. [\[CrossRef\]](#) [\[PubMed\]](#)
68. Shimizu, K.; Hicks, J.K.; Huang, T.P.; Keller, N.P. Pka, Ras and RGS Protein Interactions Regulate Activity of AflR, a Zn(II)2Cys6 Transcription Factor in *Aspergillus Nidulans*. *Genetics* **2003**, *165*, 1095–1104. [\[CrossRef\]](#) [\[PubMed\]](#)
69. Baquião, A.C.; Rodrigues, A.G.; Lopes, E.L.; Tralamazza, S.M.; Zorzete, P.; Correa, B. Expression of Genes by Aflatoxigenic and Non-aflatoxigenic Strains of *Aspergillus Flavus* Isolated from Brazil Nuts. *Foodborne Pathog. Dis.* **2016**, *13*, 434–440. [\[CrossRef\]](#) [\[PubMed\]](#)
70. Caceres, I.; El Khoury, R.; Bailly, S.; Oswald, I.P.; Puel, O.; Bailly, J.D. Piperine Inhibits Aflatoxin B1 Production in *Aspergillus Flavus* by Modulating Fungal Oxidative Stress Response. *Fungal Genet. Biol.* **2017**, *107*, 77–85. [\[CrossRef\]](#) [\[PubMed\]](#)
71. Roze, L.V.; Laivenieks, M.; Hong, S.Y.; Wee, J.; Wong, S.S.; Vanos, B.; Awad, D.; Ehrlich, K.C.; Linz, J.E. Aflatoxin Biosynthesis Is a Novel Source of Reactive Oxygen Species—A Potential Redox Signal to Initiate Resistance to Oxidative Stress? *Toxins* **2015**, *7*, 1411–1430. [\[CrossRef\]](#) [\[PubMed\]](#)
72. Ehrlich, K.C.; Li, P.; Scharfenstein, L.; Chang, P.K. HypC, the Anthrone Oxidase Involved in Aflatoxin Biosynthesis. *Appl. Environ. Microbiol.* **2010**, *76*, 3374–3377. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Zhou, R.; Linz, J.E. Enzymatic Function of the Nor-1 Protein in Aflatoxin Biosynthesis in *Aspergillus Parasiticus*. *Appl. Environ. Microbiol.* **1999**, *65*, 5639–5641. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Yabe, K.; Nakamura, Y.; Nakajima, H.; Ando, Y.; Hamasaki, T. Enzymatic Conversion of Norsolorinic Acid to Averufin in Aflatoxin Biosynthesis. *Appl. Environ. Microbiol.* **1991**, *57*, 1340–1345. [\[CrossRef\]](#) [\[PubMed\]](#)
75. Yu, J.; Chang, P.K.; Cary, J.W.; Bhatnagar, D.; Cleveland, T.E. AvnA, a Gene Encoding a Cytochrome P-450 Monooxygenase, Is Involved in the Conversion of Averantin to Averufin in Aflatoxin Biosynthesis in *Aspergillus Parasiticus*. *Appl. Environ. Microbiol.* **1997**, *63*, 1349–1356. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Cary, J.W.; Montalbano, B.G.; Ehrlich, K.C. Promoter Elements Involved in the Expression of the *Aspergillus Parasiticus* Aflatoxin Biosynthesis Pathway Gene AvnA. *Biochim. Biophys. Acta—Gene Struct. Expr.* **2000**, *1491*, 7–12. [\[CrossRef\]](#) [\[PubMed\]](#)
77. Chang, P.K.; Yu, J.; Ehrlich, K.C.; Boue, S.M.; Montalbano, B.G.; Bhatnagar, D.; Cleveland, T.E. AdhA in *Aspergillus Parasiticus* Is Involved in Conversion of 5'-Hydroxyaverantin to Averufin. *Appl. Environ. Microbiol.* **2000**, *66*, 4715–4719. [\[CrossRef\]](#) [\[PubMed\]](#)
78. Sakuno, E.; Wen, Y.; Hatabayashi, H.; Arai, H.; Aoki, C.; Yabe, K.; Nakajima, H. *Aspergillus Parasiticus* Cyclase Catalyzes Two Dehydration Steps in Aflatoxin Biosynthesis. *Appl. Environ. Microbiol.* **2005**, *71*, 2999–3006. [\[CrossRef\]](#) [\[PubMed\]](#)
79. Yu, J.; Woloshuk, C.P.; Bhatnagar, D.; Cleveland, T.E. Cloning and Characterization of AvfA and OmtB Genes Involved in Aflatoxin Biosynthesis in Three *Aspergillus* Species. *Gene* **2000**, *248*, 157–167. [\[CrossRef\]](#) [\[PubMed\]](#)
80. Wen, Y.; Hatabayashi, H.; Arai, H.; Kitamoto, H.K.; Yabe, K. Function of the CypX and MoxY Genes in Aflatoxin Biosynthesis in *Aspergillus Parasiticus*. *Appl. Environ. Microbiol.* **2005**, *71*, 3192–3198. [\[CrossRef\]](#) [\[PubMed\]](#)

81. Yu, J.; Bhatnagar, D.; Ehrlich, K.C. Aflatoxin Biosynthesis. *Rev. Iberoam. Micol.* **2002**, *19*, 191–200. [\[PubMed\]](#)
82. Henry, K.M.; Townsend, C.A. Ordering the Reductive and Cytochrome P450 Oxidative Steps in Demethylsterigmatocystin Formation Yields General Insights into the Biosynthesis of Aflatoxin and Related Fungal Metabolites. *J. Am. Chem. Soc.* **2005**, *127*, 3724–3733. [\[CrossRef\]](#) [\[PubMed\]](#)
83. Cary, J.W.; Ehrlich, K.C.; Bland, J.M.; Montalbano, B.G. The Aflatoxin Biosynthesis Cluster Gene, AflX, Encodes an Oxidoreductase Involved in Conversion of Versicolorin A to Demethylsterigmatocystin. *Appl. Environ. Microbiol.* **2006**, *72*, 1096–1101. [\[CrossRef\]](#) [\[PubMed\]](#)
84. Ehrlich, K.C.; Montalbano, B.; Boué, S.M.; Bhatnagar, D. An Aflatoxin Biosynthesis Cluster Gene Encodes a Novel Oxidase Required for Conversion of Versicolorin A to Sterigmatocystin. *Appl. Environ. Microbiol.* **2005**, *71*, 8963–8965. [\[CrossRef\]](#) [\[PubMed\]](#)
85. Ehrlich, K.C.; Chang, P.K.; Scharfenstein, L.L.; Cary, J.W.; Crawford, J.M.; Townsend, C.A. Absence of the Aflatoxin Biosynthesis Gene, NorA, Allows Accumulation of Deoxyaflatoxin B1 in *Aspergillus Flavus* Cultures. *FEMS Microbiol. Lett.* **2010**, *305*, 65–70. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Yu, J.; Chang, P.K.; Ehrlich, K.C.; Cary, J.W.; Montalbano, B.; Dyer, J.M.; Bhatnagar, D.; Cleveland, T.E. Characterization of the Critical Amino Acids of an *Aspergillus Parasiticus* Cytochrome P-450 Monooxygenase Encoded by OrdA That Is Involved in the Biosynthesis of Aflatoxins B1, G1, B2, and G2. *Appl. Environ. Microbiol.* **1998**, *64*, 4834–4841. [\[CrossRef\]](#) [\[PubMed\]](#)
87. Yu, J. Current Understanding on Aflatoxin Biosynthesis and Future Perspective in Reducing Aflatoxin Contamination. *Toxins* **2012**, *4*, 1024–1057. [\[CrossRef\]](#) [\[PubMed\]](#)
88. Madbouly, A.K.; Rashad, Y.M.; Ibrahim, M.I.M.; Elazab, N.T. Biodegradation of Aflatoxin B1 in Maize Grains and Suppression of Its Biosynthesis-Related Genes Using Endophytic *Trichoderma Harzianum* AYM3. *J. Fungi* **2023**, *9*, 209. [\[CrossRef\]](#) [\[PubMed\]](#)
89. Medina; Rodríguez, A.; Sultan, Y.; Magan, N. Climate Change Factors and *Aspergillus Flavus*: Effects on Gene Expression, Growth and Aflatoxin Production. *World Mycotoxin J.* **2015**, *8*, 171–179. [\[CrossRef\]](#)
90. Yu, J.; Chang, P.K.; Cary, J.W.; Wright, M.; Bhatnagar, D.; Cleveland, T.E.; Payne, G.A.; Linz, J.E. Comparative Mapping of Aflatoxin Pathway Gene Clusters in *Aspergillus Parasiticus* and *Aspergillus Flavus*. *Appl. Environ. Microbiol.* **1995**, *61*, 2365–2371. [\[CrossRef\]](#) [\[PubMed\]](#)
91. Tan, D.D.H.S.; Sigler, L.; Gibas, C.; Fong, I.W. Disseminated Fungal Infection in a Renal Transplant Recipient Involving *Macrophomina Phaseolina* and *Scytalidium Dimidiatum*: Case Report and Review of Taxonomic Changes among Medically Important Members of the Botryosphaeriaceae. *Med. Mycol.* **2008**, *46*, 285–292. [\[CrossRef\]](#) [\[PubMed\]](#)
92. Cheng, Y.; Tang, X.; Gao, C.; Li, Z.; Chen, J.; Guo, L.; Wang, T.; Xu, J. Molecular Diagnostics and Detection of Oomycetes on Fiber Crops. *Plants* **2020**, *9*, 769. [\[CrossRef\]](#) [\[PubMed\]](#)
93. Aslam, S.; Tahir, A.; Aslam, M.F.; Alam, M.W.; Shedayi, A.A.; Sadia, S. Recent Advances in Molecular Techniques for the Identification of Phytopathogenic Fungi—A Mini Review. *J. Plant Interact.* **2017**, *12*, 493–504. [\[CrossRef\]](#)
94. Al-masari, S.M.S. Isolation and Identification of the Most Important Aflatoxin B1-Producing Species in Najaf Governorate Respiratory Patients. *Int. J. Health Sci.* **2022**, *6*, 7684–7692. [\[CrossRef\]](#)
95. Mohamed, H.M.A.; Mubarak, A.G.; Khalifa, F.A. Molecular Detection of Aflatoxigenic *A. Flavus* in Chicken Liver with a Special Reference to Aflatoxin B1. *J. Adv. Vet. Res.* **2024**, *14*, 270–275.
96. Khalid, S.; Hussain, N.; Imran, M. Detection of Aflatoxigenicity of *Aspergillus Flavus*, Based on Potential Gene Marker, from Food and Feed Samples. *J. Food Saf.* **2018**, *38*, e12448. [\[CrossRef\]](#)
97. Fakruddin, M.; Chowdhury, A.; Hossain, M.N.; Ahmed, M.M. Characterization of Aflatoxin Producing *Aspergillus Flavus* from Food and Feed Samples. *Springerplus* **2015**, *4*, 159. [\[CrossRef\]](#) [\[PubMed\]](#)
98. Hussain, A.; Afzal, A.; Irfan, M.; Malik, K.A. Molecular Detection of Aflatoxin Producing Strains of *Aspergillus Flavus* from Peanut (*Arachis Hypogaea*). *Turk. J. Agric.—Food Sci. Technol.* **2015**, *3*, 335. [\[CrossRef\]](#)
99. Anidah; Rahayu, W.P.; Nurjanah, S.; Retnowati, I. MOLECULAR CHARACTERIZATION OF *Aspergillus Flavus* TOXIGENICITY IN AGRICULTURAL COMMODITIES IN INDONESIA. *Biotropia* **2023**, *30*, 195–205. [\[CrossRef\]](#)
100. El-Dawy, E.G.A.M.; Gherbawy, Y.A.; Hussein, M.A. Characterization of *Aspergillus* Section *Flavi* Associated with Stored Grains. *Mycotoxin Res.* **2024**, *40*, 187–202. [\[CrossRef\]](#) [\[PubMed\]](#)
101. Almusa, M.A.; Al-Otibi, F.O.; Alreshoodi, F.M.; Alsalman, S.A.; Mukhtar, L.E.; Alharbi, A.L.; Aldosari, Z.M.; Alkaleeb, M.A.; Alarjani, K.M.; Alkhanani, M.F.; et al. Detection and Molecular Characterization of Aflatoxin and Ochratoxin Produce *Aspergillus* Species in Capsicum Spices in Saudi Arabia. *Food Control* **2024**, *162*, 110377. [\[CrossRef\]](#)
102. EL-Shaer, H.; Shoukry, A.; Youness, H. Detection Aflatoxin Production by Local Isolates of *Aspergillus* spp. and Molecular Characterization. *Arch. Agric. Sci. J.* **2021**, *4*, 45–63. [\[CrossRef\]](#)
103. Trinh, H.L.T.; Anh, D.T.; Thong, P.M.; Hue, N.T. A Simple PCR for Detection of *Aspergillus Flavus* in Infected Food. *Qual. Assur. Saf. Crop. Foods* **2015**, *7*, 375–383. [\[CrossRef\]](#)
104. Haggag, W.M.; Diab, M.M.; Al-Ansary, N.A.; Ibrahim, M.I.M.; Khat tab, A.E.N.A.; Abdel-Wahhab, M.A.; Ali, M.K. Molecular Identification and Management of Mycotoxigenic Fungi in Stored Corn Grains. *Cereal Res. Commun.* **2024**, *52*, 1631–1644. [\[CrossRef\]](#)

105. Ahmad, M.M.M.; Ahmad, M.M.M.; Ali, A.; Hamid, R.; Javed, S.; Abdin, M.Z. Detection of *Aspergillus Flavus* and *Aspergillus Parasiticus* from Aflatoxin-Contaminated Peanuts and Their Differentiation Using PCR-RFLP. *Ann. Microbiol.* **2014**, *64*, 1597–1605. [\[CrossRef\]](#)
106. Ortega, S.F.; Siciliano, I.; Prencipe, S.; Gullino, M.L.; Spadaro, D. Development of PCR, LAMP and QPCR Assays for the Detection of Aflatoxigenic Strains of *Aspergillus Flavus* and *A. Parasiticus* in Hazelnut. *Toxins* **2020**, *12*, 757. [\[CrossRef\]](#) [\[PubMed\]](#)
107. Daliri, A.; Shams-Ghahfarokhi, M.; Razzaghi-Abyaneh, M. Detection of Aflatoxin B1-Producing *Aspergillus Flavus* Strains from Pistachio Orchards Soil in Iran by Multiplex Polymerase Chain Reaction Method. *Curr. Med. Mycol.* **2023**, *9*, 1–7. [\[CrossRef\]](#) [\[PubMed\]](#)
108. Chen, R.-S.S.; Tsay, J.-G.G.; Huang, Y.-F.F.; Chiou, R.Y.Y.Y. Polymerase Chain Reaction-Mediated Characterization of Molds Belonging to the *Aspergillus Flavus* Group and Detection of *Aspergillus Parasiticus* in Peanut Kernels by a Multiplex Polymerase Chain Reaction. *J. Food Prot.* **2002**, *65*, 840–844. [\[CrossRef\]](#) [\[PubMed\]](#)
109. Rodrigues, P.; Venâncio, A.; Kozakiewicz, Z.; Lima, N. A Polyphasic Approach to the Identification of Aflatoxigenic and Non-Aflatoxigenic Strains of *Aspergillus* Section *Flavi* Isolated from Portuguese Almonds. *Int. J. Food Microbiol.* **2009**, *129*, 187–193. [\[CrossRef\]](#) [\[PubMed\]](#)
110. Leharanger, A.; Paumier, D.; Orlando, B.; Bailly, S.; Valade, R. Two New QPCR Assays for Detecting and Quantifying the *Aspergillus Flavus* and *Aspergillus Parasiticus* Clades in Maize Kernels. *Plant Pathol.* **2024**, *1*, 2003–2024. [\[CrossRef\]](#)
111. Jamali, M.; Karimipour, M.; Shams-Ghahfarokhi, M.; Amani, A.; Razzaghi-Abyaneh, M. Expression of Aflatoxin Genes *AflO* (*OmtB*) and *AflQ* (*OrdA*) Differentiates Levels of Aflatoxin Production by *Aspergillus Flavus* Strains from Soils of Pistachio Orchards. *Res. Microbiol.* **2013**, *164*, 293–299. [\[CrossRef\]](#) [\[PubMed\]](#)
112. Degola, F.; Berni, E.; Dall'Asta, C.; Spotti, E.; Marchelli, R.; Ferrero, I.; Restivo, F.M. A Multiplex RT-PCR Approach to Detect Aflatoxigenic Strains of *Aspergillus Flavus*. *J. Appl. Microbiol.* **2007**, *103*, 409–417. [\[CrossRef\]](#) [\[PubMed\]](#)
113. Al-Zaban, M.I.; Alrokban, A.H.; Mahmoud, M.A. Development of a Real-Time PCR and Multiplex PCR Assay for the Detection and Identification of Mycotoxigenic Fungi in Stored Maize Grains. *Mycology* **2023**, *14*, 227–238. [\[CrossRef\]](#) [\[PubMed\]](#)
114. Luo, J.; Vogel, R.F.; Niessen, L. Rapid Detection of Aflatoxin Producing Fungi in Food by Real-Time Quantitative Loop-Mediated Isothermal Amplification. *Food Microbiol.* **2014**, *44*, 142–148. [\[CrossRef\]](#) [\[PubMed\]](#)
115. Niessen, L.; Bechtner, J.; Fodil, S.; Taniwaki, M.H.; Vogel, R.F. LAMP-Based Group Specific Detection of Aflatoxin Producers within *Aspergillus* Section *Flavi* in Food Raw Materials, Spices, and Dried Fruit Using Neutral Red for Visible-Light Signal Detection. *Int. J. Food Microbiol.* **2018**, *266*, 241–250. [\[CrossRef\]](#) [\[PubMed\]](#)
116. Luo, J.; Vogel, R.F.; Niessen, L. Development and Application of a Loop-Mediated Isothermal Amplification Assay for Rapid Identification of Aflatoxigenic Molds and Their Detection in Food Samples. *Int. J. Food Microbiol.* **2012**, *159*, 214–224. [\[CrossRef\]](#) [\[PubMed\]](#)
117. Mellikeche, W.; Ricelli, A.; Casini, G.; Gallo, M.; Baser, N.; Colelli, G.; D'Onghia, A.M. Development of Loop-Mediated Isothermal Amplification (LAMP) Assays for the Rapid Detection of Toxigenic *Aspergillus Flavus* and *A. Carbonarius* in Nuts. *Int. J. Mol. Sci.* **2024**, *25*, 3809. [\[CrossRef\]](#) [\[PubMed\]](#)
118. Arnheim, N.; Erlich, H. Polymerase Chain Reaction Strategy. *Annu. Rev. Biochem.* **1992**, *61*, 131–156. [\[CrossRef\]](#)
119. White, T.J.; Bruns, T.D.; Lee, S.B.; Taylor, J.W. Amplification and Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics. *PCR Protoc. Guide Methods Appl.* **1990**, *18*, 315–322. [\[CrossRef\]](#) [\[PubMed\]](#)
120. Zhao, J.; Kong, F.; Li, R.; Wang, X.; Wan, Z.; Wang, D. Identification of *Aspergillus Fumigatus* and Related Species by Nested PCR Targeting Ribosomal DNA Internal Transcribed Spacer Regions. *J. Clin. Microbiol.* **2001**, *39*, 2261–2266. [\[CrossRef\]](#) [\[PubMed\]](#)
121. González-Salgado, A.; Patiño, B.; Vázquez, C.; González-Jaén, M.T. Discrimination of *Aspergillus Niger* and Other *Aspergillus* Species Belonging to Section *Nigri* by PCR Assays. *FEMS Microbiol. Lett.* **2005**, *245*, 353–361. [\[CrossRef\]](#) [\[PubMed\]](#)
122. Sugita, C.; Makimura, K.; Uchida, K.; Yamaguchi, H.; Nagai, A. PCR Identification System for the Genus *Aspergillus* and Three Major Pathogenic Species: *Aspergillus Fumigatus*, *Aspergillus Flavus* and *Aspergillus Niger*. *Med. Mycol.* **2004**, *42*, 433–437. [\[CrossRef\]](#) [\[PubMed\]](#)
123. Makimura, K. Species Identification System for Dermatophytes Based on the DNA Sequences of Nuclear Ribosomal Internal Transcribed Spacer 1. *Nihon Ishinkin Gakkai Zasshi* **2001**, *42*, 61–67. [\[CrossRef\]](#) [\[PubMed\]](#)
124. Zhou, S.; Ismail, M.A.I.; Aimananda, V.; de Hoog, G.S.; Kang, Y.; Ahmed, S.A. Aflatoxin Profiles of *Aspergillus Flavus* Isolates in Sudanese Fungal Rhinosinusitis. *Med. Mycol.* **2024**, *62*, myae034. [\[CrossRef\]](#) [\[PubMed\]](#)
125. Tam, E.W.T.; Chen, J.H.K.; Lau, E.C.L.; Ngan, A.H.Y.; Fung, K.S.C.; Lee, K.C.; Lam, C.W.; Yuen, K.Y.; Lau, S.K.P.; Woo, P.C.Y. Misidentification of *Aspergillus Nomius* and *Aspergillus Tamaris* as *Aspergillus Flavus*: Characterization by Internal Transcribed Spacer, β -Tubulin, and Calmodulin Gene Sequencing, Metabolic Fingerprinting, and Matrix-Assisted Laser Desorption Ionization-Ti. *J. Clin. Microbiol.* **2014**, *52*, 1153–1160. [\[CrossRef\]](#) [\[PubMed\]](#)
126. Manonmani, H.K.; Anand, S.; Chandrashekar, A.; Rati, E.R. Detection of Aflatoxigenic Fungi in Selected Food Commodities by PCR. *Process Biochem.* **2005**, *40*, 2859–2864. [\[CrossRef\]](#)

127. Musangi, C.R.; Juma, B.S.; Mukhebi, D.W.; Isoe, E.M.; Kibiti, C.M.; Mbinda, W.M. Aspergillus Population Diversity and Its Role in Aflatoxin Contamination of Cashew Nuts from Coastal Kenya. *PLoS ONE* **2024**, *19*, e0292519. [[CrossRef](#)] [[PubMed](#)]
128. Hassan, K.I. Molecular Techniques for Detection of Aflatoxigenic Aspergillus Flavus and Determine Their Aflatoxin in Pistachios. *Iraqi J. Agric. Sci.* **2023**, *54*, 1773–1783. [[CrossRef](#)]
129. Bluhm, B.H.; Flaherty, J.E.; Cousin, M.A.; Woloshuk, C.P. Multiplex Polymerase Chain Reaction Assay for the Differential Detection of Trichothecene- and Fumonisin-Producing Species of Fusarium in Cornmeal. *J. Food Prot.* **2002**, *65*, 1955–1961. [[CrossRef](#)] [[PubMed](#)]
130. Jurado, M.; Vázquez, C.; Marín, S.; Sanchis, V.; Teresa González-Jaén, M. PCR-Based Strategy to Detect Contamination with Mycotoxigenic Fusarium Species in Maize. *Syst. Appl. Microbiol.* **2006**, *29*, 681–689. [[CrossRef](#)] [[PubMed](#)]
131. Mayer, Z.; Bagnara, A.; Färber, P.; Geisen, R. Quantification of the Copy Number of Nor-1, a Gene of the Aflatoxin Biosynthetic Pathway by Real-Time PCR, and Its Correlation to the CfU of Aspergillus Flavus in Foods. *Int. J. Food Microbiol.* **2003**, *82*, 143–151. [[CrossRef](#)] [[PubMed](#)]
132. Mitema, A.; Feto, N.A.; Okoth, S.; Rafudeen, M.S. A Tool for Reliable Detection of Aflatoxin Biosynthetic Gene Clusters in Aflatoxigenic and Atoxigenic. *bioRxiv* **2021**. [[CrossRef](#)]
133. Gachon, C.; Mingam, A.; Charrier, B. Real-Time PCR: What Relevance to Plant Studies? *J. Exp. Bot.* **2004**, *55*, 1445–1454. [[CrossRef](#)] [[PubMed](#)]
134. Shweta, S.; Madhavan, S.; Paranidharan, V.; Velazhahan, R. Detection of Aspergillus Flavus in Maize Kernels by Conventional and Real-Time PCR Assays. *Int. Food Res. J.* **2013**, *20*, 3329–3335.
135. Levin, R.E. PCR Detection of Aflatoxin Producing Fungi and Its Limitations. *Int. J. Food Microbiol.* **2012**, *156*, 1–6. [[CrossRef](#)] [[PubMed](#)]
136. Luo, Y.; Gao, W.; Doster, M.; Michailides, T.J. Quantification of Conidial Density of Aspergillus Flavus and a Parasiticus in Soil from Almond Orchards Using Real-Time PCR. *J. Appl. Microbiol.* **2009**, *106*, 1649–1660. [[CrossRef](#)] [[PubMed](#)]
137. Godet, M.; Munaut, F. Molecular Strategy for Identification in Aspergillus Section Flavi: RESEARCH LETTER. *FEMS Microbiol. Lett.* **2010**, *304*, 157–168. [[CrossRef](#)] [[PubMed](#)]
138. Rodríguez, A.; Rodríguez, M.; Luque, M.I.; Martín, A.; Córdoba, J.J. Real-Time PCR Assays for Detection and Quantification of Aflatoxin-Producing Molds in Foods. *Food Microbiol.* **2012**, *31*, 89–99. [[CrossRef](#)] [[PubMed](#)]
139. Notomi, T.; Okayama, H.; Masubuchi, H.; Yonekawa, T.; Watanabe, K.; Amino, N.; Hase, T. Loop-Mediated Isothermal Amplification of DNA. *Nucleic Acids Res.* **2000**, *28*, e63. [[CrossRef](#)] [[PubMed](#)]
140. Tao, F.; Yao, H.; Hruska, Z.; Burger, W.; Rajasekaran, K.; Bhatnagar, D.; Burger, L.W.; Rajasekaran, K.; Bhatnagar, D. Recent Development of Optical Methods in Rapid and Non-Destructive Detection of Aflatoxin and Fungal Contamination in Agricultural Products. *Trends Anal. Chem.* **2018**, *100*, 65–81. [[CrossRef](#)]
141. Karunakaran, C.; Madasamy, T.; Sethy, N.K. Enzymatic Biosensors. In *Biosensors and Bioelectronics*; Elsevier: Amsterdam, The Netherlands, 2015; pp. 133–204.
142. Alhadrami, H.A. Biosensors: Classifications, Medical Applications, and Future Prospective. *Biotechnol. Appl. Biochem.* **2018**, *65*, 497–508. [[PubMed](#)]
143. Alonso-Lomillo, M.A.; Domínguez-Renedo, O.; del Torno-de Román, L.; Arcos-Martínez, M.J. Horseradish Peroxidase-Screen Printed Biosensors for Determination of Ochratoxin A. *Anal. Chim. Acta* **2011**, *688*, 49–53. [[CrossRef](#)] [[PubMed](#)]
144. Chandran, K.; Rajkumar, R.; Bhargava, K. Introduction to Biosensors. In *Biosensors and Bioelectronics*; Elsevier: Amsterdam, The Netherlands, 2015; pp. 1–68, ISBN 9780128031001.
145. Amine, A.; Arduini, F.; Moscone, D.; Palleschi, G. Recent Advances in Biosensors Based on Enzyme Inhibition. *Biosens. Bioelectron.* **2016**, *76*, 180–194. [[CrossRef](#)] [[PubMed](#)]
146. Ispas, C.R.; Crivat, G.; Andreescu, S. Review: Recent Developments in Enzyme-Based Biosensors for Biomedical Analysis. *Anal. Lett.* **2012**, *45*, 168–186. [[CrossRef](#)]
147. Biosensors, I.; Asal, M.; Özen, Ö.; Şahinler, M.; Polatoğlu, İ. Recent Developments in Enzyme, DNA and Immuno-Based Biosensors. *Sensors* **2018**, *18*, 1924. [[CrossRef](#)] [[PubMed](#)]
148. Zachetti, V.G.L.; Granero, A.M.; Robledo, S.N.; Zon, M.A.; Fernández, H. Development of an Amperometric Biosensor Based on Peroxidases to Quantify Citrinin in Rice Samples. *Bioelectrochemistry* **2013**, *91*, 37–43. [[CrossRef](#)] [[PubMed](#)]
149. Moscone, D.; Arduini, F.; Amine, A. A Rapid Enzymatic Method for Aflatoxin B Detection. *Methods Mol. Biol.* **2011**, *739*, 217–235. [[CrossRef](#)] [[PubMed](#)]
150. Egbunike, G.N.; Ikegwuonu, F.I. Effect of Aflatoxicosis on Acetylcholinesterase Activity in the Brain and Adenohypophysis of the Male Rat. *Neurosci. Lett.* **1984**, *52*, 171–174. [[CrossRef](#)] [[PubMed](#)]
151. Xu, X.; Liu, X.; Li, Y.; Ying, Y. A Simple and Rapid Optical Biosensor for Detection of Aflatoxin B1 Based on Competitive Dispersion of Gold Nanorods. *Biosens. Bioelectron.* **2013**, *47*, 361–367. [[CrossRef](#)] [[PubMed](#)]
152. Homola, J.; Yee, S.S.; Gauglitz, G. Surface Plasmon Resonance Sensors: Review. *Sens. Actuators B Chem.* **1999**, *54*, 3–15. [[CrossRef](#)]

153. Daly, S.J.; Keating, G.J.; Dillon, P.P.; Manning, B.M.; O’Kennedy, R.; Lee, H.A.; Morgan, M.R. Development of Surface Plasmon Resonance-Based Immunoassay for Aflatoxin B(1). *J. Agric. Food Chem.* **2000**, *48*, 5097–5104. [[CrossRef](#)] [[PubMed](#)]
154. Goryacheva, I.Y.; De Saeger, S.; Eremin, S.A.; Van Peteghem, C. Immunochemical Methods for Rapid Mycotoxin Detection: Evolution from Single to Multiple Analyte Screening: A Review. *Food Addit. Contam.* **2007**, *24*, 1169–1183. [[CrossRef](#)] [[PubMed](#)]
155. Piringçi, Ş.Ş.; Ertekin, Ö.; Laguna, D.E.; Özen, F.Ş.; Öztürk, Z.Z.; Öztürk, S. Label-Free QCM Immunosensor for the Detection of Ochratoxin A. *Sensors* **2018**, *18*, 1161. [[CrossRef](#)] [[PubMed](#)]
156. Peltomaa, R.; Benito-Peña, E.; Moreno-Bondi, M.C. Bioinspired Recognition Elements for Mycotoxin Sensors. *Anal. Bioanal. Chem.* **2018**, *410*, 747–771. [[CrossRef](#)] [[PubMed](#)]
157. Pfeiffer, F.; Mayer, G. Selection and Biosensor Application of Aptamers for Small Molecules. *Front. Chem.* **2016**, *4*, 25. [[CrossRef](#)] [[PubMed](#)]
158. McKeague, M.; Bradley, C.R.; De Girolamo, A.; Visconti, A.; Miller, J.D.; Derosa, M.C. Screening and Initial Binding Assessment of Fumonisin b(1) Aptamers. *Int. J. Mol. Sci.* **2010**, *11*, 4864–4881. [[CrossRef](#)] [[PubMed](#)]
159. Chen, X.; Bai, X.; Li, H.; Zhang, B. Aptamer-Based Microcantilever Array Biosensor for Detection of Fumonisin B-1. *RSC Adv.* **2015**, *5*, 35448–35452. [[CrossRef](#)]
160. Sabet, F.S.; Hosseini, M.; Khabbaz, H.; Dadmehr, M.; Ganjali, M.R. FRET-Based Aptamer Biosensor for Selective and Sensitive Detection of Aflatoxin B1 in Peanut and Rice. *Food Chem.* **2017**, *220*, 527–532. [[CrossRef](#)] [[PubMed](#)]
161. Lu, Z.; Chen, X.; Wang, Y.; Zheng, X.; Li, C.M. Aptamer Based Fluorescence Recovery Assay for Aflatoxin B1 Using a Quencher System Composed of Quantum Dots and Graphene Oxide. *Microchim. Acta* **2015**, *182*, 571–578. [[CrossRef](#)]
162. Yao, Y.; Wang, H.; Wang, X.; Wang, X.; Li, F. Development of a Chemiluminescent Aptasensor for Ultrasensitive and Selective Detection of Aflatoxin B1 in Peanut and Milk. *Talanta* **2019**, *201*, 52–57. [[CrossRef](#)] [[PubMed](#)]
163. Yang, C.; Wang, Y.; Marty, J.-L.; Yang, X. Aptamer-Based Colorimetric Biosensing of Ochratoxin A Using Unmodified Gold Nanoparticles Indicator. *Biosens. Bioelectron.* **2011**, *26*, 2724–2727. [[CrossRef](#)] [[PubMed](#)]
164. Wu, J.; Chu, H.; Mei, Z.; Deng, Y.; Xue, F.; Zheng, L.; Chen, W. Ultrasensitive One-Step Rapid Detection of Ochratoxin A by the Folding-Based Electrochemical Aptasensor. *Anal. Chim. Acta* **2012**, *753*, 27–31. [[CrossRef](#)] [[PubMed](#)]
165. Guo, Z.; Ren, J.; Wang, J.; Wang, E. Single-Walled Carbon Nanotubes Based Quenching of Free FAM-Aptamer for Selective Determination of Ochratoxin A. *Talanta* **2011**, *85*, 2517–2521. [[CrossRef](#)] [[PubMed](#)]
166. Chen, X.; Huang, Y.; Duan, N.; Wu, S.; Ma, X.; Xia, Y.; Zhu, C.; Jiang, Y.; Wang, Z. Selection and Identification of SsDNA Aptamers Recognizing Zearalenone. *Anal. Bioanal. Chem.* **2013**, *405*, 6573–6581. [[CrossRef](#)] [[PubMed](#)]
167. Duan, H.; Li, Y.; Shao, Y.; Huang, X.; Xiong, Y. Multicolor Quantum Dot Nanobeads for Simultaneous Multiplex Immunochromatographic Detection of Mycotoxins in Maize. *Sens. Actuators B Chem.* **2019**, *291*, 411–417. [[CrossRef](#)]
168. Guo, X.; Wen, F.; Zheng, N.; Luo, Q.; Wang, H.; Wang, H.; Li, S.; Wang, J. Development of an Ultrasensitive Aptasensor for the Detection of Aflatoxin B1. *Biosens. Bioelectron.* **2014**, *56*, 340–344. [[CrossRef](#)] [[PubMed](#)]
169. Chen, J.; Wen, J.; Zhuang, L.; Zhou, S. An Enzyme-Free Catalytic DNA Circuit for Amplified Detection of Aflatoxin B1 Using Gold Nanoparticles as Colorimetric Indicators. *Nanoscale* **2016**, *8*, 9791–9797. [[CrossRef](#)] [[PubMed](#)]
170. Poeta, E.; Liboà, A.; Mistrali, S.; Núñez-Carmona, E.; Sberveglieri, V. Nanotechnology and E-Sensing for Food Chain Quality and Safety. *Sensors* **2023**, *23*, 8429. [[CrossRef](#)] [[PubMed](#)]
171. Garbacz, M.; Malec, A.; Duda-Saternus, S.; Suchorab, Z.; Guz, Ł.; Łagód, G. Methods for Early Detection of Microbiological Infestation of Buildings Based on Gas Sensor Technologies. *Chemosensors* **2020**, *8*, 7. [[CrossRef](#)]
172. Tomić, M.; Šetka, M.; Vojkúvka, L.; Vallejos, S. VOCs Sensing by Metal Oxides, Conductive Polymers, and Carbon-Based Materials. *Nanomaterials* **2021**, *11*, 552. [[CrossRef](#)] [[PubMed](#)]
173. Xu, S.; Wang, H.; Liang, X.; Lu, H. Research Progress on Methods for Improving the Stability of Non-Destructive Testing of Agricultural Product Quality. *Foods* **2024**, *13*, 3917. [[CrossRef](#)] [[PubMed](#)]
174. Okechukwu, V.O.; Mavumengwana, V.; Hümmelgen, I.A.; Mamo, M.A. Concomitant in Situ FTIR and Impedance Measurements To Address the 2-Methylcyclopentanone Vapor-Sensing Mechanism in MnO₂-Polymer Nanocomposites. *ACS Omega* **2019**, *4*, 8324–8333. [[CrossRef](#)] [[PubMed](#)]
175. Okechukwu, V.O.; Mavumengwana, V.; Mamo, M.A. A Direct Relationship between the Sensitivity of the Sensors and the Intensity of IR CO₂ Peak in in Situ FTIR-LCR Meter Chemi-Impedance SnO₂-Carbon Nanoparticles Polymer-Based Sensors in the Detection of Organic Compounds Vapor. *AIP Adv.* **2021**, *11*, 105203. [[CrossRef](#)]
176. Onyinye Okechukwu, V.; Olayiwola Idris, A.; Umukoro, E.H.; Azizi, S.; Maaza, M. Exploring the Contribution of Intelligent Nanomaterials in Gas Sensing. *ChemistrySelect* **2024**, *9*, e202304703. [[CrossRef](#)]
177. Malepe, L.; Ndinteh, D.T.; Ndungu, P.; Mamo, M.A. Selective Detection of Methanol Vapour from a Multicomponent Gas Mixture Using a CNPs/ZnO@ZIF-8 Based Room Temperature Solid-State Sensor. *RSC Adv.* **2022**, *12*, 27094–27108. [[CrossRef](#)] [[PubMed](#)]
178. Kaur, N.; Singh, M.; Comini, E. Materials Engineering Strategies to Control Metal Oxides Nanowires Sensing Properties. *Adv. Mater. Interfaces* **2022**, *9*, 2101629. [[CrossRef](#)]

179. Feng, Y.; Liu, Y.; Dai, H.; Deng, J. Review and Perspectives of Enhancement in the Catalytic Stability for the Complete Combustion of CO, CH₄, and Volatile Organic Compounds. *Energy Fuels* **2023**, *37*, 3590–3604. [\[CrossRef\]](#)
180. Chu, S.; Wang, E.; Feng, F.; Zhang, C.; Jiang, J.; Zhang, Q.; Wang, F.; Bing, L.; Wang, G.; Han, D. A Review of Noble Metal Catalysts for Catalytic Removal of VOCs. *Catalysts* **2022**, *12*, 1543. [\[CrossRef\]](#)
181. Wei, Y.; Han, C.; Yu, Z. An Environment Safety Monitoring System for Agricultural Production Based on Artificial Intelligence, Cloud Computing and Big Data Networks. *J. Cloud Comput.* **2023**, *12*, 83. [\[CrossRef\]](#)
182. Ueno, T.; Nagano, T. Fluorescent Probes for Sensing and Imaging. *Nat. Methods* **2011**, *8*, 642–645. [\[CrossRef\]](#) [\[PubMed\]](#)
183. He, H.J.; Sun, D.W. Hyperspectral Imaging Technology for Rapid Detection of Various Microbial Contaminants in Agricultural and Food Products. *Trends Food Sci. Technol.* **2015**, *46*, 99–109. [\[CrossRef\]](#)
184. Maupin, L.M.; Clements, M.J.; White, D.G. Evaluation of the MI82 Corn Line as a Source of Resistance to Aflatoxin in Grain and Use of BGYF as a Selection Tool. *Plant Dis.* **2003**, *87*, 1059–1066. [\[CrossRef\]](#) [\[PubMed\]](#)
185. Wu, Q.; Xu, H. Design and Development of an On-Line Fluorescence Spectroscopy System for Detection of Aflatoxin in Pistachio Nuts. *Postharvest Biol. Technol.* **2020**, *159*, 111016. [\[CrossRef\]](#)
186. Doster, M.A.; Michailides, T.J. Production of Bright Greenish Yellow Fluorescence in Figs Infected by *Aspergillus* Species in California Orchards. *Plant Dis.* **1998**, *82*, 669–673. [\[CrossRef\]](#) [\[PubMed\]](#)
187. Kalkan, H.; Beriat, P.; Yardimci, Y.; Pearson, T.C. Detection of Contaminated Hazelnuts and Ground Red Chili Pepper Flakes by Multispectral Imaging. *Comput. Electron. Agric.* **2011**, *77*, 28–34. [\[CrossRef\]](#)
188. Wicklow, D.T. Influence of *Aspergillus Flavus* Strains on Aflatoxin and Bright Greenish Yellow Fluorescence of Corn Kernels. *Plant Dis.* **1999**, *83*, 1146–1148. [\[CrossRef\]](#) [\[PubMed\]](#)
189. Sugiyama, J.; Fujita, K.; Tsuta, M.; Kushiro, M. Detection of Deoxynivalenol in Wheat Flour Using Fluorescence Fingerprint. *Procedia Food Sci.* **2011**, *1*, 1146–1151. [\[CrossRef\]](#)
190. Sajjadi, S.M.; Abdollahi, H.; Rahmanian, R.; Bagheri, L. Quantifying Aflatoxins in Peanuts Using Fluorescence Spectroscopy Coupled with Multi-Way Methods: Resurrecting Second-Order Advantage in Excitation–Emission Matrices with Rank Overlap Problem. *Spectrochim. Acta Part A Mol. Biomol. Spectrosc.* **2016**, *156*, 63–69. [\[CrossRef\]](#) [\[PubMed\]](#)
191. Liao, W.; Lu, X. Determination of Chemical Hazards in Foods Using Surface-Enhanced Raman Spectroscopy Coupled with Advanced Separation Techniques. *Trends Food Sci. Technol.* **2016**, *54*. [\[CrossRef\]](#)
192. Dies, H.; Siampani, M.; Escobedo, C.; Docoslis, A. Direct Detection of Toxic Contaminants in Minimally Processed Food Products Using Dendritic Surface-Enhanced Raman Scattering Substrates. *Sensors* **2018**, *18*, 2726. [\[CrossRef\]](#) [\[PubMed\]](#)
193. Law, J.W.-F.; Ab Mutalib, N.-S.; Chan, K.-G.; Lee, L.-H. Rapid Methods for the Detection of Foodborne Bacterial Pathogens: Principles, Applications, Advantages and Limitations. *Front. Microbiol.* **2014**, *5*, 770. [\[CrossRef\]](#) [\[PubMed\]](#)
194. Wang, K.; Chen, L.; Ma, X.; Ma, L.; Chou, K.C.; Cao, Y.; Khan, I.U.H.; Gözl, G.; Lu, X. *Arcobacter* Identification and Species Determination Using Raman Spectroscopy Combined with Neural Networks. *Appl. Environ. Microbiol.* **2020**, *86*. [\[CrossRef\]](#) [\[PubMed\]](#)
195. Jaafreh, S.; Valler, O.; Kreyenschmidt, J.; Günther, K.; Kaul, P. In Vitro Discrimination and Classification of Microbial Flora of Poultry Using Two Dispersive Raman Spectrometers (Microscope and Portable Fiber-Optic Systems) in Tandem with Chemometric Analysis. *Talanta* **2019**, *202*, 411–425. [\[CrossRef\]](#) [\[PubMed\]](#)
196. Yan, S.; Wang, S.; Qiu, J.; Li, M.; Li, D.; Xu, D.; Li, D.; Liu, Q. Raman Spectroscopy Combined with Machine Learning for Rapid Detection of Food-Borne Pathogens at the Single-Cell Level. *Talanta* **2021**, *226*, 122195. [\[CrossRef\]](#) [\[PubMed\]](#)
197. Srey, S.; Jahid, I.; Ha, S.-D. Biofilm Formation in Food Industries: A Food Safety Concern. *Food Control* **2013**, *31*, 572–585. [\[CrossRef\]](#)
198. Oliver, J.D. The Viable but Nonculturable State in Bacteria. *J. Microbiol.* **2005**, *43 Spec No*, 93–100. [\[PubMed\]](#)
199. Achadu, O.J.; Abe, F.; Suzuki, T.; Park, E.Y. Molybdenum Trioxide Nanocubes Aligned on a Graphene Oxide Substrate for the Detection of Norovirus by Surface-Enhanced Raman Scattering. *ACS Appl. Mater. Interfaces* **2020**, *12*, 43522–43534. [\[CrossRef\]](#) [\[PubMed\]](#)
200. Guo, Z.; Wang, M.; Barimah, A.O.; Chen, Q.; Li, H.; Shi, J.; El-Seedi, H.R.; Zou, X. Label-Free Surface Enhanced Raman Scattering Spectroscopy for Discrimination and Detection of Dominant Apple Spoilage Fungus. *Int. J. Food Microbiol.* **2021**, *338*, 108990. [\[CrossRef\]](#) [\[PubMed\]](#)
201. Neng, J.; Zhang, Q.; Sun, P. Application of Surface-Enhanced Raman Spectroscopy in Fast Detection of Toxic and Harmful Substances in Food. *Biosens. Bioelectron.* **2020**, *167*, 112480. [\[CrossRef\]](#) [\[PubMed\]](#)
202. Chauhan, R.; Singh, J.; Sachdev, T.; Basu, T.; Malhotra, B.D. Recent Advances in Mycotoxins Detection. *Biosens. Bioelectron.* **2016**, *81*, 532–545. [\[CrossRef\]](#) [\[PubMed\]](#)
203. Xue, Z.; Zhang, Y.; Yu, W.; Zhang, J.; Wang, J.; Wan, F.; Kim, Y.; Liu, Y.; Kou, X. Recent Advances in Aflatoxin B1 Detection Based on Nanotechnology and Nanomaterials—A Review. *Anal. Chim. Acta* **2019**, *1069*, 1–27. [\[CrossRef\]](#) [\[PubMed\]](#)

204. Goud, K.Y.; Kailasa, S.K.; Kumar, V.; Tsang, Y.F.; Lee, S.E.; Gobi, K.V.; Kim, K.-H. Progress on Nanostructured Electrochemical Sensors and Their Recognition Elements for Detection of Mycotoxins: A Review. *Biosens. Bioelectron.* **2018**, *121*, 205–222. [[CrossRef](#)] [[PubMed](#)]
205. Peng, Z.; Chen, L.; Zhu, Y.; Huang, Y.; Hu, X.; Wu, Q.; Nüssler, A.K.; Liu, L.; Yang, W. Current Major Degradation Methods for Aflatoxins: A Review. *Trends Food Sci. Technol.* **2018**, *80*, 155–166. [[CrossRef](#)]
206. Xu, W.; Mao, N.; Zhang, J. Graphene: A Platform for Surface-Enhanced Raman Spectroscopy. *Small* **2013**, *9*, 1206–1224. [[CrossRef](#)] [[PubMed](#)]
207. Bell, S.E.J.; Charron, G.; Cortés, E.; Kneipp, J.; de la Chapelle, M.L.; Langer, J.; Procházka, M.; Tran, V.; Schlücker, S. Towards Reliable and Quantitative Surface-Enhanced Raman Scattering (SERS): From Key Parameters to Good Analytical Practice. *Angew. Chem. Int. Ed.* **2020**, *59*, 5454–5462. [[CrossRef](#)] [[PubMed](#)]
208. Jiao, T.; Mehedi Hassan, M.; Zhu, J.; Ali, S.; Ahmad, W.; Wang, J.; Lv, C.; Chen, Q.; Li, H. Quantification of Deltamethrin Residues in Wheat by Ag@ZnO NFs-Based Surface-Enhanced Raman Spectroscopy Coupling Chemometric Models. *Food Chem.* **2021**, *337*, 127652. [[CrossRef](#)] [[PubMed](#)]
209. Zhang, X.; Wu, D.; Zhou, X.; Yu, Y.; Liu, J.; Hu, N.; Wang, H.; Li, G.; Wu, Y. Recent Progress in the Construction of Nanozyme-Based Biosensors and Their Applications to Food Safety Assay. *TrAC Trends Anal. Chem.* **2019**, *121*, 115668. [[CrossRef](#)]
210. Zhai, W.; You, T.; Ouyang, X.; Wang, M. Recent Progress in Mycotoxins Detection Based on Surface-Enhanced Raman Spectroscopy. *Compr. Rev. Food Sci. Food Saf.* **2021**, *20*, 1887–1909. [[CrossRef](#)] [[PubMed](#)]
211. Wei, X.; Xu, N.; Wu, D.; He, Y. Determination of Branched-Amino Acid Content in Fermented Cordyceps Sinensis Mycelium by Using FT-NIR Spectroscopy Technique. *Food Bioprocess Technol.* **2014**, *7*, 184–190. [[CrossRef](#)]
212. Hyunjung, M.; Byoung-Kwan, C. Spectroscopic Techniques for Nondestructive Detection of Fungi and Mycotoxins in Agricultural Materials: A Review. *J. Biosyst. Eng.* **2015**, *40*, 67–77. [[CrossRef](#)]
213. Siripatrawan, U.; Makino, Y. Monitoring Fungal Growth on Brown Rice Grains Using Rapid and Non-Destructive Hyperspectral Imaging. *Int. J. Food Microbiol.* **2015**, *199*, 93–100. [[CrossRef](#)] [[PubMed](#)]
214. Burks, C.S.; Dowell, F.E.; Xie, F. Measuring Fig Quality Using Near-Infrared Spectroscopy. *J. Stored Prod. Res.* **2000**, *36*, 289–296. [[CrossRef](#)] [[PubMed](#)]
215. Moschetti, R.; Monarca, D.; Cecchini, M.; Haff, R.P.; Contini, M.; Massantini, R. Detection of Mold-Damaged Chestnuts by Near-Infrared Spectroscopy. *Postharvest Biol. Technol.* **2014**, *93*, 83–90. [[CrossRef](#)]
216. Rodriguez-Saona, L.E.; Khambaty, F.M.; Fry, F.S.; Dubois, J.; Calvey, E.M. Detection and Identification of Bacteria in a Juice Matrix with Fourier Transform–Near Infrared Spectroscopy and Multivariate Analysis. *J. Food Prot.* **2004**, *67*, 2555–2559. [[CrossRef](#)] [[PubMed](#)]
217. Durmuş, E.; Bilgi, A.S.; Ortac, G.; Kalkan, H.; Taşdemir, K. Detection of Black Mold Infected Figs by Using Transmittance Spectroscopy. In Proceedings of the 2015 7th Workshop on Hyperspectral Image and Signal Processing: Evolution in Remote Sensing (WHISPERS), Tokyo, Japan, 2–5 June 2015; pp. 1–4.
218. Durmuş, E.; Güneş, A.; Kalkan, H. Detection of Aflatoxin and Surface Mould Contaminated Figs by Using Fourier Transform Near-infrared Reflectance Spectroscopy. *J. Sci. Food Agric.* **2017**, *97*, 317–323. [[CrossRef](#)] [[PubMed](#)]
219. Goetz, A.F.H.; Vane, G.; Solomon, J.E.; Rock, B.N. Imaging Spectrometry for Earth Remote Sensing. *Science* **1985**, *228*, 1147–1153. [[CrossRef](#)] [[PubMed](#)]
220. Jiang, L.; Zhu, B.; Jing, H.; Chen, X.; Rao, X.; Tao, Y. Gaussian Mixture Model-Based Walnut Shell and Meat Classification in Hyperspectral Fluorescence Imagery. *Trans. ASABE* **2007**, *50*, 153–160. [[CrossRef](#)]
221. Lefcourt, A.M.; Kim, M.S.; Chen, Y.R. Detection of fecal contamination in apple calyx by multispectral laser-induced fluorescence. *Trans. ASAE* **2005**, *48*, 1587–1594. [[CrossRef](#)]
222. Chu, X.; Wang, W.; Ni, X.; Zheng, H.; Zhao, X.; Zhang, R.; Li, Y. Growth Identification of *Aspergillus Flavus* and *Aspergillus Parasiticus* by Visible/Near-Infrared Hyperspectral Imaging. *Appl. Sci.* **2018**, *8*, 513. [[CrossRef](#)]
223. Senthilkumar, T.; Jayas, D.S.; White, N.D.G.; Fields, P.G.; Gräfenhan, T. Near-Infrared (NIR) Hyperspectral Imaging: Theory and Applications to Detect Fungal Infection and Mycotoxin Contamination in Food Products. *Indian J. Entomol.* **2016**, *78*, 91–99. [[CrossRef](#)]
224. Soni, A.; Reis, M.M.; Brightwell, G. Hyperspectral Imaging and Machine Learning in Food Microbiology: Developments and Challenges in Detection of Bacterial, Fungal, and Viral Contaminants. *Compr. Rev. Food Sci. Food Saf.* **2022**, *3717–3745*. [[CrossRef](#)] [[PubMed](#)]
225. Parrag, V.; Gillay, Z.; Kovács, Z.; Zitek, A.; Böhm, K.; Hinterstoisser, B.; Krska, R.; Sulyok, M.; Felföldi, J.; Firtha, F.; et al. Application of Hyperspectral Imaging to Detect Toxigenic *Fusarium* Infection on Cornmeal. *Prog. Agric. Eng. Sci.* **2020**, *16*, 51–60. [[CrossRef](#)]
226. Bauriegel, E.; Giebel, A.; Geyer, M.; Schmidt, U.; Herppich, W.B. Early Detection of *Fusarium* Infection in Wheat Using Hyperspectral Imaging. *Comput. Electron. Agric.* **2011**, *75*, 304–312. [[CrossRef](#)]

227. Shahin, M.; Symons, S. Detection of Fusarium Damage in Canadian Wheat Using Visible/near-Infrared Hyperspectral Imaging. *J. Food Meas. Charact.* **2012**, *6*, 3–11. [\[CrossRef\]](#)
228. Mathew, T.; Manickavasagan, A.; Lankapalli, R.; Jayas, D. Near Infrared (NIR) Hyperspectral Imaging to Classify Fungal Infected Date Fruits. *J. Stored Prod. Res.* **2014**, *59*, 306–313. [\[CrossRef\]](#)
229. Hruska, Z.; Yao, H.; Kincaid, R.; Darlington, D.; Brown, R.L.; Bhatnagar, D.; Cleveland, T.E. Fluorescence Imaging Spectroscopy (FIS) for Comparing Spectra from Corn Ears Naturally and Artificially Infected with Aflatoxin Producing Fungus. *J. Food Sci.* **2013**, *78*, T1313–20. [\[CrossRef\]](#) [\[PubMed\]](#)
230. Yao, H.; Hruska, Z.; Kincaid, R.; Brown, R.; Cleveland, T.; Bhatnagar, D. Correlation and Classification of Single Kernel Fluorescence Hyperspectral Data with Aflatoxin Concentration in Corn Kernels Inoculated with *Aspergillus Flavus* Spores. *Food Addit. Contam. Part A Chem. Anal. Control Expo. Risk Assess.* **2010**, *27*, 701–709. [\[CrossRef\]](#) [\[PubMed\]](#)
231. Menesatti, P.; Antonucci, F.; Pallottino, F.; Giorgi, S.; Matere, A.; Nocente, F.; Pasquini, M.; D'Egidio, M.G.; Costa, C. Laboratory vs. in-Field Spectral Proximal Sensing for Early Detection of Fusarium Head Blight Infection in Durum Wheat. *Biosyst. Eng.* **2013**, *114*, 289–293. [\[CrossRef\]](#)
232. Inglis, A.; Parnell, A.C.; Subramani, N.; Doohan, F.M. Machine Learning Applied to the Detection of Mycotoxin in Food: A Systematic Review. *Toxins* **2024**, *16*, 268. [\[CrossRef\]](#) [\[PubMed\]](#)
233. Tian, X.; Yao, J.; Yu, H.; Wang, W.; Huang, W. Early Contamination Warning of Aflatoxin B1 in Stored Maize Based on the Dynamic Change of Catalase Activity and Data Fusion of Hyperspectral Images. *Comput. Electron. Agric.* **2024**, *217*, 108615. [\[CrossRef\]](#)
234. Zhi, A.; Shi, S.; Li, Q.; Shen, F.; He, X.; Fang, Y.; Hu, Q.; Jiang, X. Aflatoxins Degradation and Quality Evaluation in Naturally Contaminated Rice by Dielectric Barrier Discharge Cold Plasma. *Innov. Food Sci. Emerg. Technol.* **2023**, *88*, 103426. [\[CrossRef\]](#)
235. Kępka-Borkowska, K.; Chałasiński, K.; Ogłuszka, M.; Borkowski, M.; Lepczyński, A.; Pareek, C.S.; Starzyński, R.R.; Lichwiarska, E.; Sultana, S.; Kalra, G. Current Approaches to Aflatoxin B1 Control in Food and Feed Safety: Detection, Inhibition, and Mitigation. *Int. J. Mol. Sci.* **2025**, *26*, 6534. [\[CrossRef\]](#) [\[PubMed\]](#)
236. Focker, M.; Liu, C.; Wang, X.; van der Fels-Klerx, H.J. The Use of Artificial Intelligence to Improve Mycotoxin Management: A Review. *Mycotoxin Res.* **2025**, *41*, 529–540. [\[CrossRef\]](#) [\[PubMed\]](#)
237. Saldan, N.C.; Almeida, R.T.R.R.; Avicola, A.; Porto, C.; Galuch, M.B.; Magon, T.F.S.S.; Pilau, E.J.; Svidzinski, T.I.E.E.; Oliveira, C.C. Development of an Analytical Method for Identification of *Aspergillus Flavus* Based on Chemical Markers Using HPLC-MS. *Food Chem.* **2018**, *241*, 113–121. [\[CrossRef\]](#) [\[PubMed\]](#)
238. Tan, K.-C.; Trengove, R.D.; Maker, G.L.; Oliver, R.P.; Solomon, P.S. Metabolite Profiling Identifies the Mycotoxin Alternariol in the Pathogen *Stagonospora Nodorum*. *Metabolomics* **2009**, *5*, 330–335. [\[CrossRef\]](#)
239. Fiehn, O. Combining Genomics, Metabolome Analysis, and Biochemical Modelling to Understand Metabolic Networks. *Comp. Funct. Genom.* **2001**, *2*, 155–168. [\[CrossRef\]](#) [\[PubMed\]](#)
240. Falade, T.D.O.; Chrysanthopoulos, P.K.; Hodson, M.P.; Sultanbawa, Y.; Fletcher, M.; Darnell, R.; Korie, S.; Fox, G. Metabolites Identified during Varied Doses of *Aspergillus* Species in Zea Mays Grains, and Their Correlation with Aflatoxin Levels. *Toxins* **2018**, *10*, 187. [\[CrossRef\]](#) [\[PubMed\]](#)
241. Varga, J.; Rigó, K.; Tóth, B.; Téren, J.; Kozakiewicz, Z. Evolutionary Relationship among *Aspergillus* Producing Mycotoxins. *Food Technol. Biotechnol.* **2003**, *41*, 29–36.
242. Cary, J.W.; OBrian, G.R.; Nielsen, D.M.; Nierman, W.; Harris-Coward, P.; Yu, J.; Bhatnagar, D.; Cleveland, T.E.; Payne, G.A.; Calvo, A.M. Elucidation of VeA-Dependent Genes Associated with Aflatoxin and Sclerotial Production in *Aspergillus Flavus* by Functional Genomics. *Appl. Microbiol. Biotechnol.* **2007**, *76*, 1107–1118. [\[CrossRef\]](#) [\[PubMed\]](#)
243. Amaike, S.; Keller, N.P. *Aspergillus Flavus*. *Annu. Rev. Phytopathol.* **2011**, *49*, 107–133. [\[CrossRef\]](#) [\[PubMed\]](#)
244. Chen, Z.-Y.; Brown, R.L.; Damann, K.E.; Cleveland, T.E. Identification of Unique or Elevated Levels of Kernel Proteins in Aflatoxin-Resistant Maize Genotypes Through Proteome Analysis. *Phytopathology* **2002**, *92*, 1084–1094. [\[CrossRef\]](#) [\[PubMed\]](#)
245. Moore, G.G.; Mack, B.M.; Beltz, S.B. Genomic Sequence of the Aflatoxigenic Filamentous Fungus *Aspergillus Nomius*. *BMC Genom.* **2015**, *16*, 551. [\[CrossRef\]](#) [\[PubMed\]](#)
246. Kim, J.H.; Yu, J.; Mahoney, N.; Chan, K.L.; Molyneux, R.J.; Varga, J.; Bhatnagar, D.; Cleveland, T.E.; Nierman, W.C.; Campbell, B.C. Elucidation of the Functional Genomics of Antioxidant-Based Inhibition of Aflatoxin Biosynthesis. *Int. J. Food Microbiol.* **2008**, *122*, 49–60. [\[CrossRef\]](#) [\[PubMed\]](#)
247. Yu, J.; Bhatnagar, D.; Cleveland, T.E. Chapter Eleven Genetics and Biochemistry of Aflatoxin Formation and Genomics Approach for Preventing Aflatoxin Contamination. In *Secondary Metabolism in Model Systems*; Romeo, J.T., Ed.; Elsevier: Amsterdam, The Netherlands, 2004; Volume 38, pp. 223–255, ISBN 0079-9920.
248. Dong, Z.; Chen, Y. Transcriptomics: Advances and Approaches. *Sci. China Life Sci.* **2013**, *56*, 960–967. [\[CrossRef\]](#) [\[PubMed\]](#)
249. Pedrotty, D.M.; Morley, M.P.; Cappola, T.P. Transcriptomic Biomarkers of Cardiovascular Disease. *Prog. Cardiovasc. Dis.* **2012**, *55*, 64–69. [\[CrossRef\]](#) [\[PubMed\]](#)
250. Macaulay, I.C.; Carr, P.; Gusnanto, A.; Ouwehand, W.H.; Fitzgerald, D.; Watkins, N.A. Platelet Genomics and Proteomics in Human Health and Disease. *J. Clin. Investig.* **2005**, *115*, 3370–3377. [\[CrossRef\]](#) [\[PubMed\]](#)

251. Musungu, B.M.; Bhatnagar, D.; Brown, R.L.; Payne, G.A.; OBrian, G.; Fakhoury, A.M.; Geisler, M. A Network Approach of Gene Co-Expression in the Zea Mays/Aspergillus Flavus Pathosystem to Map Host/Pathogen Interaction Pathways. *Front. Genet.* **2016**, *7*, 206. [[CrossRef](#)] [[PubMed](#)]
252. Cunha, S.C.; Faria, M.A.; Fernandes, J.O. Determination of Patulin in Apple and Quince Products by GC–MS Using 13C5–7 Patulin as Internal Standard. *Food Chem.* **2009**, *115*, 352–359. [[CrossRef](#)]
253. Bhatnagar, D.; Rajasekaran, K.; Gilbert, M.; Cary, J.W.; Magan, N. Advances in Molecular and Genomic Research to Safeguard Food and Feed Supply from Aflatoxin Contamination. *World Mycotoxin J.* **2018**, *11*, 47–72. [[CrossRef](#)]
254. Razzazi-Fazeli, E.; Rizwan, M.; Mayrhofer, C.; Nöbauer, K. The Use of Proteomics as a Novel Tool in Aflatoxin Research. In *Aflatoxins—Biochemistry and Molecular Biology*; IntechOpen: Rijeka, Croatia, 2011.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.